LETTER TO THE EDITOR

Nebular emission and the Lyman continuum photon escape fraction in CALIFA early-type galaxies *

P. Papaderos¹, J.M. Gomes¹, J.M. Vílchez², C. Kehrig², M.D. Lehnert³, B. Ziegler⁴, S. F. Sánchez^{5, 2}, B. Husemann⁶, A. Monreal-Ibero², R. García-Benito², J. Bland-Hawthorn⁷, C. Coritjo², A. de Lorenzo-Cáceres⁸, A. del Olmo², J. Falcón-Barroso^{9,10}, L. Galbany¹¹, J. Iglesias-Páramo^{2,5}, Á.R. López-Sánchez^{12,13}, I. Marquez², M. Mollá¹⁴, D. Mast^{5,2}, G. van de Ven¹⁵, L. Wisotzki⁶, and the CALIFA collaboration

(Affiliations can be found after the references)

Received 11 April 2013 / Accepted 2 June 2013

ABSTRACT

We use deep integral field spectroscopy data from the CALIFA survey to study the warm interstellar medium (wim) over the entire extent and optical spectral range of 32 nearby early-type galaxies (ETGs). We find that faint nebular emission is extended in all cases, and its surface brightness decreases roughly as $\propto r^{-\alpha}$. The large standard deviation in the derived α (1.09±0.67) argues against a universal power-law index for the radial drop-off of nebular emission in ETGs. Judging from the properties of their extranuclear component, our sample ETGs span a broad, continuous sequence with respect to their α , H α equivalent width (EW) and Lyman continuum (Ly_c) photon leakage fraction (plf). We propose a tentative subdivision into two groups: Type i ETGs are characterized by rather steep H α profiles ($\alpha \simeq 1.4$), comparatively large ($\gtrsim 1$ Å), nearly radially constant EWs, and *plf* $\simeq 0$. Photoionization by post-AGB stars appears to be the main driver of extended nebular emission in these systems, with nonthermal sources being potentially important only in their nuclei. Typical properties of type ii ETGs are shallower H α profiles ($\alpha \simeq 0.8$), very low ($\lesssim 0.5$ Å) EWs with positive radial gradients, and a mean $p l f \gtrsim 0.7$, rising to $\gtrsim 0.9$ in their centers. Such properties point to a low, and inwardly decreasing wim density and/or volume filling factor. We argue that, because of extensive Lyc photon leakage, emission-line luminosities and EWs are reduced in type ii ETG nuclei by at least one order of magnitude. Consequently, the line weakness of these ETGs is by itself no compelling evidence for their containing merely "weak" (sub-Eddington accreting) active galactic nuclei (AGN). In fact, Lyc photon escape, which has heretofore not been considered, may constitute a key element in understanding why many ETGs with prominent signatures of AGN activity in radio continuum and/or X-ray wavelengths show only faint emission lines and weak signatures of AGN activity in their optical spectra. The Lyc photon escape, in conjunction with dilution of nuclear EWs by line-of-sight integration through a triaxial stellar host, can systematically impede detection of AGN in gas-poor galaxy spheroids through optical emissionline spectroscopy, thereby leading to an observational bias. We further find that type i&ii ETGs differ little ($\lesssim 0.4$ dex) in their mean BPT line ratios, which in both cases are characteristic of LINERs and are, within their uncertainties, almost radius-independent. This potentially hints at a degeneracy of the projected, luminosity-weighted BPT ratios in the LINER regime, for the specific 3D properties of the *wim* and the ionizing photon field in ETGs.

Key words. galaxies: elliptical and lenticular, cD - galaxies: nuclei - galaxies: ISM

arXiv:1306.2338v1 [astro-ph.CO] 10 Jun 2013 1. Introduction

Even though the presence of faint nebular emission (ne) in the nuclei of many early-type galaxies (ETGs) has long been established observationally (e.g., Phillips et al. 1986; Sarzi et al. 2006, 2010; Annibali et al. 2010; Kehrig et al. 2012, hereafter K12), the nature of the dominant excitation mechanism of the warm interstellar medium (wim) in these systems remains uncertain. The low-ionization nuclear emission-line region (LINER) emission-line ratios, as a typical property of ETG nuclei, have prompted various interpretations (see, e.g., K12, Yan & Blanton 2012), including low-accretion rate active galactic nuclei (AGN; e.g., Ho 1999), fast shocks (e.g. Dopita & Sutherland 1995), and hot, evolved ($\geq 10^8$ yr) post-AGB (pAGB) stars (e.g., Trinchieri & di Serego Alighieri 1991; Binette et al. 1994; Stasińska et al.

2008). Since each of these mechanisms is tied to distinct and testable expectations on the 2D properties of the wim, the limited spatial coverage of previous single-aperture and longslit spectroscopic studies has been an important obstacle to any conclusive discrimination between them. Spatially resolved integral field spectroscopy (IFS) over the entire extent of ETGs offers an essential advantage in this respect and promises key observational constraints toward the resolution of this longstanding debate.

This Letter gives a brief summary of our results from an ongoing study of 32 ETGs, which were mapped with deep IFS over their entire extent and optical spectral range with the goal of gaining deeper insight into the 2D properties of their wim. A detailed discussion of individual objects and our methodology will be given in Gomes et al. (2013, in prep.; hereafter G13) and subsequent publications of this series. This study is based on low-spectral-resolution ($R \sim 850$) IFS cubes for 20 E and 12 S0 nearby (<150 Mpc) galaxies from the Calar Alto Legacy Integral Field Area (CALIFA) survey (Sánchez et al. 2012, Walcher et al. 2013, in prep.). These data are being made accessible to

Based on observations collected at the Centro Astronómico Hispano Alemán (CAHA) at Calar Alto, operated jointly by the Max-Planck-Institut für Astronomie (MPIA) and the Instituto de Astrofísica de Andalucía (CSIC)

the community in a fully reduced and well-documented format (Husemann et al. 2013) through successive data releases.

2. Methodology and results

The CALIFA data cubes were processed with the Porto3D pipeline (see K12 and G13 for details), which, among various other tasks, permits spaxel-by-spaxel spectral fitting of the stellar component with the population synthesis code STARLIGHT (Cid Fernandes et al. 2005) and subsequent determination of emission line fluxes and their uncertainties from the pure emissionline spectrum (i.e. the observed spectrum after subtraction of the best-fitting synthetic stellar model). For each ETG, typically ~1600 to ~3400 individual spectra with a S/N \geq 30 at 5150 Å were extracted and modeled in the spectral range 4000-6800 Å using both Bruzual & Charlot (2003, hereafter BC) and MILES (Sánchez-Blázquez et al. 2006; Vazdekis et al. 2010) simplestellar population (SSP) libraries, which comprise 34 ages between 5 Myr and 13 Gyr for three metallicities (0.008, 0.019, and 0.03), i.e., 102 elements each. After full analysis and crossinspection of the relevant output from the BC- and MILESbased models, the emission-line maps for each ETG were errorweighted and averaged spaxel-by-spaxel to reduce uncertainties.

An extra module in Porto3D permits computation of the Lyman continuum (Ly_c) ionizing photon rate corresponding to the best-fitting set of BC SSPs for each spaxel. The Ly_c output is then converted into Balmer line luminosities assuming case B recombination for an electron temperature and density of 10⁴ K and 100 cm⁻³, respectively. The same module computes the distance-independent τ ratio of the H α luminosity predicted from pAGB photoionization to the one observed (see Binette et al. 1994; Cid Fernandes et al. 2011, for equivalent quantities). The latter is optionally corrected for intrinsic extinction, assuming this to be equal to the extinction A_V in the stellar component (cf K12 and G13). Since spectral fits imply a low (≤ 0.3 mag) A_V in most cases, this correction typically has a weak effect on τ . We preferred to not base corrections of the τ ratio on nebular extinction estimates since these are consistent with A_V within their uncertainties.

We note that state-of-the-art SSP models imply that the Ly_c photon rate per unit mass from pAGB stellar populations of nearly-solar metallicity (0.008 $\leq Z \leq 0.03$) is almost independent of age, metallicity, and star formation history (e.g. Cid Fernandes et al. 2011, G13). However, substantial uncertainties stem from the fact that existing models differ from one another by a factor ~2 in the mean Ly_c output they predict for the pAGB stellar component (Cid Fernandes et al. 2011, see also, e.g., Brown et al. 2008 and Woods & Gilfanov 2013 for a discussion related to this subject). These theoretical uncertainties presumably prevent a determination of the τ ratio to a precision better than within a factor of ~2 from currently available SSP models.

Our analysis in Sects. 2.1 and 2.2 uses two complementary data sets: i) single-spaxel (sisp) determinations from fits with an absolute deviation $|O_{\lambda} - M_{\lambda}|/O_{\lambda} \le 2.6$ (cf K12), where O_{λ} is the observed spectrum and M_{λ} the fit. These are typically restricted to the central, brightest part ($\mu \le 23 g \text{ mag/}\square''$) of our sample ETGs. ii) The average of all single-spaxel determinations within isophotal annuli (isan) adapted to the morphology of the (line-free) continuum between 6390 Å and 6490 Å (cf K12). These data, which are to be considered in a *statistical sense*, go $\gtrsim 2$ mag fainter, allowing study of the azimuthally averaged properties of the *wim* in the ETG periphery.




Fig. 1. From top to bottom: $\log([O III]/H\beta)$, $\log([N II]/H\alpha)$, $\log(EW(H\alpha))$, and $\log(\tau)$ vs normalized photometric radius R^*/r_p . The gray shaded areas in panels a&b mark the mean and $\pm 1\sigma$ of the respective quantity, and in panel c the mean EW(H α) for $R^* \ge r_p$ (0.43±0.65 Å). The light-blue area in panel c depicts the range in EW(H α) that can be accounted for by pAGB photoionization models (0.1–2.4 Å). The color assigned to each ETG is related to its $<\tau >$ (cf text and Fig. 3) in ascending order, from orange to violet, and is identical in all figures.

2.1. Radial behavior of emission-line diagnostics

Figures 1a&b show the diagnostic $\log([O III]/H\beta)$ and $\log([N II]/H\alpha)$ line ratios for our sample ETGs as a function of the photometric radius R^* , normalized to the SDSS r band Petrosian_50 radius r_p . The profiles are based on isan determinations, with green error bars illustrating the 1σ dispersion (typically ~0.4 dex) of single-spaxel data points within each annulus. All galaxies show LINER-specific Baldwin et al. (1981, BPT) ratios out to their periphery, with weak (if any) gradients solely within their central part ($R^* \leq r_p$). The mean ratios for our sample (shaded regions) were determined to be 0.37 ± 0.13 for $\log([O III]/H\beta)$ and 0.34 ± 0.26 for $\log([N II]/H\alpha)$, with a standard deviation about the mean σ_N of 0.02 and 0.05.

The EW(H α) profiles (panel c) reveal a more complex pattern. For $R^* \ge r_p$, most data points fall between 0.1 Å (EW⁺_{*}) and 2.4 Å (EW⁺_{*}), in the range of predictions from pAGB photoiniza-




Fig. 2. a-c) log(EW(H α)), log([O III]/H β) and log([N II]/H α) vs log(τ) for our sample ETGs. Open squares mark the central value for each isan profile (interconnected symbols), and dots show sisp determinations.




Fig. 3. Normalized H α intensity vs log(R^*) for our sample ETGs, based on isan determinations. The diagonal lines correspond to a power-law intensity drop-off of the form log(I/I_0) $\propto -\alpha \cdot log(R^*)$, with $\alpha = 1$. The right-hand side table lists the power-law slope α and the radially averaged EW(H α) and τ for each ETG.

tion models (e.g., Binette et al. 1994; Cid Fernandes et al. 2011, G13), whereas at smaller radii the sample seems to diverge into a lower ($\leq EW_{\star}^{-}$) and upper ($\geq EW_{\star}^{+}$) branch.

The τ ratio profiles (panel d) include correction for intrinsic extinction, with vertical bars illustrating the effect that neglecting it would have. The reference line at $\log(\tau)=0$ corresponds to an equilibrium state where the Ly_c photon output from pAGB stars balances the observed H α luminosity. Values below $(\log(\tau)<0)$ or above $(\log(\tau)>0)$ that line imply, in the first case, Ly_c photon injection by an additional source (e.g., star formation, AGN, shocks) and, in the second, Ly_c photon escape with a photon leakage fraction $plf = 1-\tau^{-1}$.

Setting a tentative division line at a radially averaged $\langle \tau \rangle = 2$, we can see that our ETG sample segregates into two groups. In the first one (type i; $\langle \tau \rangle < 2$, 14 ETGs), the τ ratio shows little dependence on radius, with individual data points deviating in most cases by no more than 0.3 dex from the equality line. This suggests a moderate Ly_c leakage ($plf \leq 0.5$) and/or dominant contribution of pAGB photoinization to the excitation of the *wim*. In the second group (type ii; $\langle \tau \rangle \geq 2$, 18 ETGs), the *plf* is typically very large (≥ 0.9) within r_p , and far from negligible (≥0.6) even in the galaxy periphery. As is apparent from panel c, these two groups differ in their EW(H α), with radially averaged values <EW> of 1.82±1.04 Å (σ_N =0.28 Å) and 0.41±0.25 Å (σ_N =0.06 Å). Another salient feature is that EW profiles of type i ETGs are nearly constant beyond ~ $r_p/2$, whereas those of type ii ETGs show a tendency toward a smooth, monotonic increase out to their periphery.

Figures 2a-c display projections of some quantities of interest onto τ . Unsurprisingly, both sisp and isan data delineate a trend toward decreasing EW(H α) with increasing τ , with type i and type ii ETGs populating, respectively, the lower and upper parts of a continuous sequence (panel a). This trend is also reflected on a relation log $\langle \tau \rangle = (0.23 \pm 0.04) - (1.36 \pm 0.09) \cdot \log \langle EW(H\alpha) \rangle$ for our sample (cf right-hand side list in Fig. 3 for the $\langle \tau \rangle$ and $\langle EW \rangle$ of individual ETGs). On the log([O III]/H β) vs log(τ) plane (panel b), the two ETG groups differ only marginally from one another (log([O III]/H β) of 0.29 \pm 0.11 and 0.43 \pm 0.12), while a weak trend toward increasing log([N II]/H α) with log(τ) is apparent from panel c (0.11 \pm 0.13 and 0.52 \pm 0.17 for type i and type ii ETGs, respectively).

2.2. Radial intensity distribution of nebular emission

The radial H α intensity profiles in Fig. 3 indicate that faint *ne* is present over nearly the entire optical extent of our sample ETGs. From the Abel integral equation (see, e.g., Papaderos et al. 1996, for a discussion and solutions for various intensity profiles) it follows that, for an isotropically emitting spheric-symmetric volume, an intrinsic luminosity density distribution l(r) scaling as $\propto r^{-2}$ would be projected onto a power-law intensity profile of the form $\log(I/I_0) \propto -\alpha \log(R^*)$ with $\alpha = 1$. On the simplifying assumption that the Ly_c output from a putative AGN is internally reprocessed into *ne* with a $l(r) \propto r^{-2}$, one can invoke the α inferred from H α profile fitting as a minimum consistency check for the AGN illumination hypothesis. The mean α for our sample (1.09), obtained for $R^* \ge 3''_{...}7$ (the effective FWHM resolution of CALIFA IFS cubes) is indeed consistent with it and close to the value deduced by Yan & Blanton (2012, α =1.28) from comparison of two-aperture spectroscopic data. Nevertheless, the large standard deviation in the derived slopes ($\sigma = 0.67$) argues against a *universal* power-law index $\alpha \approx 1$ for the *ne* intensity drop-off in ETGs. It is interesting though that comparison of Figs. 1&3 suggests a tendency for type ii ETGs having shallower H α profiles ($\alpha = 0.85 \pm 0.56$; $\sigma_N = 0.13$) than type i ETGs ($\alpha = 1.40 \pm 0.67$; $\sigma_N = 0.18$).

3. Discussion and conclusions

Summarizing the evidence from Sect. 2, the ETGs studied here form a broad, continuous sequence with respect to their H α , EW(H α), and τ profiles. Adopting a radially averaged τ ratio cutoff of $\langle \tau \rangle = 2$, we tentatively subdivide our sample into two groups: Typical properties of type i ETGs are a rather steep H α drop-off ($\alpha > 1$), nearly constant EWs of $\gtrsim 1$ Å beyond r_p , and a $\langle \tau \rangle$ close to unity (0.3...2). Type ii ETGs display shallower H α profiles ($\alpha < 1$), overall very low ($\leq EW_{\star}$...0.5 Å), outwardly increasing EWs, and a large (up to ~ 20) $<\tau$ >. Despite a difference of almost 2 dex in their nuclear τ ratios, these two groups differ little (by ≤ 0.4 dex) in their mean log([O III]/H β) and $\log([N II]/H\alpha)$ BPT ratios, which in either case are characteristic of LINERs and, within their uncertainties (~0.4 dex), are radially constant. In our ETG sample, 64% of the S0 galaxies fall into the type i group, and 78% of the E galaxies fall into the type ii group. Clearly, the classification proposed here is only indicative and needs to be refined, both by obtaining better statistics and through a quantitative comparison with other ETG properties: of these, the X-ray luminosity and temperature, the α_4 and $(v/\sigma)_{\star}$ parameter, and the star formation history are all being actively investigated.

As far as type i ETGs are concerned, various lines of evidence from this study suggest, in line with a substantial body of previous work (e.g. Sarzi et al. 2010; Annibali et al. 2010; Yan & Blanton 2012, K12, among others), that pAGB photoinization is the main driver of extended *ne*, with nonthermal sources only being potentially important in nuclei: a) First, ne is not confined only to the nuclear regions but is extended out to $R^{\star} \sim 2-4r_{p}$, i.e. is co-spatial with the pAGB stellar background. b) Second, radial $H\alpha$ profiles rule out a power-law intensity drop-off with a *universal* slope $\alpha \approx 1$, as a possible signature of a *dominant* AGN contribution to the excitation of the *wim*. c) Third, the EW(H α) is nearly constant beyond $\sim r_p$, pointing to a causal relationship between *ne* and the projected stellar surface density Σ_{\star} along the line of sight. This is a plausible expectation from the pAGB photoinization scenario, further supported by the quantitative agreement between predicted and observed EWs, as well as the narrow range in τ ratios ($\simeq 1$). These presumably suggest that type i ETGs contain a sufficient amount of wim being well mixed with stars, to justify case B recombination as a first-order approximation. Conversely, it is not immediately apparent how AGN or shock excitation alone could lead to the remarkable fine tuning between EW(H α) and Σ_{\star} over $\gtrsim 2$ dex in Σ_{\star} .

The emerging picture for type ii ETGs appears more complex. The τ ratios inferred for these systems imply at face value that the bulk (70%... \geq 90%) of the Ly_c output from hot pAGB stars (*consequently, from any other discrete or diffuse ionizing source*) escapes into the intergalactic space without being reprocessed into *ne*. Admittedly, plf estimates depend on the pAGB mass inferred from spectral synthesis models. These are known to suffer from degeneracies, the amplitude, topology, and systematics of which remain almost uncharted territory. One might argue that the large number of fits per galaxy (up to ~6800, using two SSP libraries) permits eliminating uncertainties, at least as far as isan determinations are concerned. However, this argument would only apply if errors in spectral synthesis were uncorrelated and nonsystematic.

Nevertheless, in the specific context of type ii ETGs, an essentially model-independent argument for extensive Ly_c leakage

comes from the virtual absence of *ne*, despite a sizeable ionizing photon budget. Quantitatively, for the Ly_c escape interpretation to become untenable, H α fluxes in type ii ETGs would need be revised upward by up to two orders of magnitude, which can be excluded by any reasonable error budget.

The τ profiles of these ETGs consistently point toward a low, radially dependent volume-filling factor f and/or density for the wim. A further hint in the same direction comes from their positive EW gradients: For a spheric-symmetric volume, these in connection with shallow (α <1) H α profiles are only reproducible when the wim luminosity density is monotonically decreasing toward the center. Alternatively, a feature predicted (though not observed) by Sarzi et al. (2010, see their Fig. 10) are positive EW gradients in a spherical stellar host reprocessing its pAGB Ly_c output within a *planar* gas configuration. Expanding the considerations by these authors, the high *plf*'s and outwardly increasing EWs of type ii ETGs might be reconcilable for a geometry that involves an oblate distribution of tenuous/clumpy gas within a spherical stellar host. Evidently, such a geometry would per se imply pAGB UV photon escape, further reinforcing our interpretation.

Regardless of the 3D distribution of the *wim*, its high porosity and/or low f call into question the importance of shock excitation in type ii ETGs. If the *wim* is primarily composed of compact cloudlets of radius r_c , then their large mean-free-path ((2/3) r_c/f , e.g., Jog & Solomon 1992) would act toward reducing the efficiency of energy dissipation via cloud-cloud collisions and shocks.

Given our findings, it is important to ask whether the "weak" AGN interpretation of the optical emission lines is compelling anymore. In the presence of extensive Ly_c leakage, emission-line intensities and EWs in type ii ETG nuclei are lowered by factor between ~10 and ≤ 100 . Consequently, the presence of weak *ne* in these systems is not in itself proof of a "weak" (sub-Eddington accreting) AGN. In fact, the importance of Ly_c photon escape, which heretofore has not been investigated in detail, may be a key insight into resolving one of the longstanding enigmas in AGN/LINER research. It offers an *ansatz* for reconciling the fact that many ETGs with prominent signatures of strong AGN activity in radio continuum and/or X-ray wavelengths merely show weak (LINER) optical AGN signatures.

In addition, the relative distribution of gas compared to the stars is an important issue. While in a thin, face-on disk, a nuclear EW of, say, ≤ 10 Å can safely be regarded as evidence of faint *ne* (and a weak AGN), this is not necessarily the case for a *triaxial* ETG, where the *ne*-emitting gas volume may have a more limited extent than the stellar component. In the latter case, nuclear EWs are effectively lowered by the high-surface brightness screen of background and foreground stars along the optical path. In conjunction with the Ly_c photon escape, this line-of-sight dilution of the nuclear EWs will conspire to create an observational selection effect, favoring optical detection of AGN activity in oblate, face-on ETGs with atypically low *plf*'s.

Arguably, one of the most surprising results of this study are the similar mean BPT ratios of type i and type ii ETGs, despite having substantial differences in their *wim* characteristics. Perhaps the luminosity-weighted emission-line ratios projected along the line of sight "saturate" into the LINER regime for a broad range of *wim* distributions and characteristics (i.e. combinations of differing *f*, covering fractions, densities, ionizing photon mean free-path, and ionization parameters), becoming degenerate for ETGs. Circumstantial support for this hypothesis comes from radiation transfer models by Ercolano et al. (2012), who report that a subset of the *projected* emission line ratio determinations in a 3D model of the *Pillars of Creation* can mimic LINER characteristics in classical (1D) BPT diagrams. Clearly, detailed 3D radiative transfer modeling of the *wim* in ETGs, including nonequilibrium ionization effects (e.g., de Avillez & Breitschwerdt 2012), are important for understanding the nature of *wim* in ETGs. High-quality IFS data, such as those from CAL-IFA, will no doubt provide crucial observational constraints on these next-generation models.

References

- Annibali, F., Bressan, A., Rampazzo, R., et al. 2010, A&A, 519, A40
- Baldwin, J. A., Phillips, M. M., & Terlevich, R. 1981, PASP, 93, 5
- Binette, L., Magris, C. G., Stasińska, G., & Bruzual, A. G. 1994, A&A, 292, 13 Brown, T.M., Smith, E., Ferguson, H.C. et al. 2008, ApJ, 682, 319
- Bruzual, G. & Charlot, S. 2003, MNRAS, 344, 1000
- Cid Fernandes, R., Mateus, A., Sodré, L. et al. 2005, MNRAS, 358, 363
- Cid Fernandes, R., Stasińska, G., Mateus, A. et al. 2011, MNRAS, 413, 1687
- de Avillez, M.A. & Breitschwerdt, D. 2012, ApJ, 756, L3
- Ercolano, B., Dale, J.E., Gritschneder, M. et al. 2012, MNRAS, 420, 141
- Dopita, M. A. & Sutherland, R. S. 1995, ApJ, 455, 468
- Ho, L.C. 1999, Adv. Space Res., 23, 813
- Husemann, B., Jahnke, K., Sánchez, S. F. et al. 2013, A&A, 549, A87H
- Jog, C.J. & Solomon, P.M. 1992, ApJ, 387, 152
- Kehrig, C., Monreal-Ibero, A., Papaderos, P., et al. 2012, A&A, 540, A11 (K12) Papaderos, P., Loose, H.-H., Thuan, T.X., Fricke, K.J. 1996, A&AS, 120, 207
- Phillips, M.M., Jenkins, C.R., Dopita, M.A. et al. 1986, AJ, 91, 1062
- Sánchez, S. F., Kennicutt, R. C., Gil de Paz, A., et al. 2012, A&A, 538, A8
- Sánchez-Blázquez, P. et al. 2006, MNRAS, 371, 703
- Sarzi, M., Falcón-Barroso, J., Davies, R. L., et al. 2006, MNRAS, 366, 1151 Sarzi, M., Shields, J. C., Schawinski, K., et al. 2010, MNRAS, 402, 2187
- Stazi, M., Sineus, J. C., Schawniski, K., et al. 2010, MNRAS, 402, 2187 Stasińska, G., Vale Asari, N., Cid Fernandes, R. et al. 2008, MNRAS, 391, L29
- Trinchieri, G. & di Serego Alighieri, S. 1991, AJ, 101, 1647
- Vazdekis, A. et al. 2010, MNRAS, 404, 1639
- Woods, T.E. & Gilfanov, M. 2013, MNRAS, 432, 1640
- Yan, R. & Blanton, M.R. 2012, ApJ, 747, 61
- ¹ Centro de Astrofísica and Faculdade de Ciências, Universidade do Porto, Rua das Estrelas, 4150-762 Porto, Portugal
- ² Instituto de Astrofísica de Andalucía (CSIC), Glorieta de la Astronomía s/n Aptdo. 3004, E18080-Granada, Spain
- ³ GEPI, Observatoire de Paris, UMR 8111, CNRS, Université Paris Diderot, 5 place Jules Janssen, 92190 Meudon, France
- ⁴ University of Vienna, Türkenschanzstrasse 17, 1180 Vienna, Austria
 ⁵ Centro Astronómico Hispano Alemán de Calar Alto (CSIC-MPG),
- C/ Jesús Durbán Remón 2-2, E-4004 Almería, Spain
 ⁶ Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte
- 16, D-14482 Potsdam, Germany
 ⁷ Sydney Institute for Astronomy, University of Sydney, NSW 2006,
- Australia
- ⁸ School of Physics and Astronomy, University of St Andrews, North Haugh, St Andrews, KY16 9SS, UK (SUPA)
- ⁹ Instituto de Astrofísica de Canarias, Vía Láctea s/n, La Laguna, Tenerife, Spain
- ¹⁰ Departamento de Astrofísica, Universidad de La Laguna, E-38205 La Laguna, Tenerife, Spain
- ¹¹ CENTRA Centro Multidisciplinar de Astrofísica, Instituto Superior Técnico, Av. Rovisco Pais 1, 1049-001 Lisbon, Portugal
- ¹² Australian Astronomical Observatory, PO Box 915, North Ryde, NSW 1670, Australia
- ¹³ Department of Physics and Astronomy, Macquarie University, NSW 2109, Australia
- ¹⁴ Departamento de Investigación Básica, CIEMAT, Avda. Complutense 40, E-28040 Madrid, Spain
- ¹⁵ Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

Acknowledgements. We thank the anonymous referee for valuable suggestions. This paper is based on data from the Calar Alto Legacy Integral Field Area Survey, CALIFA (http://califa.caha.es), funded by the Spanish Ministery of Science under grant ICTS-2009-10, and the Centro Astronómico Hispano-Alemán. PP is supported by Ciencia 2008 Contract, funded by FCT/MCTES (Portugal) and

POPH/FSE (EC), and JMG by a Post-Doctoral grant, funded by FCT/MCTES (Portugal) and POPH/FSE (EC). PP&JMG acknowledge support by the Fundação para a Ciência e a Tecnologia (FCT) under project FCOMP-01-0124-FEDER-029170 (Reference FCT PTDC/FIS-AST/3214/2012), funded by FCT-MEC (PIDDAC) and FEDER (COMPETE). IM acknowledges support from Spanish grant AYA2010-15169 and the Junta de Andalucia through TIC-114 and the Excellence Project P08-TIC-03531. J.F-B. from the Ramón y Cajal Program, grants AYA2010-21322-C03-02 and AIB-2010-DE-00227 from the Spanish Ministry of Economy and Competitiveness (MINECO), as well as from the FP7 Marie Curie Actions of the European Commission, via the Initial Training Network DAGAL under REA grant agreement no 289313. The STARLIGHT project is supported by the Brazilian agencies CNPq, CAPES, and FAPESP. PP has enjoyed inspiring discussions on the effects of Ly_c photon leakage with Prof. Nils Bergvall (Uppsala University). We benefited from stimulating discussions with several members of the CALIFA collaboration. This research made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.