# A New Multi-wavelength Solar Telescope: Optical and Near-infrared Solar Eruption Tracer (ONSET)

C. Fang<sup>1,2</sup>, P. F. Chen<sup>1,2</sup>, Z. Li<sup>1,2</sup>, M. D. Ding<sup>1,2</sup>, Y. Dai<sup>1,2</sup>, X. Y. Zhang<sup>3</sup>, W. J. Mao<sup>4</sup>, J. P. Zhang<sup>4</sup>, T. Li<sup>4</sup>, Y. J. Liang<sup>4</sup> and H. T. Lu<sup>4</sup>

- <sup>1</sup> School of Astronomy & Space Science, Nanjing University, Nanjing 210093, China; *fangc@nju.edu.cn*
- <sup>2</sup> Key Laboratory of Modern Astronomy & Astrophysics, Nanjing University, Nanjing 210093, China
- <sup>3</sup> Yunnan Astronomical Observatory, National Astronomical Observatories, Chinese Academy of Sciences, Kunming 650011, China
- <sup>4</sup> Nanjing Institute of Astronomical Optics and Technology, CAS, China

Received [2013] [June] [10]; accepted [2013] [month] [day]

**Abstract** A new multi-wavelength solar telescope, Optical and Near-infrared Solar Eruption Tracer (ONSET) of Nanjing University, was constructed, being fabricated by Nanjing Institute of Astronomical Optics & Technology and run in cooperation with Yunnan Astronomical Observatory. ONSET is able to observe the Sun in three wavelength windows: He I 10830 Å, H $\alpha$ , and white-light at 3600 Å and 4250 Å, which are selected in order to obtain the dynamics in the corona, chromosphere, and the photosphere simultaneously. Full-disk or partial-disk solar images with a field of 10 arcmin at three wavelengths can be obtained nearly simultaneously. It is designed to trace solar eruptions with high spatial and temporal resolutions. This telescope was installed at a new solar observing site near Fuxian Lake in Yunnan Province, southwest China. The site is located at E102N24, with an altitude of 1722 m. The seeing is stable and of high quality. We give a brief description of the scientific objectives and the basic structure of the ONSET. Some preliminary results are also presented.

Key words: techniques: photometric — Sun: observation — Sun: telescope

## **1 INTRODUCTION**

Recently, owing to the development of solar physics and space weather forecast, multi-wavelength solar observations become a pressing issue (Fang, 2011). Fast evolution and fine structures of solar activities, such as flares, coronal mass ejections (CMEs), and filaments, make the high spatio-temporal resolution observations an essential target for the observations (see, e.g., Fang et al., 2008). Correspondingly, a few new solar telescopes were contructed in the past years. For example, in Europe, a telescope called Chromospheric Telescope (ChroTel) started to observe the Sun in Ca II K, H $\alpha$ , and He I 10830 Å since April 2012 (Bethge et al., 2011); in USA, the Synoptic Optical Long-term Investigation of the Sun (SOLIS) was equipped with a Full-Disk Patrol (FDP) module, which started to observe full disk images of the Sun at high cadence in Ca II K, H $\alpha$ , and He I 10830 Å since June 2011 (Pevtsov et al., 2011). Therefore, there is a gap of time zone with similar observations.

In order to fill in the gap so as to form a mini-network in monitoring the solar chromosphere and the signatures of the corona continuously, as well as to carry out researches on solar activities and Fang et al.

space weather during the solar cycle 24 in China, we decided to construct a new multi-wavelength solar telescope called Optical & Near-infrared Solar Eruption Tracer (ONSET), which can conduct multi-wavelength observations in He I 10830 Å, H $\alpha$ , and white-light at 3600 Å and 4250 Å. The operation of the telescope is done jointly by Nanjing University and Yunnan Astronomical Observatory. ONSET was fabricated by Nanjing Institute of Astronomical Optics & Technology. The project was initiated in 2005, and the telescope was installed in 2011 at a new solar observing site at the bank of Fuxian Lake, 60 km from Kunming, Yunnan Province in southwest China. The site is located at E102°57′11″N24°34′47″, with an altitude of 1722 m. The lake is about 8 km wide and 30 km long, with the average depth being 89.6 m. The seeing is stable, with the Fried parameter  $r_0$  slightly larger than 10 cm on average (Liu & Beckers, 2001).

In this paper, we describe the scientific objectives of ONSET in Section 2, and the basic structure and the specifications of the telescope are presented in Section 3. The preliminary results are given in Section 4. A summary is given in Section 5.

## **2** SCIENTIFIC OBJECTIVES

The main objectives of ONSET are to study the following important topics:

## (1) Dynamics and fast fluctuations in flares

Radiative transfer calculations indicate that different parts of the H $\alpha$  line may respond in different timescales to the rapid energy input (Canfield & Gayley, 1987). In observations, Wang et al. (2000) found that the H $\alpha$ -1.3 Å emission in a flare shows high-frequency fluctuations on a timescale of a few tenths of a second, which are temporally correlated with the hard X-ray emission variations. Similar observations were done by Radziszewski et al. (2011) for both the line center and line wings. Numerical simulations by Ding et al. (2001) and Ding (2005) verified the close relationship between the fast variations in H $\alpha$  intensity and hard X-ray emission (the latter of which characterizes the electron beam heating rate). Note that in some cases, the H $\alpha$  line center is more easily affected by thermal conduction and some secondary effects like chromospheric evaporation and Doppler shifts. Therefore, the line wings, in particular the blue wing (Wang et al., 2000), is more suitable to show the non-thermal impulsive heating.

He I 10830 Å line is also a potential diagnostic tool for thermal and nonthermal effects in solar or stellar flares (Ding et al., 2005). The He I 10830 Å line is a multiplet comprising of three components. This line, although formed in upper chromospheric layers, is very sensitive to coronal EUV irradiation. Generally speaking, there are two main ways to populate the atomic levels responsible for this line: photoionization by EUV irradiation followed by recombination and direct collisional excitation to the triplet levels (e.g., Centeno et al., 2008). During solar flares, however, a third process, collisional ionization by the electron beam followed by recombination, may be at work (Ding et al., 2005). Therefore, the line formation is complicated and one should be very cautious in interpreting the line features.

Our new telescope will have a partial-disk observation mode, with a high spatial resolution ( $\sim 1''$  or better) and high temporal resolution (better than 1 s), in order to study the fine structures in solar flares. This will help reveal the physical processes regarding magnetic reconnection, energy release, and energy transport in the solar atmosphere. In particular, combining these data with other observations in hard X-ray, EUV, and radio, etc., we can diagnose the thermal and non-thermal processes in flares (e.g., Fang et al., 1993; Henoux et al., 1993, 1995) and other activities.

## (2) Patrol of white-light flares

White-light flares (WLFs) are generally thought to correspond to a type of most energetic solar flares, with the continuum emissions from the photosphere or lower chromosphere. They present a major challenge to the flare atmospheric models and energy transport mechanisms (Neidig, 1989; Ding et al., 1999). It was suggested that all solar flares might be WLFs (Neidig, 1989; Hudson et al., 2006), which seems to be supported by the detection of white-light emission in weak flares (Matthews et al., 2003; Jess et al., 2008; Wang, 2009) and by the spectral irradiance observations (Kretzschmar, 2011). However, since the discovery of the first WLF in 1859 (Carrington, 1859), only around 150 WLF events have been reported conclusively so far. There are several reasons for it, e.g., (a) The white-light enhance-

ment in the visible continuum in a WLF is only a few percent or less, which might below the sensitivity of some telescopes; (b) The lifetime of the white-light emission is only  $\sim 1-2$  minutes, which is too short for a certain identification. On the other side, we have to be cautious when we use 1600 Å band emissions recorded by some spacecraft like TRACE and SDO/AIA to search for WLFs since this band contains emission lines, e.g., C IV. It is possible that some of the reported WLFs in 1600 Å are not real WLFs.

According to whether there is a Balmer jump or not, WLFs can be classified into type I and type II, respectively (Fang & Ding, 1995). While type I WLFs can be explained by the standard flare model, type II WLFs require in situ heating in the photosphere, say, low-atmosphere magnetic reconnection (e.g., Li et al., 1997b; Chen et al., 2001). Only with two white-light wavebands inside and outside the Balmer continuum can we distinguish which type a WLF is. For that purpose, we selected two white-light bands, i.e., 3600 Å or 4250 Å, for the ONSET. With the new telescope, we expect to observe more WLFs in the solar cycle 24, and to clarify some key questions regarding WLFs, such as how common WLFs are and what heating mechanism is responsible for each of the two types of WLFs.

#### (3) CME onset and filament activation

As the largest solar eruptions which may pose hazardous impact on the terrestrial environment or space weather, CMEs continue to attract more and more attentions (Chen, 2011). The most important question regarding CMEs is how the CME progenitor is triggered to erupt. Since a large fraction of CMEs originate from filaments (or prominences when they appear above the limb), it would be of great significance to monitor the activation of filaments. Various observations have revealed that before a CME is formed, the associated filament was already rising with a speed of ~10 km s<sup>-1</sup> (e.g., Cheng et al., 2010, and references therein), as demonstrated by MHD numerical simulations (e.g., Chen & Shibata, 2000). The detection of such early slow rise will be important for a better understanding of CME triggering process and for the space weather forecast.

For the limb events, we can detect the rise motion from the imaging observations directly. However, for the disk events, especially those near the solar disk center, the detection of the slow rise motion relies on spectrometers. One alternative way is to use simultaneous H $\alpha$  imaging observations at line center and line wings (both blue and red wings). With that, the true rising velocity of a filament on the solar disk can be measured with a high precision (e.g., Morimoto & Kurokawa, 2003).

#### (4) Moreton waves

Moreton waves are fast propagating fronts observed in the chromosphere, with an averaged velocity of ~660 km s<sup>-1</sup> (Zhang et al., 2011). They typically appear dark in the H $\alpha$  red wing and bright in the H $\alpha$  line center and blue wing. These features can be well explained by the Uchida's model in terms of a fast-mode mode wave in the corona sweeping the chromosphere (Uchida, 1968). There are still many puzzles awaiting further explanations. While Moreton waves were widely believed to be generated by the pressure pulse in solar flares, there appeared an alternative view, i.e., they are generated by the piston-driven shock as a flux rope erupts (e.g., Cliver et al., 1999; Chen et al., 2002). Another issue is the rarity of Moreton wave events. So far, it seems that less than 40 Moreton wave events have been detected since 1960. One reason is that most H $\alpha$  telescopes used the line center for observations, whereas Moreton waves are better observed near H $\alpha \pm 0.45$  Å (Chen et al., 2005a).

#### (5) He I 10830 waves and CME nowcast

One of the most important discoveries of the SOHO space mission is "EIT waves" (Thompson et al., 1998). The intriguing phenomenon sparked world-wide debates. A number of papers tried to explain EIT waves in terms of fast-mode waves (Li et al., 2012; Zheng et al., 2012), there are definitely plenty of observational features which cannot be accounted for by the fast-mode wave model, which stimulated the birth of other non-wave models (see Chen & Fang, 2012, for a review). For example, Chen et al. (2002) and Chen et al. (2005b) proposed the magnetic field-line stretching model, claiming that EIT waves are produced by the successive stretching of the magnetic field lines straddling over the erupting flux rope. Such a model can explain a variety of observations (Yang & Chen, 2010; Chen, 2011; Cheng et al., 2012; Dai et al., 2012; Shen & Liu, 2012).

EIT waves were found to be the EUV counterpart of CMEs (Chen, 2009). Therefore, their appearance can be used to nowcast CMEs, especially those directed toward the Earth. However, EIT waves can Fang et al.

be observed only in EUV wavelength, which is feasible in space only. Fortunately, the coronal brightening would result in the darkness of the near infrared line, He I 10830 Å (Centeno et al., 2008). Therefore, even on the ground, we may detect the near infrared counterparts of EIT waves (see Gilbert et al., 2004, for an example).

On the other hand, with the same formation mechanism of H $\alpha$  Moreton waves, the fast-mode coronal shock wave, when sweeping the chromosphere, might also be manifested in He I 10830 Å. Therefore, it is possible to detect two types of He I waves, the same as seen in EUV (Chen & Wu, 2011).

## (6) Filament oscillations

Perturbations are ubiquitous in the solar atmosphere, including the everlasting convective motions in the photosphere, frequent brightenings in the chromosphere, and sporadic flares in the corona. Subjected to these perturbations, filaments would oscillate accordingly. On one hand, filament oscillations provide an independent approach to diagnose the magnetic field across the filaments (Jing et al., 2003, 2006; Isobe & Tripathi, 2006; Zhang et al., 2013); On the other hand, long-lasting filament oscillations might be one of the precursors for CME eruptions (Chen et al., 2008).

## (7) Other fine structures

In the solar low atmosphere, there are a pool of small-scale eruptive events, including microflares (Fang et al., 2006a), Ellerman bombs (Chen et al., 2001; Fang et al., 2006b), and coronal bright points (CBPs, Zhang et al., 2012b). While Ellerman bombs can be directly observed at the wings of H $\alpha$  line, microflares and coronal bright points are often observed in X-ray, but sometimes they show brightenings in H $\alpha$  (Jiang et al., 2012; Zhang et al., 2012a). Besides, CBPs appear as dark points in He I 10830 Å (Li et al., 1997a).

It is generally believed that coronal holes are the source of the fast solar wind, which is important for space weather researches. Using He I 10830 Å images, we can clearly detect the coronal holes and trace their evolution.

# **3 BASIC STRUCTURE AND SPECIFICATIONS**



**Fig. 1** The ONSET building near Fuxian Lake (*left*) and the telescope inside the blossom dome (*right*). The left picture was taken during the construction of the building.

Figure 1 shows the ONSET building near Fuxian Lake (left panel) and the telescope inside the blossom dome (right panel). The ONSET telescope consists of four tubes: (1) a near-infrared vacuum tube with an aperture of 27.5 cm, observing at He I  $10830\pm4.0$  Å with a FWHM of 0.5 Å; (2) a chromospheric (H $\alpha$ ) vacuum tube with an aperture of 27.5 cm, observing at 6562.8  $\pm$  2.5 Å with a FWHM of 0.25 Å; (3) a white-light vacuum tube with an aperture of 20 cm, observing at the wavelength 3600 Å or 4250 Å with a FWHM of 15 Å; and (4) a guiding tube with an aperture of 14 cm, observing at the wavelength 5500 Å.

The former two tubes are equipped with two Lyot-type filters, while the latter two use interference filters. The first three tubes are vacuum ones, and they provide full-disk (extending up to  $1.3R_{\odot}$ ) or partial-disk images with a field of view  $10' \times 10'$ . The three wavelength windows can work independently and simultaneously. Two observation modes (full-disk and partial-disk) can be switched back and forth according to scientific requirements. All the observations are controlled by a computer and the observing program is flexible. It is planned to perform routine observations to acquire 8–10 full-disk images every 1 min in the H $\alpha$  line center and its two wings at ±0.5 Å, He I 10830 Å line center and its two wings, and two white-light wavelengths. We can also carry out partial-disk observations of solar activities with a high spatial (~1" or better) and temporal (0.1 to 1 s) resolutions .

The ONSET has one PIXIS 2048BR CCD for H $\alpha$ , with 2048×2048 pixels (the pixel size is 13.5  $\mu$ m), one PCO4000 CCD for white-light observations, with 4008×2672 pixels (the pixel size is 9  $\mu$ m), and one VersArray CCD for 10830 Å observations, with 1340×1300 pixels (the pixel size is 20  $\mu$ m). The diameters of all solar images are about 24 mm at the focal plane.



**Fig. 2** The inner assembly of the mechanical structure of the ONSET tubes. The telescope consists of four tubes, including one guiding tube and three main tubes for H $\alpha$ , He I 10830 Å, and white-light observations, respectively.

Figure 2 displays the mechanical structure of the ONSET tubes, and Figure 3 shows the optical layout of the telescope.

A series of software programs are being developed (e.g., Hao et al., 2013), including data archiving, search engine, data calibrations, automatic detections of solar activities (e.g., Moreton waves and He I 10830 waves), velocity measurements, and the remote-control system, etc.

### **4 PRELIMINARY RESULTS**

Since 2011 April, we have made test observations of the new telescope, and got some preliminary results. Figure 4 shows an H $\alpha$  partial-disk image and a white-light partial-disk image at 4250 Å, where the fine





**Fig. 3** The optical layout of the telescope (*left*) and the enlarged view of the block diagram in the left panel (*right*).



**Fig. 4** An H $\alpha$  partial-disk image (*left*) and a 4250 Å partial-disk image (*right*) observed by ONSET. The image in the left panel was taken at 07:05:25 UT on 21 April 2011 with a field of view of  $10' \times 10'$ , whereas that in the right panel was on 14:00:52 UT on 27 May 2013 with a field of view of  $210'' \times 160''$ .

structures of a sunspot and the background granules are clearly visible. It seems that the image quality is very good. In particular, a WLF was successfully observed on 2012 March 9, which is the first flare detected by the ONSET (Hao et al., 2012). Continuum emissions appear clearly at 3600 Å and 4250 Å wavebands. The peak enhancements at these two bands are 25% and 12%, respectively. This event shows clearly the evidence that the white light emission is caused by energetic particles bombarding the solar lower atmosphere.

Since the beginning of 2013, ONSET has started to make routine observations. Figure 5 illustrates an H $\alpha$  image (left panel) and a He I 10830 Å image (right panel), which is the first 10830 Å image observed ever in China.

ONSET is dedicated to a wide variety of solar activity researches, especially through joint observations with other ground-based and space-born instruments. The data from routine observations will be open and available on the internet (http://sdac.nju.edu.cn). Everyone is welcome to use the ONSET data and to help develop software programs for the data reduction.

## **5 SUMMARY**

After making efforts in the past several years, we have successfully constructed the Optical and Nearinfrared Solar Eruption Tracer (ONSET) at the new solar observing site near Fuxian Lake, which is



**Fig. 5** An H $\alpha$  (*left*) and a He I 10830 Å (*right*) image observed by ONSET. The image in the left panel was taken at 03:09:36 UT on 30 April 2013, whereas that in the right panel was at 03:39:21 UT on 13 March 2013.

currently the best site for solar observations in China. The ONSET consists of four tubes, and three of them are vacuum ones. The telescope can provide 8–10 solar images at He I 10830 Å (line center and blue/red wings), H $\alpha$  (line center and blue/red wings), and white-light at 3600 Å and 4250 Å. The wavebands were selected in order to observe the dynamics in the corona, chromosphere, and the photosphere simultaneously. With the accumulation of the observational data, it is expected to acquire the following results:

(1) Fast variations of solar flare emissions will be examined in a systematic way in order to indirectly derive nonthermal parameters in solar flares. In particular, the different response in H $\alpha$  and He I 10830 Å to nonthermal particles can be compared;

(2) A catalog of white-light flares will be compiled, with the classification of type I and type II events. The relationship between white-light enhancements and soft X-ray intensity can be scrutinized;

(3) Early slow rise motion of erupting filaments can be detected with a high cadence, which provides real-time monitoring of the onset of filament eruptions and CMEs;

(4) Both H $\alpha$  Moreton waves and "EIT waves" might be imprinted in He I 10830 Å images, which will help clarify the possible different nature of the two wavelike phenomena;

(5) Both transverse and longitudinal filament oscillations will be recorded routinely, which paves the way to prominence seismology;

(6) A large number of small-scale eruptions will be observed, including Ellerman bombs and CBPs.

The preliminary results demonstrate that the quality of the solar images is high and good enough for scientific researches. Since 2013, ONSET starts to perform routine observations. The data will be open and available on the internet. Everyone is welcome to use the ONSET data and to help develop software programs.

Acknowledgements This work was supported by the National Natural Science Foundation of China (NSFC) under Nos. 10878002, 10610099, 10933003, 10673004, and 11025314, as well as the grant from the 973 project 2011CB811402 of China.

#### References

Bethge, C., Peter, H., Kentischer, T. J., et al. 2011, A&A, 534, A105 Canfield, R. C., & Gayley, K. G. 1987, ApJ, 322, 999 Carrington, R. C. 1859, MNRAS, 20, 13 Fang et al.

- Centeno, R., Trujillo Bueno, J., Uitenbroek, H., & Collados, M. 2008, ApJ, 677, 742
- Chen, F., Ding, M. D., Chen, P. F., & Harra, L. K. 2011, ApJ, 740, 116
- Chen, P. F. 2009, ApJ, 698, L112
- Chen, P. F. 2011, Living Reviews in Solar Phys., 8, 1
- Chen, P. F., Ding, M. D., & Fang, C. 2005, Space Sci. Rev., 121, 201
- Chen, P. F., & Fang, C. 2012, EAS Publications Series, 55, 313
- Chen, P.-F., Fang, C., & Ding, M.-D. D. 2001, ChJAA (Chin. J. Astron. Astrophys.), 1, 176
- Chen, P. F., Fang, C., & Shibata, K. 2005, ApJ, 622, 1202
- Chen, P. F., Innes, D. E., & Solanki, S. K. 2008, A&A, 484, 487
- Chen, P. F., & Shibata, K. 2000, ApJ, 545, 524
- Chen, P. F., Wu, S. T., Shibata, K., & Fang, C. 2002, ApJ, 572, L99
- Chen, P. F., & Wu, Y. 2011, ApJ, 732, L20
- Cheng, X., Ding, M. D., & Zhang, J. 2010, ApJ, 712, 1302
- Cheng, X., Zhang, J., Olmedo, O., et al. 2012, ApJ, 745, L5
- Cliver, E. W., Webb, D. F., & Howard, R. A. 1999, Sol. Phys., 187, 89
- Dai, Y., Ding, M. D., Chen, P. F., & Zhang, J. 2012, ApJ, 759, 55
- Ding, M. D. 2005, Adv. Space Res., 35, 1846
- Ding, M. D., Fang, C., & Yun, H. S. 1999, ApJ, 512, 454
- Ding, M. D., Li, H., & Fang, C. 2005, A&A, 432, 699
- Ding, M. D., Qiu, J., Wang, H., & Goode, P. R. 2001, ApJ, 552, 340
- Fang, C., Hénoux, J.-C., & Gan, W. Q. 1993, A&A, 274, 917
- Fang, C., & Ding, M. D. 1995, A&AS, 110, 99
- Fang, C., Tang, Y. H., & Xu, Z. 2006, ChJAA, 6, 597
- Fang, C., Tang, Y. H., Xu, Z. et al. 2006, ApJ, 643, 1325
- Fang, C., Ding, M. D., & Chen, P. F. 2008, Physics of Solar Active Regions (in Chinese), Nanjing: Nanjing Univ. Press
- Fang, C. 2011, RAA, 11, 1377
- Gilbert, H. R., Holzer, T. E., Thompson, B. J., & Burkepile, J. T. 2004, ApJ, 607, 540
- Hao, Q., Fang, C., & Chen, P. F. 2013, Sol. Phys., 286, 385
- Hao, Q., Guo, Y., Dai, Y., Ding, M. D., Zhang, X. Y., and Fang, C. 2012, A&A, 544, L17
- Hénoux, J. -C., Fang, C., & Gan, W. Q. 1993, A&A, 274, 923
- Hénoux, J. -C., Fang, C., & Gan, W. Q. 1995, A&A, 297, 574
- Hudson, H. S., Wolfson, C. J., & Metcalf, T. R. 2006, Sol. Phys., 234, 79
- Isobe, H., & Tripathi, D. 2006, A&A, 449, L17
- Jess, D. B., Mathioudakis, M., Crockett, P. J. et al. 2008, ApJ, 688, L119
- Jiang, R.-L., Fang, C., & Chen, P.-F. 2012, ApJ, 751, 152
- Jing, J., Lee, J., Spirock, T. J., et al. 2003, ApJ, 584, L103
- Jing, J., Lee, J., Spirock, T. J., & Wang, H. 2006, Sol. Phys., 236, 97
- Kretzschmar, M. 2011, A&A, 530, A84
- Li, H., Fan, Z.-Y., & You, J.-Q. 1997, Chin. Astron. Astrophys., 21, 128
- Li, T., Zhang, J., Yang, S.-H., & Liu, W. 2012, RAA, 12, 104
- Li, X. Q., Song, M. T., Hu, F. M., & Fang, C. 1997, A&A, 320, 300
- Liu, Z., & Beckers, J. M. 2001, Solar Phys., 198, 197
- Matthews, S. A., van Driel-Gesztelyi, L., Hudson, H. S., & Nitta, N. V. 2003, A&A, 409, 1107
- Morimoto, T., & Kurokawa, H. 2003, PASJ, 55, 505
- Neidig, D. F. 1989, Sol. Phys., 121, 261
- Pevtsov, A. A., Streander, K., Harvey, J., et al. 2011, Bulletin of the American Astronomical Society, 1747
- Radziszewski, K., Rudawy, P., & Phillips, K. J. H. 2011, A&A, 535, A123
- Shen, Y., & Liu, Y. 2012, ApJ, 752, L23
- Thompson, B. J., Plunkett, S. P., Gurman, J. B., et al. 1998, Geophys. Res. Lett., 25, 2465
- Uchida, Y. 1968, Sol. Phys., 4, 30
- Wang, H.-M. 2009, RAA, 9, 127

Wang, H., Qiu, J., Denker, C., et al. 2000, ApJ, 542, 1080

Yang, H. Q., & Chen, P. F. 2010, Sol. Phys., 266, 59

Zhang, P., Fang, C., & Zhang, Q. M. 2012, Science in China G: Physics and Astronomy, 55, 907

Zhang, Q. M., Chen, P. F., Guo, Y., Fang, C., & Ding, M. D. 2012, ApJ, 746, 19

Zhang, Q. M., Chen, P. F., Xia, C., Keppens, R., & Ji, H. S. 2013, A&A, 554, 124

Zhang, Y., Kitai, R., Narukage, N., et al. 2011, PASJ, 63, 685

Zheng, R., Jiang, Y., Yang, J., et al. 2012, ApJ, 747, 67