

**TWISTED CYCLES AND TWISTED PERIOD RELATIONS
FOR LAURICELLA'S HYPERGEOMETRIC FUNCTION F_C**

YOSHIAKI GOTO

ABSTRACT. We study Lauricella's hypergeometric function F_C by using twisted (co)homology groups. We construct twisted cycles with respect to an integral representation of Euler type of F_C . These cycles correspond to 2^m linearly independent solutions to the system of differential equations annihilating F_C . Using intersection forms of twisted (co)homology groups, we obtain twisted period relations which give quadratic relations for Lauricella's F_C .

1. INTRODUCTION

Lauricella's hypergeometric series F_C of m -variables x_1, \dots, x_m with complex parameters a, b, c_1, \dots, c_m is defined by

$$F_C(a, b, c; x) = \sum_{n_1, \dots, n_m=0}^{\infty} \frac{(a, n_1 + \dots + n_m)(b, n_1 + \dots + n_m)}{(c_1, n_1) \cdots (c_m, n_m) n_1! \cdots n_m!} x_1^{n_1} \cdots x_m^{n_m},$$

where $x = (x_1, \dots, x_m)$, $c = (c_1, \dots, c_m)$, $c_1, \dots, c_m \notin \{0, -1, -2, \dots\}$ and $(c_1, n_1) = \Gamma(c_1 + n_1)/\Gamma(c_1)$. This series converges in the domain

$$D_C := \left\{ (x_1, \dots, x_m) \in \mathbb{C}^m \left| \sum_{k=1}^m \sqrt{|x_k|} < 1 \right. \right\},$$

and admits the integral representation (3). The system $E_C(a, b, c)$ of differential equations annihilating $F_C(a, b, c; x)$ is a holonomic system of rank 2^m with the singular locus S given in (1). There is a fundamental system of solutions to $E_C(a, b, c)$ in a simply connected domain in $D_C - S$, which is given in terms of Lauricella's hypergeometric series F_C with different parameters, see (2) for their expressions.

In the case of $m = 2$, the series $F_C(a, b, c; x)$ and the system $E_C(a, b, c)$ are called Appell's hypergeometric series $F_4(a, b, c; x)$ and system $E_4(a, b, c)$ of differential equations, which are studied in [6] by twisted (co)homology groups concerning with the integral representation (3) for $m = 2$. We construct twisted cycles corresponding to four solutions to $E_4(a, b, c)$ expressed by the series F_4 , and evaluate their intersection numbers. We also evaluate the intersection matrix for a basis of the twisted cohomology group. By using these results, we determine the monodromy representation of $E_4(a, b, c)$ and give twisted period relations, which are quadratic relations between two fundamental systems of E_4 with different parameters.

In this paper, we construct 2^m twisted cycles which represent elements of the m -th twisted homology group concerning with the integral representation (3). They imply integral representations of the solutions (2) expressed by the series F_C . We evaluate the intersection numbers of these 2^m twisted cycles. We also evaluate the intersection numbers of some elements of the twisted cohomology group. These

2010 *Mathematics Subject Classification.* 33C65.

Key words and phrases. hypergeometric functions, Lauricella's F_C , twisted cycles, twisted period relations.

results imply twisted period relations for two fundamental systems of E_C with different parameters, refer to Theorem 6.1 for their explicit forms.

In the study of twisted homology groups, twisted cycles given by bounded chambers are useful. However there are few bounded chambers in our case. By avoiding this difficulty, we succeed in constructing 2^m twisted cycles. We explain our idea in the construction. Our twisted homology group is defined by the multi-valued function

$$u(t) = \prod_{k=1}^m t_k^{1-c_k+b} \cdot v^{\sum_{k=1}^m c_k-a-m+1} \cdot w^{-b},$$

$$v = 1 - \sum_{k=1}^m t_k, \quad w = \prod_{k=1}^m t_k \cdot \left(1 - \sum_{k=1}^m \frac{x_k}{t_k}\right),$$

for fixed small positive real numbers x_1, \dots, x_m . Let $\{i_1, \dots, i_r\}$ be a subset of $\{1, \dots, m\}$ of cardinality r and let $\{j_1, \dots, j_{m-r}\}$ be its complement. We embed the direct product of an r -simplex $\tau_{i_1 \dots i_r}$ and an $(m-r)$ -simplex $\tau'_{j_1 \dots j_{m-r}}$ into the bounded chamber

$$\{(t_1, \dots, t_m) \in \mathbb{R}^m \mid t_1, \dots, t_m, v, w > 0\}.$$

We consider an m -dimensional twisted chain given by the image of $\tau_{i_1 \dots i_r} \times \tau'_{j_1 \dots j_{m-r}}$ with loading a branch of $u(t)$ on it. As is in Section 3.2.4 of [1], we can eliminate its boundary by regarding the image of $\tau_{i_1 \dots i_r}$ as the simplex

$$\{(s_{i_1}, \dots, s_{i_r}) \in \mathbb{R}^r \mid s_{i_1}, \dots, s_{i_r}, 1 - \sum_{p=1}^r s_{i_p} \geq \varepsilon\},$$

and that of $\tau'_{j_1 \dots j_{m-r}}$ as the simplex

$$\{(t_{j_1}, \dots, t_{j_{m-r}}) \in \mathbb{R}^{m-r} \mid t_{j_1}, \dots, t_{j_{m-r}}, 1 - \sum_{q=1}^{m-r} t_{j_q} \geq \varepsilon\},$$

where $s_{i_p} = x_{i_p}/t_{i_p}$ for $p = 1, \dots, r$ and ε is a certain positive real number. In this elimination of the boundary, we must regulate the difference of branches of $u(t)$ by a different way from the usual regularization. We realize this elimination by using the twisted homology group defined by another multi-valued function, see Section 4 for details. Our first main theorem states that this twisted cycle corresponds to the solution to $E_C(a, b, c)$ with the power function $\prod_{p=1}^r x_{i_p}^{1-c_{i_p}}$. Our construction of twisted cycles is also useful in the study on the system of differential equations annihilating Lauricella's hypergeometric series F_A . Refer to [5], for the twisted cycles corresponding to the solutions expressed by the series F_A and twisted period relations for F_A .

By our first main theorem and the proof of Lemma 4.1 in [6], it turns out that the intersection matrix becomes diagonal. Moreover, our construction and results in [8] enable us to evaluate the diagonal entries of the intersection matrix. If the intersection matrix for bases of twisted homology groups is evaluated, then the intersection numbers of some elements of twisted cohomology groups imply twisted period relations, which are originally identities among the integrals given by the pairings of elements of twisted homology and cohomology groups. Our first main theorem transforms these identities into quadratic relations among hypergeometric series F_C 's. Our second main theorem states these formulas in Section 6.

As is in [7], the irreducibility condition of the system $E_C(a, b, c)$ is known to be

$$a - \sum_{p=1}^r c_{i_p}, \quad b - \sum_{p=1}^r c_{i_p} \notin \mathbb{Z}$$

for any subset $\{i_1, \dots, i_r\}$ of $\{1, \dots, m\}$. Since our interest is in the property of solutions to $E_C(a, b, c)$ expressed in terms of the hypergeometric series F_C , we assume throughout this paper that the parameters a , b and $c = (c_1, \dots, c_m)$ satisfy the condition above and $c_1, \dots, c_m \notin \mathbb{Z}$.

2. DIFFERENTIAL EQUATIONS AND INTEGRAL REPRESENTATIONS

In this section, we collect some facts about Lauricella's F_C and the system E_C of hypergeometric differential equations annihilating it.

Notation 2.1. *Throughout this paper, the letter k always stands for an index running from 1 to m . If no confusion is possible, $\sum_{k=1}^m$ and $\prod_{k=1}^m$ are often simply denoted by \sum (or \sum_k) and \prod (or \prod_k), respectively. For example, under this convention $F_C(a, b, c; x)$ is expressed as*

$$F_C(a, b, c; x) = \sum_{n_1, \dots, n_m=0}^{\infty} \frac{(a, \sum n_k)(b, \sum n_k)}{\prod (c_k, n_k) \cdot \prod n_k!} \prod x_k^{n_k}.$$

Let ∂_k ($k = 1, \dots, m$) be the partial differential operator with respect to x_k . Lauricella's $F_C(a, b, c; x)$ satisfies hypergeometric differential equations

$$\begin{aligned} & \left[x_k(1-x_k)\partial_k^2 - x_k \sum_{\substack{1 \leq i \leq m \\ i \neq k}} x_i \partial_i \partial_k - \sum_{\substack{1 \leq i, j \leq m \\ i \neq k}} x_i x_j \partial_i \partial_j \right. \\ & \left. + (c_k - (a+b+1)x_k)\partial_k - (a+b+1) \sum_{\substack{1 \leq i \leq m \\ i \neq k}} x_i \partial_i - ab \right] f(x) = 0, \end{aligned}$$

for $k = 1, \dots, m$. The system generated by them is called Lauricella's system $E_C(a, b, c)$ of hypergeometric differential equations.

Proposition 2.2 ([7], [9]). *The system $E_C(a, b, c)$ is a holonomic system of rank 2^m with the singular locus*

$$(1) \quad S := \left(\prod_k x_k \cdot \prod_{\varepsilon_1, \dots, \varepsilon_m = \pm 1} (1 + \sum_k \varepsilon_k \sqrt{x_k}) = 0 \right) \subset \mathbb{C}^m.$$

If $c_1, \dots, c_m \notin \mathbb{Z}$, then the vector space of solutions to $E_C(a, b, c)$ in a simply connected domain in $D_C - S$ is spanned by the following 2^m elements:

$$(2) \quad f_{i_1 \dots i_r} := \prod_{p=1}^r x_{i_p}^{1-c_{i_p}} \cdot F_C \left(a+r - \sum_{p=1}^r c_{i_p}, b+r - \sum_{p=1}^r c_{i_p}, c^{i_1 \dots i_r}; x \right).$$

Here r runs from 0 to m , indices i_1, \dots, i_r satisfy $1 \leq i_1 < \dots < i_r \leq m$, and the row vector $c^{i_1 \dots i_r}$ is defined by

$$c^{i_1 \dots i_r} := c + 2 \sum_{p=1}^r (1 - c_{i_p}) e_{i_p},$$

where e_i is the i -th unit row vector of \mathbb{C}^m .

For the above i_1, \dots, i_r , we take j_1, \dots, j_{m-r} so that $1 \leq j_1 < \dots < j_{m-r} \leq m$ and $\{i_1, \dots, i_r, j_1, \dots, j_{m-r}\} = \{1, \dots, m\}$. It is easy to see that the i_p -th entry of $c^{i_1 \dots i_r}$ is $2 - c_{i_p}$ ($1 \leq p \leq r$) and the j_q -th entry is c_{j_q} ($1 \leq q \leq m-r$).

We denote the multi-index " $i_1 \dots i_r$ " by a letter I expressing the set $\{i_1, \dots, i_r\}$. Note that the solution (2) for $r = 0$ is $f(= f_\emptyset) = F_C(a, b, c; x)$.

Proposition 2.3 (Integral representation of Euler type, Example 3.1 in [1]). *For sufficiently small positive real numbers x_1, \dots, x_m , if $c_1, \dots, c_m, a - \sum c_k \notin \mathbb{Z}$, then $F_C(a, b, c; x)$ admits the following integral representation:*

$$(3) \quad F_C(a, b, c_1, \dots, c_m; x_1, \dots, x_m) \\ = \frac{\Gamma(1-a)}{\prod \Gamma(1-c_k) \cdot \Gamma(\sum c_k - a - m + 1)} \\ \cdot \int_{\Delta} \prod t_k^{-c_k} \cdot (1 - \sum t_k)^{\sum c_k - a - m} \cdot \left(1 - \sum \frac{x_k}{t_k}\right)^{-b} dt_1 \wedge \dots \wedge dt_m,$$

where Δ is the twisted cycle made by an m -simplex in Sections 3.2 and 3.3 of [1].

In fact, this cycle is one of twisted cycles constructed in Section 4.

3. TWISTED HOMOLOGY GROUPS

We review twisted homology groups and the intersection form between twisted homology groups in general situations, by referring to Chapter 2 of [1] and Chapters IV, VIII of [8].

For polynomials $P_j(t) = P_j(t_1, \dots, t_m)$ ($1 \leq j \leq n$), we set $D_j := \{t \mid P_j(t) = 0\} \subset \mathbb{C}^m$ and $M := \mathbb{C}^m - (D_1 \cup \dots \cup D_n)$. We consider a multi-valued function $u(t)$ on M defined as

$$u(t) := \prod_{j=1}^n P_j(t)^{\lambda_j}, \quad \lambda_j \in \mathbb{C} - \mathbb{Z} \quad (1 \leq j \leq n).$$

Let σ be a k -simplex in M , we define a loaded k -simplex $\sigma \otimes u$ by σ loading a branch of u on it. We denote the \mathbb{C} -vector space of finite sums of loaded k -simplexes by $\mathcal{C}_k(M, u)$, called the k -th twisted chain group. An element of $\mathcal{C}_k(M, u)$ is called a twisted k -chain. For a loaded k -simplex $\sigma \otimes u$ and a smooth k -form φ on M , the integral $\int_{\sigma \otimes u} u \cdot \varphi$ is defined by

$$\int_{\sigma \otimes u} u \cdot \varphi := \int_{\sigma} [\text{the fixed branch of } u \text{ on } \sigma] \cdot \varphi.$$

By the linear extension of this, we define the integral on a twisted k -chain.

We define the boundary operator $\partial^u : \mathcal{C}_k(M, u) \rightarrow \mathcal{C}_{k-1}(M, u)$ by

$$\partial^u(\sigma \otimes u) := \partial(\sigma) \otimes u|_{\partial(\sigma)},$$

where ∂ is the usual boundary operator and $u|_{\partial(\sigma)}$ is the restriction of u to $\partial(\sigma)$. It is easy to see that $\partial^u \circ \partial^u = 0$. Thus we have a complex

$$\mathcal{C}_{\bullet}(M, u) : \dots \xrightarrow{\partial^u} \mathcal{C}_k(M, u) \xrightarrow{\partial^u} \mathcal{C}_{k-1}(M, u) \xrightarrow{\partial^u} \dots,$$

and its k -th homology group $H_k(\mathcal{C}_{\bullet}(M, u))$. It is called the k -th twisted homology group. An element of $\ker \partial^u$ is called a twisted cycle.

By considering $u^{-1} = 1/u$ instead of u , we have $H_k(\mathcal{C}_{\bullet}(M, u^{-1}))$. There is the intersection pairing I_h between $H_m(\mathcal{C}_{\bullet}(M, u))$ and $H_m(\mathcal{C}_{\bullet}(M, u^{-1}))$ (in fact, the intersection pairing is defined between $H_k(\mathcal{C}_{\bullet}(M, u))$ and $H_{2m-k}(\mathcal{C}_{\bullet}(M, u^{-1}))$, however we do not consider the cases $k \neq m$). Let Δ and Δ' be elements of $H_m(\mathcal{C}_{\bullet}(M, u))$ and $H_m(\mathcal{C}_{\bullet}(M, u^{-1}))$ given by twisted cycles $\sum_i \alpha_i \cdot \sigma_i \otimes u_i$ and $\sum_j \alpha'_j \cdot \sigma'_j \otimes u_j^{-1}$ respectively, where u_i (resp. u_j^{-1}) is a branch of u (resp. u^{-1}) on σ_i (resp. σ'_j). Then their intersection number is defined by

$$I_h(\Delta, \Delta') := \sum_{i,j} \sum_{s \in \sigma_i \cap \sigma'_j} \alpha_i \alpha'_j \cdot (\sigma_i \cdot \sigma'_j)_s \cdot \frac{u_i(s)}{u_j(s)},$$

where $(\sigma_i \cdot \sigma'_j)_s$ is the topological intersection number of m -simplexes σ_i and σ'_j at s .

In this paper, we mainly consider

$$M := \mathbb{C}^m - (H_1 \cup \cdots \cup H_m \cup H \cup D),$$

where

$$\begin{aligned} H_k &:= (t_k = 0) \quad (1 \leq k \leq m), \quad H := (v = 0), \quad D := (w = 0), \\ v &:= 1 - \sum t_k, \quad w := \prod t_k \cdot \left(1 - \sum \frac{x_k}{t_k}\right). \end{aligned}$$

Note that w is a polynomial in t_1, \dots, t_m . We consider the twisted homology group on M with respect to the multi-valued function

$$\begin{aligned} u &:= \prod t_k^{1-c_k+b} \cdot v^{\sum c_k-a-m+1} w^{-b} \\ &= \prod t_k^{1-c_k} \cdot (1 - \sum t_k)^{\sum c_k-a-m+1} \cdot \left(1 - \sum \frac{x_k}{t_k}\right)^{-b} \end{aligned}$$

(the second equality holds under the coordination of branches). Proposition 2.3 means that the integral

$$\int_{\Delta} u \varphi, \quad \varphi := \frac{dt_1 \wedge \cdots \wedge dt_m}{\prod t_k \cdot (1 - \sum t_k)}$$

represents $F_C(a, b, c; x)$ modulo Gamma factors.

4. TWISTED CYCLES CORRESPONDING TO LOCAL SOLUTIONS $f_{i_1 \dots i_r}$

In this section, we construct 2^m twisted cycles in M corresponding to the solutions (2) to $E_C(a, b, c)$.

Let $0 \leq r \leq m$ and subsets $\{i_1, \dots, i_r\}$ and $\{j_1, \dots, j_{m-r}\}$ of $\{1, \dots, m\}$ satisfy $i_1 < \cdots < i_r$, $j_1 < \cdots < j_{m-r}$ and $\{i_1, \dots, i_r, j_1, \dots, j_{m-r}\} = \{1, \dots, m\}$.

Notation 4.1. *From now on, the letter p (resp. q) is always stands for an index running from 1 to r (resp. from 1 to $m-r$). We use the abbreviations \sum, \prod for the indices p, q as are mentioned in Notation 2.1.*

We set

$$M_{i_1 \dots i_r} = \mathbb{C}^m - \left(\bigcup_k (s_k = 0) \cup (v_{i_1 \dots i_r} = 0) \cup (w_{i_1 \dots i_r} = 0) \right),$$

where

$$v_{i_1 \dots i_r} := \prod_p s_{i_p} \cdot \left(1 - \sum_p \frac{x_{i_p}}{s_{i_p}} - \sum_q s_{j_q}\right), \quad w_{i_1 \dots i_r} := \prod_q s_{j_q} \cdot \left(1 - \sum_p s_{i_p} - \sum_q \frac{x_{j_q}}{s_{j_q}}\right)$$

are polynomials in s_1, \dots, s_m . Let $u_{i_1 \dots i_r}$ and $\varphi_{i_1 \dots i_r}$ be a multi-valued function and an m -form on $M_{i_1 \dots i_r}$ defined as

$$u_{i_1 \dots i_r} := \prod_k s_k^{C_k} \cdot v_{i_1 \dots i_r}^A \cdot w_{i_1 \dots i_r}^B, \quad \varphi_{i_1 \dots i_r} := \frac{ds_1 \wedge \cdots \wedge ds_m}{s_1 \cdots s_m v_{i_1 \dots i_r}},$$

where

$$A := \sum c_k - a - m + 1, \quad B := -b, \quad C_{i_p} := c_{i_p} - 1 - A, \quad C_{j_q} := 1 - c_{j_q} - B.$$

We construct a twisted cycle $\tilde{\Delta}_{i_1 \dots i_r}$ in $M_{i_1 \dots i_r}$ with respect to $u_{i_1 \dots i_r}$. Note that if $\{i_1, \dots, i_r\} = \emptyset$, then these settings coincide with those in the end of Section 3.

We choose positive real numbers $\varepsilon_1, \dots, \varepsilon_m$ and ε so that $\varepsilon < 1 - \sum_k \varepsilon_k$. And let x_1, \dots, x_m be small positive real numbers satisfying

$$x_k < \frac{\varepsilon_k}{m} \varepsilon$$

(for example, if

$$\varepsilon_k = \varepsilon = \frac{1}{3m}, \quad 0 < x_k < \frac{1}{9m^3},$$

these conditions hold). Thus the closed subset

$$\sigma_{i_1 \dots i_r} := \left\{ (s_1, \dots, s_m) \in \mathbb{R}^m \mid \begin{array}{l} s_{i_p} \geq \varepsilon_{i_p}, \quad 1 - \sum s_{i_p} \geq \varepsilon, \\ s_{j_q} \geq \varepsilon_{j_q}, \quad 1 - \sum s_{j_q} \geq \varepsilon \end{array} \right\}$$

is nonempty, since we have $(\varepsilon_1 + \frac{\delta}{2m}, \dots, \varepsilon_m + \frac{\delta}{2m}) \in \sigma_{i_1 \dots i_r}$, where $\delta := 1 - \sum \varepsilon_k - \varepsilon > 0$. Further, $\sigma_{i_1 \dots i_r}$ is contained in the bounded domain

$$\left\{ (s_1, \dots, s_m) \in \mathbb{R}^m \mid \begin{array}{l} s_k > 0, \quad 1 - \sum \frac{x_{i_p}}{s_{i_p}} - \sum s_{j_q} > 0, \\ 1 - \sum s_{i_p} - \sum \frac{x_{j_q}}{s_{j_q}} > 0 \end{array} \right\} \subset (0, 1)^m,$$

and is a direct product of an r -simplex and an $(m-r)$ -simplex. Indeed, $(s_1, \dots, s_m) \in \sigma_{i_1 \dots i_r}$ satisfies

$$\begin{aligned} 1 - \sum \frac{x_{i_p}}{s_{i_p}} - \sum s_{j_q} &> 1 - \frac{r}{m} \varepsilon - \sum s_{j_q} > 1 - \sum s_{j_q} - \varepsilon \geq 0, \\ 1 - \sum s_{i_p} - \sum \frac{x_{j_q}}{s_{j_q}} &> 1 - \sum s_{i_p} - \frac{m-r}{m} \varepsilon > 1 - \sum s_{i_p} - \varepsilon \geq 0. \end{aligned}$$

The orientation of $\sigma_{i_1 \dots i_r}$ is induced from the natural embedding $\mathbb{R}^m \subset \mathbb{C}^m$. We construct a twisted cycle from $\sigma_{i_1 \dots i_r}$. We may assume that $\varepsilon_k = \varepsilon$ (the above example satisfies this condition), and denote them by ε . Set $L_1 := (s_1 = 0), \dots, L_m := (s_m = 0), L_{m+1} := (1 - \sum s_{i_p} = 0), L_{m+2} := (1 - \sum s_{j_q} = 0)$, and let $U (\subset \mathbb{R}^m)$ be the bounded chamber surrounded by $L_1, \dots, L_m, L_{m+1}, L_{m+2}$, then $\sigma_{i_1 \dots i_r}$ is contained in U . Note that we do not consider the hyperplane L_{m+1} (resp. L_{m+2}), when $r = 0$ (resp. $r = m$). For $J \subset \{1, \dots, m+2\}$, we consider $L_J := \bigcap_{j \in J} L_j$, $U_J := \overline{U} \cap L_J$ and $T_J := \varepsilon$ -neighborhood of U_J . Then we have

$$\sigma_{i_1 \dots i_r} = U - \bigcup_J T_J.$$

Using these neighborhoods T_J , we can construct a twisted cycle $\tilde{\Delta}_{i_1 \dots i_r}$ in the same manner as Section 3.2.4 of [1] (notations L and U correspond to H and Δ in [1], respectively). Note that we have to consider contribution of branches of $s_{i_p}^{C_{i_p}} \cdot v_{i_1 \dots i_r}^A$, when we deal with the circle associated to L_{i_p} ($p = 1, \dots, r$). Indeed, for fixed positive real numbers s_k ($k \neq i_p$), s_{i_p} satisfying $1 - \sum \frac{x_{i_p}}{s_{i_p}} - \sum s_{j_q} = 0$ belongs to \mathbb{R} and we have

$$s_{i_p} = \frac{x_{i_p}}{1 - \sum_q s_{j_q} - \sum_{p' \neq p} \frac{x_{i_{p'}}}{s_{i_{p'}}}} < \frac{\frac{\varepsilon}{m} \varepsilon}{\varepsilon - (r-1) \frac{\varepsilon}{m}} = \frac{\varepsilon}{m-r+1} < \varepsilon.$$

Thus the exponent about this contribution is

$$C_{i_p} + A = c_{i_p} - 1.$$

The exponents about the contributions of the circles associated to $L_{j_q}, L_{m+1}, L_{m+2}$ are also evaluated as

$$C_{j_q} + B = 1 - c_{j_q}, \quad B = -b, \quad A = \sum c_k - a - m + 1,$$

respectively. We briefly explain the expression of $\tilde{\Delta}_{i_1 \dots i_r}$. For $j = 1, \dots, m+2$, let l_j be the $(m-1)$ -face of $\sigma_{i_1 \dots i_r}$ given by $\sigma_{i_1 \dots i_r} \cap T_j$, and let S_j be a positively

oriented circle with radius ε in the orthogonal complement of L_j starting from the projection of l_j to this space and surrounding L_j . Then $\tilde{\Delta}_{i_1 \dots i_r}$ is written as

$$\sigma_{i_1 \dots i_r} + \sum_{\emptyset \neq J \subset \{1, \dots, m+2\}} \prod_{j \in J} \frac{1}{d_j} \cdot \left(\left(\bigcap_{j \in J} l_j \right) \times \prod_{j \in J} S_j \right),$$

where

$$d_{i_p} := \gamma_{i_p} - 1, \quad d_{j_a} := \gamma_{j_a}^{-1} - 1, \quad d_{m+1} := \beta^{-1} - 1, \quad d_{m+2} := \alpha^{-1} \prod \gamma_k - 1,$$

and $\alpha := e^{2\pi\sqrt{-1}a}$, $\beta := e^{2\pi\sqrt{-1}b}$, $\gamma_k := e^{2\pi\sqrt{-1}c_k}$. Note that we define an appropriate orientation for each $(\bigcap_{j \in J} l_j) \times \prod_{j \in J} S_j$, see Section 3.2.4 of [1] for details.

Example 4.2. We give explicit forms of $\tilde{\Delta}$ and $\tilde{\Delta}_1$, for $m = 2$.

(i) In the case of $I = \emptyset$ ($r = 0$), we have

$$\begin{aligned} \tilde{\Delta} = & \sigma + \frac{S_1 \times l_1}{1 - \gamma_1^{-1}} + \frac{S_2 \times l_2}{1 - \gamma_2^{-1}} + \frac{S_4 \times l_4}{1 - \alpha^{-1} \gamma_1 \gamma_2} \\ & + \frac{S_1 \times S_2}{(1 - \gamma_1^{-1})(1 - \gamma_2^{-1})} + \frac{S_2 \times S_4}{(1 - \gamma_2^{-1})(1 - \alpha^{-1} \gamma_1 \gamma_2)} + \frac{S_4 \times S_1}{(1 - \alpha^{-1} \gamma_1 \gamma_2)(1 - \gamma_1^{-1})}, \end{aligned}$$

where the 1-chains l_j satisfy $\partial\sigma = l_1 + l_2 + l_4$ (see Figure 1), and the orientation of each direct product is induced from those of its components. Note that the face l_3 does not appear in this case.





FIGURE 1. $\tilde{\Delta}$ for $m = 2$.

(ii) In the case of $I = \{1\}$, we have

$$\begin{aligned} \tilde{\Delta}_1 = & \sigma_1 + \frac{S_1 \times l_1}{1 - \gamma_1} + \frac{S_2 \times l_2}{1 - \gamma_2^{-1}} + \frac{S_3 \times l_3}{1 - \beta^{-1}} + \frac{S_4 \times l_4}{1 - \alpha^{-1} \gamma_1 \gamma_2} \\ & + \frac{S_1 \times S_2}{(1 - \gamma_1)(1 - \gamma_2^{-1})} + \frac{S_2 \times S_3}{(1 - \gamma_2^{-1})(1 - \beta^{-1})} \\ & + \frac{S_3 \times S_4}{(1 - \beta^{-1})(1 - \alpha^{-1} \gamma_1 \gamma_2)} + \frac{S_4 \times S_1}{(1 - \alpha^{-1} \gamma_1 \gamma_2)(1 - \gamma_1)}, \end{aligned}$$

where the 1-chains l_j satisfy $\partial\sigma = \sum_{j=1}^4 l_j$ (see Figure 2), and the orientation of each direct product is induced from those of its components.

FIGURE 2. $\tilde{\Delta}_1$ for $m = 2$.

We consider the following integrals:

$$\begin{aligned}
F_{i_1 \dots i_r} &:= \int_{\tilde{\Delta}_{i_1 \dots i_r}} u_{i_1 \dots i_r} \varphi_{i_1 \dots i_r} \\
&= \int_{\tilde{\Delta}_{i_1 \dots i_r}} \prod_{p=1}^r s_{i_p}^{c_{i_p}-2} \cdot \prod_{q=1}^{m-r} s_{j_q}^{-c_{j_q}} \cdot \left(1 - \sum_{p=1}^r \frac{x_{i_p}}{s_{i_p}} - \sum_{q=1}^{m-r} s_{j_q} \right)^{\sum c_k - a - m} \\
&\quad \cdot \left(1 - \sum_{p=1}^r s_{i_p} - \sum_{q=1}^{m-r} \frac{x_{j_q}}{s_{j_q}} \right)^{-b} ds_1 \wedge \dots \wedge ds_m.
\end{aligned}$$

Proposition 4.3.

$$\begin{aligned}
F_{i_1 \dots i_r} &= \prod_p \Gamma(c_{i_p} - 1) \cdot \prod_q \Gamma(1 - c_{j_q}) \cdot \frac{\Gamma(\sum c_k - a - m + 1) \Gamma(1 - b)}{\Gamma(\sum c_{i_p} - a - r + 1) \Gamma(\sum c_{i_p} - b - r + 1)} \\
&\quad \cdot F_C \left(a + r - \sum_{p=1}^r c_{i_p}, b + r - \sum_{p=1}^r c_{i_p}, c^{i_1 \dots i_r}; x \right).
\end{aligned}$$

Proof. We compare the power series expansions of the both sides. Note that the coefficient of $x_1^{n_1} \dots x_m^{n_m}$ in the series expression of $F_C(a + r - \sum_p c_{i_p}, b + r - \sum_p c_{i_p}, c^{i_1 \dots i_r}; x)$ is

$$\begin{aligned}
A_{n_1 \dots n_m} &:= \frac{\Gamma(a + r - \sum_p c_{i_p} + \sum_k n_k)}{\Gamma(a + r - \sum_p c_{i_p})} \cdot \frac{\Gamma(b + r - \sum_p c_{i_p} + \sum_k n_k)}{\Gamma(b + r - \sum_p c_{i_p})} \\
&\quad \cdot \prod_p \frac{\Gamma(2 - c_{i_p})}{\Gamma(2 - c_{i_p} + n_{i_p})} \cdot \prod_q \frac{\Gamma(c_{j_q})}{\Gamma(c_{j_q} + n_{j_q})} \cdot \prod_k \frac{1}{n_k!}.
\end{aligned}$$

On the other hand, we have

$$\begin{aligned}
&\left(1 - \sum_p \frac{x_{i_p}}{s_{i_p}} - \sum_q s_{j_q} \right)^{\sum c_k - a - m} \\
&= \sum_{n_{i_1}, \dots, n_{i_r}} \frac{\Gamma(a - \sum_k c_k + m + \sum_p n_{i_p})}{\Gamma(a - \sum_k c_k + m)} \cdot \prod_p s_{i_p}^{-n_{i_p}} \cdot (1 - \sum_q s_{j_q})^{\sum c_k - a - m - \sum n_{i_p}} \cdot \prod_p x_{i_p}^{n_{i_p}}
\end{aligned}$$

and

$$\begin{aligned} & \left(1 - \sum_p s_{i_p} - \sum_q \frac{x_{j_q}}{s_{j_q}}\right)^{-b} \\ &= \sum_{n_{j_1}, \dots, n_{j_{m-r}}} \frac{\Gamma(b + \sum_q n_{j_q})}{\Gamma(b) \cdot \prod_q n_{j_q}!} \prod_q s_{j_q}^{-n_{j_q}} \cdot (1 - \sum_p s_{i_p})^{-b - \sum n_{j_q}} \cdot \prod_q x_{j_q}^{n_{j_q}}. \end{aligned}$$

When $r = 0$ (resp. $r = m$), we do not need the first (resp. second) expansion. The convergences of these power series expansions are verified as follows. We explain only the first one. By the construction of $\tilde{\Delta}_{i_1 \dots i_r}$, we have

$$x_k < \frac{\varepsilon_k}{m} \varepsilon, \quad |s_{i_p}| \geq \varepsilon_{i_p}, \quad \left|1 - \sum_q s_{j_q}\right| \geq \varepsilon.$$

Thus the uniform convergence on $\tilde{\Delta}_{i_1 \dots i_r}$ follows from

$$\left| \frac{1}{1 - \sum_q s_{j_q}} \cdot \sum_p \frac{x_{i_p}}{s_{i_p}} \right| \leq \frac{1}{|1 - \sum_q s_{j_q}|} \cdot \sum_p \frac{|x_{i_p}|}{|s_{i_p}|} < \frac{1}{\varepsilon} \cdot \sum_p \frac{1}{\varepsilon_{i_p}} \frac{\varepsilon_{i_p}}{m} \varepsilon = \frac{r}{m} \leq 1.$$

Since $\tilde{\Delta}_{i_1 \dots i_r}$ is constructed as a finite sum of loaded (compact) simplexes, we can exchange the sum and the integral in the expression of $F_{i_1 \dots i_r}$. Then the coefficient of $x_1^{n_1} \cdots x_m^{n_m}$ in the series expansion of $F_{i_1 \dots i_r}$ is

$$\begin{aligned} (4) \quad B_{n_1 \dots n_m} &:= \frac{\Gamma(a - \sum c_k + m + \sum n_{i_p})}{\Gamma(a - \sum c_k + m)} \cdot \frac{\Gamma(b + \sum n_{j_q})}{\Gamma(b)} \cdot \prod_k \frac{1}{n_k!} \\ &\quad \cdot \int_{\tilde{\Delta}_{i_1 \dots i_r}} \prod_p s_{i_p}^{c_{i_p} - n_{i_p} - 2} \cdot (1 - \sum_p s_{i_p})^{-b - \sum n_{j_q}} \\ &\quad \cdot \prod_q s_{j_q}^{-c_{j_q} - n_{j_q}} \cdot (1 - \sum_q s_{j_q})^{\sum c_k - a - m - \sum n_{i_p}} ds_1 \wedge \cdots \wedge ds_m. \end{aligned}$$

By the construction, the twisted cycle $\tilde{\Delta}_{i_1 \dots i_r}$ of this integral is identified with the usual regularization of the domain

$$\left\{ (s_1, \dots, s_m) \in \mathbb{R}^m \mid \begin{array}{l} s_{i_p} > 0, \quad 1 - \sum s_{i_p} > 0, \\ s_{j_q} > 0, \quad 1 - \sum s_{j_q} > 0 \end{array} \right\}$$

for the multi-valued function

$$\prod_p s_{i_p}^{c_{i_p} - n_{i_p} - 1} (1 - \sum_p s_{i_p})^{-b - \sum n_{j_q} + 1} \cdot \prod_q s_{j_q}^{-c_{j_q} - n_{j_q} + 1} \cdot (1 - \sum_q s_{j_q})^{\sum c_k - a - m - \sum n_{i_p} + 1}$$

on $\mathbb{C}^m - (\bigcup_k (s_k = 0) \cup (1 - \sum s_{i_p} = 0) \cup (1 - \sum s_{j_q} = 0))$. Hence the integral in (4) is equal to

$$\begin{aligned} & \frac{\prod \Gamma(c_{i_p} - n_{i_p} - 1) \cdot \Gamma(-b - \sum n_{j_q} + 1)}{\Gamma(-b + \sum c_{i_p} - \sum n_k - r + 1)} \\ & \cdot \frac{\prod \Gamma(-c_{j_q} - n_{j_q} + 1) \cdot \Gamma(\sum c_k - a - m - \sum n_{i_p} + 1)}{\Gamma(\sum c_{i_p} - a - \sum n_k - r + 1)}. \end{aligned}$$

Using the formula

$$(5) \quad \Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin(\pi z)},$$

we thus have

$$\frac{B_{n_1 \dots n_m}}{A_{n_1 \dots n_m}} = \prod_p \Gamma(c_{i_p} - 1) \cdot \prod_q \Gamma(1 - c_{j_q}) \cdot \frac{\Gamma(\sum c_k - a - m + 1) \Gamma(1 - b)}{\Gamma(\sum c_{i_p} - a - r + 1) \Gamma(\sum c_{i_p} - b - r + 1)},$$

which implies the proposition. \square

We define a bijection $\iota_{i_1 \dots i_r} : M_{i_1 \dots i_r} \rightarrow M$ by

$$\iota_{i_1 \dots i_r}(s_1, \dots, s_m) := (t_1, \dots, t_m); \quad t_{i_p} = \frac{x_{i_p}}{s_{i_p}}, \quad t_{j_q} = s_{j_q}.$$

For example, $\iota(= \iota_\emptyset)$ is the identity map on $M = M_\emptyset$, and $\iota_{1 \dots m}$ defines an involution on $M = M_{1 \dots m}$. We state our first main theorem.

Theorem 4.4. *We define a twisted cycle $\Delta_{i_1 \dots i_r}$ in M by*

$$(6) \quad \Delta_{i_1 \dots i_r} := (-1)^r (\iota_{i_1 \dots i_r})_* (\tilde{\Delta}_{i_1 \dots i_r}).$$

Then we have

$$\int_{\Delta_{i_1 \dots i_r}} \prod t_k^{-c_k} \cdot (1 - \sum t_k)^{\sum c_k - a - m} \cdot \left(1 - \sum \frac{x_k}{t_k}\right)^{-b} dt_1 \wedge \dots \wedge dt_m \\ \left(= \int_{\Delta_{i_1 \dots i_r}} u \varphi\right) = \prod_p x_{i_p}^{1 - c_{i_p}} \cdot F_{i_1 \dots i_r},$$

and hence this integral corresponds to the local solution $f_{i_1 \dots i_r}$ to $E_C(a, b, c)$ given in Proposition 2.2.

Proof. Pull-back the integral of the left hand side by $\iota_{i_1 \dots i_r}$. Indeed, we have

$$t_{i_p} = \frac{x_{i_p}}{s_{i_p}}, \quad dt_{i_p} = -\frac{x_{i_p}}{s_{i_p}^2} ds_{i_p}$$

by the definition of $\iota_{i_1 \dots i_r}$. Note that the sign $(-1)^r$ arising from the pull-back of $dt_1 \wedge \dots \wedge dt_m$ is canceled by that in (6). The second claim is followed from the first equality and Proposition 4.3. \square

Remark 4.5. (i) *The sign $(-1)^r$ in (6) implies that the orientation of the direct product $\iota_{i_1 \dots i_r}(\sigma_{i_1 \dots i_r})$ of two simplexes in $\Delta_{i_1 \dots i_r}$ is coincide with that of \mathbb{R}^m induced from the natural embedding $\mathbb{R}^m \subset \mathbb{C}^m$.*

(ii) *The twisted cycle Δ (for $r = 0$) equals to that mentioned in Proposition 2.3.*

The replacement $u \mapsto u^{-1} = 1/u$ and the construction same as $\Delta_{i_1 \dots i_r}$ give the twisted cycle $\Delta_{i_1 \dots i_r}^\vee$ which represents an element in $H_m(\mathcal{C}_\bullet(M, u^{-1}))$. We obtain the intersection numbers of the twisted cycles $\{\Delta_{i_1 \dots i_r}\}$ and $\{\Delta_{i_1 \dots i_r}^\vee\}$.

Theorem 4.6. (i) *For $I, J \subset \{1, \dots, m\}$ such that $I \neq J$, we have $I_h(\Delta_I, \Delta_J^\vee) = 0$.*

(ii) *The self-intersection number of $\Delta_{i_1 \dots i_r}$ is*

$$I_h(\Delta_{i_1 \dots i_r}, \Delta_{i_1 \dots i_r}^\vee) = (-1)^r \cdot \frac{\prod_q \gamma_{j_q} \cdot \left(\alpha - \prod_p \gamma_{i_p}\right) \left(\beta - \prod_p \gamma_{i_p}\right)}{\prod_k (\gamma_k - 1) \cdot \left(\alpha - \prod_k \gamma_k\right) (\beta - 1)}.$$

Proof. (i) Since $\Delta_{i_1 \dots i_r}$'s represent local solutions (2) to $E_C(a, b, c)$ by Theorem 4.4, this claim is followed from similar arguments to the proof of Lemma 4.1 in [6].

(ii) By using $\iota_{i_1 \dots i_r}$, the self-intersection number of $\Delta_{i_1 \dots i_r}$ is equal to that of $\tilde{\Delta}_{i_1 \dots i_r}$ with respect to the multi-valued function $u_{i_1 \dots i_r}$. To calculate this, we apply results of M. Kita and M. Yoshida (see [8]). Since we construct the twisted cycle $\tilde{\Delta}_{i_1 \dots i_r}$ from the direct product $\sigma_{i_1 \dots i_r}$ of two simplexes, the self-intersection number of $\tilde{\Delta}_{i_1 \dots i_r}$ is obtained as the product of those of simplexes. Thus we have

$$I_h(\Delta_{i_1 \dots i_r}, \Delta_{i_1 \dots i_r}^\vee) = \frac{1 - \prod_p \gamma_{i_p} \cdot \beta^{-1}}{\prod_p (1 - \gamma_{i_p}) \cdot (1 - \beta^{-1})} \cdot \frac{1 - \prod_q \gamma_{j_q}^{-1} \cdot \prod_k \gamma_k \cdot \alpha^{-1}}{\prod_q (1 - \gamma_{j_q}^{-1}) \cdot (1 - \prod_k \gamma_k \cdot \alpha^{-1})}.$$

□

5. INTERSECTION NUMBERS OF TWISTED COHOMOLOGY GROUPS

In this section, we review twisted cohomology groups and the intersection form between twisted cohomology groups in our situation, and evaluate some self-intersection numbers of twisted cocycles.

Recall that

$$M = \mathbb{C}^m - \left(\bigcup_k (t_k = 0) \cup (v = 0) \cup (w = 0) \right),$$

$$u = \prod t_k^{1-c_k+b} \cdot v^{\sum c_k - a - m + 1} w^{-b}.$$

We consider the logarithmic 1-form

$$\omega := d \log u = \frac{du}{u}.$$

We denote the \mathbb{C} -vector space of smooth k -forms on M by $\mathcal{E}^k(M)$. We define the covariant differential operator $\nabla_\omega : \mathcal{E}^k(M) \rightarrow \mathcal{E}^{k+1}(M)$ by

$$\nabla_\omega(\psi) := d\psi + \omega \wedge \psi, \quad \psi \in \mathcal{E}^k(M).$$

Because of $\nabla_\omega \circ \nabla_\omega = 0$, we have a complex

$$\mathcal{E}^\bullet(M) : \dots \xrightarrow{\nabla_\omega} \mathcal{E}^k(M) \xrightarrow{\nabla_\omega} \mathcal{E}^{k+1}(M) \xrightarrow{\nabla_\omega} \dots,$$

and its k -th cohomology group $H^k(M, \nabla_\omega)$. It is called the k -th twisted de Rham cohomology group. An element of $\ker \nabla_\omega$ is called a twisted cocycle. By replacing $\mathcal{E}^k(M)$ with the \mathbb{C} -vector space $\mathcal{E}_c^k(M)$ of smooth k -forms on M with compact support, we obtain the twisted de Rham cohomology group $H_c^k(M, \nabla_\omega)$ with compact support. By [2], we have $H^k(M, \nabla_\omega) = 0$ for all $k \neq m$. Further, by Lemma 2.9 in [1], there is a canonical isomorphism

$$j : H^m(M, \nabla_\omega) \rightarrow H_c^m(M, \nabla_\omega).$$

By considering $u^{-1} = 1/u$ instead of u , we have the covariant differential operator $\nabla_{-\omega}$ and the twisted de Rham cohomology group $H^k(M, \nabla_{-\omega})$. The intersection form I_c between $H^m(M, \nabla_\omega)$ and $H^m(M, \nabla_{-\omega})$ is defined by

$$I_c(\psi, \psi') := \int_M j(\psi) \wedge \psi', \quad \psi \in H^m(M, \nabla_\omega), \quad \psi' \in H^m(M, \nabla_{-\omega}),$$

which converges because of the compactness of the support of $j(\psi)$.

By the Poincaré duality (see Lemma 2.8 in [1]), we have

$$(7) \quad \begin{aligned} \dim H_k(\mathcal{C}_\bullet(M, u)) &= 0 \quad (k \neq m), \\ \dim H_m(\mathcal{C}_\bullet(M, u)) &= \dim H^m(M, \nabla_\omega). \end{aligned}$$

Proposition 5.1. *Let x_1, \dots, x_m be generic.*

- (i) *We have $\dim H_m(\mathcal{C}_\bullet(M, u)) = 2^m$.*
- (ii) *The twisted cycles $\{\Delta_I\}_I$ form a basis of $H_m(\mathcal{C}_\bullet(M, u))$.*
- (iii) *The integrations of $u\varphi$ on twisted cycles give an isomorphism between $H_m(\mathcal{C}_\bullet(M, u))$ and the space of local solutions to $E_C(a, b, c)$.*

Proof. We prove (i). By (7) and Theorem 2.2 in [1], we have

$$\dim H_m(\mathcal{C}_\bullet(M, u)) = (-1)^m \chi(M),$$

where $\chi(M)$ is the Euler characteristic of M . It is sufficient to show that $\chi(M) = (-1)^m \cdot 2^m$. Let $h \in \mathbb{C}[T_0, \dots, T_m]$ be a homogeneous polynomial defined by

$$h(T) := \prod_{i=0}^m T_i \cdot \left(\sum_{i=0}^m \prod_{j \neq i} T_j \right),$$

and let $D(h) = \{T \in \mathbb{P}^m \mid h(T) \neq 0\}$. Then we have $\chi(M) = \chi(D(h) - L)$ for some generic hyperplane L in \mathbb{P}^m . We consider the gradient map

$$\text{grad}(h) : D(h) \rightarrow \mathbb{P}^m; \quad [T_0 : \dots : T_m] \mapsto \left[\frac{\partial h}{\partial T_0}(T) : \dots : \frac{\partial h}{\partial T_m}(T) \right].$$

It is easy to see that the degree of $\text{grad}(h)$ is equal to 2^m . By Theorem 1 in [4], we obtain $\chi(D(h) - L) = (-1)^m \deg(\text{grad}(h)) = (-1)^m \cdot 2^m$, which shows (i). (The author thanks to J. Kaneko for pointing out this fact.)

The claim (ii) follows from (i), since the determinant of the intersection matrix $(I_h(\Delta_I, \Delta_J^\vee))$ is not zero by Theorem 4.6.

We show (iii). Let Sol be the space of local solutions to $E_C(a, b, c)$. By Theorem 4.4, integrals of $u\varphi$ on linear combinations of $\Delta_{i_1 \dots i_r}$'s are in Sol . Then (ii) implies that the linear map

$$\Phi : H_m(\mathcal{C}_\bullet(M, u)) \rightarrow Sol; \quad C \mapsto \int_C u\varphi$$

is defined. Proposition 2.2 and Theorem 4.4 imply that Φ is surjective. Therefore Φ is isomorphic because of $\dim H_m(\mathcal{C}_\bullet(M, u)) = \dim Sol = 2^m$. \square

We evaluate some intersection numbers of the twisted cocycles

$$\begin{aligned} \varphi &= \frac{dt_1 \wedge \dots \wedge dt_m}{\prod t_k \cdot (1 - \sum t_k)}, \\ \varphi' &:= \frac{dt_1 \wedge \dots \wedge dt_m}{vw} = \frac{dt_1 \wedge \dots \wedge dt_m}{\prod t_k \cdot (1 - \sum t_k) \cdot (1 - \sum \frac{x_k}{t_k})}. \end{aligned}$$

Theorem 5.2. (i) *The self-intersection number of φ is*

$$\begin{aligned} I_c(\varphi, \varphi) &= (2\pi\sqrt{-1})^m \left(\frac{1}{\sum c_k - a - m + 1} + \frac{1}{b + m - \sum c_k} \right) \cdot \sum_{\{I^{(r)}\}} \prod_{r=1}^{m-1} \frac{1}{b + r - \sum c_{i_p^{(r)}}}, \end{aligned}$$

where $\{I^{(r)}\}$ runs sequences of subsets of $\{1, \dots, m\}$, which satisfy

$$\{1, \dots, m\} \supseteq I^{(m-1)} \supseteq \dots \supseteq I^{(2)} \supseteq I^{(1)} \neq \emptyset,$$

and we write $I^{(r)} = \{i_1^{(r)}, \dots, i_r^{(r)}\}$.

(ii) *We have $I_c(\varphi, \varphi') = 0$.*

Proof. (i) The hypersurfaces H_1, \dots, H_m, H, D do not form a normal crossing divisor, because of $H_i \cap H_j \subset D$ for $i \neq j$. Thus we blow up \mathbb{C}^m along some intersections of hyperplanes so that the pole divisor of φ is normally crossing. Firstly we consider the blow up at the origin ($= H_1 \cap \dots \cap H_m$). Secondly we blow up this along $H_1 \cap \dots \cap \check{H}_k \cap \dots \cap H_m$ ($k = 1, \dots, m$). Repeat the blowing up process, lastly we blow up along $H_i \cap H_j$ ($1 \leq i < j \leq m$). For $i_1 < \dots < i_r$, $r \geq 2$, the exceptional divisor $E_{i_1 \dots i_r}$ arising from the blow up along $H_{i_1} \cap \dots \cap H_{i_r}$ has the exponent

$$\sum_{p=1}^r (1 - c_{i_p} + b) - (r-1)b = b + r - \sum_{p=1}^r c_{i_p}.$$

By expressing φ in each coordinates system, results in [10] give the self-intersection number of φ .

(ii) By the definition, the pole divisor of φ' does not contain the exceptional divisors. Hence the pole divisors of φ and φ' do not have m or more common factors, which implies $I_c(\varphi, \varphi') = 0$. \square

Remark 5.3. *Precisely speaking, to evaluate intersection numbers of twisted cocycles, we should blow up a compactification of \mathbb{C}^m (\mathbb{P}^m or $(\mathbb{P}^1)^m$) so that the pole divisor of $\omega = d \log u$ is normally crossing. Though the blowing up process in the above proof is not enough, it is shown that the exceptional divisors arising from the other blowing up processes do not appear as the components of the pole divisor of φ . Thus the proof is completed.*

Remark 5.4. *It seems difficult to evaluate the self-intersection number $I_c(\varphi', \varphi')$. For $m = 2$ (i.e., Appell's F_4), this number ($= \mathcal{I}_c(\varphi_{x,4}, \varphi_{x,4}$) in [6]) is written by the parameters and the factor of the defining equation of the singular locus (see [6]). Furthermore, the author does not find m -forms which form a basis of $H^m(M, \nabla_\omega)$.*

6. TWISTED PERIOD RELATIONS

By Theorem 4.6 and 5.2, we state our second main theorem.

Theorem 6.1 (Twisted period relations). *We have*

$$(8) \quad I_c(\varphi, \varphi) = \sum_I \frac{1}{I_h(\Delta_{i_1 \dots i_r}, \Delta_{i_1 \dots i_r}^\vee)} \cdot g_{i_1 \dots i_r} \cdot g_{i_1 \dots i_r}^\vee,$$

$$(9) \quad I_c(\varphi, \varphi') = \sum_I \frac{1}{I_h(\Delta_{i_1 \dots i_r}, \Delta_{i_1 \dots i_r}^\vee)} \cdot g_{i_1 \dots i_r} \cdot h_{i_1 \dots i_r}^\vee,$$

where

$$g_{i_1 \dots i_r} = \int_{\Delta_{i_1 \dots i_r}} u \varphi, \quad g_{i_1 \dots i_r}^\vee = \int_{\Delta_{i_1 \dots i_r}^\vee} u^{-1} \varphi, \quad h_{i_1 \dots i_r}^\vee = \int_{\Delta_{i_1 \dots i_r}^\vee} u^{-1} \varphi'.$$

Further, under the notations

$$a_{i_1 \dots i_r} := a - \sum c_{i_p} + r, \quad b_{i_1 \dots i_r} := b - \sum c_{i_p} + r, \quad \check{c}^{i_1 \dots i_r} := (2, \dots, 2) - c^{i_1 \dots i_r},$$

the equalities (8) and (9) are reduced to

(10)

$$\begin{aligned} & \sum_I (-1)^r \frac{1 - a_{i_1 \dots i_r}}{b_{i_1 \dots i_r}} \cdot F_C(a_{i_1 \dots i_r}, b_{i_1 \dots i_r}, c^{i_1 \dots i_r}; x) \cdot F_C(2 - a_{i_1 \dots i_r}, -b_{i_1 \dots i_r}, \check{c}^{i_1 \dots i_r}; x) \\ &= \frac{(1 - a + b) \cdot \prod (1 - c_k)}{bb_{1 \dots m}} \cdot \sum_{\{I^{(r)}\}} \prod_{r=1}^{m-1} \frac{1}{b_{I^{(r)}}}, \end{aligned}$$

(11)

$$\begin{aligned} & \sum_I (-1)^r (a_{i_1 \dots i_r} - 1) \cdot F_C(a_{i_1 \dots i_r}, b_{i_1 \dots i_r}, c^{i_1 \dots i_r}; x) \cdot F_C(2 - a_{i_1 \dots i_r}, 1 - b_{i_1 \dots i_r}, \check{c}^{i_1 \dots i_r}; x) \\ &= 0, \end{aligned}$$

respectively.

Proof. Because of the compatibility of intersection forms and pairings obtained by integrations (see [3]), we obtain the equalities (8) and (9). We show that (8) is reduced to (10). By Proposition 4.3 and Theorem 4.4, we have

$$\begin{aligned} g_{i_1 \dots i_r} &= \prod_{p=1}^r \Gamma(c_{i_p} - 1) \cdot \prod_{q=1}^{m-r} \Gamma(1 - c_{j_q}) \cdot \frac{\Gamma(\sum c_k - a - m + 1) \Gamma(1 - b)}{\Gamma(\sum c_{i_p} - a - r + 1) \Gamma(\sum c_{i_p} - b - r + 1)} \\ &\quad \cdot \prod_{p=1}^r x_{i_p}^{1 - c_{i_p}} \cdot F_C(a + r - \sum c_{i_p}, b + r - \sum c_{i_p}, c^{i_1 \dots i_r}; x). \end{aligned}$$

On the other hand, we can express $g_{i_1 \dots i_r}^\vee$ like this by the replacement

$$(a, b, c) \mapsto (2 - a, -b, (2, \dots, 2) - c),$$

since $u^{-1}\varphi$ is written as

$$u^{-1}\varphi = \prod t_k^{c_k - 2} \cdot (1 - \sum t_k)^{-\sum c_k + a + m - 2} \cdot \left(1 - \sum \frac{x_k}{t_k}\right)^b dt_1 \wedge \dots \wedge dt_m.$$

Thus we obtain

$$\begin{aligned} g_{i_1 \dots i_r}^\vee &= \prod_{p=1}^r \Gamma(1 - c_{i_p}) \cdot \prod_{q=1}^{m-r} \Gamma(c_{j_q} - 1) \\ &\quad \cdot \frac{\Gamma(-\sum c_k + a + m - 1) \Gamma(1 + b)}{\Gamma(-\sum c_{i_p} + a + r - 1) \Gamma(-\sum c_{i_p} + b + r + 1)} \\ &\quad \cdot \prod_{p=1}^r x_{i_p}^{c_{i_p} - 1} \cdot F_C(2 - a - r + \sum c_{i_p}, -b - r + \sum c_{i_p}, (2, \dots, 2) - c^{i_1 \dots i_r}; x). \end{aligned}$$

By the formula (5), we have

$$\begin{aligned} & \prod \Gamma(c_k - 1) \Gamma(1 - c_k) \cdot \frac{\Gamma(\sum c_k - a - m + 1) \Gamma(-\sum c_k + a + m - 1)}{\Gamma(\sum c_{i_p} - a - r + 1) \Gamma(-\sum c_{i_p} + a + r - 1)} \\ & \cdot \frac{\Gamma(1 - b) \Gamma(1 + b)}{\Gamma(\sum c_{i_p} - b - r + 1) \Gamma(-\sum c_{i_p} + b + r + 1)} \\ &= (2\pi\sqrt{-1})^m \cdot \frac{b}{\prod (1 - c_k) \cdot (a + m - 1 - \sum c_k)} \\ & \cdot (-1)^r \cdot \frac{a + r - 1 - \sum c_{i_p}}{b + r - \sum c_{i_p}} \cdot I_h(\Delta_{i_1 \dots i_r}, \Delta_{i_1 \dots i_r}^\vee). \end{aligned}$$

Hence, under the notations $a_{i_1 \dots i_r}$, $b_{i_1 \dots i_r}$ and $\tilde{c}^{i_1 \dots i_r}$, the equality (8) is reduced to

$$\begin{aligned} & \left(\frac{1}{1 - a_{1 \dots m}} + \frac{1}{b_{1 \dots m}} \right) \cdot \sum_{\{I^{(r)}\}} \prod_{r=1}^{m-1} \frac{1}{b_{I^{(r)}}} \\ &= \frac{b}{\prod (1 - c_k) \cdot (a_{1 \dots m} - 1)} \\ & \cdot \sum_I (-1)^r \frac{a_{i_1 \dots i_r} - 1}{b_{i_1 \dots i_r}} \cdot F_C(a_{i_1 \dots i_r}, b_{i_1 \dots i_r}, \tilde{c}^{i_1 \dots i_r}; x) \cdot F_C(2 - a_{i_1 \dots i_r}, -b_{i_1 \dots i_r}, \tilde{c}^{i_1 \dots i_r}; x). \end{aligned}$$

By multiplying $\frac{(1 - a_{1 \dots m}) \cdot \prod (1 - c_k)}{b}$, we obtain (10). The similar calculation shows that (9) is reduced to (11). \square

Note that (10) and (11) are generalizations of some equalities in Corollary 6.1 of [6].

ACKNOWLEDGMENTS

The author thanks Professor Keiji Matsumoto for his useful advice and constant encouragement. He also thanks Professor Jyoichi Kaneko for helpful comments.

REFERENCES

- [1] K. Aomoto and M. Kita, translated by K. Iohara, *Theory of Hypergeometric Functions* (Springer Verlag, New York, 2011).
- [2] K. Cho, A generalization of Kita and Noumi's vanishing theorems of cohomology groups of local system, *Nagoya Math. J.* **147** (1997) 63–69.
- [3] K. Cho and K. Matsumoto, Intersection theory for twisted cohomologies and twisted Riemann's period relations I, *Nagoya Math. J.* **139** (1995) 67–86.
- [4] A. Dimca and S. Papadima, Hypersurface complements, Milnor fibers and higher homotopy groups of arrangements, *Ann. of Math.* **158** (2003) 473–507.
- [5] Y. Goto, Twisted period relations for Lauricella's hypergeometric function F_A , preprint. arXiv:1310.6088.
- [6] Y. Goto and K. Matsumoto, The monodromy representation and twisted period relations for Appell's hypergeometric function F_4 , preprint. arXiv:1310.4243.
- [7] R. Hattori and N. Takayama, The singular locus of Lauricella's F_C , to appear in *J. Math. Soc. Japan*.
- [8] M. Kita and M. Yoshida, Intersection theory for twisted cycles, — II, Degenerate arrangements, *Math. Nachr.* **166** (1994), 287–304, **168** (1994) 171–190.
- [9] G. Lauricella, Sulle funzioni ipergeometriche a più variabili, *Rend. Circ. Mat. Palermo* **7** (1893) 111–158.
- [10] K. Matsumoto, Intersection numbers for logarithmic k -forms, *Osaka J. Math.* **35** (1998) 873–893.
- [11] M. Yoshida, *Hypergeometric functions, my love, -Modular interpretations of configuration spaces-* (Vieweg & Sohn, Braunschweig, 1997).

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO 060-0810, JAPAN
E-mail address: y-goto@math.sci.hokudai.ac.jp