

CONVEX SEPARABLY RATIONALLY CONNECTED COMPLETE INTERSECTIONS

KATSUHISA FURUKAWA

ABSTRACT. We give a generalization of a result of R. Pandharipande to arbitrary characteristic: We prove that, if X is a convex, separably rationally connected, smooth complete intersection in \mathbb{P}^N over an algebraically closed field of arbitrary characteristic, then X is rational homogeneous.

1. INTRODUCTION

F. Campana and T. Peternell [3] conjectured that, if X is a smooth Fano variety (over \mathbb{C}) with nef tangent bundle, then X is rational homogeneous. This was answered over \mathbb{C} affirmatively by Campana and Peternell in the case of $\dim(X) \leq 3$ [3] and in the case of $\dim(X) = 4$ with Picard number $\rho_X > 1$ [4], by N. Mok [10] and J. M. Hwang [8] for any 4-dimensional X , and by K. Watanabe in the case of $\dim(X) = 5$ with $\rho_X > 1$ [12].

We say that a variety X is *convex* if every morphism $\mu : \mathbb{P}^1 \rightarrow X$ satisfies $H^1(\mathbb{P}^1, \mu^*T_X) = 0$. R. Pandharipande [11] proved that, if X is a convex, rationally connected, smooth complete intersection in \mathbb{P}^N over \mathbb{C} , then X is rational homogeneous. Since a smooth Fano variety is rationally connected (over \mathbb{C}), and since nefness of the tangent bundle implies convexity, the result solved the conjecture for complete intersections.

Question 1.1. Does the statement of the above theorem of Pandharipande hold in arbitrary characteristic?

In this paper, we prove:

Theorem 1.2. *Let X be a convex, smooth complete intersection in \mathbb{P}^N over an algebraically closed field of arbitrary characteristic. Assume that: (i) there exists an immersion $\mathbb{P}^1 \rightarrow X$. Then X is of degree ≤ 2 ; in particular, X is rational homogeneous.*

Note that a complete intersection $X \subset \mathbb{P}^N$ satisfies the above condition (i) if one of the following conditions (ii-iv) holds: (ii) X is separably rationally connected; (iii) X is Fano; (iv) $2N - 2 - r \geq \sum_{i=1}^r d^i$ holds, where (d^1, \dots, d^r) is the type of X . (See Remark 4.2.) As a corollary, a smooth Fano complete intersection with nef tangent bundle is rational homogeneous in arbitrary characteristic.

2000 *Mathematics Subject Classification.* Primary 14E08, 14J45, 14M17.

Key words and phrases. convex, nef bundle, rational homogeneous space.

The paper is organized as follows. In §2, we first study the parameter space \mathcal{H} of complete intersections of type (d^1, \dots, d^r) and the incidence variety $I \subset \mathbb{G}(1, \mathbb{P}^N) \times \mathcal{H}$ parametrizing pairs (L, X) such that $L \subset X$; the techniques are based on [2], [5], [7], [9, V. 4]. A rational curve $\mu : \mathbb{P}^1 \rightarrow X$ is said to be *free* if $H^1(\mathbb{P}^1, \mu^*T_X \otimes \mathcal{O}_{\mathbb{P}^1}(-1)) = 0$. Next, we define the subset $J \subset I$ parametrizing pairs (L, X) such that L is non-free in X , and we give a calculation method of the defining polynomials of J . In §3, assuming all $d^i > 1$ and the product $\prod_{1 \leq i \leq r} d^i > 2$, we construct a pair (L, X) such that the space of non-free lines in X is smooth and of expected dimension at L . In these constructions, it is necessary to take care of the characteristic two case (see Remark 3.4). In consequence, we show that $\text{pr}_2|_J : J \rightarrow \mathcal{H}$ is surjective. Thus, under our assumption, every complete intersection X of type (d^1, \dots, d^r) has a non-free rational curve; then we have the proof of Theorem 1.2 in §4.

Acknowledgments. The author would like to thank Kiwamu Watanabe, who asked Question 1.1 after his talk at Tokyo Denki University at August 2013. Watanabe also asked “Does there exist a counter-example of Campana-Peternell conjecture in positive characteristic?”. The author was partially supported by JSPS KAKENHI Grant Number 25800030.

2. PARAMETER SPACE OF COMPLETE INTERSECTIONS

2.1. Incidence variety and its projection. We fix some notation. Let \mathcal{H}_d be the projectivization of the vector space $H^0(\mathbb{P}^N, \mathcal{O}(d))$, whose general members parametrize hypersurfaces of degrees d in \mathbb{P}^N . We take $\mathbf{d} = (d^1, \dots, d^r)$ with r positive integers d^1, \dots, d^r , and denote by $|\mathbf{d}| := \sum_{1 \leq i \leq r} d^i$. Let

$$\mathcal{H} := \mathcal{H}_{d^1} \times \cdots \times \mathcal{H}_{d^r},$$

whose general member $\mathbf{h} = (h^1, \dots, h^r)$ defines an $(N - r)$ -dimensional complete intersection $X \subset \mathbb{P}^N$ of type \mathbf{d} . For a homogeneous polynomial $\varphi \in H^0(\mathbb{P}^N, \mathcal{O}(d))$ and a line $L \subset \mathbb{P}^N$, we denote by $\varphi|_L$ the image of φ under the linear map $H^0(\mathbb{P}^N, \mathcal{O}(d)) \rightarrow H^0(L, \mathcal{O}(d))$.

Definition 2.1. We set

$$I = \{ (L, \mathbf{h}) \in \mathbb{G}(1, \mathbb{P}^N) \times \mathcal{H} \mid h^1|_L = \cdots = h^r|_L = 0 \},$$

the incidence variety whose general members parametrize pairs (L, X) such that $L \subset X$. We denote by

$$\text{pr}_1 : I \rightarrow \mathbb{G}(1, \mathbb{P}^N), \quad \text{pr}_2 : I \rightarrow \mathcal{H},$$

the first and second projections.

Remark 2.2. Each fiber of $I \rightarrow \mathbb{G}(1, \mathbb{P}^N)$ at L is isomorphic to $I_{d^1, L} \times \cdots \times I_{d^r, L}$, where we denote by $\mathcal{I}_L \subset \mathcal{O}_{\mathbb{P}^N}$ the ideal sheaf of $L \subset \mathbb{P}^N$ and by $I_{d, L} \subset \mathcal{H}_d$ the projectivization of $H^0(\mathbb{P}^N, \mathcal{I}_L(d))$. Note that $I_{d, L} \subset \mathcal{H}_d$ parametrizes hypersurfaces of degree d containing the line L . From the exact sequence

$$0 \rightarrow H^0(\mathbb{P}^N, \mathcal{I}_L(d)) \rightarrow H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(d)) \rightarrow H^0(L, \mathcal{O}_L(d)) \rightarrow 0,$$

we have $\text{codim}(I_{d,L}, \mathcal{H}_d) = d + 1$. Hence

$$\text{codim}(I, \mathcal{H} \times \mathbb{G}(1, \mathbb{P}^N)) = \text{codim}(I_{d^1,L} \times \cdots \times I_{d^r,L}, \mathcal{H}) = |\mathbf{d}| + r.$$

In addition,

$$(1) \quad \dim(I) - \dim(\mathcal{H}) = \dim \mathbb{G}(1, \mathbb{P}^N) - (|\mathbf{d}| + r) = 2N - 2 - |\mathbf{d}| - r.$$

We denote by

$$(2) \quad (S : T : Z_1 : Z_2 : \cdots : Z_{N-1})$$

the homogeneous coordinates on \mathbb{P}^N .

Definition 2.3. We take a line $L = (Z_1 = \cdots = Z_{N-1} = 0) \subset \mathbb{P}^N$. For a homogeneous polynomial $h \in H^0(\mathbb{P}^N, \mathcal{O}(d))$ with $h|_L = 0$, we define a homomorphism $\delta_L(h) : \mathcal{O}_L(1)^{\oplus N-1} \rightarrow \mathcal{O}_L(d)$ by

$$(f_1, \cdots, f_{N-1}) \mapsto h_{Z_1}|_L \cdot f_1 + \cdots + h_{Z_{N-1}}|_L \cdot f_{N-1},$$

where $h_{Z_j} := \partial h / \partial Z_j \in H^0(\mathbb{P}^N, \mathcal{O}(d-1))$.

Now, we take $\mathbf{h} = (h^1, \dots, h^r) \in \mathcal{H}$ such that $(L, \mathbf{h}) \in I$. Then, r homomorphisms $\delta_L(h^1), \dots, \delta_L(h^r)$ induce a homomorphism

$$\delta_L(\mathbf{h}) : \mathcal{O}_L(1)^{\oplus N-1} \rightarrow \bigoplus_{1 \leq i \leq r} \mathcal{O}_L(d^i) \cdot \varepsilon^i.$$

Remark 2.4. Let K be the ground field. The image of the linear map

$$H^0(\delta_L(\mathbf{h})(-1)) : H^0(L, \mathcal{O})^{\oplus N-1} = K^{\oplus N-1} \rightarrow \bigoplus_{1 \leq i \leq r} H^0(L, \mathcal{O}(d^i - 1)) \cdot \varepsilon^i$$

is equal to the vector subspace spanned by $N - 1$ elements

$$(3) \quad \sum_{1 \leq i \leq r} h_{Z_1}^i|_L \cdot \varepsilon^i, \dots, \sum_{1 \leq i \leq r} h_{Z_{N-1}}^i|_L \cdot \varepsilon^i.$$

Lemma 2.5. Let $(L, \mathbf{h}) \in I$ be as above. Let $X \subset \mathbb{P}^N$ be a complete intersection defined by \mathbf{h} , and assume that X is smooth along L . Then $\delta_L(\mathbf{h})$ is regarded as the natural map $N_{L/\mathbb{P}^N} \rightarrow N_{X/\mathbb{P}^N}|_L$,

Proof. We first consider the case of $r = 1$. The exact sequence $0 \rightarrow \mathcal{I}_L^2 \rightarrow \mathcal{I}_L \rightarrow N_{L/\mathbb{P}^N}^\vee \rightarrow 0$ induces the following linear map:

$$(4) \quad H^0(\mathbb{P}^N, \mathcal{I}_L(d)) \rightarrow H^0(L, N_{L/\mathbb{P}^N}^\vee \otimes \mathcal{O}(d)).$$

Let $h \in H^0(\mathbb{P}^N, \mathcal{I}_L(d))$ and let $X \subset \mathbb{P}^N$ be the hypersurface defined by h . Then the image of h under the above map gives a natural homomorphism $N_{L/\mathbb{P}^N} \rightarrow N_{X/\mathbb{P}^N}|_L \simeq \mathcal{O}_L(d)$ (see [7, Rem. 3.9]).

Since $L = (Z_1 = \cdots = Z_{N-1} = 0)$, we can write $h = h_1 Z_1 + \cdots + h_{N-1} Z_{N-1}$. Here the image of h in $H^0(L, N_{L/\mathbb{P}^N}^\vee \otimes \mathcal{O}(d))$ under (4) is given by $(h_1|_L, \dots, h_{N-1}|_L)$. On the other hand, we have $h_{Z_i}|_L = h_i|_L$. Hence $N_{L/\mathbb{P}^N} \rightarrow N_{X/\mathbb{P}^N}$ is identified with $\delta_L(h) : \mathcal{O}_L(1)^{\oplus N-1} \rightarrow \mathcal{O}_L(d)$.

Next, for any $r \geq 1$, considering the linear map

$$(5) \quad H^0(\mathbb{P}^N, \mathcal{I}_L \otimes \bigoplus_{1 \leq i \leq r} \mathcal{O}_{\mathbb{P}^N}(d^i)) \rightarrow H^0(L, N_{L/\mathbb{P}^N}^\vee \otimes \bigoplus_{1 \leq i \leq r} \mathcal{O}_L(d^i)),$$

instead of (4), we have the assertion in a similar way to the case of $r = 1$. \square

Remark 2.6. The linear map $H^0(\delta_L(\mathbf{h})(-1))$ is surjective if and only if L is free in X , i.e., $H^1(L, N_{L/X}(-1)) = 0$. This follows from $H^1(L, N_{L/\mathbb{P}^N}(-1)) = 0$ and the following exact sequence:

$$0 \rightarrow N_{L/X} \rightarrow N_{L/\mathbb{P}^N} \xrightarrow{\delta_L(\mathbf{h})} N_{X/\mathbb{P}^N}|_L \rightarrow 0.$$

Lemma 2.7. *Assume $2N - 2 - r \geq |\mathbf{d}|$. Then $\text{pr}_2 : I \rightarrow \mathcal{H}$ is surjective, which means that every complete intersection of type \mathbf{d} in \mathbb{P}^N contains a line.*

Proof. It holds, for example, in a similar way to [7, Proof of Prop. 3.15]. \square

Definition 2.8. We denote by J the set of pairs $(L, \mathbf{h}) \in I$ such that the linear map $H^0(\delta_L(\mathbf{h})(-1))$ is *not* surjective.

Assume that a complete intersection $X \subset \mathbb{P}^N$ defined by $\mathbf{h} \in \mathcal{H}$ is smooth along a line L . Then $(L, \mathbf{h}) \in J$ if and only if L is non-free in X , because of Remark 2.6.

Theorem 2.9. *Assume that $2N - 2 - r \geq |\mathbf{d}|$, $\prod_{1 \leq i \leq r} d^i > 2$, and all $d^i > 1$. Then $\text{pr}_2|_J : J \rightarrow \mathcal{H}$ is surjective.*

Remark 2.10. If $N \leq |\mathbf{d}|$, then we have $J = I$. This immediately follows from Remark 2.4. Thus $J \rightarrow \mathcal{H}$ is surjective, because of Lemma 2.7.

We denote by $J_{\mathbf{h}} := \text{pr}_1(\text{pr}_2^{-1}(\mathbf{h}) \cap J) \subset \mathbb{G}(1, \mathbb{P}^N)$. In the case of $|\mathbf{d}| \leq N - 2$, the above theorem follows from:

Proposition 2.11. *Assume that $N - 2 \geq |\mathbf{d}|$, $\prod_{1 \leq i \leq r} d^i > 2$, and all $d^i > 1$. Then there exists a pair $(L, \mathbf{h}) \in J$ such that $J_{\mathbf{h}}$ is smooth and of dimension $N - r - 2$ at the point $L \in \mathbb{G}(1, \mathbb{P}^N)$.*

Our goal is to construct an expected pair (L, \mathbf{h}) , satisfying the statement of Proposition 2.11. In the next subsection, as a preparation, we give defining polynomials of $J_{\mathbf{h}}$ on an affine open subset of $\mathbb{G}(1, \mathbb{P}^N)$.

2.2. Defining polynomials of the space of non-free lines. Let us fix $(s : t)$ as the homogeneous coordinates on \mathbb{P}^1 , and $(S : T : Z_1 : Z_2 : \cdots : Z_{N-1})$ on \mathbb{P}^N . We set $\mathbb{G}_1^\circ \subset \mathbb{G}(1, \mathbb{P}^N)$ to be the standard open subset, which is the set of lines L such that $L \cap (S = T = 0) = \emptyset$. Here $\mathbb{G}_1^\circ \simeq \mathbb{A}^{2(N-1)}$, which maps $L \in \mathbb{G}_1^\circ$ to

$$A_L = \begin{bmatrix} a_1, \dots, a_{N-1} \\ b_1, \dots, b_{N-1} \end{bmatrix} \in \mathbb{A}^{2(N-1)}$$

if the line $L \subset \mathbb{P}^N$ is equal to the image of the morphism $\mathbb{P}^1 \rightarrow \mathbb{P}^N$ defined by

$$(6) \quad (s : t) \mapsto [s \ t] \cdot [E_2 \ A_L] = (s : t : sa_1 + tb_1 : \cdots : sa_{N-1} + tb_{N-1}),$$

where E_2 is the unit matrix of size 2. We set

$$\xi : \mathbb{G}_1^\circ \times \mathbb{P}^1 \rightarrow \mathbb{P}^N$$

to be the morphism which sends $(L, (s : t)) \mapsto \begin{bmatrix} s & t \end{bmatrix} \cdot \begin{bmatrix} E_2 & A_L \end{bmatrix}$.

Let $X \subset \mathbb{P}^N$ be a complete intersection defined by homogeneous polynomials $\mathbf{h} = (h^1, \dots, h^r) \in \mathcal{H}$. The composite functions $h^i \circ \xi$ is regarded as a polynomial with variables $s, t, a_1, \dots, a_{N-1}, b_1, \dots, b_{N-1}$. Then $h^i \circ \xi$ is written as

$$f_0^i s^{d^i} + f_1^i s^{d^i-1} t + \dots + f_{d^i}^i t^{d^i},$$

with some $f_0^i, f_1^i, \dots, f_{d^i}^i \in K[a_1, \dots, a_{N-1}, b_1, \dots, b_{N-1}]$. Here $L \subset X$ if and only if

$$f_k^i(A_L) = 0 \text{ for any } 1 \leq i \leq r \text{ and } 0 \leq k \leq d^i.$$

Let $F_{\mathbf{h}} := \text{pr}_1(\text{pr}_2^{-1}(\mathbf{h}) \cap I) \subset \mathbb{G}(1, \mathbb{P}^N)$, which is equal to the set of lines lying in X . Then $F_{\mathbf{h}} \cap \mathbb{G}_1^\circ$ is equal to the zero set of $(r + |\mathbf{d}|)$ -polynomials f_k^i 's in the affine space $\mathbb{G}_1^\circ \simeq \mathbb{A}^{2(N-1)}$.

Remark 2.12. For a point x , we denote by $F_{\mathbf{h},x}$ the set of $L \in F_{\mathbf{h}}$ such that $x \in L$. Let $x = (1 : 0 : \dots : 0)$. Then $F_{\mathbf{h},x} \cap \mathbb{G}_1^\circ$ is equal to the zero set of f_k^i 's and a_1, a_2, \dots, a_{N-1} in $\mathbb{G}_1^\circ \simeq \mathbb{A}^{2(N-1)}$.

For a homogeneous polynomial $h \in K[S, T, Z_1, \dots, Z_{N-1}]$ of degree d with $h \circ \xi = f_0 s^d + f_1 s^{d-1} t + \dots + f_d t^d$, we have

$$(7) \quad \begin{aligned} (h \circ \xi)_{a_j} &= f_{0,a_j} s^d + f_{1,a_j} s^{d-1} t + \dots + f_{d,a_j} t^d, \\ (h \circ \xi)_{b_j} &= f_{0,b_j} s^d + f_{1,b_j} s^{d-1} t + \dots + f_{d,b_j} t^d, \end{aligned}$$

where $(h \circ \xi)_{a_j} := \partial(h \circ \xi) / \partial a_j$ and so on. In addition, it follows that

$$(8) \quad (h \circ \xi)_{a_j} = s \cdot (h_{Z_j} \circ \xi) \text{ and } (h \circ \xi)_{b_j} = t \cdot (h_{Z_j} \circ \xi).$$

This is shown as follows: We write the morphism ξ by the coordinates $(\eta_0 : \eta_1 : \xi_1 : \dots : \xi_{N-1})$, which are given as in the right hand side of (6). Then $(h \circ \xi)_{a_j}$ is equal to $\partial \eta_0 / \partial a_j \cdot (\partial h / \partial S) \circ \xi + \partial \eta_1 / \partial a_j \cdot (\partial h / \partial T) \circ \xi + \sum_{1 \leq l \leq N-1} \partial \xi_l / \partial a_j \cdot (\partial h / \partial Z_l) \circ \xi = s \cdot (h_{Z_j} \circ \xi)$. Similarly, $(h \circ \xi)_{b_j}$ is also obtained.

We denote by $v(h_{Z_j} \circ \xi)$ the row vector of size d which represents $h_{Z_j} \circ \xi$ with respect to the basis $s^{d-1}, s^{d-2}t, \dots, t^{d-1}$ of $H^0(\mathbb{P}^1, \mathcal{O}(d-1))$, i.e.,

$$h_{Z_j} \circ \xi = v(h_{Z_j} \circ \xi) \cdot {}^t [s^{d-1} \quad s^{d-2}t \quad \dots \quad t^{d-1}].$$

Each entry of $v(h_{Z_j} \circ \xi)$ is a polynomial with variables $a_1, \dots, a_{N-1}, b_1, \dots, b_{N-1}$.

Now, for $\mathbf{h} = (h^1, \dots, h^r)$, let us consider the $(N-1) \times |\mathbf{d}|$ matrix

$$(9) \quad M(\mathbf{h}) := \begin{bmatrix} v(h_{Z_1}^1 \circ \xi) & \dots & v(h_{Z_1}^r \circ \xi) \\ \vdots & & \vdots \\ v(h_{Z_{N-1}}^1 \circ \xi) & \dots & v(h_{Z_{N-1}}^r \circ \xi) \end{bmatrix},$$

where it follows from Remark 2.4 that $M(\mathbf{h})(A_L)$ is of rank $|\mathbf{d}|$ if and only if $H^0(\delta_L(\mathbf{h})(-1))$ is surjective. Then $J_{\mathbf{h}} \cap \mathbb{G}_1^\circ$ is the zero set in $F_{\mathbf{h}} \cap \mathbb{G}_1^\circ$ of the polynomials given as the $|\mathbf{d}| \times |\mathbf{d}|$ minors of $M(\mathbf{h})$.

For a polynomial f with varieties $a_1, \dots, a_{N-1}, b_1, \dots, b_{N-1}$, we set

$$\begin{aligned} D_a(f)_L &:= [f_{a_1}(A_L) \quad f_{a_2}(A_L) \quad \cdots \quad f_{a_{N-1}}(A_L)], \\ D_b(f)_L &:= [f_{b_1}(A_L) \quad f_{b_2}(A_L) \quad \cdots \quad f_{b_{N-1}}(A_L)]. \end{aligned}$$

We denote by $\mathbf{e}_i = [e_{i,1} \quad e_{i,2} \quad \cdots \quad e_{i,N-1}]$ the row vector of size $N-1$ such that $e_{i,i} = 1$ and $e_{i,j} = 0$ for $i \neq j$, and by $\mathbf{0}$ the zero vector of size $N-1$.

Remark 2.13. Let g_1, \dots, g_m be a minimal subset of the $|\mathbf{d}| \times |\mathbf{d}|$ minors of $M(\mathbf{h})$ such that $J_{\mathbf{h}} \cap \mathbb{G}_1^\circ$ is locally defined in $\mathbb{G}_1^\circ \simeq \mathbb{A}^{2(N-1)}$ around L by the $|\mathbf{d}| + r + m$ polynomials g_1, \dots, g_m and f_k^i ($1 \leq i \leq r$, $0 \leq k \leq d^i$). We consider the $(|\mathbf{d}| + r + m) \times 2(N-1)$ matrix

$$(10) \quad \left[\begin{array}{cc} \left(\begin{array}{c} \vdots \\ D_a(f_k^i) \\ \vdots \\ D_b(f_k^i) \end{array} \right) \\ \left(\begin{array}{c} D_a(g_1) \\ \vdots \\ D_a(g_m) \end{array} \right) \end{array} \quad \left(\begin{array}{c} \vdots \\ D_b(g_1) \\ \vdots \\ D_b(g_m) \end{array} \right) \right]_L.$$

Then $J_{\mathbf{h}}$ is smooth at L if and only if the rank of the above matrix (10) is equal to $|\mathbf{d}| + r + m$. We note that the rank can be calculated by using formulae (7), (8), (9).

Lemma 2.14. *Every irreducible component of J is of codimension $\leq N - |\mathbf{d}|$ in I .*

Proof. We consider the morphism $\Phi : I \cap (\mathbb{G}_1^\circ \times \mathcal{H}) \rightarrow \mathbb{A}^{(N-1)d}$ which sends $(L, \mathbf{h}) \mapsto M(\mathbf{h})(A_L)$. Let $M_{|\mathbf{d}|-1} \subset \mathbb{A}^{(N-1)d}$ the locus of matrix of rank $\leq |\mathbf{d}| - 1$. Then $J \cap (\mathbb{G}_1^\circ \times \mathcal{H})$ is equal to $\Phi^{-1}(M_{|\mathbf{d}|-1})$. Since $M_{|\mathbf{d}|-1}$ is of codimension $N - |\mathbf{d}|$ in $\mathbb{A}^{(N-1)d}$ (see [1, p. 67, II, §2, Prop.]), $J \cap (\mathbb{G}_1^\circ \times \mathcal{H})$ is of codimension $\leq N - |\mathbf{d}|$ in I , which implies the assertion. \square

3. CONSTRUCTION OF EXPECTED PAIRS

3.1. Hypersurfaces. In this subsection, we assume $r = 1$ and give construction of pairs satisfying the statement of Proposition 2.11.

Remark 3.1. Let $(L, h) \in \mathbb{G}(1, \mathbb{P}^N) \times \mathcal{H}_d$ satisfies $h|_L = 0$. Then the linear map $H^0(\delta_L(h)(-1)) : K^{\oplus N-1} \rightarrow H^0(L, \mathcal{O}(d-1))$ given in Definition 2.3 is not surjective if and only if

$$\psi(h_{Z_1}|_L) = \cdots = \psi(h_{Z_{N-1}}|_L) = 0$$

holds for some linear functional $\psi : H^0(\mathbb{P}^1, \mathcal{O}(d-1)) \rightarrow K$. A linear functional ψ is represented by the row vector

$$(11) \quad [c_0 \quad c_1 \quad \cdots \quad c_{d-1}] \quad (c_0, \dots, c_{d-1} \in K);$$

this means that $\psi(s^{d-1-k}t^k) = c_k$ for each k .

First we consider the case $(d, N) = (4, 6)$. Let $(S : T : Z_1 : \dots : Z_5)$ be the homogeneous coordinates on \mathbb{P}^6 , and $(s : t)$ be on \mathbb{P}^1 .

Example 3.2. Let $\psi : H^0(\mathbb{P}^1, \mathcal{O}(3)) \rightarrow K$ be a general linear functional, where we assume that $c_0 = 1$ in the expression (11). Let $X \subset \mathbb{P}^6$ be a hypersurface defined by the following homogeneous polynomial of degree 4,

$$h := h_1 Z_1 + h_2 Z_2 + h_3 Z_3 + T^2 Z_4 Z_5,$$

where we set

$$h_j := c_j S^3 - S^{3-j} T^j \quad (j = 1, 2, 3).$$

Let $L = (Z_1 = \dots = Z_5 = 0) \subset \mathbb{P}^6$. Then $(L, h) \in J$, and J_h is smooth and of dimension $N - 3 = 3$ at the point $L \in \mathbb{G}(1, \mathbb{P}^6)$. The reason is as follows.

It follows that $h_{Z_j} \circ \xi = h_j \circ \xi = c_j s^3 - s^{3-j} t^j$ ($j = 1, 2, 3$). In particular, we find $(L, h) \in J$ as in Remark 3.1. From (7), (8), since $s \cdot h_{Z_j} \circ \xi = c_j s^4 - s^{4-j} t^j$, it follows that $f_{0,a_j} = c_j$, $f_{j,a_j} = -1$, and other $f_{j',a_j} = 0$. In addition, since $t \cdot h_{Z_j} \circ \xi = c_j s^3 t - s^{3-j} t^{j+1}$, it follows that $f_{1,b_j} = c_j$, $f_{j+1,b_j} = -1$, and other $f_{j',b_j} = 0$. Hence, we have

$$\begin{bmatrix} D_a(f_0) \\ D_a(f_1) \\ D_a(f_2) \\ D_a(f_3) \\ D_a(f_4) \end{bmatrix}_L = \begin{bmatrix} c_1 \mathbf{e}_1 + c_2 \mathbf{e}_2 + c_3 \mathbf{e}_3 \\ -\mathbf{e}_1 \\ -\mathbf{e}_2 \\ -\mathbf{e}_3 \\ \mathbf{0} \end{bmatrix}, \quad \begin{bmatrix} D_b(f_0) \\ D_b(f_1) \\ D_b(f_2) \\ D_b(f_3) \\ D_b(f_4) \end{bmatrix}_L = \begin{bmatrix} \mathbf{0} \\ c_1 \mathbf{e}_1 + c_2 \mathbf{e}_2 + c_3 \mathbf{e}_3 \\ -\mathbf{e}_1 \\ -\mathbf{e}_2 \\ -\mathbf{e}_3 \end{bmatrix}.$$

Here, $[D_a(f_0) \ D_b(f_0)]_L$ is transformed to

$$[\mathbf{0} \ (c_1^2 - c_2) \mathbf{e}_1 + (c_1 c_2 - c_3) \mathbf{e}_2]$$

under elementary row operations of $[D_a(f_k) \ D_b(f_k)]_L$ with $1 \leq k \leq 4$.

On the other hand, from (9), we have

$$(12) \quad M(h) = \begin{bmatrix} c_1 & -1 & & & & \\ c_2 & & -1 & & & \\ c_3 & & & -1 & & \\ & a_5 & b_5 & & & \\ & a_4 & b_4 & & & \end{bmatrix} \sim \begin{bmatrix} & 0 & -1 & & & \\ & 0 & & -1 & & \\ & 0 & & & -1 & \\ c_2 a_5 + c_3 b_5 & & a_5 & b_5 & & \\ c_2 a_4 + c_3 b_4 & & a_4 & b_4 & & \end{bmatrix}$$

where the right hand side is a transformation under elementary column operations, and where we consider blank entries in the matrix to be filled by 0. Here $\text{rk } M(h) < 4$ if and only if 4×4 minors $g_1 = c_2 a_4 + c_3 b_4$ and $g_2 = c_2 a_5 + c_3 b_5$ are equal to zero. Now we have

$$\begin{bmatrix} D_a(g_1) \\ D_a(g_2) \end{bmatrix}_L = \begin{bmatrix} c_2 \mathbf{e}_4 \\ c_2 \mathbf{e}_5 \end{bmatrix}, \quad \begin{bmatrix} D_b(g_1) \\ D_b(g_2) \end{bmatrix}_L = \begin{bmatrix} c_3 \mathbf{e}_4 \\ c_3 \mathbf{e}_5 \end{bmatrix}.$$

Hence the matrix (10) in Remark 2.13 is of rank 7, which implies that $J_h \cap \mathbb{G}_1^\circ$, the zero set of 7 polynomials $f_0, \dots, f_4, g_1, g_2$, is smooth and of codimension 7 in $\mathbb{G}_1^\circ \simeq \mathbb{A}^{10}$ at L .

Here we have:

$$\begin{bmatrix} D_a(g_1) \\ D_a(g_2) \\ D_a(g_3) \\ D_a(g_4) \\ \vdots \\ D_a(g_{N-d}) \end{bmatrix}_L = \begin{bmatrix} c_{d-3}e_{d+1} \\ c_{d-3}e_d + c_{d-2}e_{d+2} \\ c_{d-2}e_{d+1} \\ c_{d-2}e_{d+3} \\ \vdots \\ c_{d-2}e_{N-1} \end{bmatrix}, \quad \begin{bmatrix} D_b(g_1) \\ D_b(g_2) \\ D_b(g_3) \\ D_b(g_4) \\ \vdots \\ D_b(g_{N-d}) \end{bmatrix}_L = \begin{bmatrix} c_{d-2}e_{d+1} \\ c_{d-2}e_d + c_{d-1}e_{d+2} \\ c_{d-1}e_{d+1} \\ c_{d-1}e_{d+3} \\ \vdots \\ c_{d-1}e_{N-1} \end{bmatrix}.$$

Since $c_1, \dots, c_{d-1} \in K$ are general, the matrix (10) in Remark 2.13 is of rank $N + 1$, which implies that $J_h \cap \mathbb{G}_1^\circ$ is smooth and of codimension $N + 1$ in $\mathbb{G}_1^\circ \simeq \mathbb{A}^{2(N-1)}$ at the point L .

Remark 3.4. Let the characteristic is *not* equal to 2. Then, by taking \tilde{h} to be

$$T^{d-2}(Z_d^2 + \dots + Z_{N-1}^2),$$

we can show the same statement of Example 3.3 with easier calculations. However, this does not work in characteristic 2.

3.2. Complete intersections. First we assume that $d^i \geq 3$ for some i . Then Example 3.3 is generalized to the case of $r \geq 2$. For example, we can calculate the case of $(r, N, (d^1, d^2)) = (2, 9, (4, 3))$, as follows.

Example 3.5. Let $\psi : H^0(\mathbb{P}^1, \mathcal{O}(3)) \rightarrow K$ be a general linear functional with $c_0 = 1$ in (11). Let $X \subset \mathbb{P}^9$ be a complete intersection defined by the following homogeneous polynomials h^1, h^2 of degrees 4, 3:

$$\begin{aligned} h^1 &:= h_1^1 Z_1 + h_2^1 Z_2 + h_3^1 Z_3 + T^2 Z_4 Z_5 \quad (h_j^1 := c_j S^3 - S^{3-j} T^j), \\ h^2 &:= S^2 Z_6 + T Z_7 + T^2 Z_8. \end{aligned}$$

Let $L = (Z_1 = \dots = Z_8 = 0) \subset \mathbb{P}^9$. Then $(L, (h^1, h^2)) \in J$, and $J_{(h^1, h^2)}$ is smooth and of dimension $N - r - 2 = 5$ at the point $L \in \mathbb{G}(1, \mathbb{P}^9)$. The reason is as follows.

It follows that $h_{Z_j}^1 \circ \xi = h_j^1 \circ \xi = c_j s^3 - s^{3-j} t^j$ ($j = 1, 2, 3$). Thus we find that $H^0(\delta_L(h^1)(-1))$ is not surjective as in Remark 3.1. This implies that $H^0(\delta_L(h^1, h^2)(-1))$ is also not surjective (see Remark 2.4); hence $(L, (h^1, h^2)) \in J$. The calculation of $D_a(f_k^1)_L$ and $D_b(f_k^1)_L$ with $0 \leq k \leq 4$ is the same as Example 3.2. On the other hand, from (7), (8), we have

$$\begin{bmatrix} D_a(f_0^2) \\ D_a(f_1^2) \\ D_a(f_2^2) \\ D_a(f_3^2) \end{bmatrix}_L = \begin{bmatrix} e_6 \\ e_7 \\ e_8 \\ \mathbf{0} \end{bmatrix}, \quad \begin{bmatrix} D_b(f_0^2) \\ D_b(f_1^2) \\ D_b(f_2^2) \\ D_b(f_3^2) \end{bmatrix}_L = \begin{bmatrix} \mathbf{0} \\ e_6 \\ e_7 \\ e_8 \end{bmatrix}.$$

Next, from (9), we have

$$M(h^1, h^2) = \begin{bmatrix} M' & \\ & E_3 \end{bmatrix},$$

where M' is a 5×5 matrix which is the same as the left hand side of (12). Thus, in a similar way to Example 3.2, we find that $M(h^1, h^2)$ is of rank < 7 if and only if the 7×7 minors g_1, g_2 are zero. As a result, the matrix (10) in Remark 2.13 is of rank 11, which implies that $J_{(h^1, h^2)} \cap \mathbb{G}_1^\circ$ is smooth and of codimension 11 in $\mathbb{G}_1^\circ \simeq \mathbb{A}^{16}$ at L .

We denote by $|\mathbf{d}|_{i_0} := \sum_{i=i_0}^r d^i$. Then $|\mathbf{d}| = |\mathbf{d}|_1$ and $|\mathbf{d}|_r = d^r$.

Example 3.6. Assume that $d^1 \geq 3$ and $\sum d^i \leq N - 2$. Let $\psi : H^0(\mathbb{P}^1, \mathcal{O}(d^1 - 1)) \rightarrow K$ be a general linear functional with $c_0 = 1$ in (11). Let $X \subset \mathbb{P}^N$ be a hypersurface defined by the following homogeneous polynomials h^1, \dots, h^r of degrees d^1, \dots, d^r . We set $h^1 = h_1^1 Z_1 + h_2^1 Z_2 + \dots + h_{d^1-1}^1 Z_{d^1-1} + \tilde{h}^1$, where $h_j^1 := c_j S^{d^1-1} - S^{d^1-j-1} T^j$, and where

$$\tilde{h}^1 := T^{d^1-2} (Z_{d^1} Z_{d^1+1} + Z_{d^1+2} Z_{d^1+3} + \dots + Z_{N-2-|\mathbf{d}|_2} Z_{N-1-|\mathbf{d}|_2})$$

if $N - |\mathbf{d}|$ is even, or

$$\tilde{h}^1 := S T^{d^1-3} Z_{d^1} Z_{d^1+1} + T^{d^1-2} (Z_{d^1+1} Z_{d^1+2} + Z_{d^1+3} Z_{d^1+4} + \dots + Z_{N-2-|\mathbf{d}|_2} Z_{N-1-|\mathbf{d}|_2})$$

if $N - |\mathbf{d}|$ is odd. Next we set h^2, \dots, h^r by

$$h^i = S^{d^i-1} Z_{N-|\mathbf{d}|_i} + S^{d^i-2} T Z_{N+1-|\mathbf{d}|_i} + \dots + T^{d^i-1} Z_{N-1-|\mathbf{d}|_{i+1}} \quad (2 \leq i \leq r).$$

Let $L = (Z_1 = \dots = Z_{N-1} = 0)$. Then, in a similar way to Example 3.3 and Example 3.5, one can show that $(L, (h^1, \dots, h^r)) \in J$ and that $J_{(h^1, \dots, h^r)}$ is smooth and of dimension $N - r - 2$ at the point $L \in \mathbb{G}(1, \mathbb{P}^N)$.

3.3. Complete intersections of hyperquadrics. We complete the case where $r \geq 2$ and $d^1 = \dots = d^r = 2$. For example, for $(N, r, (d^1, d^2)) = (7, 2, (2, 2))$, we have:

Example 3.7. Let $X \subset \mathbb{P}^7$ be the complete intersection defined by the following homogeneous quadric polynomials:

$$h^1 = S Z_1 + T Z_2 + Z_5 Z_6, \quad h^2 = S Z_2 + T Z_3 + Z_4 Z_5.$$

Let $L = (Z_1 = \dots = Z_6 = 0)$. Then $(L, (h^1, h^2)) \in J$, and $J_{(h^1, h^2)}$ is smooth and of dimension $N - r - 2 = 3$ at the point $L \in \mathbb{G}(1, \mathbb{P}^7)$. The reason is as follows.

It follows that

$$[h_{Z_1}^1|_L, h_{Z_1}^2|_L] = [s, 0], \quad [h_{Z_2}^1|_L, h_{Z_2}^2|_L] = [t, s], \quad [h_{Z_3}^1|_L, h_{Z_3}^2|_L] = [0, t]$$

and that other $[h_{Z_i}^1|_L, h_{Z_i}^2|_L]$'s are zero. Thus the above elements does not span the 4-dimensional vector space $H^0(L, \mathcal{O}(1)) \oplus H^0(L, \mathcal{O}(1))$. Then, from Remark 2.4, we have $(L, (h^1, h^2)) \in J$.

3.4. Surjectivity of the projection from J to \mathcal{H} . First, we consider the remained part, $|\mathbf{d}| = N - 1$:

Lemma 3.9. *If $|\mathbf{d}| = N - 1$, then the statement of Theorem 2.9 holds.*

Remark 3.10. Since all $d^i \geq 2$, we have $2r \leq |\mathbf{d}| < N$. Hence $2(N - r) = 2N - 2r > N$. Then [6, Cor. 2] implies that a smooth complete intersection $X \subset \mathbb{P}^N$ of type \mathbf{d} is simply connected.

Proof of Lemma 3.9. Assume that $\text{pr}_2(J) \neq \mathcal{H}$, and take a general $\mathbf{h} \in \mathcal{H}$ such that $\mathbf{h} \notin \text{pr}_2(J)$. Let $X \subset \mathbb{P}^N$ be a complete intersection defined by \mathbf{h} . We denote by $F_{\mathbf{h}} \subset \mathbb{G}(1, \mathbb{P}^N) := \text{pr}_1(\text{pr}_2^{-1}(\mathbf{h}))$, which is the space of lines lying in X , and consider universal family $\text{Univ} \subset F_{\mathbf{h}} \times X$ with projection $u : \text{Univ} \rightarrow X$. Here $u^{-1}(x)$ is identified with the space of lines lying in X passing through a point x . Since $\text{pr}_2^{-1}(\mathbf{h}) \cap J = \emptyset$, every line $L \in F_{\mathbf{h}}$ is free, where we have $N_{L/X} = \mathcal{O}_L^{\oplus N-1}$. Since $h^1(L, N_{L/X}(-1)) = 0$, every fiber $u^{-1}(x)$ is smooth and of dimension $h^0(L, N_{L/X}(-1)) = 0$. Hence u is étale morphism. Since X is simply connected, u is isomorphic. This contradicts that the length of $u^{-1}(x)$ is > 1 if X is general (see Remark 2.12). \square

Proof of Proposition 2.11. It follows from Examples 3.6 and 3.8. \square

Now, we have the surjectivity of $\text{pr}_2|_J : J \rightarrow \mathcal{H}$.

Proof of Theorem 2.9. The case of $|\mathbf{d}| \geq N - 1$ follows immediately from Remark 2.10 and Lemma 3.9.

Assume $|\mathbf{d}| \leq N - 2$. From Proposition 2.11, there exists a pair $(L, \mathbf{h}) \in J$ such that $\text{pr}_2^{-1}(\mathbf{h}) \cap J \simeq J_{\mathbf{h}}$ is smooth and of dimension $N - r - 2$ at (L, \mathbf{h}) . We consider the exact sequence

$$0 \rightarrow T_{(L, \mathbf{h})}(\text{pr}_2^{-1}(\mathbf{h}) \cap J) \rightarrow T_{(L, \mathbf{h})}J \xrightarrow{d_{(L, \mathbf{h})}(\text{pr}_2|_J)} T_{\mathbf{h}}\mathcal{H}.$$

From Lemma 2.14 and the equality (1) in Remark 2.2, we have

$$\begin{aligned} \dim(T_{(L, \mathbf{h})}J) - \dim(T_{(L, \mathbf{h})}(\text{pr}_2^{-1}(\mathbf{h}) \cap J)) \\ \geq (\dim(I) - (N - |\mathbf{d}|)) - (N - r - 2) = \dim(\mathcal{H}), \end{aligned}$$

which implies that the linear map $d_{(L, \mathbf{h})}(\text{pr}_2|_J)$ is surjective. Hence the morphism $\text{pr}_2|_J$ is smooth on an open neighborhood of (L, \mathbf{h}) in J , and hence, is surjective. \square

4. NON-FREE RATIONAL CURVES

From Theorem 2.9, there exists a non-free line lying in a complete intersection $X \subset \mathbb{P}^N$ in the case of $2N - 2 - r \geq |\mathbf{d}|$ and $\deg(X) > 2$. On the other hand, for $|\mathbf{d}| > N$, the following lemma is immediately obtained:

Lemma 4.1. *Let $X \subset \mathbb{P}^N$ be a smooth complete intersection of type $\mathbf{d} = (d_1, \dots, d_r)$. If $|\mathbf{d}| > N$, then every immersion $\mu : \mathbb{P}^1 \rightarrow X$ is non-free.*

Proof. Let $\mu^*T_X \simeq \bigoplus_{i=1}^{N-r} \mathcal{O}_{\mathbb{P}^1}(a_i)$ be the splitting on \mathbb{P}^1 with integers $a_1 \geq a_2 \geq \dots \geq a_{N-r}$. Taking the positive integer b with $\mu^*(\mathcal{O}_X(1)) = \mathcal{O}_{\mathbb{P}^1}(b)$, since $\bigwedge^{N-r} T_X = \mathcal{O}_X(N+1-|\mathbf{d}|)$, we have $\sum_{1 \leq i \leq N-r} a_i = b(N+1-|\mathbf{d}|) \leq 0$. Since $T_{\mathbb{P}^1} \rightarrow \mu^*T_X$ is injective, we find that $a_1 \geq 2$. Therefore $a_{i_0} < 0$ for some i_0 . Then $H^1(\mathbb{P}^1, \mu^*T_X \otimes \mathcal{O}_{\mathbb{P}^1}(-1)) \neq 0$. \square

Remark 4.2. We check that each of the conditions (ii-iv) in §1 implies the condition (i) in Theorem 1.2. First, (iv) implies (i); this is because, X contains a line if (iv) holds, as in Lemma 2.7. Next, (ii) or (iii) implies (iv), as follows. From [9, IV, Thm. 3.7], X is separably rationally connected if and only if there exists $\mu : \mathbb{P}^1 \rightarrow X$ such that μ^*T_X is ample, where the latter condition implies $|\mathbf{d}| \leq N$. On the other hand, X is Fano if and only if $|\mathbf{d}| \leq N$.

Proof of Theorem 1.2. Let $X \subset \mathbb{P}^N$ be a complete intersection of type $\mathbf{d} = (d^1, \dots, d^r)$ satisfying the assumption of Theorem 1.2. Without loss of generality, we may assume that all $d^i > 1$.

Suppose that $\prod d^i = \deg(X) > 2$. Then we can find a non-free immersion $\mu : \mathbb{P}^1 \rightarrow X$, as follows. Assume $2N-2-r \geq |\mathbf{d}|$. Taking defining polynomials $\mathbf{h} = (h^1, \dots, h^r)$ of X , from Theorem 2.9, we have that $J \cap \text{pr}_2^{-1}(\mathbf{h}) \neq \emptyset$; this means that a non-free line in X exists. Assume $2N-2-r < |\mathbf{d}|$. We may also assume $N-r \geq 2$. Then $N < |\mathbf{d}|$ holds. By the condition (i) in Theorem 1.2, an immersion exists and is non-free because of Lemma 4.1.

Let $\mu : \mathbb{P}^1 \rightarrow X$ be a non-free immersion. For the splitting $\mu^*T_X \simeq \bigoplus_{i=1}^{N-r} \mathcal{O}_{\mathbb{P}^1}(a_i)$ with $a_i \in \mathbb{Z}$, it follows $a_{i_0} \leq -1$ for some i_0 . Taking a double cover $\iota_2 : \mathbb{P}^1 \rightarrow \mathbb{P}^1$, since the splitting of $(\iota_2 \circ \mu)^*T_X$ contains $\mathcal{O}_{\mathbb{P}^1}(2a_{i_0})$, we have $H^1(\mathbb{P}^1, (\iota_2 \circ \mu)^*T_X) \neq 0$, which contradicts that X is convex. Hence $\deg(X) \leq 2$ must hold. \square

REFERENCES

- [1] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris, *Geometry of Algebraic Curves, Volume I*. Springer 1985.
- [2] W. Barth and A. Van de Ven. Fano varieties of lines on hypersurfaces. *Arch. Math. (Basel)* **31** (1978/79), 96–104.
- [3] F. Campana and T. Peternell, Projective manifolds whose tangent bundles are numerically effective, *Math. Ann.* **289** (1991), 169–187.
- [4] F. Campana, T. Peternell, 4-folds with numerically effective tangent bundles and second Betti numbers greater than one, *Manuscripta Math.* **79** (1993), no. 3-4, 225–238.
- [5] O. Debarre and L. Manivel. Sur la variété des espaces linéaires contenus dans une intersection complète. *Math. Ann.* **312** (1998), 549–574.
- [6] W. Fulton and J. Hansen, A connectedness theorem for projective varieties, with applications to intersections and singularities of mappings, *Ann. Math.* **110** (1979) 159–166.
- [7] K. Furukawa, Rational curves on hypersurfaces, *J. reine angew. Math.* **665** (2012), 157–188.
- [8] J. M. Hwang, Rigidity of rational homogeneous spaces, *Proceedings of ICM. 2006 Madrid, vol- ume II*, European Mathematical Society, 2006, 613–626.
- [9] J. Kollar, Rational curves on algebraic varieties, *Ergeb. Math. Grenzgeb. (3)* **32**, Springer-Verlag, Berlin 1996.

- [10] N. Mok, On Fano manifolds with nef tangent bundles admitting 1-dimensional varieties of minimal rational tangents, *Trans. Amer. Math. Soc.* **354** (2002), no. 7, 2639-2658.
- [11] R. Pandharipande, Convex rationally connected varieties, *Proc. Amer. Math. Soc.* **141** (2013), 1539-1543.
- [12] K. Watanabe, Fano 5-folds with nef tangent bundles and Picard numbers greater than one, *Math. Z.* **276** (2014), 39–49.

E-mail address: `katu@toki.waseda.jp`

DEPARTMENT OF MATHEMATICS, SCHOOL OF FUNDAMENTAL SCIENCE AND ENGINEERING, WASEDA UNIVERSITY, OHKUBO 3-4-1, SHINJUKU, TOKYO, 169-8555, JAPAN