

Energy-Dependent Light Quenching in CaWO_4 Crystals at mK Temperatures

R. Strauss^{1,2,a}, G. Angloher², A. Bento³, C. Bucci⁴, L. Canonica⁴, W. Carli⁹, C. Ciemniak¹, A. Erb^{1,5}, F.v. Feilitzsch¹, P. Gorla⁴, A. Gütlein¹, D. Hauff², D. Hellgartner¹, J. Jochum⁶, H. Kraus⁷, J.-C. Lanfranchi¹, J. Loebell⁶, A. Münster¹, F. Petricca², W. Potzel¹, F. Pröbst², F. Reindl², S. Roth¹, K. Rottler⁶, C. Sailer⁶, K. Schäffner⁴, J. Schieck⁸, S. Scholl⁶, S. Schönert¹, W. Seidel², M.v. Sivers¹, L. Stodolsky², C. Strandhagen⁶, A. Tanzke², M. Uffinger⁶, A. Ulrich¹, I. Usherov⁶, S. Wawoczny¹, M. Willers¹, M. Wüstrich², A. Zöller¹

¹Physik-Department, Technische Universität München, D-85748 Garching, Germany

²Max-Planck-Institut für Physik, D-80805 München, Germany

³CIUC, Departamento de Física, Universidade de Coimbra, P3004 516 Coimbra, Portugal

⁴INFN, Laboratori Nazionali del Gran Sasso, I-67010 Assergi, Italy

⁵Walther-Meißner-Institut für Tieftemperaturforschung, D-85748 Garching, Germany

⁶Physikalisches Institut, Eberhard-Karls-Universität Tübingen, D-72076 Tübingen, Germany

⁷Department of Physics, University of Oxford, Oxford OX1 3RH, United Kingdom

⁸Institut für Hochenergiephysik der Österreichischen Akademie der Wissenschaften, A-1050 Wien, Austria

⁹Maier-Leibnitz-Laboratorium, Ludwig-Maximilians-Universität München, D-85748 Garching, Germany

Received: date / Accepted: date

Abstract Scintillating CaWO_4 single crystals are a promising multi-element target for rare-event searches and are currently used in the direct Dark Matter experiment CRESST (Cryogenic Rare Event Search with Superconducting Thermometers). The relative light output of different particle interactions in CaWO_4 is quantified by Quenching Factors (QFs). These are essential for an active background discrimination and the identification of a possible signal induced by weakly interacting massive particles (WIMPs). We present the first precise measurements of the QFs of O, Ca and W at mK temperatures by irradiating a cryogenic detector with a fast neutron beam. A clear energy dependence of the QF of O and, less pronounced, of Ca was observed for the first time. Furthermore, in CRESST neutron-calibration data a variation of the QFs among different CaWO_4 single crystals was found. For typical CRESST detectors the QFs in the region-of-interest (10-40 keV) are $QF_O^{ROI} = (11.2 \pm 0.5)\%$, $QF_{Ca}^{ROI} = (5.94 \pm 0.49)\%$ and $QF_W^{ROI} = (1.72 \pm 0.21)\%$. The latest CRESST data (run32) is reanalyzed using these fundamentally new results on light quenching in CaWO_4 having moderate influence on the WIMP analysis. Their relevance for future CRESST runs and for the clarification of pre-

viously published results of direct Dark Matter experiments is emphasized.

Keywords Dark Matter · Scintillators · CaWO_4 · Cryogenic detectors · Neutron scattering

1 Introduction

Rare-event searches for Dark Matter (DM) in the form of weakly interacting massive particles (WIMPs) [1, 2] have reached impressive sensitivities during the last decade [3]. Well motivated WIMP candidates with masses m_χ between a few GeV/c^2 and a few TeV/c^2 might be detectable via nuclear recoils of few keV in terrestrial experiments [4]. While the DAMA/LIBRA [5], and recently the CoGeNT [6], CRESST [7], and the CDMS(Si) [8] experiments observed excess signals that might be interpreted as induced by DM particles with $m_\chi \sim 10 \text{ GeV}/c^2$ at WIMP-nucleon cross-sections of $\sim 10^{-4} \text{ pb}$, this scenario is ruled out by the LUX [9] and XENON100 [10] experiments, and almost excluded by the CDMS(Ge) [11,12], the EDELWEISS [13,14] and the SuperCDMS [15] experiments. It is strongly disfavoured by accelerator constraints [16,17] and in mild tension with an extended analysis [18] of published CRESST data [19].

^ae-mail: strauss@mpp.mpg.de

2 Dark Matter search with CRESST

The CRESST experiment [7] employs scintillating CaWO_4 crystals [20,21] as a multi-element target material. The key feature of a CRESST detector module is the simultaneous measurement of the recoil energy E_r by a particle interaction in the crystal (operated as cryogenic calorimeter at mK temperatures [22]) and the corresponding scintillation-light energy E_l by a separate cryogenic light absorber. Since the relative light yield $LY = E_l/E_r$ is reduced for highly ionizing particles compared to electron recoils (commonly referred to as quenching) nuclear-recoil events can be discriminated from e^-/γ and α backgrounds. The phenomenological Birks model [23] predicts this quenching effect to be stronger the higher the mass number A of the recoiling ion, which allows to distinguish, in general, between O ($A \approx 16$), Ca ($A \approx 40$) and W ($A \approx 184$) recoils. The expected WIMP-recoil spectrum - assuming coherent scattering - is completely dominated by W-scatters for $m_\chi \gtrsim 20 \text{ GeV}/c^2$. However, the light targets O and Ca make CRESST detectors particularly sensitive to low-mass WIMPs of $1 \text{ GeV} \lesssim m_\chi \lesssim 20 \text{ GeV}$. Furthermore, the knowledge of the recoil composition of O, Ca and W allows a test of the assumed A^2 -dependence of the spin-independent WIMP-nucleon cross-section [2]. In addition, background neutrons, which are mainly visible as O-scatters (from kinematics [24]), can be discriminated statistically.

3 Quenching Factors (QFs)

The mean LY of e^-/γ events (LY_γ) is energy dependent and phenomenologically parametrized as $LY_\gamma(E_r) = (p_0 + p_1 E_r)(1 - p_2 \exp(-E_r/p_3))$ [25]. By convention, $LY_\gamma(122 \text{ keV})$ is normalized to unity. The parameters p_0 , p_1 , p_2 and p_3 are derived from a maximum likelihood (ML) fit for every detector module individually. For the module used in this work the fit yields: $p_0 = 1.07$, $p_1 = -1.40 \cdot 10^{-5} \text{ keV}^{-1}$, $p_2 = 6.94 \cdot 10^{-2}$ and $p_3 = 147 \text{ keV}$ (errors are negligible for the following analysis). The exponential decrease towards lower recoil energies (quantified by p_2 and p_3) accounts for the scintillator non-proportionality [26]. The Quenching Factor (QF) of a nucleus x - in general energy dependent - is defined as $QF_x(E_r) = LY_x(E_r)/LY_{\gamma,norm}$ where LY_x is the mean LY of a nuclear recoil x and $LY_{\gamma,norm}$ is a detector-specific normalization factor which corresponds to the LY of e^-/γ events. By convention, $LY_{\gamma,norm} = LY_\gamma(E_r)/(1 - p_2 \exp(-E_r/p_3)) \approx p_0$ is used for the analysis since the scintillator non-proportionality is not observed for nuclear recoils and $p_1 \ll 1$. For typical CRESST detector modules, the uncertainties in en-

ergy and LY are well described by gaussians [7] consistent with photon-counting statistics in the energy range considered in this work.

Since the resolution of light-detectors operated in the CRESST setup at present is not sufficient to disentangle O, Ca and W recoils unambiguously, dedicated experiments to measure the QFs of CaWO_4 are necessary. Earlier attempts yield inconclusive results, in particular for the value of QF_W [27,28,29].

4 The neutron-scattering facility

4.1 Experimental setup

At the accelerator of the Maier-Leibnitz-Laboratorium (MLL) in Garching a dedicated neutron-scattering facility for precision measurements of QFs at mK temperatures was set up (see Fig. 1). A pulsed ^{11}B beam of $\sim 65 \text{ MeV}$ in bunches of 2-3 ns (FWHM) produces monoenergetic neutrons of $\sim 11 \text{ MeV}$ via the nuclear reaction $p(^{11}\text{B},n)^{11}\text{C}$ in a pressurized H_2 target [30]. These neutrons are irradiated onto a CRESST-like detector module consisting of a $\sim 10 \text{ g}$ cylindrical CaWO_4 single crystal (20 mm in diameter, 5 mm in height) and a separated Si light absorber (20 mm in diameter, $500 \mu\text{m}$ thick) [31]. Both are operated as cryogenic detectors in a dilution refrigerator at $\sim 20 \text{ mK}$ [32]. Undergoing elastic (single) nuclear scattering in CaWO_4 the neutrons are tagged at a fixed scattering angle θ in an array of 40 liquid-scintillator (EJ301) detectors which allow fast timing ($\sim 2 \text{ ns}$) and n/γ discrimination.

4.2 Working principle

Depending upon which of the three nuclei is hit a distinct amount of energy is deposited by the neutron in the crystal. Triple-coincidences between (1) a ^{11}B pulse on the H_2 target, (2) a neutron pulse in a liquid-scintillator detector and (3) a nuclear-recoil event in the CaWO_4 crystal can be extracted from the data set. A neutron time-of-flight (TOF) measurement between neutron production and detection combined with a precise phononic measurement of the energy deposition in the crystal (resolution $\sim 1 \text{ keV}$ (FWHM)) allows an identification of the recoiling nucleus. To derive the individual QF the corresponding scintillation-light output is measured simultaneously by the light detector. Since the onset uncertainty of cryodetector pulses is large ($\sim 5 \mu\text{s}$) compared to typical neutron TOFs ($\sim 50 \text{ ns}$) an offline coincidence analysis has to be performed [25].




Fig. 1 Schematic experimental setup of the neutron-scattering facility. Neutrons produced by the accelerator are scattered off a CRESST-like detector module (operated at 20 mK) and tagged in liquid-scintillator neutron detectors at a fixed scattering angle Θ .




Fig. 2 Histogram of the time difference Δt between neutron events with the correct TOF and the closest W-recoil in the CaWO_4 crystal ($E_r = 100 \pm 20$ keV). A fit to the distribution (solid black line) including a constant for the accidental background (shaded area) and a gaussian for the triple-coincidences on W (dashed red line) is shown. 158 W-scatters are identified with a signal-to-background ratio of $\sim 7 : 1$.

4.3 Measurements and results for QF_W

The experiment was optimized for the measurement of QF_W [25,33]. To enhance the number of W-scatters a scattering angle of $\Theta = 80^\circ$ was chosen due to scattering kinematics [27]. For this specific angle, the expected recoil energy of triple-coincident events is ~ 100 keV for W, ~ 450 keV for Ca, and ~ 1.1 MeV for O. In ~ 3 weeks of beam time a total of $\sim 10^8$ cryodetector pulses were recorded. Fig. 2 shows the time difference Δt between neutron events with the correct TOF identified in one of the liquid-scintillator detectors and the closest W-recoil (in time) in the CaWO_4 crystal ($E_r = 100 \pm 20$ keV). A gaussian peak of triple-coincidences on W (dashed red line) at $\Delta t \approx 0.016$ ms and a width of $\sigma_t \approx 4.8 \mu\text{s}$ (onset resolution of the cryodetector) is observed above a background due to accidental coincidences uniformly distributed in time (shaded area). Within the 2σ -bounds of the peak 158 W-scatters are identified with a signal-to-background (S/B) ratio of $\sim 7 : 1$.

Fig. 3 shows the LY distribution of these events in a histogram (black dots). The mean LY of the extracted




Fig. 3 LY histogram of the 158 events identified as triple-coincidences on W. A fit to the distribution (solid black line) is shown which includes a gaussian (dashed red line) accounting for W-scatters and the background-pdf (shaded area) describing accidental coincidences. The simultaneous ML fit including the timing distribution yields $QF_W = (1.96 \pm 0.22)\%$.

W-scatters is found at a lower value compared to the mean LY of all nuclear recoils, i.e., the (overlapping) contributions of O, Ca and W if no coincidence measurement is involved. The accidental coincidences have a LY-distribution equal to that which is modelled by a probability-density function (background-pdf) [25]. A simultaneous maximum-likelihood (ML) fit is performed including (1) the timing distribution which fixes the S/B ratio and the number of identified W-events, and (2) the LY distribution described by a gaussian (W-events) and the background-pdf. The final results are $LY_W = 0.0208 \pm 0.0024$ and $QF_W = (1.96 \pm 0.22)\%$, correspondingly (errors are dominated by statistics). Fig. 3 shows the fit to the LY distribution by the gaussian (dashed red line) and the background-pdf (shaded area).

5 Energy-dependent QF analysis

5.1 Principle of analysis

For the measurement of QF_{Ca} and QF_O no coincidence signals are necessary, instead, an analysis of the nuclear-recoil data alone is sufficient. Also for this analysis the neutron data obtained at the scattering facility were used. Commonly CRESST data is displayed in the energy-LY plane [7] giving rise to nearly horizontal bands which correspond to different types of particle interactions ($LY \approx 1$ for electron and $LY \lesssim 0.2$ for nuclear recoils). The nuclear-recoil bands of the data recorded during ~ 1 week of beam time ($\sim 5 \cdot 10^5$ pulses) are shown in Fig. 4 (2-d histogram). From kinematics using ~ 11 MeV neutrons as probes the O-recoil band extends up to ~ 2.4 MeV while the Ca- and W-bands extend up to ~ 1.05 MeV and ~ 240 keV, respectively [30]. De-




Fig. 4 Histogram of neutron-induced nuclear-recoil events plotted in the LY-energy plane. The corresponding 1σ acceptance bounds (full red lines) of O, Ca, and W as derived from the correlated ML fit (see text) are indicated.




Fig. 5 LY histograms of energy slices (20 keV in width) at 350 keV (top) and 40 keV (bottom) fitted by gaussians.

spite the strong overlap of the 3 nuclear-recoil bands the contributions of O and Ca fitted by two gaussians can be disentangled at $E_r \gtrsim 350$ keV (see Fig. 5 top) due to high statistics and a good light-detector resolution.

5.2 Results and discussion

In Fig. 6 the results for QF_O and QF_{Ca} (red error bars) derived by these independent one-dimensional (1-dim) fits are shown for selected recoil-energy slices of 20 keV in width. All parameters in the fit are left free except for the LY-resolutions which are fixed by a ML fit of the electron-recoil band [25]. While QF_O clearly rises towards lower recoil energies, this effect is less pronounced for QF_{Ca} .

Below ~ 350 keV, due to the strong overlap of the nuclear recoil bands, this simple approach fails. Instead, a correlated ML fit was performed based on the following assumptions: (1) for the mean LY of O- and Ca-scatters the phenomenological parametrization $LY_x(E_r) = LY_x^\infty (1 + f_x \cdot \exp(-E_r/\lambda_x))$ is proposed with the free parameters LY_x^∞ (LY at $E_r = \infty$), f_x (fraction of energy-dependent component) and λ_x (ex-

Table 1 Results for the free parameters LY_x^∞ , f_x and λ_x of the ML analysis. The statistical errors are given at 1σ C.L.

	LY_x^∞	f_x	λ_x
O	0.07908 ± 0.00002	0.7088 ± 0.0008	567.1 ± 0.9
Ca	0.05949 ± 0.00078	0.1887 ± 0.0022	801.3 ± 18.8

ponential decay with energy), and (2) the mean LY of W-scatters is approximated to be constant in the relevant energy range (up to ~ 240 keV) at the value precisely measured with the triple-coincidence technique ($LY_W = 0.0208 \pm 0.0024$ which corresponds to $QF_W = (1.96 \pm 0.22)\%$). These assumptions are supported by the result of the 1-dim fits (see Fig. 6), by Birks' model [23], and by a recent work [34] which predict the strength of the energy-dependence to decrease with A . The nuclear-recoil bands are cut into energy intervals of 10 keV (20 keV to 1 MeV), of 20 keV (1 MeV to 1.4 MeV) and 50 keV (above 1.4 MeV) and fitted with up to 3 gaussians depending on the recoil energy (e.g., shown in Fig. 5 bottom for $E_r = 40$ keV). Except for the assumptions mentioned above and the LY-resolution all parameters are left free in the fit. The fit converges over the entire energy range (20-1800 keV). In Table 1 the results for LY_x^∞ , f_x and λ_x are presented which correspond, e.g. at 40 keV, to $QF_O = (12.6 \pm 0.5)\%$, $QF_{Ca} = (6.73 \pm 0.43)\%$ at 1σ C.L. Errors are dominated by systematics including different choices of the LY parametrization. The final results for QF_O , QF_{Ca} and QF_W are presented in Fig. 6 and are found to be in perfect agreement with the outcome of the 1-dim fits (red error bars). Fig. 4 shows the 1σ acceptance bounds (full red lines) of O, Ca and W recoils as obtained in the correlated ML fit.

These are the first experimental results which clearly show a rise of QF_O of $\sim 28\%$ towards the ROI (10-40 keV) compared to that at a recoil energy of 500 keV. For QF_{Ca} the best fit yields a rise of $\sim 6\%$, however, the energy-dependence is less significant (see Fig. 6).

In previous works, the QFs of CaWO_4 were assumed to be constant over the entire energy range [7]. A statistical analysis shows that this simple model is clearly disfavoured. Employing a likelihood-ratio test in combination with Monte-Carlo simulations gives a p-value of $p < 10^{-5}$ for the data presented here to be consistent with constant QFs. Furthermore, the derived energy spectra of the individual recoiling nuclei agree with the expectation from incident 11 MeV neutrons while the constant QF approach provides non-physical results.




Fig. 6 Results of the correlated ML analysis for QF_O , QF_{Ca} and QF_W (solid lines). The shaded areas indicate the 1σ and 2σ bounds. For the first time a clear energy dependence of QF_O and QF_{Ca} is observed. These results are in agreement with that of the 1-dim fits of discrete energy intervals (see text) shown as red error bars. QF_W is fixed (in the correlated fit) at the value measured by the triple-coincidence technique.

6 QFs of CRESST detectors

In the present paper, using the 8 detector modules operated in the last CRESST measurement campaign (run32) an additional aspect was investigated: the variation of the quenching behaviour among *different* CaWO_4 crystals [25]. Nuclear recoils acquired during neutron calibration campaigns of CRESST run32 are completely dominated by O-scatters at $E_r \gtrsim 150$ keV (from kinematics) [7]. Despite low statistics (a factor of ~ 100 less compared to the measurement presented here) in the available data, the mean LY of O-events can be determined by a gaussian fit with a precision of $\mathcal{O}(1\%)$ for every module. In this way, the mean QF of O between 150 and 200 keV was determined individually for the 8 detector modules (index i) operated in run32 ($\overline{QF_{O,i}^*}$) and for the reference detector operated at the neutron-scattering facility ($\overline{QF_O}$). Different values of $\overline{QF_{O,i}^*}$ are observed for the CRESST detector crystals (variation by $\sim 11\%$) and for the reference crystal ($\sim 12\%$ higher than the mean of $\overline{QF_{O,i}^*}$). This variation appears to be correlated with the crystal's optical quality. The QF - which is a relative quantity - is found to be lower if a crystal has a smaller defect density and thus a higher absolute light output, i.e., the LY of nuclear recoils is less affected by an increased defect density. This is in agreement with the prediction described in a recent work [34]. In the present paper, a simple model to account for this variation is proposed: For every detector module which is to be calibrated a scaling factor ϵ_i is introduced, $\epsilon_i = \overline{QF_{O,i}^*} / \overline{QF_O}$. Then, within this model the QFs of the nucleus x can be calculated for every module by $QF_{x,i}^*(E_r) = \epsilon_i \cdot QF_x(E_r)$ where QF_x

Table 2 QF results averaged over the ROI (10-40 keV) and adjusted by the scaling factor ϵ_i for the modules Rita and Daisy, and the mean (\emptyset) of all run32 detectors (1σ errors).

	ϵ_i	$QF_O^{\text{ROI}}[\%]$	$QF_{Ca}^{\text{ROI}}[\%]$	$QF_W^{\text{ROI}}[\%]$
Rita	0.844	10.8 ± 0.5	5.70 ± 0.44	1.65 ± 0.19
Daisy	0.939	12.0 ± 0.7	6.33 ± 0.58	1.84 ± 0.24
\emptyset	0.880	11.2 ± 0.5	5.94 ± 0.49	1.72 ± 0.21

is the value precisely measured within this work. The nuclear-recoil behaviour of CRESST modules is well described by energy-dependent QFs. In Table 2 the QFs, averaged over the ROI (10-40 keV), and the scaling factor ϵ_i are listed for two selected detector modules (Rita and Daisy, with the lowest and highest absolute light output, respectively) and the mean of all 8 detector modules of run32 (\emptyset), $QF_O^{\text{ROI}} = (11.2 \pm 0.5)\%$, $QF_{Ca}^{\text{ROI}} = (5.94 \pm 0.49)\%$ and $QF_W^{\text{ROI}} = (1.72 \pm 0.21)\%$.

7 Re-analysis of latest CRESST results

We now turn to the effect of energy-dependent quenching since constant QFs as assumed in earlier CRESST publications do not sufficiently describe the behaviour of the nuclear-recoil bands. The value of QF_O in the ROI was underestimated by $\sim 8\%$ while the room temperature measurements overestimated the values of QF_{Ca} and QF_W by $\sim 7\%$ and $\sim 130\%$, respectively [7]. Therefore, the parameter space of accepted nuclear recoils is larger than assumed in earlier publications (by $\sim 46\%$) requiring a re-analysis of the published CRESST data.

During the latest measuring campaign (run32) a statistically significant signal (4.2σ) above known backgrounds was observed. If interpreted as induced by DM particles two WIMP solutions were found [7], e.g. at a mass of $m_\chi = 11.6 \text{ GeV}/c^2$ with a WIMP-nucleon cross section of $\sigma_\chi = 3.7 \cdot 10^{-5} \text{ pb}$. The dedicated ML analysis was repeated using the new QF values (\emptyset in Table 2) yielding $m_\chi = 12.0 \text{ GeV}/c^2$ and $\sigma_\chi = 3.2 \cdot 10^{-5} \text{ pb}$ at 3.9σ . Beside this moderate change of the WIMP parameters also the background composition (e^- , γ , neutrons, α 's and ^{206}Pb) is influenced. This is mainly due to the significantly lower value of QF_W which increases the leakage of ^{206}Pb recoils into the ROI (by $\sim 18\%$). The other WIMP solution is influenced similarly: m_χ changes from 25.3 to $25.5 \text{ GeV}/c^2$, σ_χ from $1.6 \cdot 10^{-6}$ to $1.5 \cdot 10^{-6} \text{ pb}$ and the significance drops slightly from 4.7 to 4.3σ .

8 Summary and Outlook

In conclusion, the first precise measurement of Q_{FW} at mK temperatures and under conditions comparable to that of the CRESST experiment was obtained at the neutron-scattering facility in Garching by an extensive triple-coincidence technique. Furthermore, the QFs of O and Ca were precisely determined by a dedicated maximum-likelihood analysis over the entire energy range (~ 20 – 1800 keV). The observed energy dependence of the QFs, which is more pronounced for lighter nuclei, has significant influence on the determination of the ROI for DM search. Analysing CRESST neutron-calibration data a variation of the QFs between different CaWO_4 crystals was observed which is related to the optical quality. By the simple model proposed above the measured QFs can be adapted to every individual crystal. The updated values of the QFs are highly relevant to disentangle the recoil composition (O, Ca and W) of a possible DM signal and, therefore, to determine the WIMP parameters. Since the separation between the O and W recoil bands is higher by $\sim 46\%$ compared to earlier assumptions, background neutrons which are mainly visible as O-scatters [24] can be discriminated more efficiently from possible WIMP-induced events. A reanalysis of the run32 data shows a moderate influence of the new QF values on the WIMP parameters.

The results obtained here are of importance for the current CRESST run (run33) and upcoming measuring campaigns. Providing a highly improved background level run33 has the potential to clarify the origin of the observed excess signal and to set competitive limits for the spin-independent WIMP-nucleon cross section in the near future.

For the planned multi-material DM experiment EURECA (European Underground Rare Event Calorimeter Array) [35] the neutron-scattering facility will be an important tool to investigate the light quenching of alternative target materials in the future.

Acknowledgements This research was supported by the DFG cluster of excellence: Origin and Structure of the Universe, the DFG Transregio 27: Neutrinos and Beyond, the Helmholtz Alliance for Astroparticle Physics, the Maier Leibnitz Laboratorium (Garching) and by the BMBF: Project 05A11WOC EURECA-XENON.

References

1. G. Bertone, D. Hooper, J. Silk, *Phys.Rept.* **405**, 279 (2005). DOI 10.1016/j.physrep.2004.08.031
2. G. Jungman, M. Kamionkowski, K. Griest, *Phys.Rept.* **267**, 195 (1996). DOI 10.1016/0370-1573(95)00058-5
3. P. Cushman, et al., (2013). URL <http://arxiv.org/abs/arXiv:1310.8327>
4. J. Lewin, P. Smith, *Astropart.Phys.* **6**, 87 (1996). DOI 10.1016/S0927-6505(96)00047-3
5. R. Bernabei, et al., *Eur.Phys.J.* **C67**, 39 (2010). DOI 10.1140/epjc/s10052-010-1303-9
6. C. Aalseth, et al., *Phys.Rev.Lett.* **106**, 131301 (2011). DOI 10.1103/PhysRevLett.106.131301
7. G. Angloher, et al., *Eur.Phys.J.* **C72**(4), 1 (2012). URL <http://dx.doi.org/10.1140/epjc/s10052-012-1971-8>
8. R. Agnese, et al., *Phys. Rev. Lett.* **111**, 251301 (2013). DOI 10.1103/PhysRevLett.111.251301
9. D. Akerib, et al., *Phys.Rev.Lett.* **112**, 091303 (2014). DOI 10.1103/PhysRevLett.112.091303
10. E. Aprile, et al., *Phys.Rev.Lett.* **109**, 181301 (2012). DOI 10.1103/PhysRevLett.109.181301
11. Z. Ahmed, et al., *Science* **327**, 1619 (2010). DOI 10.1126/science.1186112
12. Z. Ahmed, et al., *Phys.Rev.Lett.* **106**, 131302 (2011). DOI 10.1103/PhysRevLett.106.131302
13. E. Armengaud, et al., *Phys.Lett.* **B702**, 329 (2011). DOI 10.1016/j.physletb.2011.07.034
14. E. Armengaud, et al., *Phys.Rev.* **D86**, 051701 (2012). DOI 10.1103/PhysRevD.86.051701
15. R. Agnese, et al., *Phys.Rev.Lett.* **112**, 241302 (2014). DOI 10.1103/PhysRevLett.112.241302
16. G. Aad, et al., *JHEP* **1304**, 075 (2013). DOI 10.1007/JHEP04(2013)075
17. S. Chatrchyan, et al., *JHEP* **1209**, 094 (2012). DOI 10.1007/JHEP09(2012)094
18. A. Brown, S. Henry, H. Kraus, C. McCabe, *Phys. Rev. D* **85**, 021301 (2012). DOI 10.1103/PhysRevD.85.021301
19. G. Angloher, et al., *Astropart.Phys.* **31**(4), 270 (2009)
20. T.A. Edison, *Nature* **53**, 470 (1896)
21. V.B. Mikhailik, H. Kraus, S. Henry, A.J.B. Tolhurst, *Phys. Rev. B* **75**, 184308 (2007). DOI 10.1103/PhysRevB.75.184308. URL <http://link.aps.org/doi/10.1103/PhysRevB.75.184308>
22. F. Pröbst, et al., *J. Low Temp. Phys.* **100**(1-2), 69 (1995). DOI 10.1007/BF00753837. URL <http://dx.doi.org/10.1007/BF00753837>
23. J. Birks, *The Theory and Practice of Scintillation Counting* (Pergamon Press, 1964). URL <http://books.google.com/books?id=OR1RAAAAMAAJ>
24. S. Scholl, J. Jochum, *J. Phys.: Conference Series* **375**(1), 012020 (2012). URL <http://stacks.iop.org/1742-6596/375/i=1/a=012020>
25. R. Strauss, Ph.D. thesis, TU München (2013). URL <http://mediatum.ub.tum.de/node?id=1166886>
26. R. Lang, et al., (2009). URL <http://arxiv.org/abs/0910.4414>
27. T. Jagemann, et al., *Astropart.Phys.* **26**, 269 (2006). DOI 10.1016/j.astropartphys.2006.06.010
28. J. Ninkovic, et al., *Nucl.Instrum.Meth.* **A564**, 567 (2006). DOI 10.1016/j.nima.2006.04.039
29. I. Bavykina, et al., *Astropart.Phys.* **28**, 489 (2007). DOI 10.1016/j.astropartphys.2007.09.006
30. T. Jagemann, J. Jochum, F. Feilitzsch, *Nucl.Instrum.Meth.* **A551**(2-3), 245 (2005)
31. R. Strauss, et al., *J. Low Temp. Phys.* **167**(5-6), 1063 (2012). DOI 10.1007/s10909-012-0536-4. URL <http://dx.doi.org/10.1007/s10909-012-0536-4>
32. J.C. Lanfranchi, et al., *Opt. Mat.* **31**(10), 1405 (2009)
33. R. Strauss, et al., *J. Low Temp. Phys.* pp. 1–6 (2014). DOI 10.1007/s10909-013-1075-3. URL <http://dx.doi.org/10.1007/s10909-013-1075-3>

-
34. S. Roth, Ph.D. thesis, TU München (2013). URL <https://mediatum.ub.tum.de/node?id=1172019>
 35. G. Angloher, et al., Physics of the Dark Universe **3**(0), 41 (2014). DOI <http://dx.doi.org/10.1016/j.dark.2014.03.004>