

Expectations for the Higgs boson identification at the ILC

E. Boos¹, V. Bunichev¹, M. Dubinin^{1,2}, Y. Kurihara²

¹*Skobeltsyn Institute of Nuclear Physics, Moscow State University
119991, Moscow, Russia*

²*High Energy Accelerators Research Organization (KEK), Tsukuba, 305-0801
Ibaraki, Japan*

Abstract

Deviations from the standard Higgs sector generated by some new physics at an energy scale Λ could be described by an effective $SU(3)_c \times SU(2)_L \times U(1)$ invariant non-renormalizable Lagrangian terms of dimension six. A systematic study of various Higgs boson production channels ($\gamma\gamma$, ZZ , WW , $b\bar{b}$, $\tau\bar{\tau}$) at the International Linear Collider (ILC) in the SM extension by effective operators is carried out. Statistical methods are used to establish a degree of consistency for the standard Higgs sector with the forthcoming data, using the expected ILC accuracies of the Higgs boson production channels. Global fits in the two-parametric anomalous coupling space indicating to possible deviations from the standard Higgs-fermion and Higgs-gauge boson couplings are performed.

High precision studies of the electroweak interactions at LEP, SLC and Tevatron [1] with the following discovery of a Higgs-like signal at the LHC [2] and the evidence of a signal at the Tevatron [3], confirmed the Standard Model (SM) scheme in the form of $SU(2)_L \times U(1)_Y$ gauge theory with symmetry breaking generated by the Lagrangian which includes one $SU(2)$ doublet of scalar fields. No belief that this simplest possibility is correct naturally results in a desire to analyse theoretically acceptable extensions of the SM Higgs sector and compare their phenomenological consequences with mass, couplings and quantum numbers of the Higgs boson measured precisely in the major production channels at the next-generation e^+e^- collider.

”Minimal” extensions of the SM by the dimension-five [4] and the dimension-six [5, 6] effective operators, which lead to the rescaling of Higgs couplings, as well as more complicated modifications of Higgs couplings in the MSSM and nonminimal supersymmetric models in the decoupling limit [7] commonly demonstrate insignificant (of the order of a few percent) departures of the non-standard couplings from their SM limit in the most acceptable phenomenological scenarios. Precise measurements are required to study the couplings. Although ATLAS and CMS experiments are sensitive to almost all couplings (except invisible modes and $H \rightarrow c\bar{c}$) which have been analysed recently using the LHC data [8], the best sensitivity of individual production channels $g(H \rightarrow P\bar{P})/g(H \rightarrow P\bar{P})_{SM} - 1$ ($P = \gamma, W, Z, \tau, b$) estimated in a model independent way [9] using the *production* \times *decay* approximation is from 5% to 20% (these numbers are 1σ intervals at the energy 14 TeV with the accumulated luminosity 300 fb^{-1} , see [10]). Such accuracy will be undoubtedly sufficient to confirm an agreement of basic properties of the Higgs-like particle (mass, spin and parity) with the properties of the SM Higgs boson, but will be not sufficient to test the deviations at the percent level, inherent for the MSSM in the decoupling limit or models with composite Higgs boson, where the compositeness scale is of the order of 1 TeV.

Clean environment and better signal-to-background ratios of the ILC allows to measure more precisely the couplings of the Higgs boson, which can be observed in all modes including the two-jet decay modes and the invisible modes. Only two production channels are relevant, Higgsstrahlung $e^+e^- \rightarrow ZH$ and vector boson fusion (VBF) $e^+e^- \rightarrow \nu_e\bar{\nu}_e H$. Although it will be not possible to measure directly the total width of the scalar which is about 4 MeV, it can be indirectly determined separating the events with the Higgs recoiling from the Z in the Higgsstrahlung process and measuring directly the branching fractions. Measurement accuracies expected from the ILC experiments (see Table 2.4 in the ILC TDR [10]), estimated from the full detector simulation studies for a realistic accumulated luminosities at $\sqrt{s} = 250 \text{ GeV}$, 500 GeV and 1 TeV are listed in Table 1. Note that the ILC TDR expected accuracies are insignificantly different from the accuracies quoted in the Snowmass Report [11]; an independent simulation of some channels, including invisible modes, can be found in [12].

| | $\Delta(\sigma \cdot Br)/(\sigma \cdot Br)$ | | | | |
|--|---|-----------------|--|-----------------|---|
| \sqrt{s} and \mathcal{L} (P_{e^-}, P_{e^+}) | 250 fb $^{-1}$ at 250 GeV (-0.8,+0.3) | | 500 fb $^{-1}$ at 500 GeV (-0.8,+0.3) | | 1 ab $^{-1}$ at 1000 GeV (-0.8,+0.2) |
| channel | ZH | $\nu\bar{\nu}H$ | ZH | $\nu\bar{\nu}H$ | $\nu\bar{\nu}H$ |
| $H \rightarrow b\bar{b}$ | 1.1% | 10.5% | 1.8% | 0.66% | 0.47% |
| $H \rightarrow c\bar{c}$ | 7.4% | - | 12.0% | 6.2% | 7.6% |
| $H \rightarrow gg$ | 9.1% | - | 14.0% | 4.1% | 3.1% |
| $H \rightarrow WW^*$ | 6.4% | - | 9.2% | 2.6% | 3.3% |
| $H \rightarrow \tau^+\tau^-$ | 4.2% | - | 5.4% | 14.0% | 3.5% |
| $H \rightarrow ZZ^*$ | 19.0% | - | 25.0% | 8.2% | 4.4% |
| $H \rightarrow \gamma\gamma$ | 29-38% | - | 29-38% | 20-26% | 7-10% |
| $H \rightarrow \mu^+\mu^-$ | 100% | - | - | - | 32% |

Table 1: Expected accuracies for the Higgs boson production channels at the ILC from [10].

In this letter we are using an extension of the SM by the dimension-six effective operators [13] which has been developed for identification of the Higgs-like particle at the LHC. A set of the dimension-six operators in the Buchmueller-Wyler basis [5] modified by the subtraction of v.e.v.: $\Phi^\dagger\Phi \rightarrow \Phi^\dagger\Phi - v^2/2$ (see [14]) to avoid undesirable mixing in the gauge field kinetic terms, which is strongly constrained by precision electroweak data, can be reduced to a restricted set of five fermion-Higgs and vector boson-Higgs operators $O_{t\Phi}$, $O_{b\Phi}$, $O_{\tau\Phi}$ and $O_\Phi^{(1)}$, $O_{\Phi G}$, disposing the tensor structure of interaction vertices identical to the SM and dependent only on the two anomalous couplings c_V (rescales vector boson-Higgs vertices) and c_F (rescales fermion-antifermion-Higgs vertices). The anomalous couplings C_n, C_{mn} in front of the dimension-six operators O_n, O_{mn} are conformably redefined [13], for example

$$c_F = 1 + C_{t\Phi} \cdot \frac{v^2}{\Lambda^2}, \quad c_V = 1 + \frac{v^2}{2\Lambda^2} \cdot C_\Phi^{(1)}, \quad c_G = c_F + \frac{6\pi}{\alpha_s} \cdot C_{\Phi G} \cdot \frac{v^2}{\Lambda^2}, \quad \dots$$

($v = 246$ GeV, Λ is the energy scale of new physics). The last equation here means that in the approach under consideration the one-loop $H \rightarrow \gamma\gamma$ and $H \rightarrow gg$ vertices are "resolved" in the sense that the anomalous parameters c_V and c_F are included in the one-loop effective c_G and c_γ which parametrise $H \rightarrow gg$ and $H \rightarrow \gamma\gamma$ vertices. Such parametrization is different from phenomenological parametrization used by ATLAS and CMS collaborations [15], where independent anomalous couplings and total widths are simultaneously used. Statistical methods for the signal strength equal to one and the signal strength error defined by the abovementioned ILC TDR signal simulation studies are used to evaluate combined χ^2 fits in the (c_V, c_F) anomalous coupling plane. Details of the procedure for exclusion contours reconstruction in the (c_V, c_F) plane are described in [13] (see also [16]). A combination of simultane-

ous fits for all possible Higgs boson production channels is presented in Fig.1 and Fig.3, where the exclusion contours for $\Delta\chi^2 = 2.10, 4.61$ and 9.21 are reconstructed at the energies 250, 500 and 1000 GeV. Small signal strength error in the channels $e^+e^- \rightarrow ZH, \nu\bar{\nu}H$ with the following $H \rightarrow b\bar{b}$ or $H \rightarrow WW^*$ results in the most significant influence of the modes $H \rightarrow b\bar{b}$ and $H \rightarrow WW^*$ on the combined χ^2 fit. The shape of exclusion contours is most sensitive to the interplay of these two modes, which have opposite behaviour of the cross section dependence $\sigma(c_V, c_F)$ on the anomalous couplings. In order to demonstrate the sensitivity, in Fig.1a,b we separately plot the overall fit and the combined fit for all channels excluding $H \rightarrow WW^*$ (upper row of plots). Naive expectation that $H \rightarrow WW^*$ channel should be more sensitive to c_V , than to c_F , is delusive since the branching ratios are generally speaking not proportional to partial widths because of an untrivial dependence of the total width on the anomalous couplings. Despite a small cross section and low expected precision, the channel $H \rightarrow \gamma\gamma$ is very important to exclude the region in the vicinity of $(c_V, c_F) = (1, -1)$ which is acceptable for LHC global fits (see [13]). In the absence of $H \rightarrow \gamma\gamma$ data this region immediately appears for the ILC combined fit, see Fig.1, lower row of plots, due to an asymmetric dependence of $\chi^2(c_V, c_F)$ on the fermion anomalous coupling c_F , illustrated by the three-dimensional χ^2 in Fig.2a. Asymmetric behaviour with respect to c_F is a consequence of asymmetric behavior of χ^2 for the $H \rightarrow \gamma\gamma$ channel taken separately, see Fig.2b. At the energies $\sqrt{s} = 500$ GeV and $\sqrt{s} = 1000$ GeV the role of VBF increases with the cross section growth and smaller signal strength error, as illustrated by Fig.3.

Evaluation of the exclusion contours was carried out¹ for complete gauge invariant sets of diagrams $e^+e^- \rightarrow (HZ \text{ or } \nu\bar{\nu}H) \rightarrow (f\bar{f}f\bar{f} \text{ or } f\bar{f}VV)$ with the four-fermion and the two-fermion – two vector boson (including $H \rightarrow gg$) final states. Interferences play a role for LHC fits [13]. For the ILC case, some examples also had been found; e.g. in connection with LEP2 analyses it had been mentioned that the four-fermion state $\nu_e\bar{\nu}_e b\bar{b}$ was formed by both $e^+e^- \rightarrow HZ$ and $e^+e^- \rightarrow \nu_e\bar{\nu}_e H$ interfering production mechanisms, see details at LEP2 and NLC energies in [17]. Untrivial interferences between signal diagrams, not available in the *production* \times *decay* approximation, were found insignificant for our fits, although they could play a role in specific reconstructions of phase space distributions beyond the infinitely small width approximation [18].

In summary, we analysed possible degree of consistency of the SM Higgs sector with the forthcoming ILC data taking into account the main mechanisms of the Higgs boson production $e^+e^- \rightarrow HZ, \nu\bar{\nu}H$ in the framework of the SM extension by dimension-six effective operators. Combined χ^2 fits at the energies $\sqrt{s} = 250, 500$ and 1000 GeV were carried out and the exclusion regions in (c_V, c_F) anomalous coupling plane were reconstructed using an expected accuracies for the cross section measurements at the ILC, based on full detector simulation studies. The channel $H \rightarrow \gamma\gamma$ is important to exclude the region of

¹A special regime of 'table calculations' (numerical operations with multidimensional tables) has been implemented in CompHEP version 4.5 [19].

anomalous coupling plane close to $(c_V, c_F)=(1, -1)$.

Acknowledgements

The work of E.B., V.B. and M.D. was partially supported by RFBR grant 12-02-93108 and NSH grant 3042.2014.2.

References

- [1] ALEPH, DELPHI, L3 and OPAL and LEP electroweak collaborations, *Electroweak measurements in electron-positron collisions at W-boson-pair energies at LEP*, Phys. Rept. **532**, 119 (2013) (arXiv:1302.3415 [hep-ex])
ALEPH, DELPHI, L3, OPAL, SLD and LEP electroweak working group, SLD electroweak group and SLD heavy flavor group, *Precision electroweak measurements on the Z resonance*, Phys. Rept. **427**, 257 (2006) (hep-ex/0509008)
D. Waters for the CDF and DO Collaborations, *Electroweak Precision Measurements and Direct Higgs Searches at the Tevatron*, in: Proc. of XX international workshop on deep-inelastic scattering and related subjects (DIS 2012), p.37 (DOI: 10.3204/DESY-PROC-2012-02/21)
- [2] G. Aad *et al.* (ATLAS Collaboration), *Observation of a new particle in the search for the SM Higgs boson with the ATLAS detector at the LHC*, Phys. Lett. **B716**, 1 (2012) (arXiv:1207.7214[hep-ex])
S. Chatrchyan *et al.* (CMS Collaboration), *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, Phys. Lett. **B716**, 30 (2012) (arXiv:1207.7235[hep-ex])
- [3] T. Aaltonen *et al.* (CDF and D0 Collaborations), *Evidence for a particle produced in association with weak bosons and decaying to a $b\bar{b}$ quark pair in Higgs boson searches at the Tevatron*, Phys. Rev. Lett. **109**, 071804 (2012) (arXiv:1207.6436 [hep-ex])
- [4] M. Dolan, C. Englert, M. Spannowsky, *New physics in LHC Higgs boson pair-production*, arXiv:1210.8166 [hep-ph], 2012
A. Azatov, R. Contino, J. Galloway, *Model-independent bounds on a light Higgs*, JHEP **1204**, 127 (2012) (arXiv:1202.3144 [hep-ph])
R. Grober, M. Muhlleitner, *Composite Higgs boson pair production at the LHC*, JHEP **1106**, 020 (2011) (arXiv:1012.1562 [hep-ph])
R. Contino, C. Grojean, M. Moretti, F. Piccinini, R. Rattazzi, *Strong double Higgs production at the LHC*, JHEP **1005**, 089 (2010) (arXiv:1002.1011 [hep-ph])
G.F. Giudice, C. Grojean, A. Pomarol, R. Rattazzi, *The strongly interacting light Higgs*, JHEP **0706**, 045 (2007)(arXiv:hep-ph/0703164)

- [5] W. Buchmuller, D. Wyler, *Effective Lagrangian analysis of new interactions and flavour conservation*, Nucl.Phys. **B268**, 621 (1986)
- [6] J. Elias-Miro, C. Grojean, R. Gupta, D. Marzocca, *Scaling and tuning of EW and Higgs observables*, arXiv:1312.2928 [hep-ph], 2013
- R. Alonso, E. Jenkins, A. Manohar, M. Trott, *Renormalization group evolution of the SM dimension-six operators*, arXiv:1312.2014 [hep-ph], 2013
- C. Chen, S. Dawson, C. Zhang, *Electroweak effective operators and Higgs physics*, arXiv:1311.3107 [hep-ph], 2013
- I. Brivio, T. Corbett, O.J.P. Eboli, M.B. Gavela, J. Gonzalez-Garcia, L. Merlo, S. Rigolin, *Disentangling a dynamical Higgs*, arXiv:1311.1823 [hep-ph], 2013
- J. Elias-Miro, J.R. Espinosa, E.Masso, A. Pomarol, *Higgs windows to new physics through $d=6$ operators*, arXiv:1308.1879 [hep-ph], 2013
- R. Contino, M. Ghezzi, C. Grojean, M. Muhlleitner, M. Spira, *Effective Lagrangian for a light Higgs-like scalar*, JHEP **1307**, 035 (2013) (arXiv:1303.3876 [hep-ph])
- T. Corbett, O.J.P. Eboli, J. Gonzalez-Fraile, M.C. Gonzalez-Garcia, *Constraining anomalous Higgs interactions*, Phys.Rev. **D86**, 075013 (2012) (arXiv:1207.1344 [hep-ph])
- B. Grzadkowski, M. Iskrzynski, M. Misiak, J. Rosiek, *Dimension-six terms in the Standard Model Lagrangian*, JHEP **1010**, 085 (2010) (arXiv:1008.4884 [hep-ph])
- K. Whisnant, J.M. Yang, B.-L. Young, X. Zhang, *Dimension six CP-conserving operators of the third family quarks and their effects on collider observables*, Phys. Rev. **D56**, 467 (1997) (hep-ph/9702305)
- K.Hagiwara, S. Ishihara, R. Szalapski, D. Zeppenfeld, *Low-energy effects of new interactions in the electroweak boson sector*, Phys. Rev. **D48**, 2182 (1993)
- [7] H. Abramowicz *et al.* (CLIC Detector and Physics Study Collaboration), *Physics at the CLIC $e+e-$ Linear Collider – Input to the Snowmass process 2013*, arXiv:1307.5288 [hep-ex], 2013
- R. Gupta, H. Rzehak, J. Wells, *How well do we need to measure the Higgs boson mass and self-coupling?*, Phys.Rev. **D88** 055024 (2013) (arXiv:1305.6397 [hep-ph])
- M. Carena, H. Haber, H. Logan, S. Mrenna, *Distinguishing a MSSM Higgs boson from the SM Higgs boson at a linear collider*, Phys.Rev. **D65**, 055005 (2002), (Erratum-ibid. D65 (2002) 099902) (hep-ph/0106116) —

- [8] CMS Physics Analysis Summaries, *Combination of standard model Higgs boson searches and measurements of the properties of the new boson with a mass near 125 GeV*, CMS-PAS-HIG-13-005, see <https://cds.cern.ch/record/1542387>
- ATLAS Collaboration, *Measurements of Higgs boson production and couplings in diboson final states with the ATLAS detector at the LHC*, arXiv:1307.1427v1 [hep-ex]
- [9] M. Peskin, *Estimation of LHC and ILC capabilities for precision Higgs boson coupling measurements*, arXiv:1312.4974 [hep-ph], 2013
- [10] *The International Linear Collider Technical Design Report - Volume 2: Physics*, ed. by H.Baer, T.Barklow, K.Fujii, Y. Gao, A.Hoang, S.Kanemura, J.List, H.Logan, A.Nomerotski, M.Perelstein, arXiv:1306.6352 [hep-ph], 2013
- [11] D. Asner et al., *ILC Higgs White Paper*, arXiv:1310.0763 [hep-ph], 2013
- [12] T. Han, Zhen Lui, J. Sayre, *Potential precision on Higgs couplings and total width at the ILC*, arXiv:1311.7155 [hep-ph], 2013
- [13] E. Boos, V. Bunichev, M. Dubinin, Y. Kurihara, *Higgs boson signal at complete tree level in the SM extension by dimension-six operators*, Phys. Rev. **D89**, 035001 (2014) (arXiv:1309.5410 [hep-ph])
- [14] G. Passarino, *NLO inspired effective Lagrangians for Higgs physics*, Nucl.Phys. **B868**, 416 (2013) (arXiv:1209.5538 [hep-ph])
- [15] A.David, A.Denner, M.Duhrssen, M.Grazzini, C.Grojean, G.Passarino, M.Schumacher, M.Spira, G.Weiglein, M.Zanetti, *LHC HXSWG interim recommendations to explore the coupling structure of a Higgs-like particle*, arXiv:1209.0040 [hep-ph], 2012
- [16] B. Dumont, S. Fichet, G. von Gersdorff, *A Bayesian view of the Higgs sector with higher dimensional operators*, JHEP **1307**, 065 (2013) (arXiv:1304.3369 [hep-ph])
- J.R. Espinosa, C. Grojean, M. Muhlleitner, M. Trott, *First glimpses at Higgs' face*, JHEP **1212**, 045 (2012) (arXiv:1207.1717 [hep-ph])
- J. R. Espinosa, M. Muhlleitner, C. Grojean, M. Trott, *Probing for invisible Higgs decays with global fits*, JHEP **1209**, 126 (2012) (arXiv:1205.6790 [hep-ph])
- J.R. Espinosa, C. Grojean, M. Muhlleitner, M. Trott, *Fingerprinting Higgs Suspects at the LHC*, JHEP **1205**, 097 (2012) (arXiv:1202.3697 [hep-ph])
- S. Kraml, B.C. Allanach, M. Mangano, H.B. Prosper, S. Sekmen, C. Balazs, A. Barr, P. Bechtle, G. Belanger, A. Belyaev et al., *Searches for new*

- physics: Les Houches recommendations for the presentation of LHC results*, Eur.Phys.J. **C72**, 1976 (2012) (arXiv:1203.2489 [hep-ph])
- A. Azatov, R. Contino, J. Galloway, *Model-independent bounds on a light Higgs*, JHEP **1204**, 127 (2012) 127, Erratum-ibid. **1304**, 140 (2013) (arXiv:1202.3415 [hep-ph])
- [17] E. Boos, M. Dubinin, L. Dudko, *Higgs boson production under the resonance threshold at LEP-2*, Int.J.Mod.Phys. **A11**, 5015 (1996) (hep-ph/9602220)
- E. Boos, M. Sachwitz, H.J. Schreiber, S. Shichanin, *The reaction $e^+e^- \rightarrow \nu\bar{\nu}b\bar{b}$ and the Higgs signal at LEP-200 and NLC*, Int.J.Mod.Phys. **A10**, 2067 (1995)
- M. Dubinin, V. Edneral, Y. Kuhhara, Y. Shimizu, *Complete tree level calculation of the process $e^+e^- \rightarrow \nu\bar{\nu}b\bar{b}$ and the Higgs signal at LEP-200 and next linear colliders*, Phys.Lett. **B329**, 379 (1994)
- [18] M.Grunewald, G.Passarino et.al., *Four fermion production in electron-positron collisions*, Reports of the working groups on precision calculations for LEP2 physics, CERN report 2000-09-A, p. 1-135 (hep-ph/0005309)
- [19] E. Boos *et al.*, *CompHEP 4.4: automatic computations from Lagrangians to events*, Nucl. Instrum. Meth. **A534**, 250 (2004) (hep-ph/0403113)
- A. Pukhov *et al.*, *CompHEP 3.3 users' manual*, hep-ph/9908288;
see also <http://theory.npi.msu.su/comphep>




Figure 1: Exclusion contours for the combined χ^2 fit in the (c_V, c_F) plane, $\sqrt{s} = 250$ GeV. Blue, green and yellow areas correspond to $\Delta\chi^2 = 2.10, 4.61$ and 9.21 (CL of the fit is 65%, 90% and 99%), respectively. For the upper row of plots the channel $H \rightarrow \gamma\gamma$ is accounted for; its influence on the contours is very small. Lower row of plots shows the acceptable regions which appear without the $H \rightarrow \gamma\gamma$ channel. For the plots (b) and (d) the $H \rightarrow WW^*$ channel is excluded.




Figure 2: Three-dimensional plot of χ^2 as a function of (c_V, c_F) , $\sqrt{s} = 250$ GeV, (a) - all channels, (b) - only $H \rightarrow \gamma\gamma$ channel. Asymmetric narrow gullies in (b) appear due to the destructive interference of the fermion and the vector boson loop contributions. For this reason the left local minimum of the global χ^2 , figure (a), is lower than the right local minimum. The signal strength for all channels is assumed to be equal to 1 and the individual signal strength errors are very small (see Table 1).




Figure 3: Exclusion contours for the combined χ^2 fit in the (c_V, c_F) plane, $\sqrt{s} = 500$ GeV (upper row of plots) and $\sqrt{s} = 1000$ GeV (lower row of plots). Blue, green and yellow areas correspond to $\Delta\chi^2 = 2.10$, 4.61 and 9.21, respectively. For the plots (b) and (d) the $H \rightarrow WW^*$ channel is excluded.