

# On monotonicity and order-preservation for multidimensional $G$ -diffusion processes

Peng Luo\*      Guangyan Jia†

## Abstract

In this paper, we prove a comparison theorem for multidimensional  $G$ -SDEs. Moreover we obtain respectively the sufficient conditions and necessary conditions of the monotonicity and order-preservation for two multidimensional  $G$ -diffusion processes. Finally, we give some applications.

**Keywords:**  $G$ -diffusion processes,  $G$ -SDE, Comparison theorem, Monotonicity, Order-preservation.

**Mathematics Subject Classification (2010).** 60H30, 60H10.

## 1 Introduction

In the classical framework, it is known that Itô diffusions can be used to construct linear semigroups, which are known as Markov semigroups. The relationships among Itô's diffusions, Markov semigroups and infinitesimal generators have been well studied and many interesting results have been deduced. They can be summarized as follows. We suppose  $(X_t)_{t \geq 0}$  to be  $n$ -dimensional Itô diffusion

$$dX_t = b(X_t)dt + \sigma(X_t)dW_t,$$

where  $(W_t)_{t \geq 0}$  is a  $d$ -dimensional Brownian motion and  $b, \sigma$  are Lipschitz continuous functions on  $\mathbb{R}^n$ . The Markov semigroup  $P_t$  is defined by  $P_t f(x) = E[f(X_t^{0,x})]$ , where  $X_t^{0,x}$  represents the Itô process with initial condition  $x$  at initial time  $t = 0$  and  $f$  is a function defined on  $\mathbb{R}^n$ . Here  $E[\cdot]$  stands for the expectation related to a probability  $P$ . The infinitesimal generator  $L$  of the Markov semigroup, which satisfies

$$Lf(x) = \lim_{t \rightarrow 0^+} \frac{P_t f(x) - f(x)}{t}$$

for  $f$  appropriately taken such that the above limit exists, is of the following form:

$$Lf = \frac{1}{2} \text{tr}[\sigma^* \frac{\partial^2}{\partial x^2} f \sigma] + b^* \frac{\partial}{\partial x} f,$$

where " $*$ " denotes the transposition. For more details, the readers can refer to, for example, Stroock and Varadhan ([36]) and Rogers and Williams ([35]).

The monotonicity property of the semigroups associated with the corresponding diffusions was initiated by Holly ([13]) and studied by Cox ([4]) and Harris ([10]). Afterwards, Herbst and Pitt

---

\*School of Mathematics and Qilu Securities Institute for Financial Studies, Shandong University and Department of Mathematics and Statistics, University of Konstanz; pengluo1989@gmail.com. The author thanks the partial support from China Scholarship Council, and the 111 project(No. B12023).

†Qilu Securities Institute for Financial Studies, Shandong University; jiagy@sdu.edu.cn. The author thanks the partial support from the NSF of China (11171186), and the 111 project(No. B12023).

([11]) investigated the use of diffusion equations as a tool for establishing stochastic monotonicity of semigroups. Chen and Wang ([3]) continued the study on the order-preservation for multidimensional diffusion processes. One of the main results of Chen and Wang ([3]), which covers the monotonicity result in Herbst and Pitt ([11]), is as follows:

**Lemma 1.1** (Chen and Wang ([3], Theorem 1.3)) *Let  $A = \frac{1}{2}\sum_{i,j=1}^n a_{ij} \frac{\partial^2}{\partial x_i \partial x_j} + \sum_{i=1}^n l_i \frac{\partial}{\partial x_i}$  (resp.  $\bar{A} = \frac{1}{2}\sum_{i,j=1}^n \bar{a}_{ij} \frac{\partial^2}{\partial x_i \partial x_j} + \sum_{i=1}^n \bar{l}_i \frac{\partial}{\partial x_i}$ ). Assume that  $(a_{ij})$  and  $(\bar{a}_{ij})$  are nonnegative definite everywhere,  $a_{ij}, \bar{a}_{ij}, l_i, \bar{l}_i \in C(\mathbb{R}^n)$  and the martingale problems for  $A$  and  $\bar{A}$  are well posed. Let  $P_t$  (resp.  $\bar{P}_t$ ) be the Markov semigroup generated by  $A$  (resp.  $\bar{A}$ ). Then  $P_t \geq \bar{P}_t$  if and only if the following two conditions hold:*

- (a) for all  $i$  and  $j$ ,  $a_{ij} \equiv \bar{a}_{ij}$  and  $a_{ij}(x)$  depends only on  $x_i$  and  $x_j$ ;
- (b) for all  $i$ ,  $l_i(x) \geq \bar{l}_i(y)$  whenever  $x_i = y_i$  and  $x_j \geq y_j$  for all  $j \in \{1, \dots, n\}$  and  $j \neq i$ .

Peng ([27]) introduced the notion of  $g$ -expectation defined via a backward stochastic differential equation (BSDE). A  $g$ -expectation preserves most properties of the classical expectations except non-linearity since it is a nonlinear functional. Its nonlinearity can be characterized by its generator  $g$ . Zhang and Jia ([37]) constructed a nonlinear semigroup by decoupled FBSDEs. They obtained the equivalent conditions of the monotonicity and order-preservation of semigroups.

Recently, Peng systemically established a time-consistent fully nonlinear expectation theory ( see [28], [29] and [30]). As a typical and important case, Peng introduced the  $G$ -expectation theory ( see [32] and the references therein) in 2006. In the  $G$ -expectation framework ( $G$ -framework for short), the notion of  $G$ -Brownian motion and the corresponding stochastic calculus of Itô's type were established. On that basis, Gao [7] and Peng [31] have studied the existence and uniqueness of the solution of  $G$ -SDEs under a standard Lipschitz condition on its coefficients. Moreover, based on Gao [7], Bai and Lin [2] obtained the existence and uniqueness of the solution of GSDEs under some integral-Lipschitz conditions. For a recent account and development of this theory we refer the reader to ([20],[21],[22],[23],[24]). Recently, Hu et. al. ([14]) proved an existence and uniqueness result on BSDEs driven by  $G$ -Brownian motions (G-BSDEs), further (in [15]) they gave a comparison theorem for G-BSDEs. He et. al. ([12]) proved the representation theorem for generators of G-BSDEs, and then the converse comparison theorem of G-BSDEs and some equivalent results for nonlinear expectations generated by G-BSDEs.

This paper is organized as follow: In section 2, we recall some notations and results that we will use in this paper. In section 3, we give our assumptions and recall some notations and results of  $G$ -SDEs and  $G$ -BSDEs. In section 4, we obtain a comparison theorem for multidimensional  $G$ -SDEs. In section 5, we obtain respectively the sufficient conditions and necessary conditions of the monotonicity and order-preservation of two multidimensional  $G$ -diffusion processes. In section 6, some applications are given.

## 2 Preliminaries

### 2.1 Sublinear expectation, $G$ -Brownian motion and capacity

#### 2.1.1 Sublinear expectation

We present some preliminaries in the theory of sublinear expectation,  $G$ -Brownian motions and the capacity under  $G$ -framework. More details can be found in Peng [31] and Li and Peng [19].

**Definition 2.1** *Let  $\Omega$  be a given set and let  $\mathcal{H}$  be a linear space of real valued functions defined on  $\Omega$  with  $c \in \mathcal{H}$  for all constants  $c$ , and  $|X| \in \mathcal{H}$ , if  $X \in \mathcal{H}$ .  $\mathcal{H}$  is considered as the space of our random variables. A sublinear expectation  $\hat{\mathbb{E}}$  on  $\mathcal{H}$  is a functional  $\hat{\mathbb{E}} : \mathcal{H} \rightarrow \mathbb{R}$  satisfying the following*

properties: for all  $X, Y \in \mathcal{H}$ , we have

(a) *Monotonicity*: if  $X \geq Y$ , then  $\hat{\mathbb{E}}[X] \geq \hat{\mathbb{E}}[Y]$ .

(b) *Constant preserving*:  $\hat{\mathbb{E}}[c] = c, \forall c \in \mathbb{R}$ .

(c) *Sub-additivity*:  $\hat{\mathbb{E}}[X + Y] \leq \hat{\mathbb{E}}[X] + \hat{\mathbb{E}}[Y]$ .

(d) *Positive homogeneity*:  $\hat{\mathbb{E}}[\lambda X] = \lambda \hat{\mathbb{E}}[X], \forall \lambda \geq 0$ .

The triple  $(\Omega, \mathcal{H}, \hat{\mathbb{E}})$  is called a *sublinear expectation space*.  $X \in \mathcal{H}$  is called a *random variable* in  $(\Omega, \mathcal{H}, \hat{\mathbb{E}})$ . We often call  $Y = (Y_1, \dots, Y_d), Y_i \in \mathcal{H}$  a *d-dimensional random vector* in  $(\Omega, \mathcal{H}, \hat{\mathbb{E}})$ . Let us consider a space of random variables  $\mathcal{H}$  satisfying: if  $X_i \in \mathcal{H}, i = 1, \dots, d$ , then  $\varphi(X_1, \dots, X_d) \in \mathcal{H}$ , for all  $\varphi \in C_{b, Lip}(\mathbb{R}^d)$ , where  $C_{b, Lip}(\mathbb{R}^d)$  is the space of all bounded real-valued Lipschitz continuous functions.

**Definition 2.2** In a sublinear expectation space  $(\Omega, \mathcal{H}, \hat{\mathbb{E}})$  an *m-dimensional random vector*  $X = (X_1, \dots, X_m)$  is said to be *independent from another n-dimensional random vector*  $Y = (Y_1, \dots, Y_n)$  under  $\hat{\mathbb{E}}$  if for any test function  $\varphi \in C_{b, Lip}(\mathbb{R}^{m+n})$  we have

$$\hat{\mathbb{E}}[\varphi(X, Y)] = \hat{\mathbb{E}}[\hat{\mathbb{E}}[\varphi(x, Y)]_{x=X}]$$

**Definition 2.3** Let  $X_1$  and  $X_2$  be two *n-dimensional random vector* defined on sublinear expectation spaces  $(\Omega_1, \mathcal{H}_1, \hat{\mathbb{E}}_1)$  and  $(\Omega_2, \mathcal{H}_2, \hat{\mathbb{E}}_2)$ , respectively. They are called *identically distributed*, denoted by  $X_1 \stackrel{d}{=} X_2$ , if

$$\hat{\mathbb{E}}_1[\varphi(X_1)] = \hat{\mathbb{E}}_2[\varphi(X_2)], \forall \varphi \in C_{b, Lip}(\mathbb{R}^n)$$

We call  $\bar{X}$  an *independent copy of X* if  $\bar{X} \stackrel{d}{=} X$  and  $\bar{X}$  is independent from  $X$ .

**Definition 2.4 (G-normal distribution)** An *m-dimensional random vector*  $X = (X_1, \dots, X_m)$  on a sublinear expectation space  $(\Omega, \mathcal{H}, \hat{\mathbb{E}})$  is called (*centralized*) *G-normal distributed* if for any  $a, b \geq 0$

$$aX + b\bar{X} \stackrel{d}{=} \sqrt{a^2 + b^2}X$$

where  $\bar{X}$  is an independent copy of  $X$ . The letter  $G$  denotes the function

$$G(A) := \frac{1}{2} \hat{\mathbb{E}}[(AX, X)] : \mathbb{S}(d) \mapsto \mathbb{R}.$$

### 2.1.2 G-Brownian motion

**Definition 2.5 (G-Brownian motion)** Let  $G : \mathbb{S}(d) \mapsto \mathbb{R}$  be a given monotonic and sublinear function. A process  $(B(t) \in \mathcal{H})_{t \geq 0}$  on a sublinear expectation space  $(\Omega, \mathcal{H}, \hat{\mathbb{E}})$  is called a *G-Brownian motion* if the following properties are satisfied:

(a)  $B(0) = 0$ .

(b) For each  $t, s \geq 0$  the increment  $B_{t+s} - B_t \stackrel{d}{=} \sqrt{s}X$  and independent from  $(B_{t_1}, B_{t_2}, \dots, B_{t_n})$  for each  $n \in \mathbb{N}, 0 \leq t_1 \leq t_2 \leq \dots \leq t_n \leq t$ , where  $X$  is *G-normal distributed*.

We denote by  $\Omega = C_0^d(\mathbb{R}^+)$  the space of all  $\mathbb{R}^d$ -value continuous paths  $(\omega_t)_{t \in \mathbb{R}^+}$ , with  $\omega_0 = 0$ , equipped with the distance

$$\rho(\omega^1, \omega^2) := \sum_{i=1}^{\infty} 2^{-i} [(\max_{t \in [0, i]} |\omega^1(t) - \omega^2(t)|) \wedge 1].$$

We denote by  $\mathcal{B}(\Omega)$  the Borel  $\sigma$ -algebra of  $\Omega$ .

We also denote, for each  $t \in [0, \infty)$ :

$\Omega_t := \{\omega(\cdot \wedge t) : \omega \in \Omega\}$ ,  
 $\mathcal{F}_t := \mathcal{B}(\Omega_t)$ ,  
 $L^0(\Omega)$  : the space of all  $\mathcal{B}(\Omega)$ -measurable real function,  
 $L^0(\Omega_t)$  : the space of all  $\mathcal{B}(\Omega_t)$ -measurable real function,  
 $B_b(\Omega_t)$  : all bounded elements in  $L^0(\Omega)$ ,  $B_b(\Omega_t) := B_b(\Omega) \cap L^0(\Omega_t)$ ,  
 $C_b(\Omega_t)$  : all continuous elements in  $B_b(\Omega)$ ,  $C_b(\Omega_t) := B_b(\Omega) \cap L^0(\Omega_t)$ .

In Peng [31], a  $G$ -Brownian motion is constructed on a sublinear expectation space  $(\Omega, L_G^p, \hat{\mathbb{E}}, (\hat{\mathbb{E}}_t)_{t \geq 0})$  for  $p = 1$ , where  $L_G^p$  is a Banach space under the natural norm  $\|X\|_p = \hat{\mathbb{E}}[|X|^p]^{1/p}$ . In this space the corresponding canonical process  $B(t, \omega) = \omega(t)$ ,  $t \in [0, \infty)$ , for  $\omega \in \Omega$ , is a  $G$ -Brownian motion. It is proved in Denis et al.[5] that  $L^0(\Omega) \supset L_G^p(\Omega) \supset C_b(\Omega)$ , and there exists a weakly compact family  $\mathcal{P}$  of probability measures defined on  $(\Omega, \mathcal{B}(\Omega))$  such that

$$\hat{\mathbb{E}}[X] = \sup_{P \in \mathcal{P}} E_P[X], \quad X \in L_G^1(\Omega).$$

We now introduce the natural choquet capacity

$$c(A) := \sup_{P \in \mathcal{P}} P(A), \quad A \in \mathcal{B}(\Omega).$$

**Definition 2.6** A set  $A \subset \Omega$  is polar if  $c(A) = 0$ . A property holds ‘quasi-surely’ (q.s.) if it holds outside a polar set.

**Definition 2.7** A real function  $X$  on  $\Omega$  is said to be quasi-continuous if for each  $\varepsilon > 0$ , there exists an open set  $O$  with  $c(O) < \varepsilon$  such that  $X|_{O^c}$  is continuous.

Then  $L_G^p(\Omega)$  can be characterized as follows:

$$L_G^p(\Omega) = \{X \in L^0(\Omega) \mid \sup_{P \in \mathcal{P}} E_P[|X|^p] < \infty, \text{ and } X \text{ is } c \text{ quasi-surely continuous}\}.$$

We denote, for  $p > 0$ ,

$$\mathcal{L}^p := \{X \in L^0(\Omega) : \hat{\mathbb{E}}[|X|^p] = \sup_{P \in \mathcal{P}} E_P[|X|^p] < \infty\};$$

$$\mathcal{N}^p := \{X \in L^0(\Omega) : \hat{\mathbb{E}}[|X|^p] = 0\};$$

$$\mathcal{N} := \{X \in L^0(\Omega) : X=0, \text{ } c\text{-quasi surely (q.s.)}\}$$

It is seen that  $\mathcal{L}^p$  and  $\mathcal{N}^p$  are linear spaces and  $\mathcal{N}^p = \mathcal{N}$ , for each  $p > 0$ . We define the space  $L^p(\Omega) = \mathcal{L}^p / \mathcal{N}$  as the equivalence classes of  $\mathcal{L}^p$  modulo equality in  $\|\cdot\|_p$ . Similarly, we can define  $L^p(\Omega_t) = L^p(\Omega) \cap L^0(\Omega_t)$ . As usual, we do not make the distinction between classes and their representatives.

**Definition 2.8** Let  $L_b^p(\Omega)$  be the completion of  $B_b(\Omega)$  under the Banach norm  $(\hat{\mathbb{E}}[|X|^p])^{1/p}$ .

Then we have the following characterisation (see [5]): for each  $p \geq 1$ ,

$$L_b^p(\Omega) = \{X \in L^p(\Omega) : \lim_{n \rightarrow \infty} \hat{\mathbb{E}}[|X|^p \mathbf{I}_{\{|X| > n\}}] = 0\}.$$

**Definition 2.9** For  $p \geq 1$  and  $T \in \mathbb{R}^+$  be fixed. Consider the following simple type of processes:

$$M_G^{0,p}(0, T) = \{\eta := \eta(t, \omega) = \sum_{j=0}^{N-1} \xi_j(\omega) I_{[t_j, t_{j+1})}(t) \\ \forall N > 0, 0 = t_0 < \dots < t_N = T, \xi_j(\omega) \in \mathbb{L}_G^p(\Omega_{t_j}), j = 0, 1, 2, \dots, N-1\}.$$

For each  $p \geq 1$ , we denote by  $M_G^p(0, T)$  the completion of  $M_G^{0,p}(0, T)$  under the norm

$$\|\eta\|_{M_G^p(0, T)} = \left| \int_0^T \hat{\mathbb{E}}[|\eta(t)|^p] dt \right|^{1/p}.$$

**Definition 2.10** For  $p \geq 1$  and  $T \in \mathbb{R}^+$  be fixed. Consider the following simple type of processes:

$$M_b^0(0, T) = \left\{ \eta := \eta_t(\omega) = \sum_{j=0}^{N-1} \xi_j(\omega) \mathbf{I}_{[t_j, t_{j+1})}(t) \right.$$

$$\left. \forall N > 0, 0 = t_0 < \dots < t_N = T, \xi_j(\omega) \in B_b(\Omega_{t_j}), j = 0, 1, 2, \dots, N-1 \right\}.$$

For each  $p \geq 1$ , we denote by  $M_b^p(0, T)$  the completion of  $M_b^0(0, T)$  under the norm  $\|\cdot\|_{M_b^p(0, T)}$ .

**Definition 2.11** (Integration with respect to  $G$ -Brownian motion) For each  $\eta \in M_G^{0,2}(0, T)$  with the form

$$\eta_t(\omega) = \sum_{k=0}^{N-1} \xi_k(\omega) I_{[t_k, t_{k+1})}(t)$$

define

$$I(\eta) = \int_0^T \eta(s) dB(s) := \sum_{k=0}^{N-1} \xi_k (B_{t_{k+1}^N} - B_{t_k^N}).$$

The mapping  $I : M_G^{0,2}(0, T) \mapsto \mathbb{L}_G^2(\Omega_T)$  can be continuously extended to  $I : M_G^2(0, T) \mapsto \mathbb{L}_G^2(\Omega_T)$ . For each  $\eta \in M_G^2(0, T)$ , the stochastic integral is defined by

$$I(\eta) := \int_0^T \eta(s) dB_s, \quad \eta \in M_G^2(0, T).$$

We have the following general case.

**Definition 2.12** For each  $\eta \in M_b^0(0, T)$  with the form

$$\eta_t(\omega) = \sum_{k=0}^{N-1} \xi_k(\omega) \mathbf{I}_{[t_k, t_{k+1})}(t),$$

define

$$I(\eta) = \int_0^T \eta(s) dB_s := \sum_{k=0}^{N-1} \xi_k (B_{t_{k+1}^N} - B_{t_k^N}).$$

The mapping  $\mathbf{I} : M_b^0(0, T) \mapsto L^2(\Omega_T)$  can be continuously extended to  $\mathbf{I} : M_b^2(0, T) \mapsto L^2(\Omega_T)$ . For each  $\eta \in M_b^2(0, T)$ , the stochastic integral is defined by

$$I(\eta) := \int_0^T \eta(s) dB_s, \quad \eta \in M_b^2(0, T).$$

For notational simplicity, we denote by  $B^i := B^{e_i}$  the  $i$ th coordinate of the  $d$ -dimensional  $G$ -Brownian motion  $B$ , under a given orthonormal basis  $(e_1, \dots, e_d)$  of  $\mathbb{R}^d$ . We also denote by  $B_t^a := (a, B_t)$  for fixed  $a \in \mathbb{R}^d$ . Then  $(B_t^a)_{t \geq 0}$  is a 1-dimensional  $G$ -Brownian motion with  $\sigma_{aa^T}^2 = \hat{\mathbb{E}}[(a, B_1)^2]$  and  $\sigma_{-aa^T}^2 = -\hat{\mathbb{E}}[-(a, B_1)^2]$ . Let  $a$  and  $\bar{a}$  be two given vectors in  $\mathbb{R}^d$ . We can define

$$\langle B^a \rangle_t := (B_t^a)^2 - \int_0^t B_s^a dB_s^a$$

where  $\langle B^a \rangle$  is called the quadratic variation process of  $B^a$ . We can also define mutual variation process by

$$\langle B^a, B^{\bar{a}} \rangle_t := \frac{1}{4}[\langle B^a + B^{\bar{a}} \rangle_t - \langle B^a - B^{\bar{a}} \rangle_t]$$

Itô's integral with respect to  $\langle B^a \rangle$  or  $\langle B^i, B^j \rangle$  can be similarly defined. By Li and Peng [19], we have

**Lemma 2.13** *Let  $X \in M_b^p(0, T)$ . Then for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that for all  $\eta \in M_b^0(0, T)$  satisfying  $\hat{\mathbb{E}} \int_0^T |\eta_t| dt \leq \delta$  and  $|\eta_t(\omega)| \leq 1$ , we have  $\hat{\mathbb{E}} \int_0^T |X_t|^p |\eta_t| dt \leq \varepsilon$ .*

**Definition 2.14** *A stopping time  $\tau$  relative to the filtration  $(\mathcal{F}_t)$  is a map on  $\Omega$  with values in  $[0, T]$ , such that for every  $t$ ,*

$$\{\tau \leq t\} \in \mathcal{F}_t.$$

**Lemma 2.15** *For each stopping time  $\tau \in [0, T]$ , we have  $I_{[0, \tau]}(\cdot)X \in M_b^p(0, T)$ , for each  $X \in M_b^p(0, T)$ .*

**Definition 2.16** *A process  $(M_t)_{t \geq 0}$  is called a  $G$ -martingale if for each  $t \in [0, T]$ ,  $M_t \in \mathbb{L}_G^1(\Omega_t)$  and for each  $s \in [0, t]$ , we have*

$$\hat{\mathbb{E}}_s[M_t] = M_s.$$

**Corollary 2.17** *For each  $\eta \in M_G^2(0, T)$ , the process  $(\int_0^t \eta(s) dB_s)_{t \in [0, T]}$  is a  $G$ -martingale.*

### 3 $G$ -SDEs and $G$ -BSDEs

#### 3.1 $G$ -SDEs

We make use of the following assumptions on the generator  $b, h$  and  $\sigma$  of GSDE:

**(H1)**  $b, h_{ij}$  and  $\sigma_i$  are given  $\mathbb{R}^n$ -valued bounded Lipschitz continuous functions defined on  $[0, T] \times \mathbb{R}^n$  which satisfy the Lipschitz condition, i.e., there exists some constant  $K$  such that  $|\varphi(t, x) - \varphi(t, y)| \leq K|x - y|$ , for each  $t \in [0, T]$ ,  $x, y \in \mathbb{R}^n$ ,  $\varphi = b, h_{ij}$  and  $\sigma_i$  respectively,  $i, j = 1, \dots, d$ .

**(H2)**  $b, h_{ij}$  and  $\sigma_i$  are given  $\mathbb{R}^n$ -valued bounded Lipschitz continuous functions defined on  $\mathbb{R}^n$ , i.e., there exists some constant  $K$  such that  $|\varphi(x) - \varphi(y)| \leq K|x - y|$ , for each  $x, y \in \mathbb{R}^n$ ,  $\varphi = b, h_{ij}$  and  $\sigma_i$  respectively,  $i, j = 1, \dots, d$ .

We consider the following SDE driven by a  $d$ -dimensional  $G$ -Brownian motion:

$$X(t) = X(0) + \int_0^t b(s, X(s)) ds + \int_0^t h_{ij}(s, X(s)) d\langle B^i, B^j \rangle_s + \int_0^t \sigma_i(s, X(s)) dB_s^i, \quad t \in [0, T], \quad (1)$$

where the initial condition  $X_0 \in \mathbb{R}^n$  is a given constant.

**Theorem 3.1** *Under the assumption (H1), there exists a unique solution  $X \in M_G^2(0, T)$  of the stochastic differential equation (1).*

### 3.2 $G$ -BSDEs

In this subsection, we present some notations and results of  $G$ -BSDEs. More details can be found in [14] and [15].

**Definition 3.2** For fixed  $T > 0$ , let  $M_G^0(0, T)$  be the collection of processes in the following form: for a given partition  $\{t_0, \dots, t_N\} = \pi_T$  of  $[0, T]$ ,

$$\eta_t(\omega) = \sum_{j=0}^{N-1} \xi_j I_{[t_j, t_{j+1})}(t),$$

where  $\xi_j \in L_{ip}(\Omega_{t_j})$ ,  $j = 0, 1, 2, \dots, N-1$ . For  $p \geq 1$ , we denote by  $H_G^p(0, T)$ ,  $M_G^p(0, T)$  the completion of  $M_G^0(0, T)$  under the norms  $\|\eta\|_{H_G^p} = \{\hat{\mathbb{E}}[(\int_0^T |\eta_s|^2 ds)^{p/2}]\}^{1/p}$ ,  $\|\eta\|_{M_G^p} = \{\int_0^T \hat{\mathbb{E}}[|\eta_s|^p] ds\}^{1/p}$  respectively.

For each  $\eta \in M_G^1(0, T)$ , we can define the integrals  $\int_0^T \eta_t dt$  and  $\int_0^T \eta_t d\langle B^{\mathbf{a}}, B^{\bar{\mathbf{a}}}\rangle_t$  for each  $\mathbf{a}, \bar{\mathbf{a}} \in \mathbb{R}^d$ . For each  $\eta \in H_G^p(0, T; \mathbb{R}^d)$  with  $p \geq 1$ , we can define Itô's integral  $\int_0^T \eta_t dB_t$ .

Let  $S_G^0(0, T) = \{h(t, B_{t_1 \wedge t}, \dots, B_{t_n \wedge t}) : t_1, \dots, t_n \in [0, T], h \in C_{b, Lip}(\mathbb{R}^{n+1})\}$ . For  $p \geq 1$  and  $\eta \in S_G^0(0, T)$ , set  $\|\eta\|_{S_G^p} = \{\hat{\mathbb{E}}[\sup_{t \in [0, T]} |\eta_t|^p]\}^{\frac{1}{p}}$ . Denote by  $S_G^p(0, T)$  the completion of  $S_G^0(0, T)$  under the norm  $\|\cdot\|_{S_G^p}$ .

We only consider non-degenerate  $G$ -normal distribution, i.e.,

**(H3)** There exists some  $\underline{\sigma}^2 > 0$  such that  $G(A) - G(B) \geq \underline{\sigma}^2 \text{tr}[A - B]$  for any  $A \geq B$ .

We consider the following type of  $G$ -BSDEs (in this paper we always use Einstein convention):

$$\begin{aligned} Y_t &= \xi + \int_t^T f(s, Y_s, Z_s) ds + \int_t^T g_{ij}(s, Y_s, Z_s) d\langle B^i, B^j \rangle_s \\ &\quad - \int_t^T Z_s dB_s - (K_T - K_t), \end{aligned} \tag{2}$$

where

$$f(t, \omega, y, z), g_{ij}(t, \omega, y, z) : [0, T] \times \Omega_T \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$$

satisfy the following properties:

**(H4)** There exists some  $\beta > 1$  such that for any  $y, z$ ,  $f(\cdot, \cdot, y, z), g_{ij}(\cdot, \cdot, y, z) \in M_G^\beta(0, T)$ .

**(H5)** There exists some  $L > 0$  such that

$$|f(t, y, z) - f(t, y', z')| + \sum_{i,j=1}^d |g_{ij}(t, y, z) - g_{ij}(t, y', z')| \leq L(|y - y'| + |z - z'|).$$

For simplicity, we denote by  $\mathfrak{S}_G^\alpha(0, T)$  the collection of processes  $(Y, Z, K)$  such that  $Y \in S_G^\alpha(0, T)$ ,  $Z \in H_G^\alpha(0, T; \mathbb{R}^d)$ ,  $K$  is a decreasing  $G$ -martingale with  $K_0 = 0$  and  $K_T \in L_G^\alpha(\Omega_T)$ .

**Definition 3.3** Let  $\xi \in L_G^\beta(\Omega_T)$  and  $f$  satisfy (H4) and (H5) for some  $\beta > 1$ . A triplet of processes  $(Y, Z, K)$  is called a solution of equation (2) if for some  $1 < \alpha \leq \beta$  the following properties hold:

**(a)**  $(Y, Z, K) \in \mathfrak{S}_G^\alpha(0, T)$ ;

$$(b) \quad Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds + \int_t^T g_{ij}(s, Y_s, Z_s) d\langle B^i, B^j \rangle_s - \int_t^T Z_s dB_s - (K_T - K_t).$$

**Theorem 3.4** ([14]) *Assume that  $\xi \in L_G^\beta(\Omega_T)$  and  $f, g_{ij}$  satisfy (H4) and (H5) for some  $\beta > 1$ . Then equation (2) has a unique solution  $(Y, Z, K)$ . Moreover, for any  $1 < \alpha < \beta$  we have  $Y \in S_G^\alpha(0, T)$ ,  $Z \in H_G^\alpha(0, T; \mathbb{R}^d)$  and  $K_T \in L_G^\alpha(\Omega_T)$ .*

We consider the following  $G$ -BSDEs:

$$\begin{aligned} Y_t^{l,\xi} &= \xi + \int_t^T f^l(s, Y_s^{l,\xi}, Z_s^{l,\xi}) ds + \int_t^T g_{ij}^l(s, Y_s^{l,\xi}, Z_s^{l,\xi}) d\langle B^i, B^j \rangle_s \\ &\quad - \int_t^T Z_s^{l,\xi} dB_s - (K_T^{l,\xi} - K_t^{l,\xi}), \quad l = 1, 2, \end{aligned}$$

where  $g_{ij}^l = g_{ji}^l$ .

He and Hu [12] generalized the comparison theorem in [15].

**Proposition 3.5** *Let  $f^l$  and  $g_{ij}^l$  satisfy (H4) and (H5) for some  $\beta > 1$ ,  $l = 1, 2$ . If  $f^2 - f^1 + 2G((g_{ij}^2 - g_{ij}^1)_{i,j=1}^d) \leq 0$ , then for each  $\xi \in L_G^\beta(\Omega_T)$ , we have  $Y_t^{1,\xi} \geq Y_t^{2,\xi}$  for  $t \in [0, T]$ .*

### 3.3 Nonlinear Feynman-Kac Formula

In this subsection, we give the nonlinear Feynman-Kac Formula which was studied in Peng [32] for special type of  $G$ -BSDEs. We consider the following type of  $G$ -FBSDEs:

$$dX_s^{t,\xi} = b(s, X_s^{t,\xi}) ds + h_{ij}(s, X_s^{t,\xi}) d\langle B^i, B^j \rangle_s + \sigma_j(s, X_s^{t,\xi}) dB_s^j, \quad X_t^{t,\xi} = \xi, \quad (3)$$

$$\begin{aligned} Y_s^{t,\xi} &= \Phi(X_T^{t,\xi}) + \int_s^T f(r, X_r^{t,\xi}, Y_r^{t,\xi}, Z_r^{t,\xi}) dr + \int_s^T g_{ij}(r, X_r^{t,\xi}, Y_r^{t,\xi}, Z_r^{t,\xi}) d\langle B^i, B^j \rangle_r \\ &\quad - \int_s^T Z_r^{t,\xi} dB_r - (K_T^{t,\xi} - K_s^{t,\xi}), \end{aligned} \quad (4)$$

where  $b, h_{ij}, \sigma_j : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ ,  $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $f, g_{ij} : [0, T] \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$  are deterministic functions and satisfy the following conditions:

(A1)  $h_{ij} = h_{ji}$  and  $g_{ij} = g_{ji}$  for  $1 \leq i, j \leq d$ ;

(A2)  $b, h_{ij}, \sigma_j, f, g_{ij}$  are continuous in  $t$ ;

(A3) There exist a positive integer  $m$  and a constant  $L > 0$  such that

$$|b(t, x) - b(t, x')| + \sum_{i,j=1}^d |h_{ij}(t, x) - h_{ij}(t, x')| + \sum_{j=1}^d |\sigma_j(t, x) - \sigma_j(t, x')| \leq L|x - x'|,$$

$$|\Phi(x) - \Phi(x')| \leq L(1 + |x|^m + |x'|^m)|x - x'|,$$

$$\begin{aligned} &|f(t, x, y, z) - f(t, x', y', z')| + \sum_{i,j=1}^d |g_{ij}(t, x, y, z) - g_{ij}(t, x', y', z')| \\ &\leq L[(1 + |x|^m + |x'|^m)|x - x'| + |y - y'| + |z - z'|]. \end{aligned}$$



We define

$$u(t, x) := Y_t^{t,x}, \quad (t, x) \in [0, T] \times \mathbb{R}^n.$$

**Remark 3.6** *It is important to note that  $u(t, x)$  is a deterministic function of  $(t, x)$ , because  $b, h_{ij}, \sigma_j, \Phi, f, g_{ij}$  are deterministic functions and  $\tilde{B}_s := B_{t+s} - B_t$  is a  $G$ -Brownian motion.*

We now give the Feynman-Kac formula.

**Theorem 3.7** *Let  $u(t, x) := Y_t^{t,x}$  for  $(t, x) \in [0, T] \times \mathbb{R}^n$ . Then  $u(t, x)$  is the unique viscosity solution of the following PDE:*

$$\begin{cases} \partial_t u + F(D_x^2 u, D_x u, u, x, t) = 0, \\ u(T, x) = \Phi(x), \end{cases} \quad (5)$$

where

$$\begin{aligned} F(D_x^2 u, D_x u, u, x, t) = & G(H(D_x^2 u, D_x u, u, x, t)) + \langle b(t, x), D_x u \rangle \\ & + f(t, x, u, \langle \sigma_1(t, x), D_x u \rangle, \dots, \langle \sigma_d(t, x), D_x u \rangle), \end{aligned}$$

$$\begin{aligned} H_{ij}(D_x^2 u, D_x u, u, x, t) = & \langle D_x^2 u \sigma_i(t, x), \sigma_j(t, x) \rangle + 2 \langle D_x u, h_{ij}(t, x) \rangle \\ & + 2g_{ij}(t, x, u, \langle \sigma_1(t, x), D_x u \rangle, \dots, \langle \sigma_d(t, x), D_x u \rangle). \end{aligned}$$

## 4 Comparison Theorem for Multidimensional GSDEs

In the classical framework, comparison theorem for stochastic differential equations are well studied (see [1], [6], [8], [9], [18] and etc.). In particular, Geiß and Mantey ([8]) obtained a comparison theorem for multidimensional SDEs. We also refer to Lin ([21]) and Luo and Wang ([24]) for the comparison theorem of 1-dimensional  $G$ -SDEs. In this paper, we follow the idea of their proof to get our results. We consider the following SDEs driven by a  $d$ -dimensional  $G$ -Brownian motion:

$$X(t) = X(0) + \int_0^t b(s, X(s)) ds + \int_0^t h_{ij}(s, X(s)) d\langle B^i, B^j \rangle_s + \int_0^t \sigma_i(s, X(s)) dB_s^i, \quad t \in [0, T]$$

and

$$Y(t) = Y(0) + \int_0^t \bar{b}(s, Y(s)) ds + \int_0^t \bar{h}_{ij}(s, Y(s)) d\langle B^i, B^j \rangle_s + \int_0^t \sigma_i(s, Y(s)) dB_s^i, \quad t \in [0, T],$$

where the initial conditions  $X(0), Y(0) \in \mathbb{R}^n$  are given constants together with

$$X_k(0) \leq Y_k(0), \quad k = 1, \dots, n.$$

We now give a comparison theorem for multidimensional  $G$ -SDEs.

**Theorem 4.1** *Suppose that the following two conditions hold.*

**(B1)** *For any  $t \in [0, T]$ , and  $i = 1, \dots, n$ , the inequality*

$$b_i(t, x) - \bar{b}_i(t, y) + G([\langle h_{lk} \rangle_i + \langle h_{kl} \rangle_i]_{l,k=1}^d(t, x) - [\langle \bar{h}_{lk} \rangle_i + \langle \bar{h}_{kl} \rangle_i]_{l,k=1}^d(t, y)) \leq 0$$

*are fulfilled, whenever  $x_i = y_i$  and  $x_j \leq y_j$  for all  $j \neq i$ .*

(B2)  $b, h_{ij}, \sigma_i$  and  $\bar{b}, \bar{h}_{ij}, \sigma_i$  satisfy (H1) and  $(\sigma_i)_k$  depends only on  $x_k$ , for each  $k = 1, \dots, n$ ,  $i, j = 1, \dots, d$ , i.e.,

$$|(\sigma_i)_k(t, x) - (\sigma_i)_k(t, y)| \leq K|x_k - y_k|$$

for all  $t \in [0, T]$ ,  $x, y \in \mathbb{R}^n$ .

Then for all  $t \in [0, T]$ ,

$$X_k(t) \leq Y_k(t) \quad k = 1, \dots, n \quad q.s.$$

**Proof.** We first proof the theorem under the following condition (B1') instead of (B1).

(B1') For any  $t \in [0, T]$  and  $i = 1, \dots, n$  the inequality

$$b_i(t, x) - \bar{b}_i(t, y) + G([(h_{lk})_i + (h_{kl})_i]_{l,k=1}^d(t, x) - [(\bar{h}_{lk})_i + (\bar{h}_{kl})_i]_{l,k=1}^d(t, y)) < 0$$

are fulfilled, whenever  $x_i = y_i$  and  $x_j \leq y_j$  for all  $j \neq i$ .

Define the stopping times

$$\tau_j = \inf\{t > 0 : X_j(t) > Y_j(t)\} \wedge T, \quad j = 1, \dots, n$$

and

$$\tau := \tau_1 \wedge \dots \wedge \tau_n.$$

Obviously,  $X_j(\tau_j) = Y_j(\tau_j)$  and  $X_j(\tau) \leq Y_j(\tau)$ ,  $j = 1, \dots, n$ . Because of condition (B1'), the continuity of  $b, h_{ij}, \bar{b}, \bar{h}_{ij}, i, j = 1, 2, \dots, d$ , and the continuity of  $X$  and  $Y$  there exists a stopping time  $T \geq \kappa > \tau$  q.s. defined on  $\{\tau < T\}$  such that

$$\begin{aligned} & b_j(t, X(s)) - \bar{b}_j(t, Y_1(s), \dots, Y_{j-1}(s), X_j(s), Y_{j+1}(s), \dots, Y_d(s)) + G([(h_{lk})_j + (h_{kl})_j]_{l,k=1}^d(t, X(s)) \\ & - [(\bar{h}_{lk})_j + (\bar{h}_{kl})_j]_{l,k=1}^d(t, Y_1(s), \dots, Y_{j-1}(s), X_j(s), Y_{j+1}(s), \dots, Y_d(s))) < 0 \end{aligned} \quad (6)$$

on  $\{\tau_j = \tau < T\}$  for all  $s \in [\tau, \kappa]$  q.s.. Actually, we can define

$$\begin{aligned} \kappa_1 := \inf\{t > \tau : & b_j(t, X(s)) - \bar{b}_j(t, Y_1(s), \dots, Y_{j-1}(s), X_j(s), Y_{j+1}(s), \dots, Y_d(s)) \\ & + G([(h_{lk})_j + (h_{kl})_j]_{l,k=1}^d(t, X(s)) \\ & - [(\bar{h}_{lk})_j + (\bar{h}_{kl})_j]_{l,k=1}^d(t, Y_1(s), \dots, Y_{j-1}(s), X_j(s), Y_{j+1}(s), \dots, Y_d(s))) > 0\} \wedge T, \end{aligned}$$

then we take  $\kappa = \frac{\tau + \kappa_1}{2}$ . It is easy to check that  $\kappa$  satisfies the above condition.

Now we define  $\rho(x) = Kx$  for  $x \geq 0$ . Then  $\int_{0+} \rho^{-2}(u)du = \infty$ . Hence there exists a strictly increasing sequence  $\{a_n\}_{n=0}^\infty$  such that  $a_0 = 1$ ,  $\lim_{n \rightarrow \infty} a_n = 0$  (actually,  $a_n = \frac{2}{n(n+1)K^2+2}$ ) and

$$\int_{a_n}^{a_{n-1}} \rho^{-2}(u)du = n, \quad \text{for all } n \geq 1.$$

Let  $\psi_n$  be a continuous function such that its support is contained in  $(a_n, a_{n-1})$ ,  $0 \leq \psi_n(u) \leq 2\rho^{-2}(u)n^{-1}$  and  $\int_{a_n}^{a_{n-1}} \psi_n(u)du = 1$ . Put

$$\varphi_n(x) = \begin{cases} 0, & x \leq 0, \\ \int_0^x \int_0^y \psi_n(u)dudy, & x > 0. \end{cases}$$

One can easily see that  $\varphi_n$  is twice continuously differentiable,  $\varphi_n(0) = 0$  for  $x \leq 0$ ,  $0 \leq \varphi'_n(x) \leq 1$  and  $\varphi_n(x) \uparrow x^+$  as  $n \rightarrow \infty$ . Assume

$$c(\{\tau < T\}) > 0. \quad (7)$$

An application of  $G$ -Itô's formula (see [19]) yields

$$\begin{aligned} & \varphi_n(X_k((\tau+t) \wedge \kappa) - Y_k((\tau+t) \wedge \kappa)) \\ = & \varphi_n(X_k(\tau) - Y_k(\tau)) \\ & + \int_{\tau}^{(\tau+t) \wedge \kappa} \varphi'_n(X_k(s) - Y_k(s)) [b_k(s, X(s)) - \bar{b}_k(s, Y(s))] ds \\ & + \int_{\tau}^{(\tau+t) \wedge \kappa} \varphi'_n(X_k(s) - Y_k(s)) [(h_{ij})_k(s, X(s)) - (\bar{h}_{ij})_k(s, Y(s))] d\langle B^i, B^j \rangle_s \\ & + \int_{\tau}^{(\tau+t) \wedge \kappa} \varphi'_n(X_k(s) - Y_k(s)) [(\sigma_i)_k(s, X(s)) - (\sigma_i)_k(s, Y(s))] dB_s^i \\ & + \frac{1}{2} \int_{\tau}^{(\tau+t) \wedge \kappa} \varphi''_n(X_k(s) - Y_k(s)) [(\sigma_i)_k(s, X(s)) - (\sigma_i)_k(s, Y(s))]^2 d\langle B^i \rangle_s \\ = & \varphi_n(X_k(\tau) - Y_k(\tau \wedge T)) \\ & + \int_{\tau}^{(\tau+t) \wedge \kappa} \varphi'_n(X_k(s) - Y_k(s)) [b_k(s, X(s)) - \bar{b}_k(s, Y(s))] \\ & + G([(h_{ij})_k + (h_{ji})_k]_{i,j=1}^d(s, X(s)) - [(\bar{h}_{ij})_k + (\bar{h}_{ji})_k]_{i,j=1}^d(s, Y(s))] ds \\ & + V_{(\tau+t) \wedge \kappa} - V_{\tau} \\ & + \int_{\tau}^{(\tau+t) \wedge \kappa} \varphi'_n(X_k(s) - Y_k(s)) [(\sigma_i)_k(s, X(s)) - (\sigma_i)_k(s, Y(s))] dB_s^i \\ & + \frac{1}{2} \int_{\tau \wedge T}^{(\tau+t) \wedge \kappa} \varphi''_n(X_k(s) - Y_k(s)) [(\sigma_i)_k(s, X(s)) - (\sigma_i)_k(s, Y(s))]^2 d\langle B^i \rangle_s \end{aligned}$$

where

$$\begin{aligned} V_t = & \int_0^t \varphi'_n(X_k(s) - Y_k(s)) [(h_{ij})_k(s, X(s)) - (\bar{h}_{ij})_k(s, Y(s))] d\langle B^i, B^j \rangle_s \\ & - \int_0^t \varphi'_n(X_k(s) - Y_k(s)) G([(h_{ij})_k + (h_{ji})_k]_{i,j=1}^d(s, X(s)) - [(\bar{h}_{ij})_k + (\bar{h}_{ji})_k]_{i,j=1}^d(s, Y(s))] ds. \end{aligned}$$

It is easy to check that  $(V_t)_{0 \leq t \leq T}$  is a decreasing process. Thus

$$\begin{aligned} & \varphi_n(X_k((\tau+t) \wedge \kappa) - Y_k((\tau+t) \wedge \kappa)) \\ \leq & \varphi_n(X_k(\tau) - Y_k(\tau)) \\ & + \int_{\tau}^{(\tau+t) \wedge \kappa} \varphi'_n(X_k(s) - Y_k(s)) [b_k(s, X(s)) - \bar{b}_k(s, Y(s))] \\ & + G([(h_{ij})_k + (h_{ji})_k]_{i,j=1}^d(s, X(s)) - [(\bar{h}_{ij})_k + (\bar{h}_{ji})_k]_{i,j=1}^d(s, Y(s))] ds \\ & + \int_{\tau}^{(\tau+t) \wedge \kappa} \varphi'_n(X_k(s) - Y_k(s)) [(\sigma_i)_k(s, X(s)) - (\sigma_i)_k(s, Y(s))] dB_s^i \\ & + \frac{1}{2} \int_{\tau}^{(\tau+t) \wedge \kappa} \varphi''_n(X_k(s) - Y_k(s)) [(\sigma_i)_k(s, X(s)) - (\sigma_i)_k(s, Y(s))]^2 d\langle B^i \rangle_s \\ = & S_1(n) + S_2(n) + S_3(n) + S_4(n). \end{aligned}$$

Obviously, from the construction it follows that  $S_1(n) = 0$ ,  $n = 1, 2, \dots$ ,

$$\hat{\mathbb{E}}[S_3(n)\mathbf{1}_{\{\tau_k=\tau\}}] = 0.$$

From (B2) we derive

$$\hat{\mathbb{E}}[|S_4(n)|] \leq \frac{Ct}{n}$$

Relation (6) implies

$$\begin{aligned} & S_2(n)\mathbf{1}_{\{\tau_k=\tau\}} \\ & \leq \mathbf{1}_{\{\tau_k=\tau\}} \int_{\tau}^{(\tau+t)\wedge\kappa} \varphi'_n(X_k(s) - Y_k(s))[b_i(t, X(s)) - \bar{b}_i(t, Y_1(s), \dots, Y_{k-1}(s), X_k(s), Y_{k+1}(s), \dots, Y_d(s))] \\ & + G([(h_{ij})_k + (h_{ij})_k]_{i,j=1}^d(t, X(s)) \\ & - [(\bar{h}_{ij})_k + (\bar{h}_{ij})_k]_{i,j=1}^d(t, Y_1(s), \dots, Y_{k-1}(s), X_k(s), Y_{k+1}(s), \dots, Y_d(s)))] ds \\ & + \mathbf{1}_{\{\tau_k=\tau\}} \int_{\tau}^{(\tau+t)\wedge\kappa} \varphi'_n(X_k(s) - Y_k(s))[\bar{b}_i(t, Y_1(s), \dots, Y_{k-1}(s), X_k(s), Y_{k+1}(s), \dots, Y_d(s)) - \bar{b}_k(s, Y(s))] \\ & + G([\bar{h}_{ij})_k + (\bar{h}_{ij})_k]_{i,j=1}^d(s, Y_1(s), \dots, Y_{k-1}(s), X_k(s), Y_{k+1}(s), \dots, Y_d(s)) - [(\bar{h}_{ij})_k + (\bar{h}_{ij})_k]_{i,j=1}^d(s, Y(s))] ds \\ & \leq \mathbf{1}_{\{\tau_k=\tau\}} K \int_{\tau}^{(\tau+t)\wedge\kappa} [X_k(s) - Y_k(s)]^+ ds \\ & \leq \mathbf{1}_{\{\tau_k=\tau\}} K \int_0^t [X_k((\tau+s)\wedge\kappa) - Y_k((\tau+s)\wedge\kappa)]^+ ds. \end{aligned}$$

Consequently, as  $n \rightarrow \infty$  we arrive at

$$\hat{\mathbb{E}}[(X_k((\tau+t)\wedge\kappa) - Y_k((\tau+t)\wedge\kappa))^+ \mathbf{1}_{\{\tau_k=\tau\}}] \leq K \int_0^t \hat{\mathbb{E}}[(X_k((\tau+s)\wedge\kappa) - Y_k((\tau+s)\wedge\kappa))^+ \mathbf{1}_{\{\tau_k=\tau\}}] ds.$$

In view of Gronwall's inequality this implies

$$\hat{\mathbb{E}}[(X_k((\tau+t)\wedge\kappa) - Y_k((\tau+t)\wedge\kappa))^+ \mathbf{1}_{\{\tau_k=\tau\}}] = 0$$

and hence

$$X_k((\tau+t)\wedge\kappa) \leq Y_k((\tau+t)\wedge\kappa) \quad q.s.$$

on  $\{\tau_k = \tau\}$  for all  $t \in [\tau, \kappa]$ . This contradicts (7).

Now we consider the condition (B1). Let  $\epsilon > 0$  be arbitrarily chosen and define

$$b_k^\epsilon := b_k - \epsilon, \quad k = 1, \dots, n.$$

From (B1) it follows immediately that  $b^\epsilon$  satisfies condition (B1'). Consequently, we get for the corresponding solutions  $X^\epsilon$  and  $Y$  the relation

$$X_k^\epsilon(t) \leq Y_k(t) \quad q.s.$$

for all  $t \in [0, T]$ ,  $k = 1, \dots, n$ . Choose a strictly decreasing sequence  $(\epsilon_m)_{m \geq 1}$  with  $\lim_{m \rightarrow \infty} \epsilon_m = 0$ . By the same arguments as above we get

$$X_k^{\epsilon_1}(t) \leq X_k^{\epsilon_2}(t) \leq \dots \leq Y_k(t) \quad q.s.$$

as well as

$$X_k^{\epsilon_1}(t) \leq X_k^{\epsilon_2}(t) \leq \dots \leq X_k(t) \quad q.s.$$

for all  $t \in [0, T]$ ,  $k = 1, \dots, n$ .

Define

$$\tilde{X}_k(t) := \lim_{m \rightarrow \infty} X_k^{\epsilon_m}(t)$$

for each  $t \in [0, T]$ . Obviously,

$$\tilde{X}_k(t) \leq Y_k(t) \quad q.s.$$

for all  $t \in [0, T]$ ,  $k = 1, \dots, n$ . Moreover, we have,

$$\begin{aligned} \hat{\mathbb{E}}\left[\sup_{0 \leq t \leq T} |X^{\epsilon_m}(t) - X(t)|^2\right] &= \hat{\mathbb{E}}\left[\sup_{0 \leq t \leq T} \left| \int_0^t (b^{\epsilon_m}(s, X^{\epsilon_m}(s)) - b(s, X(s))) ds \right. \right. \\ &\quad \left. \left. + \int_0^t (h_{ij}(s, X^{\epsilon_m}(s)) - h_{ij}(s, X(s))) d\langle B^i, B^j \rangle \right. \right. \\ &\quad \left. \left. + \int_0^t (\sigma_i(s, X^{\epsilon_m}(s)) - \sigma_i(s, X(s))) dB_s^i \right|^2\right] \\ &\leq 3\left(\hat{\mathbb{E}}\left[\sup_{0 \leq t \leq T} \left| \int_0^t (b^{\epsilon_m}(s, X^{\epsilon_m}(s)) - b(s, X(s))) ds \right|^2\right] \right. \\ &\quad \left. + \hat{\mathbb{E}}\left[\sup_{0 \leq t \leq T} \left| \int_0^t (h_{ij}(s, X^{\epsilon_m}(s)) - h_{ij}(s, X(s))) d\langle B^i, B^j \rangle \right|^2\right] \right. \\ &\quad \left. + \hat{\mathbb{E}}\left[\sup_{0 \leq t \leq T} \left| \int_0^t (\sigma_i(s, X^{\epsilon_m}(s)) - \sigma_i(s, X(s))) dB_s^i \right|^2\right] \right). \end{aligned}$$

Applying BDG-inequality, we get

$$\begin{aligned} \hat{\mathbb{E}}\left[\sup_{0 \leq t \leq T} |X^{\epsilon_m}(t) - X(t)|^2\right] &\leq C(\epsilon_m^2 + \int_0^T \hat{\mathbb{E}}[|X^{\epsilon_m}(t) - X(t)|^2] dt) \\ &\leq C(\epsilon_m^2 + \int_0^T \hat{\mathbb{E}}[\sup_{0 \leq s \leq t} |X^{\epsilon_m}(s) - X(s)|^2] dt). \end{aligned}$$

By Gronwall's inequality, we have

$$\hat{\mathbb{E}}\left[\sup_{0 \leq t \leq T} |X^{\epsilon_m}(t) - X(t)|^2\right] \leq C|\epsilon_m|^2.$$

Up to a subsequence, still denoted as  $(X^{\epsilon_m})_{m \geq 1}$ , we have

$$\lim_{m \rightarrow \infty} X^{\epsilon_m}(t) = X(t) \quad q.s.$$

for all  $t \in [0, T]$ . This ends the proof. ■

**Remark 4.2** *In general, the condition (B1) in the above theorem can not be replaced by the following condition:*

(B1'')

$$\bar{b}_i(t, y) - b_i(t, x) + G([\bar{h}_{lk}]_i + (\bar{h}_{kl})_i]_{l,k=1}^d(t, y) - [(h_{lk})_i + (h_{kl})_i]_{l,k=1}^d(t, x) \geq 0$$

are fulfilled, whenever  $x_i = y_i$  and  $x_j \leq y_j$  for all  $j \neq i$ .

In fact, let  $b_1 = h_1 = \bar{b}_1 = \bar{h}_1 = h_2 = \bar{b}_2 = 0$ ,  $b_2 = \frac{\bar{\sigma}^2 + \underline{\sigma}^2}{2}$ ,  $\bar{h}_2 = 1$  and  $x = \bar{x} = 0$ , we consider 1-dimensional  $G$ -Brownian motion. One can show that  $b, h, \bar{b}, \bar{h}$  satisfy condition (B''). However we can not deduce that  $\frac{\bar{\sigma}^2 + \underline{\sigma}^2}{2}t \leq \langle B \rangle_t$ ,  $q.s.$  whenever  $\underline{\sigma}^2 < \bar{\sigma}^2$ .

## 5 Stochastic monotonicity and order-preservation

Lin ([20]) defined the infinitesimal generator of  $G$ -SDEs and obtained the representation theorem under the Lipschitz condition. Similarly, we can obtain the relationships among  $G$ -Itô's diffusions, Markov semigroups and infinitesimal generators. They can be summarized as follows. We suppose  $(X_t)_{t \geq 0}$  to be  $n$ -dimensional  $G$ -Itô diffusion

$$dX_t = X_0 + b(X_s)ds + h_{ij}(X_s)d\langle B^i, B^j \rangle_s + \sigma_j(X_s)dB_s^j, \quad t \in [0, T],$$

where  $(B_t)_{t \geq 0}$  is a  $d$ -dimensional  $G$ -Brownian motion and  $b, h, \sigma$  are Lipschitz continuous functions on  $\mathbb{R}^n$ . The Markov semigroup  $\mathcal{E}_t$  is defined by  $\mathcal{E}_t f(x) = \hat{\mathbb{E}}[f(X_t^{0,x})]$ , where  $X_t^{0,x}$  represents the  $G$ -Itô process with initial condition  $x$  at initial time  $t = 0$  and  $f$  is a function defined on  $\mathbb{R}^n$ . The infinitesimal generator  $L$  of the Markov semigroup, which satisfies

$$Lf(x) = \lim_{t \rightarrow 0^+} \frac{\mathcal{E}_t f(x) - f(x)}{t}$$

for  $f$  appropriately taken such that the above limit exists, is of the following form:

$$Lf = \langle \partial_x f, b \rangle + G(\langle \partial_x f, h \rangle + \langle \partial_{xx}^2 f \sigma, \sigma \rangle)$$

where  $\langle \partial_x f, h \rangle + \langle \partial_{xx}^2 f \sigma, \sigma \rangle$  is a  $d \times d$  symmetric matrix in  $\mathbb{S}^d(\mathbb{R})$ , defined by:

$$\begin{aligned} \langle \partial_x f, h \rangle + \langle \partial_{xx}^2 f \sigma, \sigma \rangle &:= [\langle \partial_x f, h_{ij} + h_{ji} \rangle + \langle \partial_{xx}^2 f \sigma_i, \sigma_j \rangle]_{i,j=1}^d, \\ L &= \sum_{i=1}^n b_i \frac{\partial}{\partial x_i} + G([\sum_{i=1}^n (h_{lk} + h_{kl})_i \frac{\partial}{\partial x_i} + \sum_{i,j=1}^n \sigma_{il} \sigma_{jk} \frac{\partial^2}{\partial x_i \partial x_j}]_{l,k=1}^d). \end{aligned}$$

Now we introduce the following definitions, which are similar to that in Herbst and Pitt ([11]) and Chen and Wang ([3]). Let " $\leq$ " denote the usual semi-order in  $\mathbb{R}^n$ .

(1) A measurable function  $f$  is called monotone if

$$f(x) \leq f(\bar{x}) \quad \text{for all } x \leq \bar{x}.$$

Denote by  $\mathcal{M}$  the set of all bounded Lipschitz continuous monotone functions.

(2) For two semigroups  $\{\mathcal{E}_t\}_{0 \leq t \leq T}$  and  $\{\bar{\mathcal{E}}_t\}_{0 \leq t \leq T}$ , we write  $\mathcal{E}_t \geq \bar{\mathcal{E}}_t$ , if for all  $f \in \mathcal{M}$ , for all  $x \geq \bar{x}$  and  $0 \leq t \leq T$ ,

$$\mathcal{E}_t f(x) \geq \bar{\mathcal{E}}_t f(\bar{x}).$$

If in addition,  $\mathcal{E}_t = \bar{\mathcal{E}}_t$ , we call  $\mathcal{E}_t$  monotone.

Let

$$Lf = \langle \partial_x f, b \rangle + G(\langle \partial_x f, h \rangle + \langle \partial_{xx}^2 f \sigma, \sigma \rangle),$$

$$\bar{L}f = \langle \partial_x f, \bar{b} \rangle + G(\langle \partial_x f, \bar{h} \rangle + \langle \partial_{xx}^2 f \bar{\sigma}, \bar{\sigma} \rangle),$$

$$\bar{L}'f = \langle \partial_x f, \bar{b} \rangle + G(\langle \partial_x f, \bar{h} \rangle + \langle \partial_{xx}^2 f \sigma, \sigma \rangle)$$

and let  $\{\mathcal{E}_t\}_{0 \leq t \leq T}$ ,  $\{\bar{\mathcal{E}}_t\}_{0 \leq t \leq T}$  and  $\{\bar{\mathcal{E}}_t'\}_{0 \leq t \leq T}$  be the semigroup generated by  $L, \bar{L}$  and  $\bar{L}'$  respectively. And we always assume that  $b, h_{ij}, \sigma_i$  and  $\bar{b}, \bar{h}_{ij}, \bar{\sigma}_i$  satisfy (H2) for each  $i, j = 1, \dots, d$ .

We have the following results.

**Theorem 5.1** *Suppose the following conditions hold:*

(C1) *for all  $i, j$ ,  $\sigma_{li}\sigma_{kj}$  depends only on  $x_i$  and  $x_j$ ,  $l, k = 1, \dots, d$ .*

(C2) *for all  $i$ ,  $b_i(x) - b_i(y) + G([\frac{(h_{l,k})_i(x) + (h_{k,l})_i(x)}{2m}]_{l,k=1}^d - [\frac{(h_{l,k})_i(y) + (h_{k,l})_i(y)}{2m}]_{l,k=1}^d) \leq 0$  whenever  $x \leq y$  with  $x_i = y_i$ .*

then  $\mathcal{E}_t$  is monotone

**Proof.** Suppose (C1) and (C2) hold. By setting  $\bar{b} = b$  and  $\bar{h} = h$  in Theorem (4.1), we have  $\forall x \leq \bar{x}$ ,  $X_t^{0,x} \leq X_t^{0,\bar{x}}$  q.s.. Then by the monotonicity of  $f$ , the results follows. ■

**Theorem 5.2** *If  $\mathcal{E}_t$  is monotone, then the following conditions hold:*

(C1) *for all  $i, j$ ,  $\sigma_{li}\sigma_{kj}$  depends only on  $x_i$  and  $x_j$ ,  $l, k = 1, \dots, d$ .*

(C2') *for all  $i$ ,  $b_i(x) - b_i(y) + G([\frac{(h_{l,k})_i(x) + (h_{k,l})_i(x)}{2m}]_{l,k=1}^d - [\frac{(h_{l,k})_i(y) + (h_{k,l})_i(y)}{2m}]_{l,k=1}^d) \geq 0$  whenever  $x \geq y$  with  $x_i = y_i$ .*

**Proof.** Suppose that  $\mathcal{E}_t$  is monotone,

$$\lim_{t \rightarrow 0^+} \frac{1}{t} (\mathcal{E}_t f(x) - f(x)) = \langle \partial_x f, b \rangle + G(\langle \partial_x f, h \rangle + \langle \partial_{xx}^2 f \sigma, \sigma \rangle). \quad (8)$$

(a) For given  $i$ , we take  $x^{(1)} \leq x^{(2)}$  with  $x_i^{(1)} = x_i^{(2)}$  and a sequence of functions  $f_m \in \mathcal{M} \cap C_b^\infty$  ( $m \in \mathbb{N}$ ) such that  $f_m(x) = (x_i - x_i^{(1)} + 1)^{2m+1}$  in a neighborhood of  $\{x^{(1)}, x^{(2)}\}$ . We have  $\lim_{t \rightarrow 0} \frac{1}{t} (\mathcal{E}_t f_m(x^{(1)}) - f_m(x^{(1)})) \leq \lim_{t \rightarrow 0} \frac{1}{t} (\mathcal{E}_t f_m(x^{(2)}) - f_m(x^{(2)}))$ , i.e.,

$$\begin{aligned} & \langle \partial_x f(x^{(1)}), b(x^{(1)}) \rangle + G(\langle \partial_x f(x^{(1)}), h(x^{(1)}) \rangle + \langle \partial_{xx}^2 f(x^{(1)}) \sigma(x^{(1)}), \sigma(x^{(1)}) \rangle) \\ & \leq \langle \partial_x f(x^{(2)}), b(x^{(2)}) \rangle + G(\langle \partial_x f(x^{(2)}), h(x^{(2)}) \rangle + \langle \partial_{xx}^2 f(x^{(2)}) \sigma(x^{(2)}), \sigma(x^{(2)}) \rangle). \end{aligned}$$

Then we get

$$\begin{aligned} & G([\frac{(h_{l,k})_i(x^{(2)}) + (h_{k,l})_i(x^{(2)})}{2m} + \sigma_{li}(x^{(2)})\sigma_{ki}(x^{(2)})]_{l,k=1}^d \\ & \quad - [\frac{(h_{l,k})_i(x^{(1)}) + (h_{k,l})_i(x^{(1)})}{2m} + \sigma_{li}(x^{(1)})\sigma_{ki}(x^{(1)})]_{l,k=1}^d) \\ & \geq G([\frac{(h_{l,k})_i(x^{(2)}) + (h_{k,l})_i(x^{(2)})}{2m} + \sigma_{li}(x^{(2)})\sigma_{ki}(x^{(2)})]_{j,k=1}^d \\ & \quad - G([\frac{(h_{l,k})_i(x^{(1)}) + (h_{k,l})_i(x^{(1)})}{2m} + \sigma_{li}(x^{(1)})\sigma_{ki}(x^{(1)})]_{l,k=1}^d) \\ & \geq \frac{1}{2m} [b_i(x^{(1)}) - b_i(x^{(2)})]. \end{aligned}$$

So  $[\sigma_{li}(v)\sigma_{ki}(v)]_{l,k=1}^d \geq [\sigma_{li}(u)\sigma_{ki}(u)]_{l,k=1}^d$  as  $m \rightarrow \infty$ . Replacing  $f_m$  with  $(x_i - x_i^{(1)} - 1)^{2m+1}$  in a neighborhood of  $\{x^{(1)}, x^{(2)}\}$ , we obtain the inverse inequality. Therefore,  $[\sigma_{li}(v)\sigma_{ki}(v)]_{l,k=1}^d = [\sigma_{li}(u)\sigma_{ki}(u)]_{l,k=1}^d$ .

(b) For given  $i, j$   $i \neq j$ , we take  $x^{(1)} \leq x^{(2)}$  with  $x_i^{(1)} = x_i^{(2)}$ ,  $x_j^{(1)} = x_j^{(2)}$  and a sequence of functions  $f_m \in \mathcal{M} \cap C_b^\infty$  ( $m \in \mathbb{N}$ ) such that  $f_m(x) = (x_i + x_j - x_i^{(1)} - x_j^{(1)} + 1)^{2m+1}$  in a neighborhood of  $\{x^{(1)}, x^{(2)}\}$ . We have  $\lim_{t \rightarrow 0} \frac{1}{t} (\mathcal{E}_t f_m(x^{(1)}) - f_m(x^{(1)})) \leq \lim_{t \rightarrow 0} \frac{1}{t} (\mathcal{E}_t f_m(x^{(2)}) - f_m(x^{(2)}))$ , i.e.,

$$\begin{aligned} & \langle \partial_x f(x^{(1)}), b(x^{(1)}) \rangle + G(\langle \partial_x f(x^{(1)}), h(x^{(1)}) \rangle + \langle \partial_{xx}^2 f(x^{(1)}) \sigma(x^{(1)}), \sigma(x^{(1)}) \rangle) \\ & \leq \langle \partial_x f(x^{(2)}), b(x^{(2)}) \rangle + G(\langle \partial_x f(x^{(2)}), h(x^{(2)}) \rangle + \langle \partial_{xx}^2 f(x^{(2)}) \sigma(x^{(2)}), \sigma(x^{(2)}) \rangle). \end{aligned}$$

Then we get

$$\begin{aligned}
& G\left(\left[\frac{(h_{l,k})_i(x^{(2)}) + (h_{k,l})_i(x^{(2)}) + (h_{l,k})_j(x^{(2)}) + (h_{k,l})_j(x^{(2)})}{2m} + \sigma_{li}(x^{(2)})\sigma_{kj}(x^{(2)})\right]_{l,k=1}^d\right. \\
& \quad \left. - \left[\frac{(h_{l,k})_i(x^{(1)}) + (h_{k,l})_i(x^{(1)}) + (h_{l,k})_j(x^{(1)}) + (h_{k,l})_j(x^{(1)})}{2m} + \sigma_{li}(x^{(1)})\sigma_{kj}(x^{(1)})\right]_{l,k=1}^d\right) \\
& \geq G\left(\left[\frac{(h_{l,k})_i(x^{(2)}) + (h_{k,l})_i(x^{(2)}) + (h_{l,k})_j(x^{(2)}) + (h_{k,l})_j(x^{(2)})}{2m} + \sigma_{li}(x^{(2)})\sigma_{kj}(x^{(2)})\right]_{j,k=1}^d\right) \\
& \quad - G\left(\left[\frac{(h_{l,k})_i(x^{(1)}) + (h_{k,l})_i(x^{(1)}) + (h_{l,k})_j(x^{(1)}) + (h_{k,l})_j(x^{(1)})}{2m} + \sigma_{li}(x^{(1)})\sigma_{kj}(x^{(1)})\right]_{l,k=1}^d\right) \\
& \geq \frac{1}{2m}[b_i(x^{(1)}) + b_j(x^{(1)}) - b_i(x^{(2)}) - b_j(x^{(2)})].
\end{aligned}$$

So  $[\sigma_{li}(v)\sigma_{kj}(v)]_{l,k=1}^d \geq [\sigma_{li}(u)\sigma_{kj}(u)]_{l,k=1}^d$  as  $m \rightarrow \infty$ . Replacing  $f_m$  with  $(x_i + x_j - x_i^{(1)} - x_j^{(1)} + 1)^{2m+1}$  in a neighborhood of  $\{x^{(1)}, x^{(2)}\}$ , we obtain the inverse inequality. Therefore,  $[\sigma_{li}(v)\sigma_{kj}(v)]_{l,k=1}^d = [\sigma_{li}(u)\sigma_{kj}(u)]_{l,k=1}^d$ . Thus (C1) holds.

- (c) For given  $i$ , we take  $x^{(1)} \leq x^{(2)}$  with  $x_i^{(1)} = x_i^{(2)}$  and  $f_m \in \mathcal{M} \cap C_b^\infty$  ( $m \in \mathbb{N}$ ) such that  $f(x) = x_i$  in a neighborhood of  $\{x^{(1)}, x^{(2)}\}$ . By (8), we have

$$b_i(x^{(1)}) + G([\sigma_{li}(v)\sigma_{kj}(v)]_{l,k=1}^d) \leq b_i(x^{(2)}) + G([\sigma_{li}(u)\sigma_{kj}(u)]_{l,k=1}^d).$$

Thus by the subadditivity, (C2') holds.

■

**Theorem 5.3** *If  $\mathcal{E}_t \geq \bar{\mathcal{E}}_t$  then the following two conditions hold:*

- (D1) *for all  $i, j$ ,  $\sigma_{il}\sigma_{jk} \equiv \bar{\sigma}_{il}\bar{\sigma}_{jk}$  and  $\sigma_{il}\sigma_{jk}$  depends only on  $x_i$  and  $x_j$ ,  $l, k = 1, \dots, d$ .*  
(D2) *for all  $i$ ,  $b_i(x) - \bar{b}_i(y) + G([\sigma_{li}(x)\sigma_{kj}(x)]_{l,k=1}^d - [\bar{\sigma}_{li}(y)\bar{\sigma}_{kj}(y)]_{l,k=1}^d) \geq 0$  whenever  $x \geq y$  with  $x_i = y_i$ .*

To prove the above theorem, we need some Lemmas.

**Lemma 5.4** *If  $\mathcal{E}_t \geq \bar{\mathcal{E}}_t$ , then  $Lf(x) \geq \bar{L}(y)$  for all  $x \geq y$  and  $f \in \mathcal{M} \cap C_b^\infty$  with  $f(x) = f(y)$ .*

**Proof.** Without loss of generality, assume that  $f \geq 0$ . Choose  $m > 0$  such that  $\{z : |z| < m\}$  contains  $x$  and  $y$  and take  $h \in C_b^\infty$  such that

$$1 \geq h(z) = \begin{cases} 1, & \text{if } |z| \leq m, \\ 0, & \text{if } |z| \geq m + 1, \\ > 0, & \text{otherwise.} \end{cases}$$

Set

$$f_1 = hf + a(1 - h), \quad f_2 = hf,$$

where  $a$  is a constant larger than the upper bound of  $f$ . Then  $f_1, f_2 \in C_0^\infty$ ,  $f_1 \geq f \geq f_2$  and  $f_1 = f = f_2$  on the set  $\{z : |z| < m\}$ . Since

$$\mathcal{E}_t f(x) \geq \bar{\mathcal{E}}_t f(y), \quad f(x) = f(y),$$

we have

$$\frac{1}{t}[\mathcal{E}_t f_1(x) - f_1(x)] \geq \frac{1}{t}[\bar{\mathcal{E}}_t f_2(y) - f_2(y)].$$

The assertion now follows by letting  $t \downarrow 0$ . ■



**Lemma 5.5** *If  $\mathcal{E}_t \geq \bar{\mathcal{E}}_t$ , then (D2) holds.*

**Proof.** For given  $i$ , let  $u \leq v$  with  $u_i = v_i$ . Choose  $f \in \mathcal{M} \cap C_b^\infty$  such that in a neighborhood of  $\{u, v\}$ ,

$$f(x) = x_i.$$

Then by Lemma (5.4), we get

$$b_i(v) - \bar{b}_i(u) + G([\frac{(h_{l,k})_i(v) + (h_{k,l})_i(v)}{2m}]_{l,k=1}^d - [\frac{(\bar{h}_{l,k})_i(u) + (\bar{h}_{k,l})_i(u)}{2m}]_{l,k=1}^d) \geq 0.$$

■

**Lemma 5.6** *If  $\mathcal{E}_t \geq \bar{\mathcal{E}}_t$ , then (D1) holds.*

**Proof.** The proof consists of two steps.

- (1) For given  $i$ , let  $u \leq v$  with  $u_i = v_i$ . Choose  $f_m \in \mathcal{M} \cap C_b^\infty$  ( $m \in \mathbb{N}$ ) such that in a neighborhood of  $\{u, v\}$ ,

$$f_m(x) = (x_i - u_i + 1)^{2m+1}.$$

By Lemma (5.4), we have

$$\begin{aligned} & G([\frac{(h_{l,k})_i(v) + (h_{k,l})_i(v)}{2m} + \sigma_{il}(v)\sigma_{ik}(v)]_{l,k=1}^d - [\frac{(\bar{h}_{l,k})_i(u) + (\bar{h}_{k,l})_i(u)}{2m} + \bar{\sigma}_{il}(u)\bar{\sigma}_{ik}(u)]_{l,k=1}^d) \\ & \geq G([\frac{(h_{l,k})_i(v) + (h_{k,l})_i(v)}{2m} + \sigma_{il}(v)\sigma_{ik}(v)]_{j,k=1}^d) - G([\frac{(\bar{h}_{l,k})_i(u) + (\bar{h}_{k,l})_i(u)}{2m} + \bar{\sigma}_{il}(u)\bar{\sigma}_{ik}(u)]_{l,k=1}^d) \\ & \geq \frac{1}{2m}[\bar{b}_i(u) - b_i(v)]. \end{aligned}$$

Since  $m$  is arbitrary, we deduce that:

$$[\sigma_{il}(v)\sigma_{ik}(v)]_{l,k=1}^d \geq [\bar{\sigma}_{il}(u)\bar{\sigma}_{ik}(u)]_{l,k=1}^d.$$

Replacing  $f_m$  with  $(x_i - u_i - 1)^{2m+1}$  in the neighborhood of  $\{u, v\}$ , we obtain the inverse inequality. Therefore  $\sigma_{il}(v)\sigma_{ik}(v) = \bar{\sigma}_{il}(u)\bar{\sigma}_{ik}(u)$ .

- (2) For given  $i \neq j$  and  $u \leq v$ :  $u_i = v_i$ ,  $u_j = v_j$ , choose  $f_m \in \mathcal{M} \cap C_b^\infty$  ( $m \in \mathbb{N}$ ) such that in a neighborhood of  $\{u, v\}$ ,

$$f_m(x) = (x_i + x_j - u_i - u_j + 1)^{2m+1}.$$

By (1) and Lemma (5.4), we get

$$\begin{aligned} & G([\frac{(h_{l,k})_i(v) + (h_{k,l})_i(v) + h_{l,k}j(v) + (h_{k,l})_j(v)}{2m} + \sigma_{il}(v)\sigma_{jk}(v)]_{l,k=1}^d \\ & \quad - [\frac{(\bar{h}_{l,k})_i(u) + (\bar{h}_{k,l})_i(u) + \bar{h}_{l,k}j(u) + (\bar{h}_{k,l})_j(u)}{2m} + \bar{\sigma}_{il}(u)\bar{\sigma}_{jk}(u)]_{l,k=1}^d) \\ & \geq G([\frac{(h_{l,k})_i(v) + (h_{k,l})_i(v) + (h_{l,k})_i(v) + (h_{k,l})_j(v)}{2m} + \sigma_{il}(v)\sigma_{jk}(v)]_{j,k=1}^d) \\ & \quad - G([\frac{(\bar{h}_{l,k})_i(u) + (\bar{h}_{k,l})_i(u) + (\bar{h}_{l,k})_i(u) + (\bar{h}_{k,l})_j(u)}{2m} + \bar{\sigma}_{il}(u)\bar{\sigma}_{jk}(u)]_{l,k=1}^d) \\ & \geq \frac{1}{2m}[\bar{b}_i(u) + \bar{b}_j(u) - b_i(v) - b_j(v)] \end{aligned}$$

and

$$[\sigma_{il}(v)\sigma_{jk}(v)]_{l,k=1}^d \geq [\bar{\sigma}_{il}(u)\bar{\sigma}_{jk}(u)]_{l,k=1}^d.$$

Similarly, we have the inverse inequality and hence  $\sigma_{il}(v)\sigma_{jk}(v) = \bar{\sigma}_{il}(u)\bar{\sigma}_{jk}(u)$ .

■

**Proof of theorem 5.3.** The proof follows directly from Lemma(5.5) and Lemma (5.6). ■

**Corollary 5.7** If  $\mathcal{E}_t \geq \bar{\mathcal{E}}'_t$  then the following two conditions hold:

(D3) for all  $i, j$ ,  $\sigma_{il}\sigma_{jk}$  depends only on  $x_i$  and  $x_j$ ,  $l, k = 1, \dots, d$ .

(D4) for all  $i$ ,  $b_i(x) - \bar{b}_i(y) + G([\!(h_{l,k})_i(x) + (h_{k,l})_i(x)\!]_{l,k=1}^d - [\!(\bar{h}_{l,k})_i(y) + (\bar{h}_{k,l})_i(y)\!]_{l,k=1}^d) \geq 0$  whenever  $x \geq y$  with  $x_i = y_i$ .

**Corollary 5.8** Suppose the following two conditions hold:

(D3) for all  $i, j$ ,  $\sigma_{il}\sigma_{jk}$  depends only on  $x_i$  and  $x_j$ ,  $l, k = 1, \dots, d$ .

(D4') for all  $i$ ,  $\bar{b}_i(x) - b_i(y) + G([\!(\bar{h}_{l,k})_i(x) + (\bar{h}_{k,l})_i(x)\!]_{l,k=1}^d - [\!(h_{l,k})_i(y) + (h_{k,l})_i(y)\!]_{l,k=1}^d) \leq 0$  whenever  $x \leq y$  with  $x_i = y_i$ .

then  $\mathcal{E}_t \geq \bar{\mathcal{E}}'_t$ .

**Proof.** Suppose (D3) and (D4') hold. By Theorem (4.1), we have  $\forall x \geq \bar{x}$ ,  $X_t^{0,x} \geq \bar{X}_t^{0,\bar{x}}$  q.s.. Then by the monotonicity of  $f$ , the results follows. ■

**Corollary 5.9** If  $\mathcal{E}_t \geq \bar{\mathcal{E}}_t$  then the following hold:

(D1) for all  $i, j$ ,  $\sigma_{il}\sigma_{jk} \equiv \bar{\sigma}_{il}\bar{\sigma}_{jk}$  and  $\sigma_{il}\sigma_{jk}$  depends only on  $x_i$  and  $x_j$ ,  $l, k = 1, \dots, d$

(D2') for all  $x, K \in \mathbb{R}^n$ ,  $K \geq 0$ ,  $K^*(b(x) - \bar{b}(x)) + G([\!K^*((h_{l,k})(x) + (h_{k,l})(x))\!]_{l,k=1}^d - [K^*((\bar{h}_{l,k})_i(x) + (\bar{h}_{k,l})_i(x))\!]_{l,k=1}^d) \geq 0$ .

**Proof.** First, we suppose that  $\mathcal{E}_t \geq \bar{\mathcal{E}}_t$ . Then (D1) holds directly from Theorem (5.3). For fixed  $\bar{x} \in \mathbb{R}^n$ ,  $K \in \mathbb{R}^n$ ,  $K \geq 0$ , we take  $f \in \mathcal{M} \cap C_b^\infty$  such that in a neighborhood of  $\bar{x}$ ,  $f(x) = \sum_{i=1}^n K_i(x_i - \bar{x}_i)$ . Then we have  $\lim t \rightarrow 0 \frac{1}{t}(\mathcal{E}_t f(\bar{x}) - f(\bar{x})) \geq \lim t \rightarrow 0 \frac{1}{t}(\bar{\mathcal{E}}_t f(\bar{x}) - f(\bar{x}))$ . Thus (D2') holds. ■

**Theorem 5.10** Assume (H3) and assume that  $\sigma\sigma^*$  (or resp.  $\bar{\sigma}\bar{\sigma}^*$ ) is uniformly positive definite, i.e., there exists a constant  $\beta > 0$ , such that for all  $y \in \mathbb{R}^n$ ,  $x \in \mathbb{R}^n$ ,  $y^*\sigma(x)\sigma^*(x)y \geq \beta|y|^2$ . If one of  $\mathcal{E}_t$  and  $\bar{\mathcal{E}}_t$  is monotone, if the following hold:

(D1) for all  $i, j$ ,  $\sigma_{il}\sigma_{jk} \equiv \bar{\sigma}_{il}\bar{\sigma}_{jk}$  and  $\sigma_{il}\sigma_{jk}$  depends only on  $x_i$  and  $x_j$ ,  $l, k = 1, \dots, d$

(D5) for all  $x, K \in \mathbb{R}^n$ ,  $K \geq 0$ ,  $K^*(\bar{b}(x) - b(x)) + G([\!K^*((\bar{h}_{l,k})(x) + (\bar{h}_{k,l})(x))\!]_{l,k=1}^d - [K^*((h_{l,k})_i(x) + (h_{k,l})_i(x))\!]_{l,k=1}^d) \leq 0$ .

then  $\mathcal{E}_t \geq \bar{\mathcal{E}}_t$ .

**Proof.** We now suppose (D1) and (D5) hold. Without lost of generalization, we assume  $\bar{\mathcal{E}}$  is monotone. Here we denote  $(\sigma\sigma^*)^{-1}\sigma$  by  $\Sigma$ . Let  $f \in \mathcal{M}$  and we consider the following  $G$ -SDE and  $G$ -BSDEs:

$$\bar{X}_t^{0,x} = x + \int_0^t \sigma_i(\bar{X}_r^{0,x}) dB_r^i, \quad t \in [0, T],$$

$$\begin{aligned} Y_s^{0,x,t,f} &= f(\bar{X}_t^{0,x}) + \int_s^t b^*(\bar{X}_r^{0,x}) \Sigma(\bar{X}_r^{0,x}) Z_r^{0,x,t,f} dr + \int_s^t h_{ij}^*(\bar{X}_r^{0,x}) \Sigma(\bar{X}_r^{0,x}) Z_r^{0,x,t,f} d\langle B^i, B^j \rangle_r \\ &\quad - \int_s^t Z_r^{0,x,t,f} dB_r - (K_s - K_t), \quad s \in [0, t] \end{aligned}$$

and

$$\begin{aligned} \bar{Y}_s^{0,x,t,f} &= f(\bar{X}_t^{0,x}) + \int_s^t \bar{b}^*(\bar{X}_r^{0,x}) \Sigma(\bar{X}_r^{0,x}) \bar{Z}_r^{0,x,t,f} dr + \int_s^t \bar{h}_{ij}^*(\bar{X}_r^{0,x}) \Sigma(\bar{X}_r^{0,x}) \bar{Z}_r^{0,x,t,f} d\langle B^i, B^j \rangle_r \\ &\quad - \int_s^t \bar{Z}_r^{0,x,t,f} dB_r - (\bar{K}_s - \bar{K}_t), \quad s \in [0, t]. \end{aligned}$$

We have the following results:  $u(t, x) := \mathcal{E}_t f(x) = Y_0^{0,x,t,f}$  is the unique viscosity solution of the following PDE:

$$\begin{cases} \partial_t u - Lu = 0, \\ u(0, x) = f(x). \end{cases} \quad (9)$$

$\bar{u}(t, x) := \bar{\mathcal{E}}_t = \bar{Y}_0^{0,x,t,f}$  is the unique viscosity solution of the following PDE:

$$\begin{cases} \partial_t \bar{u} - \bar{L}\bar{u} = 0, \\ \bar{u}(0, x) = f(x). \end{cases} \quad (10)$$

By Theorem 6.4.3 in Krylov [17] (see also Theorem 4.4 in Appendix C in Peng [31]), there exists a constant  $\alpha \in (0, 1)$  such that for each  $\kappa > 0$ ,

$$\|\bar{u}\|_{C^{1+\alpha/2, 2+\alpha}([0, T] \times \mathbb{R}^n)} < \infty.$$

Let  $\hat{u}(t, x) = \bar{u}(T - t, x)$  and apply  $G$ -Itô's formula to  $\hat{u}(s, \bar{X}_s)$  for  $s \in [0, T - \kappa]$

$$\begin{aligned} \hat{u}(s, \bar{X}_s^{0,x}) &= \hat{u}(T - \kappa, \bar{X}_{T-\kappa}^{0,x}) + \int_s^{T-\kappa} \bar{b}^*(\bar{X}_r^{0,x}) \Sigma(\bar{X}_r^{0,x}) \sigma^*(\bar{X}_r^{0,x}) \partial_x \hat{u}(r, \bar{X}_r^{0,x}) dr \\ &\quad + \int_s^{T-\kappa} \bar{h}_{ij}^*(\bar{X}_r^{0,x}) \Sigma(\bar{X}_r^{0,x}) \sigma^*(\bar{X}_r^{0,x}) \partial_x \hat{u}(r, \bar{X}_r^{0,x}) d\langle B^i, B^j \rangle_r \\ &\quad - \int_s^{T-\kappa} \sigma^*(\bar{X}_r^{0,x}) \partial_x \hat{u}(r, \bar{X}_r^{0,x}) dB_r - (K'_{T-\kappa} - K'_s) \end{aligned}$$

where

$$\begin{aligned} K_s &= \frac{1}{2} \int_0^s \sigma^*(\bar{X}_r^{0,x}) \partial_{xx}^2 \hat{u}(r, \bar{X}_r^{0,x}) \sigma(\bar{X}_r^{0,x}) d\langle B \rangle_r + \int_s^{T-\kappa} \bar{h}^*(\bar{X}_r^{0,x}) \Sigma(\bar{X}_r^{0,x}) \sigma^*(\bar{X}_r^{0,x}) \partial_x \hat{u}(r, \bar{X}_r^{0,x}) d\langle B \rangle_r \\ &\quad - \int_0^s G(\langle \partial_x \hat{u}(r, \bar{X}_r^{0,x}), \bar{h}^*(\bar{X}_r^{0,x}) \rangle + \langle \partial_{xx}^2 \hat{u}(r, \bar{X}_r^{0,x}) \sigma(\bar{X}_r^{0,x}), \sigma(\bar{X}_r^{0,x}) \rangle) dr \end{aligned}$$

is a non-increasing  $G$ -martingale. By the uniqueness of solutions of  $G$ -BSDEs,  $\bar{Z}_r^{0,x,t,f} = \sigma^*(X_r^{0,x}) \partial_x \hat{u}(t, X_t^{0,x})$  for all  $0 \leq r \leq t \leq T - \kappa$ . By the assumption that  $\bar{\mathcal{E}}_t$  is monotone, we have  $\bar{u}$  (then  $\hat{u}$ ) is nondecreasing in  $x$ . Thus  $\partial_x \hat{u} \geq 0$ . By condition (D5) and the comparison of  $G$ -BSDEs, we have  $Y_r^{0,x,t,f} \geq \bar{Y}_r^{0,x,t,f}$  for all  $0 \leq r \leq t \leq T - \kappa$ . Particularly, we have  $\mathcal{E}_t f(x) \geq \bar{\mathcal{E}}_t f(x)$ . By the monotonicity of  $\bar{\mathcal{E}}_t$ , for all  $x \geq \bar{x}$ ,  $\mathcal{E}_t f(x) \geq \bar{\mathcal{E}}_t f(x) \geq \bar{\mathcal{E}}_t f(\bar{x})$ . Thus  $\mathcal{E}_t \geq \bar{\mathcal{E}}_t$  for all  $t \in [0, T]$ . By continuity, we have  $\mathcal{E}_t \geq \bar{\mathcal{E}}_t$  for all  $t \in [0, T]$ . ■

## 6 Applications to PDEs

In this section, we will give some applications of the above results to a special type of PDEs. We assume that  $b, h_{ij}, \sigma_i$  and  $\bar{b}, \bar{h}_{ij}, \bar{\sigma}_i$  satisfy (H2) for each  $i, j = 1, \dots, d$ . We have the following results. We will omit the proofs since the results hold in view of Theorem 5.1, Theorem 5.2, Corollary 5.9, Theorem 5.10 and the nonlinear Feynman-Kac formula in section 3.3.

**Theorem 6.1** Let  $u$  be the unique viscosity solution of PDE (9), if (C1) and (C2) hold, then  $u$  is nondecreasing in  $x$  for all  $f \in \mathcal{M}$ .

**Theorem 6.2** Let  $u$  be the unique viscosity solution of PDE (9), if  $u$  is nondecreasing in  $x$  for all  $f \in \mathcal{M}$ , then (C1) and (C2') hold.

**Theorem 6.3** Let us consider PDE (9) and PDE (10), if  $u(t, x) \geq \bar{u}(t, \bar{x})$  for all  $t \in [0, T]$ ,  $f \in \mathcal{M}$  and  $x \geq \bar{x}$ , then (D1), (D2), and (D2') hold.

**Theorem 6.4** Assume (H3) and assume that  $\sigma\sigma^*$  (or resp.  $\bar{\sigma}\bar{\sigma}^*$ ) is uniformly positive definite. Suppose that (C1) and (C2) hold for  $b, h, \sigma$  or  $\bar{b}, \bar{h}, \bar{\sigma}$ . Let us consider PDE (9) and PDE (10), if (D1) and (D5) hold, then  $u(t, x) \geq \bar{u}(t, \bar{x})$  for all  $t \in [0, T]$ ,  $f \in \mathcal{M}$  and  $x \geq \bar{x}$ .

## Acknowledgment

## References

- [1] Anderson, W.J. (1972) Local Behaviour of solutions of stochastic integral equations, *Trans. Amer. Math. Soc.* **164** 309-321.
- [2] Bai, X. and Lin Y. (2010) On the existence and uniqueness of solutions to stochastic differential equations driven by  $G$ -Brownian motion with integral-Lipschitz coefficients, in arXiv:1002.1046.
- [3] CHEN, M. and WANG, F. (1993) On order-preservation and positive correlations for multidimensional diffusion processes, *Probability Theory and Related Fields* **95** 421-428.
- [4] Cox, J.T. (1984) An alternative proof of a correlation inequality of Harris, *Annals of Probability*. **12**, 272-273.
- [5] DENIS, L., HU, M. and PENG, S. (2011) Function spaces and capacity related to a sublinear expectation: application to  $G$ -Brownian motion pathes. *Potential Analysis*. **34(2)** 139-161.
- [6] Gal'cuk, L.I. and Davis, M.H.A. (1982) A note on a comparison theorem for equations with differential equations, *Stochastics*. **6** 147-149.
- [7] GAO, F. (2009) Pathwise properties and homomorphic flows for stochastic differential equations driven by  $G$ -Brownian motion, *Stochastic Processes and their Applications*. **119** 3356-3382.
- [8] GEIß, G. and MANTHEY, R. (1994) Comparison theorems for stochastic differential equations in finite and infinite dimensions, *Stochastic Processes and their Applications*. **53**, 23-35.
- [9] Gihman, I.V. and Skorohod, A.V. (1972) Stochastic differential equations, Springer-Verlag, Berlin.
- [10] Harris, T.E. (1977) A correlation inequality for Markov processes in partially ordered state spaces, *Annals of Probability*. **5**, 451-454.
- [11] HERBST, I. and Pitt, L. (1991) Diffusion equation techniques in stochastic monotonicity and positive correlations *Probability Theory and Related Fields*. **87** 275-312.
- [12] He, K. and Hu, M. (2013) Representation theorem for generators of BSDEs driven by  $G$ -Brownian motion and its applications, *Abstract and Applied Analysis*, vol. 2013, Article ID 342038, 10 pages. doi:10.1155/2013/342038.

- [13] Holley, R. (1974) Remarks on the FKG inequality, *Commun. Math. Phys.* **36**, 227-231.
- [14] Hu, M., Ji, S., Peng, S. and Song, Y. (2014) Backward stochastic differential equations driven by  $G$ -Brownian motion, *Stochastic Processes and their Applications* 124:759-784.
- [15] Hu, M., Ji, S., Peng S., and Song, Y. (2014) Comparison theorem, Feynman-Kac formula and Girsanov transformation for BSDEs driven by  $G$ -Brownian motion, *Stochastic Processes and their Applications* 124:1170-1195.
- [16] Ikeda, N. and Watababe, S. (1977) A comparison theorem for solutions of stochastic differential equations and its applications, *Osaka J. Math.* **14** 619-633.
- [17] Krylov, N.V. (1987) Nonlinear Parabolic and Elliptic Equations of the Second Order, Reidel Publishing Company. (Original Russian Version by Nauka, Moscow, 1985).
- [18] Ladde, G.S. and Lakshmikantham, V. (1980) Random Differential Inequalities, Academic Press, New York.
- [19] LI, X. and PENG, S. (2011) Stopping times and related Itô calculus with  $G$ -Brownian motion, *Stochastic Processes and their Applications.* **121**, 1492–1508.
- [20] LIN, X. (2013) The Stability of Stochastic Systems Driven by  $G$ -Brownian motion. Doctoral thesis. Shandong University.
- [21] LIN, Q. (2013) Some properties of stochastic differential equations driven by  $G$ -Brownian motion. *Acta Mathematica Sinica, English Series.* Vol. 29, 923–942.
- [22] LIN, Q. (2013) Differentiability of stochastic differential equations driven by  $G$ -Brownian motion. *Science China Mathematics.* Vol. 56, 1087–1107.
- [23] LIN, Y. (2013) Stochastic differential equations driven by  $G$ -Brownian motion with reflecting boundary. *Electron. J. Probab.* **18**: **9**, 1–23.
- [24] LUO, P. and WANG, F. (2014) Stochastic differential equations driven by  $G$ -Brownian motion and ordinary differential equations, *Stochastic Processes and their Applications.* **124**, 3869–3885.
- [25] PENG, S. (1991) Probabilistic interpretation for systems of quasilinear parabolic partial differential equation. *Stochastics Stochastics Rep.* **37** 61–74.
- [26] PENG, S. (1992) A nonlinear Feynman-Kac formula and applications. *Control theory, stochastic analysis and applications (Hangzhou, 1991)*. 173–184. World Sci. Publ., River Edge, NJ.
- [27] PENG, S. (1997) BSDE and related  $g$ -expectation. In: El Karoui, N., Mazliak, L. (Eds.), Backward Stochastic Differential Equations. In: Pitman Research Notes in Mathematics Series, Vol. 264. pp. 141-159.
- [28] PENG, S. (2004) Filtration consistent nonlinear expectations and evaluations of contingent claims, *Acta Mathematicae Applicatae Sinica, English Series.* **20(2)** 1–24.
- [29] Peng, S. (2005) *Nonlinear expectations and nonlinear Markov chains*, *Chin. Ann. Math.* **26B(2)** 159–184.
- [30] Peng, S. (2009) *Survey on normal distributions, central limit theorem, Brownian motion and the related stochastic calculus under sublinear expectations*, *Science in China Series A: Mathematics.* **52(7)**: 1391-1411.

- [31] PENG, S. (2010) Nonlinear expectations and stochastic calculus under uncertainty. Preprint (arxiv:1002.4546v1). 2010.
- [32] PENG, S. (2010) Backward stochastic differential equation, nonlinear expectation and their applications. *Proceedings of the International Congress of Mathematicians Hyderabad*. India.
- [33] PENG, S. (2011). Note on viscosity solution of path-dependent PDE and G-martingales lecture notes. Preprint (arxiv:1106.1144v2).
- [34] REVUZ, D. and YOR, M. (1999) Continuous Martingales and Brownian Motion, 3rd ed. Springer, Berlin.
- [35] ROGERS, L.C.G. and WILLIAMS, D. (1994) Diffusions, Markov Processes, and Martingales, second ed.. In: Foundations, John Wiley & Sons Ltd., Chichester.
- [36] Stroock, D.W. and VARADHAN, S.R.S. (1982) Multidimensional Diffusion Equations. Springer, Berlin Heidelberg New York.
- [37] ZHANG, N. and JIA, G. (2013) Stochastic monotonicity and order-preservation for a type of nonlinear semigroups, *Statistic and Probability Letters*. **83** 422-429.