

Relativistic corrections to the Kompaneets equation

Satoshi Nozawa

Josai Junior College, 1-1 Keyakidai, Sakado-shi, Saitama, 350-0295, Japan

Yasuharu Kohyama

Advanced Simulation Technology of Mechanics Co. Ltd, 2-3-13 Minami, Wako-shi, Saitama, 351-0104, Japan

Abstract

We study the Sunyaev–Zeldovich effect for clusters of galaxies. We explore the relativistic corrections to the Kompaneets equation in terms of two different expansion approximation schemes, namely, the Fokker-Planck expansion approximation and delta function expansion approximation. We show that two expansion approximation formalisms are equivalent under the Thomson approximation, which is extremely good approximation for the CMB photon energies. This will clarify the situation for existing theoretical methods to analyse observation data.

Keywords: Sunyaev-Zeldovich effect, cosmic microwave background, galaxy clusters, relativistic corrections

1. Introduction

The Sunyaev–Zeldovich (SZ) effect [1, 2], which arises from the Compton scattering of the cosmic microwave background (CMB) photons by hot electrons in clusters of galaxies (CG), provides a useful method for studies of cosmology. For the reviews, for example, see Refs. [3] and [4]. The original SZ formula has been derived from the Kompaneets equation [5] in the non-relativistic approximation. However, X-ray observations, for example, by Allen et al. [6] have revealed the existence of high-temperature CG such as $k_B T_e \simeq 20\text{keV}$. For such high-temperature CG, the relativistic corrections will become important.

The theoretical studies on the relativistic corrections have been done by several groups. Wright [7] and Rephaeli [8] have done the pioneering work to the SZ effect for the CG. Challinor & Lasenby [9] and Itoh, Kohyama & Nozawa [10] have adopted a relativistically covariant formalism to describe the Compton scattering process and have obtained higher order relativistic corrections to the thermal SZ effect in the form of the Fokker–Planck expansion approximation. Nozawa, Itoh & Kohyama [11] have extended their method to the case where the CG is moving with a peculiar velocity β_c with respect to the CMB frame and have obtained the relativistic corrections to the kinematical SZ effect. Itoh, Nozawa & Kohyama [12] have also applied the covariant formalism to the polarization SZ effect [13]. The effect of the motion of the observer was also studied, for example, by Refs. [14] and [15].

On the other hand, Brown & Preston [16] studied the leading order relativistic corrections to the Kompaneets

equation by calculating the Boltzmann equation in the delta function expansion approximation. Although their expansion scheme is quite different from the previous schemes [9, 10], their leading order corrections on the SZ effect reproduced the previous results [9, 10].

In this paper, we study the SZ effect for clusters of galaxies. We explore the relativistic corrections to the Kompaneets equation in terms of two different expansion approximation schemes, namely, the Fokker-Planck expansion approximation [17] and delta function expansion approximation [16]. We show that two expansion approximation formalisms are equivalent under the Thomson approximation. This will clarify the situation for existing theoretical methods to analyse observation data.

This paper is organized as follows. In Section 2, we study the Boltzmann equation in the Fokker-Planck expansion approximation under the Thomson approximation [17]. We derive analytic expressions for the Boltzmann equation which is identical to Ref. [18]. In Section 3, we investigate the Boltzmann equation in the delta function expansion approximation [16]. Starting from the expression in the delta function expansion approximation, we derive the same expression as the Fokker-Planck expansion approximation. Finally, concluding remarks are given in Section 4.

2. Fokker-Planck expansion formalism

2.1. Boltzmann equation

Let us consider that both the CG and observer are fixed to the CMB frame. Then, the Boltzmann equation for the electron-photon scattering in the CMB frame is written

Email address: snozawa@josai.ac.jp (Satoshi Nozawa)

by:

$$\frac{dn(\omega)}{dt} = -2 \int d^3p d^3p' d^3k' W [n(\omega) \{1 + n(\omega')\} \times f(E) - n(\omega') \{1 + n(\omega)\} f(E')] , \quad (1)$$

$$W = \frac{\delta^4(p + k - p' - k') \alpha^2 \bar{X}}{(2\pi)^3 2\omega\omega' EE'} , \quad (2)$$

where $k = (\omega, \mathbf{k})$ and $k' = (\omega', \mathbf{k}')$ are the initial and final CMB photon momenta, respectively, and $p = (E, \mathbf{p})$ and $p' = (E', \mathbf{p}')$ are the momenta for electrons. In Eq. (1), $n(\omega)$ and $f(E)$ denote the distribution functions for the CMB photons and electrons in the CG, respectively. In Eq. (2), α is the fine structure constant and \bar{X} is the invariant transition probability of the Compton scattering, which will be discussed later in this section. In this paper, we use the natural unit $\hbar = c = 1$, unless otherwise stated explicitly.

According to Ref. [17], Eq. (1) is simplified in the Thomson approximation as follows:

$$\frac{dn_{\text{NK}}(x)}{d\tau} = \frac{3}{64\pi^2} \int d^3p p_e(E) \frac{1}{\gamma^2} \int d\Omega_{k'} \bar{X}_A \times \frac{1 - \beta\mu}{(1 - \beta\mu')^2} [n(x') - n(x)] , \quad (3)$$

$$\frac{x'}{x} \approx \frac{1 - \beta\mu}{1 - \beta\mu'} , \quad (4)$$

$$d\tau = N_e \sigma_T dt , \quad (5)$$

where $x = \omega/k_B T_{\text{CMB}}$, $x' = \omega'/k_B T_{\text{CMB}}$, $\sigma_T = 8\pi\alpha^2/(3m^2)$ and $f(E) = N_e \pi^2 p_e(E)$. In Eq. (3), $\beta \equiv \mathbf{p}/E$ is the velocity of the initial electron, $\gamma = 1/\sqrt{1 - \beta^2}$, $\mu = \hat{\beta} \cdot \hat{\mathbf{k}}$ and $\mu' = \hat{\beta} \cdot \hat{\mathbf{k}}'$. The invariant transition probability \bar{X} is reduced to

$$\bar{X}_A = 2 - \frac{2(1 - \cos\Theta)}{\gamma^2(1 - \beta\mu)(1 - \beta\mu')} + \frac{(1 - \cos\Theta)^2}{\gamma^4(1 - \beta\mu)^2(1 - \beta\mu')^2} , \quad (6)$$

$$\cos\Theta = \mu\mu' + \sqrt{1 - \mu^2}\sqrt{1 - \mu'^2} \cos(\phi_k - \phi_{k'}) . \quad (7)$$

Note that \bar{X}_A does not depend on ω and ω' . Note also that the Thomson approximation used in deriving Eqs. (3) and (4) is an extremely good approximation for the CMB photon energies.

Applying the Fokker-Planck expansion to $n(x')$, namely,

$$n(x') = n(x) + \sum_{\ell=1}^{\infty} \frac{1}{\ell!} (x' - x)^\ell n^{(\ell)}(x) , \quad (8)$$

one can rewrite Eq. (3) as follows:

$$\frac{dn_{\text{NK}}(x)}{d\tau} = \frac{3}{64\pi^2} \int d^3p p_e(E) \frac{1}{\gamma^2} \int d\Omega_{k'} \bar{X}_A I_{\text{NK}}(x) , \quad (9)$$

$$I_{\text{NK}}(x) = \frac{1 - \beta\mu}{(1 - \beta\mu')^2} \sum_{\ell=1}^{\infty} \frac{1}{\ell!} \left(\beta \frac{\mu' - \mu}{1 - \beta\mu'} \right)^\ell x^\ell n^{(\ell)}(x) , \quad (10)$$

where $n^{(\ell)}(x)$ is the n -th order derivative of $n(x)$.

2.2. Calculation in the electron rest frame

Calculation of Eq. (9) can be simplified by applying the following Lorentz transformations between angles in the CMB and electron rest frames [17]:

$$\mu = \frac{-\mu_0 + \beta}{1 - \beta\mu_0} , \quad (11)$$

$$\mu' = \frac{-\mu'_0 + \beta}{1 - \beta\mu'_0} , \quad (12)$$

which give

$$\bar{X}_A = 1 + \cos^2 \Theta_0 , \quad (13)$$

$$\cos \Theta_0 = \mu_0 \mu'_0 + \sqrt{1 - \mu_0^2} \sqrt{1 - \mu_0'^2} \cos(\phi_k - \phi_{k'}) . \quad (14)$$

Inserting Eqs. (11)–(14) into Eq. (9), one has

$$\frac{dn_{\text{NK}}(x)}{d\tau} = \int_0^\infty dp p^2 p_e(E) F_{\text{NK}}(x) , \quad (15)$$

$$F_{\text{NK}}(x) = \sum_{\ell=1}^{\infty} F_\ell(\beta) x^\ell n^{(\ell)}(x) , \quad (16)$$

$$F_\ell(\beta) = \int_{-1}^{+1} d\mu_0 \int_{-1}^{+1} d\mu'_0 f_\ell(\beta, \mu_0, \mu'_0) , \quad (17)$$

$$f_\ell(\beta, \mu_0, \mu'_0) = \frac{3}{16\ell!} \frac{\beta^\ell}{\gamma^4(1 - \beta\mu_0)^{\ell+3}} (\mu_0 - \mu'_0)^\ell \times [1 + \mu_0^2 \mu_0'^2 + \frac{1}{2}(1 - \mu_0^2)(1 - \mu_0'^2)] . \quad (18)$$

Equation (17) can be integrated analytically. We summarize the results up to $\mathcal{O}(\beta^{10})$ as follows:

$$F_1(\beta) = \frac{4}{3}\beta^2 + \frac{4}{3}\beta^4 + \frac{4}{3}\beta^6 + \frac{4}{3}\beta^8 + \frac{4}{3}\beta^{10} , \quad (19)$$

$$F_2(\beta) = \frac{1}{3}\beta^2 + \frac{26}{15}\beta^4 + \frac{47}{15}\beta^6 + \frac{68}{15}\beta^8 + \frac{89}{15}\beta^{10} , \quad (20)$$

$$F_3(\beta) = \frac{14}{25}\beta^4 + \frac{172}{75}\beta^6 + \frac{26}{5}\beta^8 + \frac{232}{25}\beta^{10} , \quad (21)$$

$$F_4(\beta) = \frac{7}{150}\beta^4 + \frac{17}{25}\beta^6 + \frac{1709}{630}\beta^8 + \frac{10958}{1575}\beta^{10} , \quad (22)$$

$$F_5(\beta) = \frac{44}{525}\beta^6 + \frac{7892}{11025}\beta^8 + \frac{4472}{1575}\beta^{10} , \quad (23)$$

$$F_6(\beta) = \frac{11}{3150}\beta^6 + \frac{239}{2450}\beta^8 + \frac{117}{175}\beta^{10} , \quad (24)$$

$$F_7(\beta) = \frac{128}{19845}\beta^8 + \frac{1832}{19845}\beta^{10} , \quad (25)$$

$$F_8(\beta) = \frac{16}{99225}\beta^8 + \frac{724}{99225}\beta^{10} , \quad (26)$$

$$F_9(\beta) = \frac{2}{6615}\beta^{10} , \quad (27)$$

$$F_{10}(\beta) = \frac{1}{198450}\beta^{10} . \quad (28)$$

Note that Eq. (16) with Eqs. (19)–(28) is identical to Eq. (37) with Eq. (41)–(45) of Nozawa & Kohyama [18].

3. Delta function expansion formalism

3.1. Boltzmann equation

In this section, we first review the prescription used in Brown & Preston [16]. Let us start with the following identity relation:

$$\frac{d^3 p'}{2E'} = d^4 p' \delta(p'^2 - m^2). \quad (29)$$

Inserting Eq. (29) into Eq. (1) and eliminating the 4-dimensional delta function, one has

$$\begin{aligned} \frac{dn(\omega)}{dt} = & - \int \frac{d^3 p}{(2\pi)^3 E^2} \int d\Omega_{k'} \int d\omega' \frac{\omega'}{\omega} \delta(\omega' - \omega \\ & + \beta(\omega\mu - \omega'\mu') + \frac{\omega\omega'}{E}(1 - \cos\Theta)) \\ & \times \alpha^2 \bar{X} [n(\omega) \{1 + n(\omega')\} f(E) \\ & - n(\omega') \{1 + n(\omega)\} f(E')]. \end{aligned} \quad (30)$$

In the Thomson approximation $\gamma\omega/m \ll 1$, Eq. (30) is simplified as follows:

$$\begin{aligned} \frac{dn_{\text{BP}}(x)}{d\tau} = & \frac{3}{64\pi^2} \int d^3 p p_e(E) \frac{1}{\gamma^2} \int d\Omega_{k'} \bar{X}_A \\ & \times \int dx' \delta(x' - x + \beta(x\mu - x'\mu')) G(x'), \end{aligned} \quad (31)$$

$$G(x') = \frac{x'}{x} [n(x') - n(x)]. \quad (32)$$

In Eq. (31), the delta function is formally expanded by

$$\delta(x' - x + \mathcal{O}(x')) = \sum_{\ell=0}^{\infty} \frac{1}{\ell!} \mathcal{O}^\ell(x') \delta^{(\ell)}(x' - x), \quad (33)$$

$$\mathcal{O}(x') = \beta(x\mu - x'\mu'), \quad (34)$$

where $\delta^{(\ell)}(x' - x)$ is the ℓ -th order derivative of $\delta(x' - x)$ in terms of x' .

Inserting Eq. (33) into Eq. (31), one finally has

$$\frac{dn_{\text{BP}}(x)}{d\tau} = \frac{3}{64\pi^2} \int d^3 p p_e(E) \frac{1}{\gamma^2} \int d\Omega_{k'} \bar{X}_A I_{\text{BP}}(x), \quad (35)$$

$$I_{\text{BP}}(x) = \sum_{\ell=0}^{\infty} \frac{1}{\ell!} \int dx' \delta^{(\ell)}(x' - x) G(x') \mathcal{O}^\ell(x'). \quad (36)$$

In Ref. [16], Eq. (30) was calculated with the delta function identities defined in their Appendix C up to fourth-order derivatives.

Thus, we have shown the expressions for the Boltzmann equation under the Thomson approximation in two different schemes. Although the expressions for $I_{\text{NK}}(x)$ and $I_{\text{BP}}(x)$ are quite different each other, we will show in the next subsection that these formalisms are indeed identical.

3.2. Equivalence of the formalisms

In order to show the equivalence of two formalisms, we start with $I_{\text{BP}}(x)$ of Eq. (36) and derive the same expression as $I_{\text{NK}}(x)$ of Eq. (10). Let us first introduce the definition for the ℓ -th order derivative of the delta function:

$$\int dx \delta^{(\ell)}(x - a) \phi(x) = (-1)^\ell \int dx \delta(x - a) \phi^{(\ell)}(x), \quad (37)$$

where $\phi(x)$ is an arbitrary function. Then, one can rewrite Eq. (36) as follows:

$$I_{\text{BP}}(x) = \sum_{\ell=0}^{\infty} \frac{(-1)^\ell}{\ell!} \int dx' \delta(x' - x) [G(x') \mathcal{O}^\ell(x')]^{(\ell)}. \quad (38)$$

Equation (38) is further simplified with the relation:

$$\begin{aligned} [G(x') \mathcal{O}^\ell(x')]^{(\ell)} = & \sum_{j=0}^{\ell} \ell C_j G^{(j)}(x') \\ & \times \frac{\ell!}{j!} (-\beta\mu')^{\ell-j} \mathcal{O}^j(x'). \end{aligned} \quad (39)$$

Inserting Eq. (39) into Eq. (38), exchanging the order of summations for j and ℓ , and introducing a new index i by $\ell = j + i$, one obtains

$$\begin{aligned} I_{\text{BP}}(x) = & \sum_{j=0}^{\infty} \frac{(-1)^j}{j!} \sum_{i=0}^{\infty} {}_{\ell} C_j (\beta\mu')^i \\ & \times \int dx' \delta(x' - x) G^{(j)}(x') \mathcal{O}^j(x') \\ = & \sum_{j=0}^{\infty} \frac{(-1)^j}{j!} \frac{\mathcal{O}^j(x)}{(1 - \beta\mu')^{j+1}} \int dx' \delta(x' - x) G^{(j)}(x'). \end{aligned} \quad (40)$$

In deriving the last equality of Eq. (40), we used

$$\sum_{i=0}^{\infty} {}_{\ell} C_j (\beta\mu')^i = \frac{1}{(1 - \beta\mu')^{j+1}}, \quad (41)$$

and we took $\mathcal{O}^j(x)$ outside of the integral with a familiar property of the delta function.

The j -th order derivative of $G(x')$ is calculated with Eq. (32) as follows:

$$G^{(j)}(x') = \frac{x'}{x} n^{(j)}(x') + \frac{j}{x} n^{(j-1)}(x'), \quad (42)$$

where $n^{(0)}(x') \equiv G(x')$. Inserting Eqs. (34) and (42) into Eq. (40), one finally obtains

$$I_{\text{BP}}(x) = \frac{1 - \beta\mu}{(1 - \beta\mu')^2} \sum_{j=1}^{\infty} \frac{1}{j!} \left(\beta \frac{\mu' - \mu}{1 - \beta\mu'} \right)^j x^j n^{(j)}(x), \quad (43)$$

where $j = 0$ term was zero because of the relation:

$$\int dx' \delta(x' - x) G(x') = 0. \quad (44)$$

Thus, one finds that two formalisms are equivalent in the Thomson approximation

Before closing this section it should be noted as follows: In Nagirner & Poutanen [19], a similar discussion was made between the delta function expansion formalism and the Fokker-Planck expansion formalism, where the Maxwellian electron distribution function was expanded in terms $\theta_e \equiv k_B T_e / mc^2$ and the delta function, and the $O(\theta_e)$ terms were retained. On the other hand, the equivalence of the two formalisms in this paper is more general, which is independent of the electron distribution functions.

4. Conclusion

We studied the SZ effect in the Thomson approximation. In Section 2, we investigated the Boltzmann equation for the photon distribution function in the Fokker-Planck expansion approximation. We derived the analytic formula for the Boltzmann equation which is identical to that obtained in Nozawa & Kohyama [18].

In Section 3, we studied the Boltzmann equation for the photon distribution function in the delta function expansion approximation [16]. Starting from the expression in the delta function expansion approximation, we derived the same expression as the Fokker-Planck expansion approximation. We conclude that two expansion approximation formalisms are equivalent under the Thomson approximation, which is extremely good approximation for the CMB photon energies. This will clarify the situation for existing theoretical methods to analyse observation data.

Acknowledgements

We thank our referee for a valuable suggestion.

References

- [1] Ya. B. Zeldovich, R. A. Sunyaev, *Ap&SS* 4 (1969) 301.
- [2] R. A. Sunyaev, Ya. B. Zeldovich, *MNRAS* 190 (1980) 413.
- [3] M. Birkinshaw, *Phys. Rep.* 310 (1999) 97.
- [4] J. E. Carlstrom, G. P. Holder, E. D. Reese, *ARA&A* 40 (2002) 643.
- [5] A. S. Kompaneets, *Sov. Phys.—JETP* 31 (1956) 876.
- [6] S. W. Allen, R. W. Schmidt, A. C. Fabian, *MNRAS* 335 (2002) 256
- [7] E. L. Wright, *ApJ* 232 (1979) 348.
- [8] Y. Rephaeli, *ApJ* 445 (1995) 33.
- [9] A. Challinor, A. Lasenby, *ApJ* 499 (1998) 1.
- [10] N. Itoh, Y. Kohyama, S. Nozawa, *ApJ* 502 (1998) 7.
- [11] S. Nozawa, N. Itoh, Y. Kohyama, *ApJ* 508 (1998) 17.
- [12] N. Itoh, S. Nozawa, Y. Kohyama, *ApJ* 533 (2000) 588.
- [13] R. A. Sunyaev, Ya. B. Zeldovich, *Astrophys. Space Phys. Rev.* 1 (1981) 1.
- [14] J. Chluba, G. Hütsi, R. A. Sunyaev, *A&A* 434 (2005) 811.
- [15] S. Nozawa, N. Itoh, Y. Kohyama, *A&A* 440 (2005) 39.
- [16] L. S. Brown, D. L. Preston, *Astroparticle Phys.* 35 (2012) 742.
- [17] S. Nozawa, Y. Kohyama, *Phys. Rev. D* 79 (2009) 083005.
- [18] S. Nozawa, Y. Kohyama, *MNRAS* 441 (2014) 3018.
- [19] D. I. Nagirner, J. Poutanen, *Astrophys. Space Phys. Rev.* 9 (1994) 1.