Reduction of a family of ideals

Tomasz Rodak

Abstract. In the paper we prove that there exists a simultaneous reduction of one-parameter family of \mathfrak{m}_n -primary ideals in the ring of germs of holomorphic functions. As a corollary we generalize the result of A. Płoski [6] on the semicontinuity of the Łojasiewicz exponent in a multiplicity-constant deformation.

Mathematics Subject Classification (2010). Primary 14B07; Secondary 14C17, 13H15.

Keywords. Reduction of an ideal, Hilbert-Samuel multiplicity, Łojasiewicz exponent.

1. Introduction

Let R be a ring and I an ideal. We say that an ideal J is a *reduction of* I if it satisfies the following condition:

 $J \subset I$, and for some r > 0 we have $I^{r+1} = JI^r$.

The notion of reduction is closely related to the notions of *Hilbert-Samuel multiplicity* and *integral closure* of an ideal.

Recall that if (R, \mathfrak{m}) is a Noetherian local ring of dimension n and I is an \mathfrak{m} -primary ideal of R, then the *Hilbert-Samuel multiplicity* of I is given by the formula

$$e(I) = n! \lim_{k \to \infty} \frac{\text{length}_R R/I^k}{k^n}.$$

For the multiplicity theory in local rings see for example [5] or [2].

Let I be an ideal in a ring R. An element $x \in R$ is said to be *integral* over I if there exists an integer n and elements $a_k \in I^k$, k = 1, ..., n, such that

$$x^{n} + a_{1}x^{n-1} + \dots + a_{n} = 0.$$

This research was partially supported by the Polish OPUS Grant No $2012/07/{\rm B}/{\rm ST1}/03293.$

The set of all elements of R that are integral over I is called the *integral* closure of I, and is denoted \overline{I} . If $I = \overline{I}$ then I is called *integrally closed*. It is well known that \overline{I} is an ideal.

The relationship between the above notions is given in the following Theorem due to D. Rees:

Theorem 1 (Rees, [2, Cor. 1.2.5, Thm. 11.3.1]). Let (R, \mathfrak{m}) be a formally equidimensional Noetherian local ring and let $J \subset I$ be two \mathfrak{m} -primary ideals. Then the following conditions are equivalent:

- 1. J is a reduction of I;
- 2. e(I) = e(J);
- 3. $\overline{I} = \overline{J}$.

It is an important fact that a reduction of an ideal is often generated by a system of parameters. More precisely we have

Theorem 2 ([5, Theorem 14.14]). Let (R, \mathfrak{m}) be a d-dimensional Noetherian local ring, and suppose that $k = R/\mathfrak{m}$ is an infinite field; let $I = (u_1, \ldots, u_s)$ be an \mathfrak{m} -primary ideal. Then there exist a finite number of polynomials $D_{\alpha} \in$ $k[Z_{ij}; 1 \leq i \leq d, 1 \leq j \leq s], 1 \leq \alpha \leq \nu$ such that if $y_i = \sum a_{ij}u_j, i = 1, \ldots, d$ and at least one of $D_{\alpha}(\overline{a}_{ij}; 1 \leq i \leq d, 1 \leq j \leq s) \neq 0$, then the ideal $(y_1, \ldots, y_d)R$ is a reduction of I and $\{y_1, \ldots, y_d\}$ is a system of parameters of R.

Let $(\mathcal{O}_n, \mathfrak{m}_n)$ be the ring of germs of holomorphic functions $(\mathbb{C}^n, 0) \to \mathbb{C}$. The aim of this note is to prove the following:

Theorem 3. Let $F = F_t(x) = F(x,t)$: $(\mathbb{C}^n \times \mathbb{C}, 0) \to (\mathbb{C}^m, 0)$ be a holomorphic map. Assume that $(F_t)\mathcal{O}_n$ is an \mathfrak{m}_n -primary ideal for all t. Then there exists a complex linear map $\pi : \mathbb{C}^m \to \mathbb{C}^n$ such that for all t the ideal $(\pi \circ F_t)\mathcal{O}_n$ is a reduction of $(F_t)\mathcal{O}_n$.

In the next section we get as a corollary that if the above family $(F_t) \mathcal{O}_n$ is of constant multiplicity then the Łojasiewicz exponent in this family is a lower semicontinuos function of t. A. Płoski proved this result under additional restriction m = n but with space of parameters of arbitrary dimension.

The proof of Theorem 3 is based on some geometric property of Hilbert-Samuel multiplicity, given in section 3.

2. Semicontinuity of the Łojasiewicz exponent

Let (R, \mathfrak{m}) be a local ring and let I be an \mathfrak{m} -primary ideal. By the *Lojasiewicz* exponent $\mathcal{L}(I)$ of I we define the infimum of

$$\left\{\frac{p}{q}:\mathfrak{m}^p\subset\overline{I^q}\right\}.$$

It was proved in [3] that if $F: (\mathbb{C}^n, 0) \to (\mathbb{C}^m, 0)$ is a holomorphic map with an isolated zero at the origin and $I := (F) \mathcal{O}_n$, then $\mathcal{L}(I)$ is an optimal exponent ν in the inequality

$$|F(x)| \ge C |x|^{\nu},$$

where C is some positive constant and x runs through sufficiently small neighbourhood of $0 \in \mathbb{C}^n$.

Lemma 4. Let (R, \mathfrak{m}) be a Noetherian local ring. If I is an \mathfrak{m} -primary ideal of R and J is a reduction of I then $\mathcal{L}(I) = \mathcal{L}(J)$.

Proof. Obviously $\mathcal{L}(I) \leq \mathcal{L}(J)$. Assume that $\mathfrak{m}^p \subset \overline{I^q}$. Since J is a reduction of I, then also J^q is a reduction of I^q [2, Prop. 8.1.5]. Thus $\overline{J^q} = \overline{I^q}$ by Theorem 1, which gives $\mathfrak{m}^p \subset \overline{J^q}$. This proves the inequality $\mathcal{L}(J) \leq \mathcal{L}(I)$ and ends the proof.

Corollary 5 (A. Płoski for m = n, [6]). Let $F: (\mathbb{C}^n \times \mathbb{C}, 0) \to (\mathbb{C}^m, 0)$ be a holomorphic map. Put $I_t := (F_t) \mathcal{O}_n$. If the function $t \mapsto e(I_t)$ is constant and finite then the function $t \mapsto \mathcal{L}(I_t)$ is lower semicontinuos.

Proof. By Theorem 3 there exists a linear map $\pi : \mathbb{C}^m \to \mathbb{C}^n$ such that $J_t := (\pi \circ F_t) \mathcal{O}_n$ is a reduction of I_t for all t. Thus $\mathcal{L}(J_t) = \mathcal{L}(I_t)$ and $e(J_t) = e(I_t)$ by Theorem 1 and Lemma 4. Consequently $t \mapsto e(J_t)$ is constant and finite and the assertion follows from the case m = n proved by A. Płoski. \Box

3. Improper intersection multiplicity

Let I be an \mathfrak{m}_n -primary ideal of \mathcal{O}_n and let f_1, \ldots, f_m be its generators. We put $f = (f_1, \ldots, f_m)$. It is well known that if m = n then

$$e(I) = \dim_{\mathbb{C}} \mathcal{O}_n / I.$$

On the other hand, if m > n then we may define so-called *improper in*tersection multiplicity $i_0(I)$ of I as the improper intersection multiplicity $i(\operatorname{graph} f \cdot (\mathbb{C}^n \times \{0\}); (0,0))$ of $\operatorname{graph} f$ and $\mathbb{C}^n \times \{0\}$ at the point $(0,0) \in \mathbb{C}^n \times \mathbb{C}^m$ (see [1]).

Let C_f be the (Whitney) tangent cone of the germ of the image of f at the origin. The following observation is due to S. Spodzieja.

Theorem 6 ([7]). The number $i_0(I)$ is well defined. Moreover, if $\pi : \mathbb{C}^m \to \mathbb{C}^l$ is a linear map such that ker $\pi \cap C_f = \{0\}$, then the ideal J generated by $\pi \circ f$ is \mathfrak{m}_n -primary and we have $i_0(I) = i_0(J)$. If additionally l = n then $i_0(I) = e(J)$.

Corollary 7. If I is an \mathfrak{m}_n -primary ideal in \mathcal{O}_n , then $i_0(I) = e(I)$.

Proof. Let $I = (f_1, \ldots, f_m)\mathcal{O}_n$. By Theorems 2 and 6 there exists linear combinations $g_i = \sum a_{ij}f_j$, $i = 1, \ldots, n$ such that $J = (g_1, \ldots, g_n)\mathcal{O}_n$ is a reduction of I, $\{g_1, \ldots, g_n\}$ is a system of parameters of \mathcal{O}_n and $i_0(I) = i_0(J) = e(J)$. From Theorem 1 we get e(I) = e(J). This ends the proof. \Box

Corollary 8. If $\pi : \mathbb{C}^m \to \mathbb{C}^l$ is a linear map such that ker $\pi \cap C_f = \{0\}$, then the ideal J generated by $\pi \circ f$ is a reduction of I.

Proof. We have $J \subset I$ and e(J) = e(I). This and Theorem 1 give the assertion.

4. Elementary blowing-up

Here we recall the notion of an elementary blowing-up after [4].

Let $U \subset \mathbb{C}^n$ be an open and connected neighbourhood of $0 \in \mathbb{C}^n$; let $f = (f_0, \ldots, f_m) \neq 0$ be a sequence of holomorphic functions on U. Put $S = \{x \in U : f(x) = 0\}$ and

$$E(f) = \{ (x, u) \in U \times \mathbb{P}^m : f_i(x)u_j = f_j(x)u_i, i, j = 0, \dots, m \},\$$

where $u = [u_0 : \cdots : u_m] \in \mathbb{P}^m$.

Let Y be the closure of $E(f) \setminus S$ in $U \times \mathbb{P}^m$. The natural projection

 $\pi:Y\to U$

is called the *(elementary)* blowing-up of U by means of f_0, \ldots, f_m . The analytic subset S is called a *centre of the blowing-up* and its inverse image $\pi^{-1}(S) \subset Y$ is called the *exceptional set* of the blowing-up.

Proposition 9. Under above notations we have:

- 1. Y is an analytic subset of $U \times \mathbb{P}^m$;
- 2. π is proper, its range is U and the restriction $\pi_{|Y\setminus\pi^{-1}(S)}$ is a biholomorphism onto $U\setminus S$;
- 3. Y is irreducible;
- 4. The exceptional set $\pi^{-1}(S)$ is analytic in $U \times \mathbb{P}^m$ and it is of pure dimension n-1.

Proof. Although the above proposition is well known, we think that point (4) is worth proving. Let us consider the analytic map

$$F \colon U \times \mathbb{P}^m \ni (x, u) \mapsto (f(x), u) \in \mathbb{C}^{m+1} \times \mathbb{P}^m.$$

Let y_0, \ldots, y_m be coordinates in \mathbb{C}^{m+1} . If we denote by $\pi_{m+1} \colon \Pi_{m+1} \to \mathbb{C}^{m+1}$ the blowing-up of \mathbb{C}^{m+1} by means of y_0, \ldots, y_m then for the restriction $\tilde{f} = F_{|Y}$ we get the following commutative diagram of analytic maps:

$$Y \xrightarrow{\widetilde{f}} \Pi_{m+1}$$
$$\downarrow^{\pi} \qquad \qquad \downarrow^{\pi_{m+1}}$$
$$U \xrightarrow{f} \mathbb{C}^{m+1}$$

Take $(x_0, u_0) \in \pi^{-1}(0)$. Let $\Omega \subset \prod_{m+1}$ be a neighbourhood of $(0, u_0)$, $h: \Omega \to \mathbb{C}$ an analytic function such that

$$\pi_{m+1}^{-1}(0) \cap \Omega = \{(y, u) \in \Omega : h(y, u) = 0\}$$

Let $\widetilde{\Omega} \subset Y$ be a neighbourhood of (x_0, u_0) such that $\widetilde{f}(\widetilde{\Omega}) \subset \Omega$. Since $\widetilde{f}^{-1}(\pi_{m+1}^{-1}(0)) = \pi^{-1}(S)$ we get

$$\pi^{-1}(S) \cap \widetilde{\Omega} = \{ (x, u) \in \widetilde{\Omega} : h \circ \widetilde{f}(x, u) = 0 \}.$$

Thus there exists a neighbourhood $\Delta \subset U \times \mathbb{P}^m$ of (x_0, u_0) and an analytic set $V \subset \Delta$ of pure dimension n + m - 1 such that

$$\pi^{-1}(S) \cap \Delta = V \cap Y \cap \Delta.$$

This gives

$$\dim_{(x_0,u_0)} \pi^{-1}(S) \ge \dim_{(x_0,u_0)} Y - 1 = n - 1.$$

Since Y is irreducible and $\pi^{-1}(S) \subsetneq Y$ we get that $\dim_p \pi^{-1}(S) = n-1$ for any $p \in \pi^{-1}(S)$. This ends the proof. \Box

5. Proof of Theorem 3

Lemma 10. Let $F: (\mathbb{C}^n \times \mathbb{C}, 0) \to (\mathbb{C}^{m+1}, 0), m \ge n$ be a holomorphic map. Assume that 0 is an isolated point of $F_t^{-1}(0)$ for $|t| < \delta$. Then there exists $\delta > \epsilon > 0$ and a complex line $V \subset \mathbb{C}^{m+1}$, such that $V \cap C_{F_t} = \{0\}$ for $|t| < \epsilon$.

Proof. Let $F: U \to \mathbb{C}^{m+1}$, where $U \subset \mathbb{C}^n \times \mathbb{C}$ is a connected neighbourhood of the origin. Put $S = \{(z,t) \in U : F(z,t) = 0\}$ and let $\pi: U \times \mathbb{P}^m \supset Y \to U$ be the elementary blowing-up of U by F. By Proposition 9 its exceptional set $E := \pi^{-1}(S)$ is an analytic set of pure dimension n. Let \mathcal{E} be a set of those irreducible components W of E for which origin in \mathbb{C}^{n+1} is an accumulation point of $\pi(W) \cap (\{0\} \times \mathbb{C})$. Then \mathcal{E} is finite. Denote by $\widetilde{C_{F_t}}$ the image of the cone C_{F_t} in \mathbb{P}^m . Observe that

$$\{(0,t)\} \times \widetilde{C_{F_t}} \subset \bigcup \mathcal{E}, \quad |t| < \delta.$$

On the other hand for any $W \in \mathcal{E}$ we have

 $\dim W \cap (\{0\} \times \mathbb{P}^m) \leqslant n - 1 < m.$

Thus there exists $\epsilon > 0$ and an open set $G \subset \mathbb{P}^m$ such that

$$(\{(0,t)\} \times G) \cap [\mathcal{E} = \emptyset, \quad 0 < |t| < \epsilon$$

As a result if V is a line in \mathbb{C}^{m+1} corresponding to some point in G then $V \cap C_{F_t} = \{0\}$ for $0 < |t| < \epsilon$. Since G is not a subset of C_{F_0} we get the assertion.

Proof of Theorem 3. Induction on m. In the case m = n there is nothing to prove. Let us assume that the assertion is true for some $m \ge n$ and let $F: (\mathbb{C}^n \times \mathbb{C}, 0) \to (\mathbb{C}^{m+1}, 0)$ be a holomorphic map such that the ideals $(F_t) \mathcal{O}_n$ are \mathfrak{m}_n -primary. By Lemma 10 there exists $\epsilon > 0$ and a linear mapping $\pi': \mathbb{C}^{m+1} \to \mathbb{C}^m$ such that ker $\pi' \cap C_{F_t} = \{0\}$ for $|t| < \epsilon$. Thus, by Corollary 8 the ideal $(\pi' \circ F_t) \mathcal{O}_n$ is a reduction of $(F_t) \mathcal{O}_n$. On the other hand, by induction hypothesis, there exists a linear map $\pi'': \mathbb{C}^m \to \mathbb{C}^n$ such that $(\pi'' \circ \pi' \circ F_t) \mathcal{O}_n$ is a reduction of $(\pi' \circ F_t) \mathcal{O}_n$ for small t. Thus if we put $\pi := \pi'' \circ \pi'$ we get the assertion. \Box

Tomasz Rodak

References

- R. Achilles, P. Tworzewski, and T. Winiarski. On improper isolated intersection in complex analytic geometry. Ann. Polon. Math., 51:21–36, 1990.
- [2] Craig Huneke and Irena Swanson. Integral closure of ideals, rings, and modules, volume 336 of London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 2006.
- [3] M. Lejeune-Jalabert and B. Teissier. Séminaire Lejeune-Teissier: Cloture Integrale des Ideaux et equisingularite : Chapitre 1. Universite Scientifique et Medicale de Grenoble. Laboratoire de mathématiques pures associe au C.N.R.S., 1974.
- [4] Stanisław Łojasiewicz. Introduction to complex analytic geometry. Birkhäuser Verlag, Basel, 1991. Translated from the Polish by Maciej Klimek.
- [5] Hideyuki Matsumura. Commutative ring theory, volume 8 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, second edition, 1989. Translated from the Japanese by M. Reid.
- [6] A. Płoski. Semicontinuity of the Łojasiewicz exponent. Univ. Iagel. Acta Math., 48:103–110, 2010.
- [7] S. Spodzieja. Multiplicity and the Łojasiewicz exponent. Ann. Polon. Math., 73(3):257–267, 2000.

Tomasz Rodak University of Łódź Faculty of Mathematics and Computer Science S. Banacha 22, 90-238 Łódź, Poland e-mail: rodakt@math.uni.lodz.pl