

# Variable Speed of Light Cosmology, Primordial Fluctuations and Gravitational Waves

J. W. Moffat

Perimeter Institute for Theoretical Physics, Waterloo, Ontario N2L 2Y5, Canada  
and  
Department of Physics and Astronomy, University of Waterloo, Waterloo,  
Ontario N2L 3G1, Canada

February 9, 2019

PACS: 98.80.C; 04.20.G; 04.40.-b

## Abstract

A variable speed of light (VSL) cosmology is developed with a spontaneous breaking of Lorentz invariance in the early universe. A non-minimal electromagnetic coupling to curvature and the resulting quantum electrodynamic vacuum polarization dispersive medium can produce  $c \gg c_0$  in the early universe, where  $c_0$  is the measured speed of light today. Higher derivative curvature contributions to the effective gravitational action and quantum gravity vacuum polarization can produce a dispersive medium and a large increase in the speed of gravitational waves  $c_g \gg c_{g0}$  in the early universe, where  $c_{g0}$  is the speed of gravitational waves today. The initial value problems of cosmology are solved: the horizon and flatness problems. The model predicts primordial scalar and tensor fluctuation spectral indices  $n_s = 0.96$  and  $n_t = -0.04$ , respectively.

## 1 Introduction

Although inflationary models have been successful in fitting cosmological data [1, 2, 3], there have been issues raised about the fundamental consequences of the models [4]. In particular, the need for chaotic and eternal inflation models has raised the specter of a multiverse cosmological scenario [5, 6]. The question as to whether such a scenario can be falsified by observations and the lack of predictability of inflation models has been a cause for concern. Moreover, the standard single-field inflation models suffer from significant fine-tuning, such as the requirement of a slow-rolling potential and the fine-tuning needed to fit the magnitude of the CMB amplitude [7, 8, 9, 10, 11, 12].

We shall consider a model based on the idea that the speed of light  $c$  can have a significantly larger value in the very early universe [13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25]. This assumption leads to a resolution of the horizon and flatness problems, thereby solving the initial value problem in early universe cosmology. A bimetric variable speed of light (VSL) model [26, 27, 28, 29, 30] has been proposed with two metric tensors. One metric and its light cone describe a varying speed of light and a constant speed of gravitational waves, while the other metric and its light cone describe a constant speed of light and a varying speed of gravitational waves. This model cannot produce an observed value for relic gravitational waves and a non-zero tensor mode spectral index  $n_t$ .

In the following, we will formulate a version of VSL based on an earlier VSL model with one metric. In Section 2, we postulate a gravitational action in which the speed of light  $c = c(x)$  is a dynamical field. The total action also contains an action for a minimally coupled scalar “seed” field  $\phi$ , which will produce quantum primordial fluctuations. It also contains an action inducing a spontaneous violation of Lorentz invariance [13, 14, 31, 32, 33] by means of a non-vanishing vacuum expectation value of a vector field  $\psi_\mu$ . The Lorentz group  $SO(3,1) \rightarrow O(3) \times R$  where  $R$  is the absolute preferred time and corresponds to the

comoving time  $t$  in the preferred frame associated with the Friedmann-Lemaître-Robertson-Walker (FLRW) metric. In Section 3, we develop the VSL cosmology and we show how the model can solve the horizon and flatness problems in initial value cosmology. In Section 4, we will demonstrate how a large speed of light  $c \gg c_0$ , where  $c_0$  is the present measured speed of light, and a large speed of gravitational waves  $c_g \gg c_{0g}$  where  $c_{0g}$  is the speed of gravitational waves today are produced fractions of seconds after the big bang, induced by non-minimal electromagnetic and gravitational couplings to the curvature of spacetime. Section 5 presents the calculations of the power spectra and spectral indices for primordial scalar density perturbations and tensor gravitational waves. In Section 6 we end with concluding comments.

The VSL cosmology can remove the fine-tuning of the initial values in the standard big-bang cosmology and fit the available observational data, such as an almost scale invariant, adiabatic and Gaussian scalar matter power spectrum and the potential observation of a gravitational wave power spectrum. As a viable alternative to standard inflationary models, it can relieve the need for fine-tuning present in these models and not require a multiverse scenario in the form of eternal inflation.

## 2 The action and field equations

We adopt the following action:

$$S = S_G + S_\psi + S_\phi + S_M, \quad (1)$$

where

$$S_G = \frac{1}{16\pi G} \int d^4x \sqrt{-g} \Phi(R + 2\Lambda). \quad (2)$$

Here,  $g = \det(g_{\mu\nu})$ ,  $R = g^{\mu\nu} R_{\mu\nu}$ ,  $\Lambda$  is the cosmological constant and  $\Phi(x) = c^4(x)$ . The action  $S_\psi$  is given by

$$S_\psi = \int d^4x \sqrt{-g} \left[ -\frac{1}{4} B^{\mu\nu} B_{\mu\nu} - W(\psi_\mu) \right], \quad (3)$$

where  $\psi_\mu$  is a vector field,  $B_{\mu\nu} = \partial_\mu \psi_\nu - \partial_\nu \psi_\mu$ , and  $W(\psi_\mu)$  is a potential. The scalar field action  $S_\phi$  is defined by

$$S_\phi = \frac{c_0^4}{16\pi G} \int d^4x \sqrt{-g} \left[ \frac{1}{2} (\partial_\mu \phi \partial^\mu \phi) - V(\phi) \right], \quad (4)$$

where  $\phi$  is a dimensionless scalar field,  $V(\phi)$  is a potential and  $c_0$  is the measured speed of light today. The matter action is  $S_M = S_M(\phi_m, g_{\mu\nu})$  where  $\phi_m$  denotes matter fields.

The energy-momentum tensor is

$$T_{\mu\nu} = T_{M\mu\nu} + T_{\psi\mu\nu} + T_{\phi\mu\nu}, \quad (5)$$

where

$$\frac{1}{\sqrt{-g}} \frac{\delta S_X}{\delta g^{\mu\nu}} = -\frac{1}{2} T_{X\mu\nu}, \quad [X = M, \psi, \phi]. \quad (6)$$

The  $T_{\psi\mu\nu}$  and  $T_{\phi\mu\nu}$  are given by

$$T_{\psi\mu\nu} = B_{\mu\alpha} B^\alpha{}_\nu - \frac{1}{4} g_{\mu\nu} B^{\alpha\beta} B_{\alpha\beta} - \frac{\partial W(\psi_\mu)}{\partial g^{\mu\nu}} + \frac{1}{2} g_{\mu\nu} W(\psi_\mu), \quad (7)$$

and

$$T_{\phi\mu\nu} = \frac{c_0^4}{16\pi G} \left( \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi - 2 \frac{\partial V(\phi)}{\partial g^{\mu\nu}} + \frac{1}{2} g_{\mu\nu} V(\phi) \right). \quad (8)$$

Setting the cosmological constant  $\Lambda = 0$ , the variation of the action with respect to  $g^{\mu\nu}$  yields the field equation:

$$G_{\mu\nu} = \frac{8\pi G}{\Phi} T_{\mu\nu} + \frac{\nabla_\mu \partial_\nu \Phi}{\Phi} - g_{\mu\nu} \frac{\nabla^\alpha \nabla_\alpha \Phi}{\Phi}, \quad (9)$$

where  $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R$  and where  $\nabla_\mu$  is the covariant derivative with respect to the metric  $g_{\mu\nu}$ . We also obtain the field equations:

$$\nabla_\mu (B^{\mu\nu}) - \frac{\partial W(\psi_\mu)}{\partial \psi_\nu} = 0, \quad (10)$$

and

$$g^{\mu\nu}\nabla_\mu\nabla_\nu\phi + \frac{\partial V(\phi)}{\partial\phi} = 0. \quad (11)$$

The energy-momentum tensor  $T^{\mu\nu}$  satisfies the conservation law:

$$\nabla_\nu T^{\mu\nu} = 0. \quad (12)$$

In the action  $S$  we have made the speed of light  $c$  a dynamical degree of freedom, in addition to the dynamical degrees of freedom associated with the metric  $g_{\mu\nu}$  and the fields  $\psi_\mu$  and  $\phi$ .

We must now require that local Lorentz invariance and diffeomorphism invariance are violated, so that the speed of light  $c = c(t)$  cannot be made constant by a coordinate transformation as is the case in GR. Let us choose  $W(\psi_\mu)$  to be of the form of a ‘‘Mexican hat’’ potential [13, 14]:

$$W(\psi_\mu) = -\frac{1}{2}\mu^2\psi_\mu\psi^\mu + \frac{1}{4}\lambda(\psi_\mu\psi^\mu)^2, \quad (13)$$

where  $\lambda > 0$  and  $\mu^2 > 0$ . If  $W$  has a minimum at

$$v_\mu \equiv \psi_\mu = \langle 0|\psi_\mu|0\rangle, \quad (14)$$

then the spontaneously broken solution is given by

$$v^2 \equiv \psi^\mu\psi_\mu = \frac{\mu^2}{\lambda}. \quad (15)$$

We choose the ground state to be described by the timelike vector:

$$\psi_\mu^{(0)} = \delta_{\mu 0}v = \delta_{\mu 0}\left(\frac{\mu^2}{\lambda}\right)^{1/2}. \quad (16)$$

The homogeneous Lorentz group  $SO(3,1)$  is broken down to the spatial rotation group  $O(3)$ . The three rotation generators  $J_i$  ( $i = 1, 2, 3$ ) leave the vacuum invariant,  $J_i v_i = 0$ , while the Lorentz boost generators  $K_i$  break the vacuum symmetry  $K_i v_i \neq 0$ . The spontaneous breaking of the Lorentz and diffeomorphism symmetries produces massless Nambu-Goldstone modes and massive particle modes [34]. The spontaneous breaking of Lorentz invariance and diffeomorphism invariance has selected a preferred frame and direction of time.

### 3 Initial value conditions and cosmology

We will work in our application to cosmology in the preferred frame in which the metric is of the FLRW form:

$$ds^2 = c^2 dt^2 - a^2 \left[ \frac{dr^2}{1 - Kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right], \quad (17)$$

where  $K$  is the Gaussian curvature of space and  $K = 0, +1, -1$  (in units of  $[\text{length}]^{-2}$ ) for flat, closed and open models, respectively. The metric has the group symmetry  $O(3) \times R$  with a *preferred* comoving time  $t$ . The energy-momentum tensor  $T_{\mu\nu}$  will be described by a perfect fluid:

$$T^{\mu\nu} = \left( \rho + \frac{p}{c^2} \right) u^\mu u^\nu - pg^{\mu\nu}, \quad (18)$$

where  $u^\mu = dx^\mu/ds$  is the fluid element four-velocity and  $p$  are the matter density and pressure, respectively.

As in inflationary models, the horizon problem is solved in our VSL model [13]. Consider a locally flat patch with the line element:

$$ds^2 = c^2 dt^2 - (dx^i)^2. \quad (19)$$

From the geodesic equation for light travel  $ds^2 = 0$ , we get

$$dt^2 = \frac{1}{c^2}(dx^i)^2. \quad (20)$$

In the limit when  $c \rightarrow \infty$  in an early phase of the universe,  $dt^2 \rightarrow 0$  and the Minkowski light cone is squashed. Now all points in an expanding bubble near the beginning of the universe will be in causal communication with one another. The horizon scale is determined by

$$d_H = ca(t) \int_0^t \frac{dt'}{a(t')}. \quad (21)$$

Let us assume that for  $t > t_c$  we have  $c = c_0$ . Then for  $t < t_c$  we get  $d_H \rightarrow \infty$  as  $c \rightarrow \infty$  and in the phase  $t < t_c$  all points in the expanding spacetime will have been in causal communication.

The Friedmann equations are given in our model by

$$H^2 + \frac{Kc^2}{a^2} = \frac{8\pi G\rho}{3} - 4H\frac{\dot{c}}{c}, \quad (22)$$

and

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left( \rho + 3\frac{p}{c^2} \right) - 2H\frac{\dot{c}}{c} - 6\frac{\dot{c}^2}{c^2} - 2\frac{\ddot{c}}{c}, \quad (23)$$

where  $H = \dot{a}/a$  and  $\rho = \rho_M + \rho_\psi + \rho_\phi$ .

In standard big-bang cosmology with  $c = c_0$ :

$$|\Omega(10^{-43}\text{sec}) - 1| \sim \mathcal{O}(10^{-60}), \quad |\Omega(1\text{sec}) - 1| \sim \mathcal{O}(10^{-16}), \quad (24)$$

where  $\Omega = \rho/\rho_c$ ,  $\rho_c = 3H^2/8\pi G$  and the Planck time  $t_{\text{PL}} \sim 10^{-43}$  sec. This implies that the radius of curvature  $R_{\text{curv}}$  at the Planck time was very large compared to the Hubble radius  $R_H = c_0/H$ :

$$R_{\text{curv}}(10^{-43}\text{sec}) \sim 10^{30}R_H, \quad R_{\text{curv}}(1\text{sec}) \sim 10^8R_H. \quad (25)$$

This means that the universe in standard big-bang cosmology was very special at the Planck time. The universe has survived some  $10^{60}$  Planck times without re-collapsing or becoming curvature dominated.

We obtain from (22):

$$\Omega - 1 = \frac{Kc^2}{\dot{a}^2} - \frac{4}{H} \frac{\dot{c}}{c}, \quad (26)$$

where  $\Omega = 8\pi G\rho/3H^2$ . In order to resolve the flatness problem, we separate out the initial value  $\Omega_i - 1$  and the final value  $\Omega_0 - 1$  in (26):

$$\Omega_0 - 1 = \frac{Kc_0^2}{\dot{a}_0^2} \quad (27)$$

and

$$\Omega_i - 1 = \frac{Kc_i^2}{\dot{a}_i^2} - \frac{4}{H_i} \frac{\dot{c}_i}{c_i}. \quad (28)$$

Dividing (27) by (28) yields

$$\Omega_0 = 1 + \frac{KH_i\dot{a}_i^2c_0^2}{\dot{a}_0^2c_i^2(KH_i - 4\dot{c}_i\dot{a}_i^2/c_i^3)} |(\Omega_i - 1)|. \quad (29)$$

We have for  $c_i = c_0$ :

$$\Omega_0 = 1 + \frac{\dot{a}_i^2}{\dot{a}_0^2} |(\Omega_i - 1)|. \quad (30)$$

If  $\dot{a}_i/\dot{a}_0 < 10^{-5}$  corresponding to an early phase of inflationary acceleration and  $|\Omega_i - 1| \sim \mathcal{O}(1)$ , then  $\Omega_0 = 1$  to a high accuracy [35]. This solves the flatness problem in inflationary models<sup>3</sup>.

In our VSL model, the flatness problem can also be solved. When in an early phase:

$$c_i^3 \gg 4\dot{c}_i\dot{a}_i^2, \quad c_i^2 \gg \frac{\dot{a}_i^2c_0^2}{\dot{a}_0^2} |(\Omega_i - 1)|, \quad (31)$$

then from (29)  $\Omega_0 = 1$  is predicted to a high accuracy.

<sup>3</sup>By implementing a measure Gibbons and Turok [42] find that the probability of  $N_e$  e-folds of inflation is of order  $\exp(-3N_e)$ .

During the radiation dominated phase with  $a(t) \propto t^{1/2}$  the radiation density is given by

$$\rho_r = \frac{4\sigma}{c} T^4 = \frac{b}{c^3} T^4, \quad (32)$$

where  $\sigma$  is the Stefan-Boltzmann constant,  $b = 8\pi^5 k_B^4 / 15h^3$  and  $k_B$  and  $h$  are Boltzmann's constant and Planck's constant, respectively. Thus,  $\rho_r \rightarrow 0$  as  $c \rightarrow \infty$  and from this we can deduce that  $\Omega_i = 8\pi G \rho_r / 3H_i^2$  in (29) is sufficiently diluted, so that for a large enough initial value of  $c_i$  we have  $\Omega_0 = 1$  to a high degree of accuracy.

In the VSL model, *traces of an initial inhomogeneity will be sufficiently smoothed out* as the universe expands. However, in inflation models bubbles of inflation can be produced that do not inflate sufficiently and these will generate a non-uniformity problem as the universe expands to the present day [4].

We have adopted the scenario that in the very early universe the speed of light  $c(t)$  has a large value during a short time duration when  $t_i < t < t_c$ , and it has the value  $c = c_0$  for  $t > t_c$  when the Friedmann equations and the cosmology are described by GR.

Let us assume for our scalar fluctuation "seed" field  $\phi$  that in (11)  $V(\phi) = 0$ , yielding

$$\ddot{\phi} + 3H\dot{\phi} = 0. \quad (33)$$

The solution to this equation is

$$\dot{\phi} = \sqrt{12}B \left( \frac{a_*}{a} \right)^3, \quad (34)$$

where  $B$  is a constant and  $a_*$  is a reference value for  $a$ . For large  $\dot{\phi}$  the kinetic contribution to  $\rho_\phi \propto \frac{1}{2}(\dot{\phi})^2$  will dominate the matter densities  $\rho_M$  and  $\rho_\psi$  and the Friedmann equation (22) for  $c = c_0$  becomes

$$H^2 + \frac{Kc_0^2}{a^2} = \frac{1}{12}\dot{\phi}^2. \quad (35)$$

Substituting the solution (34), we get

$$H^2 + \frac{Kc_0^2}{a^2} = B^2 \left( \frac{a_*}{a} \right)^6. \quad (36)$$

Because the field  $\phi$  dominates in the early universe, we can neglect the spatial curvature for  $c = c_0$  and we obtain the approximate solution given by

$$a(t) = a_*(3Bt)^{1/3}, \quad H(t) = \frac{1}{3t}. \quad (37)$$

The equation of state for a massless scalar field gives for the exponent  $n$  in  $a(t) \propto t^n$  the value  $n = 2/3(1+w)$ , so for  $n = 1/3$  we get  $w = 1$ . For  $w = 1$  the kinetic term  $\frac{1}{2}(\dot{\phi})^2$  dominates the potential term  $V(\phi)$ .

We observe that there is no explicit source for the field  $\phi$  in Eq.(11), which implies that as the universe expands the field  $\phi$  becomes increasingly diluted to the point where it has unobservable effects at present. By performing a post-Newtonian expansion of the gravitational field in the solar system, we have:  $g_{00} \approx 1 + 2GM_\odot/c_0^2 1 \text{ AU}$ . If the contribution of the field  $\phi$  is to be significant in the solar system, then  $2GM_\odot/c_0^2 1 \text{ AU} \propto \dot{\phi}^2$ . However, as the universe expands we have according to (34) that  $\dot{\phi} \sim 1/a^3$ , so that the effects of the scalar field  $\phi$  will become unobservable in the present universe.

## 4 Electromagnetic-Gravitational Origin of Variable Speed of Light

We are concerned with the physics that determines the speed of light. It is inadequate to just postulate *ad hoc* that the speed of light changes without a fundamental physical reason for its change and magnitude. Moreover, we should distinguish between the speed of light  $c_{EM}$  deduced from Maxwell's vacuum equations and the speed of light  $c_{ST}$  corresponding to the constant that synthesizes space and time. The basic mechanism for a changing  $c$  can originate with a generalization of Maxwell's equations. Drummond and Hathrell [36, 37, 38, 39] applied generalized Maxwell equations to cosmology, leading to a varying speed

of light in an FLRW spacetime. They showed that light propagates at a speed greater than  $c_0$  when vacuum polarization in QED acts as a dispersive medium in the presence of tidal gravitational forces with a refractive index  $n_{\text{refrac}}$ .

The gauge invariant action for the electromagnetic field in the presence of a gravitational field is defined by

$$S_{EM} = \int d^4x \sqrt{-g} \mathcal{L}_{EM}, \quad (38)$$

where [40]:

$$\mathcal{L}_{EM} = -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} + \frac{1}{4} \sigma R F^{\mu\nu} F_{\mu\nu} + \frac{1}{2} \chi R_{\mu\nu} F^{\mu\rho} F^\nu{}_\rho + \frac{1}{4} \omega R_{\mu\nu\rho\sigma} F^{\mu\nu} F^{\rho\sigma} + \zeta \nabla_\mu F^{\mu\nu} \nabla_\sigma F^\sigma{}_\nu - j^\mu A_\mu. \quad (39)$$

We have  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ ,  $A_\mu$  and  $j^\mu$  are the electromagnetic potential and charge current density, respectively, and  $R_{\mu\nu\rho\sigma}$  is the Riemann curvature tensor. The constant parameters  $\sigma, \chi, \omega$  and  $\zeta$  have dimension [length]<sup>2</sup>. The fifth term in (39) survives in flat Minkowski space, for it represents off-mass-shell effects in the vacuum polarization. In the following, we will omit this contribution to the action and the equation of motion.

The gravitationally modified equation of motion is given by

$$\nabla_\mu F^{\mu\nu} + \nabla_\mu [4\sigma R F^{\mu\nu} + 2\chi (R^\mu{}_\sigma F^{\sigma\nu} - R^\nu{}_\sigma F^{\sigma\mu}) + 4\omega R^{\mu\nu}{}_{\sigma\tau} F^{\sigma\tau}] = 0. \quad (40)$$

The remaining Maxwell equation is

$$\nabla_\rho F_{\mu\nu} + \nabla_\mu F_{\nu\rho} + \nabla_\nu F_{\rho\mu} = 0. \quad (41)$$

In an FLRW spacetime, Drummond and Hathrell derived the expression [36]:

$$k^2 = \frac{11\xi^2}{c_0^2} \left( \frac{\dot{a}^2 + Kc_0^2}{a^2} - \frac{\ddot{a}}{a} \right) (k^\mu u_\mu)^2 = \frac{44\xi^2}{c_0^2} \left[ \pi G \left( \rho + \frac{p}{c_0^2} \right) \right] (k^\mu u_\mu)^2, \quad (42)$$

where  $k^2 = (k^0)^2 - |\mathbf{k}|^2$ ,  $(k^\mu u_\mu)^2 = (k^0)^2$  and  $c/c_0 = k^0/|\mathbf{k}|$ . Moreover, we have

$$\xi = \left( \frac{\alpha}{90\pi} \right)^{1/2} \lambda_c, \quad (43)$$

where  $\alpha = e^2/\hbar c_0^2$  and  $\lambda_c = \hbar/m_e c_0$  are the fine-structure constant and the Compton wavelength of the electron, respectively.

For the radiation dominated phase of the universe  $w = 1/3$ ,  $a \propto t^{1/2}$  and  $\rho \sim 1/a^4 \sim 1/t^2$ , while for the equation of state  $w = 1$  in the  $\phi$  field dominated phase of the universe  $a \propto t^{1/3}$  and  $\rho \sim 1/a^6 \sim 1/t^2$ . We get from (42) the speed of light:

$$c = \left( 1 - \frac{11}{2} \frac{t_*^2}{t^2} \right)^{-1/2} c_0, \quad (44)$$

where  $t_* = \xi/c_0 = 6.6 \times 10^{-24}$  sec. The speed of light  $c \rightarrow \infty$  as  $t \rightarrow \sqrt{11/2} t_*$ . This occurs for  $t \gtrsim 1.5 \times 10^{-23}$  sec after the big bang, corresponding to an energy  $\sim 2 \times 10^5$  TeV. The maximum LHC energy of 14 TeV corresponds to  $t_{\text{LHC}} \sim 4 \times 10^{-15}$  sec after the big bang. The expression (44) for  $c$  fails for  $t \lesssim 2 \times 10^{-23}$  sec, signaling the breakdown of the QED vacuum polarization calculation based on the effective Lagrangian (39). Near the time  $t \gtrsim 1.5 \times 10^{-23}$  sec, we can have a large increase in the speed of light,  $c \gtrsim 10^{30} c_0$ , adequate to solve the horizon and flatness problems and stretch the tiny quantum perturbative density fluctuations  $\delta\phi$  to super-horizon scale.

The superluminal speed of gravitational waves in the very early universe has been derived [41] by adopting a non-minimal coupling of the graviton to higher derivative curvature and the resulting tidal forces on the graviton. In the early universe near the Planck energy, the gravitational fine structure constant  $\alpha_g = Gm^2/\hbar c_0 \sim 1$ , for a given mass  $m$ , allowing for sufficiently large quantum gravity (QG) vacuum polarization contributions comparable to the those for the photon, obtained from the QED fine structure coupling strength  $\alpha = e^2/\hbar c_0$ . These QG vacuum polarization contributions will induce a dispersive medium with a refractive

index. In the radiation dominated era  $\rho \propto 1/t^2$ , we expect that near the Planck time  $t_{\text{PL}} = 5.4 \times 10^{-43}$  sec and the Planck density  $\rho_{\text{PL}} = c_0^5/\hbar G^2 = 5.2 \times 10^{93} \text{ g cm}^{-3}$ , we have

$$c_g = \left(1 - \frac{\zeta_g^2}{t^2}\right)^{-1/2} c_{g0}, \quad (45)$$

where  $\zeta_g$  is a constant determined by the QG vacuum polarization calculation using the gravitational fine structure constant  $\alpha_g = Gm^2/\hbar c_0$ . Near the Planck energy  $mc_0^2 = (\hbar c_0^5/G)^{1/2}$ , we will have  $\alpha_g \sim \mathcal{O}(1)$  and QG vacuum polarization can be comparable in size to the QED contributions. For  $t \gtrsim \zeta_g$  we can have a large increase in the speed of gravitational waves,  $c_g \gtrsim 10^{30} c_{g0}$ . This will stretch the primordial gravitational waves to a super-horizon size,  $\lambda_g = c_g/\nu_g$ , permitting them to be observed in B-polarization experiments.

It is clear that the large increases in  $c$  and  $c_g$  induced by the refractive vacuum polarization medium and the non-minimal coupling of the electromagnetic and gravitational fields to curvature, can provide a fundamental explanation for how the large speeds of light and gravitational waves in the early universe can be obtained from quantum field theory. This sets VSL cosmology on a firmer footing by removing the *ad hoc* assumption that  $c$  and  $c_g$  possess large enough increases in the early universe and justifying a VSL cosmological model.

## 5 Primordial fluctuations and gravitational waves

Let us consider in our VSL model the mechanism for the generation of cosmological fluctuations and the growth of large structures without inflation. In inflationary models, scalar quantum fluctuations oscillate until their wavelengths become equal to the Hubble radius  $R_H = c_0/H$ . When they pass beyond  $R_H$  the oscillations are damped and the fluctuation modes are “frozen” as classical fluctuations with amplitude  $\delta\phi \sim H/2\pi$  [35]. In an inflationary spacetime the wavelengths of quantum field fluctuations  $\delta\phi$  are stretched by rapid expansion:

$$\lambda_f \propto a(t) \propto \exp(Ht), \quad (46)$$

where  $H$  is approximately constant. Short-wavelength fluctuations are quickly redshifted by the inflationary expansion until their wavelengths are larger than the size of the horizon  $R_H$ .

In our VSL model, the wavelengths of the quantum field fluctuations  $\delta\phi$  are stretched by the short duration large increase of  $c$ :

$$\lambda_f = \frac{c}{\nu_f}, \quad (47)$$

where  $\nu_f$  is the frequency. The redshifted wavelengths of the short-wavelength fluctuations become larger than the horizon and are frozen in as classical fluctuations. The amplitudes of quantum modes are calculated at the horizon crossing, when the wavelength of a mode is equal to  $a/k = c_0/H$ . For an inflationary epoch with 60 e-folds of inflation of the cosmic scale  $a$ , the initial matter fluctuations and gravitational waves with wavelengths  $\lambda_i$  are stretched to their wavelengths today by the amount:

$$\lambda_0 \sim \exp(\mathcal{N})\lambda_i, \quad (48)$$

where  $\mathcal{N}$  is the number of e-folds of inflation,  $\mathcal{N} \geq 60$ . In the VSL model, the initial fluctuation wavelengths  $\lambda_i$  are stretched by an equivalent amount:

$$\lambda_0 \sim \mathcal{Q}\lambda_i, \quad (49)$$

where  $\mathcal{Q} \gtrsim 10^{30}$ .

The fluctuations are of two kinds: scalar matter quantum fluctuations and tensor gravitational wave fluctuations. Let us consider first the scalar fluctuations. We will consider a scenario in which we assume that  $V(\phi) = 0$ . The fluctuations  $\delta\phi(t, \mathbf{x})$  about a cosmological background will be considered in the comoving frame with the minimally-coupled Klein-Gordon field equation:

$$\frac{d^2\delta\phi_{\mathbf{k}}}{dt^2} + 3H\frac{d\delta\phi_{\mathbf{k}}}{dt} + \frac{c_0^2 k^2}{a^2}\delta\phi_{\mathbf{k}} = 0, \quad (50)$$

where  $k = |\mathbf{k}|$  and

$$\delta\phi(\mathbf{x}) = (2\pi)^{-2/3} \int d^3\mathbf{k} \exp[-i\mathbf{k} \cdot \mathbf{x}] \delta\phi_{\mathbf{k}}. \quad (51)$$

By substituting the solution (37) into (50), we get the equation of motion:

$$\frac{d^2\delta\phi_{\mathbf{k}}}{dt^2} + \frac{1}{t} \frac{d\delta\phi_{\mathbf{k}}}{dt} + \frac{c_0^2 k^2}{a_*^2 (3Bt)^{2/3}} \delta\phi_{\mathbf{k}} = 0. \quad (52)$$

This equation has the general Bessel function solution [28]:

$$\delta\phi_{\mathbf{k}}(y) = A_1 J_0(y) + A_2 Y_0(y), \quad (53)$$

where  $A_1$  and  $A_2$  are constant coefficients and

$$y = \frac{c_0 k}{2a_* H_\ell} \left( \frac{a}{a_*} \right)^2. \quad (54)$$

Here,  $H_\ell = (3B)^{2/3}/3a_*$  and we define  $H_\ell = c_0/\ell_0$ , where  $\ell_0$  is the length scale at which the normalized wave function is in its ground state.

We can obtain from (53) the normalized plane wave solution:

$$\delta\phi_{\mathbf{k}}(t_k) = \left( \frac{\kappa c_0^2 \hbar}{(2\pi a_k)^3 \omega_k} \right)^{1/2} \cos(\omega_k t_k - \mathbf{k} \cdot \mathbf{x} + \delta), \quad (55)$$

where  $\kappa = 16\pi G/c_0^4$  and

$$\omega_k = \frac{c_0 k}{a_k}. \quad (56)$$

This solution is approximately equivalent to the one we get from adopting a flat Minkowski spacetime. We assume that after the fluctuation modes cross the horizon they are in a classical state in the Minkowski spacetime<sup>4</sup>.

The scale at which the fluctuation mode exists is given by the condition:

$$a_k = k\ell_0. \quad (57)$$

From this condition, we get from (54):

$$y_k = \gamma k^3, \quad (58)$$

where  $y_k$  is the value of  $y$  evaluated for  $a = a_k$  and  $\gamma = \ell_0^3/2a_*^3$ .

If we assume that we should use (55) as initial data for the classical solution of (50), then we will match not only the initial perturbation, but also its time derivative. Keeping only the dominant contribution as  $y \rightarrow 0$  and  $\omega_k = c_0 k/a_k$  gives

$$\delta\phi_{\mathbf{k}} \approx \left( \frac{9\kappa c_0^2 \hbar}{(2\pi a_k)^3 32\omega_k} \right)^{1/2} \cos(\omega_k t_k - \mathbf{k} \cdot \mathbf{x} + \delta) \ln(y_k) J_0(y_k). \quad (59)$$

From (56), (58) and the Planck length,  $\ell_{\text{PL}} = (G\hbar/c_0^3)^{1/2} \sim 10^{-33}$  cm, we obtain the density perturbation power spectrum:

$$\mathcal{P}_{\delta\phi} = \frac{9}{2(2\pi)^3} \left( \frac{\ell_{\text{PL}}^2}{\ell_0^2} \right) \ln^2(y_k). \quad (60)$$

---

<sup>4</sup>Hollands and Wald [43] considered a model which did not have a VSL or inflationary mechanism to stretch the quantum fluctuation wavelengths to super-horizon lengths. They assumed that the classical fluctuations were born super-horizon, were frozen at a length scale  $\ell_0$  and were described by a plane wave in its ground state. Due to the lack of a VSL or inflationary dilution of the radiation density, the value of the radiation density at the epoch when the fluctuations were born was unphysically large [44]. As demonstrated in Eq.(32), the VSL mechanism as in the case of inflation dilutes the radiation density.



Recalling that  $\dot{\phi}^2 \sim 12H^2$ , we obtain the curvature power spectrum:

$$\mathcal{P}_{\mathcal{R}}(k) = \left(\frac{H^2}{\dot{\phi}^2}\right) \mathcal{P}_{\delta\phi} = \frac{3}{8(2\pi)^3} \left(\frac{\ell_{\text{PL}}^2}{\ell_0^2}\right) \ln^2(y_k). \quad (61)$$

Moreover, we have

$$\mathcal{P}_{\mathcal{R}}(k) = 2\pi^2 A_s \left(\frac{k}{k_*}\right)^{n_s-1}, \quad (62)$$

where  $A_s$  is the scalar fluctuation amplitude and  $n_s$  is the spectral index. The power spectrum (61) is scale invariant except for the factor  $\ln^2(y_k)$ . The non-scale invariant contribution results from matching the initial state and its time derivative, and that the Bessel function  $Y_0(y_k)$  is logarithmically divergent when  $y_k \rightarrow 0$ . This will lead to a slight deviation from a scale invariant spectrum.

The scalar mode spectral index is given by

$$n_s = 1 + \frac{d \ln \mathcal{P}_{\mathcal{R}}}{d \ln k}. \quad (63)$$

From (58) and (61), we get

$$n_s = 1 + \frac{6}{\ln(y_k)}. \quad (64)$$

The running of the spectral index is calculated from

$$\alpha_s = dn_s/d \ln k, \quad (65)$$

which yields

$$\alpha_s = -\frac{1}{2}(1 - n_s)^2. \quad (66)$$

We have in the large scale limit the anisotropic amplitude:

$$\delta_H = \frac{2}{5} \sqrt{\mathcal{P}_{\mathcal{R}}} \approx \frac{2}{5} \left(\frac{3}{8(2\pi)^3}\right)^{1/2} \left(\frac{\ell_{\text{PL}}}{\ell_0}\right) |\ln(y_k)|. \quad (67)$$

By adopting

$$\frac{2}{5} \left(\frac{3}{8(2\pi)^3}\right)^{1/2} |\ln(y_k)| \sim \mathcal{O}(1), \quad (68)$$

we have  $\delta_H \sim \ell_{\text{PL}}/\ell_0$ . Fixing the length scale  $\ell$  to be  $\ell_0 \sim 10^5 \ell_{\text{PL}} \sim 10^{-28}$  cm, which is of the order of the grand unification scale, we can match the amplitude of the observed CMB fluctuations,  $\delta_H \sim 10^{-5}$ .

For the value  $\ln(y_k) \sim -150$ , we obtain the result for the spectral index:

$$n_s \sim 0.96. \quad (69)$$

This is in good agreement with the result obtained by Planck Mission [1, 2, 3]:

$$n_s = 0.9603 \pm 0.0073. \quad (70)$$

For the running of the spectral index  $n_s$ , we get

$$\alpha_s \sim -8 \times 10^{-4}, \quad (71)$$

which is in approximate agreement with the Planck Mission result:

$$\alpha_s = -0.013 \pm 0.0090. \quad (72)$$

In inflationary models the derivation of the power spectrum and the spectral index depend sensitively on the shape of the inflaton potential  $V(\phi_{\text{inflaton}})$  and its derivatives with respect to  $\phi$ . The condition of a slow-roll potential is required to produce enough e-folds of inflation. This is not the case in our VSL derivation of the power spectrum and the spectral index. Our derivation of the scalar fluctuation power

spectrum does not depend sensitively on the shape of the potential  $V(\phi)$ . This can reduce the VSL model dependence and associated fine-tuning problems.

We now turn to the spectrum of relic gravitational waves. The conformally flat background metric is

$$ds^2 = a^2(\eta)(c^2 d\eta^2 - (dx^i)^2). \quad (73)$$

In our model the wavelength of gravity waves is given by  $\lambda_g = c_g/\nu_g$ , where  $\nu_g$  is the frequency of a gravitational wave. During the phase when  $c_g \gg c_0$  gravitational waves are generated and their wavelengths are stretched and cross the horizon. As the universe expands, the amplitude of the gravitational wave spectrum passes back into the observable universe and we can observe a B-polarization with a non-zero spectral index  $n_t$  and ratio  $r$ =tensor/scalar.

The tensor perturbations can be expressed as

$$ds^2 = a^2[c^2 d\eta^2 - (\delta_{ij} + 2h_{ij})dx^i dx^j], \quad (74)$$

where

$$h_{ij} = \int \frac{d^3\mathbf{k}}{(2\pi)^{3/2}} \sum_{\lambda=1}^2 \psi_{\mathbf{k},\lambda} e_{ij}(\mathbf{k}, \lambda \exp(i\mathbf{k} \cdot \mathbf{x})), \quad (75)$$

and where  $e_{ij}(\mathbf{k}, \lambda)$  is a polarization tensor satisfying  $e_{ij} = e_{ji}$ ,  $e_{ii} = 0$ ,  $k_i e_{ij} = 0$  and  $e_{ij}(\mathbf{k}, \lambda) e_{ij}^*(\mathbf{k}, \mu) = \delta_{\lambda\mu}$ . Moreover, we have

$$\langle \psi_{\mathbf{k},\lambda}, \psi_{\mathbf{l},\lambda}^* \rangle = 2\pi^2 \mathcal{P}_{\delta\psi} \delta(\mathbf{k} - \mathbf{l}), \quad (76)$$

where  $\mathcal{P}_{\delta\psi}$  is the gravitational wave spectrum.

The gravitational tensor component can be expressed as the superposition of two scalar polarization wave modes:

$$h_{+,\times}^{\text{tensor}} = \phi_{+,\times} = \text{scalar}, \quad (77)$$

where  $+$ ,  $\times$  refer to the longitudinal and transverse polarization modes, respectively. The modes obey the scalar equation of motion at the horizon and super-horizon scales when  $c = c_0$ :

$$\frac{d^2 \phi_{+,\times,\mathbf{k}}}{dt^2} + 3H \frac{d\phi_{+,\times,\mathbf{k}}}{dt} + \frac{c_0^2 k^2}{a^2} \phi_{+,\times,\mathbf{k}} = 0. \quad (78)$$

The sizes of the second and third terms depend on the magnitude of  $c_0 k/aH$ . If  $c_0 k/aH \rightarrow 0$ , then the gravitational wave mode is outside the horizon and we can neglect the third term and the solution to the wave equation approaches a constant. If  $c_0 k/aH$  is large, the mode is inside the horizon and the second term becomes sub-leading. The mode then undergoes a damped oscillation and decays as  $1/a$ . The tensor modes of interest are outside the horizon with constant values determined by the primordial distribution, generated at a sub-horizon scale during the phase when  $c_g$  has a large value  $c_g \gg c_0$ . During the radiation and matter domination epochs the modes gradually re-enter the horizon and damp away. Only the tensor modes which entered the horizon just before the surface of last scattering lead to important effects in the CMB.

The tensor mode power spectrum is

$$\mathcal{P}_t(k) = \mathcal{P}_\psi(k_p) \left( \frac{k}{k_p} \right)^{n_t}, \quad (79)$$

where  $k_p = 0.004 \text{ Mpc}^{-1}$  is the pivotal scale. By solving the scalar wave equation (78) in the same manner as was done for the scalar matter fluctuations, we obtain the power spectrum:

$$\mathcal{P}_t(k) = k^3 \mathcal{P}_{\phi_{+,\times}}(k) = \frac{N}{(2\pi)^3} \left( \frac{\ell_{\text{PL}}^2}{\ell_0^2} \right) \ln^2(y_k), \quad (80)$$

where  $N$  is a constant. As in the case of the scalar matter fluctuations, the gravitational wave mode fluctuations are scale invariant up to the slight scale breaking factor  $\ln^2(y_k)$ .

The tensor mode spectral index is given by

$$n_t = \frac{d \ln \mathcal{P}_t}{d \ln k}. \quad (81)$$

This yields the result

$$n_t = \frac{6}{\ln(y_k)}. \quad (82)$$

Choosing, as before for the scalar perturbations, the value  $\ln(y_k) \sim -150$  we obtain

$$n_t = -0.04. \quad (83)$$

This spectral index (tilt) is red which agrees with the standard inflationary model result when the Null Energy Condition is violated,  $\dot{H} < 0$ .

If we now adopt the BICEP2 B-polarization result  $r = 0.2_{-0.05}^{+0.07}$  [45]<sup>5</sup>, then we get  $r/n_t = -5$ , which is close to the single-field inflationary model consistency condition determined by the slow-roll parameter,  $\epsilon = -\dot{H}/H^2$ , related to the equation of state  $p = \epsilon - \rho$ . We have  $r = 16\epsilon$  and  $n_t = -2\epsilon$  giving  $r/n_t = -8$ , which is satisfied irrespective of the form of the single-field inflationary potential.

## 6 Conclusions

We have formulated a VSL model in which the homogeneous Lorentz group  $SO(3, 1)$  is spontaneously broken to the rotation group  $O(3)$  by the non-zero vacuum expectation value  $\langle 0|\psi_\mu|0\rangle$ . This determines a preferred time  $t$  in the cosmological model corresponding to the comoving time in an FLRW spacetime. The phase in the very early universe when  $c \gg c_0$  and  $c_g \gg c_{g0}$  are induced by the vacuum polarization in QED and QG, generated by a non-minimal coupling between the electromagnetic field and curvature [36, 38, 39] and by non-minimal coupling to higher derivative curvature.

In contrast to the inflationary scenario, our VSL model prediction of the almost scale invariant, Gaussian fluctuation spectra for matter and relic gravitational waves does not rely on determining the shape of a potential and its derivatives. The model can relieve the fine-tuning that is inevitably a consequence of inflationary models. Once the short duration phase of  $c \gg c_0$  and  $c_g \gg c_{g0}$  has taken place and the wavelengths of the initial primordial quantum fluctuations have been stretched through the horizon, then a classical solution of the wave equation for the fluctuations can be employed to generate the power spectrum and the spectral indices of the scalar and tensor fluctuation modes. Our VSL model can produce the non-zero scalar and tensor spectral indices  $n_s = 0.96$  and  $n_t = -0.04$ . Combined with the tensor/scalar ratio potentially observed by BICEP2,  $r = 0.2_{-0.05}^{+0.07}$ , we obtain the ratio  $r/n_t = -5$ , which is close to the single inflation field consistency condition  $r/n_t = -8$ .

Since the ordered phase in the spontaneous symmetry breaking of Lorentz invariance is at a much lower entropy than the restored, disordered symmetry phase and due to the existence of a domain determined by the direction of the vev,  $\langle 0|\psi_\mu|0\rangle$ , a natural explanation is given for the cosmological arrow of time and the origin of the second law of thermodynamics [13, 14, 47]. The ordered state of low entropy in the symmetry broken phase with  $c \gg c_0$ , becomes a state of high entropy in the symmetry restored disordered phase with  $c = c_0$ . The spontaneous symmetry breaking of the gravitational vacuum leads to a manifold with the structure  $O(3) \times R$ , in which time appears as an absolute external time parameter. The vev  $\langle 0|\psi_\mu|0\rangle$  points in a chosen direction of time to break the symmetry of the vacuum creating an arrow of time.

## Acknowledgments

I thank Andrew Liddle, Jerome Martin, Michael Clayton, Viktor Toth and Martin Green for helpful discussions. This research was generously supported by the John Templeton Foundation. Research at the Perimeter Institute for Theoretical Physics is supported by the Government of Canada through industry Canada and by the Province of Ontario through the Ministry of Research and Innovation (MRI). I thank the Institut d'Astrophysique de Paris (IAP) where this research was completed for their hospitality.

---

<sup>5</sup>The B-polarization result claiming to have observed relic gravitational waves has been disputed because of foreground dust contamination [46]. Further experimental data for B-polarization signals are required to decide whether relic gravitational waves can be observed.

## References

- [1] P. A. R. Ade et al., Planck Collaboration, arXiv:1303.5076 [astro-ph].
- [2] P. A. R. Ade et al., Planck Collaboration), arXiv:1303.5082 [astro-ph].
- [3] P. A. R. Ade et al., Planck Collaboration, arXiv:1303.5084 [astro-ph].
- [4] A. Ijjas, P. J. Steinhardt, and A. Loeb, Phys. Lett., **B723**, 261, (2013), arXiv:1304.2785 [astro-ph].
- [5] A. H. Guth, D. I. Kaiser and Y. Nomura, Phys. Lett., **B733**, 112 (2014), arXiv:1312.7619 [astro-ph].
- [6] A. Ijjas, P. J. Steinhardt and A. Loeb, arXiv:1402.6980 [astro-ph.CO].
- [7] A. H. Guth, Phys. Rev., D 23, 347 (1981).
- [8] A. Linde, Phys. Lett., B 108, 389 (1982).
- [9] A. D. Linde, *Particle Physics and Inflationary Cosmology: Contemporary Concepts in Physics*, Harwood Academic Publishers, Chur, Switzerland, 1990.
- [10] S. W. Hawking and I. Moss, Phys. Lett., B 110 35 (1982).
- [11] A. Albrecht and P. Steinhardt, Phys. Rev. Lett., 48, 1220 (1982).
- [12] James R. Liddle and David H. Lyth, *Cosmological Inflation and Large Scale Structure*, Cambridge University Press, New York, 2000.
- [13] J. W. Moffat, Int. J. Mod. Phys., **D2**, 351 (1993), arXiv:9211020 [gr-qc].
- [14] J. W. Moffat, Found. Phys., **23**, 411 (1993), arXiv:9209001 [gr-qc].
- [15] A. Albrecht and J. Maguijeo, Phys. Rev., **D59**, 043516 (1999), arXiv:9811018 [astro-ph].
- [16] J. D. Barrow, Phys. Rev., **D59**, 043515 (1999), arXiv:9811022 [astro-ph].
- [17] J. D. Barrow and J. Magueijo, Class. Quant. Grav., **16**, 1435 (1999), arXiv:9901049 [astro-ph].
- [18] J. Magueijo, Rep. on Prog. in Phys., **66** (11), 2025 (2003), arXiv:0305457 [astro-ph].
- [19] P. Teyssandier, Ann. Fond. Broglie **29**, 173 (2004) [gr-qc].
- [20] G. F. R. Ellis and J. P. Uzan, Am. J. Phys. **73**, 240 (2005).
- [21] H. Shojaie and M. Farhoudi, Can. J. Phys., **84**, 933 (2006), arXiv:0407096 [gr-qc].
- [22] G. F. R. Ellis, Gen. Rel. Grav. **39**, 511 (2007), arXiv:070375 [astro-ph].
- [23] H. Shojaie and M. Farhoudi, Can. J. Phys., **85**, 1395 (2007), arXiv:0406027 [gr-qc].
- [24] J. Magueijo and J. Moffat, Gen. Rel. Grav., **40**, 1797 (2008), arXiv:0705.4507 [gr-qc].
- [25] M. Roshan, M. Nuri and F. Shojai, Phys. Lett. **B672**, 197 (2009), arXiv:0901.3191 [gr-qc].
- [26] M. A. Clayton and J. W. Moffat, Phys. Lett., **B460**, 263 (1999), arXiv:9812481 [gr-qc].
- [27] M. A. Clayton and J. W. Moffat, Phys. Lett., **B477** 269 (2000). arXiv:9910112 [gr-qc].
- [28] M. A. Clayton and J. W. Moffat, JCAP 0307:004 (2003), arXiv:0304058 [gr-qc].
- [29] J. W. Moffat, Int. J. Mod. Phys., **D12**, 281 (2002), arXiv:0202012 [gr-gc].
- [30] J. Magueijo, Phys. Rev., **D79** 043525 (2009), arXiv:0807.1689 [gr-qc].
- [31] J. W. Moffat, Int. J. Mod. Phys., **D12**, 1279 (2003), arXiv:0211167 [hep-th].

- [32] J. W. Moffat, *Class. Quant. Grav.*, **27**, 135016 (2010), arXiv:0905.1668 [hep-th].
- [33] R. Bluhm, *The Springer Handbook of Spacetime, Springer-Verlag 2013*, arXiv:1302.1150.
- [34] R. Bluhm, Shu-Hong Fung and V. A. Kostelecký, *Phys. Rev.*, **D77**, 065020 (2008), arXiv:0712.4119 [hep-th].
- [35] V. Mukhanov, *Physical Foundations of Cosmology*, Cambridge University Press, Cambridge 2005.
- [36] I. T. Drummond and S. J. Hathrell, *Phys. Rev.*, **D22**, 343 (1980).
- [37] I. V. Khriplovich, *Phys. Lett.*, **B346**, 251 (1995).
- [38] G. M. Shore, *Nucl. Phys.*, **B460**, 379 (1996), arXiv:9504041 [gr-qc].
- [39] T. J. Hollowood and G. M. Shore, *Int. J. Mod. Phys.*, **D21**, 1241003 (2012), arXiv:1205.3291 [hep-th].  
This paper contains a list of earlier published papers by the authors.
- [40] S. Deser and P. van Nieuwenhuizen, *Phys. Rev.*, **D10**, 401 (1974).
- [41] J. W. Moffat, arXiv:1406.2609 [astro-ph.CO].
- [42] G. W. Gibbons and N. Turok, *Phys. Rev.*, **D77**, 063516 (2008), arXiv:0609095 [hep-th].
- [43] S. Hollands and R. M. Wald, *Gen. Rel. Grav.*, **34**, 2043 (2002), arXiv:0205058 [hep-th].
- [44] L. Kofman, A. Linde and V. Mukhanov, *JHEP* 0210:057 (2002), arXiv:hep-th/0206088.
- [45] P. A. R. Ade et al., BICEP2 Collaboration, *Phys. Rev. Lett.*, **112**, 241101, arXiv:1403.3985 [astro-ph].
- [46] R. Adam et al., Planck Collaboration, arXiv:1409.5738 [astro-ph.CO].
- [47] J. W. Moffat, arXiv: 1407.1026 [gr-qc].