50/500 or 100/1000 debate is not about the time frame – Reply to Rosenfeld

Richard Frankham

Department of Biological Sciences, Macquarie University, NSW 2109, Australia Tel.: +61 2 9850 8186; fax: +61 2 9850 8245. E-mail address: richard.frankham@mq.ed.au

Corey J.A. Bradshaw and Barry W. Brook

The Environment Institute and School of Earth and Environmental Sciences, The University of Adelaide, South Australia 5005, Australia E-mail addresses: corey.bradshaw@adelaide.edu.au and barry.brook@adelaide.edu.au

The Letter from Rosenfeld (2014) in response to Jamieson and Allendorf (2012) and Frankham et al. (2014) and related papers is misleading in places and requires clarification and correction, as follows:

1. "Census population size (the MVP) may be anywhere from 5 to 10 times the effective population size": this is far too low for highly fecund species, such as fish where it is typically 1000 or more (Palstra & Ruzzante 2008; Frankham et al. 2014).

2. "Frankham and associates ... goals focus on recovery targets needed for longterm persistence in perpetuity, rather than short-term prevention of extinction...": Incorrect: the $N_e = 50$ (or 100) that is a major focus of our paper (including in the title) deals explicitly with avoiding inbreeding depression in the short-term and thus avoiding immediate extinctions. Furthermore, the five-generations duration we recommend is most definitively 'short-term'. Frankham and colleagues have also done many laboratory and modelling experiments on inbreeding and extinction and reviewed the field several times, concluding that the inbreeding avoidance we recommend applies to short-term management time frames (see Frankham 2005; Frankham et al. 2010 for references).

1

3. "Frankham et al. (2014) do not appear to dispute the case made by Jamieson and Allendorf (2012) that most endangered species are rapidly declining as a consequence of human impacts ... that increase mortality and that the effects of inbreeding and reduced evolutionary potential are secondary.": this is misleading; some are, but typically it is a synergy of deterministic and stochastic effects (including genetic ones) that cannot be arbitrarily disentangled, as established by several independent approaches (described and referenced in Frankham et al. 2014). First, population viability analyses for many real, threatened species revealed that inclusion of inbreeding depression in stochastic-demographic models resulted in median reductions of 30-40% in median times to extinction – such reductions are clearly not "secondary" (Brook et al. 2002; O'Grady et al. 2006). Second, the related view that other factors typically drive species to extinction before genetic factors can impact them has been refuted, based on > 170 comparisons of genetic diversity in threatened and closely related non-threatened species (Spielman et al. 2004; Evans & Sheldon 2008; Flight 2010). Third, empirical field studies, where relative contributions could be partitioned, have also demonstrated large effects of inbreeding and loss of genetic diversity on extinction risk (Newman & Pilson 1997; Saccheri et al. 1998; Nieminen et al. 2001; Vilas et al. 2006). Fourth, gene flow into inbred populations with low genetic diversity typically leads to large genetic rescue effects on fitness, especially in natural outbreeding species (Tallmon et al. 2004; Frankham, unpublished data). Fifth, small populations of self-incompatible species with so few S alleles that they are functionally extinct become capable of sexual reproduction when outcrossing adds new S alleles, as observed in the Illinois population of the Lakeside daisy and in Florida ziziphus (DeMauro 1993; Weekley et al. 2002; Gitzendanner et al. 2012).

4. "... much of this debate concerns the time scale over which to plan for recovery and persistence ... rather than the science": while there are substantial areas of agreement between the groups, Jamieson and Allendorf disagree with our revision from 50/500 to 100/1000 (Franklin et al. 2014).

5. "However, long-term PVAs that account for evolutionary potential are not at all incompatible with PVAs over shorter time horizons that identify priority threats to persistence and interim recovery targets." We did not say otherwise.

2

6. Neither our paper, nor our previous work related to the issue (e.g., Traill et al. 2010) "... implicitly link long-term MVPs in the thousands to a triage approach that may explicitly write off extremely rare species that are costly to recover", as suggested by Jamieson and Allendorf (2012) and repeated by Rosenfeld (2014). Frankham et al. (2014) did not discuss triage, which is essentially "smart decision making" (Bottrill et al. 2008).

7. Finally, framing the question of short- or long-term management of extinction risk as mutually exclusive goals is moot if the ultimate outcome is extinction. Separating the two time frames is therefore fallacious.

References

- Bottrill, M.C., Joseph, L.N., Carwardine, J., Bode, M., Cook, C., Game, E.T.,
 Grantham, H., Kark, S., Linke, S., McDonald-Madden, E., Pressey, R.L.,
 Walker, S., Wilson, K.A., Possingham, H.P., 2008. Is conservation triage just
 smart decision making? *Trends in Ecology and Evolution* 23, 649-654.
 doi:<u>10.1016/j.tree.2008.07.007</u>
- Brook, B.W., Tonkyn, D.W., O'Grady, J.J., Frankham, R., 2002. Contribution of inbreeding to extinction risk in threatened species. *Conservation Ecology* 6(1), 16. <u>www.consecol.org/vol16/iss11/art16</u>.
- Demauro, M.M., 1993. Relationship of breeding system to rarity in the Lakeside daisy (*Hymenoxys acaulis* var. *glabra*). *Conservation Biology* 7, 542-550. doi:10.1046/j.1523-1739.1993.07030542.x
- Evans, S.R., Sheldon, B.C., 2008. Interspecific patterns of genetic diversity in birds: correlations with extinction risk. *Conservation Biology* 22, 1016-1025. doi:10.1111/j.1523-1739.2008.00972.x
- Flight, P.A., 2010. Phylogenetic comparative methods strengthen evidence for reduced genetic diversity among endangered tetrapods. *Conservation Biology* 24, 1307-1315. doi:<u>10.1111/j.1523-1739.2010.01498.x</u>
- Frankham, R., 2005. Genetics and extinction. *Biological Conservation* 126, 131-140. doi:<u>10.1016/j.biocon.2005.05.002</u>

- Frankham, R., Ballou, J.D., Briscoe, D.A., 2010. <u>Introduction to Conservation</u> <u>Genetics, 2nd edition</u>. Cambridge University Press, Cambridge, U.K.
- Frankham, R., Bradshaw, C.J.A., Brook, B.W., 2014. Genetics in conservation management: revised recommendations for the 50/500 rules, Red List criteria and population viability analyses. *Biological Conservation* 170, 56-63. doi:10.1016/j.biocon.2013.12.036
- Franklin, I.R., Allendorf, F.W., Jamieson, I.G., 2014. The 50/500 rule is still valid Reply to Frankham et al. *Biological Conservation*. doi:10.1016/j.biocon.2014.05.004
- Gitzendanner, M.A., Weekley, C.W., Germain-Aubrey, C.C., Soltis, D.E., Soltis, P.S., 2012. Microsatellite evidence for high clonality and limited genetic diversity in *Ziziphus celata* (Rhamnaceae), an endangered, self-incompatible shrub endemic to the Lake Wales Ridge, Florida, USA. *Conservation Genetics* 13, 223-234. doi:10.1007/s10592-011-0287-9
- Jamieson, I.G., Allendorf, F.W., 2012. How does the 50/500 rule apply to MVPs? *Trends in Ecology and Evolution* 27, 578-584. doi:<u>10.1016/j.tree.2012.07.001</u>
- Newman, D., Pilson, D., 1997. Increased probability of extinction due to decreased genetic effective population size: experimental populations of *Clarkia pulchella*. *Evolution* 51, 354-362. doi:10.2307/2411107
- Nieminen, M., Singer, M.C., Fortelius, W., Schöps, K., Hanksi, I., 2001. Experimental confirmation that inbreeding depression increases extinction risk in butterfly populations. *American Naturalist* 157, 237-244. doi:<u>10.1086/318630</u>
- O'Grady, J.J., Brook, B.W., Reed, D.H., Ballou, J.D., Tonkyn, D.W., Frankham, R., 2006. Realistic levels of inbreeding depression strongly affect extinction risk in wild populations. *Biological Conservation* 133, 42-51.
 doi:10.1016/j.biocon.2006.05.016
- Palstra, F.P., Ruzzante, D.E., 2008. Genetic estimates of contemporary effective population size: what can they tell us about the importance of genetic stochasticity for wild population persistence? *Molecular Ecology* 17, 3428-3447. doi:10.1111/j.1365-294X.2008.03842.x
- Rosenfeld, J.S., 50/500 or 100/1000? Reconciling short- and long-term recovery targets and MVPs. *Biological Conservation*. doi:<u>10.1016/j.biocon.2014.05.005</u>

- Saccheri, I., Kuussaari, M., Kankare, M., Vikman, P., Fortelius, W., Hanski, I., 1998. Inbreeding and extinction in a butterfly metapopulation. *Nature* 392, 491-494. doi:<u>10.1038/33136</u>
- Spielman, D., Brook, B.W., Frankham, R., 2004. Most species are not driven to extinction before genetic factors impact them. *Proceedings of the National Academy of Sciences of the USA* 101, 15261-15264. doi:10.1073/pnas.0403809101
- Tallmon, D.A., Luikart, G., Waples, R.S., 2004. The alluring simplicity and the complex reality of genetic rescue. *Trends in Ecology and Evolution* 19, 489-496. doi:10.1016/j.tree.2004.07.003
- Traill, L.W., Brook, B.W., Frankham, R., Bradshaw, C.J.A., 2010. Pragmatic population viability targets in a rapidly changing world. *Biological Conservation* 143, 28-34. doi:<u>10.1016/j.biocon.2009.09.001</u>
- Vilas, C., Miguel, E.S., Amaro, R., Garcia, C., 2006. Relative contribution of inbreeding depression and eroded adaptive diversity to extinction risk in small populations of shore campion. *Conservation Biology* 20, 229-238. doi:<u>10.1111/j.1523-1739.2005.00275.x</u>
- Weekley, C.W., Kubisiak, T.L., Race, T.M., 2002. Genetic impoverishment and cross-incompatibility in remnant genotypes of *Ziziphus celata* (Rhamnaceae), a rare shrub endemic to the Lake Wales Ridge, Florida. *Biodiversity and Conservation* 11, 2027-2046. doi:<u>10.1023/A:1020810800820</u>