L_p -CONVERGENCE OF HIGHER ORDER HERMITE OR HERMITE-FEJÉR INTERPOLATION POLYNOMIALS WITH EXPONENTIAL-TYPE WEIGHTS

HEE SUN JUNG¹ AND RYOZI SAKAI²

ABSTRACT. Let $\mathbb{R} = (-\infty, \infty)$, and let $Q \in C^1(\mathbb{R}) : \mathbb{R} \to \mathbb{R}^+ = [0, \infty)$ be an even function, which is an exponent. We consider the weight $w_\rho(x) =$ $|x|^{\rho}e^{-Q(x)}, \rho \ge 0, x \in \mathbb{R}$, and then we can construct the orthonormal polynomials $p_n(w_\rho^2; x)$ of degree n for $w_\rho^2(x)$. In this paper we obtain L_p -convergence theorems of even order Hermite-Fejér interpolation polynomials at the zeros $\{x_{k,n,\rho}\}_{k=1}^n$ of $p_n(w_\rho^2; x)$.

MSC; 41A05, 41A10

Keywords; higher order Hermite-Fejér interpolation polynomials

1. INTRODUCTION

Let $\mathbb{R} = (-\infty, \infty)$, and let $Q \in C^1(\mathbb{R}) : \mathbb{R} \to \mathbb{R}^+ = [0, \infty)$ be an even function. Consider the weight $w(x) = \exp(-Q(x))$ and define for $\rho > -\frac{1}{2}$,

(1.1)
$$w_{\rho}(x) := |x|^{\rho} w(x), \quad x \in \mathbb{R}$$

Suppose that $\int_0^\infty x^n w_\rho^2(x) dx < \infty$ for all $n = 0, 1, 2, \ldots$ Then, we can construct the orthonormal polynomials $p_{n,\rho}(x) = p_n(w_\rho^2; x)$ of degree *n* for $w_\rho^2(x)$, that is,

$$\int_{-\infty}^{\infty} p_{n,\rho}(x) p_{m,\rho}(x) w_{\rho}^{2}(x) dx = \delta_{mn} (\text{Kronecker's delta}).$$

Denote

$$p_{n,\rho}(x) = \gamma_n x^n + \dots, \gamma_n = \gamma_{n,\rho} > 0,$$

and the zeros of $p_{n,\rho}(x)$ by

$$-\infty < x_{n,n,\rho} < x_{n-1,n,\rho} < \ldots < x_{2,n,\rho} < x_{1,n,\rho} < \infty.$$

Let \mathcal{P}_n denote the class of polynomials with degree at most n. For $f \in C(\mathbb{R})$ we define the higher order Hermite-Fejér interpolation polynomial $L_n(\nu, f; x)$ based at the zeros $\{x_{k,n,\rho}\}_{k=1}^n$ as follows:

$$L_n^{(i)}(\nu, f; x_{k,n,\rho}) = \delta_{0,i} f(x_{k,n,\rho}) \text{ for } k = 1, 2, \dots, n \text{ and } i = 0, 1, \dots, \nu - 1.$$

 $L_n(1, f; x)$ is the Lagrange interpolation polynomial, $L_n(2, f; x)$ is the ordinary Hermite-Fejér interpolation polynomial, and $L_n(4, f; x)$ is the Krilov-Stayermann

Date: August 21, 2018.

polynomial. The fundamental polynomials $h_{k,n,\rho}(\nu; x) \in \mathcal{P}_{\nu n-1}$ for the higher order Hermite-Fejér interpolation polynomial $L_n(\nu, f; x)$ are defined as follows:

(1.2)
$$\begin{aligned} h_{k,n,\rho}(\nu;x) &= l_{k,n,\rho}^{\nu}(x) \sum_{i=0}^{\nu-1} e_i(\nu,k,n) (x - x_{k,n,\rho})^i, \\ l_{k,n,\rho}(x) &= \frac{p_n(w_{\rho}^2;x)}{(x - x_{k,n,\rho})p'_n(w_{\rho}^2;x_{k,n,\rho})}, \\ h_{k,n,\rho}(\nu;x_{p,n,\rho}) &= \delta_{k,p}, \quad h_{k,n,\rho}^{(i)}(\nu;x_{p,n,\rho}) = 0, \\ k,p &= 1, 2, \dots, n, i = 1, 2, \dots, \nu - 1. \end{aligned}$$

Using them, we can write $L_n(\nu, f; x)$ as follows:

$$L_n(\nu, f; x) = \sum_{k=1}^n f(x_{k,n,\rho}) h_{k,n,\rho}(\nu; x).$$

Furthermore, we extend the operator $L_n(\nu, f; x)$. Let l be a non-negative integer, and let $l \leq \nu - 1$. For $f \in C^l(\mathbb{R})$ we define the (l, ν) -order Hermite-Fejér interpolation polynomials $L_n(l, \nu, f; x) \in \mathcal{P}_{\nu n-1}$ as follows: For each $k = 1, 2, \ldots, n$,

$$L_n(l,\nu,f;x_{k,n,\rho}) = f(x_{k,n,\rho}),$$

$$L_n^{(j)}(l,\nu,f;x_{k,n,\rho}) = f^{(j)}(x_{k,n,\rho}), \quad j = 1, 2, \dots, l,$$

$$L_n^{(j)}(l,\nu,f;x_{k,n,\rho}) = 0, \quad j = l+1, l+2, \dots, \nu-1.$$

Especially, $L_n(0,\nu, f; x)$ is equal to $L_n(\nu, f; x)$, and for each $P \in \mathcal{P}_{\nu n-1}$ we see $L_n(\nu-1,\nu,P;x) = P(x)$. The fundamental polynomials $h_{s,k,n,\rho}(l,\nu;x) \in \mathcal{P}_{\nu n-1}, k = 1, 2, \ldots, n$, of $L_n(l,\nu,f;x)$ are defined by

(1.3)
$$\begin{aligned} h_{s,k,n,\rho}(l,\nu;x) &= l_{k,n,\rho}^{\nu}(x) \sum_{i=s}^{\nu-1} e_{s,i}(\nu,k,n)(x-x_{k,n,\rho})^i, \\ h_{s,k,n,\rho}^{(j)}(l,\nu;x_{p,n,\rho}) &= \delta_{s,j}\delta_{k,p}, j, s = 0, 1, \dots, \nu-1, p = 1, 2, \dots, n. \end{aligned}$$

Then we have

$$L_n(l,\nu,f;x) = \sum_{k=1}^n \sum_{s=0}^l f^{(s)}(x_{k,n,\rho}) h_{s,k,n,\rho}(l,\nu;x).$$

For the ordinary Hermite and Hermite-Fejér interpolation polynomial $L_n(1, 2, f; x)$, $L_n(2, f; x)$ and the related approximation process, Lubinsky [15] gave some interesting convergence theorems.

In [7] we obtained uniform convergence theorems with respect to the interpolation polynomials $L_n(\nu, f; x)$ and $L_n(l, \nu, f; x)$ with even integer ν . In this paper we will give the L_p -convergence theorems of $L_n(\nu, f; x)$ and $L_n(l, \nu, f; x)$ with even order ν . If we consider the higher order Hermite-Fejér interpolation polynomial $L_n(\nu, f; x)$ on a finite interval, then we can see a remarkable difference between the cases of an odd number ν and of an even number ν , for example, as between the Lagrange interpolation polynomial $L_n(1, f; x)$ and the Hermite-Fejér interpolation polynomial $L_n(2, f; x)$ ([17]-[22]). We can also see a similar phenomenon in the cases of the infinite intervals ([7]-[14]). In this paper, we consider the even case in L_p -norm. We will discuss the odd case in L_p -norm elsewhere.

Here, we give the class of weights which is treated in this paper. We say that $f : \mathbb{R} \to \mathbb{R}^+$ is quasi-increasing if there exists C > 0 such that $f(x) \leq Cf(y), 0 < x < y$. In the following, we introduce the class of weights defined in [16].

Definition 1.1 (see [16]). Let $Q : \mathbb{R} \to \mathbb{R}^+$ be a continuous even function satisfying the following properties:

(a) Q'(x) is continuous in \mathbb{R} and Q(0) = 0.

- (b) Q''(x) exists and is positive in $\mathbb{R} \setminus \{0\}$.
- (c) $\lim_{x\to\infty} Q(x) = \infty$.
- (d) The function

$$T(x) := \frac{xQ'(x)}{Q(x)}, \quad x \neq 0$$

is quasi-increasing in $(0,\infty)$, with

$$T(x) \ge \Lambda > 1, \quad x \in \mathbb{R}^+ \setminus \{0\}.$$

(e) There exists $C_1 > 0$ such that

$$\frac{Q^{\prime\prime}(x)}{|Q^{\prime}(x)|} \le C_1 \frac{|Q^{\prime}(x)|}{Q(x)}, \quad a.e. \quad x \in \mathbb{R} \setminus \{0\}.$$

Then we say that $w = \exp(-Q)$ is in the class $\mathcal{F}(C^2)$. Besides, if there exists a compact subinterval $J(\ni 0)$ of \mathbb{R} and $C_2 > 0$ such that

$$\frac{Q''(x)}{|Q'(x)|} \ge C_2 \frac{|Q'(x)|}{Q(x)}, \quad a.e. \quad x \in \mathbb{R} \setminus J,$$

then we say that $w = \exp(-Q)$ is in the class $\mathcal{F}(C^2+)$. If T(x) is bounded, then w is called a Freud-type weight, and if T(x) is unbounded, then w is Erdös-type weight.

Some typical examples in $\mathcal{F}(C^2+)$ are given as follows:

Example 1.2 (see [16]). (1) For $\alpha > 1$ and a non-negative integer ℓ , we put

$$Q(x) = Q_{\ell,\alpha}(x) := \exp_{\ell}(|x|^{\alpha}) - \exp_{\ell}(0),$$

where for $\ell \ge 1$,

$$\exp_{\ell}(x) := \exp(\exp(\exp(\cdots \exp x) \dots)) \quad (\ell\text{-times})$$

and $\exp_0(x) := x$.

(2) For $m \ge 0$, $\alpha \ge 0$ with $\alpha + m > 1$, we put

(1.4)
$$Q(x) = Q_{\ell,\alpha,m}(x) := |x|^m \{ \exp_\ell(|x|^\alpha) - \alpha^* \exp_\ell(0) \},$$

where for $\ell > 0$ we suppose $\alpha^* = 0$ if $\alpha = 0$; $\alpha^* = 1$ otherwise. For $\ell = 0$ we suppose m > 1 and $\alpha = 0$. Note that $Q_{\ell,0,m}$ is a Freud-type weight. (3) For $\alpha > 1$, we put

$$Q(x) = Q_{\alpha}(x) := (1 + |x|)^{|x|^{\alpha}} - 1.$$

To consider the higher order Hermite-Fejér interpolation polynomial we define a class of further strengthened weights for $\nu \ge 2$ than Definition 1.1 as follows:

Definition 1.3 (cf.[7]). Let $w(x) = \exp(-Q(x)) \in \mathcal{F}(C^2+)$, and let $\nu \ge 2$ be an integer. Assume that Q(x) is a ν -times continuously differentiable function on \mathbb{R} and satisfies the following:

(a) $Q^{(\nu+1)}(x)$ exists and $Q^{(i)}(x), 0 \le i \le \nu+1$, are positive for x > 0.

(b) There exist constants $C_i > 0$ such that

$$\left|Q^{(i+1)}(x)\right| \le C_i \left|Q^{(i)}(x)\right| \frac{|Q'(x)|}{Q(x)}, \quad x \in \mathbb{R} \setminus \{0\}, \quad i = 1, 2, \dots \nu.$$

(c) There exist $0 \le \delta < 1$ and $c_1 > 0$ such that

(1.5)
$$Q^{(\nu+1)}(x) \le C\left(\frac{1}{x}\right)^{\delta}, \quad x \in (0, c_1].$$

Then we say that $w(x) = \exp(-Q(x))$ is in the class $\mathcal{F}_{\nu}(C^2+)$. (d) Suppose one of the following:

- (d-1) Q'(x)/Q(x) is quasi-increasing on a certain positive interval $[c_2, \infty)$.
- (d-2) $Q^{(\nu+1)}(x)$ is non-decreasing on a certain positive interval $[c_2, \infty)$.
- (d-3) There exist constants C > 0 and $0 \le \delta < 1$ such that $Q^{(\nu+1)}(x) \le C(1/x)^{\delta}$ on $(0, \infty)$.

Then we write $w(x) = \exp(-Q(x)) \in \tilde{\mathcal{F}}_{\nu}(C^2+).$

In this paper we treat the weight $w_{\rho}(x) = |x|^{\rho} \exp(-Q(x)), \ \rho \ge 0$ of type (1.1). For $w_{\rho}(x) = |x|^{\rho} \exp(-Q(x)), \ w(x) = \exp(-Q(x)) \in \tilde{\mathcal{F}}_{\nu}(C^2+)$, we write $w_{\rho} \in \tilde{\mathcal{F}}_{\nu,\rho}(C^2+)$.

In the following, we give specific examples of $w \in \tilde{\mathcal{F}}_{\nu}(C^2+)$.

Example 1.4 (cf.[6, Theorem 3.1]). Let ν be a positive integer, and let $Q_{\ell,\alpha,m}$ be defined in (1.4).

(1) Let *m* and α be non-negative even integers with $m + \alpha > 1$. Then $w(x) = \exp(-Q_{\ell,\alpha,m}) \in \mathcal{F}_{\nu}(C^2+)$, and one has the following.

(a) If $\ell > 0$, then we see that $Q'_{\ell,\alpha,m}(x)/Q_{\ell,\alpha,m}(x)$ is quasi-increasing on a certain positive interval (c_1, ∞) , and $Q_{\ell,0,m}(x)$ is non-decreasing on $(0, \infty)$.

(b) If $\ell = 0$, then we see that $Q_{0,0,m}(x), m \ge 2$, is non-decreasing on $(0,\infty)$. Hence $w(x) = \exp(-Q_{\ell,\alpha,m}) \in \tilde{\mathcal{F}}_{\nu}(C^2+)$. (2) Let $m + \alpha - \mu \ge 0$. Then $w(x) = \exp(-Q_{\ell,\alpha}) \in \mathcal{F}(C^2+)$ and one has the

(2) Let $m + \alpha - \nu > 0$. Then $w(x) = \exp(-Q_{\ell,\alpha,m}) \in \mathcal{F}_{\nu}(C^2+)$, and one has the following.

(c) If $\ell \ge 2$ and $\alpha > 0$, then there exists a constant $c_1 > 0$ such that $Q'_{\ell,\alpha,m}(x)/Q_{\ell,\alpha,m}(x)$ is quasi-increasing on (c_1, ∞) .

(d) Let $\ell = 1$. If $\alpha \ge 1$, then there exists a constant $c_2 > 0$ such that $Q'_{1,\alpha,m}(x)/Q_{1,\alpha,m}(x)$ is quasi-increasing on (c_2, ∞) , and if $0 < \alpha < 1$, then $Q'_{1,\alpha,m}(x)/Q_{1,\alpha,m}(x)$ is quasi-decreasing on (c_2, ∞) .

(e) Let $\ell = 1$, and $0 < \alpha < 1$, then $Q_{1,\alpha,m}^{(\nu+1)}(x)$ is non-decreasing on a certain positive interval on (c_2, ∞) . Hence $w(x) = \exp(-Q_{\ell,\alpha,m}) \in \tilde{\mathcal{F}}_{\nu}(C^2+)$.

Example 1.5 ([6, Theorem 3.5]). Let ν be a positive integer and $\alpha > \nu$. Then $w(x) = \exp(-Q_{\alpha}(x))$ belongs to $\mathcal{F}_{\nu}(C^2+)$. Moreover, there exists a positive constant $c_2 > 0$ such that $Q'_{\alpha}(x)/Q_{\alpha}(x)$ is quasi-increasing on (c_2, ∞) . Hence $w(x) = \exp(-Q_{\alpha}) \in \tilde{\mathcal{F}}_{\nu}(C^2+)$.

In Section 2, we report the L_p -convergence theorems. We write some lemmas to prove the theorems and then we prove them in Section 3.

In what follows we abbreviate several notations as $x_{k,n} := x_{k,n,\rho}, h_{kn}(x) := h_{k,n,\rho}(\nu, x), \ l_{kn}(x) := l_{k,n,\rho}(x), \ h_{skn}(x) := h_{s,k,n,\rho}(\nu, x) \ \text{and} \ p_n(x) := p_{n,\rho}(x) \ \text{if}$ there is no confusion. For arbitrary nonzero real valued functions f(x) and g(x), we write $f(x) \sim g(x)$ if there exist constants $C_1, C_2 > 0$ independent of x such that $C_1g(x) \leq f(x) \leq C_2g(x)$ for all x. For arbitrary positive sequences $\{c_n\}_{n=1}^{\infty}$ and $\{d_n\}_{=1}^{\infty}$ we define $c_n \sim d_n$ similarly.

2. Theorems

We use the following notations.

(1) Mhaskar-Rakhmanov-Saff numbers a_x :

$$x = \frac{2}{\pi} \int_0^1 \frac{a_x u Q'(a_x u)}{(1 - u^2)^{1/2}} du, \quad x > 0.$$

(2)

(2.1)
$$\varphi_u(x) = \begin{cases} \frac{|x|}{u} \frac{1 - \frac{|x|}{a_{2u}}}{\sqrt{1 - \frac{|x|}{a_u} + \delta_u}}, & |x| \le a_u; \\ \varphi_u(a_u), & a_u < |x|, \end{cases}$$

where

$$\delta_u = (uT(a_u))^{-2/3}, \quad u > 0.$$

In the rest of this paper, we assume the following:

Assumption 2.1. Let the weight $w_{\rho} \in \tilde{\mathcal{F}}_{\nu,\rho}(C^2+), \ \rho \ge 0$. (a) If T(x) is bounded, then we suppose that for some C > 0

$$Q(x) \ge C|x|^2$$

and for δ in (1.5),

$$(2.2) a_n \le C n^{1/(1+\nu-\delta)}.$$

(b) There exist $0 \le \gamma < 1$ and $C(\gamma) > 0$ such that

(2.3)
$$T(a_n) \le C(\gamma) n^{\gamma}$$

here, if T(x) is bounded, that is, the weight w is a Freud weight, then we set $\gamma = 0$. We put

(2.4)
$$\varepsilon_n := \begin{cases} \frac{a_n}{n}, & \frac{T(a_n)}{a_n} < 1; \\ \frac{1}{n^{1-\gamma}}, & \frac{T(a_n)}{a_n} \ge 1. \end{cases}$$

Lemma 2.2 ([4, Theorem 1.4]). (1) Let $w = \exp(-Q) \in \mathcal{F}(C^2)$, and let T(x) be unbounded. Then, for any $\eta > 0$ there exists $C(\eta) > 0$ such that for $t \ge 1$,

(2.5)
$$a_t \le C(\eta) t^{\eta}.$$

(2) Let $\lambda := C_1$ be the constant in Definition 1.1 (e), that is,

$$\frac{Q''(x)}{Q'(x)} \le \lambda \frac{Q'(x)}{Q(x)}, \quad a.e. \quad x \in \mathbb{R} \setminus \{0\}.$$

If $1 < \lambda$, and if η is defined in (2.5), then there exists $C(\lambda, \eta)$ such that

$$T(a_t) \le C(\lambda, \eta) t^{\frac{2(\eta+\lambda-1)}{\lambda+1}},$$

and if $\lambda \leq 1 + \eta$, then there exists $C(\lambda, \eta)$ such that

(2.6)
$$T(a_t) \le C(\lambda, \eta) t^{4\eta}, \quad t \ge 1.$$

Remark 2.3. (1) If T(x) is unbounded, then for any L > 0 we have $Q(x) \ge C|x|^L([24, \text{Lemma 3.2}])$ and (2.2) holds ([4, Theorem 1.4]).

(2) (2.3) holds for

$$\gamma = \frac{2(\eta + \lambda - 1)}{\lambda + 1}, \quad 0 < \lambda < 3.$$

(3) Let us denote the MRS-number a_n and the function T(x) for $Q_{r,\alpha,m}$ by $a_{n,r,\alpha,m}$ and $T_{r,\alpha,m}$ respectively. Then $T_{r,\alpha,m}(a_{n,r,\alpha,m})$ satisfies (2.3). In fact, we obtain from [23, Proposition 4] and [16, Example 2 and (1.34)],

$$a_{n,r,\alpha,m} \sim (1 + \log_r^+ n)^{1/\alpha}$$

where

$$\log_r^+(x) = \begin{cases} \log(\log(\log(\ldots\log x))) \ (r \text{ times}), & \text{ if } x > \exp_r(0), \\ 0, & \text{ otherwise }, \end{cases}$$

and for every $0 < \gamma < 1$,

$$T_{r,\alpha,m}(a_{n,r,\alpha,m}) \sim (1 + \log_r^+ n)(1 + \log_{r-1}^+ n)(1 + \log^+ n) < n^{\gamma}$$

by

$$T_{r,\alpha,m}(a_{n,r,\alpha,m}) = m + T_{r,\alpha,0}(a_{n,r,\alpha,m}).$$

(4) If the weight $w = \exp(-Q) \in \mathcal{F}(C^2+)$ satisfies

$$\mu = \lim \inf_{x \to \infty} \frac{Q(x)Q''(x)}{(Q'(x))^2} = \lim \sup_{x \to \infty} \frac{Q(x)Q''(x)}{(Q'(x))^2},$$

then we say that the weight w is regular. For the regular weight w we have

$$\mu = \lim_{x \to \infty} \frac{Q(x)Q''(x)}{(Q'(x))^2} = 1,$$

therefore, we obtain (2.6) for any fixed $\eta > 0$ (see [24, Corollary 5.5]). We note that all of examples in Example 1.4 are regular. (5) (2.5) means

$$0 < C \le \frac{n}{a_n} \left(\frac{T(a_n)}{a_n}\right)^{\nu-1}.$$

Let

$$X_{n}(\nu, f; x) := \sum_{k=1}^{n} f(x_{k,n}) l_{kn}^{\nu}(x) \sum_{i=0}^{\nu-2} e_{i}(\nu, k, n) (x - x_{k,n})^{i},$$

$$Y_{n}(\nu, f; x) := \sum_{k=1}^{n} f(x_{k,n}) l_{kn}^{\nu}(x) e_{\nu-1}(\nu, k, n) (x - x_{k,n})^{\nu-1},$$

$$Z_{n}(l, \nu, f; x) := \sum_{k=1}^{n} \sum_{s=1}^{l} f^{(s)}(x_{k,n}) l_{kn}^{\nu}(x) \sum_{i=s}^{\nu-1} e_{si}(\nu, k, n) (x - x_{k,n})^{i}.$$

Then we know

$$L_{n}(\nu, f; x) = X_{n}(\nu, f; x) + Y_{n}(\nu, f; x),$$

and

$$L_n(l, \nu, f; x) = L_n(\nu, f; x) + Z_n(l, \nu, f; x).$$

Let

(2.7)
$$\Phi(x) := \frac{1}{(1+Q(x))^{2/3}T(x)}$$

$$\Phi(x) \sim \frac{Q(x)^{\frac{1}{3}}}{xQ'(x)}.$$

Moreover, if we define

(2.8)
$$\Phi_n(x) := \max\left\{\delta_n, 1 - \frac{|x|}{a_n}\right\}, \quad n = 1, 2, 3, ...,$$

then we have the following:

Lemma 2.4 ([7, Lemma 3.4]). For $x \in \mathbb{R}$ we have

$$\Phi(x) \le C\Phi_n(x), n \ge 1.$$

We have a chain of results as follows. In this section, we let $w_{\rho} \in \tilde{\mathcal{F}}_{\nu,\rho}(C^2+)$, $\rho \ge 0$, and we suppose Assumption 2.1. In addition, we let $\alpha > 0, \Delta > -1$, $1 . Let <math>C_f$ be a positive constant depending only on f, and let C > 0 be a constant.

Proposition 2.5. Let $\nu = 2, 3, 4, ...,$ and let

(2.9)
$$\Delta \ge \frac{1}{p} - \min\left\{1, \alpha\right\}.$$

For $f \in C(\mathbb{R})$ satisfying

(2.10)
$$|f(x)|(1+|x|)^{\alpha} \left\{ \Phi^{-\frac{3}{4}}(x)w_{\rho}(x) \right\}^{\nu} \leq C_f, \quad x \in \mathbb{R},$$

we have

(2.11)
$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} X_n(\nu, f; x) \right\|_{L_p(\mathbb{R})} \le CC_f.$$

Proposition 2.6. Let $\nu = 2, 4, 6, ...$ and assume that (2.9) holds. For $f \in C(\mathbb{R})$ satisfying

$$(2.12) \quad |f(x)|(1+|x|)^{\alpha} \left\{ \Phi^{-\frac{3}{4}}(x)w_{\rho}(x) \right\}^{\nu} \left(|Q'(x)| + \frac{1}{|x|} \right) \le C_f, \quad x \in \mathbb{R} \setminus \{0\}$$

we have

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} Y_n(\nu, f; x) \right\|_{L_p(\mathbb{R})} \le CC_f \varepsilon_n(a_n + \log n),$$

where ε_n is defined by (2.4).

Proposition 2.7. Let $\nu = 2, 3, 4, ...$ and assume that (2.9) holds. For $f \in C^{l}(\mathbb{R})$, $0 \leq l \leq \nu - 1$ satisfying

(2.13)
$$|f^{(s)}(x)|(1+|x|)^{\alpha} \left\{ \Phi^{-\frac{3}{4}}(x)w_{\rho}(x) \right\}^{\nu} \leq C_f, \quad x \in \mathbb{R}, \ s = 1, 2, ..., l,$$

we have

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} Z_n(l,\nu,f;x) \right\|_{L_p(\mathbb{R})}$$

$$\leq C_f \frac{a_n^2 \log n}{n} \left\{ \begin{array}{cc} 1, & \Delta p < 1; \\ \frac{\log a_n}{a_n}, & \Delta p = 1; \\ \frac{1}{a_n}, & \Delta p > 1. \end{array} \right.$$

Proposition 2.8. Let $\nu = 2, 3, 4, ...$ and assume that (2.9) holds. Let $P \in \mathcal{P}_{\nu n-1}$ be fixed. Then, we have

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} \left(L_n(\nu, P; x) - P(x) \right) \right\|_{L_p(\mathbb{R})} \to 0 \quad as \quad n \to \infty.$$

Proposition 2.9. Let $\nu = 2, 4, 6, ...$ and assume that (2.9) holds. Let $P \in \mathcal{P}_{\nu n-1}$ be fixed. Then, we have

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} (X_n(\nu, P; x) - P(x)) \right\|_{L_p(\mathbb{R})} \to 0 \quad as \quad n \to \infty.$$

Remark 2.10. Let $f \in C(\mathbb{R})$ satisfy that for given $0 < \eta < 1$ and $\alpha \ge 0$, (2.14)

$$|f(x)|(1+|x|)^{\alpha}w^{\nu-\eta}(x)\left\{|Q'(x)|+\frac{1}{|x|}\right\} \le C_f, \quad x \in \mathbb{R} \quad \text{and} \quad \lim_{x \to 0} \frac{f(x)}{x} \le C_f,$$

where C_f is a constant depending only on f. Then,

(1) (2.10) and (2.12) hold.

(2) Let $f \in C^{l}(\mathbb{R})$ for a certain $0 \leq l \leq \nu - 1$. Then (2.13) holds.

(3) Let (2.14) be satisfied, then we see f(0) = 0.

Theorem 2.11. Let $\nu = 2, 4, 6, ...$ and assume that (2.9) holds. For f(x) satisfying (2.14) we have

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} \left(L_n(\nu, f; x) - f(x) \right) \right\|_{L_p(\mathbb{R})} \to 0 \quad as \quad n \to \infty.$$

Theorem 2.12. Let $\nu = 2, 4, 6, ...$ and assume that (2.9) holds. For f(x) satisfying (2.14) and (2.13), we have

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} \left(L_n(l,\nu,f;x) - f(x) \right) \right\|_{L_p(\mathbb{R})} \to 0 \quad as \quad n \to \infty.$$

Remark 2.13. From the formulas (2.2), (2.4) and (2.5), we have

$$\varepsilon_n(a_n + \log n) \to 0$$
 as $n \to \infty$,

and

$$\frac{a_n^2 \log n}{n} \begin{cases} 1, & \Delta p < 1;\\ \frac{\log a_n}{a_n}, & \Delta p = 1; \to 0 \quad \text{as} \quad n \to \infty.\\ \frac{1}{a_n}, & \Delta p > 1 \end{cases}$$

3. Lemmas and Proof of Theorems

For the coefficients $e_{si}(\nu, k, n)(e_i(\nu, k, n) := e_{0i}(\nu, k, n))$ in (1.2) or (1.3) we have the following estimates.

Lemma 3.1 ([3, Theorem 2.6]). Let $w(x) = \exp(-Q(x)) \in \mathcal{F}(C^2+)$. We have the following. For each $s = 0, 1, ..., \nu - 1$ and $i = s, s + 1, ..., \nu - 1$,

$$e_0(\nu, k, n) = 1, \quad |e_{si}(\nu, k, n)| \le C \left\{ \frac{n}{(a_{2n}^2 - x_{k,n}^2)^{1/2}} \right\}^{i-s}$$

•

Lemma 3.2 ([5, Theorem 4.4 and Lemma 3.7 (3.20)]). Let $w_{\rho} \in \tilde{\mathcal{F}}_{\nu,\rho}(C^2+)$. If $x_{kn} \neq 0$ and $|x_{k,n}| \leq a_n(1+\delta_n)$, then $e_0(\nu, k, n) = 1$ and for $i = 1, 2, ..., \nu - 1$,

$$|e_i(\nu,k,n)| \le C \left\{ \frac{T(a_n)}{a_n} + |Q'(x_{k,n})| + \frac{1}{x_{k,n}} \right\}^{\langle i \rangle} \left\{ \frac{n}{a_{2n} - |x_{k,n}|} + \frac{T(a_n)}{a_n} \right\}^{i - \langle i \rangle},$$

where

$$\langle i \rangle = \begin{cases} 1, & \text{if } i \text{ is odd,} \\ 0, & \text{if } i \text{ is even.} \end{cases}$$

For $x_{kn} = 0$, we see $e_0(\nu, k, n) = 1$ and

$$|e_i(\nu,k,n)| \le C\left(\frac{n}{a_n}\right)^i \quad i = 1, 2, \dots, \nu - 1.$$

We have the following.

Lemma 3.3 ([1, Theorem 2.4]). Let $w_{\rho} \in \tilde{\mathcal{F}}_{\nu,\rho}(C^2+)$ ($\rho \ge 0$), $0 , and <math>\beta \in \mathbb{R}$. Then given r > 1, there exist $C, n_0, \alpha > 0$ such that for $n \ge n_0$ and $P \in \mathcal{P}_n$,

$$\left\| (Pw)(x)|x|^{\beta} \right\|_{L_{p}(a_{rn} \le |x|)} \le \exp(-Cn^{\alpha}) \left\| (Pw)(x)|x|^{\beta} \right\|_{L_{p}(L^{\frac{a_{n}}{n}} \le |x| \le a_{n}(1-L\eta_{n}))}.$$

Lemma 3.4. (1) Let $P \in \mathcal{P}_{\nu n-1}$. Then, we have

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} P(x) \right\|_{L_p(a_{2n} \le |x|)}$$

$$\le C e^{-C_1 n^{\eta}} \left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} P(x) \right\|_{L_p(|x| \le a_{2n})}.$$

So, if

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x)(|x|+1)^{\rho} \right\}^{\nu} P(x) \right\|_{L_{p}(\mathbb{R})} < \infty,$$

 $then \ we \ have$

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x) w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} P(x) \right\|_{L_p(a_{2n} \le |x|)} \to 0 \quad as \quad n \to \infty.$$

Proof. (1) Now, we denote the Mhaskar-Rakhmanov-Saff numbers for the exponents Q(x) and $\nu Q(x)$ by $a_n(Q)$ and $a_n(\nu Q)$ respectively. Then, we have $a_n(Q) = a_{\nu n}(\nu Q)$. From (2.7) we note that $(1 + |x|)^{\Delta} \Phi(x)^{-3\nu/4}$ is quasi-increasing. Then

applying Lemma 3.3 with $w^{\nu}_{\nu\rho}(x) = \exp(-\nu Q(x))|x|^{\nu\rho}$, we have

$$\begin{split} & \left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} P(x) \right\|_{L_p(a_{2n}(Q) \le |x|)} \\ & \le C \left\| w^{\nu}(x) \left(|x| + \frac{a_n}{n} \right)^{\nu\rho} P(x) \right\|_{L_p(a_{2\nu n}(\nu Q) \le |x|)} \\ & \le C e^{-C(\nu n)^{\eta}} \left\| w^{\nu}(x) \left(|x| + \frac{a_n}{n} \right)^{\nu\rho} P(x) \right\|_{L_p(|x| \le a_{2\nu n}(\nu Q))} \\ & \le C \frac{e^{-C\nu^{\eta}n^{\eta}}}{(1+a_{2n}(Q))^{-\Delta} \Phi(a_{2n}(Q))^{3\nu/4}} \\ & \times \left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)(w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} P(x) \right\|_{L_p(|x| \le a_{2n}(Q))} \\ & \le C e^{-C_1 n^{\eta}} \left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} P(x) \right\|_{L_p(|x| \le a_{2n}(Q))} \\ & \to 0 \quad \text{as} \quad n \to \infty, \end{split}$$

because that $(1 + a_{2n}(Q))^{\Delta} \Phi(a_{2n}(Q))^{-3\nu/4}$ has order n^s for some s > 0.

In the rest of this section we let $w_{\rho} \in \tilde{\mathcal{F}}_{\nu,\rho}(C^2+)$, $\rho \ge 0$, and we suppose Assumption 2.1. In addition, we let $\alpha > 0, \Delta > -1$, $1 , and <math>C_f$ be a positive constant depending only on f.

To simplify the proofs of theorems we use the results of [7]. In what follows we use the following notation. Let $x_{0,n} := x_{1,n} + \varphi_n(x_{1,n})$, and $x_{n+1,n} = -x_{0,n}$. When $|x| \leq x_{0,n}$, we define

(3.1)
$$x_{m,n} := x_{m(x),n}; \quad |x - x_{m,n}| = \min_{0 \le j \le n} |x - x_{j,n}|.$$

If $|x - x_{j+1,n}| = |x - x_{j,n}|$, then we set $x_{m,n} := x_{j,n}$. If $x_{0,n} < x$, then we put m = 0. And if $x < x_{n+1,n}$, then we put m = n + 1. Here, we note that there exists a constant $\delta > 0$ such that $|x - x_{m,n}| \le \delta \varphi_n(x_{m,n})$.

Lemma 3.5 (cf.[7]). Let m := m(x) be defined by (3.1). (1) If (2.10) holds, then, we have (3.2)

$$\left\{\Phi^{\frac{3}{4}}(x)w(x)\left(|x|+\frac{a_n}{n}\right)^{\rho}\right\}^{\nu}|X_n(\nu,f;x)| \le C_f\sum_{j=0}^n(1+|x_{j,n}|)^{-\alpha}\sum_{i=0}^{\nu-2}\left(\frac{1}{1+|m-j|}\right)^{\nu-i}$$

(2) If (2.12) holds, then, we have (3.3)

$$\left\{\Phi^{\frac{3}{4}}(x)w(x)\left(|x|+\frac{a_n}{n}\right)^{\rho}\right\}^{\nu}|Y_n(\nu,f;x)| \le C_f\varepsilon_n\sum_{j=0}^n(1+|x_{j,n}|)^{-\alpha}\left(\frac{1}{1+|m-j|}\right).$$

(3) If (2.13) holds, then, we have

(3.4)
$$\left\{ \Phi^{\frac{3}{4}}(x)w(x)\left(|x|+\frac{a_n}{n}\right)^{\rho} \right\}^{\nu} |Z_n(l,\nu,f;x)| \\ \leq C_f \frac{a_n}{n} \sum_{j=0}^n (1+|x_{j,n}|)^{-\alpha} \sum_{i=0}^{\nu-1} \left(\frac{1}{1+|m-j|}\right)^{\nu-i} \right\}$$

Proof. In [7, Propositions 3.7, 3.8 and 3.9] we got the similar estimations in $L_{\infty}(\mathbb{R})$ -space. Now, in there we can exchange f(x) with $f(x)(1+|x|)^{\alpha}$, and then we obtain the results in $L_p(\mathbb{R})$ by the similar methods as the proofs of [7, Propositions 3.7, 3.8 and 3.9] (see [7, p.16-p.23]).

Lemma 3.6. (1) [7, Lemma 4.3] Uniformly for $n \ge 1$,

(3.5)
$$\sup_{x \in \mathbb{R}} |p_{n,\rho}(x)w(x)| \left(|x| + \frac{a_n}{n}\right)^{\rho} \Phi^{\frac{1}{4}}(x) \le Ca_n^{-\frac{1}{2}}.$$

(2) [2, Theorem 2.5(a)] Uniformly for $n \ge 1$,

(3.6)
$$|p'_{n,\rho}w|(x_{j,n})\left(|x_{j,n}|+\frac{a_n}{n}\right)^{\rho} \sim \varphi_n(x_{j,n})^{-1}[a_n^2-x_{j,n}^2]^{-1/4}.$$

Lemma 3.7. Let $w(x) = \exp(-Q(x)) \in \mathcal{F}(C^2+)$. For the zeros $x_{j,n} = x_{j,n,\rho}$, we have the following.

(1) [2, Theorem 2.2, (b)] For $n \ge 1$ and $1 \le j \le n-1$,

$$x_{j,n} - x_{j+1,n} \sim \varphi_n(x_{j,n}),$$

and [2, Lemma A.1 (A.3)]

$$\varphi_n(x_{j,n}) \sim \varphi_n(x_{j+1,n}).$$

(2) [2, Theorem 2.2, (a)] For the minimum positive zero $x_{[n/2],n}$ ([n/2] is the largest integer $\leq n/2$), we have

$$x_{[n/2],n} \sim a_n n^{-1},$$

and for large enough n,

$$1 - \frac{x_{1,n}}{a_n} \sim \delta_n.$$

(3) [2, Lemma 4.7]

$$b_n = \frac{\gamma_{n-1}}{\gamma_n} \sim a_n \sim x_{1,n}$$

Lemma 3.8. Let $w = \exp(-Q) \in \mathcal{F}(C^2+)$. Let L > 0 be fixed. Then we have the following.

(a) ([16, Lemma 3.5, (a)]) Uniformly for t > 0,

 $a_{Lt} \sim a_t.$

(b) ([16, Lemma 3.5, (b)]) Uniformly for t > 0,

$$Q^{(j)}(a_{Lt}) \sim Q^{(j)}(a_t), \quad j = 0, 1.$$

Moreover,

$$T(a_{Lt}) \sim T(a_t).$$

(c) ([16, Lemma 3.11 (3.52)]) Uniformly for t > 0,

$$\left|1 - \frac{a_{Lt}}{a_t}\right| \sim \frac{1}{T(a_t)}$$

From Lemma 3.6 it is sufficient that we estimate Proposition 2.5, 2.6 or 2.7 for only $|x| \leq a_{2n}$.

Proof of Proposition 2.5. First we set

$$X_{n}(\nu, f; x) = \sum_{j=1}^{n} f(x_{j,n}) l_{jn}^{\nu}(x) \sum_{i=0}^{\nu-2} e_{i}(\nu, j, n) (x - x_{j,n})^{i}$$

=
$$: \sum_{j \ ; |x_{j,n}| \ge \frac{a_{n}}{3}} + \sum_{j \ ; |x_{j,n}| < \frac{a_{n}}{3}} =: X_{1,n}(\nu, f; x) + X_{2,n}(\nu, f; x).$$

Using (3.2), we see

$$\left| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x) w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} X_{1,n}(\nu, f; x) \right|$$

$$\leq C_f (1+|x|)^{-\Delta} \sum_{|x_{j,n}| \ge \frac{1}{3}a_n} (1+|x_{j,n}|)^{-\alpha} \sum_{i=0}^{\nu-2} \left(\frac{1}{1+|m-j|} \right)^{\nu-i}$$

$$\leq C C_f a_n^{-\alpha} (1+|x|)^{-\Delta}.$$

Therefore, we have by (2.9)

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} X_{1,n}(\nu, f; x) \right\|_{L_p(|x| \le a_{2n})}$$

$$\leq C_f a_n^{-\alpha} \left\| (1+|x|)^{-\Delta} \right\|_{L_p(|x| \le a_{2n})}$$

$$\leq C_f a_n^{-\alpha} \left\{ \begin{array}{l} a_n^{\frac{1}{p} - \Delta}, & \Delta p < 1, \\ \log a_n, & \Delta p = 1, \\ 1, & \Delta p > 1 \end{array} \right. \leq C_f.$$

Now, we will estimate for $X_{2,n}(\nu, f, x)$. For |x| < 1, since we have from (3.2)

$$\left\{\Phi^{\frac{3}{4}}(x)w(x)\left(|x|+\frac{a_n}{n}\right)^{\rho}\right\}^{\nu}|X_{2,n}(\nu,f;x)| \le C_f,$$

we see

(3.7)
$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} |X_{2,n}(\nu, f; x)| \right\|_{L_p(|x| \le 1)} \le C_f.$$

Let $|x| \ge 1$. We divide it into two sums as follows:

$$\begin{split} X_{2,n}(\nu,f;x) &=: \quad \sum_{\substack{j; \ |x_{j,n}| < \frac{a_n}{3}, \\ |x-x_{j,n}| \geqslant \frac{|x|}{2}}} + \sum_{\substack{j; \ |x_{jn}| < \frac{a_n}{3}, \\ |x-x_{j,n}| > \frac{|x|}{2}}} =: X_{2,n}^{[1]}(\nu,f;x) + X_{2,n}^{[2]}(\nu,f;x). \end{split}$$

Using (2.10), Lemma 3.7(2), (3.5) and (3.6) with $\varphi_n(x_{j,n}) \sim a_n/n$, we see that

(3.8)
$$\left\{ \Phi^{\frac{3}{4}}(x)w(x)\left(|x| + \frac{a_n}{n}\right)^{\rho} \right\}^{\nu} \left| f(x_{j,n})l_{jn}^{\nu}(x) \right|$$

$$\leq C_f (1 + |x_{j,n}|)^{-\alpha} \left| \frac{\left\{ \Phi^{\frac{3}{4}}(x_{j,n})\Phi^{\frac{3}{4}}(x)w(x)\left(|x| + \frac{a_n}{n}\right)^{\rho} \right\} p_n(x)}{p'_n(x_{j,n})w(x_{j,n})(\frac{a_n}{n} + |x_{j,n}|)^{\rho}} \right|^{\nu}$$

$$\leq CC_f (1 + |x_{j,n}|)^{-\alpha} \left(\frac{a_n}{n}\right)^{\nu}$$
and Lemme 2.1

and Lemma 3.1

$$\sum_{i=0}^{\nu-2} e_i(\nu, j, n) (x - x_{j,n})^i \le C \sum_{i=0}^{\nu-2} \left(\frac{n}{a_n}\right)^i |x|^{i-\nu} \le C \left(\frac{n}{a_n}\right)^{\nu-2} |x|^{-2}.$$

Then, we have

$$\left| \left\{ \Phi^{\frac{3}{4}}(x)w(x)\left(|x| + \frac{a_n}{n}\right)^{\rho} \right\}^{\nu} X_{2,n}^{[1]}(\nu, f; x) \right|$$

$$\leq C_f \sum_{\substack{j \ ; |x_{j,n}| < \frac{a_n}{3}, \\ |x - x_{j,n}| \geqslant \frac{|x|}{2}}} (1 + |x_{j,n}|)^{-\alpha} \left(\frac{a_n}{n}\right)^2 |x|^{-2} \leq CC_f \frac{a_n^2}{n} \frac{1}{x^2}$$

Therefore, we have, using $\Delta > -1$,

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} X_{2,n}^{[1]}(\nu, f; x) \right\|_{L_p(1 \le |x| \le a_{2n})}$$

$$(3.9) \leq C_f \frac{a_n^2}{n} \left\| (1+|x|)^{-(\Delta+2)} \right\|_{L_p(1 \le |x| \le a_{2n})} \le CC_f \frac{a_n^2}{n} \le CC_f$$

(note (2.2) and (2.5)). Since, for $|x - x_{j,n}| < \frac{|x|}{2}$, we see $|x_{j,n}| \sim |x|$, the inequality (3.2) implies

$$(3.10) \quad \left\{ \Phi^{\frac{3}{4}}(x)w(x)\left(|x| + \frac{a_n}{n}\right)^{\rho} \right\}^{\nu} X_{2,n}^{[2]}(\nu, f; x) \\ \leq CC_f \sum_{\substack{j; \ |x_{j,n}| < \frac{a_n}{3}, \\ |x - x_{j,n}| < \frac{|x|}{2}}} (1 + |x_{j,n}|)^{-\alpha} \sum_{i=0}^{\nu-2} \left(\frac{1}{1 + |m-j|}\right)^{\nu-i} \leq CC_f (1 + |x|)^{-\alpha}.$$

Therefore, we obtain by (2.9)

(3.11)
$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} X_{2,n}^{[2]}(\nu, f; x) \right\|_{L_p(1 \le |x| \le a_{2n})}$$

 $\le CC_f \left\| (1+|x|)^{-(\alpha+\Delta)} \right\|_{L_p(|x| \le a_{2n})} \le CC_f a_n^{\frac{1}{p}-(\alpha+\Delta)} \le CC_f.$

Hence, from (3.7), (3.8), (3.9), (3.11) and Lemma 3.4 we conclude (2.11) \Box *Proof of Proposition 2.6.* We repeat the methods of the proof of Proposition 2.5.

$$Y_n(\nu,f;x) =: \sum_{j; |x_{j,n}| \ge \frac{a_n}{3}} + \sum_{j; |x_{j,n}| < \frac{a_n}{3}} =: Y_{1,n}(\nu,f;x) + Y_{2,n}(\nu,f;x).$$

Then we have by (3.3), (2.9), (2.2) and (2.5)

$$\begin{aligned} \left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} Y_{1,n}(\nu, f; x) \right\|_{L_p(|x| \le a_{2n})} \\ &\leq C_f a_n^{-\alpha} \varepsilon_n \log n \left\| (1+|x|)^{-\Delta} \right\|_{L_p(|x| \le a_{2n})} \\ &\leq CC_f a_n^{-\alpha} \varepsilon_n \log n \begin{cases} a_n^{\frac{1}{p}-\Delta}, & \Delta p < 1; \\ \log a_n, & \Delta p = 1; \\ 1, & \Delta p > 1 \end{cases} \\ &\leq CC_f \varepsilon_n \log n \begin{cases} 1, & \Delta p < 1; \\ \frac{\log a_n}{a_n}, & \Delta p = 1; \\ \frac{1}{a_n}, & \Delta p > 1. \end{cases} \end{aligned}$$

Now, we estimate $Y_{2,n}(\nu, f; x)$. For $|x| \leq 1$ we obtain that from (3.3),

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} Y_{2,n}(\nu, f; x) \right\|_{L_p(|x| \le 1)} \le CC_f \varepsilon_n \log n.$$

Let $|x| \ge 1$ and

$$Y_{2,n}(\nu, f; x) =: \sum_{\substack{j \ ; |x_{j,n}| < \frac{a_n}{3}, \\ |x - x_{j,n}| \geqslant \frac{|x|}{2} > 0}} + \sum_{\substack{j \ ; |x_{j,n}| < \frac{a_n}{3}, \\ |x - x_{j,n}| < \frac{|x|}{2}}} =: Y_{2,n}^{[1]} + Y_{2,n}^{[2]}$$

Then we have similarly to (3.10) and (3.11),

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} Y_{2,n}^{[2]}(\nu, f; x) \right\|_{L_p(1 \le |x| \le a_{2n})}$$

$$\leq CC_f \varepsilon_n \log n \left\| (1+|x|)^{-(\alpha+\Delta)} \right\|_{L_p(|x| \le a_{2n})} \le CC_f a_n^{\frac{1}{p}-(\alpha+\Delta)} \varepsilon_n \log n \le CC_f \varepsilon_n \log n$$

(see (2.9)). Finally, we estimate $Y_{2,n}^{[1]}$. Let $|x_{j,n}| < \frac{a_n}{3}$ and $|x - x_{j,n}| \ge \frac{|x|}{2}$. Then, from Lemma 3.2, (2.12), (3.5) and (3.6), we have

$$(3.12) \qquad \left\{ \Phi^{\frac{3}{4}}(x)w(x)\left(|x| + \frac{a_n}{n}\right)^{\rho} \right\}^{\nu} \left| f(x_{j,n})l_{jn}^{\nu}(x) \right| |x - x_{j,n}|^{\nu-1}e_{\nu-1}(\nu, k, n) \\ \leq CC_f (1 + |x_{j,n}|)^{-\alpha} \left\{ |Q'(x_{j,n})| + \frac{1}{|x_{j,n}|} \right\}^{-1} \\ \times \left| \frac{\left\{ \Phi^{\frac{3}{4}}(x_{j,n})\Phi^{\frac{3}{4}}(x)w(x)\left(|x| + \frac{a_n}{n}\right)^{\rho} \right\} p_n(x)}{p'_n(x_{j,n})w(x_{j,n})(\frac{a_n}{n} + |x_{j,n}|)^{\rho}} \right|^{\nu} \frac{1}{|x - x_{j,n}|} \\ \times \left\{ \frac{T(a_n)}{a_n} + |Q'(x_{k,n})| + \frac{1}{x_{k,n}} \right\} \left\{ \frac{n}{a_{2n} - |x_{k,n}|} \right\}^{\nu-2} \\ \leq CC_f (1 + |x_{j,n}|)^{-\alpha} \left\{ |Q'(x_{j,n})| + \frac{1}{|x_{j,n}|} \right\}^{-1} \\ \times \Phi^{\frac{3\nu}{4}}(x_{j,n})\varphi_n^{\nu}(x_{j,n}) \left(1 - \frac{|x_{j,n}|}{a_n} \right)^{\frac{\nu}{4}} \frac{1}{|x - x_{j,n}|} \\ \times \left\{ \frac{T(a_n)}{a_n} + |Q'(x_{k,n})| + \frac{1}{x_{k,n}} \right\} \left\{ \frac{n}{a_{2n} - |x_{k,n}|} \right\}^{\nu-2}.$$

On the other hand, from (2.1), (2.8) and $|x_{j,n}| < a_n/3$, we easily see

(3.13)
$$\Phi_n^{\frac{3\nu}{4}}(x_{j,n})\varphi_n^{\nu}(x_{j,n})\left(1-\frac{|x_{j,n}|}{a_n}\right)^{\frac{\nu}{4}}\left(\frac{n}{a_{2n}-|x_{j,n}|}\right)^{\nu-2} \le C\left(\frac{a_n}{n}\right)^2.$$

Moreover, since $|Q'(x_{j,n})| + \frac{1}{|x_{j,n}|} \ge C > 0$ for some constant C, we see (3.14)

$$\left\{\frac{T(a_n)}{a_n} + |Q'(x_{j,n})| + \frac{1}{|x_{j,n}|}\right\} \left\{ |Q'(x_{j,n})| + \frac{1}{|x_{j,n}|} \right\}^{-1} \le C \max\left\{1, \frac{T(a_n)}{a_n}\right\}$$

and since $|x - x_{j,n}| \ge \frac{|x|}{2}$, we have

(3.15)
$$\frac{1}{|x - x_{j,n}|} \le \frac{2}{|x|}.$$

Thus, using (3.12), (3.13), (3.14) and (3.15), we can estimate for $Y_{2,n}^{[1]}(\nu, f; x)$ as follows:

$$\left\{ \Phi^{\frac{3}{4}}(x)w(x)\left(|x|+\frac{a_{n}}{n}\right)^{\rho} \right\}^{\nu}Y_{2,n}^{[1]}(\nu,f;x) \\
\leq CC_{f} \max\left\{ 1,\frac{T(a_{n})}{a_{n}} \right\} \frac{1}{|x|} \left(\frac{a_{n}}{n}\right)^{2} \sum_{\substack{j; \ |x_{j,n}| < \frac{a_{n}}{3}, \\ |x-x_{j,n}| \geqslant \frac{|x|}{2}}} (1+|x_{j,n}|)^{-\alpha} \\
\leq CC_{f} \max\left\{ 1,\frac{T(a_{n})}{a_{n}} \right\} \frac{1}{|x|} \frac{a_{n}^{2}}{n} \leq CC_{f} a_{n} \varepsilon_{n} \frac{1}{|x|}.$$

Therefore we have by (2.9)

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} Y_{2,n}^{[1]}(\nu, f; x) \right\|_{L_p(1 \le |x| \le a_{2n})}$$

$$\le CC_f a_n \varepsilon_n \left\| (1+|x|)^{-(\Delta+1)} \right\|_{L_p(1 \le |x| \le a_{2n})} \le CC_f a_n \varepsilon_n a_n^{\frac{1}{p}-(\Delta+1)} \le CC_f a_n \varepsilon_n.$$

Proof of Proposition 2.7. Using (3.4), we have from (2.2) and (2.5)

$$\begin{aligned} \left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} Z_n(l,\nu,f;x) \right\|_{L_p(\mathbb{R})} \\ &\leq CC_f \frac{a_n \log n}{n} \left\| (1+|x|)^{-\Delta} \right\|_{L_p(|x| \le a_{2n})} \\ &\leq CC_f \frac{a_n \log n}{n} \begin{cases} a_n^{\frac{1}{p} - \Delta}, & \Delta p < 1; \\ \log a_n, & \Delta p = 1; \le CC_f \frac{a_n^2 \log n}{n} \begin{cases} 1, & \Delta p < 1; \\ \frac{\log a_n}{a_n}, & \Delta p = 1; \\ 1, & \Delta p > 1 \end{cases} \\ \end{aligned}$$

Proof of Proposition 2.8. Since we see for a fixed polynomial $P \in \mathcal{P}_{\nu n-1}$

$$L_n(\nu, P) - P = L_n(\nu, P) - L_n(\nu - 1, \nu, P) = Z_n(\nu - 1, \nu, P),$$

we can obtain from Proposition 2.7,

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} \left\{ L_n(\nu, P) - P \right\} \right\|_{L_p(\mathbb{R})}$$

$$\leq CC_f \frac{a_n^2 \log n}{n} \begin{cases} 1, & \Delta p < 1; \\ \frac{\log a_n}{a_n}, & \Delta p = 1; \rightarrow 0 \quad \text{as} \quad n \rightarrow \infty \\ \frac{1}{a_n}, & \Delta p > 1. \end{cases}$$

(note Remark 2.13).

Proof of Proposition 2.9. Since we see for a fixed polynomial $P \in \mathcal{P}_{\nu n-1}$

$$X_n(\nu, P) - P = L_n(\nu, P) - P + Y_n(\nu, P)$$

and $\varepsilon_n(a_n + \log n) \to 0$ as $n \to 0$, we have the result from Proposition 2.6 and Proposition 2.8.

Proof of Theorem 2.11. Let $\varepsilon > 0$ be taken arbitrarily. Then there exists a polynomial P such that

(3.16)
$$\left\| (f(x) - P(x)) \left(1 + |x|\right)^{\alpha} \left\{ \Phi^{-\frac{3}{4}}(x) w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu - \frac{\eta}{2}} \right\|_{L_{\infty}(\mathbb{R})} < \varepsilon.$$

In fact, by (2.14) we see

$$\lim_{|x| \to \infty} \left| f(x) w^{\nu - \eta}(x) \right| = 0,$$

hence as [7, (4.55)] we obtain (3.16). Then we also have

(3.17)
$$\left\| (f(x) - P(x)) (1 + |x|)^{\alpha} \left\{ \Phi^{-\frac{3}{4}}(x) w_{\rho}(x) \right\}^{\nu} \right\|_{L_{\infty}(\mathbb{R})} < C\varepsilon,$$

and from

$$\left\| (1+|x|)^{\Delta-\alpha} \left\{ \Phi^{-\frac{3}{4}}(x) w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu/2} \right\|_{L_{\infty}(\mathbb{R})} < \infty,$$

with (3.16), we have

(3.18)
$$\left\| (f(x) - P(x)) (1 + |x|)^{\Delta} \left\{ \Phi^{-\frac{3}{4}}(x) w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} \right\|_{L_{\infty}(\mathbb{R})} < C\varepsilon.$$

Now, we see

$$|L_n(\nu, f) - f| \le |X_n(\nu, f - P)| + |X_n(\nu, P) - P| + |P - f| + |Y_n(\nu, f)|.$$

From Proposition 2.5 with $C_{f-P} = C\varepsilon$ and (3.17), we know

(3.19)
$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} X_n(\nu, f-P) \right\|_{L_p(\mathbb{R})} \le C\varepsilon.$$

Then, using (3.18), (3.19), Proposition 2.9 and Proposition 2.6, we conclude

$$\left\| (1+|x|)^{-\Delta} \left\{ \Phi^{\frac{3}{4}}(x)w(x) \left(|x| + \frac{a_n}{n} \right)^{\rho} \right\}^{\nu} (L_n(\nu, f) - f) \right\|_{L_p(\mathbb{R})} \le C\varepsilon.$$

Hence, we have the result.

Proof of Theorem 2.12. If we split $|L_n(l,\nu,f) - f|$ as follows:

$$|L_n(l,\nu,f) - f| \le |L_n(\nu,f) - f| + |Z_n(l,\nu,f)|,$$

then the result is proved from Theorem 2.11 and Proposition 2.7.

References

- [1] H. S. Jung and R. Sakai, Inequalities with exponential weights, JCAM 212(2008) 359-373.
- H. S. Jung and R. Sakai, Orthonormal polynomials with exponential-type weights, J. Approx. Theory 152(2008) 215-238.
- [3] H. S. Jung and R. Sakai, Markov-Bernstein inequality and Hermite-Fejér interpolation for exponential-type weights, J. Approx. Theory 162(2010), 1381-1397.
- [4] H. S. Jung and R. Sakai, Derivatives of integrating functions for orthonormal polynomials with exponential-type weights, J. Inequalities and Applications, Vol. 2009, Article ID 528454, 22 pages.
- [5] H. S. Jung and R. Sakai, Derivatives of orthonormal polynomials and coefficients of Hermite-Fejér interpolation polynomial with exponential-type weights, J. Inequalities and Applications, Vol. 2010, Article ID 816363, 29 pages.
- [6] H. S. Jung and R. Sakai, Specific examples of exponential weights, Commun. Korean Math. Soc. 24 (2009), No.2, 303-319.

- [7] H. S. Jung, G. Nakamura, R. Sakai and N. Suzuki, Convergence or divergence of higher order Hermite or Hermite-Fejér interpolation polynomials with exponential-type weights, ISRN Mathematical Analysis, Volume 2012, Article ID 904169, 31 pages, doi:10.5402/904169
- [8] T. Kasuga and R. Sakai, Uniform or Mean Convergence of Hermite-Fejér Interpolation of Higher Order for Freud Weights, J. Approx. Theory 101, 330-358(1999).
- [9] T. Kasuga and R. Sakai, Orthonormal polynomials with generalized Freud-type weights, J. Approx. Theory 121(2003), 13-53.
- [10] T. Kasuga and R. Sakai, Orthonormal Polynomials for Generalized Freud-type Weights and Higher Order Hermite-Fejér Interpolation Polynomials, J. Approx. Theory 127(2003), 1-38.
- [11] T. Kasuga and R. Sakai, Orthonormal polynomials for Laguerre-type weights, Far East J. Math. Sci. (FJMS), Vol. 15 No.1(2004), 95-105.
- [12] T. Kasuga and R. Sakai, Conditions for Uniform or Mean Convergence Hermite-Fejér Interpolation of Higher Order for Generalized Freud Weights, FJMS Vol.19(2005), 145-199.
- [13] Y. Kanjin and R. Sakai, Pointwise convergence of Hermite-Fejér interpolation of higher order for Freud weights, Tohoku Math. J. 46(1994), 181-206.
- [14] Y. Kanjin and R. Sakai, Convergence of the Derivatives of Hermite-Fejér Interpolation of Higher Order Based at the Zeros of Freud Polynomials Weights, J. Approx. Theory 80(1995), 378-389.
- [15] D. S. Lubinsky, Hermite-Fejér interpolation and associated product integration rules on the real line: The L_{∞} Theory, J. Approx. Theory 70(1992), 284-334.
- [16] A. L. Levin and D. S. Lubinsky, Orthogonal Polynomials for Exponential Weights, Springer, New York, 2001.
- [17] R. Sakai, Hermite-Fejér interpolation, Collo. Math. Soci. Janos Bolyai "Approx. Theory" (Hungary Kecskemet)(1990), 591-601.
- [18] R. Sakai, The degree of approximation of differentiable by Hermite interpolation polynomials, Acta Math. Acad. Sci. Hungar., 58(1991), 9-11.
- [19] R. Sakai, Hermite-Fejér interpolation prescribing higher order derivatives, Progress in Approximation Theory (edited by P. Nevai and A. Pinkus), Academic Press, Inc.(1991) 731-759.
- [20] R. Sakai, Certain bounded Hermite-Fejér interpolation polynomials operator, Acta Math. Acad. Sci. Hungar., 59(1992), 111-114.
- [21] R. Sakai and P. Vértesi, Hermite-Fejér interpolation of higher order, V, Studia Sci. Math. Hungarica 28(1993), 87-97.
- [22] R. Sakai and P. Vértesi, Hermite-Fejér interpolation of higher order, W, Studia Sci. Math. Hungarica 28(1993), 379-386.
- [23] R. Sakai and N. Suzuki, Farvard-type theorem and Jackson-type theorem, Pioneer J. Math. vol 3. no.1, 2011, 1-16.
- [24] R. Sakai and N. Suzuki, Mollification of exponential weights and its application to the Markov-Bernstein inequality, Pioneer J. Math. vol 7. no.1, 2013, 83-101.

 $^{1}\mathrm{Department}$ of Mathematics Education, Sungkyunkwan University, Seoul 110-745, Republic of Korea.

E-mail address: hsun90@skku.edu

²DEPARTMENT OF MATHEMATICS, MEIJO UNIVERSITY, NAGOYA 468-8502, JAPAN. *E-mail address*: ryozi@crest.ocn.ne.jp