Identifying Local and Quasilocal Conserved Quantities in Integrable Systems

Marcin Mierzejewski, ¹ Peter Prelovšek, ^{2,3} and Tomaž Prosen²

¹Institute of Physics, University of Silesia, 40-007 Katowice, Poland ²Faculty of Mathematics and Physics, University of Ljubljana, SI-1000 Ljubljana, Slovenia ³J. Stefan Institute, SI-1000 Ljubljana, Slovenia

We outline a procedure for counting and identifying a complete set of local and quasilocal conserved operators in integrable lattice systems. The method yields a systematic generation of all independent, conserved quasilocal operators related to time-average of local operators with a support on up to M consecutive sites. As an example we study the anisotropic Heisenberg spin-1/2 chain and show that the number of independent conserved operators grows linearly with M. Besides the known local operators there exist novel quasilocal conserved quantities in all the parity sectors. The existence of quasilocal conserved operators is shown also for the isotropic Heisenberg model. Implications for the anomalous relaxation of quenched systems are discussed as well.

PACS numbers: 72.10.-d,75.10.Pq, 05.60.Gg,05.70.Ln

Introduction. One expects that the long-time properties of generic systems are consistent with the Gibbs ensemble [1-4]. It means that the steady states are determined by very few conserved quantities (CQ), e.g., the total energy, the total spin and the particle number. However, in the integrable models there exist a macroscopic number of local CQ which can explain some anomalous transport properties [5]. Various studies have recently suggested that steady states of integrable systems [6–9] are fully specified by these local CQ. This conjecture is known as the generalized Gibbs ensemble [10–14] (GGE) and has been well established in systems which can be mapped on noninteracting particles [15]. It is quite evident that an application of this concept or its possible extension to other integrable systems relies on completeness of the set of CQ [16–18].

Let us consider the well known class of integrable models of strongly interacting particles on one-dimensional lattice with L sites. It includes anisotropic Heisenberg spin-1/2 model (XXZ model), the Hubbard model and the supersymmetric t-J model. In this Letter we focus on the first one while the procedure and arguments are generally applicable. For XXZ model there exist a straightforward procedure [19, 20] to construct a set of translationally invariant CQ, $Q_m = \sum_j q_j^m$, where q_j^m are local operators spanning at most m sites starting from j. Such local CQ commute with the Hamiltonian and with each other. On the other hand, it has been recognised that these CQ are not enough to account for the dissipationless spin transport [5, 21–31] neither for absence of decay of other correlations functions [17, 32] nor the kinetic energy [33]. A partial resolution of the dilemma was in finding novel quasilocal conserved quantities (QLCQ), constructed for XXZ model by one of the present authors [34–36] and others [37], which have direct overlap with the spin current operator. These particular QLCQ have recently been used to extend the validity of GGE to cases of current-carrying states [16]. Furthermore, results on quantum quenches [17, 32] suggested that new QLCQ

(unrelated to Q_m) should exist also in other symmetry sectors.

In this Letter we present a systematic procedure for generating local QC and QLCQ. It is based on the construction of time-averaged operators A emerging from a general local A spanning up to M sites. Such operators by construction commute with H but are in general highly nonlocal and do not have any meaningful thermodynamic limit $L \to \infty$. Hence we use an appropriate Hilbert-Schmidt inner product to select the relevant independent QLCQ. For arbitrary model parameters we find clear evidence for the existence of QLCQ in the even spinflip parity sector, while odd parity QLCQ are found only in the easy-plane regime. On the other hand, the total number of CQ and QLCQ appears to scale linearly with M being close to the number of CQ in a noninteracting system, 2(M-1), and far below the maximum number of mutually commuting (diagonal) operators $\propto 2^{M}$. Such a finding can have important consequences for relevance of extended GGE

 $\it The\ method.-$ For concreteness we focus on a paradigmatic example of integrable quantum systems, the XXZ model

$$H = J \sum_{j=1}^{L} \left[\frac{1}{2} (S_j^+ S_{j+1}^- + S_j^- S_{j+1}^+) + \Delta S_j^z S_{j+1}^z \right], \quad (1)$$

where $S_j^{\pm,z}$ are spin-1/2 operators and Δ is the anisotropy. To avoid degenerate states we use twisted boundary conditions $S_{j+L}^z \equiv S_j^z, S_{j+L}^\pm \equiv e^{\pm i\phi}S_j^\pm$ introducing flux $\phi \neq 0$ (see Appendix).

Within the space of all translationally invariant traceless observables, named as A_L , we introduce the following (Hilbert-Schmidt) inner product

$$(A|B) = \frac{1}{L} \frac{1}{2^L} \operatorname{tr} A^{\dagger} B, \qquad (2)$$

which is equivalent to the infinite temperature $(\beta \to 0)$ correlation $(A|B) = \langle A^{\dagger}B \rangle_{\beta=0}/L$. The normalisation is

chosen such that extensive local observables, like Q_m , have finite norm $\|A\|^2 = (A|A)$ in the thermodynamic limit $L \to \infty$. We consider finiteness of this norm in the thermodynamic limit as a general definition of either locality or quasilocality of an observable A (see also Ref. [38]). We define a subspace \mathcal{A}_L^m of local translationally invariant observables with support of size m. In particular, we can define the basis of \mathcal{A}_L^m (m-local basis) to be composed of operators

$$O_{\underline{s}} = \sum_{j} \sigma_{j}^{s_1} \sigma_{j+1}^{s_2} \cdots \sigma_{j+m-1}^{s_m}, \tag{3}$$

where $\sigma_j^z \equiv 2S_j^z, \sigma_j^{\pm} \equiv \sqrt{2}S_j^{\pm}, \sigma_j^0 \equiv 1, \underline{s} = (s_1, \dots, s_m), s_j \in \{+, -, z, 0\}$ while $s_{1,m} \in \{+, -, z\}$. Note that for given m there are $N_m = 3 \times 4^{m-2} \times 3$ different $O_{\underline{s}}$, and $\dim \mathcal{A}_L^m = N_m$ for $m \leq L/2$. Definitions (2,3) imply that operators are orthonormal, i.e. $(O_s|O_{s'}) = \delta_{s,s'}$.

Let us define the time-average of operator $A \in \mathcal{A}_L^m$

$$\bar{A} = \lim_{\tau \to \infty} \frac{1}{\tau} \int_0^{\tau} dt e^{iHt} A e^{-iHt} = \sum_{n,n'}^{E_n = E_{n'}} \langle n|A|n' \rangle |n\rangle \langle n'|,$$
(4)

which by construction gives a conserved operator $[H, \bar{A}] = 0$. In principle, \bar{A} can be reexpressed, using exact diagonalization $H|n\rangle = E_n|n\rangle$ via Eq. (4), in terms of operators (3). On this basis one could decide whether the operator \bar{A} is local ($\in \mathcal{B}_L^M \equiv \bigoplus_{l=1}^M \mathcal{A}_L^l$ for some L-independent $M \geq m$), or quasilocal (with convergent sum of operators $\in \mathcal{A}_L^l$ with increasing l), or generic nonlocal. Such a direct approach is, however, tedious and less transparent and in following we use a different protocol.

Picking M>0, we calculate the complete set of $\bar{O}_{\underline{s}}$ of $D_M=\sum_{m=1}^M N_m=3\times 4^{M-1}$ (where $N_1=3$) traceless operators spanning \mathcal{B}_L^M . To answer how many of $\bar{O}_{\underline{s}}$ are local or might be quasilocal, and are as well independent we evaluate Hermitian positive-definite $D_M\times D_M$ matrix

$$K_{\underline{s},\underline{s}'} = (\bar{O}_{\underline{s}}|\bar{O}_{\underline{s}'}) = (O_{\underline{s}}|\bar{O}_{\underline{s}'}) \tag{5}$$

which can be considered as the generalized stiffness matrix (at $\beta \to 0$) in analogy with the (spin) current J_s stiffness $D = \beta(\bar{J}_s|\bar{J}_s)$. Orthonormal eigenvectors $u_{l,\underline{s}}$ of matrix K corresponding to eigenvalues $\lambda_l > 0$ generate linearly independent conserved operators Q_l'

$$Q'_{l} = \sum_{\underline{s}} u_{l,\underline{s}} \bar{O}_{\underline{s}} = \sum_{\underline{s}} v_{l,\underline{s}} O_{\underline{s}} + Q_{l}^{\perp}, \tag{6}$$

where the operator Q_l^{\perp} has support on more than M sites, hence $(Q_l^{\perp}|O_s)=0$. Calculating the inner product of Q_l' with $O_{\underline{s}'}$ and utilizing Eq. (5) one finds that $v_{l,\underline{s}}=\lambda_l u_{l,\underline{s}}$. Substituting this result back into Eq. (6) and calculating the (squared) norm of operators on both hand sides one finds that

$$\lambda_l = \lambda_l^2 + \|Q_l^{\perp}\|^2. \tag{7}$$

Local CQ with support on up to M sites $(\|Q_l^\perp\|=0)$ has $\lambda_l=1$ strictly independent of L. Contrary to this, $\lambda_l|_{L\to\infty}>0$ corresponding to QLCQ $(\|Q_l^\perp\|>0)$ are always smaller than 1 gradually approaching unity with growing M. The objective of our study is to establish how the number of CQ and QLCQ depends on M.

Symmetries. The matrix K can be decomposed in terms of the symmetries of the system. Within the XXZ model (1), one CQ is the magnetisation $S_{tot}^z = \sum_i S_i^z$ implying that one may consider S_{tot}^z preserving subspace $[S^z_{tot},O_{\underline{s}}]=0,$ i.e., subset of $O_{\underline{s}}$ with the constraint that the number of s = + equals the number of s = - in the sequence \underline{s} , reducing the dimensions D_M . More interesting is the spin-flip \mathbb{Z}^2 symmetry, generated by the parity operator $P = \Pi_j(S_j^+ + S_j^-)$. m-local operator spaces \mathcal{A}_L^m can then be decomposed into even(E)/odd(O), with the bases generated by sets $\{O_{\underline{s}} \pm PO_{\underline{s}}P\}$. Remarkably, all known local conserved operators Q_m are even, while it has been shown recently [34–36] that odd QLCQ exist for $|\Delta| < 1$ which determine the properties of the spin current. Furthermore, XXZ model is time-reversal invariant implying that the time-averaging matrix K is real symmetric and its eigenoperators $\sum_{s} v_{l,\underline{s}} O_{\underline{s}}$ can be classified as real (R) or imaginary (I), being spanned separately by the bases $O_{\underline{s}} + O_{\underline{s}}^{\dagger}$, $i(O_{\underline{s}} - O_{\underline{s}}^{\dagger})$. Hence one ends up with four orthogonal sectors: RE, RO, IE, IO.

Numerical procedure and results. In order to reduce the computational effort we restrict our calculations to the Hilbert subspace with $S_{tot}^z=0$. This requires a straightforward modification of the scalar product in Eq. (2) where the number of states 2^L should be replaced by $\binom{L}{L/2}$. Although the set $\{O_{\underline{s}}\}$ is not orthonormal within the chosen subspace, the overlap matrix $N_{\underline{s},\underline{s}'} = (O_{\underline{s}}|O_{\underline{s}'})$ remains real, symmetric and positive-definite. Since it can be diagonalized by an orthogonal matrix V one can introduce an orthonormal set of operators O_{μ} = $\theta_{\mu}^{-1/2} \sum_{s} V_{\underline{s},\mu} O_{\underline{s}}$, where θ_{μ} are eigenvalues of N. The numerical calculations have been performed via full ED of systems with sizes L = 10 - 20 and the boundary condition twisted by $\phi = 10^{-4}$. For given M and L we start the procedure generating first all operators O_s then the orthonormal ones O_{μ} and finally the time–averaged \bar{O}_{μ} . At the end the matrix $K_{\mu,\nu} = (\bar{O}_{\mu}|\bar{O}_{\nu})$ is evaluated and diagonalized, leading to the eigenvalues $\lambda_l \in [0, 1]$. The number of non-vanishing eigenvalues is the quantity that we are looking for: the number of local or quasilocal mutually orthogonal conserved operators.

Let us first consider a more generic case as a test of the method. Adding to the XXZ hamiltonian (1) a next nearest neighbor interaction term $H'=\Delta_2\sum_j S_{j+2}^z S_j^z$ the model becomes nonintegrable. In this case we expect the existence of a single CQ (besides S_{tot}^z) which is the full Hamiltonian, i.e. $Q_2=H+H'$. In Fig. 1 we show the results for eigenvalues λ_l vs. $1/L, L=10,\ldots,18$ for both E/O sectors, choosing parameters $\Delta=0.5, \Delta_2=0.5$.

This implies the expectation that $\lambda_1 = 1$ being independent of L for $M \geq 3$, as confirmed in Fig. 1. All other λ_l in both sectors vanish with increasing L, predominantly exponentially, $\lambda_l \propto \exp(-\zeta L)$. There are exceptions decaying as $\lambda \propto 1/L$ which might be related to powers of local operators, e.g., $(Q_2)^2$ which are however nonlocal quantities [39].

On the other hand, for the integrable XXZ model one finds several CQ and QLCQ as shown in the three topmost rows of Fig.1 as well as in Fig.2. The latter figure demonstrates in more detail how the spectra of the ma- $\operatorname{trix} K$ depend on M and L. Results in Fig.1 show that the strictly local CO exist only in the even sector. These are exactly the well known CQ described in Refs. [19, 20]. All the other (novel) CQ are quasilocal. In the easy-axis, $\Delta > 1$, or isotropic, $\Delta = 1$, cases QLCQ exist only in the even parity sector. We have confirmed the latter observation carrying out a finite-size scaling of trK for the anisotropy $\Delta \geq 1$ (not shown). We have found a very clear vanishing of tr K for $L \to \infty$, in the odd sector at any fixed M, what excludes existence of QLCQ in these sectors. On the contrary, in the easy-plane regime, $|\Delta| < 1$, QLCQ exist in all parity and time-reversal sec-

While the initial operators $O_{\underline{s}}$ as well as the orthonormal ones O_{μ} have support on not more than M sites, the subsequent time-averaging may extend their support beyond this value. Therefore, before counting the number of QLCQ one should exclude the possibility that O_{μ} can be expanded solely in terms of higher local CQ having support on more than M sites. In order to estimate this contribution we have calculated the first nontrivial eigenvalue of $\tilde{K}_{\mu,\nu} = \sum_{m=2}^{\infty} (O_{\mu}|Q_m)(Q_m|O_{\nu})/(Q_m|Q_m)$ (for computation we truncate at m = 18), which would agree with $K_{\mu,\nu}$ under the assumption that the orthogonal local operators Q_m are a complete set, i.e. that A = $A \equiv \sum_{m} (Q_m|A)Q_m$. These leading eigenvalues of $\tilde{K}_{\mu,\nu}$, which are shown in Fig. 2 as open circles, are always well below eigenvalues of unprojected $K_{\mu,\nu}$ clearly indicating that the set of known Q_m is incomplete also in the even sector. The central question is how the number of CQ and QLCQ grows with the size of the support M and, in particular, whether this growth is linear as in the case of noninteracting particles, where one finds 2M-2 local operators, $Q_{m+1}^R=\sum_k 2\cos(mk)n_k=\sum_j (c_{j+m}^\dagger c_j+1)$ H.c.), $Q_{m+1}^{I} = \sum_{k} 2\sin(mk)n_{k} = i\sum_{j}(c_{j+m}^{\dagger}c_{j} - \text{H.c.}),$ $m = 1, \ldots, M-1$, rewritable into the spin language via $c_{j+m}^{\dagger}c_l=(1/2)\sigma_{j+m}^+\sigma_{j+m-1}^z\cdots\sigma_{j+1}^z\sigma_j^-.$

Since the eigenvalues $\lambda_l \leq 1$, the total number of CQ and QLCQ is obviously bounded from below by $\operatorname{tr} K$. Moreover, as all CQ with support on up to M sites correspond to $\lambda_l = 1$, $\operatorname{tr} K$ should gradually approach the number of QC and QLCQ for large enough M. In Fig. 3 we show $\operatorname{tr} K$ obtained after the 1/L size scaling. For comparison we show also the number of nonzero λ_l

counted directly after carrying out a finite-size scaling of individual leading eigenvalues. The number of directly counted CQ and QLCQ as well as trK increase linearly with M. We find it particularly surprising and suggestive that the in easy-plane regime trK is very close to 2(M-1), i.e. to the number of CQ in the systems of noninteracting particles. However, contrary to the latter systems trK in XXZ model is not equally distributed among the parity sectors. Half of the total trK comes from the known (even) QC, approximately one quarter originates from even QLQC and one quarter from odd QLCQ. In the isotropic and Ising regimes ($\Delta \geq 1$) QLQC in the odd sector disappear, while the total number of CQ and QLCQ in the even sector is evidently larger than (M-1)in noninteracting case. Our central observation concerns the main difference between interacting and noninteracting integrable systems. We have found that this difference does not consists in the number of QC (which is extensive in both cases) but rather in their locality and symmetry.

Identifying the conserved quantities.— Besides known $Q_m, m=2,3\ldots$ which are all in the even sector we did not find any other strictly local CQ. As expected we also confirmed the existence of the QLCQ in the odd sector which has been analytically constructed [34–37] and used in the extended GGE previously [16]. Furthermore, very weak double scaling of eigenvectors of the matrix K with, first, increasing L, and then, increasing M, allows us to clearly identify novel QLCQ. In the isotropic case $\Delta=1$ and ER symmetry sector, the first nontrivial QLCQ Q' reads approximately, writing all terms to order m=4, $Q' \propto \sum_j (\mathbf{S}_{j+2} \cdot \mathbf{S}_j + \mathbf{S}_{j+3} \cdot \mathbf{S}_j + (\mathbf{S}_{j+3} \cdot \mathbf{S}_{j+2})(\mathbf{S}_{j+1} \cdot \mathbf{S}_j) - 2(\mathbf{S}_{j+3} \cdot \mathbf{S}_{j+1})(\mathbf{S}_{j+2} \cdot \mathbf{S}_j) + \dots$ See Appendix for details. Steady state after a linear quench.—Let us assume that at time t=0 the Hamiltonian is quenched from $H-X_a$ to the integrable H, where $X_a \in \mathcal{A}_L$ is considered as a perturbation. The system is initially in the Gibbs state

at time t=0 the Hamiltonian is quenched from $H-X_a$ to the integrable H, where $X_a \in \mathcal{A}_L$ is considered as a perturbation. The system is initially in the Gibbs state $\rho_a = \exp[-\beta(H-X_a)]/Z$, where $Z = \operatorname{Tr} \exp[-\beta(H-X_a)]$, and it relaxes towards a steady state $\overline{\rho}_a$. Since \overline{X}_a commutes with H one easily finds the following linear expansion

$$\overline{\exp[-\beta(H - X_a)]} \simeq \exp(-\beta H)(1 + \beta \overline{X_a}). \tag{8}$$

Calculating the trace over the eigenstates of H one obtains an analogous approximation for the partition function $Z = Z_T(1 + \beta \langle X_a \rangle)$. Here $Z_T = \text{Tr}[\exp(-\beta H)]$ and the average $\langle ... \rangle$ is defined for the thermal state $\rho_T = \exp(-\beta H)/Z_T$. At the end we obtain the linear approximation for the steady state:

$$\overline{\rho}_a = \rho_T [1 + \beta (\overline{X}_a - \langle X_a \rangle)] \tag{9}$$

Next, we consider some other extensive observable $X_b \in \mathcal{A}_L$ and a related intensive one X_b/L . We study whether the (possibly) nonthermal $\overline{\rho}_a$ and the thermal ρ_T states can be distinguished by the measurement of X_b/L . We




FIG. 1. (Color online) The size (1/L) scaling of leading eigenvalues λ_l of matrix K with M=6, corresponding to symmetric (red circles) or antisymmetric (blue crosses) eigenoperators with respect to time reversal. Left/right column shows even/odd parity sectors, while rows indicate different regimes of integrable (upper three rows) and non-integrable (lower row) with parameters indicated in the panel. Dashed lines indicate 1/L extrapolation to TL which in some cases provide clear indication of existence of QLCQ $\lambda_l|_{L\to\infty}>0$, beyond the local eigenoperators with $\lambda_l=1$.

calculate

$$\beta K_{ab} = \text{Tr}\left[(\overline{\rho}_a - \rho_T) \frac{X_b}{L} \right] = \frac{\beta}{L} (\langle \overline{X}_a \overline{X}_b \rangle - \langle \overline{X}_a \rangle \langle \overline{X}_b \rangle).$$
(10)

In the high–temperature regime $(\beta \to 0)$ and for traceless operators X the above correlation matrix K coincides with the matrix of inner products in Eq. (5).

Conclusions.— We have presented a systematic procedure which allows to establish the existence of local and quasi local CQ in 1D many-body models with short range interactions. In spite of limitations of our results to finite sizes $L \leq 20$, explicitly performed only within the (unpolarised) subspace $S^z_{tot} = 0$, they allow for some firm conclusions for XXZ model. The method confirms besides the known strictly local CQ, being all in the even sector, also even and odd QLCQ. In the metallic regime $\Delta < 1$ the odd QLCQ are consistent with the finite




FIG. 2. (Color online) Dependence of leading eigenvalues λ_l of K in RE sector for isotropic HM $\Delta=1$. Different panels indicate decreasing support sizes M=6,5,4,3, while decreasing sizes of points and colors indicate the system size L=20 (orange), 18 (red), 16 (green), 14 (cyan), 12 (blue), 10 (brown). Extrapolated $L\to\infty$ values are indicated with crosses if in the range of the plot. Open circles are explained in the text.





FIG. 3. (Color online) Left: The extrapolated $1/L \to 0$ value of ${\rm tr} K$ versus support size M for different regimes of XXZ model: $\Delta=1/2$ (blue squares), where contributions for even/odd parity sector are shown separately with crosses/disks, $\Delta=1$ (red circles), $\Delta=3/2$ (diamonds). Right: similar plot for the number of non-vanishing extrapolated eigenvalues λ_l . Long/short dashed lines indicate number of known local CQ for interacting (M-1)/non-interacting (2M-2) cases.

spin stiffness D>0 and analytical construction [34–37]. The novel QLCQ in the even sector exist in the whole $\Delta>0$ range and can explain at least part of deviations from GGE observed so far in quenched spin systems at $\Delta>1$ [17, 32]. There are clear indications for the existence of further QLCQ which emerge with increasing support size M but for $M\leq 6$ are not yet sufficiently converged to determine their explicit form. It is nevertheless plausible that in quenched or driven systems the major role in thermalisation will be related to CQ and QLCQ with smaller supports M as identified in our study. Here the most important conclusion is that the number of local and quasilocal CQ appears to scale linearly with M, for $|\Delta|<1$, being approximately 2(M-1) similar to a






FIG. 4. Projections of Q' ($\Delta=1$, RE sector) on the basis vectors O_s as defined in Eq. (6) in the main text. The explicit form of O_s for s=1,...,16 is shown in the text. The shaded/unshaded backgrounds from the left to right side of figure mark the sets of O_s with support on 2, 3, 4 and 5 sites, respectively. Results for $s\leq 16$ are magnified in the lower panel.

model of noninteracting fermions. Our results allow for a meaningful extension of GGE incorporating the full set of QLCQ.

M.M. acknowledges support from the DEC-2013/09/B/ST3/01659 project of the Polish National Science Center. P.P. and T.P. acknowledge support by the program P1-0044 and projects J1-4244 (P.P.) and J1-5349, N1-0025 (T.P.) of the Slovenian Research Agency.

APPENDIX

In the Appendix we give some technical details of our study of the leading local and quasilocal conserved quantities (QLCQ) of the isotropic Heisenberg model. We also demonstrate that while the twisted boundary conditions facilitate the numerical calculations by lifting degeneracy of eigenstates, they have negligible influence on the numerical results, provided that the related flux ϕ is small enough.

Leading QLCQ.— Here we study the leading QLCQ in the real even—parity (RE) sector of the isotropic Heisenberg model $(\Delta=1)$ and we denote this QLCQ by Q'. The norm of Q' is shown in Fig. 2 in the main text as the 4th eigenvalue for M=6 as well as the 3rd eigenvalue for M=5 or M=4. In Fig. 4 we show projections, v_{ls} , of Q' on the basis operators O_s as defined in Eq. (6) in the main text. For the sake of completeness we list also explicit form of all O_s with support on up to 4 sites:

$$O_1 = \sqrt{2} \sum_{i} (S_i^- S_{i+1}^+ + H.c.)$$
 (11)

$$O_2 = 4\sum_i S_i^z S_{i+1}^z \tag{12}$$

$$O_3 = -\sqrt{2} \sum_{i} (S_i^- S_{i+2}^+ + H.c.)$$
 (13)

$$O_4 = 4\sum_{i} S_i^z S_{i+2}^z \tag{14}$$

$$O_5 = 4\sqrt{2} \sum_{i} (S_i^- S_{i+1}^z S_{i+2}^z S_{i+3}^+ + H.c.)$$
 (15)

$$O_6 = -4\sqrt{2} \sum_{i} (S_i^- S_{i+1}^z S_{i+2}^+ S_{i+3}^z + H.c.)$$
 (16)

$$O_7 = 2\sqrt{2} \sum_{i} (S_i^- S_{i+2}^+ S_{i+1}^- S_{i+3}^+ + H.c.)$$
 (17)

$$O_8 = \sqrt{2} \sum_{i} S_i^- S_{i+3}^+ + H.c.$$
 (18)

$$O_9 = -2\sqrt{2} \sum_{i} (S_i^- S_{i+1}^+ S_{i+2}^- S_{i+3}^+ + H.c.)$$
 (19)

$$O_{10} = 4\sqrt{2} \sum_{i} S_{i}^{-} S_{i+1}^{+} S_{i+2}^{z} S_{i+3}^{z}$$
 (20)

$$O_{11} = 4\sum_{i} S_i^z S_{i+3}^z + H.c. (21)$$

$$O_{12} = -4\sqrt{2} \sum_{i} (S_i^z S_{i+1}^- S_{i+2}^z S_{i+3}^+ + H.c.)$$
 (22)

$$O_{13} = 4\sqrt{2} \sum_{i} (S_i^z S_{i+1}^- S_{i+2}^+ S_{i+3}^z + H.c.)$$
 (23)

$$O_{14} = 4\sqrt{2} \sum_{i} (S_i^z S_{i+1}^z S_{i+2}^- S_{i+3}^+ + H.c.)$$
 (24)

$$O_{15} = 16 \sum_{i} S_{i}^{z} S_{i+1}^{z} S_{i+2}^{z} S_{i+3}^{z}$$
 (25)

$$O_{16} = 2\sqrt{2} \sum_{i} (S_{i}^{+} S_{i+1}^{-} S_{i+2}^{-} S_{i+3}^{+} + H.c)$$
 (26)

In order to establish Q' in the thermodynamic limit one should carry out two subsequent finite—size scalings of v_{ls} : first $L \to \infty$ and then $M \to \infty$. Results shown in the upper panel in Fig. 4 indicate that the range of

accessible L and M is insufficient to carry out such scaling for all s. However, the projections of Q' on the basis operators with support on up to 4 sites (s=1,...,16) only weakly depend on M and L, as shown in the lower panel in the same figure. We stress that the most important (mostly studied) physical quantities have support on a few lattice sites only. In order to study the influence of Q' on the dynamics of these quantities in the high-temperature regime, it is sufficient to specify v_{ls} only for these Q_s which have the same support as the investigated quantities. Hence, the projections shown in the lower panel in Fig. 4 are of key importance.

In order to get a deeper insight into the structure of QLCL we are focusing on the first nontrivial Q' in the case of SU(2) symmetric (isotropic) XXX model $(\Delta = 1)$. Using a simple group-theoretic argument one realises that the operators Q' should also be SU(2) invariants. In fact, they should be SU(2) scalars, unless they correspond do degenerate eigenvalues of the matrix K – which we have checked that it never takes place for finite M. As the only translationally invariant SU(2) scalar is the Hamiltonian itself $H = J \sum_{j} \mathbf{S_{j+1}} \cdot \mathbf{S_{j}}$, this immediately implies that the projections of Q' on O_1 and O_2 should be vanishingly small, in the limit $L \to \infty$, as clearly demonstrated in Fig. 4. We note that SU(2) invariance is expected to be restored only in the thermodynamic limit as we work with the microcanonical ensemble of zero total spin projection $\sum_j S_j^z$, which is (strictly) not SU(2) invariant for finite L. In other words, nonvanishing projections on O_1 or O_2 would imply that ether Q' is not SU(2) invariant or it is not orthogonal to the Hamiltonian $\sum_{i} \mathbf{S}_{i+1} \mathbf{S}_{i}$. The general form of SU(2)-invariant Q' with all terms up to order m=4

$$Q' = \sum_{i} [\alpha \mathbf{S}_{i+2} \cdot \mathbf{S}_{i} + \beta (\mathbf{S}_{i+3} \cdot \mathbf{S}_{i+2}) (\mathbf{S}_{i+1} \cdot \mathbf{S}_{i})$$

$$+ \gamma (\mathbf{S}_{i+3} \cdot \mathbf{S}_{i+1}) (\mathbf{S}_{i+2} \cdot \mathbf{S}_{i}) +$$

$$+ \delta (\mathbf{S}_{i+3} \cdot \mathbf{S}_{i}) (\mathbf{S}_{i+2} \cdot \mathbf{S}_{i+1}) + \zeta (\mathbf{S}_{i+3} \cdot \mathbf{S}_{i})] + ...,$$

$$(27)$$

can be expressed in terms of basis operators O_s

$$Q' = \frac{1}{4} (\alpha O_4 - \sqrt{2}\alpha O_3 + \zeta O_{11} + \sqrt{2}\zeta O_8)$$

$$+ \frac{\sqrt{2}}{16} [(\delta + \gamma)O_7 - (\beta + \delta)O_9 + (\beta + \gamma)O_{16}]$$

$$+ \frac{\sqrt{2}}{16} [\beta(O_{10} + O_{14}) - \gamma(O_6 + O_{12})$$

$$+ \delta(O_5 + O_{13})] + \frac{1}{16} (\beta + \delta + \gamma)O_{15}. \tag{28}$$

Numerical results for $v_{l3}, ..., v_{l16}$ shown in Fig. 4 show that Eq. (28) indeed holds true confirming SU(2) invariant form of Q' for isotropic Heisenberg model. We have also found the following approximate values of parame-




FIG. 5. The size (1/L) scaling of leading eigenvalues λ_l of matrix K with M=5 obtained for RE sector of anisotropic Heisenberg. Fig. 1 in the main text shows analogous results for M=6. Crosses and squares show results obtained for twisted and period boundary conditions, respectively.

ters in Eq. (27):

$$\alpha \simeq \beta \simeq \zeta,$$
 (29)

$$\gamma \simeq -2\alpha,\tag{30}$$

$$|\delta| \ll |\alpha|. \tag{31}$$

Twisted boundary conditions.— In order to avoid degeneracy of energy levels we used in the computations twisted boundary conditions $S_{j+L}^z \equiv S_j^z, S_{j+L}^\pm \equiv e^{\pm i\phi}S_j^\pm$ introducing flux $\phi = 10^{-4}$. As immediately follows from Eq. (4) in the main text, $\phi \neq 0$ facilitates the numerical calculations focused on the time-averaged operators. In Fig. (5) we compared results obtained for $\phi = 0$ and $\phi = 10^{-4}$ which confirm that such small values of ϕ have negligible influence on the eigenvalues of the K matrix.

- S. Goldstein, J. L. Lebowitz, R. Tumulka, and N. Zanghì, Phys. Rev. Lett. 96, 050403 (2006).
- [2] N. Linden, S. Popescu, A. J. Short, and A. Winter, Phys. Rev. E 79, 061103 (2009).
- [3] A. Riera, C. Gogolin, and J. Eisert, Phys. Rev. Lett. 108, 080402 (2012).
- [4] A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, Rev. Mod. Phys. 83, 863 (2011).
- [5] X. Zotos, F. Naef, and P. Prelovsek, Phys. Rev. B 55, 11029 (1997).
- [6] S. Hershfield, Phys. Rev. Lett. 70, 2134 (1993).
- [7] B. Doyon and N. Andrei, Phys. Rev. B 73, 245326 (2006).
- [8] L. F. Santos, A. Polkovnikov, and M. Rigol, Phys. Rev. Lett. 107, 040601 (2011).
- [9] M. Mierzejewski, T. Prosen, D. Crivelli, and P. Prelovšek, Phys. Rev. Lett. 110, 200602 (2013).
- [10] M. Rigol, V. Dunjko, V. Yurovsky, and M. Olshanii, Phys. Rev. Lett. 98, 050405 (2007).

- [11] M. Kollar, F. A. Wolf, and M. Eckstein, Phys. Rev. B 84, 054304 (2011).
- [12] A. C. Cassidy, C. W. Clark, and M. Rigol, Phys. Rev. Lett. 106, 140405 (2011).
- [13] C. Gogolin, M. P. Müller, and J. Eisert, Phys. Rev. Lett. 106, 040401 (2011).
- [14] M. Fagotti, M. Collura, F. H. L. Essler, and P. Calabrese, Phys. Rev. B 89, 125101 (2014).
- [15] M. Rigol, Phys. Rev. E **90**, 031301 (2014).
- [16] M. Mierzejewski, P. Prelovšek, and T. Prosen, Phys. Rev. Lett. 113, 020602 (2014).
- [17] B. Pozsgay, Μ. Mestyán, M. Α. Werner, Kormos, G. Zaránd, G. Takács, and Phys. Rev. Lett. 113, 117203 (2014).
- Goldstein N. Andrei, and Phys. Rev. A **90**, 043625 (2014).
- [19] M. Tetelman, Sov. Phys. JETP **55**, 306 (1982).
- [20] M. P. Grabowski and P. Mathieu, Ann. Phys. (N.Y.) 243, 299 (1995).
- [21] X. Zotos and P. Prelovšek, Phys. Rev. B 53, 983 (1996).
- [22] X. Zotos, Phys. Rev. Lett. 82, 1764 (1999).
- [23] J. Benz, T. Fukui, A. Klümper, and C. Scheeren, Journal of the Physical Society of Japan 74, 181 (2005), http://journals.jps.jp/doi/pdf/10.1143/JPSJS.74S.181.
- [24] F. Heidrich-Meisner, A. Honecker, and W. Brenig, The European Physical Journal Special Topics 151, 135 (2007). arXiv:0812.3096v1 (2008).

- [25] M. Rigol В. S. Shastry, and Phys. Rev. B 77, 161101 (2008).
- [26] J. Herbrych, P. Prelovšek, and Χ. Zotos, Phys. Rev. B 84, 155125 (2011).
- M. Žnidarič, Phys. Rev. Lett. 106, 220601 (2011).
- [28] J. Sirker, R. G. Pereira, and I. Affleck, Phys. Rev. Lett. 103, 216602 (2009).
- [29] R. Steinigeweg, J. Gemmer, and W. Brenig, Phys. Rev. Lett. 112, 120601 (2014).
- T. Prosen, Phys. Rev. Lett. 107, 137201 (2011).
- R. Steinigeweg, J. Herbrych, P. Prelovšek, and M. Mierzejewski, Phys. Rev. B 85, 214409 (2012).
- [32] B. J. De Nardis, M. Brockmann, Wouters, Fioretto, M. Rigol, and J.-S. Phys. Rev. Lett. 113, 117202 (2014).
- and P. Prelovšek, [33] R. Steinigeweg, J. Herbrych, Phys. Rev. E 87, 012118 (2013).
- [34] T. Prosen, Phys. Rev. Lett. **106**, 217206 (2011).
- [35] T. Prosen and Ilievski, Phys. Rev. Lett. 111, 057203 (2013).
- T. Prosen, Nuclear Physics B 886, 1177 (2014).
- [37] R. G. Pereira, V. Pasquier, J. Sirker, and I. Affleck, Journal of Statistical Mechanics: Theory and Experiment 2014, P09
- M. Olshanii, Phys. Rev. Lett. 114, 060401 (2015).
- [39] M. Hawkins, Μ. Long, and X. Zotos.