

# On solutions of the Pauli equation in non-static de Sitter metrics

E. M. Ovsiyuk, K. V. Kazmerchuk  
 Mozyr State Pedagogical University named after I.P. Shamyakin  
 e.ovsiyuk@mail.ru, kristinash2@mail.ru

October 23, 2018

## Abstract

A particle with spin 1/2 is investigated both in expanding and oscillating cosmological de Sitter models. It is shown that these space-time geometries admit existence of the non-relativistic limit in the covariant Dirac equation. Procedure for transition to the Pauli approximation is conducted in the equations in the variables  $(t, r)$ , obtained after separating the angular dependence of  $(\theta, \phi)$  from the wave function. The non-relativistic systems of equations in the variables  $(t, r)$  is solved exactly in both models. The constructed wave functions do not represent stationary states with fixed energy, however the corresponding probability density does not depend on the time.

## 1 Introduction

De Sitter and anti de Sitter models take attraction during long time in the context of development of the quantum theory in curved space-time. In particular, long history has the task of exact solving the wave equations for fields with different spins [1]–[12]; the main attention was focused of relativistic wave equations.

In the present paper we will examine the nonrelativistic Pauli approximation for spin 1/2 particle in non-static metrics of de Sitter models: we demonstrate how transition to Pauli approximation can be performed in non-static de Sitter's metrics, and we will construct exact solutions for these non-relativistic systems in both de Sitter models: expanding and oscillating ones.

## 2 Pauli equation in expanding de Sitter Universe

Generally covariant Dirac equation in orthogonal coordinates [13]

$$[i\gamma^k (e_{(k)}^\alpha \partial_\alpha + B_k) - m] \Psi = 0,$$

$$B_k(x) = \frac{1}{2} e_{(k);\alpha}^\alpha(x) = \frac{1}{2} \frac{1}{\sqrt{-g}} \frac{\partial}{\partial x^\alpha} \sqrt{-g} e_{(k)}^\alpha$$

being specified for the non-static de Sitter metric

$$x^\alpha = (t, r, \theta, \phi), \quad dS^2 = dt^2 - \cosh^2 t [dr^2 + \sin^2 r (d\theta^2 + \sin^2 \theta d\phi^2)]$$

and the corresponding tetrad

$$\begin{aligned} e_{(0)}^\alpha &= (1, 0, 0, 0), & e_{(3)}^\alpha &= \left(0, \frac{1}{\cosh t}, 0, 0\right), \\ e_{(2)}^\alpha &= \left(0, 0, 0, \frac{1}{\cosh t \sin r \sin \theta}\right), & e_{(1)}^\alpha &= \left(0, 0, \frac{1}{\cosh t \sin r}, 0\right) \end{aligned}$$

takes the form

$$\left[ i\gamma^0 \cosh t \left( \partial_t + \frac{3}{2} \tanh t \right) + \frac{1}{\sin r} \Sigma_{\theta\phi} + i\gamma^3 \left( \partial_r + \frac{1}{\tan r} \right) - m \cosh t \right] \Psi = 0,$$

where

$$\Sigma_{\theta,\phi} = i\gamma^1 \partial_\theta + \gamma^2 \frac{i\partial_\phi + i\sigma^{12} \cos \theta}{\sin \theta}.$$

Separating a simple factor in the wave function

$$\Psi(x) = \frac{1}{\sin r} \frac{1}{\cosh^{3/2} t} \varphi(x),$$

we obtain a simpler equation

$$\left[ i\gamma^0 \cosh t \frac{\partial}{\partial t} + i\gamma^3 \frac{\partial}{\partial r} + \frac{1}{\sin r} \Sigma_{\theta\phi} - m \cosh t \right] \varphi = 0.$$

To diagonalize the total angular moment on the wave functions, we are to use the following substitution [14] (Wigner's functions [15] are referred as  $D_{-m,\sigma}^j(\phi, \theta, 0) \equiv D_\sigma$ ):

$$\varphi_{jm}(x) = \begin{vmatrix} f_1(t, r) & D_{-1/2} \\ f_2(t, r) & D_{+1/2} \\ f_3(t, r) & D_{-1/2} \\ f_4(t, r) & D_{+1/2} \end{vmatrix}. \quad (2.1)$$

With the use of recurrent relations [15]

$$\begin{aligned} \partial_\theta D_{+1/2} &= a D_{-1/2} - b D_{+3/2}, \\ \frac{-m - 1/2 \cos \theta}{\sin \theta} D_{+1/2} &= -a D_{-1/2} - b D_{+3/2}, \\ \partial_\theta D_{-1/2} &= b D_{-3/2} - a D_{+1/2}, \\ \frac{-m + 1/2 \cos \theta}{\sin \theta} D_{-1/2} &= -b D_{-3/2} - a D_{+1/2}, \end{aligned}$$

where

$$a = \frac{j + 1/2}{2}, \quad b = \frac{1}{2} \sqrt{(j - 1/2)(j + 3/2)},$$

we find the action of the angular operator  $\Sigma_{\theta\phi}$

$$\Sigma_{\theta,\phi} \varphi_{jm}(x) = i\nu \begin{vmatrix} -f_4(t, r) & D_{-1/2} \\ +f_3(t, r) & D_{+1/2} \\ +f_2(t, r) & D_{-1/2} \\ -f_1(t, r) & D_{+1/2} \end{vmatrix}, \quad (2.2)$$

where  $\nu = j + 1/2$ . Then produce four equations in variables  $(t, r)$ :

$$\begin{aligned} i \cosh t \frac{\partial}{\partial t} f_3 - i \frac{\partial}{\partial r} f_3 - i \frac{\nu}{\sin r} f_4 - m \cosh t f_1 &= 0, \\ i \cosh t \frac{\partial}{\partial t} f_4 + i \frac{\partial}{\partial r} f_4 + i \frac{\nu}{\sin r} f_3 - m \cosh t f_2 &= 0, \\ i \cosh t \frac{\partial}{\partial t} f_1 + i \frac{\partial}{\partial r} f_1 + i \frac{\nu}{\sin r} f_2 - m \cosh t f_3 &= 0, \\ i \cosh t \frac{\partial}{\partial t} f_2 - i \frac{\partial}{\partial r} f_2 - i \frac{\nu}{\sin r} f_1 - m \cosh t f_4 &= 0. \end{aligned} \quad (2.3)$$

Usual  $P$ -reflection operator in Cartesian basis of the tetrad  $\hat{\Pi}_C = i\gamma^0 \otimes \hat{P}$

$$\hat{\Pi}_{C.} = \begin{vmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \\ i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{vmatrix} \otimes \hat{P}, \quad \hat{P}(\theta, \phi) = (\pi - \theta, \phi + \pi)$$

after translating it to spherical tetrad will take the form

$$\hat{\Pi}_{sph} = \begin{vmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{vmatrix} \otimes \hat{P}. \quad (2.4)$$

From eigenvalue equation  $\hat{\Pi}_{sph}\Psi_{jm} = \Pi \Psi_{jm}$  with the us of the known identity [15]

$$\hat{P} D_{-m,\sigma}^j(\phi, \theta, 0) = (-1)^j D_{-m,-\sigma}^j(\phi, \theta, 0)$$

we derive linear restrictions of the functions:

$$\Pi = \delta(-1)^{j+1}, \quad \delta = \pm 1, \quad f_4 = \delta f_1, \quad f_3 = \delta f_2.$$

Thus, the wave function for states with fixed parity takes the form

$$\varphi(x)_{jm\delta} = \begin{vmatrix} f_1(t, r) D_{-1/2} \\ f_2(t, r) D_{+1/2} \\ \delta f_2(t, r) D_{-1/2} \\ \delta f_1(t, r) D_{+1/2} \end{vmatrix}. \quad (2.5)$$

Allowing for (2.5), from (2.3) we get more simple equations

$$\begin{aligned} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sin r} \right) f + \left( i \cosh t \frac{\partial}{\partial t} + \delta m \cosh t \right) g &= 0, \\ \left( \frac{\partial}{\partial r} - \frac{\nu}{\sin r} \right) g - \left( i \cosh t \frac{\partial}{\partial t} - \delta m \cosh t \right) f &= 0, \end{aligned} \quad (2.6)$$

where instead of  $f_1(t, r)$  and  $f_2(t, r)$  their linear combinations are used:

$$f(t, r) = \frac{f_1 + f_2}{\sqrt{2}}, \quad g(t, r) = \frac{f_1 - f_2}{i\sqrt{2}}.$$

Now, let us perform nonrelativistic approximation in the system (2.6) (here we will adhere the method exposed in [13, 14]). First, we separate the rest energy by the formal change

$$i \frac{\partial}{\partial t} \implies M + i \frac{\partial}{\partial t}.$$

The cases  $\delta = \pm 1$  should be considered separately:

$$\delta = +1,$$

$$\begin{aligned} \frac{1}{\cosh t} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sin r} \right) f + \left( M + i \frac{\partial}{\partial t} + M \right) g &= 0, \\ \frac{1}{\cosh t} \left( \frac{\partial}{\partial r} - \frac{\nu}{\sin r} \right) g - \left( M + i \frac{\partial}{\partial t} - M \right) f &= 0; \end{aligned} \quad (2.7)$$

$\delta = -1$ ,

$$\begin{aligned} \frac{1}{\cosh t} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sin r} \right) f + \left( M + i \frac{\partial}{\partial t} - M \right) g &= 0, \\ \frac{1}{\cosh t} \left( \frac{\partial}{\partial r} - \frac{\nu}{\sin r} \right) g - \left( M + i \frac{\partial}{\partial t} + M \right) f &= 0; \end{aligned} \quad (2.8)$$

the term  $i\partial_t$  should be neglected in comparing with  $2M$ . Equations (2.7), (2.8) give respectively  
 $\delta = +1$ ,

$$\begin{aligned} \frac{1}{\cosh t} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sin r} \right) f + 2Mg &= 0, \\ \frac{1}{\cosh t} \left( \frac{\partial}{\partial r} - \frac{\nu}{\sin r} \right) g - i \frac{\partial}{\partial t} f &= 0; \end{aligned}$$

$\delta = -1$ ,

$$\begin{aligned} \frac{1}{\cosh t} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sin r} \right) f + i \frac{\partial}{\partial t} g &= 0, \\ \frac{1}{\cosh t} \left( \frac{\partial}{\partial r} - \frac{\nu}{\sin r} \right) g - 2Mf &= 0. \end{aligned}$$

In each case, we derive an equation for the big component (the concomitant equation permits to express the small component via the big one)

$\delta = +1$ ,

$$\begin{aligned} i \frac{\partial}{\partial t} f &= -\frac{1}{2M} \frac{1}{\cosh^2 t} \left( \frac{\partial}{\partial r} - \frac{\nu}{\sin r} \right) \left( \frac{\partial}{\partial r} + \frac{\nu}{\sin r} \right) f, \\ g &= -\frac{1}{2M} \frac{1}{\cosh t} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sin r} \right) f; \end{aligned} \quad (2.9)$$

$\delta = -1$ ,

$$\begin{aligned} i \frac{\partial}{\partial t} g &= -\frac{1}{2M} \frac{1}{\cosh t} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sin r} \right) \left( \frac{\partial}{\partial r} - \frac{\nu}{\sin r} \right) g, \\ f &= \frac{1}{2M} \frac{1}{\cosh^2 t} \left( \frac{\partial}{\partial r} - \frac{\nu}{\sin r} \right) g. \end{aligned} \quad (2.10)$$

Recall that the Pauli wave function is constructed from relativistic one in accordance with

$$\varphi_{jm\delta}(x) = \begin{vmatrix} f_1 D_{-1/2} \\ f_2 D_{+1/2} \\ \delta f_2 D_{-1/2} \\ \delta f_1 D_{+1/2} \end{vmatrix} \implies \Psi_{jm\delta}^{Pauli}(x) = \begin{vmatrix} f_1 D_{-1/2} + \delta f_2 D_{-1/2} \\ f_2 D_{+1/2} + \delta f_1 D_{+1/2} \end{vmatrix}. \quad (2.11)$$

For different parities, eq. (2.11) gives respectively

$\delta = +1$ ,

$$\Psi_{jm,+1}^{Pauli}(t, r, \theta, \phi) = \begin{vmatrix} f_1 D_{-1/2} + f_2 D_{-1/2} \\ f_2 D_{+1/2} f_1 D_{+1/2} \end{vmatrix} = \begin{vmatrix} f(t, r) D_{-1/2} \\ f(t, r) D_{+1/2} \end{vmatrix};$$

$\delta = -1$ ,

$$\Psi_{jm,-1}^{Pauli}(t, r, \theta, \phi) = \begin{vmatrix} f_1 D_{-1/2} - f_2 D_{-1/2} \\ f_2 D_{+1/2} - f_1 D_{+1/2} \end{vmatrix} = \begin{vmatrix} -ig(t, r) D_{-1/2} \\ -ig(t, r) D_{+1/2} \end{vmatrix}.$$

Note that according to (2.9) and (2.10), non-relativistic functions  $f(t, r)$  and  $g(t, r)$  obey similar but slightly different equations:

$\delta = +1$ ,

$$i \frac{\partial f}{\partial t} = -\frac{1}{2M} \frac{1}{\cosh^2 t} \left( \frac{\partial^2}{\partial r^2} - \frac{\nu^2 + \nu \cos r}{\sin^2 r} \right) f; \quad (2.12)$$

$\delta = -1$ ,

$$i \frac{\partial g}{\partial t} = -\frac{1}{2M} \frac{1}{\cosh^2 t} \left( \frac{\partial^2}{\partial r^2} - \frac{\nu^2 - \nu \cos r}{\sin^2 r} \right) g. \quad (2.13)$$

In (2.12), (2.13), the variables are separated by the substitutions

$$f(t, r) = f(t)f(r), \quad g(t, r) = g(t)g(r);$$

in this way we get

$\delta = +1$ ,

$$i \cosh^2 t \frac{1}{f(t)} \frac{d}{dt} f(t) = -\frac{1}{2M} \frac{1}{f(r)} \left( \frac{d^2}{dr^2} - \frac{\nu^2 - \nu \cos r}{\sin^2 r} \right) f(r) = E;$$

$\delta = -1$ ,

$$i \cosh^2 t \frac{1}{g(t)} \frac{d}{dt} g(t) = -\frac{1}{2M} \frac{1}{g(r)} \left( \frac{d^2}{dr^2} - \frac{\nu^2 + \nu \cos r}{\sin^2 r} \right) g(r) = E.$$

The final separated equations are

$\delta = +1$ ,

$$\begin{aligned} i \cosh^2 t \frac{1}{f(t)} \frac{d}{dt} f(t) = E &\implies f(t) = e^{-iE \tanh t}, \\ -\frac{1}{2M} \frac{1}{f(r)} \left( \frac{d^2}{dr^2} - \frac{\nu^2 + \nu \cos r}{\sin^2 r} \right) f(r) = E; \end{aligned}$$

$\delta = -1$ ,

$$\begin{aligned} i \cosh^2 t \frac{1}{g(t)} \frac{d}{dt} g(t) = E &\implies g(t) = e^{-iE \tanh t}, \\ -\frac{1}{2M} \frac{1}{g(r)} \left( \frac{d^2}{dr^2} - \frac{\nu^2 - \nu \cos r}{\sin^2 r} \right) g(r) = E. \end{aligned}$$

Note the symmetry between equations for  $f(r)$  and  $g(r)$ :  $\nu \implies -\nu$ ; therefore it suffices to examine in detail only one case and then to employ the formal change. For definiteness, we will consider the variant  $\delta = +1$ :

$$\left( \frac{d^2}{dr^2} - \frac{\nu^2 + \nu \cos r}{\sin^2 r} + 2ME \right) f(r) = 0. \quad (2.14)$$

Because the 3-subspace in de Sitter space-time coincides with the compact spherical Riemann model, the motion if the variable  $r$  must be quantized; besides, we must assume that  $2ME > 0$ . In eq. (2.14), it is convenient to use a new variable,  $z = \cos r$ ,  $z \in (-1, +1)$ :

$$(1 - z^2) \frac{d^2 f}{dz^2} - z \frac{df}{dz} - \left( \nu \frac{z}{1 - z^2} + \frac{\nu^2}{1 - z^2} - 2ME \right) f = 0,$$

and then to introduce another variable

$$y = \frac{1 - z}{2} = \frac{1 - \cos r}{2},$$

$$y(1 - y) \frac{d^2 f}{dy^2} + \left( \frac{1}{2} - y \right) \frac{df}{dy} - \left[ \frac{\nu(\nu + 1)}{4y} + \frac{\nu(\nu - 1)}{4(1 - y)} - 2ME \right] f = 0.$$

With the substitution  $f = y^A(1 - y)^B F$ , we get

$$y(1 - y) \frac{d^2 F}{dy^2} + \left[ 2A + \frac{1}{2} - (2A + 2B + 1)y \right] \frac{dF}{dy}$$

$$- \left[ (A + B)^2 - \frac{2A(2A - 1) - \nu(\nu + 1)}{4y} - \frac{2B(2B - 1) - \nu(\nu - 1)}{4(1 - y)} - 2ME \right] F = 0.$$

At restrictions

$$2A = \nu + 1, -\nu; \quad 2B = -\nu + 1, +\nu,$$

the above equation simplifies

$$y(1 - y) \frac{d^2 F}{dy^2} + \left[ 2A + \frac{1}{2} - (2A + 2B + 1)y \right] \frac{dF}{dy} - [(A + B)^2 - 2ME] F = 0,$$

and coincides with the equation of hypergeometric type

$$y(1 - y) F'' + [(c - (a + b + 1)y) F' - ab F] = 0,$$

where

$$c = 2A + \frac{1}{2}, \quad a + b = 2A + 2B, \quad ab = (A + B)^2 - 2ME;$$

from whence it follows

$$a = A + B - \sqrt{2ME}, \quad b = A + B + \sqrt{2ME}.$$

From physical considerations, we should construct solutions in polynomials and they should be associated with discrete spectrum of energy  $2ME > 0$ . Appropriate solutions are possible at positive  $A$  and  $B$ :

$$2A = +\nu + 1 = j + 3/2, \quad 2B = +\nu = j + 1/2,$$

polynomial condition provides us with the quantization rule

$$a = -n, \quad n = 0, 1, 2, \dots \quad 2ME = (j + 1 + n)^2;$$

corresponding radial functions are

$$f(y) = y^{(\nu+1)/2}(1 - y)^{\nu/2} F(a, b, c, y),$$

$$b = 2(j + 1) + n, \quad c = j + 2 \quad (\nu = j + 1/2).$$

With the use of the mentioned symmetry between  $f(r)$  and  $g(r)$ , one gets description of states with other parity

$$g(y) = y^{A'}(1-y)^{B'}G(y), \\ 2A' = -\nu + 1, \nu, \quad 2B' = \nu + 1, -\nu;$$

at positive  $A'$  and  $B'$ :

$$2A' = \nu = j + 1/2, \quad c' = j + 1 = c - 1, \\ 2B' = +\nu + 1 = j + 3/2, \quad A' + B' = j + 1, \\ a' = -n = a, \quad b' = 2(j + 1) + n = b, \\ g(y) = y^{\nu/2}(1-y)^{(\nu+1)/2}F(a, b, c - 1, y).$$

Thus, solutions of the Pauli equation in non-static de Sitter metrics have been constructed  
 $\delta = +1$ ,

$$\Psi_{Ejm,+1}^{Pauli}(t, r, \theta, \phi) = e^{-iE\tanh t}f(r) \begin{vmatrix} D_{-m,-1/2}^j(\phi, \theta, 0) \\ D_{-m,+1/2}^j(\phi, \theta, 0) \end{vmatrix}, \\ f(r) = \left(\sin \frac{r}{2}\right)^{j+3/2} \left(\cos \frac{r}{2}\right)^{j+1/2} F(-n, n + 2j + 2, j + 2, \sin^2 \frac{r}{2});$$

$\delta = -1$ ,

$$\Psi_{Ejm,-1}^{Pauli}(t, r, \theta, \phi) = e^{-iE\tanh t}g(r) \begin{vmatrix} D_{-m,-1/2}^j(\phi, \theta, 0) \\ D_{-m,+1/2}^j(\phi, \theta, 0) \end{vmatrix}, \\ g(r) = \left(\sin \frac{r}{2}\right)^{j+1/2} \left(\cos \frac{r}{2}\right)^{j+3/2} F(-n, n + 2j + 2, j + 1, \sin^2 \frac{r}{2}).$$

Spectral parameter  $E$  (it does not represent the energy of stationary states) is quantized according to the rule (the same for state with different parities)

$$2ME = (j + 1 + n)^2.$$

### 3 Pauli equation in the oscillating anti de Sitter Universe

In the non-static anti de Sitter metric

$$x^\alpha = (t, r, \theta, \phi), \quad dS^2 = dt^2 - \cos^2 t [dr^2 + \sinh^2 r (d\theta^2 + \sin^2 \theta d\phi^2)]$$

and the corresponding diagonal tetrad

$$e_{(0)}^\alpha = (1, 0, 0, 0), \quad e_{(3)}^\alpha = \left(0, \frac{1}{\cos t}, 0, 0\right), \\ e_{(2)}^\alpha = \left(0, 0, 0, \frac{1}{\cos t \sinh r \sin \theta}\right), \quad e_{(1)}^\alpha = \left(0, 0, \frac{1}{\cos t \sinh r}, 0\right),$$

generally covariant Dirac equation takes the form

$$\left[ i\gamma^0 \cos t (\partial_t - \frac{3}{2} \tan t) + \frac{1}{\sinh r} \Sigma_{\theta\phi} + i\gamma^3 (\partial_r + \frac{1}{\tanh r}) - m \cos t \right] \Psi = 0.$$

Separating a simper multiplier in the wave function

$$\Psi(x) = \frac{1}{\sinh r} \frac{1}{\cos^{3/2} t} \varphi(x),$$

we obtain a more simple form

$$\left[ i\gamma^0 \cos t \frac{\partial}{\partial t} + i\gamma^3 \frac{\partial}{\partial r} + \frac{1}{\sinh r} \Sigma_{\theta\phi} - m \cos t \right] \varphi = 0.$$

Using the technique of  $D$ -Wigner functions (2.1)–(2.2), we obtain the radial equation

$$\begin{aligned} i \cos t \frac{\partial}{\partial t} f_3 - i \frac{\partial}{\partial r} f_3 - i \frac{\nu}{\sinh r} f_4 - m \cos t f_1 &= 0, \\ i \cos t \frac{\partial}{\partial t} f_4 + i \frac{\partial}{\partial r} f_4 + i \frac{\nu}{\sinh r} f_3 - m \cos t f_2 &= 0, \\ i \cos t \frac{\partial}{\partial t} f_1 + i \frac{\partial}{\partial r} f_1 + i \frac{\nu}{\sinh r} f_2 - m \cos t f_3 &= 0, \\ i \cos t \frac{\partial}{\partial t} f_2 - i \frac{\partial}{\partial r} f_2 - i \frac{\nu}{\sinh r} f_1 - m \cos t f_4 &= 0. \end{aligned}$$

Using the operator of spatial parity (2.4)–(2.5), we get more simple equations

$$\begin{aligned} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sinh r} \right) f + \left( i \cos t \frac{\partial}{\partial t} + \delta m \cos t \right) g &= 0, \\ \left( \frac{\partial}{\partial r} - \frac{\nu}{\sinh r} \right) g - \left( i \cos t \frac{\partial}{\partial t} - \delta m \cos t \right) f &= 0. \end{aligned} \quad (3.1)$$

Now, let us perform the nonrelativistic approximation in the system (3.1). First, we separate the rest energy. The cases  $\delta = \pm 1$  should be considered separately:

$$\delta = +1,$$

$$\begin{aligned} \frac{1}{\cos t} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sinh r} \right) f + \left( M + i \frac{\partial}{\partial t} + M \right) g &= 0, \\ \frac{1}{\cos t} \left( \frac{\partial}{\partial r} - \frac{\nu}{\sinh r} \right) g - \left( M + i \frac{\partial}{\partial t} - M \right) f &= 0; \end{aligned}$$

$$\delta = -1,$$

$$\begin{aligned} \frac{1}{\cos t} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sinh r} \right) f + \left( M + i \frac{\partial}{\partial t} - M \right) g &= 0, \\ \frac{1}{\cos t} \left( \frac{\partial}{\partial r} - \frac{\nu}{\sinh r} \right) g - \left( M + i \frac{\partial}{\partial t} + M \right) f &= 0. \end{aligned}$$

These system give respectively (the term  $i\partial_t$  should be neglected in comparing with  $2M$ )

$$\delta = +1,$$

$$\begin{aligned} \frac{1}{\cos t} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sinh r} \right) f + 2Mg &= 0, \\ \frac{1}{\cos t} \left( \frac{\partial}{\partial r} - \frac{\nu}{\sinh r} \right) g - i \frac{\partial}{\partial t} f &= 0; \end{aligned}$$

$\delta = -1$ ,

$$\begin{aligned} \frac{1}{\cos t} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sinh r} \right) f + i \frac{\partial}{\partial t} g &= 0, \\ \frac{1}{\cos t} \left( \frac{\partial}{\partial r} - \frac{\nu}{\sinh r} \right) g - 2Mf &= 0. \end{aligned}$$

For both cases, we derive equations

$\delta = +1$ ,

$$\begin{aligned} i \frac{\partial}{\partial t} f &= -\frac{1}{2M} \frac{1}{\cos^2 t} \left( \frac{\partial}{\partial r} - \frac{\nu}{\sinh r} \right) \left( \frac{\partial}{\partial r} + \frac{\nu}{\sinh r} \right) f, \\ g &= -\frac{1}{2M} \frac{1}{\cos t} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sinh r} \right) f; \end{aligned}$$

$\delta = -1$ ,

$$\begin{aligned} i \frac{\partial}{\partial t} g &= -\frac{1}{2M} \frac{1}{\cos^2 t} \left( \frac{\partial}{\partial r} + \frac{\nu}{\sinh r} \right) \left( \frac{\partial}{\partial r} - \frac{\nu}{\sinh r} \right) g, \\ f &= \frac{1}{2M} \frac{1}{\cos t} \left( \frac{\partial}{\partial r} - \frac{\nu}{\sinh r} \right) g. \end{aligned}$$

Nonrelativistic functions  $f(t, r)$  and  $g(t, r)$  obey similar equations

$\delta = +1$ ,

$$i \frac{\partial}{\partial t} f = -\frac{1}{2M} \frac{1}{\cos^2 t} \left( \frac{\partial^2}{\partial r^2} - \frac{\nu^2 + \nu \cosh r}{\sinh^2 r} \right) f; \quad (3.2)$$

$\delta = -1$ ,

$$i \frac{\partial}{\partial t} g = -\frac{1}{2M} \frac{1}{\cos^2 t} \left( \frac{\partial^2}{\partial r^2} - \frac{\nu^2 - \nu \cosh r}{\sinh^2 r} \right) g. \quad (3.3)$$

In (3.2), (3.3) the variables are separated by the substitutions

$$f(t, r) = f(t)f(r), \quad g(t, r) = g(t)g(r),$$

in this way we get

$\delta = +1$ ,

$$i \cos^2 t \frac{1}{f(t)} \frac{d}{dt} f(t) = -\frac{1}{2M} \frac{1}{f(r)} \left( \frac{d^2}{dr^2} - \frac{\nu^2 + \nu \cosh r}{\sinh^2 r} \right) f(r) = E; \quad (3.4)$$

$\delta = -1$ ,

$$i \cos^2 t \frac{1}{g(t)} \frac{d}{dt} g(t) = -\frac{1}{2M} \frac{1}{g(r)} \left( \frac{d^2}{dr^2} - \frac{\nu^2 - \nu \cosh r}{\sinh^2 r} \right) g(r) = E. \quad (3.5)$$

The final separated equations are

$\delta = +1$ ,

$$\begin{aligned} i \cos^2 t \frac{1}{f(t)} \frac{d}{dt} f(t) = E &\implies f(t) = e^{-iE \tan t}, \\ -\frac{1}{2M} \frac{1}{f(r)} \left( \frac{d^2}{dr^2} - \frac{\nu^2 + \nu \cosh r}{\sinh^2 r} \right) f(r) &= E; \end{aligned}$$

$\delta = -1$ ,

$$\begin{aligned} i \cos^2 t \frac{1}{g(t)} \frac{d}{dt} g(t) &= E \implies g(t) = e^{-iE \tan t}, \\ -\frac{1}{2M} \frac{1}{g(r)} \left( \frac{d^2}{dr^2} - \frac{\nu^2 - \nu \cosh r}{\sinh^2 r} \right) g(r) &= E. \end{aligned}$$

Due to the symmetry between equations for  $f(r)$  and  $g(r)$ :  $\nu \implies -\nu$ ; it suffices to examine in detail only one case and then to employ the formal change. For definiteness, we will consider the variant  $\delta = +1$ :

$$\left( \frac{d^2}{dr^2} - \frac{\nu^2 + \nu \cosh r}{\sinh^2 r} + 2ME \right) f(r) = 0. \quad (3.6)$$

In eq. (3.6), it is convenient to use the new variable  $z = \cosh r$ ,  $z \in (1, +\infty)$ :

$$(1 - z^2) \frac{d^2 f}{dz^2} - z \frac{df}{dz} - \left( \nu \frac{z}{1 - z^2} + \frac{\nu^2}{1 - z^2} + 2ME \right) f = 0,$$

and then to introduce another variable

$$\begin{aligned} y &= \frac{1 - z}{2} = \frac{1 - \cosh r}{2}, \\ y(1 - y) \frac{d^2 f}{dy^2} + \left( \frac{1}{2} - y \right) \frac{df}{dy} - \left[ \frac{\nu(\nu + 1)}{4y} + \frac{\nu(\nu - 1)}{4(1 - y)} + 2ME \right] f &= 0. \end{aligned}$$

With the substitution  $f = y^A(1 - y)^B F$ , we get

$$\begin{aligned} y(1 - y) \frac{d^2 F}{dy^2} + \left[ \frac{1}{2} + 2A - (1 + 2A + 2B)y \right] \frac{dF}{dy} \\ - \left[ (A + B)^2 - \frac{2A(2A - 1) - \nu(\nu + 1)}{4y} - \frac{2B(2B - 1) - \nu(\nu - 1)}{4(1 - y)} + 2ME \right] F = 0. \end{aligned}$$

At additional restriction

$$A = -\frac{1}{2}\nu, \quad \frac{1}{2} + \frac{1}{2}\nu; \quad B = \frac{1}{2} - \frac{1}{2}\nu, \quad \frac{1}{2}\nu$$

the above equation simplifies

$$y(1 - y) \frac{d^2 F}{dy^2} + \left[ \frac{1}{2} + 2A - (1 + 2A + 2B)y \right] \frac{dF}{dy} - [(A + B)^2 + 2ME] F = 0,$$

and coincides with the equation of hypergeometric type for  $F(a, b, c, y)$  with

$$c = 2A + \frac{1}{2}, \quad a = A + B - i\sqrt{2ME}, \quad b = A + B + i\sqrt{2ME}.$$

Corresponding solutions are

$$a = \frac{1}{2} + \nu - i\sqrt{2ME}, \quad b = \frac{1}{2} + \nu + i\sqrt{2ME}, \quad c = \frac{3}{2} + \nu = j + 2,$$

$$a = 1 + j - i\sqrt{2ME}, \quad b = 1 + j + i\sqrt{2ME}, \quad f(r) = C y^{(\nu+1)/2} (1 - y)^{\nu/2} F(a, b, c, y).$$

Allowing for the mentioned symmetry between  $f(r)$  and  $g(r)$ , one gets description of states with other parity:

$$\begin{aligned} g &= C' y^{A'} (1-y)^{B'} G, \quad A' = \frac{1}{2} \nu, \frac{1}{2} - \frac{1}{2} \nu, \quad B' = \frac{1}{2} + \frac{1}{2} \nu, -\frac{1}{2} \nu, \\ a' &= 1 + j - i\sqrt{2ME}, \quad b' = 1 + j + i\sqrt{2ME}, \quad c' = j + 1 = c - 1, \\ g(r) &= C' y^{\nu/2} (1-y)^{(\nu+1)/2} F(a, b, c-1, y). \end{aligned}$$

Thus, solutions of the Pauli equation in non-static anti de Sitter metrics have been constructed

$$\begin{aligned} \delta = +1, \quad \Psi_{Ejm,+1}^{Pauli}(t, r, \theta, \phi) &= e^{-iE \tan t} f(r) \begin{vmatrix} D_{-m, -1/2}^j(\phi, \theta, 0) \\ D_{-m, +1/2}^j(\phi, \theta, 0) \end{vmatrix}, \\ f(r) &= C \left( \frac{1 - \cosh r}{2} \right)^{(\nu+1)/2} \left( \frac{1 + \cosh r}{2} \right)^{\nu/2} F \left( -n, n + 2\nu + 1, \nu + \frac{3}{2}, \frac{1 - \cosh r}{2} \right); \\ \delta = -1, \quad \Psi_{Ejm,-1}^{Pauli}(t, r, \theta, \phi) &= e^{-iE \tan t} g(r) \begin{vmatrix} D_{-m, -1/2}^j(\phi, \theta, 0) \\ D_{-m, +1/2}^j(\phi, \theta, 0) \end{vmatrix}, \\ g(r) &= C' \left( \frac{1 - \cosh r}{2} \right)^{\nu/2} \left( \frac{1 + \cosh r}{2} \right)^{(\nu+1)/2} F \left( -n, n + 2\nu + 1, \nu + \frac{1}{2}, \frac{1 - \cosh r}{2} \right). \end{aligned}$$

### Acknowledgements

This work was supported by the Fund for Basic Researches of Belarus, F 13K-079, within the cooperation framework between Belarus and Ukraine, and by the Fund for Basic Researches of Belarus, F 14ARM-021, within the cooperation framework between Republic of Belarus and Republic of Armenia.

The authors are grateful to V.M. Red'kov for advice and help.

## References

- [1] P.A.M. Dirac. The electron wave equation in the de Sitter space. *Ann. Math.* **36**, 657–669 (1935).
- [2] K. Goto. Wave equations in de Sitter space. *Progr. Theor. Phys.* **6**, 1013–1014 (1951).
- [3] O. Nachtmann. Quantum theory in de-Sitter space. *Commun. Math. Phys.* **6**, 1–16 (1967).
- [4] N.A. Chernikov, E.A. Tagirov. Quantum Theory of Scalar Field in de Sitter Space-Time. *Ann. Inst. H. Poincaré. A.* **9**, 109–141 (1968).
- [5] G. Börner, H.P. Dürr. Classical and quantum theory in de Sitter space. *Nuovo Cim. A.* **64**, 669–713 (1969).
- [6] D. Lohiya, N. Panchapakesan. Particle emission in the de Sitter universe for massless fields with spin. *J. Phys. A.* **12**, 533–539 (1979).
- [7] F. Riordan. Solutions of the Dirac equation in finite de Sitter space. *Nuovo Cim. B.* **20**, 309–325 (1974).
- [8] V.S. Ouchik. On the Hawking radiation of spin 1/2 particles in the de Sitter space-time. *Class. Quantum Grav.* **2**, 539–543 (1985).

- [9] Takashi Mishima, Akihiro Nakayama. Particle production in de Sitter spacetime. *Progr. Theor. Phys.* **77**, 218–222 (1987).
- [10] H. Suzuki, E. Takasugi. Absorption Probability of De Sitter Horizon for Massless Fields with Spin. *Mod. Phys. Lett. A.* **11**, 431–436 (1996).
- [11] V.M. Red'kov, E.M. Ovsiyuk. On exact solutions for quantum particles with spin  $S = 0, 1/2, 1$  and de Sitter event horizon. *Ricerche di matematica.* **60**, 57–88 (2011).
- [12] V.M Red'kov, E.M. Ovsiyuk, G.G. Krylov. Hawking Radiation in de Sitter Space: Calculation of the Reflection Coefficient for Quantum Particles. *Vestnik RUDN.* **4**, 153–169 (2012).
- [13] V.M. Red'kov. Particle fields in Riemannian space and the Lorentz group. Minsk, 2009 (in Russian).
- [14] V.M Red'kov. Tetrad formalism, spherical symmetry and the basis of Schrödinger. Minsk, 2010 (in Russian).
- [15] D.A. Varshalovich, A.N. Moskalev, V.K. Hersonskiy. Quantum theory of angular moment. Leningrad, 1975 (in Russian).