

Conformal Correlation Functions in the Brownian Loop Soup

Federico Camia,^{★♠} Alberto Gandolfi,^{★♣} and Matthew Kleban^{★♡}

[★]*New York University Abu Dhabi, United Arab Emirates*

[♠]*VU University, Amsterdam, the Netherlands*

[♣]*University of Florence, Italy*

[♡]*Center for Cosmology and Particle Physics, Department of Physics, New York University, United States of America*

E-mail: federico.camia@nyu.edu, albertogandolfi@nyu.edu,
kleban@nyu.edu

ABSTRACT: We define and study a set of operators that compute statistical properties of the Brownian Loop Soup, a conformally invariant gas of random Brownian loops (Brownian paths constrained to begin and end at the same point) in two dimensions. We prove that the correlation functions of these operators have many of the properties of conformal primaries in a conformal field theory, and compute their conformal dimension. The dimensions are real and positive, but have the novel feature that they vary continuously as a periodic function of a real parameter. We comment on the relation of the Brownian Loop Soup to the free field, and use this relation to establish that the central charge of the Loop Soup is twice its intensity.

Contents

1	Introduction	1
1.1	The Brownian Loop Soup	1
1.2	Motivation	3
1.3	Correlation functions in the Brownian Loop Soup	4
2	Summary and results	6
3	Correlators of the layering and winding operators	8
3.1	Correlators of layering operator	8
3.2	The 1-point function in the layering model	9
3.3	The winding operator	10
3.4	The 2-point function in the layering model	11
4	Conformal covariance of the n-point functions	14
4.1	The layering model in finite domains	15
4.2	The winding model in finite domains	19
4.3	The layering model in the plane	20
5	Dimension of the exponential winding operator	22
6	Central charge and relation to the free boson	24
7	Conclusions	25
8	Appendix. The Brownian loop measure: two lemmas	26

1 Introduction

1.1 The Brownian Loop Soup

Take a handful of loops of various sizes and sprinkle them onto a flat surface. The position where each loop lands is uniformly random, independent of any loops already in place. Each loop is Brownian – a Brownian motion constrained to begin and end at the same “root” point, but otherwise with no restriction – and characterized by a “time”




Figure 1: Two Brownian loops, each of time length $t = 1$.

length t that is linearly related to its average area (*cf.* Fig. 1). The distribution in t is $\sim dt/t^2$, so that there are many more small loops than large, and is chosen to ensure invariance under scale transformations. The overall density of loops is characterized by a single parameter: the “intensity” $\lambda > 0$. This random ensemble of loops is called the Brownian Loop Soup (BLS) and was introduced in [1].

More precisely, the BLS is a Poissonian random collection of loops in a planar domain D with intensity measure $\lambda\mu_D^{loop}$, where $\lambda > 0$ is a constant and μ_D^{loop} is the restriction to D of the *Brownian loop measure*

$$\mu^{loop} = \int_{\mathbb{C}} \int_0^\infty \frac{1}{2\pi t^2} \mu_{z,t}^{br} dt d\mathbf{A}(z), \quad (1.1)$$

where \mathbf{A} denotes area and $\mu_{z,t}^{br}$ is the complex Brownian bridge measure with starting point z and duration t . We note that the Brownian loop measure should be interpreted as a measure on “unrooted” loops, that is, loops without a specified “root” point. (Formally, unrooted loops are equivalence classes of rooted loops—the interested reader is referred to [1] for the details.) For ease of notation, the μ^{loop} -measure of a set $\{\dots\}$ will be denoted $\mu^{loop}(\dots) \equiv \mu^{loop}(\{\dots\})$.

The BLS turns out to be not just scale invariant, but fully conformally invariant. For sufficiently low intensities λ , the intersecting loops form clusters whose outer boundaries are distributed like Conformal Loop Ensembles (CLEs)[2]. CLEs are the unique ensembles of planar, non-crossing and non-self-crossing loops satisfying a natural conformal restriction property that is conjecturally satisfied by the continuum scaling limits of interfaces in two-dimensional models from statistical physics. The loops of a CLE_κ are forms of SLE_κ (the Schramm-Loewner Evolution with parameter κ [3]). The




Figure 2: The Brownian loop soup for intensities $\lambda = 2.5$ (left panel) and $\lambda = .5$ (right panel). The plots were made with both a “UV” cutoff on short timelengths and an “IR” cutoff on long timelengths. The true BLS is a scale-invariant fractal that covers every point with probability 1.

CLEs generated by the BLS correspond to values of κ between $8/3$ and 4 . For example, the collection of outermost interfaces in a planar critical Ising model in a finite domain with plus boundary condition is conjectured to converge to CLE_3 in the scaling limit.

In this paper we will define and compute certain statistical correlation functions that characterize aspects of the BLS distribution. We will focus on two types of information: the number of distinct loops that surround (or “cover” if one thinks of the loops as being filled in) a given point or set of points, and the net number of windings of all the loops around a given point or points (see Fig. 3). In both cases, we find results consistent with the correlation functions of primary operators in a conformal field theory.

1.2 Motivation

We have several motivations for this work. In [4] one of us considered a similar model, where instead of Brownian loops one sprinkles disks. The 2- and 3-point correlation functions of certain operators in that model behave like those of a conformal field theory (CFT), and with a novel set of conformal dimensions. This is of interest because the model of [4] was derived in [5] as an approximation to the asymptotic distribution of bubble nucleations in theories of eternal inflation, a theory for which there is some




Figure 3: A stylized Brownian loop. The numbers indicate the winding numbers of the loop that contribute additively to N_w , while the green shaded region is the interior of the loop (the set of points disconnected from infinity by the loop) that contributes ± 1 (where the sign is a Boolean variable assigned randomly to each loop) to the layering number N_ℓ .

reason to believe a CFT dual may exist [6–8]. However, [4] computed the 4-point function exactly, and it suffered from a deficiency: it was not smooth as the position of the fourth point crossed the circle connecting the other three.

The origin of the non-analyticity in [4] is probably the fact that although the disk distribution is invariant under global conformal transformations, it is not locally conformally invariant (since disks do not map to disks). By contrast the BLS distribution is fully conformally invariant. Therefore we expect the analogous correlation functions to be better behaved, perhaps defining a healthy CFT that could be related to the physics of de Sitter spacetime and eternal inflation.

Another motivation for considering CFTs related to the BLS is the relation between the BLS and SLE [3], which in turn is related to a large class of conformally invariant models ranging from percolation [9–12] to the Ising model [13].

If in fact the BLS correlation functions we study arise from or define a CFT, it does not seem to be one that is currently known, and it has several interesting and novel features. As in [4], the conformal dimensions of the primary operators are real and positive, but vary continuously and as a *periodic* function of a real parameter β . As we will see this periodicity arises because the operators are of the form $e^{i\beta N}$, where N is integer valued.

1.3 Correlation functions in the Brownian Loop Soup

As mentioned above, we will study the correlation functions (*i.e.* expectation values) of two distinct types of operators in the BLS.

Layering number: The first type is closely related to the operators considered in [4] (which we refer to as the “disk model”). For each loop, define the interior as the set

of points inside the outermost edge of the loop (including any isolated “islands” that might appear inside due to self-intersections; *cf.* Fig. 3). Points in this set are “covered” by the loop. For a point z , consider an operator $N(z)$ that counts the number of distinct loops that cover the point z , so that $N(z)$ is the number of “layers” at z .

One difficulty arises immediately. Because the BLS is conformally (and therefore scale-) invariant, any given point of the plane is covered by infinitely many loops with probability one. Since $N(\cdot) \geq 0$, this means that, for any fixed z , $\langle N(z) \rangle$ diverges with probability one. Precisely the same difficulty arises in the disk model, where it was dealt with by adding another, identical and independent copy of the distribution, and then counting the difference in the number of disks of each type that cover z .¹ We will follow a very similar procedure here, assigning to each loop a random Boolean value and then defining the layering operator $N_\ell(z) \equiv N_+(z) - N_-(z)$.

Winding number: The other operator we will discuss is $N_w(z)$, which counts the total number of windings of all loops around a point z (Fig. 3). This makes use of the fact that Brownian loops have an orientation (they grow in a particular direction as the time t increases). Since the winding number can be positive or negative, it is not necessary to include another copy of the distribution or compute a difference of two values.

A natural physical interpretation of winding in the BLS is as follows. If each loop represents the configuration of a string at some instant of time in a 2+1 dimensional spacetime, winding number counts the number of units of the flux the string is charged under (see *e.g.* [15]).

In both cases we will focus on exponentials of these operators times imaginary coefficients (correlators of the number operators themselves are afflicted by logarithmic divergences, like massless fields in two dimensions). Due to the similarity to free-field vertex operators we will denote these as

$$V_\beta(z) \equiv e^{i\beta N(z)},$$

where N can be either a layering or winding number operator.²

Our paper is written for a mixed audience of mathematicians and physicists. In most of the paper we present rigorous proofs of our results. Sections 5 and 6, where we perform “physics-style” calculations, are the exceptions. The appendix, Section 8,

¹In this disk model this can be thought of as counting the number of bubble transitions that have affected the point z , in a model where the field has a discrete shift symmetry and there is a simple rule for bubble collisions (namely the one discussed in [14]).

²A special case of the winding model on a lattice was considered by Le Jan (see Section 6 of [16]).

is dedicated to two important lemmas about the Brownian loop measure (1.1) which are used several times in the rest of the paper. A reader uninterested in our methods may simply read Sec. 2 for a summary of results, and Sec. 7 for our conclusions.

2 Summary and results

Our main results relate to correlation functions (*i.e.* expectation values of products) of exponentials of the winding and layering operators in the BLS. Specifically, we establish the following:

- For both versions in finite domains D , correlators of $n \in \mathbb{N}$ exponential operators

$$\langle \prod_j V_{\beta_j}(z_j) \rangle_{\delta, D} = \langle e^{i \sum_j \beta_j N(z_j)} \rangle_{\delta, D} \quad (2.1)$$

exist as long as a short-time cutoff $\delta > 0$ on the loops is imposed, and

$$\lim_{\delta \rightarrow 0} \frac{\langle \prod_{j=1}^n V_{\beta_j}(z_j) \rangle_{\delta, D}}{\prod_{j=1}^n \delta^{2\Delta(\beta_j)}} \equiv \phi_D(z_1, \dots, z_n; \beta_1, \dots, \beta_n) \equiv \phi_D(\mathbf{z}; \boldsymbol{\beta})$$

exists and is finite. Moreover, if D' is another finite domain and $f : D \rightarrow D'$ is a conformal map such that $z'_1 = f(z_1), \dots, z'_n = f(z_n)$, then

$$\phi_{D'}(\mathbf{z}'; \boldsymbol{\beta}) = \prod_{j=1}^n |f'(z_j)|^{-2\Delta(\beta_j)} \phi_D(\mathbf{z}; \boldsymbol{\beta}),$$

where the $\Delta(\beta)$ is defined below. This is the behavior expected for a conformal primary operator.

- For both versions in infinite volume, correlators of n exponential operators

$$\langle \prod_j V_{\beta_j}(z_j) \rangle_{\delta} = \langle e^{i \sum_j \beta_j N(z_j)} \rangle_{\delta} \quad (2.2)$$

vanish. However, in the case of the layering model, one can remove the short-time cutoff and still obtain a nontrivial limit by imposing the following “charge conservation” condition, satisfied mod 2π ,

$$\sum_j \beta_j = 2\pi k, \quad k \in \mathbb{Z}. \quad (2.3)$$

- The correlators (2.2) of layering operators in the plane are finite and non-zero when (2.3) is satisfied, so long as the loop soup is cut off at short times δ (no long-time cutoff is necessary).

- In the case of 2 points, assuming (2.3), the $\delta \rightarrow 0$ limit of the renormalized layering operator correlators in the plane (2.2) can be explicitly computed up to an overall multiplicative constant. The result is

$$\phi_{\mathbb{C}}(z_1, z_2; \beta_1, \beta_2) = C_2 \left| \left(\frac{1}{z_1 - z_2} \right)^{\Delta(\beta_1) + \Delta(\beta_2)} \right|^2,$$

where C_2 is a constant (Theorem 3.1).

- The $\delta \rightarrow 0$ limit of the renormalized 3-point function for the layering model in the plane, assuming (2.3), is

$$\begin{aligned} \phi_{\mathbb{C}}(z_1, z_2, z_3; \beta_1, \beta_2, \beta_3) = \\ C_3 \left| \left(\frac{1}{|z_1 - z_2|} \right)^{\Delta(\beta_1) + \Delta(\beta_2) - \Delta(\beta_3)} \left(\frac{1}{|z_1 - z_3|} \right)^{\Delta(\beta_1) + \Delta(\beta_3) - \Delta(\beta_2)} \left(\frac{1}{|z_2 - z_3|} \right)^{\Delta(\beta_2) + \Delta(\beta_3) - \Delta(\beta_1)} \right|^2, \end{aligned}$$

where C_3 is a constant (Theorem 4.5).

- The conformal dimensions $\Delta(\beta)$ differ for the two types of operators. For the layering number,

$$\Delta_{\ell}(\beta) = \frac{\lambda}{10}(1 - \cos \beta).$$

For the winding number,

$$\Delta_w(\beta) = \lambda\beta(2\pi - \beta)/8\pi^2,$$

where this formula is valid for $0 \leq \beta < 2\pi$, and $\Delta_w(\beta)$ is periodic under $\beta \rightarrow \beta + 2\pi$.

Open questions: Our results leave a number of questions to be answered.

- We have not determined whether the correlators of the winding operators converge in the infinite plane. The missing piece is the winding number analog of the fact that the BLS is thin [17] (see Sec. 3.4). It is plausible that these correlators do not in fact converge without an explicit long-distance regulator even when charge conservation (2.3) is satisfied.
- We have not established that the n -point correlators in either model are analytic functions of the z_i , although we expect this is the case.
- We expect that these correlators define some kind of conformal field theory. What conformal field theory is it?

- The disk model of [4] could be thought of as the late-time distribution of bubbles produced by a first-order phase transition in de Sitter spacetime. Is there an analogous physical interpretation of the BLS?

3 Correlators of the layering and winding operators

3.1 Correlators of layering operator

As discussed in Sec. 1.3, Model 1 is defined by randomly assigning a Boolean variable to each loop in the Brownian loop soup. Alternatively, one can think of this as two independent Brownian loop soups, each with a Poisson distribution $P_{\lambda_{+(-)}, \mu^{loop}}$ with intensity measure $\lambda_{+(-)}\mu^{loop}$, where we take $\lambda_+ = \lambda_- = \lambda/2$. (This follows from the fact that the collection of all loops from a BLS of intensity λ_+ and an independent one of intensity λ_- is distributed like a BLS with intensity $\lambda_+ + \lambda_-$.)

Denote by $N_{+(-)}(z)$ the number of loops γ in the first (respectively, second) class such that the point $z \in \mathbb{C}$ is separated from infinity by the image of γ in \mathbb{C} . If $\bar{\gamma}$ is the “filled-in” loop γ , then this condition becomes $z \in \bar{\gamma}$, or z is covered by γ . We are interested in the layering field N_ℓ , with $N_\ell(z) = N_+(z) - N_-(z)$. This is purely formal as both $N_{+(-)}(z)$ are infinite with probability one for any z . They are infinite for two reasons: both because there are infinitely many large loops surrounding z (infrared, or IR, divergence), and because there are infinitely many small loops around z (ultraviolet, or UV, divergence).

We will consider correlators of the exponential operator $V_\beta = e^{i\beta N_\ell(z)}$, and show that there are choices of β that remove the IR divergence and a normalization which removes the UV divergence. Specifically, we are interested in the correlators $V_\beta(z_1, \dots, z_n) \equiv \prod_{j=1}^n V_{\beta_j}(z_j) = e^{i \sum_{j=1}^n \beta_j N_\ell(z_j)}$ and their moments

$$\langle V_\beta(z_1, \dots, z_n) \rangle \equiv \mathbb{E}_\lambda(V_\beta(z_1, \dots, z_n))$$

where $z_j \in \mathbb{C}$, $\beta = (\beta_1, \dots, \beta_n) \in \mathbb{R}^n$, and the expected value \mathbb{E}_λ is taken with respect to the distribution $P_{\lambda, \mu^{loop}} \otimes P_{1/2}$, where $P_{\lambda, \mu^{loop}}$ is the Poisson distribution with intensity measure $\lambda\mu^{loop}$ and $P_{1/2}$ is the Bernoulli distribution with parameter 1/2 (remember that each loop belongs to one of two classes with equal probability) or, equivalently, with respect to two independent copies of the Brownian Loop Soup with equal intensities $\lambda/2$.

As the field N_ℓ has both IR and UV divergences, we get a meaningful definition by introducing cutoffs which restrict the loops to have diameter³ within some δ and

³The diameter of a Brownian loop is the largest distance between two points on its outer boundary.

$R \in \mathbb{R}^+$, $\delta < R$: let $\mu_{\delta,R}^{loop}(\cdot) = \mu^{loop}(\cdot \cap \{\gamma : \delta \leq \text{diam}(\gamma) < R\})$ and consider the correlators

$$\langle V_\beta(z_1, \dots, z_n) \rangle_{\delta,R} \equiv \mathbb{E}_{\lambda,\delta,R} \left(e^{i \sum_{j=1}^n \beta_j N_\ell(z_j)} \right),$$

where the expectation $\mathbb{E}_{\lambda,\delta,R}$ is as the \mathbb{E}_λ above with μ^{loop} replaced by $\mu_{\delta,R}^{loop}$.

3.2 The 1-point function in the layering model

In this section we explicitly compute the 1-point function in the presence of IR and UV cutoffs. Replacing the area of a filled Brownian loop of time length 1 with that of a disk of radius 1, the result reproduces the 1-point function in the disk model [4].

Lemma 3.1. *For all $z \in \mathbb{C}$, we have that*

$$\langle V_\beta(z) \rangle_{\delta,R} = \left(\frac{R}{\delta} \right)^{-\frac{\lambda}{5}(1-\cos \beta)}.$$

Proof. In this and other similar results that follow, we first compute $\langle V_\beta(z) \rangle_{\delta,R} = \mathbb{E}_{\lambda,\delta,R} (e^{i\beta N_\ell(z)})$ in terms of probabilities such as $\alpha_{z,\delta,R} = \mu_{\delta,R}(\gamma : z \in \bar{\gamma})$ (see (3.1)). Such calculations can be performed by using two independent copies of the BLS, with random distributions of loops indicated by M_1 and M_2 respectively; then writing $N_\ell(z) = \int \mathbb{1}_{\bar{\gamma} \ni z} (dM_1 - dM_2)$, where $\mathbb{1}_A$ is the indicator function of A ; next, observing that the properties of the Poisson distribution imply that $\mathbb{E}_{\lambda,\delta,R} (e^{i\beta \int \mathbb{1}_{\bar{\gamma} \ni z} (dM_1 - dM_2)}) = e^{-\int (1 - e^{i\beta \mathbb{1}_{\bar{\gamma} \ni z}}) d(\lambda \mu^{loop})}$; and, finally, computing the integral in this last expression. We now give a detailed proof based on the representation with randomly colored loops.

Recall that with IR and UV cutoffs in place, the field $N_\ell(z)$ can be realized as follows. Let η be a realization of loops, and let $\{X_\gamma\}_{\gamma \in \eta}$ be a collection of independent Bernoulli symmetric random variables taking values in $\{-1, 1\}$. The quantity

$$N_\ell(z) = \sum_{\gamma \in \eta, z \in \bar{\gamma}, \delta \leq \text{diam}(\gamma) < R} X_\gamma \equiv \sum^* X_\gamma$$

is finite $P_{\lambda,\mu^{loop}}$ almost surely, since $\mu^{loop}\{\gamma : z \in \bar{\gamma}, \delta \leq \text{diam}(\gamma) < R\} = \mu_{\delta,R}\{\gamma : z \in \bar{\gamma}\} < \infty$ (see [18]). Now,

$$\begin{aligned} \langle V_\beta(z) \rangle_{\delta,R} &= \mathbb{E}_{\lambda,\delta,R} (e^{i\beta N_\ell(z)}) \\ &= \sum_{k=0}^{\infty} \mathbb{E}_{\lambda,\delta,R} (e^{i\beta N_\ell(z)} | \mathcal{L}_k) P_{\lambda,\mu_{\delta,R}}(\mathcal{L}_k), \end{aligned}$$

where $\mathcal{L}_k = \{\eta : |\{\gamma \in \eta : z \in \bar{\gamma}, \delta \leq \text{diam}(\gamma) < R\}| = k\}$. If X denotes a (± 1) -valued symmetric random variable,

$$\mathbb{E}_{\lambda,\delta,R} (e^{i\beta \sum^* X_\gamma} | \mathcal{L}_k) = (E(e^{i\beta X}))^k = (\cos \beta)^k.$$

Therefore, for $\alpha_{z,\delta,R} = \mu_{\delta,R}(\gamma : z \in \bar{\gamma})$, we have that

$$\langle V_\beta(z) \rangle_{\delta,R} = \sum_{k=0}^{\infty} (\cos \beta)^k \frac{(\lambda \alpha_{z,\delta,R})^k}{k!} e^{-\lambda \alpha_{z,\delta,R}} \quad (3.1)$$

$$= e^{-\lambda \alpha_{z,\delta,R} (1 - \cos \beta)}. \quad (3.2)$$

Moreover, by Lemma 8.1 in Sec. 8,

$$\alpha_{z,\delta,R} = \frac{1}{5} \log \frac{R}{\delta},$$

which implies

$$\langle V_\beta(z) \rangle_{\delta,R} = \left(\frac{R}{\delta} \right)^{-\frac{\lambda}{5}(1 - \cos(\beta))},$$

as claimed. \square

3.3 The winding operator

To define the second model, let $N_w(z)$ denote the total winding number about the point z of all loops in a Brownian Loop Soup; as for the layering operators, this is a formal definition as $N_w(z)$ might have divergences. Consider again the correlators $V_\beta(z_1, \dots, z_n) = e^{i \sum_{j=1}^n \beta_j N_w(z_j)}$ and their moments $\langle V_\beta(z_1, \dots, z_n) \rangle = E(V_\beta(z_1, \dots, z_n))$ where $z_j \in \mathbb{C}$, $\beta = (\beta_1, \dots, \beta_n) \in \mathbb{R}^n$, and the expected value is taken with respect to the BLS distribution. Denoting by $P_{\lambda, \mu^{loop}}$ the Poisson distribution with intensity measure $\lambda \mu^{loop}$, and restricting the loops to have diameter between some δ and $R \in \mathbb{R}^+$, with $\delta < R$, we let $\mu_{\delta,R}^{loop}(\cdot) = \mu^{loop}(\cdot \cap \{\gamma : \delta \leq \text{diam}(\gamma) < R\})$ and consider the correlators

$$\langle V_\beta(z_1, \dots, z_n) \rangle_{\delta,R} = \mathbb{E}_{\lambda, \delta, R} \left(e^{i \sum_{j=1}^n \beta_j N_w(z_j)} \right).$$

We now explicitly compute the 1-point function in the presence of IR and UV cutoffs for the winding model.

Lemma 3.2. *For all $z \in \mathbb{C}$, we have that*

$$\langle V_\beta(z) \rangle_{\delta,R} = \left(\frac{R}{\delta} \right)^{-\lambda \frac{\beta(2\pi - \beta)}{4\pi^2}}, \quad (3.3)$$

where the formula is valid for $\beta \in [0, 2\pi)$, and for $\beta \notin [0, 2\pi)$, in the right hand side, β should be replaced by $(\beta \bmod 2\pi)$.

Proof. For a point z and a loop γ , let $\theta_\gamma(z)$ indicate the winding number of γ around z . Moreover, for $\mathbf{k} \in (\mathbb{N} \cup \{0\})^\mathbb{N}$ let

$$\mathcal{L}_{\mathbf{k}} = \{\eta : |\{\gamma \in \eta : z \in \bar{\gamma}, \delta \leq \text{diam}(\gamma) < R, |\theta_\gamma(z)| = m\}| = k_m \text{ for all } m \in \mathbb{N}\}.$$

If a loop γ has $|\theta_\gamma(z)| = m$, then the winding number $\theta_\gamma(z)$ is $\pm m$ with equal probability under $P_{\lambda, \mu^{loop}}$. Finally, using Lemma 8.2 from Sec. 8, for $m \geq 1$ we have

$$\begin{aligned} \alpha_{z, \delta, R, m} &\equiv \mu_{\delta, R}^{loop}(\gamma : z \in \bar{\gamma}, |\theta_\gamma(z)| = m) \\ &= \mu^{loop}(\gamma : z \in \bar{\gamma}, \delta \leq \text{diam}(\gamma) < R, |\theta_\gamma(z)| = m) \\ &= \frac{1}{\pi^2 m^2} \log \frac{R}{\delta}. \end{aligned}$$

For all $\mathbf{k} \in (\mathbb{N} \cup \{0\})^\mathbb{N}$ we have

$$P_{\lambda, \mu_{\delta, R}}(\mathcal{L}_{\mathbf{k}}) = \prod_{m=1}^{\infty} \frac{(\lambda \alpha_{z, \delta, R, m})^{k_m}}{k_m!} e^{-\lambda \alpha_{z, \delta, R, m}},$$

as for different m 's the sets of loops with those winding numbers are disjoint. Hence with IR and UV cutoffs in place, denoting by $E_{\lambda, \delta, R}$ the expectation with respect to the Poisson distribution $P_{\lambda, \mu_{\delta, R}^{loop}}$ with intensity measure $\mu_{\delta, R}^{loop}$, we have, for all z ,

$$\begin{aligned} \langle V_\beta(z) \rangle_{\delta, R} &= E_{\lambda, \delta, R} (e^{i\beta N_w(z)}) \\ &= \sum_{\mathbf{k} \in (\mathbb{N} \cup \{0\})^\mathbb{N}} E_{\lambda, \delta, R} (e^{i\beta N_w(z)} | \mathcal{L}_{\mathbf{k}}) P_{\lambda, \mu_{\delta, R}}(\mathcal{L}_{\mathbf{k}}) \\ &= \sum_{\mathbf{k} \in (\mathbb{N} \cup \{0\})^\mathbb{N}} \prod_{m=1}^{\infty} (\cos(m\beta))^{k_m} \frac{(\lambda \alpha_{z, \delta, R, m})^{k_m}}{k_m!} e^{-\lambda \alpha_{z, \delta, R, m}} \\ &= \prod_{m=1}^{\infty} e^{-\lambda \alpha_{z, \delta, R, m} (1 - \cos(m\beta))} \\ &= \left(\frac{R}{\delta} \right)^{-\lambda \sum_{m=1}^{\infty} \frac{1}{\pi^2 m^2} (1 - \cos(m\beta))} = \left(\frac{R}{\delta} \right)^{-\lambda \frac{\beta(2\pi - \beta)}{4\pi^2}}, \end{aligned}$$

where the β on the right hand side of the last equality is to be interpreted modulo 2π . \square

This coincides with the result (5.3) computed using physics path integral methods.

3.4 The 2-point function in the layering model

We now analyze the 2-point function when the IR cutoff is removed by the charge conservation condition (2.3).

Theorem 3.1. *If $\beta_1 + \beta_2 = 2k\pi$ with $k \in \mathbb{Z}$, there is a positive constant $C_2 < \infty$ such that, for all $z_1 \neq z_2$,*

$$\lim_{R \rightarrow \infty} \langle V_{\beta_1}(z_1) V_{\beta_2}(z_2) \rangle_{\delta, R} = C_2 \left(\frac{|z_1 - z_2|}{\delta} \right)^{-\frac{\lambda}{5}(2 - \cos \beta_1 - \cos \beta_2)}.$$

As a consequence,

$$\lim_{\delta \rightarrow 0} \lim_{R \rightarrow \infty} \frac{\langle V_{\beta_1}(z_1) V_{\beta_2}(z_2) \rangle_{\delta, R}}{\delta^{\frac{\lambda}{5}(2 - \cos \beta_1 - \cos \beta_2)}} = C_2 |z_1 - z_2|^{-\frac{\lambda}{5}(2 - \cos \beta_1 - \cos \beta_2)}.$$

Proof. Letting $d \equiv |z_1 - z_2|$, for given β_1 and β_2 , and $d \geq \delta$, we have that

$$\begin{aligned} \langle V_{\beta_1}(z_1) V_{\beta_2}(z_2) \rangle_{\delta, R} &= \langle e^{i(\beta_1 N_{\ell}(z_1) + \beta_2 N_{\ell}(z_2))} \rangle_{\delta, R} \\ &= \langle e^{i(\beta_1 + \beta_2) N_{12}} \rangle_{d, R} \langle e^{i\beta_1 N_1} \rangle_{\delta, R} \langle e^{i\beta_2 N_2} \rangle_{\delta, R}, \end{aligned}$$

where N_{12} is the number of loops that cover both z_1 and z_2 , and N_1 (N_2) is the number of loops that cover z_1 but not z_2 (z_2 but not z_1 , resp.). The two-point function factorizes because the sets of loops contributing to N_{12} , N_1 and N_2 are disjoint; the δ is replaced by d in the first factor in the second line because a loop covering both z_1 and z_2 must have diameter at least d .

As in the 1-point function calculation, we can write

$$\langle e^{i(\beta_1 + \beta_2) N_{12}} \rangle_{d, R} = \sum_{n=0}^{\infty} (\cos(\beta_1 + \beta_2))^n P_{\lambda, \mu_{d, R}^{loop}}(N_{12} = n) = e^{-\lambda \alpha_{d, R}(z_1, z_2)(1 - \cos(\beta_1 + \beta_2))},$$

where $\alpha_{d, R}(z_1, z_2) \equiv \mu_{d, R}^{loop}(\gamma : z_1, z_2 \in \bar{\gamma})$. Similarly, if $\alpha_{\delta, R}(z_1, \neg z_2) = \mu_{\delta, R}^{loop}(\gamma : z_1 \in \bar{\gamma}, z_2 \notin \bar{\gamma})$ and $\alpha_{\delta, R}(\neg z_1, z_2)$ is correspondingly defined, then

$$\langle e^{i\beta_1 N_1} \rangle_{\delta, R} = e^{-\lambda \alpha_{\delta, R}(z_1, \neg z_2)(1 - \cos \beta_1)}$$

and

$$\langle e^{i\beta_2 N_2} \rangle_{\delta, R} = e^{-\lambda \alpha_{\delta, R}(\neg z_1, z_2)(1 - \cos \beta_2)}.$$

Combining the three terms we obtain

$$\begin{aligned} &\langle e^{i(\beta_1 N_{\ell}(z_1) + \beta_2 N_{\ell}(z_2))} \rangle_{\delta, R} \\ &= e^{-\lambda \alpha_{d, R}(z_1, z_2)(1 - \cos(\beta_1 + \beta_2))} e^{-\lambda \alpha_{\delta, R}(z_1, \neg z_2)(1 - \cos \beta_1)} e^{-\lambda \alpha_{\delta, R}(\neg z_1, z_2)(1 - \cos \beta_2)}. \end{aligned}$$

It is easy to see that $\lim_{R \rightarrow \infty} \alpha_{d, R}(z_1, z_2) = \infty$. (This follows from the scale invariance of μ^{loop} by considering an increasing – in size – sequence of disjoint, concentric annuli

around z_1 and z_2 that are scaled versions of each other.) Hence, in order to remove the IR cutoff, we must impose (2.3) and set $\beta_1 + \beta_2 = 2k\pi$, so that $1 - \cos(\beta_1 + \beta_2) = 0$.

Assuming that $\beta_1 + \beta_2 = 2k\pi$, we are left with

$$\left\langle e^{i(\beta_1 N_\ell(z_1) + \beta_2 N_\ell(z_2))} \right\rangle_{\delta, R} = e^{-\lambda \alpha_{\delta, R}(z_1, \neg z_2)(1 - \cos \beta_1) - \lambda \alpha_{\delta, R}(\neg z_1, z_2)(1 - \cos \beta_2)}. \quad (3.4)$$

To remove the infrared cutoff, we use the fact that the loop is *thin*: If $z_1 \neq z_2$, $\mu^{loop}(\gamma : z_1 \in \bar{\gamma}, z_2 \notin \bar{\gamma}, \text{diam}(\gamma) \geq \delta) < \infty$ for any $\delta > 0$ (see [17], Lemma 4). By the obvious monotonicity of $\alpha_{\delta, R}(z_1, \neg z_2)$ in R , this implies that

$$\lim_{R \rightarrow \infty} \alpha_{\delta, R}(z_1, \neg z_2) = \mu^{loop}\{\gamma \in \eta : z_1 \in \bar{\gamma}, z_2 \notin \bar{\gamma}, \text{diam}(\gamma) \geq \delta\} \equiv \alpha_\delta(z_1, \neg z_2).$$

By scale, rotation and translation invariance of the Brownian loop measure μ^{loop} , $\alpha_\delta(z_1, \neg z_2)$ can only depend on the ratio $x = d/\delta$, so we can introduce the notation $\alpha(x) \equiv \alpha_\delta(z_1, \neg z_2)$. The function α has the following properties, which are also immediate consequences of the scale, rotation and translation invariance of the Brownian loop measure.

- $\alpha(x) = \alpha_\delta(0, \neg z)$ for any z such that $|z| = d$.
- For $\sigma \geq 1$, if $\delta < d$, letting $\alpha_{\delta, R}(z) \equiv \alpha_{z, \delta, R} = \mu_{\delta, R}(\gamma : z \in \bar{\gamma})$,

$$\begin{aligned} \alpha(\sigma x) &= \alpha_\delta(0, \neg \sigma z) \\ &= \alpha_{\sigma\delta}(0, \neg \sigma z) + \alpha_{\delta, \sigma\delta}(0, \neg \sigma z) \\ &= \alpha(x) + \alpha_{\delta, \sigma\delta}(0) = \alpha(x) + \alpha_{1, \sigma}(0). \end{aligned} \quad (3.5)$$

Now let

$$G(x) \equiv \left\langle e^{i(\beta_1 N_\ell(z_1) + \beta_2 N_\ell(z_2))} \right\rangle_\delta \equiv \lim_{R \rightarrow \infty} \left\langle e^{i(\beta_1 N_\ell(z_1) + \beta_2 N_\ell(z_2))} \right\rangle_{\delta, R};$$

using (3.4) and the definition of the function α , we can write

$$G(x) = e^{-\lambda \alpha(x)(2 - \cos \beta_1 - \cos \beta_2)}.$$

Then, for $\sigma \geq 1$,

$$G(\sigma x) = e^{-\lambda \alpha_{1, \sigma}(0)(2 - \cos \beta_1 - \cos \beta_2)} G(x).$$

Using Lemma 8.1, we have that

$$\alpha_{1, \sigma}(0) = \frac{1}{5} \log \sigma.$$

It then follows that, for $\sigma \geq 1$,

$$G(\sigma x) = \sigma^{-\frac{\lambda}{5}(2 - \cos \beta_1 - \cos \beta_2)} G(x). \quad (3.6)$$

For $0 < \sigma < 1$, (3.5) implies

$$\alpha(\sigma x) = \alpha(x) - \alpha_{1,1/\sigma}(0).$$

But since

$$\alpha_{1,1/\sigma}(0) = -\frac{1}{5} \log \sigma,$$

equation (3.6) is unchanged when $0 < \sigma < 1$.

The fact that (3.6) is valid for all $\sigma > 0$ immediately implies that

$$G(x) = C_2 x^{-\frac{\lambda}{5}(2 - \cos \beta_1 - \cos \beta_2)}$$

for some constant $C_2 > 0$. \square

4 Conformal covariance of the n -point functions

We now analyze the n -point functions for general $n \geq 1$ and their conformal invariance properties. In bounded domains $D \subset \mathbb{C}$, we show, for both models, how to remove the UV cutoff $\delta > 0$ by dividing by $\delta^{2\sum_{j=1}^n \Delta_j}$, with the appropriate Δ_j 's. We also show that this procedure leads to conformally covariant functions of the domain D . The scaling with δ originates from the fact that loops with diameter less than δ can only wind around a single point in the limit $\delta \rightarrow 0$, and so for these small loops the n -point function reduces to the product of 1-point functions.

In Section 4.3, we deal with the layering model in the full plane, \mathbb{C} , and show that, together with the UV cutoff $\delta > 0$, we can also remove the IR cutoff $R < \infty$, provided we impose the condition $\sum_{j=1}^n \beta_j \in 2\pi\mathbb{Z}$ (cf. (2.3)). We refer to this condition as “charge conservation” because—apart from the periodicity—it is reminiscent of momentum or charge conservation for the vertex operators of the free boson.

Just as for the two-point function, in the layering model the IR convergence (given “charge conservation”) is due to the finiteness of the total mass of the loops which cover some points but not others; this is basically the property that the soup of outer boundaries of a Brownian loop soup is *thin* in the language of Nacu and Werner [17]. We did not prove the analogous finiteness for the winding model, so in that case we are not able to prove that the IR cutoff can be removed.

4.1 The layering model in finite domains

In the theorem below, we let $\left\langle \prod_{j=1}^n V_{\beta_j}(z_j) \right\rangle_{\delta, D} = \mathbb{E}_{\lambda, \delta, D} \left(\prod_{j=1}^n e^{i\beta_j N_\ell(z_j)} \right)$ denote the expectation of the product $\prod_{j=1}^n e^{i\beta_j N_\ell(z_j)}$ with respect to a loop soup in D with intensity $\lambda > 0$ containing only loops of diameter at least $\delta > 0$, that is, with respect to the distribution $P_{\lambda, \mu_{\delta, D}^{loop}} \otimes P_{1/2}$, where $P_{\lambda, \mu_{\delta, D}^{loop}}$ is the Poisson distribution with intensity measure $\mu_{\delta, D}^{loop} = \mu_D^{loop} \mathbb{1}_{\{\text{diam}(\gamma) \geq \delta\}} = \mu^{loop} \mathbb{1}_{\{\gamma \subset D, \text{diam}(\gamma) \geq \delta\}}$ and $P_{1/2}$ is the Bernoulli distribution with parameter $1/2$ (remember that each loop belongs to one of two classes with equal probability).

Theorem 4.1. *If $n \in \mathbb{N}$, $D \subset \mathbb{C}$ is bounded and $\boldsymbol{\beta} = (\beta_1, \dots, \beta_n)$, then*

$$\lim_{\delta \rightarrow 0} \frac{\left\langle \prod_{j=1}^n V_{\beta_j}(z_j) \right\rangle_{\delta, D}}{\delta^{\lambda \frac{1}{5} \sum_{j=1}^n (1 - \cos \beta_j)}} \equiv \phi_D(z_1, \dots, z_n; \boldsymbol{\beta})$$

exists and is finite and real. Moreover, if D' is another bounded subset of \mathbb{C} and $f : D \rightarrow D'$ is a conformal map such that $z'_1 = f(z_1), \dots, z'_n = f(z_n)$, then

$$\phi_{D'}(z'_1, \dots, z'_n; \boldsymbol{\beta}) = \prod_{j=1}^n |f'(z_j)|^{-\frac{\lambda}{5}(1 - \cos \beta_j)} \phi_D(z_1, \dots, z_n; \boldsymbol{\beta}).$$

The proof of the theorem will make use of the following lemma, where $B_\delta(z)$ denotes the disc of radius δ centered at z , $\bar{\gamma}$ denotes the complement of the unique unbounded component of $\mathbb{C} \setminus \gamma$, and where $o(1)$ denotes a quantity “smaller than $O(1)$ ”, i.e. that tends to zero as $\delta \rightarrow 0$.

Lemma 4.2. *Let $D, D' \subset \mathbb{C}$ and let $f : D \rightarrow D'$ be a conformal map. For $n \geq 1$, assume that $z_1, \dots, z_n \in D$ are distinct and that $z'_1 = f(z_1), \dots, z'_n = f(z_n)$, and let $s_j = |f'(z_j)|$ for $j = 1, \dots, n$. Then we have that, for each $j = 1, \dots, n$,*

$$\begin{aligned} & \mu_{D'}^{loop}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset f(B_\delta(z_j)), z'_k \notin \bar{\gamma} \forall k \neq j) \\ & - \mu_{D'}^{loop}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset B_{s_j \delta}(z'_j), z'_k \notin \bar{\gamma} \forall k \neq j) = o(1) \text{ as } \delta \rightarrow 0. \end{aligned}$$

Proof. Let $B_{in}(z'_j)$ denote the largest (open) disc centered at z'_j contained inside $f(B_\delta(z_j)) \cap B_{s_j \delta}(z'_j)$, and $B_{out}(z'_j)$ denote the smallest disc centered at z'_j containing

$f(B_\delta(z_j)) \cup B_{s_j\delta}(z'_j)$. A moment of thought reveals that, for δ sufficiently small,

$$\begin{aligned}
& |\mu_{D'}^{loop}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset f(B_\delta(z_j)), z'_k \notin \bar{\gamma} \forall k \neq j) \\
& - \mu_{D'}^{loop}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset B_{s_j\delta}(z'_j), z'_k \notin \bar{\gamma} \forall k \neq j)| \\
& = \mu_{D'}^{loop}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset f(B_\delta(z_j)) \cap s_j B_\delta(z'_j), \bar{\gamma} \subset f(B_\delta(z_j)) \cup s_j B_\delta(z'_j), z'_k \notin \bar{\gamma} \forall k \neq j) \\
& \leq \mu_{D'}^{loop}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset B_{in}(z'_j), \bar{\gamma} \subset B_{out}(z'_j)) \\
& = c \log \frac{\text{diam}(B_{out}(z'_j))}{\text{diam}(B_{in}(z'_j))}, \tag{4.1}
\end{aligned}$$

where $c < \infty$ is a positive constant and the last equality follows from Proposition 3 of [18]. Note that, when $D' = \mathbb{C}$, the quantities above involving $\mu_{\mathbb{C}}^{loop}$ are bounded because of the fact that the Brownian loop soup is *thin* [17].

Since f is analytic, for every $w \in \partial B_\delta(z_j)$, we have that

$$|f(w) - z'_j| = s_j\delta + O(\delta^2),$$

which implies that

$$\lim_{\delta \rightarrow 0} \frac{\text{diam}(B_{out}(z'_j))}{\text{diam}(B_{in}(z'_j))} = 1.$$

In view of (4.1), this concludes the proof of the lemma. \square

Proof of Theorem 4.1. We first show that the limit is finite. Using the notation of the previous section, we let η denote a realization of loops and $\{X_\gamma\}_{\gamma \in \eta}$ a collection of independent Bernoulli symmetric random variables taking values in $\{-1, 1\}$. Moreover, let $[n] \equiv \{1, \dots, n\}$, let \mathcal{K} denote the space of assignments of a nonnegative integer to each nonempty subset S of $\{z_1, \dots, z_n\}$, and for $S \subset \{z_1, \dots, z_n\}$, let $I_S \subset [n]$ be the set of indices such that $k \in I_S$ if and only if $z_k \in S$. We have that

$$\begin{aligned}
\left\langle \prod_{j=1}^n V_{\beta_j}(z_j) \right\rangle_{\delta, D} &= \mathbb{E}_{\lambda, \delta, D} \left(e^{i \sum_{j=1}^n \beta_j N_\ell(z_j)} \right) \\
&= \sum_{\mathbf{k} \in \mathcal{K}} \mathbb{E}_{\lambda, \delta, D} \left(e^{i \sum_{j=1}^n \beta_j N_\ell(z_j)} | \mathcal{L}_{\mathbf{k}} \right) P_{\lambda, \mu_{\delta, D}^{loop}}(\mathcal{L}_{\mathbf{k}})
\end{aligned}$$

where $\mathcal{L}_{\mathbf{k}} = \{\eta : \forall S \subset \{z_1, \dots, z_n\}, S \neq \emptyset, |\{\gamma \in \eta : z_j \in \bar{\gamma} \ \forall j \in I_S, z_j \notin \bar{\gamma} \ \forall j \notin I_S\}| = \mathbf{k}(S)\}$. With probability one with respect to $P_{\lambda, \mu_{\delta, D}^{loop}}$, we have that for each $j = 1, \dots, n$,

$$N_\ell(z_j) = \sum_{\gamma: z_j \in \bar{\gamma}, \text{diam}(\gamma) \geq \delta} X_\gamma = \sum_{S \subset \{z_1, \dots, z_n\}: z_j \in S} \sum_{\gamma: S \subset \bar{\gamma}, S^c \subset \bar{\gamma}^c, \text{diam}(\gamma) \geq \delta} X_\gamma.$$

With the notation $\sum^S \equiv \sum_{\gamma: S \subset \bar{\gamma}, S^c \subset \bar{\gamma}^c, \text{diam}(\gamma) \geq \delta}$, and letting X denote a (± 1) -value symmetric random variable, we have that

$$\begin{aligned}
\mathbb{E}_{\lambda, \delta, R} \left(e^{i \sum_{j=1}^n \beta_j N_\ell(z_j)} | \mathcal{L}_k \right) &= \mathbb{E}_{\lambda, \delta, R} \left(e^{i \sum_{j=1}^n \beta_j \sum_{S \subset \{z_1, \dots, z_n\}: z_j \in S} \sum^S X_\gamma} | \mathcal{L}_k \right) \\
&= \mathbb{E}_{\lambda, \delta, R} \left(e^{i \sum_{S \subset \{z_1, \dots, z_n\}} \sum^S (\sum_{j \in I_S} \beta_j) X_\gamma} | \mathcal{L}_k \right) \\
&= \prod_{S \subset \{z_1, \dots, z_n\}, S \neq \emptyset} \left(E \left(e^{i (\sum_{j \in I_S} \beta_j) X} \right) \right)^{\mathbf{k}(S)} \\
&= \prod_{S \subset \{z_1, \dots, z_n\}, S \neq \emptyset} \left(\cos \left(\sum_{j \in I_S} \beta_j \right) \right)^{\mathbf{k}(S)}.
\end{aligned}$$

Next, given $S \subset \{z_1, \dots, z_n\}$ with $|S| \geq 2$, let $\alpha_D(S) \equiv \mu_D(\gamma : S \subset \bar{\gamma}, S^c \subset \bar{\gamma}^c)$ and $\alpha_{\delta, D}(z_j) \equiv \mu_D(\gamma : \text{diam}(\gamma) \geq \delta, z_j \in \bar{\gamma}, z_k \notin \bar{\gamma} \forall k \neq j)$. Furthermore, let I_S be the set of indices such that $k \in I_S$ if and only if $z_k \in S$. Let $m = \min_{i, j: i \neq j} |z_i - z_j| \wedge \min_i \text{dist}(z_i, \partial D)$ and note that, when $\delta < m$, we can write

$$\begin{aligned}
\left\langle \prod_{j=1}^n V_{\beta_j}(z_j) \right\rangle_{\delta, D} &= \sum_{\mathbf{k} \in \mathcal{K}} \prod_{S \subset \{z_1, \dots, z_n\}, |S| > 1} \left(\cos \left(\sum_{k \in I_S} \beta_k \right) \right)^{\mathbf{k}(S)} \frac{(\lambda \alpha_D(S))^{\mathbf{k}(S)}}{(\mathbf{k}(S))!} e^{-\lambda \alpha_D(S)} \\
&\quad \prod_{j=1}^n (\cos \beta_j)^{\mathbf{k}(z_j)} \frac{(\lambda \alpha_{\delta, D}(z_j))^{\mathbf{k}(z_j)}}{(\mathbf{k}(z_j))!} e^{-\lambda \alpha_{\delta, D}(z_j)} \\
&= \prod_{S \subset \{z_1, \dots, z_n\}, |S| > 1} \exp \left[-\lambda \alpha_D(S) \left(1 - \cos \left(\sum_{k \in I_S} \beta_k \right) \right) \right] \\
&\quad \prod_{j=1}^n \exp [-\lambda \alpha_{\delta, D}(z_j) (1 - \cos \beta_j)]. \tag{4.2}
\end{aligned}$$

For every $j = 1, \dots, n$, using Lemma 8.1, we have that

$$\begin{aligned}
\alpha_{\delta, D}(z_j) &= \mu_D^{\text{loop}}(\gamma : \text{diam}(\gamma) \geq \delta, z_j \in \bar{\gamma}, z_k \notin \bar{\gamma} \forall k \neq j) \\
&= \mu_D^{\text{loop}}(\gamma : m > \text{diam}(\gamma) \geq \delta, z_j \in \bar{\gamma}) \\
&\quad + \mu_D^{\text{loop}}(\gamma : \text{diam}(\gamma) \geq m, z_j \in \bar{\gamma}, z_k \notin \bar{\gamma} \forall k \neq j) \\
&= \frac{1}{5} \log \frac{m}{\delta} + \alpha_{m, D}(z_j).
\end{aligned}$$

Therefore, we obtain

$$\begin{aligned}
\lim_{\delta \rightarrow 0} \frac{\left\langle \prod_{j=1}^n V_{\beta_j}(z_j) \right\rangle_{\delta, D}}{\delta^{\frac{\lambda}{5} \sum_{j=1}^n (1 - \cos \beta_j)}} &= \prod_{S \subset \{z_1, \dots, z_n\}, |S| > 1} \exp \left[-\lambda \alpha_D(S) \left(1 - \cos \left(\sum_{k \in I_S} \beta_k \right) \right) \right] \\
&\quad m^{-\frac{\lambda}{5} \sum_{j=1}^n (1 - \cos \beta_j)} e^{-\sum_{j=1}^n \lambda \alpha_{m, D}(z_j) (1 - \cos \beta_j)} \\
&= m^{-\frac{\lambda}{5} \sum_{j=1}^n (1 - \cos \beta_j)} \exp \left[-\lambda \sum_{j=1}^n \alpha_{m, D}(z_j) (1 - \cos \beta_j) \right] \\
&\quad \exp \left[-\lambda \sum_{S \subset \{z_1, \dots, z_n\}, |S| > 1} \alpha_D(S) \left(1 - \cos \left(\sum_{k \in I_S} \beta_k \right) \right) \right] \\
&\equiv \phi_D(z_1, \dots, z_n; \boldsymbol{\beta}).
\end{aligned}$$

This concludes the first part of the proof.

To prove the second part of the theorem, using (4.2), we write

$$\begin{aligned}
\left\langle \prod_{j=1}^n V_{\beta_j}(z_j) \right\rangle_{\delta, D} &= \exp \left[-\lambda \sum_{S \subset \{z_1, \dots, z_n\}, |S| \geq 2} \alpha_D(S) \left(1 - \cos \left(\sum_{k \in I_S} \beta_k \right) \right) \right] \\
&\quad \prod_{j=1}^n \exp [-\lambda \alpha_{\delta, D}(z_j) (1 - \cos \beta_j)]. \tag{4.3}
\end{aligned}$$

For each $S \subset \{z_1, \dots, z_n\}$ with $|S| \geq 2$, $\alpha_D(S)$ is invariant under conformal transformations, that is, if $f : D \rightarrow D'$ is a conformal map from D to another bounded domain D' , and $S' = \{z'_1, \dots, z'_n\}$, where $z'_1 = f(z_1), \dots, z'_n = f(z_n)$, then $\alpha_{D'}(S') = \alpha_D(S)$. Therefore, the first exponential term in (4.3) is also invariant under conformal transformations. This implies that, for δ sufficiently small,

$$\frac{\left\langle \prod_{j=1}^n V_{\beta_j}(z_j) \right\rangle_{\delta, D}}{\left\langle \prod_{j=1}^n V_{\beta_j}(z'_j) \right\rangle_{\delta, D'}} = \prod_{j=1}^n \exp \left\{ -\lambda [\alpha_{\delta, D}(z_j) - \alpha_{\delta, D'}(z'_j)] (1 - \cos \beta_j) \right\}.$$

Writing

$$\begin{aligned}
\alpha_{\delta, D}(z_j) &= \mu_D^{\text{loop}}(\gamma : \text{diam}(\gamma) \geq \delta, z_j \in \bar{\gamma}, \bar{\gamma} \subset B_\delta(z_j)) \\
&\quad + \mu_D^{\text{loop}}(\gamma : z_j \in \bar{\gamma}, \bar{\gamma} \not\subset B_\delta(z_j), z_k \notin \bar{\gamma} \forall k \neq j)
\end{aligned}$$

and noticing that $\mu_D^{\text{loop}}(\gamma : \text{diam}(\gamma) \geq \delta, z_j \in \bar{\gamma}, \bar{\gamma} \subset B_\delta(z_j)) = \mu_{D'}^{\text{loop}}(\gamma : \text{diam}(\gamma) \geq \delta, z'_j \in \bar{\gamma}, \bar{\gamma} \subset B_\delta(z'_j))$ (where we have assumed, without loss of generality, that δ is so

small that $B_\delta(z_j) \subset D$ and $B_\delta(z'_j) \subset D'$, we have that

$$\begin{aligned} \alpha_{\delta,D}(z_j) - \alpha_{\delta,D'}(z'_j) &= \mu_D^{\text{loop}}(\gamma : z_j \in \bar{\gamma}, \bar{\gamma} \not\subset B_\delta(z_j), z_k \notin \bar{\gamma} \forall k \neq j) \\ &\quad - \mu_{D'}^{\text{loop}}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset B_\delta(z'_j), z'_k \notin \bar{\gamma} \forall k \neq j). \end{aligned}$$

To evaluate this difference, using conformal invariance, we write

$$\begin{aligned} &\mu_D^{\text{loop}}(\gamma : z_j \in \bar{\gamma}, \bar{\gamma} \not\subset B_\delta(z_j), z_k \notin \bar{\gamma} \forall k \neq j) \\ &= \mu_{D'}^{\text{loop}}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset f(B_\delta(z_j)), z'_k \notin \bar{\gamma} \forall k \neq j). \end{aligned}$$

Letting $s_j = |f'(z_j)|$ and using (8.1) and Lemma 8.1 from the appendix, we can write

$$\begin{aligned} &\alpha_{\delta,D}(z_j) - \alpha_{\delta,D'}(z'_j) \\ &= \mu_{D'}^{\text{loop}}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset f(B_\delta(z_j)), z'_k \notin \bar{\gamma} \forall k \neq j) - \mu_{D'}^{\text{loop}}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset B_\delta(z'_j), z'_k \notin \bar{\gamma} \forall k \neq j) \\ &= \mu_{D'}^{\text{loop}}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset f(B_\delta(z_j)), z'_k \notin \bar{\gamma} \forall k \neq j) - \mu_{D'}^{\text{loop}}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset B_{s_j\delta}(z'_j), z'_k \notin \bar{\gamma} \forall k \neq j) \\ &\quad - [\mu_{D'}^{\text{loop}}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset B_\delta(z'_j), z'_k \notin \bar{\gamma} \forall k \neq j) - \mu_{D'}^{\text{loop}}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset B_{s_j\delta}(z'_j), z'_k \notin \bar{\gamma} \forall k \neq j)] \\ &= \mu_{D'}^{\text{loop}}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset f(B_\delta(z_j)), z'_k \notin \bar{\gamma} \forall k \neq j) - \mu_{D'}^{\text{loop}}(\gamma : z'_j \in \bar{\gamma}, \bar{\gamma} \not\subset B_{s_j\delta}(z'_j), z'_k \notin \bar{\gamma} \forall k \neq j) \\ &\quad - \frac{1}{5} \log s_j. \end{aligned}$$

Using Lemma 4.2, we obtain $\alpha_{\delta,D}(z_j) - \alpha_{\delta,D'}(z'_j) = -\frac{1}{5} \log |f'(z_j)| + o(1)$ as $\delta \rightarrow 0$, which gives

$$\frac{\langle V_\beta(z_1, \dots, z_n) \rangle_{\delta,D}}{\langle V_\beta(z'_1, \dots, z'_n) \rangle_{\delta,D'}} = e^{-o(1)} \prod_{j=1}^n |f'(z_j)|^{\frac{\lambda}{5}(1-\cos \beta_j)} \text{ as } \delta \rightarrow 0.$$

Letting $\delta \rightarrow 0$ concludes the proof. \square

4.2 The winding model in finite domains

As above, let $N_w(z)$ denote the total number of windings of all loops of a given soup around $z \in \mathbb{C}$. We have the following theorem.

Theorem 4.3. *If $n \in \mathbb{N}$, $D \subset \mathbb{C}$ is bounded and $\boldsymbol{\beta} = (\beta_1, \dots, \beta_n)$, then*

$$\lim_{\delta \rightarrow 0} \frac{\langle e^{i\beta_1 N_w(z_1)} \dots e^{i\beta_n N_w(z_n)} \rangle_{\delta,D}}{\delta^{\frac{\lambda}{4\pi^2} \sum_{j=1}^n \beta_j (2\pi - \beta_j)}} \equiv \psi_D(z_1, \dots, z_n; \boldsymbol{\beta})$$

exists and is finite and real. Moreover, if D' is another bounded subset of \mathbb{C} and $f : D \rightarrow D'$ is a conformal map such that $z'_1 = f(z_1), \dots, z'_n = f(z_n)$, then

$$\psi_{D'}(z'_1, \dots, z'_n; \boldsymbol{\beta}) = \prod_{j=1}^n |f'(z_j)|^{-\lambda \frac{\beta_j (2\pi - \beta_j)}{4\pi^2}} \psi_D(z_1, \dots, z_n; \boldsymbol{\beta}),$$

where, in the exponent, the β_j 's are to be interpreted modulo 2π .

Proof. The proof is analogous to that of Theorem 4.1. Let $\alpha_D(S; k_{i_1}, \dots, k_{i_l}) := \mu_D(\gamma : \theta_\gamma(z_{i_j}) = k_{i_j} \text{ for each } z_{i_j} \in S \text{ and } S^c \subset \bar{\gamma}^c)$, and $\alpha_{\delta,D}(z_j; k) := \mu_D(\gamma : z_j \in \bar{\gamma}, \theta_\gamma(z_j) = k, z_k \notin \bar{\gamma} \forall k \neq j)$. With this notation we can write

$$\begin{aligned} & \langle e^{\beta_1 N_w(z_1)} \dots e^{\beta_n N_w(z_n)} \rangle_{\delta,D} \\ &= \exp \left[-\lambda \sum_{l=2}^n \sum_{\substack{S \subset \{z_1, \dots, z_n\} \\ |S|=l}} \sum_{k_{i_1}, \dots, k_{i_l} = -\infty}^{\infty} \alpha_D(S; k_{i_1}, \dots, k_{i_l}) (1 - \cos(k_{i_1} \beta_{i_1} + \dots + k_{i_l} \beta_{i_l})) \right] \\ & \quad \prod_{j=1}^n \exp \left[-\lambda \sum_{k=-\infty}^{\infty} \alpha_{\delta,D}(z_j; k) (1 - \cos(k \beta_j)) \right], \end{aligned}$$

For each $S \subset \{z_1, \dots, z_n\}$ with $|S| = l \geq 2$, $\alpha_D(S; k_{i_1}, \dots, k_{i_l})$ is invariant under conformal transformations; therefore,

$$\frac{\langle e^{i\beta_1 N_w(z_1)} \dots e^{i\beta_n N_w(z_n)} \rangle_{\delta,D}}{\langle e^{i\beta_1 N_w(z'_1)} \dots e^{i\beta_n N_w(z'_n)} \rangle_{\delta,D'}} = \prod_{j=1}^n \exp \left\{ -\lambda \sum_{k=-\infty}^{\infty} [\alpha_{\delta,D}(z_j; k) - \alpha_{\delta,D'}(z'_j; k)] (1 - \cos(k \beta_j)) \right\}.$$

Proceeding as in the proof of Theorem 4.1, but using Lemma 8.2 instead of Lemma 8.1, gives

$$\alpha_{\delta,D}(z_j; k) - \alpha_{\delta,D'}(z'_j; k) = -c_k \log |f'(z_j)| + o(1) \text{ as } \delta \rightarrow 0,$$

where $c_k = \frac{1}{2\pi^2 k^2}$ for $k \in \mathbb{Z} \setminus \{0\}$ and $c_0 = 1/30$.

This, together with the observation, already used at the end of Section 3.3, that $\sum_{k=-\infty}^{\infty} c_k (1 - \cos(k\beta)) = \frac{\beta(2\pi - \beta)}{4\pi^2}$ (where, on the right hand side, β should be interpreted modulo 2π), readily implies the statement of the theorem. \square

4.3 The layering model in the plane

Recall that $\left\langle \prod_{j=1}^n V_{\beta_j}(z_j) \right\rangle_{\delta,R}$ denotes the expectation of the product $\prod_{j=1}^n e^{i\beta_j N_\ell(z_j)}$ with respect to a loop soup in \mathbb{C} with intensity $\lambda > 0$ containing only loops γ of diameter $0 < \delta \leq \text{diam}(\gamma) < R < \infty$.

Theorem 4.4. *If $n \in \mathbb{N}$ and $\boldsymbol{\beta} = (\beta_1, \dots, \beta_n)$ with $|\boldsymbol{\beta}| = \sum_{j=1}^n \beta_j \in 2\pi\mathbb{Z}$, then*

$$\lim_{\delta \rightarrow 0, R \rightarrow \infty} \frac{\left\langle \prod_{j=1}^n V_{\beta_j}(z_j) \right\rangle_{\delta,R}}{\delta^{\frac{\lambda}{5} \sum_{j=1}^n (1 - \cos \beta_j)}} \equiv \phi_{\mathbb{C}}(z_1, \dots, z_n; \boldsymbol{\beta})$$

exists and is finite and real. Moreover, if $f : \mathbb{C} \rightarrow \mathbb{C}$ is a conformal map such that $z'_1 = f(z_1), \dots, z'_n = f(z_n)$, then

$$\phi_{\mathbb{C}}(z'_1, \dots, z'_n; \boldsymbol{\beta}) = \prod_{j=1}^n |f'(z_j)|^{-\frac{\lambda}{5} (1 - \cos \beta_j)} \phi_{\mathbb{C}}(z_1, \dots, z_n; \boldsymbol{\beta}).$$

Proof sketch. The beginning of the proof proceeds like that of Theorem 4.1 until equation (4.2), leading to the following equation:

$$\left\langle \prod_{j=1}^n V_{\beta_j}(z_j) \right\rangle_{\delta, R} = \prod_{S \subset \{z_1, \dots, z_n\}, 1 < |S| < n} \exp \left[-\lambda \alpha_R(S) \left(1 - \cos \left(\sum_{k \in I_S} \beta_k \right) \right) \right] \prod_{j=1}^n \exp [-\lambda \alpha_{\delta, R}(z_j) (1 - \cos \beta_j)],$$

where $\alpha_R(S) \equiv \mu_{\mathbb{C}}(\gamma : S \subset \bar{\gamma}, S^c \subset \bar{\gamma}^c, \text{diam}(\gamma) < R)$, for $S \subset \{z_1, \dots, z_n\}$ with $2 \leq |S| < n$, and $\alpha_{\delta, R}(z_j) \equiv \mu_{\mathbb{C}}(\gamma : \delta \leq \text{diam}(\gamma) < R, z_j \in \bar{\gamma}, z_k \notin \bar{\gamma} \forall k \neq j)$, and where I_S denotes the set of indices such that $k \in I_S$ if and only if $z_k \in S$.

Note, in the equation above, the condition $|S| < n$ in the first product on the right hand side; this condition comes from the fact that the term $-\lambda \alpha_R(S)$ with $S = \{z_1, \dots, z_n\}$ is multiplied by $1 - \cos(\sum_{k=1}^n \beta_k) = 0$, where we have used the ‘‘charge conservation’’ condition $|\beta| = \sum_{j=1}^n \beta_j \in 2\pi\mathbb{Z}$.

For every $j = 1, \dots, n$, using Lemma 8.1, we have that

$$\begin{aligned} \alpha_{\delta, R}(z_j) &= \mu_{\mathbb{C}}(\gamma : \delta \leq \text{diam}(\gamma) < R, z_j \in \bar{\gamma}, z_k \notin \bar{\gamma} \forall k \neq j) \\ &= \mu_{\mathbb{C}}(\gamma : m > \text{diam}(\gamma) \geq \delta, z_j \in \bar{\gamma}) \\ &\quad + \mu_D(\gamma : m \leq \text{diam}(\gamma) < R, z_j \in \bar{\gamma}, z_k \notin \bar{\gamma} \forall k \neq j) \\ &= \frac{1}{5} \log \frac{m}{\delta} + \alpha_{m, R}(z_j). \end{aligned}$$

Now note that monotonicity and the fact that the Brownian loop soup is *thin* [17] imply that $\alpha_{m, \mathbb{C}}(z_j) \equiv \lim_{R \rightarrow \infty} \alpha_{m, R}(z_j)$ and $\alpha_{\mathbb{C}}(S) \equiv \lim_{R \rightarrow \infty} \alpha_R(S)$, for $S \subset \{z_1, \dots, z_n\}$ with $2 \leq |S| < n$, exist and are bounded. After letting $R \rightarrow \infty$, the proof proceeds like that of Theorem 4.1, with $D = D' = \mathbb{C}$. \square

We have already seen the behavior of the 2-point function in the layering model in Section 3.4; the theorem below deals with the 3-point function.

Theorem 4.5. *Let $z_1, z_2, z_3 \in \mathbb{C}$ be three distinct points, then we have that*

$$\phi_{\mathbb{C}}(z_1, z_2, z_3; \beta_1, \beta_2, \beta_3) = C_3 \left| \left(\frac{1}{|z_1 - z_2|} \right)^{\Delta_I(\beta_1) + \Delta_I(\beta_2) - \Delta_I(\beta_3)} \left(\frac{1}{|z_1 - z_3|} \right)^{\Delta_I(\beta_1) + \Delta_I(\beta_3) - \Delta_I(\beta_2)} \left(\frac{1}{|z_2 - z_3|} \right)^{\Delta_I(\beta_2) + \Delta_I(\beta_3) - \Delta_I(\beta_1)} \right|^2$$

for some constant C_3 .

Proof. Theorem 4.4 implies that the 3-point function in the full plane transforms covariantly under conformal maps. This immediately implies the theorem following standard argument (see, *e.g.*, [19]). We briefly sketch those arguments below for the reader's convenience.

Scale invariance, rotation invariance, and translation invariance immediately imply that there are constants C_{abc} such that

$$\phi_{\mathbb{C}}(z_1, z_2, z_3; \beta_1, \beta_2, \beta_3) = \sum C_{abc} z_{12}^{-a} z_{13}^{-b} z_{23}^{-c}, \quad (4.4)$$

where $z_{ij} = |z_i - z_j|$ and the sum is over all triplets $a, b, c \geq 0$ satisfying $a + b + c = 2(\Delta_l(\beta_1) + \Delta_l(\beta_2) + \Delta_l(\beta_3))$ (the constraint on the exponents a, b, c follows from Theorem 4.4 applied to scale transformations).

Now let f be a conformal transformation from \mathbb{C} to \mathbb{C} ; f is then a Möbius transformation and has the form $f(z) = \frac{Az+B}{Cz+D}$, with $f'(z) = \frac{AD-BC}{(Cz+D)^2}$. Letting $\gamma_j \equiv |f'(z_j)|^{-1}$, if $\tilde{z} = f(z)$, it is easy to check that $\tilde{z}_{ij} = \gamma_i^{-1/2} \gamma_j^{-1/2} z_{ij}$. Using this fact and Theorem 4.4, we have that

$$\begin{aligned} \phi_{\mathbb{C}}(\tilde{z}_1, \tilde{z}_2, \tilde{z}_3; \beta_1, \beta_2, \beta_3) &= \left(\gamma_1^{\Delta_l(\beta_1)} \gamma_2^{\Delta_l(\beta_2)} \gamma_3^{\Delta_l(\beta_3)} \right)^2 \sum C_{abc} z_{12}^{-a} z_{13}^{-b} z_{23}^{-c} \\ &= \left(\gamma_1^{\Delta_l(\beta_1)} \gamma_2^{\Delta_l(\beta_2)} \gamma_3^{\Delta_l(\beta_3)} \right)^2 \sum C_{abc} \frac{\tilde{z}_{12}^{-a} \tilde{z}_{13}^{-b} \tilde{z}_{23}^{-c}}{\gamma_1^{a/2+b/2} \gamma_2^{a/2+c/2} \gamma_3^{b/2+c/2}}. \end{aligned}$$

For this last expression to be of the correct form (4.4), the γ 's need to cancel; this immediately leads to the relations $a = 2(\Delta_l(\beta_2) + \Delta_l(\beta_3) - \Delta_l(\beta_1))$, $b = 2(\Delta_l(\beta_1) + \Delta_l(\beta_3) - \Delta_l(\beta_2))$, $c = 2(\Delta_l(\beta_1) + \Delta_l(\beta_2) - \Delta_l(\beta_3))$. \square

5 Dimension of the exponential winding operator

In this section we compute the conformal dimension of the exponential winding number operator $e^{i\beta N_w(z)}$ for the BLS in the plane using non-rigorous path integral methods.

The measure for a single Brownian loop of time length t rooted at x is given by the path integral for a free particle in two Euclidean dimensions, with the path $y(\tau)$ constrained to begin and end at x :

$$\int_{y(0)=x}^{y(t)=x} [d^2y] e^{-\frac{1}{2} \int_0^t d\tau |\dot{y}|^2}, \quad (5.1)$$

where $\dot{y} \equiv \partial y / \partial \tau$, y, x are complex coordinates, and the path integral measure is Brownian normalized so that $\int_x^z [d^2y] e^{-\frac{1}{2} \int_0^t d\tau |\dot{y}|^2} = e^{-|x-z|^2/2t} / (2\pi t)$.

To construct the Brownian loop soup in the plane, we should integrate the root point x over the plane with uniform measure, integrate over t with measure dt/t ,

and sum over n -loop sectors weighted by λ^n and divided by $n!$ since the loops are indistinguishable:

$$\begin{aligned} Z_\lambda &= 1 + \sum_{n=1}^{\infty} \frac{\lambda^n}{n!} \prod_{k=1}^n \int_{\delta^2}^{R^2} \frac{dt_k}{t_k} \int d^2 x_k \int_{y_k(0)=x_k}^{y_k(t_k)=x_k} [d^2 y_k] \exp\left(-\frac{1}{2} \int_0^{t_k} d\tau_k |\dot{y}_k|^2\right) \\ &= \exp\left(\lambda \int_{\delta^2}^{R^2} \frac{dt}{t} \int d^2 x \int_{y(0)=x}^{y(t)=x} [d^2 y] e^{-\frac{1}{2} \int_0^t d\tau |\dot{y}|^2}\right). \end{aligned} \quad (5.2)$$

This result is closely related to (3.1) of [4].

We wish to compute the 1-point function $\langle e^{i\beta N_w(z)} \rangle$, where $\langle \dots \rangle$ represents the average over the distribution defined by (5.2). Since N_w is integer the result should be invariant under $\beta \rightarrow \beta + 2\pi n$. Furthermore for every configuration there is a mirror configuration where $N \rightarrow -N$, so $\langle e^{i\beta N(z)} \rangle$ is real and invariant under $\beta \rightarrow -\beta$. Because we are in the plane and charge conservation cannot be satisfied, the 1-point function will diverge as a power of the ratio of the cutoffs R/δ , but the power tells us the dimension of the operator we wish to compute (*cf.* (3.3)).

To compute $\langle e^{i\beta N_w(z)} \rangle$ we can insert $e^{i\beta N_w(z)}$ in the path integral for each loop in (5.2). By translation invariance in the plane we can set $z = 0$. Then this insertion corresponds to adding a term $i\beta \int_0^t d\tau \dot{\phi}(\tau)/2\pi$ to the single-particle ‘‘lagrangian’’ in the exponential of (5.1), where $\phi(\tau) \bmod 2\pi = \arg(y(\tau))$ is the angular position of the path $y(\tau)$ relative to the origin $z = 0$. Therefore

$$\langle e^{i\beta N_w(z)} \rangle = Z_\lambda^{-1} \exp\left(\lambda \int_{\delta^2}^{R^2} \frac{dt}{t} \int d^2 x \int_{y(0)=x}^{y(t)=x} [d^2 y] e^{\int_0^t d\tau [-\frac{1}{2}(\dot{y})^2 + i\beta\dot{\phi}/2\pi]}\right).$$

Apart from the integral over t , the quantity in the exponential is related to the canonical partition function at inverse temperature t of a charged particle in the field of a single magnetic monopole of charge $\beta/2\pi$ at the origin (n -point functions involving products of exponentials $e^{i\beta_n N_w(z_n)}$ would correspond to multiple magnetic monopoles with charge $\beta_n/2\pi$ at the locations z_n). To compute it, one can quantize the Hamiltonian for a particle in the monopole field, then take the trace in the energy basis [20], or perform the position-space path integral directly and obtain a result in terms of a sum over Bessel functions [21]. The final result is (see for instance [22]):

$$\int d^2 x \int_{y(0)=x}^{y(t)=x} [d^2 y] e^{-\int_0^t d\tau \frac{1}{2}(\dot{y})^2} \left(e^{i\beta \int_0^t d\tau \dot{\phi}/2\pi} - 1 \right) = -\beta(2\pi - \beta)/8\pi^2$$

valid for $0 \leq \beta < 2\pi$ and periodic in $\beta \simeq \beta + 2\pi n$. From this we obtain

$$\langle e^{i\beta N_w(z)} \rangle = \exp\left(-\lambda \frac{\beta(2\pi - \beta)}{8\pi^2} \int_{\delta^2}^{R^2} \frac{dt}{t}\right) = \left(\frac{R}{\delta}\right)^{-\lambda\beta(2\pi - \beta)/4\pi^2} \quad (5.3)$$

in agreement with (3.3).

6 Central charge and relation to the free boson

The partition function (5.2) has a simple interpretation for the case $\lambda = 1/2$. It corresponds precisely to the partition function for a free, massless, real bosonic field in two (Euclidean) dimensions:

$$\ln Z_{\text{boson}} = -\frac{1}{2} \log \det H = \frac{1}{2} \int_0^\infty \frac{dt}{t} \text{Tr} e^{-tH} = \frac{1}{2} \int_0^\infty \frac{dt}{t} \int d^2x \langle x | e^{-tH} | x \rangle,$$

where $H = \square$ is the Laplace operator. Expressing the heat kernel $\langle x | e^{-tH} | y \rangle$ using the path integral (5.1) completes the identification with (5.2).

An analogous relation between the partition function of the discrete Gaussian free field (the lattice analog of the free field) and that of the random walk loop soup (the lattice analog of the BLS) can be derived easily; namely $Z_{DGGFF}^D = \left(\frac{\pi}{2}\right)^{|D|} Z_{1/2}^D$, where Z_{DGGFF}^D is the partition function of the discrete Gaussian free field in D with zero boundary condition, $Z_{1/2}^D$ is the partition function of the random walk loop soup in D with intensity $1/2$, and $|D|$ is the number of vertices in D (see, for example [23], particularly Sections 2.1 and 2.2 and Exercise 2.5). This result may be related to work of Le Jan, who demonstrated that for $\lambda = k/2$, the occupation field for the BLS can be identified with the sum of squares of k copies of a free field [16]. It suggests that for general λ , the BLS might give meaning to the notion of a fractional power of a free field.

Because the massless boson is a CFT with central charge $c = 1$, the central charge of the BLS appears to be $c = 2\lambda$ for the case $\lambda = 1/2$. Given the form of (5.2), one can express the partition function of the BLS for arbitrary λ in terms of that for $\lambda = 1/2$: $Z_\lambda = (Z_{1/2})^{2\lambda}$. The same relation can be rigorously proved to hold for the random walk loop soup mentioned earlier (see equation (2.2) in Section 2.1 of [23]). This leads to the conclusion that the relation $c(\lambda) = 2\lambda$ between the intensity λ and the central charge c of the BLS holds for all λ .

This agrees with the central charge of the SLE corresponding to the ensemble of BLS cluster boundaries. Indeed, conformal field theory considerations lead to the formula

$$c(\kappa) = \frac{(3\kappa - 8)(6 - \kappa)}{2\kappa},$$

but it is also known [2] that, when $8/3 < \kappa \leq 4$, the ensemble of cluster boundaries of a BLS with intensity

$$\lambda = \frac{(3\kappa - 8)(6 - \kappa)}{4\kappa} = \frac{c(\kappa)}{2}$$

is a CLE_κ . (For example, a BLS with $\lambda = 1/4$ gives a CLE_3 , with central charge $1/2$.)

We note that most of the existing literature, including [2], contains an error in the correspondence between κ and the loop soup intensity λ . The error can be traced back to the choice of normalization of the (infinite) Brownian loop measure μ^{loop} , which determines the constant in front of the log in Lemma 8.1 below.⁴ With the normalization used in this paper, which coincides with the one in the original definition of the Brownian loop soup [1], for a given $8/3 < \kappa \leq 4$, the corresponding value of the loop soup intensity λ is half of that given in [2].

7 Conclusions

There is still only a partial understanding of the rich connection between the conformal field theories studied by physicists and conformally invariant stochastic models such as the BLS or SLE. Physicists are often interested in CFTs defined by a Lagrangian, and the correlation functions of primary operators, the spectrum of conformal dimensions, and the central charge are the main objects of study and interest. By contrast, conformal stochastic models are often defined and studied using very different methods and with different goals.

Here we have tried to take a few steps towards strengthening this connection. Using rigorous methods, we defined a set of quantities in the BLS and proved that their expectation values behave like the correlation functions of primary operators. While we have not established it, we expect that these correlation functions may define a conformal field theory. If so, it has several novel features, such as a periodic spectrum of conformal dimensions.

There are many basic questions that remain to be answered. Does this approach to the BLS in fact define a CFT? If so, have we found the complete set of primary operators? What is the stress-energy tensor? Is the theory reflection positive and/or modular invariant? Is it unique in some sense?

An even more ambitious set of questions relates to eternal inflation, albeit in 2+1 dimensions. The model of [4] was proposed as a toy model for the late-time evolution of an eternally inflating spacetime, but the lack of analyticity in the 4-point function of the (analog of the) layering operator derailed it as a putative CFT. Does the BLS solve this problem, and if so, could it help define a CFT dual to eternal inflation? What type of object in 2+1 dimensional de Sitter space produces the BLS as its late-time distribution? Is there a natural generalization to higher dimensions?

⁴We thank Greg Lawler for discussions on this topic.

8 Appendix. The Brownian loop measure: two lemmas

In this appendix we prove two important lemmas which are used several times in the rest of the paper. The lemmas concern the μ^{loop} -measures of certain sets of loops, where, as in the rest of the paper, μ^{loop} is the intensity measure used in the definition of the Brownian loop soup (see equation (1.1)).

Lemma 8.1. *Let $z \in \mathbb{C}$, then*

$$\mu^{loop}(\gamma : z \in \bar{\gamma}, \delta \leq \text{diam}(\gamma) < R) = \frac{1}{5} \log \frac{R}{\delta}.$$

Proof. Since $\mu^{loop}(\gamma : \text{diam}(\gamma) = R) = 0$, we have that

$$\begin{aligned} & \mu^{loop}(\gamma : z \in \bar{\gamma}, \delta \leq \text{diam}(\gamma) < R) - \mu^{loop}(\gamma : z \in \bar{\gamma}, \gamma \not\subset B_{z,\delta}, \gamma \subset B_{z,R}) \\ &= \mu^{loop}(\gamma : z \in \bar{\gamma}, \text{diam}(\gamma) \geq \delta, \gamma \subset B_{z,\delta}) - \mu^{loop}(\gamma : z \in \bar{\gamma}, \text{diam}(\gamma) \geq R, \gamma \subset B_{z,R}), \end{aligned}$$

where $B_{z,a}$ is a disk of radius a around z , and $\gamma \not\subset B_{z,\delta}$ indicates that the image of γ is not fully contained in $B_{z,\delta}$. By the scale invariance of μ^{loop} , the last two terms are identical, so that

$$\begin{aligned} \mu^{loop}(\gamma : z \in \bar{\gamma}, \delta \leq \text{diam}(\gamma) < R) &= \mu^{loop}(\gamma : z \in \bar{\gamma}, \gamma \not\subset B_{z,\delta}, \gamma \subset B_{z,R}) \\ &= c \log \frac{R}{\delta} \end{aligned} \tag{8.1}$$

for some positive constant $c < \infty$, where the last equality follows from Proposition 3 of [18] and the fact that the Brownian loop measure satisfies the conformal restriction property. In order to determine the constant c , we use the fact that, for any $r > 0$,

$$\mu^{loop}(\gamma : z \in \bar{\gamma}, 1 \leq \text{diam}(\gamma) < e^r) = \mu^{loop}(\gamma : z \in \bar{\gamma}, 1 \leq t_\gamma < e^{2r}). \tag{8.2}$$

(Equation (8.2) is essentially a consequence of Brownian scaling and can be proved using standard techniques. The interested reader can consult, for example, Appendix B of [23]).

Following [18], we compute the right hand side of (8.2) using the definition of Brownian loop measure and translation invariance:

$$\begin{aligned} \mu^{loop}(\gamma : 0 \in \bar{\gamma}, 1 \leq t_\gamma < e^{2r}) &= \int_{\mathbb{C}} \int_1^{e^{2r}} \frac{1}{2\pi t^2} \mu_{z,t}^{br}(\{\gamma : 0 \in \bar{\gamma}\}) dt d\mathbf{A}(z) \\ &= \int_{\mathbb{C}} \int_1^{e^{2r}} \frac{1}{2\pi t^2} \mu_{0,t}^{br}(\{\gamma : z \in \bar{\gamma}\}) dt d\mathbf{A}(z) \\ &= \int_1^{e^{2r}} \frac{1}{2\pi t^2} \mathbb{E}_{0,t}^{br} \left(\int_{\mathbb{C}} \mathbb{1}_{\{\gamma : z \in \bar{\gamma}\}} d\mathbf{A}(z) \right) dt \\ &= \int_1^{e^{2r}} \frac{1}{2\pi t} \mathbb{E}_{0,1}^{br} \left(\int_{\mathbb{C}} \mathbb{1}_{\{\gamma : z \in \bar{\gamma}\}} d\mathbf{A}(z) \right) dt, \end{aligned}$$

where $\mathbb{E}_{0,t}^{br}$ denotes expectation with respect to a complex Brownian bridge of time length t started at the origin, and where, in the last equality, we have used the fact that

$$\mathbb{E}_{0,t}^{br} \left(\int_{\mathbb{C}} \mathbb{1}_{\{\gamma: z \in \bar{\gamma}\}} d\mathbf{A}(z) \right) = t \mathbb{E}_{0,1}^{br} \left(\int_{\mathbb{C}} \mathbb{1}_{\{\gamma: z \in \bar{\gamma}\}} d\mathbf{A}(z) \right)$$

because of scaling. The expected area of a “filled-in” Brownian bridge, computed in [24], is

$$\mathbb{E}_{0,1}^{br} \left(\int_{\mathbb{C}} \mathbb{1}_{\{\gamma: z \in \bar{\gamma}\}} d\mathbf{A}(z) \right) = \frac{\pi}{5},$$

so that

$$\mu^{loop}(\gamma : z \in \bar{\gamma}, 1 \leq t_\gamma < e^{2r}) = \frac{r}{5}. \quad (8.3)$$

Using (8.3) and (8.2), we obtain

$$\mu^{loop}(\gamma : z \in \bar{\gamma}, 1 \leq \text{diam}(\gamma) < e^r) = \frac{r}{5}.$$

Comparing this to (8.1) gives $c = 1/5$. \square

Lemma 8.2. *Let $z \in \mathbb{C}$ and $k \in \mathbb{Z} \setminus \{0\}$, then*

$$\mu^{loop}(\gamma : \gamma \text{ has winding number } k \text{ around } z, \delta \leq \text{diam}(\gamma) < R) = \frac{1}{2\pi^2 k^2} \log \frac{R}{\delta}.$$

Proof. It is easy to check that the measure on loops surrounding the origin induced by μ^{loop} , but restricted to loops that wind k times around a given $z \in \mathbb{C}$, satisfies the conformal restriction property. Therefore, as in the proof of Lemma 8.1, we have

$$\begin{aligned} & \mu^{loop}(\gamma : \gamma \text{ has winding number } k \text{ around } z, \delta \leq \text{diam}(\gamma) < R) \\ &= \mu^{loop}(\gamma : \gamma \text{ has winding number } k \text{ around } z, \gamma \not\subset B_{z,\delta}, \gamma \subset B_{z,R}) \\ &= c_k \log \frac{R}{\delta}, \end{aligned}$$

for some positive constant $c_k < \infty$, where in the last equality we have used Proposition 3 of [18]. In order to find the constants c_k , we can proceed as in the proof of Lemma 8.1, using the fact that the expected area of a “filled-in” Brownian loop winding k times around the origin was computed in [24] and is equal to $1/2\pi k^2$ for $k \in \mathbb{Z} \setminus \{0\}$ (and $\pi/30$ for $k = 0$). \square

Acknowledgements

It is a pleasure to thank M. Bauer, D. Bernard, B. Duplantier, B. Freivogel, P. Kleban, G. Lawler, T. Lupu, Y. Le Jan, M. Lis, M. Porrati, A. Sokal, and S. Storage for discussions. The work of FC is supported in part by the Netherlands Organization for Scientific Research (NWO) through grant Vidi 639.032.916. The work of MK is supported in part by the NSF through grant PHY-1214302.

References

- [1] G. F. LAWLER and W. WERNER, “The Brownian loop soup,” *Probab. Theory Related Fields* **128** no. 4, (2004) 565–588. <http://dx.doi.org/10.1007/s00440-003-0319-6>.
- [2] S. SHEFFIELD and W. WERNER, “Conformal loop ensembles: the Markovian characterization and the loop-soup construction,” *Ann. of Math. (2)* **176** no. 3, (2012) 1827–1917. <http://dx.doi.org/10.4007/annals.2012.176.3.8>.
- [3] O. SCHRAMM, “Scaling limits of loop-erased random walks and uniform spanning trees [mr1776084],” in *Selected works of Oded Schramm. Volume 1, 2*, Sel. Works Probab. Stat., pp. 791–858. Springer, New York, 2011. http://dx.doi.org/10.1007/978-1-4419-9675-6_27.
- [4] B. FREIVOGEL and M. KLEBAN, “A Conformal Field Theory for Eternal Inflation,” *JHEP* **0912** (2009) 019, [arXiv:0903.2048](https://arxiv.org/abs/0903.2048) [hep-th].
- [5] B. FREIVOGEL, M. KLEBAN, A. NICOLIS, and K. SIGURDSON, “Eternal Inflation, Bubble Collisions, and the Disintegration of the Persistence of Memory,” *JCAP* **0908** (2009) 036, [arXiv:0901.0007](https://arxiv.org/abs/0901.0007) [hep-th].
- [6] J. M. MALDACENA, “The Large N limit of superconformal field theories and supergravity,” *Int.J.Theor.Phys.* **38** (1999) 1113–1133, [arXiv:hep-th/9711200](https://arxiv.org/abs/hep-th/9711200) [hep-th].
- [7] A. STROMINGER, “The dS / CFT correspondence,” *JHEP* **0110** (2001) 034, [arXiv:hep-th/0106113](https://arxiv.org/abs/hep-th/0106113) [hep-th].
- [8] L. SUSSKIND, “The Census taker’s hat,” [arXiv:0710.1129](https://arxiv.org/abs/0710.1129) [hep-th].
- [9] S. SMIRNOV, “Critical percolation in the plane: conformal invariance, Cardy’s formula, scaling limits,” *C. R. Acad. Sci. Paris Sér. I Math.* **333** no. 3, (2001) 239–244. [http://dx.doi.org/10.1016/S0764-4442\(01\)01991-7](http://dx.doi.org/10.1016/S0764-4442(01)01991-7).
- [10] F. CAMIA and C. M. NEWMAN, “Two-Dimensional Critical Percolation: The Full Scaling Limit,” *Communications in Mathematical Physics* **268** (Nov., 2006) 1–38, [math/0605035](https://arxiv.org/abs/math/0605035).

- [11] F. CAMIA and C. M. NEWMAN, “Critical percolation exploration path and SLE_6 : a proof of convergence,” *Probab. Theory Related Fields* **139** no. 3-4, (2007) 473–519. <http://dx.doi.org/10.1007/s00440-006-0049-7>.
- [12] F. CAMIA and C. M. NEWMAN, “ $SLE(6)$ and $CLE(6)$ from Critical Percolation,” in *Probability, geometry and integrable systems*, M. PINSKY and B. BIRNIR, eds., pp. 103–130. Cambridge University Press, Cambridge, 2008.
- [13] D. CHELKAK, H. DUMINIL-COPIN, C. HONGLER, A. KEMPPAINEN, and S. SMIRNOV, “Convergence of Ising interfaces to Schramm’s SLE curves,” *C. R. Math. Acad. Sci. Paris* **352** no. 2, (2014) 157–161. <http://dx.doi.org/10.1016/j.crma.2013.12.002>.
- [14] R. EASTHER, J. GIBLIN, JOHN T., L. HUI, and E. A. LIM, “A New Mechanism for Bubble Nucleation: Classical Transitions,” *Phys.Rev.* **D80** (2009) 123519, [arXiv:0907.3234](https://arxiv.org/abs/0907.3234) [hep-th].
- [15] M. KLEBAN, K. KRISHNAIYENGAR, and M. PORRATI, “Flux Discharge Cascades in Various Dimensions,” *JHEP* **1111** (2011) 096, [arXiv:1108.6102](https://arxiv.org/abs/1108.6102) [hep-th].
- [16] Y. LE JAN, *Markov paths, loops and fields*, vol. 2026 of *Lecture Notes in Mathematics*. Springer, Heidelberg, 2011. <http://dx.doi.org/10.1007/978-3-642-21216-1>. Lectures from the 38th Probability Summer School held in Saint-Flour, 2008, École d’Été de Probabilités de Saint-Flour. [Saint-Flour Probability Summer School].
- [17] Ş. NACU and W. WERNER, “Random soups, carpets and fractal dimensions,” *J. Lond. Math. Soc. (2)* **83** no. 3, (2011) 789–809. <http://dx.doi.org/10.1112/jlms/jdq094>.
- [18] W. WERNER, “The conformally invariant measure on self-avoiding loops,” *J. Amer. Math. Soc.* **21** no. 1, (2008) 137–169. <http://dx.doi.org/10.1090/S0894-0347-07-00557-7>.
- [19] P. DI FRANCESCO, P. MATHIEU, and D. SÉNÉCHAL, *Conformal field theory*. Graduate Texts in Contemporary Physics. Springer-Verlag, New York, 1997. <http://dx.doi.org/10.1007/978-1-4612-2256-9>.
- [20] D. AROVAS, J. SCHRIEFFER, F. WILCZEK, and A. ZEE, “Statistical Mechanics of Anyons,” *Nucl.Phys.* **B251** (1985) 117–126.
- [21] C. C. GERRY and V. A. SINGH, “Feynman path-integral approach to the aharonov-bohm effect,” *Phys. Rev. D* **20** (Nov, 1979) 2550–2554. <http://link.aps.org/doi/10.1103/PhysRevD.20.2550>.
- [22] J. DESBOIS and S. OUVRY, “Algebraic and arithmetic area for m planar Brownian paths,” *Journal of Statistical Mechanics: Theory and Experiment* **5** (May, 2011) 24, [arXiv:1101.4135](https://arxiv.org/abs/1101.4135) [math-ph].
- [23] F. CAMIA, “Brownian Loops and Conformal Fields,” *ArXiv e-prints* (Jan., 2015) , [arXiv:1501.04861](https://arxiv.org/abs/1501.04861) [math.PR].

- [24] C. GARBAN and J. A. T. FERRERAS, “The Expected Area of the Filled Planar Brownian Loop is $\pi/5$,” *Communications in Mathematical Physics* **264** (June, 2006) 797–810, [math/0504496](#).