PFAFFIAN OF APPELL'S HYPERGEOMETRIC SYSTEM F₄ IN TERMS OF THE INTERSECTION FORM OF TWISTED COHOMOLOGY GROUPS

YOSHIAKI GOTO, JYOICHI KANEKO, AND KEIJI MATSUMOTO

ABSTRACT. We study a Pfaffian of the system of differential equations annihilating Appell's hypergeometric series $F_4(a, b, c; x)$ by twisted cohomology groups associated with integrals representing solutions to this system. We simplify its connection matrix by the pull-back under a double cover of the complement of the singular locus. We express the simplified connection matrix in terms of the intersection form of the twisted cohomology groups.

1. INTRODUCTION

Appell's hypergeometric system $\mathcal{F}_4(a, b, c)$ is generated by differential equations (2.2) annihilating Appell's hypergeometric series $F_4(a, b, c; x)$ of $x = (x_1, x_2)$ defined by (2.1), where a, b and $c = (c_1, c_2)$ are parameters. It is a holonomic system of rank 4 with singular locus

$$S = \{(x_1, x_2) \in \mathbb{C}^2 \mid x_1 x_2 (x_1^2 + x_2^2 - 2x_1 x_2 - 2x_1 - 2x_2 + 1) = 0\} \cup L_{\infty},$$

where L_{∞} is the line at infinity in the projective plane \mathbb{P}^2 .

In this paper, we study a Pfaffian of the system $\mathcal{F}_4(a, b, c)$ by a twisted cohomology group $H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla)$ associated to integrals (2.3) representing solutions to this system. We regard the Pfaffian of $\mathcal{F}_4(a, b, c)$ as a connection ∇_X of a vector bundle over $X = \mathbb{P}^2 - S$ with fiber $H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla)$ over $x \in X$. We find a frame of this vector bundle and represent the connection as $\Xi = \Xi^1 dx_1 + \Xi^2 dx_2$ in Theorem 4.1 in terms of 4×4 -matrices Ξ^1 and Ξ^2 . The connection matrix satisfies the integrability condition $d\Xi = \Xi \wedge \Xi$ and has no apparent singularity, but Ξ_1 and Ξ_2 are complicated and $d\Xi \neq O$. To make the connection matrix simple, we consider a double cover Y of X defined by a map

$$\mathbb{C}^2 \ni (y_1, y_2) \mapsto (x_1, x_2) = (y_1(1 - y_2), y_2(1 - y_2)) \in \mathbb{C}^2.$$

Date: October 10, 2018.

²⁰¹⁰ Mathematics Subject Classification. 33C65, 58A17.

Key words and phrases. Appell's hypergeometric system of differential equations, Pfaffian system, Twisted cohomology group .

It induces the pull-back bundle and the pull-back connection ∇_Y . By changing a frame of the pull-back bundle, we express the pull-back connection ∇_Y by $\widehat{\Xi}$ in terms of logarithmic 1-forms of the variables y_1 and y_2 in Theorem 5.1. In particular, we have $d\widehat{\Xi} = \widehat{\Xi} \wedge \widehat{\Xi} = O$. It is studied in [Kat] that the pull-back of $F_4(a, b, c; x)$ under a similar map satisfies a Pfaffian equation equivalent to ours.

There is the intersection pairing \mathcal{I}_c between $H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla)$ and $H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla^{\vee})$, which is defined by the dual derivative ∇^{\vee} of ∇ . We have the dual vector bundle over X and the dual connection ∇_X^{\vee} satisfying

$$d_x \mathcal{I}_c(\varphi, \varphi') = \mathcal{I}_c(\nabla_X \varphi, \varphi') + \mathcal{I}_c(\varphi, \nabla_X^{\vee} \varphi'),$$

where φ and φ' are a section of the vector bundle and that of its dual, respectively, and d_x is the exterior derivative with respect to x_1 and x_2 . We can not find a frame of $H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla)$ satisfying

(1.1)
$$d_x \mathcal{I}_c(\varphi_i, \varphi_i^{\vee}) = 0$$

 $\mathbf{2}$

for any $1 \leq i, j \leq 4$, where $\varphi_i \in H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla)$ and φ_j^{\vee} is the image of φ_j under the natural map from $H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla)$ to $H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla^{\vee})$. The intersection pairing is defined between the pull-back bundles, and the frame of the pull-back bundle used in Theorem 5.1 satisfies the relations (1.1) for any $1 \leq i, j \leq 4$. The existence of the frame enables us to represent the pull-back connection ∇_Y by the intersection form \mathcal{I}_c in Theorem 5.2. We remark that this expression is not given in terms of matrices.

The monodromy representation of $\mathcal{F}_4(a, b, c)$ is initially given in [Kan] by a twisted homology group associated to integrals (2.3), and reconsidered in [GM] by the intersection pairing. In [GM], we characterize the circuit transform of a loop turning around each component of Sby the intersection form and a subspace of vanishing cycles as x approaches the component. In Lemma 5.4, we characterize the coefficient matrix of the logarithmic 1-form corresponding to a component of \widetilde{S} in $\widehat{\Xi}$ by the intersection form and a subspace of vanishing forms as xapproaches the component, where \widetilde{S} is the preimage of S under the double cover.

Appell's hypergeometric system $\mathcal{F}_4(a, b, c)$ is generalized to Lauricella's hypergeometric system $\mathcal{F}_C(a, b, c)$ of rank 2^m with m variables. For this system, we have twisted (co)homology groups associated to integrals representing solutions. The monodromy representation of $\mathcal{F}_C(a, b, c)$ is expressed in terms of intersection form between twisted homology groups in [G]. However, we have not seen that the system $\mathcal{F}_C(a, b, c)$ admits a Pfaffian system with a simple expression of the

pull-back connection under some covering map. We think that it is important to find a covering map and a frame of its pull-back bundle satisfying the relations (1.1) for any $1 \le i, j \le 2^m$.

For studies of Pfaffians of Lauricella's hypergeometric systems \mathcal{F}_A and \mathcal{F}_D of *m*-variables, it is easy to find such frames without considering covering maps. Their Pfaffians are expressed in terms of intersection forms between twisted cohomology groups associated to integrals representing solutions, refer to [M1] and [M2].

2. Appell's hypergeometric function F_4

In [ApKa], Appell's hypergeometric series $F_4(a, b, c; x)$ of variables x_1, x_2 with parameters $a, b, c = (c_1, c_2)$ is defined by

(2.1)
$$F_4(a, b, c_1, c_2; x_1, x_2) = \sum_{n_1, n_2=0}^{\infty} \frac{(a, n_1 + n_2)(b, n_1 + n_2)}{(c_1, n_1)(c_2, n_2)n_1!n_2!} x_1^{n_1} x_2^{n_2},$$

where $c_1, c_2 \neq 0, -1, -2, \ldots$ and $(a, k) = a(a+1)\cdots(a+k-1) = \Gamma(a+k)/\Gamma(a)$. This series converges in the domain

$$\mathbb{D} = \{ x = (x_1, x_2) \in \mathbb{C} \mid \sqrt{|x_1|} + \sqrt{|x_2|} < 1 \},\$$

and satisfies differential equations

$$\left[x_1(1-x_1)\partial_1^2 - x_2^2\partial_2^2 - 2x_1x_2\partial_1\partial_2 + \{c_1 - (a+b+1)x_1\}\partial_1 - (a+b+1)x_2\partial_2 - ab \right] f(x) = 0,$$

(2.2)

$$\begin{bmatrix} x_2(1-x_2)\partial_2^2 - x_1^2\partial_1^2 - 2x_1x_2\partial_1\partial_2 \\ +\{c_2 - (a+b+1)x_2\}\partial_2 - (a+b+1)x_1\partial_1 - ab \end{bmatrix} f(x) = 0.$$

The system generated by them is called Appell's hypergeometric system $\mathcal{F}_4(a, b, c_1, c_2)$ of differential equations. This system is of rank 4 with singular locus

$$S = \{(x_1, x_2) \in \mathbb{C}^2 \mid x_1 x_2 R(x) = 0\} \cup L_{\infty} \subset \mathbb{P}^2,$$

where L_{∞} is the line at infinity and

$$R(x) = x_1^2 + x_2^2 - 2x_1x_2 - 2x_1 - 2x_2 + 1.$$

We use the following integral representation of solutions of $\mathcal{F}_4(a, b, c)$

(2.3)
$$\int_{\Delta} u(s,x) \frac{ds_1 \wedge ds_2}{s_1 s_2},$$

where

4

$$u(s,x) = s_1^{\lambda_1} s_2^{\lambda_2} Q(s)^{\lambda_3} L(s,x)^{\lambda_4},$$

$$Q = Q(s) = s_1 s_2 - s_1 - s_2, \quad L = L(s,x) = 1 - s_1 x_1 - s_2 x_2,$$

$$\lambda_1 = a - c_2 + 1, \ \lambda_2 = a - c_1 + 1, \ \lambda_3 = c_1 + c_2 - a - 2, \ \lambda_4 = -b,$$

and a 2-chain Δ loading a branch of u(s, x) is a twisted 2-cycle. Refer to [AoKi] for twisted cycles and twisted homology groups.

Remark 2.1. It is convenient for the study of a Pfaffian system of $\mathcal{F}_4(a, b, c)$ to use the multi-valued function u(s, x) in stead of

$$u(t,x) = t_1^{1-c_1} t_2^{1-c_2} (1-t_1-t_2)^{c_1+c_2-a-1} (1-\frac{x_1}{t_1}-\frac{x_2}{t_2})^{-b}$$

used in [GM]. We have u(s, x) by the change of variables $(t_1, t_2) = (1/s_1, 1/s_2)$ and the replacement $a \mapsto a + 1$ for u(t, x).

3. Twisted cohomology group

We regard the parameters a, b and $c = (c_1, c_2)$ as indeterminants and we set

$$a_{00} = a, a_{10} = a - c_1 + 1, a_{01} = a - c_2 + 1, a_{11} = a - c_1 - c_2 + 2,$$

 $b_{00} = b, b_{10} = b - c_1 + 1, b_{01} = b - c_2 + 1, b_{11} = b - c_1 - c_2 + 2.$

We assume that

(3.1)
$$a_{ij}, b_{ij} \notin \mathbb{Z} \quad (i, j \in \mathbb{Z}_2 = \{0, 1\})$$

when we assign complex values to the parameters. Recall that

$$\lambda_1 = a - c_2 + 1, \ \lambda_2 = a - c_1 + 1, \ \lambda_3 = c_1 + c_2 - a - 2, \ \lambda_4 = -b.$$

In this section, we regard vector spaces as defined over the rational function field $\mathbb{C}(\lambda) = \mathbb{C}(\lambda_1, \ldots, \lambda_4) = \mathbb{C}(a, b, c_1, c_2).$

We set

$$\mathfrak{X} = \left\{ (s, x) \in \mathbb{C}^2 \times X \middle| s_1 s_2 Q(s) L(s, x) \neq 0 \right\} \subset (\mathbb{P}^1 \times \mathbb{P}^1) \times \mathbb{P}^2,$$

where X is the complement of the singular locus S of $\mathcal{F}_4(a, b, c)$ in \mathbb{P}^2 . There is a natural projection

$$\mathfrak{p}:\mathfrak{X}\ni(s,x)\mapsto x\in X.$$

For any fixed $x \in X$, we have

$$\mathbb{C}_x^2 = \mathfrak{p}^{-1}(x) = \{ s = (s_1, s_2) \in \mathbb{C}^2 \mid s_1 s_2 Q(s) L(s, x) \neq 0 \}$$

and an inclusion map

$$i_x : \mathbb{C}^2_x \ni s \mapsto (s, x) \in \mathfrak{X}.$$

We denote the $\mathbb{C}(\lambda)$ -algebra of rational functions on \mathbb{P}^2 with poles only along S by $\mathcal{O}(X)$. We denote the vector space of rational k-forms on $(\mathbb{P}^1 \times \mathbb{P}^1) \times \mathbb{P}^2$ with poles only along the complement of \mathfrak{X} by $\Omega^k(\mathfrak{X})$ and the subspace of $\Omega^{i+j}(\mathfrak{X})$ consisting elements which are *i*-forms with respect to the variables s_1, s_2 by $\Omega^{i,j}(\mathfrak{X})$.

We set

$$\omega = d_s \log(u(s,x)) = \frac{\lambda_1 ds_1}{s_1} + \frac{\lambda_2 ds_2}{s_2} + \frac{\lambda_3 d_s Q(s)}{Q(s)} + \frac{\lambda_4 d_s L(s,x)}{L(s,x)},$$

$$\omega_X = d_x \log(u(s,x)) = -\frac{\lambda_4 s_1 dx_1}{L(s,x)} - \frac{\lambda_4 s_2 dx_2}{L(s,x)},$$

where d_s and d_x are the exterior derivative with respect to s_1, s_2 and to x_1, x_2 , respectively. Note that $\omega \in \Omega^{1,0}(\mathfrak{X})$ and $\omega_X \in \Omega^{0,1}(\mathfrak{X})$. By a twisted exterior derivation $\nabla = d_s + \omega \wedge$ on \mathfrak{X} , we define quotient spaces

$$H^{k}(\Omega^{\bullet,0}(\mathfrak{X}),\nabla) = \ker\left(\nabla: \Omega^{k,0}(\mathfrak{X}) \to \Omega^{k+1,0}(\mathfrak{X})\right) \big/ \nabla\left(\Omega^{k-1,0}(\mathfrak{X})\right)$$

as $\mathcal{O}(X)$ -modules, where k = 0, 1, 2 and we regard $\Omega^{-1,0}(\mathfrak{X})$ as the zero vector space.

For a fixed x, the inclusion map ι_x induces a natural map from $H^k(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$ to the rational twisted cohomology group

$$H^{k}(\Omega^{\bullet}(\mathbb{C}^{2}_{x}), \nabla) = \ker \left(\nabla : \Omega^{k}(\mathbb{C}^{2}_{x}) \to \Omega^{k+1}(\mathbb{C}^{2}_{x}) \right) / \nabla \left(\Omega^{k-1}(\mathbb{C}^{2}_{x}) \right)$$

on \mathbb{C}_x^2 with respect to the twisted exterior derivative induced from ∇ . Here $\Omega^k(\mathbb{C}_x^2)$ is the vector space of rational k-forms with poles only along the complement of \mathbb{C}_x^2 in $\mathbb{P}^1 \times \mathbb{P}^1$. The structure of $H^k(\Omega^{\bullet}(\mathbb{C}_x^2), \nabla)$ is known as follows.

Fact 3.1 ([AoKi], [C]). (i) We have

$$\dim H^k(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla) = \begin{cases} 4 & \text{if } k = 2, \\ 0 & \text{if } k = 0, 1. \end{cases}$$

(ii) There is a canonical isomorphism

$$\begin{aligned} \jmath_x : H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla) &\to H^2(\mathcal{E}^{\bullet}_c(\mathbb{C}^2_x), \nabla) \\ &= \ker(\nabla : \mathcal{E}^2_c(\mathbb{C}^2_x) \to \mathcal{E}^3_c(\mathbb{C}^2_x)) / \nabla(\mathcal{E}^1_c(x)), \end{aligned}$$

where $\mathcal{E}_{c}^{k}(\mathbb{C}_{x}^{2})$ is the vector space of smooth k-forms with compact support in \mathbb{C}_{x}^{2} .

We have a twisted exterior derivation $\nabla^{\vee} = d_s - \omega \wedge$ for $-\omega$ and

$$\begin{aligned} H^2(\Omega^{\bullet,0}(\mathfrak{X}),\nabla^{\vee}) &= \Omega^{2,0}(\mathfrak{X}) / \nabla^{\vee}(\Omega^{1,0}(\mathfrak{X})), \\ H^2(\Omega^{\bullet}(\mathbb{C}^2_x),\nabla^{\vee}) &= \Omega^2(\mathbb{C}^2_x) / \nabla^{\vee}(\Omega^1(\mathbb{C}^2_x)). \end{aligned}$$

For any fixed $x \in X$, we define the intersection form between $H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla)$ and $H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla^{\vee})$ by

$$\mathcal{I}_c(\varphi_x,\varphi'_x) = \int_{\mathbb{C}^2_x} j_x(\varphi_x) \wedge \varphi'_x \in \mathbb{C}(\lambda),$$

where $\varphi_x, \varphi'_x \in \Omega^2(\mathbb{C}^2_x)$, j_x is given in Fact 3.1. This integral converges since $j_x(\varphi_x)$ is a smooth 2-form on \mathbb{C}^2_x with compact support. It is bi-linear over $\mathbb{C}(\lambda)$.

We take four elements $\varphi_1, \ldots, \varphi_4$ of $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$:

$$\varphi_1 = \frac{ds_{12}}{s_1 s_2}, \qquad \varphi_2 = \frac{x_1 ds_{12}}{s_2 L(s, x)},$$
$$\varphi_3 = \frac{x_2 ds_{12}}{s_1 L(s, x)}, \quad \varphi_4 = \frac{ds_{12}}{Q(s) L(s, x)},$$

where $ds_{12} = ds_1 \wedge ds_2$.

Proposition 3.1. For a fixed $x \in X$, the numbers $\mathcal{I}_c(i_x^*(\varphi_i), i_x^*(\varphi_j))$ $(1 \leq i, j \leq 4)$ are $(2\pi\sqrt{-1})^2 C_{ij}$, where

$$\begin{split} C_{11} &= \frac{1}{\lambda_{123}} \left(\frac{1}{\lambda_1} + \frac{1}{\lambda_2} \right) + \frac{1}{\lambda_{134}^-} \left(\frac{1}{\lambda_0} + \frac{1}{\lambda_2} \right) + \frac{1}{\lambda_{234}^-} \left(\frac{1}{\lambda_0} + \frac{1}{\lambda_1} \right) \\ &= \frac{1}{a_{00}} \left(\frac{1}{a_{01}} + \frac{1}{a_{10}} \right) + \frac{1}{b_{10}} \left(\frac{1}{b_{11}} + \frac{1}{a_{10}} \right) + \frac{1}{b_{01}} \left(\frac{1}{b_{11}} + \frac{1}{a_{01}} \right) , \\ C_{12} &= \frac{-1}{\lambda_{134}^-} \left(\frac{1}{\lambda_0} + \frac{1}{\lambda_2} \right) = \frac{-1}{b_{10}} \left(\frac{1}{b_{11}} + \frac{1}{a_{10}} \right) , \\ C_{13} &= \frac{-1}{\lambda_{234}^-} \left(\frac{1}{\lambda_0} + \frac{1}{\lambda_1} \right) = \frac{-1}{b_{01}} \left(\frac{1}{b_{11}} + \frac{1}{a_{01}} \right) , \\ C_{14} &= 0, \\ C_{22} &= \left(\frac{1}{\lambda_0} + \frac{1}{\lambda_2} \right) \left(\frac{1}{\lambda_4} + \frac{1}{\lambda_{134}^-} \right) = \left(\frac{1}{b_{11}} + \frac{1}{a_{10}} \right) \left(\frac{-1}{b_{00}} + \frac{1}{b_{10}} \right) , \\ C_{23} &= \frac{-1}{\lambda_0 \lambda_4} = \frac{1}{b_{11}b_{00}} , \\ C_{24} &= 0, \\ C_{33} &= \left(\frac{1}{\lambda_0} + \frac{1}{\lambda_1} \right) \left(\frac{1}{\lambda_4} + \frac{1}{\lambda_{234}^-} \right) = \left(\frac{1}{b_{11}} + \frac{1}{a_{01}} \right) \left(\frac{-1}{b_{00}} + \frac{1}{b_{01}} \right) , \\ C_{34} &= 0, \\ C_{44} &= \frac{2}{\lambda_3 \lambda_4 R(x)} = \frac{2}{a_{11}b_{00}R(x)} , \\ C_{ji} &= C_{ij} \quad for \quad i < j, \\ \lambda_0 &= -(\lambda_1 + \lambda_2 + 2\lambda_3 + \lambda_4) = b_{11}, \quad \lambda_{123} = \lambda_1 + \lambda_2 + \lambda_3 = a_{00}, \\ \lambda_{134}^- &= -(\lambda_1 + \lambda_3 + \lambda_4) = b_{10}, \\ \lambda_{134}^- &= -(\lambda_2 + \lambda_3 + \lambda_4) = b_{01}. \end{split}$$

The matrix $C = (C_{ij})_{i,j}$ is symmetric and its determinant is

$$\det(C) = \frac{4\lambda_3}{\lambda_0\lambda_1\lambda_2\lambda_4^3\lambda_{123}\lambda_{134}^-\lambda_{234}^-R(x)} = \left(\prod_{i,j=0,1}\frac{1}{a_{ij}b_{ij}}\right)\frac{4a_{11}^2}{b_{00}^2R(x)}.$$

Proof. For a fixed x, we blow up $\mathbb{P}^1 \times \mathbb{P}^1 (\supset \mathbb{C}^2_x)$ at two points (0,0) and (∞,∞) . We tabulate the residue of the pull back of ω to this space in Table 1, where E_0 and E_∞ are exceptional divisors coming from the blow up of $\mathbb{P}^1 \times \mathbb{P}^1$ at the points (0,0) and (∞,∞) , respectively. By

divisor	λ 's	a_{ij}, b_{ij}	a, b, c_1, c_2
E_{∞}	λ_0	b_{11}	$b - c_1 - c_2 + 2$
$s_1 = 0$	λ_1	a_{01}	$a - c_2 + 1$
$s_2 = 0$	λ_2	a_{10}	$a - c_1 + 1$
Q(s) = 0	λ_3	$-a_{11}$	$-a + c_1 + c_2 - 2$
L(s,x) = 0	λ_4	$-b_{00}$	-b
E_0	λ_{123}	a_{00}	a
$s_1 = \infty$	λ_{134}^{-}	b_{10}	$b - c_1 + 1$
$s_2 = \infty$	λ_{234}^-	b_{01}	$b - c_2 + 1$

TABLE 1. Residues for components of the pole divisor of ω

using these data, follow the proof of Theorem 5.1 in [GM].

Note that the matrix C is well-defined and $\det(C) \neq 0$ for any $x \in X$ under our assumption. The natural map from $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$ to $H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla)$ is surjective by Fact 3.1.

Corollary 3.1. We can regard the $\mathcal{O}(X)$ -modules $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$ and $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla^{\vee})$ as vector bundles

$$\bigcup_{x\in X} H^2(\mathscr{Q}^{\bullet}(\mathbb{C}^2_x), \nabla), \quad \bigcup_{x\in X} H^2(\mathscr{Q}^{\bullet}(\mathbb{C}^2_x), \nabla^{\vee}),$$

over X. The classes of $\varphi_1, \ldots, \varphi_4$ form frames of $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$ and $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla^{\vee})$.

Proof. We have only to prove that the natural map from $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$ to $H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla)$ is injective. We show that if an element $\varphi \in \Omega^{2,0}(\mathfrak{X})$ satisfying $i_x^*(\varphi) = 0$ as an element of $H^2(\Omega^{\bullet}(\mathbb{C}^2_x), \nabla)$ for any fixed $x \in X$ then φ belongs to $\nabla(\Omega^{1,0}(\mathfrak{X}))$. There exists $\psi_x \in \Omega^1(\mathbb{C}^2_x)$ such that $\nabla \psi_x = i_x^*(\varphi)$ for any x. Since this differential equation is of variables s_1, s_2 with parameters x_1, x_2, ψ_x can be globally extended to ψ . Hence we have $\psi \in \Omega^{1,0}(\mathfrak{X})$ such that $\nabla(\psi) = \varphi$. 8




FIGURE 1. Pole divisor of ω

By Proposition 3.1 and Corollary 3.1, the intersection form \mathcal{I}_c is regarded as a map from $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla) \times H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla^{\vee})$ to $\mathcal{O}(X)$.

4. Pfaffian system of $\mathcal{F}_4(a, b, c)$

For any $\varphi \in H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$, we have (4.1) $d_x \int_{\Delta} u(s,x)\varphi = \int_{\Delta} (d_x u(s,x) \wedge \varphi + u(s,x)d_x\varphi) = \int_{\Delta} u(s,x)(\nabla_X \varphi),$

where a 2-chain Δ loading a branch of u(s, x) is a twisted 2-cycle. Thus the exterior derivative d_x on X induces the connection $\nabla_X = d_x + \omega_X \wedge$

$$\nabla_X: H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla) \to H^2(\Omega^{\bullet,1}(\mathfrak{X}), \nabla) = \Omega^{2,1}(\mathfrak{X})/\nabla(\Omega^{1,1}(\mathfrak{X})).$$

By considering 1/u(s, x) instead of u(s, x), we also have the connection $\nabla_X^{\vee} = d_x - \omega_X \wedge$

 $\nabla_X^{\vee}: H^2(\varOmega^{\bullet,0}(\mathfrak{X}), \nabla^{\vee}) \to H^2(\varOmega^{\bullet,1}(\mathfrak{X}), \nabla^{\vee}) = \varOmega^{2,1}(\mathfrak{X})/\nabla^{\vee}(\varOmega^{1,1}(\mathfrak{X})).$

Proposition 4.1. The connection ∇_X is compatible with the intersection form \mathcal{I}_c , i.e., they satisfy

$$d_x \mathcal{I}_c(\varphi, \varphi') = \mathcal{I}_c(\nabla_X \varphi, \varphi') + \mathcal{I}_c(\varphi, \nabla_X^{\vee} \varphi').$$

Proof. It is enough to show this identity in any small simply connected domain U in X. We have

$$d_{x}\mathcal{I}_{c}(\varphi,\varphi') = d_{x} \int_{\mathbb{C}_{x}^{2}} u(s,x) j_{x}(\varphi) \wedge \frac{\varphi'}{u(s,x)}$$
$$= \int_{\mathbb{C}_{x}^{2}} d_{x}[u(s,x) j_{x}(\varphi)] \wedge \frac{\varphi'}{u(s,x)} + \int_{\mathbb{C}_{x}^{2}} u(s,x) j_{x}(\varphi) \wedge d_{x} \frac{\varphi'}{u(s,x)}$$
$$= \int_{\mathbb{C}_{x}^{2}} \nabla_{X}(j_{x}(\varphi)) \wedge \varphi' + \int_{\mathbb{C}_{x}^{2}} j_{x}(\varphi) \wedge \nabla_{X}^{\vee} \varphi',$$

since $j_x(\varphi)$ is with compact support for any point $x \in U$. By following the proof of Lemma 7.2 in [M1], we can show that $\nabla_X(j_x(\varphi))$ is cohomologue to $j_x(\nabla_X(\varphi))$. Hence we have

$$\int_{\mathbb{C}^2_x} \nabla_X(j_x(\varphi)) \wedge \varphi' = \int_{\mathbb{C}^2_x} j_x(\nabla_X(\varphi)) \wedge \varphi' = \mathcal{I}_c(\nabla_X \varphi, \varphi'),$$

which completes the proof.

Since $\nabla_X \varphi \in \Omega^{2,1}(\mathfrak{X})$, there exist Ξ_i^1 and Ξ_i^2 in $\mathcal{O}(X)$ such that

$$\nabla_X \varphi = dx_1 \wedge \sum_{i=1}^4 \Xi_i^1 \varphi_i + dx_2 \wedge \sum_{i=1}^4 \Xi_i^2 \varphi_i.$$

By calculating Ξ_i^1 and Ξ_i^2 for $\varphi = \varphi_1, \ldots, \varphi_4$, we represent the connection ∇_X as

$$\nabla_X {}^t(\varphi_1, \dots, \varphi_4) = \Xi \land {}^t(\varphi_1, \dots, \varphi_4), \quad \Xi = dx_1 \Xi^1 + dx_2 \Xi^2,$$

where Ξ^1 and Ξ^2 are 4×4 -matrices over the algebra $\mathcal{O}(X)$. By (4.1), the vector valued function

$$F(x) = {}^{t} \left(\int_{\Delta} u(s, x) \varphi_{1}, \dots, \int_{\Delta} u(s, x) \varphi_{4} \right)$$

satisfies a system of differential equations

$$d_x F(x) = \Xi \wedge F(x).$$

Let us determine Ξ^1 and Ξ^2 . A straightforward calculation implies the following.

Lemma 4.1.

10

$$\begin{aligned} \nabla_X(\varphi_1) &= dx_1 \wedge \frac{-\lambda_4 ds_{12}}{s_2 L} + dx_2 \wedge \frac{-\lambda_4 ds_{12}}{s_1 L} \\ &= \frac{dx_1}{x_1} \wedge (-\lambda_4)\varphi_2 + \frac{dx_2}{x_2} \wedge (-\lambda_4)\varphi_3, \\ \nabla_X(\varphi_2) &= dx_1 \wedge \frac{(1 - \lambda_4 s_1 x_1 - s_2 x_2) ds_{12}}{s_2 L^2} + dx_2 \wedge \frac{(1 - \lambda_4) x_1 ds_{12}}{L^2}, \\ \nabla_X(\varphi_3) &= dx_1 \wedge \frac{(1 - \lambda_4) x_2 ds_{12}}{L^2} + dx_2 \wedge \frac{(1 - s_1 x_1 - \lambda_4 s_2 x_2) ds_{12}}{s_1 L^2}, \\ \nabla_X(\varphi_4) &= dx_1 \wedge \frac{(1 - \lambda_4) s_1 ds_{12}}{Q L^2} + dx_2 \wedge \frac{(1 - \lambda_4) s_2 ds_{12}}{Q L^2}. \end{aligned}$$

To obtain Ξ^1 and Ξ^2 , we express

$$\frac{ds_{12}}{L^2}, \quad \frac{s_1 ds_{12}}{s_2 L^2}, \quad \frac{s_2 ds_{12}}{s_1 L^2}, \quad \frac{ds_{12}}{s_2 L^2}, \quad \frac{ds_{12}}{s_1 L^2}, \quad \frac{s_1 ds_{12}}{Q L^2}, \quad \frac{s_2 ds_{12}}{Q L^2},$$

in terms of $\varphi_1, \ldots, \varphi_4$.

Lemma 4.2. As elements of $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$, we have

$$\frac{2(\lambda_4 - 1)x_1x_2}{L^2}ds_{12} = (\lambda_1 + \lambda_2 + \lambda_3)\varphi_1 + (\lambda_2 - \lambda_4)\varphi_2 + (\lambda_1 - \lambda_4)\varphi_3 + \lambda_3(1 - x_1 - x_2)\varphi_4,$$

$$\frac{(\lambda_4 - 1)x_1s_1}{s_2L^2}ds_{12} = \frac{\lambda_1 + \lambda_3 + 1}{x_1}\varphi_2 + \lambda_3\varphi_4,$$

$$\frac{(\lambda_4 - 1)x_2s_2}{s_1L^2}ds_{12} = \frac{\lambda_2 + \lambda_3 + 1}{x_2}\varphi_3 + \lambda_3\varphi_4.$$

Proof. Straightforward calculations imply

$$\begin{aligned} \nabla \left(\frac{x_1 ds_1 - x_2 ds_2}{L} + \frac{ds_1}{s_1} - \frac{ds_2}{s_2} \right) \\ &= \frac{2(\lambda_4 - 1)x_1 x_2 ds_{12}}{L^2} \\ &- [(\lambda_1 + \lambda_2 + \lambda_3)\varphi_1 + (\lambda_2 - \lambda_4)\varphi_2 + (\lambda_1 - \lambda_4)\varphi_3 + \lambda_3(1 - x_1 - x_2)\varphi_4], \\ &\nabla \left(-\frac{s_1 ds_2}{s_2 L} \right) = \frac{(\lambda_4 - 1)x_1 s_1}{s_2 L^2} ds_{12} - \left[\frac{\lambda_1 + \lambda_3 + 1}{x_1} \varphi_2 + \lambda_3 \varphi_4 \right], \\ &\nabla \left(\frac{s_2 ds_1}{s_1 L} \right) = \frac{(\lambda_4 - 1)x_2 s_2}{s_1 L^2} ds_{12} - \left[\frac{\lambda_2 + \lambda_3 + 1}{x_2} \varphi_3 + \lambda_3 \varphi_4 \right], \end{aligned}$$

which shows this lemma.

Lemma 4.3. As elements of $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$, we have

$$\begin{aligned} \frac{2(\lambda_4 - 1)x_2}{s_1L^2} ds_{12} &= (\lambda_1 + \lambda_2 + \lambda_3)\varphi_1 + (\lambda_2 - \lambda_4)\varphi_2 \\ &+ (\lambda_1 + 2\lambda_2 + 2\lambda_3 + \lambda_4)\varphi_3 + \lambda_3(1 - x_1 + x_2)\varphi_4, \\ \frac{2(\lambda_4 - 1)x_1}{s_2L^2} ds_{12} &= (\lambda_1 + \lambda_2 + \lambda_3)\varphi_1 + (2\lambda_1 + \lambda_2 + 2\lambda_3 + \lambda_4)\varphi_2 \\ &+ (\lambda_1 - \lambda_4)\varphi_3 + \lambda_3(1 + x_1 - x_2)\varphi_4. \end{aligned}$$

Proof. Note that

$$\left(\frac{1}{s_1L^2} - \frac{x_1}{L^2} - \frac{s_2x_2}{s_1L^2}\right) ds_{12} = \frac{ds_{12}}{s_1L} = \frac{\varphi_3}{x_2},$$
$$\left(\frac{1}{s_2L^2} - \frac{x_2}{L^2} - \frac{s_1x_1}{s_2L^2}\right) ds_{12} = \frac{ds_{12}}{s_2L} = \frac{\varphi_2}{x_1}.$$

Use Lemma 4.2.

Lemma 4.4.

$$\begin{aligned} \frac{s_1 ds_{12}}{QL^2} &= \frac{1 - x_1 + x_2}{R(x)} \varphi_4 + \frac{(\lambda_1 + \lambda_2 + 2\lambda_3 + \lambda_4)(1 + x_1 - x_2)x_2}{\lambda_3 R(x)} \frac{ds_{12}}{L^2} \\ &- \frac{2\lambda_1 x_2}{\lambda_3 R(x)} \frac{ds_{12}}{s_1 L^2} - \frac{\lambda_2 (1 - x_1 - x_2)}{\lambda_3 R(x)} \frac{ds_{12}}{s_2 L^2}, \\ \frac{s_1 ds_{12}}{QL^2} &= \frac{1 + x_1 - x_2}{R(x)} \varphi_4 + \frac{(\lambda_1 + \lambda_2 + 2\lambda_3 + \lambda_4)(1 - x_1 + x_2)x_1}{\lambda_3 R(x)} \frac{ds_{12}}{L^2} \\ &- \frac{\lambda_1 (1 - x_1 - x_2)}{\lambda_3 R(x)} \frac{ds_{12}}{s_1 L^2} - \frac{2\lambda_2 x_1}{\lambda_3 R(x)} \frac{ds_{12}}{s_2 L^2}. \end{aligned}$$

Proof. Set

$$\eta_0 = \frac{ds_{12}}{QL^2}, \quad \eta_1 = \frac{s_1 ds_{12}}{QL^2}, \quad \eta_2 = \frac{s_2 ds_{12}}{QL^2}.$$

There is a relation

$$\eta_0 - x_1\eta_1 - x_2\eta_2 = \frac{(1 - s_1x_1 - s_2x_2)ds_{12}}{QL^2} = \varphi_4$$

among them. We have

$$\begin{aligned} \nabla\left(\frac{x_1ds_1+x_2ds_2}{L^2}\right) \\ &= \lambda_3(x_1-x_2)\eta_0 - \lambda_3x_1\eta_1 + \lambda_3x_2\eta_2 + (\frac{\lambda_1x_2}{s_1L^2} - \frac{\lambda_2x_1}{s_2L^2})ds_{12}, \\ \nabla\left(\frac{ds_1-ds_2}{L^2} + \frac{(x_1+x_2)(-s_2ds_1+s_1ds_2)}{L^2}\right) \\ &= 2\lambda_3\eta_0 + \lambda_3(x_1+x_2-1)\eta_1 + \lambda_3(x_1+x_2-1)\eta_2 \\ &- \left(\frac{\lambda_1}{s_1L^2} + \frac{\lambda_2}{s_2L^2} - \frac{(\lambda_1+\lambda_2+2\lambda_3+\lambda_4)(x_1+x_2)}{L^2}\right)ds_{12}, \end{aligned}$$

which are zero as elements of $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$. By regarding these relations as linear equations of variables η_0, η_1, η_2 , solve them.

Remark 4.1. The form $\eta_0 = ds_{12}/QL^2$ in the proof of Lemma 4.4 is expressed as

$$\eta_0 = \frac{1 - x_1 - x_2}{R(x)} \varphi_4 + \frac{2(\lambda_1 + \lambda_2 + 2\lambda_3 + \lambda_4)x_1x_2}{\lambda_3 R(x)} \frac{ds_{12}}{L^2} - \frac{\lambda_1 x_2(1 + x_1 - x_2)}{\lambda_3 R(x)} \frac{ds_{12}}{s_1 L^2} - \frac{\lambda_2 x_1(1 - x_1 + x_2)}{\lambda_3 R(x)} \frac{ds_{12}}{s_2 L^2}.$$

Lemmas 4.1, 4.2, 4.3 and 4.4 yield the representation of ∇_X with respect to the frame ${}^t(\varphi_1, \ldots, \varphi_4)$.

Theorem 4.1. With respect to the frame ${}^{t}(\varphi_1, \ldots, \varphi_4)$, the connection ∇_X is represented as

$$\nabla_X {}^t(\varphi_1,\ldots,\varphi_4) = \Xi \wedge {}^t(\varphi_1,\ldots,\varphi_4),$$

where $\Xi = \Xi^1 dx_1 + \Xi^2 dx_2$ and

$$\Xi^{1} = \begin{pmatrix} 0 & -\frac{\lambda_{4}}{x_{1}} & 0 & 0\\ 0 & -\frac{\lambda_{1}+\lambda_{3}}{x_{1}} & 0 & -\lambda_{3}\\ -\frac{\lambda_{1}+\lambda_{2}+\lambda_{3}}{2x_{1}} & -\frac{\lambda_{2}-\lambda_{4}}{2x_{1}} & -\frac{\lambda_{1}-\lambda_{4}}{2x_{1}} & -\frac{\lambda_{3}(1-x_{1}-x_{2})}{2x_{1}}\\ \Xi^{1}_{4,1} & \Xi^{1}_{4,2} & \Xi^{1}_{4,3} & \Xi^{1}_{4,4} \end{pmatrix}$$

,

$$\Xi^{2} = \begin{pmatrix} 0 & 0 & -\frac{\lambda_{4}}{x_{2}} & 0 \\ -\frac{\lambda_{1} + \lambda_{2} + \lambda_{3}}{2x_{2}} & -\frac{\lambda_{2} - \lambda_{4}}{2x_{2}} & -\frac{\lambda_{1} - \lambda_{4}}{2x_{2}} & -\frac{\lambda_{3}(1 - x_{1} - x_{2})}{2x_{2}} \\ 0 & 0 & -\frac{\lambda_{2} + \lambda_{3}}{x_{2}} & -\lambda_{3} \\ \Xi^{2}_{4,1} & \Xi^{2}_{4,2} & \Xi^{2}_{4,3} & \Xi^{2}_{4,4} \end{pmatrix},$$

$$\Xi_{4,1}^{1} = \frac{\lambda_{1} + \lambda_{2} + \lambda_{3}}{\lambda_{3}} \left(\frac{\lambda_{1} - \lambda_{2}}{R(x)} - \frac{(\lambda_{1} + 2\lambda_{3} + \lambda_{4})(1 + x_{1} - x_{2})}{2x_{1}R(x)} \right),$$

$$\Xi_{4,2}^{1} = -\frac{(\lambda_{2} - \lambda_{4})(\lambda_{2} + 2\lambda_{3} + \lambda_{4})}{\lambda_{3}R(x)}$$

$$\begin{aligned} &+ \frac{\lambda_1 \lambda_2 + \lambda_1 \lambda_4 + \lambda_2 \lambda_4 + 2\lambda_3 \lambda_4 + \lambda_4^2}{2\lambda_3} \frac{1 - x_1 - x_2}{x_1 R(x)}, \\ \Xi_{4,3}^1 &= \frac{\lambda_1 \lambda_2 + \lambda_1 \lambda_4 + \lambda_2 \lambda_4 + 2\lambda_3 \lambda_4 + \lambda_4^2}{\lambda_3 R(x)} \\ &- \frac{(\lambda_1 - \lambda_4)(\lambda_1 + \lambda_4 + 2\lambda_3)}{2\lambda_3} \frac{1 - x_1 - x_2}{x_1 R(x)}, \\ \Xi_{4,4}^1 &= -(\lambda_1 - 2\lambda_3 - 3\lambda_4 + 2) \frac{x_1}{2R(x)} + (\lambda_1 - \lambda_4 + 1) \frac{1 + x_2}{R(x)} \\ &- (\lambda_1 + 2\lambda_3 + \lambda_4) \frac{(x_2 - 1)^2}{2x_1 R(x)}, \end{aligned}$$

$$\begin{split} \Xi_{4,1}^2 &= \frac{\lambda_1 + \lambda_2 + \lambda_3}{\lambda_3} \left(\frac{\lambda_2 - \lambda_1}{R(x)} - \frac{(\lambda_2 + 2\lambda_3 + \lambda_4)(1 - x_1 + x_2)}{2x_2 R(x)} \right), \\ \Xi_{4,2}^2 &= \frac{\lambda_1 \lambda_2 + \lambda_1 \lambda_4 + \lambda_2 \lambda_4 + 2\lambda_3 \lambda_4 + \lambda_4^2}{\lambda_3 R(x)} \\ &\quad - \frac{(\lambda_2 - \lambda_4)(\lambda_2 + 2\lambda_3 + \lambda_4)}{2\lambda_3} \frac{1 - x_1 - x_2}{x_2 R(x)}, \\ \Xi_{4,3}^2 &= -\frac{(\lambda_1 - \lambda_4)(\lambda_1 + \lambda_4 + 2\lambda_3)}{\lambda_3 R(x)} \\ &\quad + \frac{\lambda_1 \lambda_2 + \lambda_1 \lambda_4 + \lambda_2 \lambda_4 + 2\lambda_3 \lambda_4 + \lambda_4^2}{2\lambda_3} \frac{1 - x_1 - x_2}{x_2 R(x)}, \\ \Xi_{4,4}^2 &= -(\lambda_2 - 2\lambda_3 - 3\lambda_4 + 2) \frac{x_2}{2 R(x)} + (\lambda_2 - \lambda_4 + 1) \frac{x_1 + 1}{R(x)} \\ &\quad - (\lambda_2 + 2\lambda_3 + \lambda_4) \frac{(x_1 - 1)^2}{2x_2 R(x)}. \end{split}$$

Remark 4.2. Though the connection matrix Ξ is not closed, it satisfies the integrability condition, *i.e.*,

$$\Xi \wedge \Xi = (\Xi^1 \Xi^2 - \Xi^2 \Xi^1) dx_1 \wedge dx_2$$
$$= d_x \Xi = \left(-\frac{\partial}{\partial x_2} \Xi^1 + \frac{\partial}{\partial x_1} \Xi^2 \right) dx_1 \wedge dx_2 \neq O.$$

Remark 4.3. We give some expressions of elements of $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$ in terms of $\varphi_1, \ldots, \varphi_4$:

$$\begin{aligned} \nabla(-\frac{ds_1}{s_1} + \frac{ds_2}{s_2}) &= \frac{\lambda_3 ds_{12}}{Q} - \left[-(\lambda_1 + \lambda_2 + \lambda_3)\varphi_1 + \lambda_4\varphi_2 + \lambda_4\varphi_3\right], \\ \nabla(\frac{ds_2}{s_2}) &= \frac{\lambda_3 ds_{12}}{s_1 Q} - \left[-(\lambda_1 + \lambda_3)\varphi_1 + \lambda_4\varphi_2\right], \\ \nabla(-\frac{ds_1}{s_1}) &= \frac{\lambda_3 ds_{12}}{s_2 Q} - \left[-(\lambda_2 + \lambda_3)\varphi_1 + \lambda_4\varphi_3\right]. \end{aligned}$$

Remark 4.4. We define a function $f_1(x)$ by $\int_{\Delta} u(s, x)\varphi_1$ for a twisted cycle Δ loading a branch of u(s, x). This function satisfies

$$\begin{aligned} x_1 \frac{\partial f_1(x)}{\partial x_1} &= -\lambda_4 \int_{\Delta} u(s, x)\varphi_2, \quad x_2 \frac{\partial f_1(x)}{\partial x_2} = -\lambda_4 \int_{\Delta} u(s, x)\varphi_3, \\ x_1 x_2 \frac{\partial^2 f_1(x)}{\partial x_1 \partial x_2} &= \frac{\lambda_4}{2} \Big[\lambda_{123} \int_{\Delta} u(s, x)\varphi_1 + (\lambda_2 - \lambda_4) \int_{\Delta} u(s, x)\varphi_2 \\ &+ (\lambda_1 - \lambda_4) \int_{\Delta} u(s, x)\varphi_3 + \lambda_3 (1 - x_1 - x_2) \int_{\Delta} u(s, x)\varphi_4 \Big] \end{aligned}$$

Thus the vector-valued function

$$F_{\partial}(x) = {}^{t} \left(f_{1}(x), x_{1} \frac{\partial f_{1}(x)}{\partial x_{1}}, x_{2} \frac{\partial f_{1}(x)}{\partial x_{2}}, x_{1} x_{2} \frac{\partial^{2} f_{1}(x)}{\partial x_{1} \partial x_{2}} \right)$$

satisfies

$$d_x F_{\partial}(x) = (G_{\partial} \Xi G_{\partial}^{-1} + d_x G_{\partial} \ G_{\partial}^{-1}) F_{\partial}(x),$$

where

$$G_{\partial} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -\lambda_4 & 0 & 0 \\ \frac{\lambda_{123}\lambda_4}{2} & \frac{(\lambda_2 - \lambda_4)\lambda_4}{2} & \frac{(\lambda_1 - \lambda_4)\lambda_4}{2} & \frac{\lambda_3\lambda_4(1 - x_1 - x_2)}{2} \end{pmatrix},$$

$$G_{\partial}^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -\lambda_4^{-1} & 0 & 0 \\ 0 & 0 & -\lambda_4^{-1} & 0 \\ \frac{\lambda_{123}}{\lambda_3(1 - x_1 - x_2)} & \frac{\lambda_2 - \lambda_4}{\lambda_3\lambda_4(1 - x_1 - x_2)} & \frac{\lambda_1 - \lambda_4}{\lambda_3\lambda_4(1 - x_1 - x_2)} \end{pmatrix}.$$

Note that the matrix G_{∂} does not belongs to $GL_4(\mathcal{O}(X))$.

5. Connection matrix in terms of intersection form

We define an affine variety

$$\widetilde{X} = \{ (x_1, x_2, x_3) \in \mathbb{C}^3 \mid x_3^2 = R(x) \},\$$

which is regarded as the double cover of \mathbb{C}^2 branching along the divisor $R(x_1, x_2) = 0$ by the projection

$$pr: \widetilde{X} \ni (x_1, x_2, x_3) \to (x_1, x_2) \in \mathbb{C}^2.$$

Note that

$$\begin{pmatrix} \frac{1+x_1-x_2-x_3}{2} \end{pmatrix} \begin{pmatrix} \frac{1+x_1-x_2+x_3}{2} \end{pmatrix} = x_1, \\ \begin{pmatrix} \frac{1-x_1+x_2-x_3}{2} \end{pmatrix} \begin{pmatrix} \frac{1-x_1+x_2+x_3}{2} \end{pmatrix} = x_2, \\ \sim \end{pmatrix}$$

for $(x_1, x_2, x_3) \in \widetilde{X}$. Thus the preimages of lines $x_1 = 0$ and $x_2 = 0$ in \mathbb{C}^2 under the projection *pr* are expressed as equations

$$\frac{1+x_1-x_2-x_3}{2} = 0, \quad \frac{1-x_1+x_2-x_3}{2} = 1,$$

and

$$\frac{1+x_1-x_2-x_3}{2} = 1, \quad \frac{1-x_1+x_2-x_3}{2} = 0$$

in \widetilde{X} respectively. By a map

$$\widetilde{X} \ni (x_1, x_2, x_3) \mapsto (y_1, y_2) = \left(\frac{1 + x_1 - x_2 - x_3}{2}, \frac{1 - x_1 + x_2 - x_3}{2}\right) \in \mathbb{C}^2,$$

 \widetilde{X} is bi-holomorphic to \mathbb{C}^2 ; its inverse is expressed as

$$\mathbb{C}^2 \ni (y_1, y_2) \mapsto (x_1, x_2, x_3) = (y_1(1-y_2), (1-y_1)y_2, 1-y_1-y_2) \in \widetilde{X}.$$

Though (x_1, x_2) are not valid as local coordinates on the set

$$\left\{ (x_1, x_2, x_3) \in \widetilde{X} \middle| \frac{\partial}{\partial x_2} (x_3^2 - R(x)) = 2x_3 = 0 \right\},\$$

we can use $(y_1, y_2) = \left(\frac{1+x_1-x_2-x_3}{2}, \frac{1-x_1+x_2-x_3}{2}\right)$ as a global coordinates system on \widetilde{X} . The covering transformation

 $\rho: (x_1, x_2, x_3) \mapsto (x_1, x_2, -x_3)$

of $pr:\widetilde{X}\to X$ is represented as

$$(y_1, y_2) \mapsto (1 - y_2, 1 - y_1).$$

The ramification locus of pr is

$$\{(y_1, y_2) \in \mathbb{C}^2 \mid y_1 + y_2 = 1\}$$

16 YOSHIAKI GOTO, JYOICHI KANEKO, AND KEIJI MATSUMOTO

and the preimage of the singular locus of $\mathcal{F}_4(a, b, c)$ in \mathbb{C}^2 under the projection pr is

$$\{(y_1, y_2) \in \mathbb{C}^2 \mid [y_1(1-y_2)] \cdot [y_2(1-y_1)] \cdot (1-y_1-y_2) = 0\}.$$

We set

$$Y = \{y = (y_1, y_2) \in \mathbb{C}^2 \mid y_1(1 - y_1)y_2(1 - y_2)(1 - y_1 - y_2) \neq 0\} \subset \mathbb{P}^2,$$

$$\widetilde{S} = \mathbb{P}^2 - Y.$$

Note that \widetilde{S} and Y are invariant as sets under the action of ρ . By the vector bundle $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$ over X and the projection $pr: Y \to X$, we have the vector bundle

$$pr^*H^2(\Omega^{\bullet,0}(\mathfrak{X}),\nabla) = \{(y,\varphi) \in Y \times H^2(\Omega^{\bullet,0}(\mathfrak{X}),\nabla) \mid pr(y) = p(\varphi)\}.$$

Its frame ${}^{t}(\widetilde{\varphi}_{1},\ldots,\widetilde{\varphi}_{4})$ is given by the pull-backs

$$\widetilde{\varphi}_i = pr^*(\varphi_i) \quad (i = 1, \dots, 4)$$

of $\varphi_1, \ldots, \varphi_4$ in $H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$ under the projection pr. The connection $\nabla_Y = pr^* \nabla_X$ of $pr^* H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$ for the frame ${}^t(\widetilde{\varphi}_1, \ldots, \widetilde{\varphi}_4)$ is defined by the pull-back of Ξ under the projection pr.

Corollary 5.1. With respect to the frame ${}^{t}(\widetilde{\varphi}_{1},\ldots,\widetilde{\varphi}_{4})$, the connection ∇_{Y} is expressed as

$$\widetilde{\Xi} \wedge {}^t (\widetilde{\varphi}_1, \dots, \widetilde{\varphi}_4), \quad \widetilde{\Xi} = \widetilde{\Xi}^1 dy_1 + \widetilde{\Xi}^2 dy_2,$$

where

$$\widetilde{\Xi}^{1} = \begin{pmatrix} 0 & \frac{-\lambda_{4}}{y_{1}} & \frac{-\lambda_{4}}{y_{1}-1} & 0 \\ -\frac{\lambda_{123}}{2(y_{1}-1)} & \frac{-\lambda_{2}+\lambda_{4}}{2(y_{1}-1)} - \frac{\lambda_{1}+\lambda_{3}}{y_{1}} & \frac{-\lambda_{1}+\lambda_{4}}{2(y_{1}-1)} & -\frac{\lambda_{3}}{2} - \frac{\lambda_{3}y_{2}}{2(y_{1}-1)} \\ -\frac{\lambda_{123}}{2y_{1}} & \frac{-\lambda_{2}+\lambda_{4}}{2y_{1}} & \frac{-\lambda_{1}+\lambda_{4}}{2y_{1}} - \frac{\lambda_{2}+\lambda_{3}}{y_{1}-1} & \frac{\lambda_{3}}{2} + \frac{\lambda_{3}(y_{2}-1)}{2y_{1}} \\ \widetilde{\Xi}_{41}^{1} & \widetilde{\Xi}_{42}^{1} & \widetilde{\Xi}_{43}^{1} & \widetilde{\Xi}_{44}^{1} \end{pmatrix},$$

$$\widetilde{\Xi}^{2} = \begin{pmatrix} 0 & \frac{-\lambda_{4}}{y_{2}-1} & \frac{-\lambda_{4}}{y_{2}} & 0 \\ -\frac{\lambda_{123}}{2y_{2}} & \frac{-\lambda_{2}+\lambda_{4}}{2y_{2}} - \frac{\lambda_{1}+\lambda_{3}}{y_{2}-1} & \frac{-\lambda_{1}+\lambda_{4}}{2y_{2}} & \frac{\lambda_{3}}{2} + \frac{\lambda_{3}(y_{1}-1)}{2}y_{2} \\ -\frac{\lambda_{123}}{2(y_{2}-1)} & \frac{-\lambda_{2}+\lambda_{4}}{2(y_{2}-1)} & \frac{-\lambda_{1}+\lambda_{4}}{2(y_{2}-1)} - \frac{\lambda_{2}+\lambda_{3}}{y_{2}} & -\frac{\lambda_{3}}{2} - \frac{\lambda_{3}y_{1}}{2(y_{2}-1)} \\ \widetilde{\Xi}_{41}^{2} & \widetilde{\Xi}_{42}^{2} & \widetilde{\Xi}_{43}^{2} & \widetilde{\Xi}_{44}^{2} \end{pmatrix},$$

$$\begin{split} \widetilde{\Xi}_{41}^{1} &= \frac{(\lambda_{3} - \lambda_{134}^{-})\lambda_{123}}{(2\lambda_{3}y_{1}(y_{2} - 1))} - \frac{(\lambda_{3} - \lambda_{234}^{-})\lambda_{123}}{(2\lambda_{3}(y_{1} - 1)y_{2})} \\ &+ \frac{(\lambda_{3} - \lambda_{234}^{-})\lambda_{123}}{(2\lambda_{3}y_{2}(y_{1} + y_{2} - 1))} - \frac{(2\lambda_{3} + \lambda_{4} + \lambda_{1})\lambda_{123}}{(2\lambda_{3}(y_{2} - 1)(y_{1} + y_{2} - 1))}, \\ \widetilde{\Xi}_{42}^{1} &= \frac{(\lambda_{2} + \lambda_{4})(\lambda_{1} + \lambda_{4})}{(2\lambda_{3}(y_{2} - 1)(y_{1} + y_{2} - 1))} - \frac{(\lambda_{2} - \lambda_{4})(\lambda_{3} - \lambda_{234}^{-})}{(2\lambda_{3}y_{2}(y_{1} - 1))} + \frac{(\lambda_{2} - \lambda_{4})(\lambda_{3} - \lambda_{234}^{-})}{(2\lambda_{3}y_{2}(y_{1} + y_{2} - 1))}, \\ \widetilde{\Xi}_{43}^{1} &= \frac{(\lambda_{2} + \lambda_{4})(\lambda_{1} + \lambda_{4})}{(2\lambda_{3}y_{2}(y_{1} - 1))} - \frac{(\lambda_{2} + \lambda_{4})(\lambda_{1} + \lambda_{4})}{(2\lambda_{3}y_{2}(y_{1} + y_{2} - 1))} - \frac{\lambda_{4}}{(2\lambda_{3}y_{2}(y_{1} + y_{2} - 1))} \\ &+ \frac{\lambda_{4}}{y_{2}(y_{1} - 1)} - \frac{(\lambda_{1} - \lambda_{4})(\lambda_{3} - \lambda_{134}^{-})}{(2\lambda_{3}(y_{2} - 1)(y_{1} + y_{2} - 1))} + \frac{(\lambda_{1} - \lambda_{4})(\lambda_{3} - \lambda_{134}^{-})}{(2\lambda_{3}(y_{2} - 1)y_{1})}, \\ \widetilde{\Xi}_{41}^{2} &= \frac{(2\lambda_{4} + 2\lambda_{3} - 1)}{(2\lambda_{3}(y_{1} - 1))} - \frac{\lambda_{3} - \lambda_{134}^{-}}{(2\lambda_{3}y_{1}(y_{1} + y_{2} - 1))} \\ &- \frac{(\lambda_{2} + 2\lambda_{3} + \lambda_{4})\lambda_{123}}{(2\lambda_{3}y_{1}(y_{2} - 1))} + \frac{(\lambda_{2} + 2\lambda_{3} + \lambda_{4})\lambda_{123}}{(2\lambda_{3}(y_{1} - 1)(y_{1} + y_{2} - 1))} \\ &- \frac{(\lambda_{2} + 2\lambda_{3} + \lambda_{4})\lambda_{123}}{(2\lambda_{3}(y_{1} - 1)(y_{1} + y_{2} - 1)} - \frac{\lambda_{4}}{(y_{1} + \lambda_{4})} \\ \widetilde{\Xi}_{42}^{2} &= \frac{(\lambda_{2} + \lambda_{4})(\lambda_{1} + \lambda_{4})}{(2\lambda_{3}(y_{2} - 1)y_{1}} - \frac{(\lambda_{2} - \lambda_{4})(\lambda_{3} - \lambda_{234}^{-})}{(2\lambda_{3}(y_{1} - 1)(y_{1} + y_{2} - 1)} + \frac{(\lambda_{2} - \lambda_{4})(\lambda_{3} - \lambda_{234}^{-})}{(2\lambda_{3}y_{2}(y_{1} - 1))} \\ &+ \frac{\lambda_{4}}{(y_{2} - 1)y_{1}} - \frac{(\lambda_{2} - \lambda_{4})(\lambda_{3} - \lambda_{234}^{-})}{(2\lambda_{3}(y_{1} - 1)(y_{1} + y_{2} - 1)} + \frac{(\lambda_{4} - \lambda_{4})(\lambda_{3} - \lambda_{234}^{-})}{(\lambda_{3}y_{2}(y_{1} - 1))} \\ &- \frac{\lambda_{4}}{(y_{2}(y_{1} - 1)} - \frac{(\lambda_{1} - \lambda_{4})(\lambda_{3} - \lambda_{134}^{-})}{(2\lambda_{3}(y_{2} - 1)y_{1}} + \frac{(\lambda_{1} - \lambda_{4})(\lambda_{3} - \lambda_{134}^{-})}{(2\lambda_{3}y_{2}(y_{1} - 1))} \\ &- \frac{\lambda_{4}}{(y_{2} - 1)y_{1}} - \frac{(\lambda_{2} - \lambda_{4})(\lambda_{1} - \lambda_{234}^{-})}{(2\lambda_{3}(y_{1} - 1)(y_{1} + y_{2} - 1)} + \frac{\lambda_{4}}{(\lambda_{3} - \lambda_{234}^{-})}{(\lambda_{3} - \lambda_{234}^{-})}, \\ \widetilde{\Xi}_{42}^{2} &= -\frac{($$

Proof. By using

$$\begin{aligned} pr^*(x_1) &= y_1(1-y_2), & pr^*(x_1) &= (1-y_1)y_2, \\ pr^*(dx_1) &= (1-y_2)dy_1 - y_1dy_2, & pr^*(dx_2) &= -y_2dy_1 + (1-y_1)dy_2, \\ pr^*(R(x)) &= (1-y_1-y_2)^2, \end{aligned}$$

we have only to calculate the pull back of Ξ under the projection pr. \Box

Let $\mathcal{O}(Y)$ be the $\mathbb{C}(\lambda)$ -algebra of rational functions on \mathbb{P}^2 with poles only along the complement of Y. We change the frame ${}^{t}(\widetilde{\varphi}_{1},\ldots,\widetilde{\varphi}_{4})$ of the vector bundle $pr^*H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$ over Y as

$$f(\widehat{\varphi}_1, \dots, \widehat{\varphi}_4) = G^t(\widetilde{\varphi}_1, \dots, \widetilde{\varphi}_4),$$

$$G = \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \\ & & & 1 - y_1 - y_2 \end{pmatrix} \in GL_4(\mathcal{O}(Y)).$$

The connection matrix $\widehat{\Xi}$ of $pr^*H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$ with respect to the frame ${}^{t}(\widehat{\varphi}_{1},\ldots,\widehat{\varphi}_{4})$ is given by the gauge transformation

$$G \widetilde{\Xi} G^{-1} + d_y G G^{-1}$$

of $\widetilde{\Xi}$. Straightforward calculations imply the following.

Theorem 5.1. The connection matrix $\widehat{\Xi}$ is $\widehat{\Xi}^{1}\frac{dy_{1}}{y_{1}} + \widehat{\Xi}^{2}\frac{dy_{2}}{y_{2}} + I_{3,1}\widehat{\Xi}^{2}I_{3,1}^{-1}\frac{dy_{1}}{y_{1}-1} + I_{3,1}\widehat{\Xi}^{1}I_{3,1}^{-1}\frac{dy_{2}}{y_{2}-1} + \widehat{\Xi}^{3}\frac{dy_{1}+dy_{2}}{y_{1}+y_{2}-1},$ where

The connection matrix $\widehat{\Xi}$ satisfies

$$d_y\widehat{\Xi} = \widehat{\Xi} \wedge \widehat{\Xi} = O$$

Remark 5.1. The gauge transformation by G changes the non-closed connection matrix $\widehat{\Xi}$ into the closed connection matrix $\widehat{\Xi}$.

We express the connection ∇_Y in terms of intersection form \mathcal{I}_c . Since the map $pr: Y \to X$ is locally isomorphic, we have

$$\mathcal{I}_c(\widetilde{\varphi}_i,\widetilde{\varphi}_j) = \mathcal{I}_c(\varphi_i,\varphi_j), \quad (1 \le i,j \le 4).$$

We give the intersection numbers for the frame ${}^{t}(\widehat{\varphi}_{1},\ldots,\widehat{\varphi}_{4})$.

Corollary 5.2. The intersection numbers $\mathcal{I}_c(\widehat{\varphi}_i, \widehat{\varphi}_j)$ $(1 \le i, j \le 4)$ are $(2\pi\sqrt{-1})^2 \widehat{C}_{ij}$, where

$$\widehat{C}_{ij} = \begin{cases} C_{ij}, & \text{if } 1 \le i, j \le 3, \\ \frac{2}{\lambda_3 \lambda_4} = \frac{2}{a_{11} b_{00}}, & \text{if } (i, j) = (4, 4), \\ 0, & \text{otherwise}, \end{cases}$$

 C_{ij} are in Proposition 3.1.

Proof. It is clear by the transformation G,

$$pr^*(R(x)) = (1 - y_1 - y_2)^2,$$

and Proposition 3.1.

Let \widehat{C} be the matrix $(\widehat{C}_{ij})_{1 \leq i,j \leq 4}$. This matrix is symmetric and $\widehat{C}^{\vee} = \widehat{C}$, where \widehat{C}^{\vee} is given by the replacement $\lambda_i \to -\lambda_i$ $(i = 1, \ldots, 4)$ for every entry of \widehat{C} .

Lemma 5.1. The connection matrix $\widehat{\Xi}$ satisfies

$$\widehat{\Xi}^{\vee} = -\widehat{\Xi}, \quad \widehat{\Xi} \, \widehat{C} + \widehat{C}^{\ t} \widehat{\Xi}^{\vee} = O,$$

where $\widehat{\Xi}^{\vee}$ is given by the replacement $\lambda_i \to -\lambda_i$ for $\widehat{\Xi}$.

Proof. We can easily check this lemma by Theorem 5.1 and Corollary 5.2. The second equality is also obtained by Proposition 4.1 from $d_y \hat{C} = O$.

Lemma 5.2. (i) The eigenvalues of $\widehat{\Xi}^1$ are 0 and $-(\lambda_1 + \lambda_3) = 1 - c_1$ and each of the eigenspaces is 2-dimensional. Its $(1 - c_1)$ -eigenspace is spanned by the row vectors

$$e_2 = (0, 1, 0, 0), \quad (\frac{\lambda_{123}}{\lambda_3}, \frac{\lambda_2 - \lambda_4}{\lambda_3}, \frac{\lambda_1 - \lambda_4}{\lambda_3}, 1).$$

(ii) The eigenvalues of $\widehat{\Xi}^2$ are 0 and $-(\lambda_2 + \lambda_3) = 1 - c_2$ and each of the eigenspaces is 2-dimensional. Its $(1 - c_2)$ -eigenspace is spanned by the row vectors

$$e_3 = (0, 0, 1, 0), \quad (\frac{\lambda_{123}}{\lambda_3}, \frac{\lambda_2 - \lambda_4}{\lambda_3}, \frac{\lambda_1 - \lambda_4}{\lambda_3}, 1).$$

20 YOSHIAKI GOTO, JYOICHI KANEKO, AND KEIJI MATSUMOTO

(iii) The eigenvalues of $\widehat{\Xi}^3$ are 0 and $2(\lambda_3 + \lambda_4) = 2(c_1 + c_2 - a - b - 2)$ and the 0-eigenspace of $\widehat{\Xi}^3$ is 3-dimensional and the $2(\lambda_3 + \lambda_4)$ eigenspace of $\widehat{\Xi}^3$ is 1-dimensional. Its $2(c_1 + c_2 - a - b - 2)$ eigenvector is $e_4 = (0, 0, 0, 1)$.

Proof. We can easily check this lemma by Theorem 5.1.

We set

$$e_5 = \left(\frac{\lambda_{123}}{\lambda_3}, \frac{\lambda_2 - \lambda_4}{\lambda_3}, \frac{\lambda_1 - \lambda_4}{\lambda_3}, 1\right), \quad e_6 = \left(\frac{\lambda_{123}}{\lambda_3}, \frac{\lambda_2 - \lambda_4}{\lambda_3}, \frac{\lambda_1 - \lambda_4}{\lambda_3}, -1\right),$$

and

$$\widehat{\varphi}_5 = \frac{\lambda_{123}}{\lambda_3} \widehat{\varphi}_1 + \frac{\lambda_2 - \lambda_4}{\lambda_3} \widehat{\varphi}_2 + \frac{\lambda_1 - \lambda_4}{\lambda_3} \widehat{\varphi}_3 + \widehat{\varphi}_4,$$
$$\widehat{\varphi}_6 = \frac{\lambda_{123}}{\lambda_3} \widehat{\varphi}_1 + \frac{\lambda_2 - \lambda_4}{\lambda_3} \widehat{\varphi}_2 + \frac{\lambda_1 - \lambda_4}{\lambda_3} \widehat{\varphi}_3 - \widehat{\varphi}_4,$$

corresponding to the vectors e_5 and e_6 , respectively.

Lemma 5.3. (i) The forms $\widehat{\varphi}_2$ and $\widehat{\varphi}_5$ vanish as $y_1 \to 0$. The forms $\widehat{\varphi}_2$ and $\widehat{\varphi}_6$ vanish as $y_2 \to 1$.

- (ii) The forms $\widehat{\varphi}_3$ and $\widehat{\varphi}_5$ vanish as $y_2 \to 0$. The forms $\widehat{\varphi}_3$ and $\widehat{\varphi}_6$ vanish as $y_1 \to 1$.
- (iii) The form $\widehat{\varphi}_4$ vanishes as $y_1 + y_2 \rightarrow 1$.

Proof. (i) Since $\widehat{\varphi}_2 = y_1(1-y_2)ds_{12}/(s_2L(s,y))$, it vanishes as $y_1 \to 0$ and as $y_2 \to 1$. We consider the image of $2(\lambda_4 - 1)x_1x_2ds_{12}/L^2$ under the map pr^* . By Lemma 4.2, it is

$$\lambda_{123}\widehat{\varphi}_1 + (\lambda_2 - \lambda_4)\widehat{\varphi}_2 + (\lambda_1 - \lambda_4)\widehat{\varphi}_3 + \frac{\lambda_3(1 - y_1 - y_2 + 2y_1y_2)}{1 - y_1 - y_2}\widehat{\varphi}_4.$$

It is clear that this element vanishes and its last term converges to $\lambda_3 \widehat{\varphi}_4$ (resp. $-\lambda_3 \widehat{\varphi}_4$) as $y_1 \to 0$ (resp. $y_2 \to 1$). Thus $\lambda_3 \widehat{\varphi}_5$ vanishes as $y_1 \to 0$ and $\lambda_3 \widehat{\varphi}_6$ vanishes as $y_2 \to 1$.

(ii) Similarly we can show the statements.

(iii) Since

$$\widehat{\varphi}_4 = \frac{(1 - y_1 - y_2)ds_{12}}{Q(s)L(s, y)},$$

it vanishes as $y_1 + y_2 \rightarrow 1$.

We set

$$\widehat{C}_{1} = \begin{pmatrix} \mathcal{I}_{c}(\widehat{\varphi}_{2}, \widehat{\varphi}_{2}) & \mathcal{I}_{c}(\widehat{\varphi}_{2}, \widehat{\varphi}_{5}) \\ \mathcal{I}_{c}(\widehat{\varphi}_{5}, \widehat{\varphi}_{2}) & \mathcal{I}_{c}(\widehat{\varphi}_{5}, \widehat{\varphi}_{5}) \end{pmatrix} = \begin{pmatrix} \frac{(\lambda_{3} - \lambda_{134}^{-})(\lambda_{1} + \lambda_{3})}{\lambda_{0}\lambda_{2}\lambda_{4}\lambda_{134}^{-}} & \frac{-2(\lambda_{1} + \lambda_{3})}{\lambda_{0}\lambda_{3}\lambda_{4}} \\ \frac{-2(\lambda_{1} + \lambda_{3})}{\lambda_{0}\lambda_{3}\lambda_{4}} & \frac{-4(\lambda_{2} + \lambda_{3})(\lambda_{1} + \lambda_{3})}{\lambda_{0}\lambda_{3}^{2}\lambda_{4}} \end{pmatrix},$$

$$\widehat{C}_{2} = \begin{pmatrix} \mathcal{I}_{c}(\widehat{\varphi}_{3}, \widehat{\varphi}_{3}) & \mathcal{I}_{c}(\widehat{\varphi}_{3}, \widehat{\varphi}_{5}) \\ \mathcal{I}_{c}(\widehat{\varphi}_{5}, \widehat{\varphi}_{3}) & \mathcal{I}_{c}(\widehat{\varphi}_{5}, \widehat{\varphi}_{5}) \end{pmatrix} = \begin{pmatrix} \frac{(\lambda_{3} - \lambda_{234}^{-})(\lambda_{2} + \lambda_{3})}{\lambda_{0}\lambda_{1}\lambda_{4}\lambda_{234}^{-}} & \frac{-2(\lambda_{2} + \lambda_{3})}{\lambda_{0}\lambda_{3}\lambda_{4}} \\ \frac{-2(\lambda_{2} + \lambda_{3})}{\lambda_{0}\lambda_{3}\lambda_{4}} & \frac{-4(\lambda_{2} + \lambda_{3})(\lambda_{1} + \lambda_{3})}{\lambda_{0}\lambda_{3}^{2}\lambda_{4}} \end{pmatrix}.$$

We have

$$\begin{pmatrix} \mathcal{I}_c(\widehat{\varphi}_2, \widehat{\varphi}_2) & \mathcal{I}_c(\widehat{\varphi}_2, \widehat{\varphi}_6) \\ \mathcal{I}_c(\widehat{\varphi}_6, \widehat{\varphi}_2) & \mathcal{I}_c(\widehat{\varphi}_6, \widehat{\varphi}_6) \end{pmatrix} = \widehat{C}_1, \begin{pmatrix} \mathcal{I}_c(\widehat{\varphi}_3, \widehat{\varphi}_3) & \mathcal{I}_c(\widehat{\varphi}_3, \widehat{\varphi}_6) \\ \mathcal{I}_c(\widehat{\varphi}_6, \widehat{\varphi}_3) & \mathcal{I}_c(\widehat{\varphi}_6, \widehat{\varphi}_6) \end{pmatrix} = \widehat{C}_2.$$

Lemma 5.4. The matrix $\widehat{\Xi}^i$ (i = 1, 2, 3) is expressed in terms of \widehat{C} and its eigenvectors with non-zero eigenvalue as

$$\begin{split} \widehat{\Xi}^{1} &= -(\lambda_{1} + \lambda_{3})\widehat{C}(\ {}^{t}e_{2},\ {}^{t}e_{5})(\widehat{C}_{1})^{-1} \left(\begin{array}{c} e_{2} \\ e_{5} \end{array}\right), \\ \widehat{\Xi}^{2} &= -(\lambda_{2} + \lambda_{3})\widehat{C}(\ {}^{t}e_{3},\ {}^{t}e_{5})(\widehat{C}_{2})^{-1} \left(\begin{array}{c} e_{3} \\ e_{5} \end{array}\right), \\ \widehat{\Xi}^{3} &= 2(\lambda_{3} + \lambda_{4})\widehat{C}\ {}^{t}e_{4}(\widehat{C}_{44})^{-1}e_{4}. \end{split}$$

Proof. We claim that

$$v\widehat{C}^{t}w^{\vee} = 0$$

for any eigenvector v of $\widehat{\Xi}^i$ with non-zero eigenvalue α and for any eigenvector w of $\widehat{\Xi}^i$ with eigenvalue 0. Lemma 5.1 implies

$$\widehat{\Xi}^i \widehat{C} = -\widehat{C} \ {}^t (\widehat{\Xi}^i)^{\vee}.$$

Since $\alpha \neq 0$ and

$$\begin{aligned} \alpha(v\widehat{C} \ {}^{t}w^{\vee}) &= (v\widehat{\Xi}^{i})\widehat{C} \ {}^{t}w^{\vee} = v(\widehat{\Xi}^{i}\widehat{C}) \ {}^{t}w^{\vee} = -v(\widehat{C} \ {}^{t}(\widehat{\Xi}^{i})^{\vee}) \ {}^{t}w^{\vee} \\ &= -v\widehat{C} \ {}^{t}(w\widehat{\Xi}^{i})^{\vee} = 0, \end{aligned}$$

we have this claim. Lemma 5.2 together with this claim gives the expression of the matrix $\widehat{\Xi}^i$ in this lemma.

Theorem 5.2. The connection ∇_Y of $pr^*H^2(\Omega^{\bullet,0}(\mathfrak{X}), \nabla)$ is expressed in terms of the intersection form \mathcal{I}_c as

$$\begin{split} \nabla_{Y}(\widehat{\varphi}) &= \frac{dy_{1}}{y_{1}} \wedge (1-c_{1})(\mathcal{I}_{c}(\widehat{\varphi},\widehat{\varphi}_{2}),\mathcal{I}_{c}(\widehat{\varphi},\widehat{\varphi}_{5}))(\widehat{C}_{1})^{-1}\left(\begin{array}{c}\widehat{\varphi}_{2}\\\widehat{\varphi}_{5}\end{array}\right) \\ &+ \frac{dy_{2}}{y_{2}} \wedge (1-c_{2})(\mathcal{I}_{c}(\widehat{\varphi},\widehat{\varphi}_{3}),\mathcal{I}_{c}(\widehat{\varphi},\widehat{\varphi}_{5}))(\widehat{C}_{2})^{-1}\left(\begin{array}{c}\widehat{\varphi}_{3}\\\widehat{\varphi}_{5}\end{array}\right) \\ &+ \frac{dy_{1}}{y_{1}-1} \wedge (1-c_{2})(\mathcal{I}_{c}(\widehat{\varphi},\widehat{\varphi}_{3}),\mathcal{I}_{c}(\widehat{\varphi},\widehat{\varphi}_{6}))(\widehat{C}_{2})^{-1}\left(\begin{array}{c}\widehat{\varphi}_{3}\\\widehat{\varphi}_{6}\end{array}\right) \\ &+ \frac{dy_{2}}{y_{2}-1} \wedge (1-c_{1})(\mathcal{I}_{c}(\widehat{\varphi},\widehat{\varphi}_{2}),\mathcal{I}_{c}(\widehat{\varphi},\widehat{\varphi}_{6}))(\widehat{C}_{1})^{-1}\left(\begin{array}{c}\widehat{\varphi}_{2}\\\widehat{\varphi}_{6}\end{array}\right) \\ &+ \frac{dy_{1}+dy_{2}}{y_{1}+y_{2}-1} \wedge 2(c_{1}+c_{2}-a-b-2)\mathcal{I}_{c}(\widehat{\varphi},\widehat{\varphi}_{4})(\widehat{C}_{44})^{-1}\widehat{\varphi}_{4}. \end{split}$$

Proof. Note that the linear transformation

$$\widehat{\varphi} \mapsto (1 - c_1)(\mathcal{I}_c(\widehat{\varphi}, \widehat{\varphi}_2), \mathcal{I}_c(\widehat{\varphi}, \widehat{\varphi}_5))(\widehat{C}_1)^{-1} \left(\begin{array}{c} \widehat{\varphi}_2\\ \widehat{\varphi}_5 \end{array}\right)$$

is represented matrix $\widehat{\Xi}^1$ with respect to the frame ${}^t(\widehat{\varphi}_1, \ldots, \widehat{\varphi}_4)$. The eigen space of $I_{3,1}\widehat{\Xi}^1I_{3,1}^{-1}$ with non-zero eigenvalue is spanned by e_2 and e_6 , which correspond to $\widehat{\varphi}_2$ and $\widehat{\varphi}_6$, respectively. Thus the linear transformation

$$\widehat{\varphi} \mapsto (1 - c_1)(\mathcal{I}_c(\widehat{\varphi}, \widehat{\varphi}_2), \mathcal{I}_c(\widehat{\varphi}, \widehat{\varphi}_6))(\widehat{C}_1)^{-1} \begin{pmatrix} \widehat{\varphi}_2 \\ \widehat{\varphi}_6 \end{pmatrix}$$

is represented by $I_{3,1}\widehat{\Xi}^1 I_{3,1}^{-1}$ with respect to ${}^t(\widehat{\varphi}_1,\ldots,\widehat{\varphi}_4)$. Similarly, we have representation matrices $\widehat{\Xi}^2$ and $I_{3,1}\widehat{\Xi}^2 I_{3,1}^{-1}$. The linear transformation

$$2(c_1+c_2-a-b-2)\mathcal{I}_c(\widehat{\varphi},\widehat{\varphi}_4)(\widehat{C}_{44})^{-1}\widehat{\varphi}_4$$

is represented by $\widehat{\Xi}^3$ with respect to ${}^t(\widehat{\varphi}_1, \ldots, \widehat{\varphi}_4)$. Theorem 5.1 and Lemma 5.4 yield this theorem.

References

- [ApKa] P. Appell and J. Kampé de Fériet, *Fonctions hypergéométriques et hyper-sphériques: polynomes d'Hermite*, Gauthier-Villars, Paris, 1926.
- [AoKi] K. Aomoto and M. Kita, translated by K. Iohara, *Theory of Hypergeometric Functions*, (Springer Verlag, Now York, 2011).
- [C] K. Cho, A generalization of Kita and Noumi's vanishing theorems of cohomology groups of local system, Nagoya Math. J., 147 (1997), 63–69.

- [G] Y. Goto, The monodromy representation of Lauricella's hypergeometric function F_C , preprint 2014, arXiv:1403.1654.
- [GM] Y. Goto and K. Matsumoto, The monodromy representation and twisted period relations for Appell's hypergeometric function F_4 , to appear in Nagoya Math. J.
- [Kan] J. Kaneko, Monodromy group of Appell's system (F_4) , Tokyo J. Math., 4 (1981), 35–54.
- [Kat] M. Kato, A Pfaffian system of Appell's F_4 , Bull. Coll. Edu. Univ. Ryukyus, **33** (1988), 331 -334.
- [M1] K. Matsumoto, Monodromy and Pfaffian of Lauricella's F_D in terms of the intersection forms of twisted (co)homology groups, Kyushu J. Math. **67** (2013), 367–387.
- [M2] _____, Pfaffian of Lauricella's hypergeometric system F_A , preprint 2015, arXiv:1502.00334.

(Goto) DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, KOBE UNIVERSITY, KOBE 657-8501, JAPAN

E-mail address: y-goto@math.kobe-u.ac.jp

(Kaneko) DEPARTMENT OF MATHEMATICAL SCIENCE, UNIVERSITY OF THE RYUKYUS, NISHIHARA, OKINAWA, 903-0213, JAPAN

E-mail address: kaneko@math.u-ryukyu.ac.jp

(Matsumoto) DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAP-PORO 060-0810, JAPAN

E-mail address: matsu@math.sci.hokudai.ac.jp