## OPTICAL SPECTROSCOPIC OBSERVATIONS OF BLAZARS AND $\gamma$ -RAY BLAZAR CANDIDATES IN THE SLOAN DIGITAL SKY SURVEY DATA RELEASE NINE

F. MASSARO<sup>1</sup>, N. MASETTI<sup>3</sup>, R. D'ABRUSCO<sup>4</sup>, A. PAGGI<sup>4</sup>, & S. FUNK<sup>2</sup>

version October 9, 2018: fm

#### ABSTRACT

We present an analysis of the optical spectra available in the Sloan Digital Sky survey data release nine (SDSS DR9) for the blazars listed in the ROMA-BZCAT and for the  $\gamma$ -ray blazar candidates selected according to their IR colors. First, we adopt a statistical approach based on MonteCarlo simulations to find the optical counterparts of the blazarslisted in the ROMA-BZCAT catalog. Then we crossmatched the SDSS spectroscopic catalog with our selected samples of blazars and  $\gamma$ -ray blazar candidates searching for those with optical spectra available to classify our blazar-like sources and, whenever possible, to confirm their redshifts. Our main objectives are determining the classification of uncertain blazars listed in the ROMA-BZCAT and discovering new gamma-ray blazars. For the ROMA-BZCAT sources we investigated a sample of 84 blazars confirming the classification for 20 of them and obtaining 18 new redshift estimates. For the  $\gamma$ -ray blazars, indicated as potential counterparts of unassociated *Fermi* sources or with uncertain nature, we established the blazar-like nature of 8 out the 27 sources analyzed and confirmed 14 classifications.

Subject headings: methods: statistical - galaxies: active - quasars: general - surveys - radiation mechanisms: non-thermal

#### 1. INTRODUCTION

According to the well assessed unification scenario of the active galactic nuclei (AGN; e.g., Antonucci 1993; Urry & Padovani 1995) blazars are radio loud sources, featuring compact radio cores combined with a "flat" radio spectra that extends from frequencies below  $\sim 1$ GHz (e.g., Massaro et al. 2013a; Massaro et al. 2013b; Nori et al. 2013) up to the submillimeter band (e.g., Giommi et al. 2012). They are characterized by a variable, non-thermal, continuum and exhibit a typical double bumped spectral energy distribution (SED), and represent the largest known population of  $\gamma$ -ray sources (e.g., Abdo et al. 2010a; Ackermann et al. 2011) proving the most relevant contribution to the extragalactic  $\gamma$ -ray background (e.g., Mukherjee et al. 1997; Abdo et al. 2010b).

Blazars are generally classified on the basis of their optical spectra and divided in two main classes: i) BL Lac objects, labeled as BZBs according to the nomenclature of the ROMA-BZCAT<sup>5</sup> (Massaro et al. 2009; Massaro et al. 2011a) when presenting featureless optical spectra and ii) flat spectrum radio quasars (hereinafter BZQs) having a typical quasar-like optical appearance but also featuring high and variable optical polarization. In particular, blazars are classified as BZB if the rest-frame equivalent width of their optical features is lower than 5 Å (Stickel et al. 1991; Stoke et al. 1991;

<sup>5</sup> http://www.asdc.asi.it/bzcat/

Laurent-Muehleisen et al. 1999; Landoni et al. 2013).

As recently discovered using the WISE all-sky survey (Wright et al. 2010), blazars show by peculiar infrared (IR) colors (Massaro et al. 2011b) mostly due to their non-thermal continuum that allowed to distinguish them from other classes of active galaxies (e.g., D'Abrusco et al. 2012; Massaro et al. 2012a). This IR property was also interpreted as due to the lack of observational signatures form a dusty torus in the case of BZBs (e.g., Plotkin et al. 2012).

The variable, non-thermal emission of both BZBs and BZQs, extending from radio up to TeV energies, is interpreted as arising from high-energy particles accelerated in a relativistic jet oriented along to the line of sight, whereas relativistic effects amplifies both their luminosity and the amplitude of their variability (Blandford & Rees 1978; Giommi et al. 2013).

Recently, we searched for blazar-like objects as potential counterparts of the unidentified  $\gamma$ -ray sources (UGSs) observed with *Fermi* (Abdo et al. 2010a; Nolan et al. 2012) with several methods based on the IR colors alone (Massaro et al. 2012b; D'Abrusco et al. 2013) or combined with other multifrequency observations, as radio (Massaro et al. 2013d) or X-ray properties (Paggi et al. 2013). We also explored the use of low radio frequency observations (i.e., below  $\sim 1$  GHz) as an alternative possibility to find blazar-like counterparts (e.g., Massaro et al. 2013a; Nori et al. 2013) for the UGSs listed in the second *Fermi*-Large Area Telescope (LAT) catalog (2FGL, Nolan et al. 2012) in addition to other multifrequency analysis (e.g., Mirabal & Halpern 2009: Ackermann et al. 2012: Cowperthwaite et al. 2013; Masetti et al. 2013). A11 these investigations provided several lists of gamma-ray blazar candidates that has to be confirmed and classified via optical spectroscopy.

Here we investigate the optical spectra of two

<sup>&</sup>lt;sup>1</sup> Yale Center for Astronomy and Astrophysics, Physics Department, Yale University, PO Box 208120, New Haven, CT 06520-8120, USA

<sup>&</sup>lt;sup>2</sup> SLAC National Laboratory and Kavli Institute for Particle Astrophysics and Cosmology, 2575 Sand Hill Road, Menlo Park, CA 94025, USA $^{3}$ INAF - Istituto di Astrofisica Spaziale e Fisica Cosmica di

Bologna, via Gobetti 101, 40129, Bologna, Italy

Harvard - Smithsonian Astrophysical Observatory, 60 Garden Street, Cambridge, MA 02138, USA

blazar samples that lie in the footprint of the spectroscopic catalog of Sloan Digital Sky Survey data release 9 (SDSS DR9, Ahn et al. 2012). The first sample includes all the ROMA-BZCAT sources that have an uncertain classification, uncertain redshift estimates or have been classified as BL Lac candidates due to the lack of an optical spectrum in literature (Massaro et al. 2011a). The second sample lists all the  $\gamma$ -ray blazar candidates, that were identified as potential counterparts of UGSs in our previous analyses, and for which SDSS spectra are now available. This study is complementary to on going spectroscopic campaigns planned to investigate blazar optical properties (e.g., Sbarufatti et al. 2005; Sbarufatti et al. 2009; Plotkin et al. 2010; Paggi et al. 2014).

The paper is organized as follows. In Section 2 we present the statistical approach adopted to determine the optical counterparts of the ROMA-BZCAT sources in the SDSS DR9 catalog while in Section 3 we describe the samples considered in our analysis. In Section 4 the results of the spectroscopic analysis are illustrated while, finally, Section 5 is devoted to our summary and conclusions. For our numerical results, we use cgs units unless stated otherwise.

#### 2. SPATIAL ASSOCIATIONS

The ROMA-BZCAT was mainly compiled on the basis of radio, optical and X-ray surveys and the blazar coordinates reported are not uniform. The positional accuracy is generally less than <1'' but it could reach  $\sim 5''$ , corresponding to the typical uncertainty on the radio positions of the NVSS, for those sources with radio flux densities close to the survey limit (Condon et al. 1998). Since the positional uncertainties for each source are not reported in the ROMA-BZCAT to identify the SDSS optical counterparts of the ROMA-BZCAT blazars we adopted the following a statistical approach.

First we computed the total number of ROMA-BZCAT blazars that lie within the footprint of the SDSS, corresponding to 1820 sources. For each blazar, we counted the total number of optical counterparts in the SDSS N(R) present within circular regions of variable radius R in the range between 0" and 10". To be conservative in our analysis, we only included in the N(R) calculation SDSS sources having the flags: CLASS\_OBJECT (i.e., mode) and CODE\_MISC (i.e., clean) both equal to 1<sup>6</sup>.

We then created 100 mock realizations of the ROMA-BZCAT by shifting each blazar position in a random direction of the sky by a fixed length of 30''. The shift used to create the mock ROMA-BZCAT catalogs were chosen not too distant from the original ROMA-BZCAT location and within the SDSS footprint. This guarantees to obtain fake catalogs with a sky distribution similar to the original ROMA-BZCAT and to crossmatch each fake catalog and the SDSS taking into account the local density distribution of the optical sources. The total number of blazars in each mock realization is also preserved being equal to that of the ROMA-BZCAT sources that lie in the SDSS footprint. For each mock realization of the ROMA-BZCAT we counted the number of associations with the SDSS occurring at angular separations R smaller than 10". Then we computed the mean num-



FIG. 1.— The values of  $\Delta \lambda(R)$  (red circles) and  $\Delta N(R)$  (black squares) as function of the angular separation R. We restricted the x axis to R values below 10". Our choice of  $R_A$  is marked by the vertical dashed line. It occurs at the first R value for which  $\Delta \lambda(R) \simeq \Delta N(R)$ .

ber  $\lambda(R)$  of these fake associations, averaged over the 100 mock ROMA-BZCAT catalogs, verifying that  $\lambda(R)$  has a Poissonian distribution. Increasing the radius by  $\Delta R = 0''.2$ , we also calculated the difference  $\Delta \lambda(R)$  as:

$$\Delta \lambda(R) = \lambda(R + \Delta R) - \lambda(R), \qquad (1)$$

In Figure 1 we show the comparison between  $\Delta N(R)$ and  $\Delta \lambda(R)$ . For radii larger than  $R_A = 1''.8$  the  $\Delta \lambda(R)$  curve superimposes that of  $\Delta N(R)$  indicating that ROMA-BZCAT-SDSS cross-matches could occur by chance at angular separations larger than  $R_A$ . Thus we choose 1''.8 as to the maximum angular separation between the ROMA-BZCAT and the SDSS position at which we consider the optical source a reliable counterpart of the blazar in the ROMA-BZCAT.

Finally the chance probability of spurious associations  $p(R_A)$  was calculated as the ratio between the number of real associations  $N(R_A)$  and the average of those found in the mock realizations of the ROMA-BZCAT  $\lambda(R_A)$ , corresponding to a value of ~1% (see also Maselli et al. 2010; Massaro et al. 2011b; D'Abrusco et al. 2013, for additional details on  $p(R_A)$ ).

### 3. SAMPLE SELECTION

We adopted the value of  $R_A$  of 1".8 to search for the optical counterparts of the blazar-like sources in our two samples within the available spectra of the SDSS DR9 to confirm the source nature and whenever possible to estimate the redshift. Then we analyzed two samples of sources that lie in the footprint of the SDSS dr9 and with optical spectra available defined in the following.

1. The total number of ROMA-BZCAT sources having a optical counterpart in the SDSS DR9 spectroscopic catalog within  $R_A$  is 219. Among them there are 50 blazars with an uncertain redshift estimate and additional 34 sources classified as BL Lac candidates for which optical spectra were not available in literature while ROMA-BZCAT v4.1

<sup>6</sup> http://cas.sdss.org/dr7/sp/help/browser/browser.asp?n=PhotoObj&t=U was prepared. These 84 ROMA-BZCAT sources,

with a unique correspondence in the SDSS DR9 spectroscopic catalog within 1''.8, constitute our first sample investigated.

2. The second sample lists 15 blazar-like sources with uncertain classification and/or uncertain z estimates included in the 2FGL (Nolan et al. 2012) and in the Second *Fermi*-LAT AGN Catalog (2LAC Ackermann et al. 2011) plus additional 12 sources identified as potential counterparts of UGSs according to their peculiar IR colors in our recent analyses (e.g., Massaro et al. 2012a; Massaro et al. 2012b; Massaro et al. 2013c). All these 27 sources also have a unique correspondence in the SDSS DR9 spectroscopic catalog within 1".8.

#### 4. RESULTS

We visually inspected all the optical spectra available for the sources in our samples to avoid misleading classifications due to artifacts of the SDSS automatic analysis, and if necessary, for example to confirm a redshift estimate, we downloaded and analyzed the raw data.

We remark that for the BZB classification we adopted the criterion described in Laurent-Muehleisen et al. (1999), measuring the rest-frame equivalent widths of the emission and/or absorption lines whenever they are detectable above the continuum (see also the recent analyses performed by Sbarufatti et al. 2006; Landoni et al. 2013). In addition, we also adopted the criterion developed by Massaro et al. (2013c) to classify BL Lac object based on the SDSS (u-r) color, that supersedes the one based on the Ca H&K break contrast originally introduced by Stoke et al. (1991). Thus, for each source, we computed the absorption corrected (ur) color equal to  $(u-r)_{obs} - 0.81 * \bar{A}_r$ , where  $A_r$  is the Galactic extinction in the R band, and we considered BL Lac objects only those sources with (u-r) < 1.4 (see also Maselli et al. 2013). We assign a BZB classification only to sources having both a "featureless" spectra and the (u-r) color lower than 1.4.

### 4.1. ROMA-BZCAT blazars

In the first sample of 84 ROMA-BZCAT blazars with SDSS DR9 spectra, we found that there are 3 BZQs, all associated to *Fermi* sources in the 2FGL and in the 2LAC (Nolan et al. 2012; Ackermann et al. 2011), with an uncertain z estimate. Our analysis of their optical SDSS spectra allowed us to confirm both their nature and their redshifts. In addition to these BZQs, there are 47 sources out of the 84 listed in the first sample classified as BZB according to the ROMA-BZCAT but with an uncertain redshift estimate. We found that 9 of them have good SDSS spectra from which we obtained a z measurement. Unfortunately none of these 9 is detected in the  $\gamma$ -rays.

The remaining 34 sources out of 84 objects in the first sample are indeed classified as BL Lac candidates, 5 of them being associated to *Fermi* sources in the 2LAC. We confirmed the BL Lac nature for 20 of them, including all the *Fermi* sources and providing a new redshift estimate for 6. The remaining 14 sources were classified as: normal galaxies (8), type 2 Seyfert galaxies (5), according to the criteria described in Winkler (1992) plus 1 source having still an uncertain nature, mostly resembling a type 2 AGN.

All our results are summarized in Table 2, where we report the ROMA-BZCAT and the SDSS names, together with the results of our analysis (i.e., classification and redshift estimates when possible) and their (u-r) colors. Blazars that are associated to *Fermi* sources are also indicated. Uncertain values of redshifts are indicated with a question mark (?); they are due to the poor signal to noise of few SDSS archival spectra or to the presence of only a single emission/absorption feature. Then, in Figure 2 we show the optical spectrum of one of the BL Lac classified from our analysis together with two cases of wrong classifications and a quasar.

### 4.2. $\gamma$ -ray blazar candidates

The second sample of  $\gamma$ -ray blazar candidates selected according to our IR based procedures and having SDSS spectra available lists: 27 sources. Fifteen blazar-like sources were already present in the 2LAC but with uncertain classification or uncertain redshift estimates. Among them we found 7 having quasar-like optical spectra and being classified as BZQs, 2 also with new z estimates, 7 BZBs including 2 sources with measured redshifts and 1 misclassified object: SDSS J122011.88+020342.2, associated with 2FGLJ1219.7+0201 that appears to be a Seyfert galaxy rather than BZQ.

For 2FGLJ1023.6+3947, associated to the SDSS J102333.50+395312.7 source, the we obtained a redshift of 1.3328 instead of 1.254 reported in the 2LAC and for the BL Lac object SDSS J110021.05+401928.0 counterpart of 2FGLJ1100.9+4014, we were not able to find any optical feature to confirm the 2LAC redshift of 0.225. Among these 15 sources there is SDSS J222329.57+010226.6, associated to the AGN of uncertain type 2FGLJ223.4+0104 and selected in Cowperthwaite et al. (2013) as a  $\gamma$ -ray blazar candidate that we confirmed as a BL Lac at unknown redshift.

The remaining 12 sources were all selected as  $\gamma$ -ray blazar candidates in our previous analyses of their IR colors (Massaro et al. 2012b; Massaro et al. 2013c; Massaro et al. 2013a; Paggi et al. 2013). We found that 3 sources, all with new z estimates, out of 12 have a quasar like spectrum, similar to those of the BZQs, plus one uncertain due to noisy SDSS spectrum (i.e., SDSS J015852.77+010132.8). Then there are 4 confirmed BL Lac objects while the remaining 4 sources are indeed contaminants of the association methods (1 star and 4 Seyfert galaxies).

All these results are reported in Table 3 in the same order as discussed above, where we also indicate the previous classification of each source.

#### 5. SUMMARY AND CONCLUSIONS

We performed an analysis of the archival optical spectra present in the SDSS DR9 (Ahn et al. 2012) for two selected samples of blazars and  $\gamma$ -ray blazar candidates to confirm their nature and whenever possible to estimate their redshifts.

First, we adopted a statical approach to find the the SDSS optical cross-matches of the sources in our sample. Then, we analyzed a first sample of 84 blazars listed in the ROMA-BZCAT as BL Lac candidates or

as BL Lac objects and flat spectrum radio quasars with uncertain redshift estimates and a second sample of 27  $\gamma$ -ray blazar candidates selected according to their peculiar IR colors or with uncertain classification (e.g., D'Abrusco et al. 2013; Massaro et al. 2013c; Massaro et al. 2013d, and references therein).

On the basis of the SDSS spectra, we confirmed the redshifts for 3 flat spectrum radio quasars (all  $\gamma$ -ray sources detected by *Fermi*) and measured the z for 9 additional BL Lacs investigated. Then, we have been able to classify 34 BL Lac candidates listed in the ROMA-BZCAT, 20 of them appearing as BL Lac objects, providing new z estimates for 6 BL Lac objects. These spectroscopic information, even if available for a small fraction of the whole ROMA-BZCAT catalog will be essential to refine its future releases as well as those of the *Fermi* catalogs.

For the second sample listing 27  $\gamma$ -ray blazar candidates we found a total of 11 BZBs (2 with new z measurements) and 11 BZQs having new redshift estimates for 2 of them. The remaining 5 sources did not appear to have the typical blazar-like optical spectrum. All our results are summarized in Table 4.

We thank our anonymous referee for many helpful comments which greatly improved this manuscript. We are grateful to our colleague E. Palazzi that helped us in the data reduction of some SDSS spectra. The work is supported by the NASA grants NNX12AO97G and NNX13AP20G. Part of this work is based on archival data, software or on-line services provided by the ASI Science Data Center. This research has made use of data obtained from the high-energy Astrophysics Science Archive Research Center (HEASARC) provided by NASA's Goddard Space Flight Center; the SIM-BAD database operated at CDS, Strasbourg, France; the NASA/IPAC Extragalactic Database (NED) operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration. TOP- $CAT^7$  (Taylor 2005) for the preparation and manipulation of the tabular data and the images. Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is http://www.sdss.org/. The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

### REFERENCES

Abdo, A. A. et al. 2010a ApJS 188 405 Abdo, A. A. et al. 2010b ApJ, 720, 435 Ackermann, M. et al. 2011a ApJ, 743, 171 Ackermann, M. et al. 2012 ApJ, 753, 83 Ahn et al. 2012, ApJS, 203, 21 Antonucci, R. 1993 ARA&A, 31, 473 Blandford, R.D. & Rees, M. J. 1978, in Proc. Pittsburgh Conf. on BL Lac objects, (Pittsburgh, PA: Univ. Pittsburgh), 328 Cowperthwaite, P. S. et al. 2013 AJ, 146, 110 Condon, J. J. et al. 1998, AJ, 115, 1693 D'Abrusco, R. et al. 2012 ApJ, 748, 68 D'Abrusco, R. et al. 2013 ApJS, 206, 12 Giommi, P. et al. 2012 A&A, 541A, 160 Giommi et al. 2013 MNRAS, 431, 1914 Landoni, M. et al. 2013 AJ, 145, 114 Laurent-Muehleisen, S.A. et al., 1999, ApJ, 525, 127 Maselli, A. et al. 2010 A&A, 512A, 74 Maselli, A. et al. 2013 ApJS, 206, 17 Masetti, N. et al. 2013 A&A, 559, A58 Massaro, E. et al. 2009 A&A, 495, 691 Massaro, E. et al. 2011a "Multifrequency Catalogue of Blazars (3rd Edition)", ARACNE Editrice, Rome, Italy Massaro, F. et al. 2011b ApJ, 740L, 48 Massaro, F. et al. 2012a ApJ, 750, 138

Massaro, F. et al. 2012b ApJ, 752, 61 Massaro, E. et al. 2012 MNRAS, 422, 2322 Massaro, F. et al. 2013a ApJS, 207, 4 Massaro, F. et al. 2013b ApJS, 208, 15 Massaro, F. et al. 2013c ApJS, 206, 13 Massaro, F. et al. 2013d ApJS, 209, 10 Massaro, F. et al. 2013e ApJS, 207, 16 Mirabal, N. 2009 ApJ, 701, 129 Mukherjee, R. et al., 1997 ApJ, 490, 116 Nolan et al. 2012 ApJS, 199, 31 Nori, M. et al. 2014 ApJS, 212, 3 Paggi, A. et al. 2013 ApJS, 209, 9 Paggi, A. et al. 2014 AJ, 147, 112 Plotkin, R. M. 2010 AJ, 139, 390 Plotkin, R. M. et al. 2012 ApJ, 745L, 27 Sbarufatti et al. 2005, AJ, 129, 559 Sbarufatti et al. 2006, AJ, 132, 1 Sbarufatti, B. et al. 2009 AJ, 137, 337 Stickel, M. et al. 1991 ApJ, 374, 431 Stocke et al. 1991, ApJS, 76, 813 Taylor, M. B. 2005, ASP Conf. Ser., 347, 29 Urry, C. M., & Padovani, P. 1995, PASP, 107, 803 Winkler H. 1992 MNRAS, 257, 677 Wright, E. L. et al. 2010 AJ, 140, 1868

<sup>&</sup>lt;sup>7</sup> http://www.star.bris.ac.uk/~mbt/topcat/



FIG. 2.— Upper left panel) The BL Lac object SDSSJ103220.28+030949.2 at redshift 0.3232. Upper right panel) The Seyfert 2 galaxy SDSSJ002608.37-000547.0 previously misclassified as a BL Lac candidate Lower left panel) The normal galaxy SDSSJ122604.12+260427.9 spectrum at redshift z=0.176. Lower right panel) The z=1.398 quasar SDSSJ132700.86+221050.1. The man in spectral emission and/or absorption features are marked in each figure.

# F. Massaro et al. 2014

TABLE 1 Association of ROMA-BZCAT sources.

DZCAM	apaa	DZCAT	apaa	DZCAT	apaa	
BZCAT	SDSS	BZCAT	SDSS	BZCAT	SDSS	u-r
name	counterpart	class	class	redshift	redshift	
$BZQJ0310 + 3814^{*}$	J031049.87 + 381453.8	FSRQ	QSO	0.816?	0.816	1.02
$BZQJ0830 + 2410^{*}$	J083052.08 + 241059.8	FSRQ	QSO	0.939?	0.939	0.42
$BZQJ1327 + 2210^{*}$	J132700.86 + 221050.1	FSRQ	QSO	1.4?	1.398	0.44
BZBJ0001-0011	J000121.46-001140.3	BL Lac	BL Lac	0.462?	0.462	0.67
BZBJ0100-0055	J010058.19-005547.7	BL Lac	BL Lac	0.67 > ?	?	0.98
BZBJ0127-0821	J012716.31-082128.8	BL Lac	BL Lac	0.36?	0.75?	1.06
BZBJ0141-0928*	J014125.83-092843.7	BL Lac	BL Lac	0.73?	0.03?	0.96
BZBJ0731+2804	J073152.73+280432.9	BL Lac	BL Lac	0.25?	0.248	1.2
BZBJ0755+3726	J075523.11+372618.7	BL Lac	BL Lac	0.606?	?	0.9
BZBJ0801+1336	J080115.01 + 133642.2	BL Lac	BL Lac	1.042?	?	0.77
BZBJ0818+4222*	J081815.99 + 422245.4	BL Lac	BL Lac	0.53??	?	0.98
BZBJ0823+2223	$1082324.75 \pm 222303.2$	BL Lac	BL Lac	0.951?	?	0.87
BZBJ0840+3440	J084013.03 + 344026.9	BL Lac	BL Lac	0.451?	?	1.07
$BZB10856 \pm 2057^*$	$108563974 \pm 2057433$	BL Lac	BL Lac	0.18?	0.2?	0.83
BZBJ0926+5411	1092638 87+541126 5	BL Lac	BL Lac	0.841?	0.8?	0.81
BZB10940±2603	1094014 72+260330 0	BL Lac	BL Lac	0.498??	2	1 11
BZBJ0940+2003	1005127 81   010210 2	BL Lac	BL Lac	0.49811	2	0.75
BZBJ0991+0102	1100656 46 245445 1	BL Lac	BL Lac	0.5021		0.75
BZBJ1000+3434	1101010 04 000057 1	BL Lac	BL Lac	0.6121		0.49
BZBJ1012+0630	J101213.34+063057.1	BL Lac	BL Lac	0.5187	: 2	0.69
BZBJ1031+5053	J103118.51+505335.8	BL Lac	BL Lac	0.36177	2 2222	0.45
BZBJ1032+0309	J103220.28+030949.2	BL Lac	BL Lac	0.323?	0.3232	0.64
BZBJ1100+4019*	J110021.05+401928.0	BL Lac	BL Lac	0.225?	?	-0.02
BZBJ1107+5010	J110704.78+501037.9	BL Lac	BL Lac	0.706?	0.7062	0.99
BZBJ1117+2548	J111740.39 + 254846.5	BL Lac	BL Lac	0.36?	?	0.63
BZBJ1120+4212*	J112048.06+421212.4	BL Lac	BL Lac	0.124??	?	0.34
BZBJ1132+0034*	J113245.62+003427.7	BL Lac	BL Lac	1.223??	?	0.89
BZBJ1136+1601	J113617.53+160152.2	BL Lac	BL Lac	0.574?	0.5734	0.8
BZBJ1138+4113	J113812.15+411352.8	BL Lac	BL Lac	0.574?	0.5740?	1.1
BZBJ1211+2242	J121158.63+224232.9	BL Lac	BL Lac	0.455?	0.4527	0.53
BZBJ1219+0446	J121944.97 + 044622.4	BL Lac	BL Lac	0.489?	?	0.72
$BZBJ1231+2847^{*}$	J123143.57 + 284749.7	BL Lac	BL Lac	0.236?	?	0.68
BZBJ1237+3020	J123705.61 + 302005.1	BL Lac	BL Lac	?	?	0.36
BZBJ1238+4431	J123826.01 + 443137.1	BL Lac	BL Lac	0.312?	0.3121	1.34
BZBJ1239+4132	J123922.73 + 413251.4	BL Lac	BL Lac	0.16??	?	0.53
BZBJ1247+4423	J124700.72+442318.8	BL Lac	BL Lac	0.6?	0.73?	0.51
BZBJ1255 + 3848	J125555.40 + 384811.3	BL Lac	BL Lac	0.559?	?	0.81
BZBJ1328+1145	J132833.56 + 114520.5	BL Lac	BL Lac	0.49?	?	-0.06
BZBJ1401+3611	J140138.72+361121.9	BL Lac	BL Lac	0.507?	0.5064?	1.21
BZBJ1404+0402*	J140450.90 + 040202.1	BL Lac	BL Lac	0.344??	?	0.45
BZBJ1423+1412	J142330.67+141247.9	BL Lac	BL Lac	0.769??	0.7687?	0.6
BZBJ1436+4129	J143627.16+412932.3	BL Lac	BL Lac	0.404?	?	2.74
BZBJ1436+5639*	J143657.72+563924.8	BL Lac	BL Lac	0.15?	?	0.47
BZBJ1443+2515	J144334.40 + 251558.2	BL Lac	BL Lac	0.529??	0.5295	0.8
BZBJ1456+5048	J145603.64+504825.9	BL Lac	BL Lac	0.479?	?	0.52
BZBJ1506+0814*	J150644.47+081400.6	BL Lac	BL Lac	0.376?	?	0.52
BZBJ1553+0601	J155331.06 + 060143.8	BL Lac	BL Lac	0.619?	0.6189	0.98
BZBJ1603+1105	J160341.93+110548.7	BL Lac	BL Lac	0.143??	?	0.94
BZBJ1623+2841	$J_{162332,25+284128,7}$	BL Lac	BL Lac	0.377?	?	0.46
BZBJ1652+3632	$1165248 44 \pm 363212 5$	BL Lac	BL Lac	0.648?	0.6470?	0.96
BZBJ1701+3954	J170124.63+395437 0	BL Lac	BL Lac	0.507?	?	0.27
		100				
ol. (1) ROMA-BZC Col. (2) SDSS name	of the optical counterpar	·t.				
Col. (3) ROMA-BZC	AT classification.					
ol. (4) SDSS spectr	oscopic classification bas	sed on our	analysis.			
ol. (5) Redshift est	imate reported in the RC	MA-BZCA	T. Questio	n mark ind	licates unce	rtain es
ol. (b) Redshift est	imate derived from our a lor.	.naiysis.Qu	estion marl	(indicates	uncertain e	stimate
(1) SDSS u=r CO	101.					

	Association of ROMA-BZCAT sources.					
SDSS	BZCAT	SDSS	BZCAT	SDSS	u-r	Classification
counterpart	class	class	redshift	redshift		flag
J002608.37-000547.0	BL Lac Can.	Sy2	0.107	0.106	2.16	no
J010908.17 + 181607.5	BL Lac Can.	BL Lac	0.145	?	0.7	yes
J025315.60-012405.3	BL Lac Can.	BL Lac	?	?	0.67	yes
J075445.66 + 482350.7	BL Lac Can.	BL Lac	?	?	0.93	yes

TABLE 2

Į	name	counterpart	class	class	redshift	redshift		flag
	BZBJ0026-0005	J002608.37-000547.0	BL Lac Can.	Sy2	0.107	0.106	2.16	no
	BZBJ0109+1816*	J010908.17+181607.5	BL Lac Can.	BL Lac	0.145	?	0.7	yes
	BZBJ0253-0124	J025315.60-012405.3	BL Lac Can.	BL Lac	?	?	0.67	yes
	BZBJ0754+4823*	J075445.66 + 482350.7	BL Lac Can.	BL Lac	?	?	0.93	yes
	BZBJ0814 + 0856	J081421.66 + 085706.1	BL Lac Can.	galaxy	0.23??	0.24	7.33	no
	BZBJ0829+1754	J082904.82+175415.8	BL Lac Can.	Sy2	0.089	0.0895	2.12	no
	BZBJ0831 + 5400	J083100.36 + 540023.2	BL Lac Can.	Sy2	?	0.0617	3.06	no
	BZBJ0839+4015	J083903.08 + 401545.6	BL Lac Can.	galaxy	0.194	0.1941	2.43	no
	BZBJ0905 + 4705	J090536.44 + 470546.3	BL Lac Can.	type2	0.174	0.1736	2.33	no
	BZBJ0912 + 4235	J091227.22 + 423545.1	BL Lac Can.	galaxy	0.266	0.2662	4.0	no
	BZBJ0933+0003	J093310.57 + 000323.5	BL Lac Can.	BL Lac	?	?	0.45	yes
	BZBJ0944 + 5557	J094441.47 + 555752.9	BL Lac Can.	BL Lac	?	?	0.93	yes
	BZBJ1007 + 5023	J100710.44 + 502356.4	BL Lac Can.	Sy2	0.133	0.1326	1.87	no
	BZBJ1057 + 2303	J105723.09 + 230318.7	BL Lac Can.	BL Lac	0.379	0.3782	1.07	yes
	BZBJ1058 + 2817	J105829.89 + 281746.3	BL Lac Can.	BL Lac	?	0.4793?	0.89	yes
	BZBJ1100 + 4210	J110020.99 + 421053.1	BL Lac Can.	galaxy	0.323	0.3229	1.75	no
	BZBJ1110 + 3539	J111056.83 + 353907.2	BL Lac Can.	BL Lac	?	0.61?	0.72	yes
	BZBJ1111 + 3452	J111130.90 + 345203.2	BL Lac Can.	BL Lac	0.212	?	0.42	yes
	BZBJ1152 + 2837	J115210.70+283721.3	BL Lac Can.	BL Lac	?	0.4412	1.17	yes
	BZBJ1153+3823	J115210.70+283721.3	BL Lac Can.	Sy2	?	0.4098	0.99	no
	BZBJ1224+2239	J122401.03 + 223939.5	BL Lac Can.	BL Lac	?	0.4821	0.88	yes
	BZBJ1226 + 2604	J122604.12 + 260427.9	BL Lac Can.	galaxy	0.176	0.1761	2.0	no
	BZBJ1243+3627*	J124312.73+362743.9	BL Lac Can.	BL Lac	?	?	0.45	yes
	BZBJ1253+3826	J125300.95 + 382625.7	BL Lac Can.	BL Lac	0.372	0.3707	1.08	yes
	BZBJ1311 + 0853	J131155.76 + 085340.9	BL Lac Can.	BL Lac	0.469	0.4694	0.85	yes
	$BZBJ1314 + 2348^{*}$	J131443.80 + 234826.7	BL Lac Can.	BL Lac	?	0.15?	0.72	yes
	BZBJ1341+3716	J134138.66 + 371644.8	BL Lac Can.	galaxy	0.17	0.1745	2.93	no
	BZBJ1404 + 2701	J140436.82 + 270141.0	BL Lac Can.	galaxy	0.136	0.1383	2.67	no
	BZBJ1410+2820*	J141029.56 + 282055.6	BL Lac Can.	BL Lac	?	?	0.58	yes
	BZBJ1426 + 2415	J142645.52 + 241523.0	BL Lac Can.	BL Lac	0.055??	0.36?	0.4	yes
	BZBJ1437 + 4717	J143716.14 + 471726.3	BL Lac Can.	BL Lac	?	?	0.68	yes
	BZBJ2129 + 0035	$J212940.67 {+} 003527.4$	BL Lac Can.	BL Lac	0.425	0.4264	-0.1	yes
	BZBJ2227 + 0037	J222758.13 + 003705.4	BL Lac Can.	BL Lac	?	?	0.76	yes
	BZBJ2319-0116	J231952.83-011626.8	BL Lac Can.	galaxy	?	0.2835	1.72	no

BZCAT

Col. (1) ROMA-BZCAT name. Col. (2) SDSS name of the optical counterpart. Col. (3) ROMA-BZCAT classification. Col. (4) SDSS spectroscopic classification based on our analysis. Col. (6) Redshift estimate derived from our analysis.Question mark indicates uncertain estimates. Col. (7) u-r color . Col. (8) Classification flag: (yes) marks sources that have been classified on the basis of our analysis.

TABLE 3 Association of  $\gamma$ -ray blazar candidates (00-24 HH).

Source	SDSS	SDSS	SDSS	u-r	Classification
name	counterpart	class	redshift		flag
2FGLJ0323.6-0108	J032343.62-011146.1	BL Lac	?	0.58	yes
2FGLJ0924.0+2819	J092351.52 + 281525.1	QSO	0.7442	0.57	yes
2FGLJ0950.1 + 4554	J095011.82 + 455320.0	BL Lac	0.3994	0.99	yes
2FGLJ1017.0+3531	J101810.97 + 354239.4	QSO	1.2280	0.42	yes
2FGLJ1023.6+3947	J102333.50 + 395312.7	QSO	1.3328	0.23	yes
2FGLJ1100.9+4014	J110021.05 + 401928.0	BL Lac	?	0.43	yes
2FGLJ1219.7+0201	J122011.88 + 020342.2	Sy1.8	0.2402	-0.02	no
2FGLJ1222.4+0413	J122222.54 + 041315.7	QSO	0.9642	0.44	yes
2FGLJ1301.6+3331	J130129.15+333700.3	QSO	1.0084	0.75	yes
2FGLJ1310.9+0036	J131106.47 + 003510.0	BL Lac	?	0.62	yes
2FGLJ1351.4+1115	J135120.84 + 111453.0	BL Lac	?	0.56	yes
2FGLJ1332.7+4725	J133245.24 + 472222.6	QSO	0.6687	0.74	yes
2FGLJ1442.0+4352	J144207.15 + 434836.7	BL Lac	?	0.78	yes
2FGLJ1522.0+4348	J152149.61 + 433639.2	QSO	2.1677	0.35	yes
2FGLJ2223.4+0104+	J222329.57+010226.6	BL Lac	0.29?	0.49	yes
2FGLJ0158.4+0107	J015852.77+010132.8	QSO?	1.61?	0.95	yes
2FGLJ0440.5+2554	J043947.48 + 260140.8	star	0.	-1.56	no
2FGLJ0823.0+4041	J082257.55 + 404149.7	QSO	0.8655	0.58	yes
1FGLJ0835.4+0936	J083543.21+093717.9	BL Lac	0.35?	0.79	yes
2FGLJ0844.9+6214	J084406.82 + 621458.4	Sy1.9	0.1208	3.13	no
2FGLJ1129.5+3758	J112903.24+375656.7	BL Lac	?	1.14	yes
2FGLJ1209.6+4121	J120922.78+411941.3	BL Lac	?	0.75	yes
1FGLJ1422.7+3743	J142304.61+373730.5	BL Lac	?	0.88	yes
2FGLJ1612.0+1403	J161118.10 + 140328.7	QSO	0.5855	0.15	yes
2FGLJ1614.8+4703	J161541.21 + 471111.7	Sy2	0.1986	1.31	no
2FGLJ1627.8+3219	J162800.39 + 322414.0	QSO	0.9051	0.69	yes
1FGLJ2117.8+0016	J211817.39 + 001316.7	Sy1.5	0.4629	0.79	no

Col. (1) Source name.
Col. (2) SDSS name of the optical counterpart.
Col. (3) SDSS spectroscopic classification based on our analysis.
Col. (4) Redshift estimate derived from our analysis. Question mark indicates uncertain estimates.
(<sup>+</sup>): 2FGLJ2223.4+0104 is the source indicated by Cowperthwaite et al. (2013).
Col. (6) Classification flag: (yes) marks sources that have been classified on the basis of our analysis.

# F. Massaro et al. 2014

TABLE 4 SUMMARY.

Sample	Tot.	BZBs	BZQs	candidates	$\operatorname{new} z$
$(1^{st})$ : ROMA-BZCAT	84	47(47)	3(3)	34(20)	18
$(2^{nd})$ : $\gamma$ -ray blazar candidates	27	7(7)	8(7)	12(4)	4

Col. (2) Total umber of sources listed in the sample.
Col. (3) Number of sources classified as BZBs; those confirmed by our analysis are in parenthesis.
Col. (4) Number of sources classified as BZQs; those confirmed by our analysis are in parenthesis.
Col. (5) Number of blazar candidates; those confirmed by our analysis are in parenthesis.
Col. (6) Sources with new z estimates.