

A HIGHLY ECCENTRIC 3.9-MILLISECOND BINARY PULSAR IN THE GLOBULAR CLUSTER NGC 6652

MEGAN E. DECESAR

Physics Department, University of Wisconsin-Milwaukee, 1900 East Kenwood Boulevard, Milwaukee, WI 53211, USA

SCOTT M. RANSOM

National Radio Astronomy Observatory, Charlottesville, VA 22903, USA

DAVID L. KAPLAN

Physics Department, University of Wisconsin-Milwaukee, 1900 East Kenwood Boulevard, Milwaukee, WI 53211, USA

PAUL S. RAY

Space Science Division, Naval Research Laboratory, Washington, DC 20375-5352, USA

AARON M. GELLER

Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA) and Department of Physics and Astronomy, Northwestern University, 2145

Sheridan Road, Evanston, IL 60208, USA and

Department of Astronomy and Astrophysics, University of Chicago, 5640 S. Ellis Avenue, Chicago, IL 60637, USA

Draft version June 11, 2015

ABSTRACT

We present the Robert C. Byrd Green Bank Telescope discovery of the highly eccentric binary millisecond pulsar PSR J1835–3259A in the *Fermi* Large Area Telescope-detected globular cluster NGC 6652. Timing over one orbit yields the pulse period 3.89 ms, orbital period 9.25 d, eccentricity ~ 0.95 , and an unusually high companion mass of $0.74 M_{\odot}$ assuming a $1.4 M_{\odot}$ pulsar. We caution that the lack of data near periastron prevents a precise measurement of the eccentricity, and that further timing is necessary to constrain this and the other orbital parameters. From tidal considerations, we find that the companion must be a compact object. This system likely formed through an exchange encounter in the dense cluster environment. Our initial timing results predict the measurements of at least two post-Keplerian parameters with long-term phase-connected timing: the rate of periastron advance $\dot{\omega} \sim 0.1^{\circ} \text{ yr}^{-1}$, requiring 1 yr of phase connection; and the Einstein delay $\gamma_{\text{GR}} \sim 10$ ms, requiring 2–3 yr of timing. For an orbital inclination $i > 50^{\circ}$, a measurement of $\sin i$ is also likely. PSR J1835–3259A thus provides an opportunity to measure the neutron star mass with high precision; to probe the cluster environment; and, depending on the nature of the companion, to investigate the limits of general relativity.

Subject headings:

1. INTRODUCTION

Globular clusters (GCs) are efficient producers of low-mass X-ray binaries (LMXBs) and their descendant millisecond pulsars (MSPs; Papitto et al. 2013, and references therein): orders of magnitude more MSPs and LMXBs exist, by mass, in clusters than in the Galactic field (Camilo & Rasio 2005). The dense GC environment heightens the probability of stellar interactions (parameterized by encounter rate γ ; Verbunt & Freire 2014), increasing the likelihood of forming new binaries and of existing binaries gaining new companions. Systems that rarely (if ever) form through known binary evolutionary channels in the field can in principle form through such stellar interactions in GCs, for example: sub-ms pulsars; highly eccentric binaries; or unusual binary systems like MSP-main sequence (Pallanca et al. 2010, and references therein), MSP-MSP, or MSP-black hole (MSP-BH) binaries (Ransom 2008). Such systems would allow astrophysical studies that may not otherwise be possible, e.g., strong-field tests of gravity with MSP-MSP or MSP-BH binaries.

The *Fermi* Large Area Telescope (LAT) has found MSPs to be nearly ubiquitous γ -ray emitters; GeV emission from GCs (Abdo et al. 2010; Tam et al. 2011) may originate from

the clusters' MSP populations (e.g., Venter et al. 2009). The LAT-detected GCs included NGC 6388 and NGC 6652 (Abdo et al. 2010), neither of which contained any known MSPs, but whose γ -ray luminosities implied large MSP populations. NGC 6388 is particularly interesting due to its high encounter rate (e.g., Freire et al. 2008; Maxwell et al. 2012); NGC 6652 may also have a higher encounter rate than previously thought (Noyola & Gebhardt 2006). The presence of a MSP population is supported by the number of X-ray sources, including LMXBs, in both clusters (at least two in NGC 6652 and five in NGC 6388: Maxwell et al. 2012; Stacey et al. 2012). We searched these clusters for radio pulsars; here we report on the discovery and timing of a highly eccentric binary MSP in NGC 6652.

2. OBSERVATIONS AND PULSAR SEARCH ANALYSIS

We observed NGC 6388 and NGC 6652 (Table 1) with the National Radio Astronomy Observatory¹ Robert C. Byrd Green Bank Telescope (GBT) and the Green Bank Ultimate

¹ The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

Pulsar Processing Instrument (GUPPI²) backend (DuPlain et al. 2008), at S band (2 GHz) in coherent search mode. The data were taken with 2048 spectroscopic channels and an effective bandwidth of 700 MHz (accounting for excised radio frequency interference), with coherent dedispersion at the clusters’ predicted dispersion measure (DM) values (Table 1). The observing log and minimum detectable flux densities S_{\min} for an assumed 10% duty cycle are given in Table 2.

The data were analyzed using PRESTO³ (Ransom 2001). Time series were dedispersed at 1900 DMs from 0–691 cm^{-3} pc for NGC 6652 and 5456 DMs between 0–800 cm^{-3} pc for NGC 6388, and were searched for periodicities. We searched for accelerated signals over $z = \pm 200$ Fourier bins (cf. Ransom et al. 2001), corresponding to maximum line-of-sight accelerations between ± 300 – 3000 cm s^{-2} for a 5 ms pulsar.

3. DISCOVERY AND INITIAL TIMING ANALYSIS OF PSR J1835–3259A

We discovered PSR J1835–3259A (hereafter NGC6652A) in the direction of NGC 6652 (DeCesar et al. 2011), with the fundamental frequency at an acceleration of 11.1 cm s^{-2} ($z = 9$). Figure 1 shows the PRESTO discovery plot, and Table 2 contains estimates of the 2 GHz flux density S_2 . We discuss the unexpectedly low DM value of 63.35 cm^{-3} pc below (Section 4.1).

We fit the Doppler-shifted P and \dot{P} (Table 2) with a phase-incoherent orbital model Freire et al. (2001), using a routine by R. Lynch (private communication) employing MPFIT⁴, and found a very eccentric orbit ($e > 0.7$). Starting with this orbital model, we ran TEMPO⁵ iteratively on the pulse times of arrival (TOAs; Table 2) to converge on a family of timing solutions. We phase-connected the first five observations; we did not observe the pulsar at periastron, between observations 5 and 6, so we allowed the phase between these observations to remain arbitrary (i.e., we kept a JUMP between these observations’ sets of TOAs). Using the DD model (Damour & Deruelle 1985, 1986), we find $e = 0.968$. “Faking” phase connection by removing the JUMP yields $e = 0.950$; alternatively, including arbitrary phase JUMPs between all TOA sets yields $e \approx 0.8$, which we take to be the lowest possible e of this system.

The best-fit DD timing model parameters are in Table 3, with fit residuals in Figure 2. The systematics in the residuals are present in all our fits, including those with JUMPs between all observations; we attribute them to parameter, and therefore phase, uncertainties resulting from the lack of TOAs through periastron. We stress that the initial timing parameters in Table 3 belong to a family of solutions—a unique determination of the MSP’s timing solution requires further observations, especially through periastron passage. If further timing confirms the parameters, then NGC6652A will be the most eccentric binary MSP known to date.

4. DISCUSSION

We adopt timing parameters from the $e = 0.950$ model, pulsar mass $m_p = 1.4 M_\odot$, and cluster parameters from Harris (1996, 2010 edition) for all calculations, unless otherwise stated.

² <http://dx.doi.org/10.1117/12.790003>

³ <http://www.cv.nrao.edu/~sransom/presto/>

⁴ <http://cars9.uchicago.edu/software/python/mpfit.html>

⁵ <http://tempo.sourceforge.net/>

4.1. Cluster Membership

The discrepancy between the discovery and predicted DMs (63.35 and 190 cm^{-3} pc, respectively) initially led us to question the MSP’s cluster association (DeCesar et al. 2011). However, the Cordes & Lazio model commonly has uncertainties of a factor 0.5–2, and sometimes larger. The measured DM is consistent with the low optical reddening $E_{B-V} = 0.10 \pm 0.02$ (Ortolani et al. 1994) and estimated X-ray absorption column $N_{\text{H}} \sim 5.5 \times 10^{20} \text{ cm}^{-2}$ (Predehl & Schmitt 1995) toward NGC 6652.

The high e of NGC6652A is much more probable in a GC than in the field due to the high probability of stellar encounters (Camilo & Rasio 2005), discussed further below. Additionally, given the beamwidth of the GBT at S-band ($6.3'$), we estimate a $\approx 0.2\%$ chance of finding an unassociated MSP coincident with NGC 6652 (assuming an isotropic distribution of known galactic MSPs). We conclude that the MSP is almost certainly a cluster member.

4.2. Nature of the Companion

The minimum companion mass (orbital inclination $i = 90^\circ$) is $m_{c,\min} \approx 0.74 M_\odot$ (Table 3). Comparing with the Australia Telescope National Facility (ATNF) Pulsar Catalog⁶ shows that the companion is unusually massive; it may be a main sequence (MS) or evolved star, or a compact object. Based on the cluster’s age (11.7 ± 1.6 Gyr; Chaboyer et al. 2000), the main-sequence turn-off mass is $\approx 0.8 M_\odot$ (Stacey et al. 2012). For $i < 70^\circ$, $m_c > 0.8 M_\odot$, limiting the range of inclinations for which an unevolved MS companion is possible (c.f. Freire et al. 2007).

For a non-compact companion, significant tides at periastron will circularize the orbit. The circularization, or dissipation, timescale t_D for an eccentric binary system can be estimated as (Socrates et al. 2012, and references therein)

$$t_D \equiv \frac{m_c a_F^8}{3k_L \tau G m_p^2 R^5} \quad (1)$$

where $a_F \equiv a \sin i (1 - e^2)$, τ is the constant tidal lag time of the companion, k_L is the Love number, and R is the companion’s radius. For high e , the tidal quality factor Q is related to τ by Equation 23 of Socrates et al. (2012). We estimate k_L to be between 0.05–0.15 for both MS and WD companions, based on calculations with Modules for Experiments in Stellar Astrophysics (MESA; Paxton et al. 2011; Brooker & Olle 1955). For $Q = 10^6$, the circularization timescales are $\sim \text{Myr}$ for a MS companion and $\sim 10^4$ Gyr for a WD companion. We conclude that the companion is a compact object, whose mass and nature will be constrained through further timing.

The merger timescale from gravitational wave-driven inspiral depends on e as $t_{\text{merge}} \propto (1 - e^2)^{7/2}$ (Peters 1964). For $i = 90^\circ$ ($m_{c,\min} = 0.74 M_\odot$) and $e = 0.95$, $t_{\text{merge}} \approx 12$ Gyr; varying e yields a range of $t_{\text{merge}} \sim 1$ Gyr ($e = 0.975$) to > 100 Gyr ($e \leq 0.9$). The system may therefore be disrupted (Section 4.4.2) before it has time to merge. In the event of a merger, when the system comes into contact, the outcome will depend on the exact nature of the binary. Stable mass transfer will be possible for $q \equiv m_c/m_p < 2/3$ ($i > 50^\circ$), forming an ultra-compact X-ray binary and possibly an isolated MSP. For larger inclinations, the mass transfer will be unstable; while a black hole would form from accretion-induced collapse (AIC; Giacomazzo & Perna 2012, and references

⁶ <http://www.atnf.csiro.au/people/pulsar/psrcat/>

therein) if the system mass exceeds the maximum NS mass, it is unclear whether substantial mass would be ejected from the system during unstable mass transfer, preventing AIC (L. Bildsten, *private comm.*). An eventual merger of this system may result in a long-GRB-like, calcium-rich transient (e.g., King et al. 2007) if the companion is a massive WD, or a short GRB for a NS companion (e.g., Grindlay et al. 2006).

4.3. Post-Keplerian Parameters and Mass Constraints

Finding m_c and m_p requires measurements of at least two post-Keplerian (PK) parameters (we employ the general relativistic formalism of Damour & Taylor 1992). Our preliminary timing solution predicts that the rate of change of the longitude of periastron passage $\dot{\omega}$ (i.e., the orbital precession rate) is $\dot{\omega} > 0.08^\circ \text{ yr}^{-1}$, the Einstein delay $\gamma_{\text{GR}} > 10 \text{ ms}$, and $\dot{P}_b > 6 \times 10^{-12} \text{ s s}^{-1}$. Because of the high e , we will measure $\dot{\omega}$ with high precision: from simulations assuming the $e = 0.95$ orbital model parameters, we find that we will measure $\dot{\omega}$ with $> 100\sigma$ significance after one year of timing, yielding the total system mass ($\dot{\omega} \propto M_{\text{tot}}^{2/3}$, where $M_{\text{tot}} = m_p + m_c$) and constraints on m_p and m_c . Knowledge of the pulsar position (requiring 1 yr of timing or an interferometric detection) would yield a measurement of $\dot{\omega}$ with one month of phase-connected timing. Our simulations also show that γ_{GR} will be measured with 10% uncertainty with 2.5 yr of phase connection.

We may also measure one Shapiro delay parameter, $s = \sin i$. For $i > 50^\circ$, the timing residuals from s are significantly larger than the $\approx 20 \mu\text{s}$ uncertainties in the pulse TOAs we used to build the timing model. Statistically, it is most likely that the MSP companion is a WD, requiring $i > 40^\circ$ for $m_c < 1.4 M_\odot$. Even a marginal detection of Shapiro delay will yield a precise s because $\dot{\omega}$ and s are nearly orthogonal in the mass-mass diagram (Lynch et al. 2012). With these two PK measurements, we would precisely measure m_p and m_c . We note that the very precise mass of PSR J1807–2500B (NGC6544B) was measured in this way (Lynch et al. 2012).

4.4. System Origin

The vast majority of field MSP binaries have circular orbits (cf. Champion et al. 2008) from dissipation during the mass-transfer phase (Phinney 1992); known eccentric systems in the field are either double NSs (with eccentricity coming from a second supernova (SN) kick; e.g., Brandt & Podsiadlowski 1995), disrupted triples (Champion et al. 2008), or possibly NS-He WD binaries with circumbinary disks (Antoniadis 2014, and references therein). In contrast, a number of the MSP binaries in GCs are substantially eccentric (e.g., Freire et al. 2008; Lynch et al. 2012)⁷, with a likely origin in dynamical encounters (e.g., Verbunt & Freire 2014). The highest- e binary MSP currently known, PSR J0514–4002A (NGC1851A; Freire et al. 2007), has $e = 0.888$ and an unusually massive ($m_c > 0.96 M_\odot$) companion, similar to NGC6652A. Here we consider the plausibility of several mechanisms through which NGC6652A could have gained its high e .

4.4.1. Possible Formation Mechanisms

There are several ways to form a high-eccentricity system like NGC6652A. An initially circular orbit may gain eccentricity from 3-body encounters with other stars in the GC (Rasio & Heggie 1995). For a double neutron star (DNS), the

eccentricity could have been imparted on the system by the SN kick of a massive companion (Brandt & Podsiadlowski 1995, and references therein). The system could also have formed through an exchange encounter, in which the original companion was ejected from the system and the third body became the new companion (e.g., Verbunt & Freire 2014). In this case, the new companion can be any type of compact object.

The SN kick is ruled out by observational evidence that all known radio pulsars with NS companions have spin periods of 20–100 ms (Tauris 2011), suggesting that MSPs cannot be fully recycled by short-lived, massive companions. The first scenario is plausible, as using Equation 5 of Rasio & Heggie (1995), we find that $\approx 11.8 \text{ Gyr}$ (comparable to the GC age) of non-exchange three-body interactions would be needed for a binary in an initially circular orbit to gain $e = 0.95$. However, the exchange encounter scenario seems most natural, and we discuss this mechanism in more detail below.

Other scenarios for the origin of the binary’s eccentricity include a physical collision between a MSP and a giant star (e.g., Freire et al. 2007), or a triple system in which the outermost companion is pumping the eccentricity of the inner binary (e.g., B1620–26; Thorsett et al. 1999). These mechanisms cannot be excluded *a priori*, but are outside the scope of this letter.

4.4.2. Dynamical Formation Through an Exchange Encounter

We consider a dynamical encounter resulting in a companion exchange, using FEWBODY (Fregeau et al. 2004) to simulate a particular scenario. As a progenitor system, we take the current most common MSP binary in GCs: a MSP in a circular 2 d orbit with a low-mass companion. We chose $m_c = 0.2 M_\odot$, which follows from a binary period of 2 d using the period-core mass relation from Tauris & Savonije (1999) for Pop II stars. We simulated 5000 encounters between this binary and a third body, drawing the incoming velocities from a Maxwellian distribution⁸ using $\sigma = 10 \text{ km s}^{-1}$ distributed between 0 and $30a$ (where a is the binary’s semi-major axis). For the third body we assume a WD with $0.7 M_\odot$.

Approximately 70% of the encounters result in an exchange, with the low-mass companion ejected and an eccentric binary remaining. The new binary has a range of eccentricities strongly biased toward high values, with 66% of the new systems having $e > 0.8$, but energies comparable to that of the progenitor. The orbit has expanded due to the factor of 3.1 increase in m_c , leading to a factor of $3.1^{3/2}$ increase in P_b . Therefore, systems with $e \approx 1$ and $P_b \approx 10 \text{ d}$ are naturally formed through this mechanism. If no exchange happened, then the binary remains close to the 2 d initial period, albeit with enhanced eccentricity.

We estimate the frequency of encounters between a particular NGC6652A-like binary and a single star in NGC 6652 using the single-binary encounter rate γ from Verbunt & Freire (2014), normalized to M4, and find $\gamma_{6652} \approx 6.7\gamma_{\text{M4}}$. A NGC6652A-like binary in M4 would encounter single stars at a rate $\xi_{1+2} \sim (\rho_{c,\text{M4}}/L_\odot)\sigma_{1+2}v_{\text{M4}}$, where ρ_c is the GC core density and σ_{1+2} is the gravitationally-focused single-binary cross-section (Equation A2, Leigh & Sills 2011). In M4, this encounter rate is $\xi_{1+2,\text{M4}} \sim 0.17 \text{ Gyr}^{-1}$; in NGC 6652,

⁸ While the velocity profile of NGC 6652 has not been measured directly, we estimate a velocity dispersion σ of about $10\text{--}15 \text{ km s}^{-1}$, scaling from globular clusters with similar physical core radii (NGC 6388, 6093, and 6441); comparable values are obtained by McLaughlin & van der Marel (2005).

⁷ Also see <http://www.naic.edu/~pfreire/GCpsr.html>.

$\xi_{1+2,6652} \sim 1 \text{ Gyr}^{-1}$. We note that the core radius of $1''.15$ measured by Noyola & Gebhardt (2006) is much smaller than that from Harris (1996, 2010 edition), yielding $\gamma_{6652} \approx 38\gamma_{M4}$ and $\xi_{1+2,6652} \sim 6 \text{ Gyr}^{-1}$. The companion exchange scenario is therefore quite plausible. The position of NGC6652A in the cluster may give additional clues to its formation (cf. Phinney & Sigurdsson 1991).

5. CONCLUSIONS

We discovered one new MSP, NGC6652A. Although NGC 6388 and NGC 6652 are expected to host substantial MSP populations, cluster MSPs are extremely faint—detecting them requires long integration times and the largest telescopes in the world. We did not find more MSPs in these GCs simply because we are sensitivity limited. NGC6652A is an intriguing source for southern-hemisphere Square Kilometer Array (SKA) precursors and eventually the SKA Mid-Frequency Aperture Array⁹.

Our timing analysis over 1.2 orbit of NGC6652A shows that the MSP is in an extremely eccentric binary system with an unusually massive compact companion. The system quite plausibly formed through an exchange encounter in the dense cluster environment. We cannot exclude all other formation mechanisms *a priori*; determining the nature of the compan-

ion will help discriminate between scenarios. Similarly, a precise position will help determine whether the binary is dynamically relaxed (and hence close to the core as expected from mass segregation) or has been kicked out of the core by a recent encounter.

With a phase-connected timing solution over ≥ 1 yr, we will uniquely determine the MSP's timing parameters and measure its position and $\dot{\omega}$ to high precision. After ≈ 2.5 yr of timing, we expect to measure γ_{GR} , allowing measurements of m_p and m_c and clarifying the companion's nature. If $i > 50^\circ$, we may also measure $\sin i$. New timing observations are underway, and will be reported upon in a subsequent paper.

The authors thank P. Arras, P. Freire, J. Fuller, V. Kalogera, and F. Rasio for helpful discussions; F. Camilo for helpful discussions and the use of his computer cluster; and the anonymous referee for useful suggestions that improved the quality of this letter. MED acknowledges funding from NSF Award No. AST-1312822 and NASA's CRESST grant No. 01526268. PSR is supported by the Chief of Naval Research (CNR). AMG is funded by a NSF Astronomy and Astrophysics Postdoctoral Fellowship under Award No. AST-1302765.

⁹ <https://www.skatelescope.org/mfaa>

REFERENCES

- Abdo, A. A., Ackermann, M., Ajello, M., et al. 2010, *A&A*, 524, A75
 Antoniadis, J. 2014, *ApJ*, 797, L24
 Brandt, N., & Podsiadlowski, P. 1995, *MNRAS*, 274, 461
 Brooker, R. A., & Olle, T. W. 1955, *MNRAS*, 115, 101
 Camilo, F., & Rasio, F. A. 2005, in *Astronomical Society of the Pacific Conference Series*, Vol. 328, *Binary Radio Pulsars*, ed. F. A. Rasio & I. H. Stairs, 147
 Chaboyer, B., Sarajedini, A., & Armandroff, T. E. 2000, *AJ*, 120, 3102
 Champion, D. J., Ransom, S. M., Lazarus, P., et al. 2008, *Science*, 320, 1309
 Cordes, J. M., & Lazio, T. J. W. 2002, *ArXiv Astrophysics e-prints*, astro-ph/0207156
 Damour, T., & Deruelle, N. 1985, *Ann. Inst. Henri Poincaré Phys. Théor.*, Vol. 43, No. 1, p. 107 - 132, 43, 107
 —, 1986, *Ann. Inst. Henri Poincaré Phys. Théor.*, Vol. 44, No. 3, p. 263 - 292, 44, 263
 Damour, T., & Taylor, J. H. 1992, *Phys. Rev. D*, 45, 1840
 de Oliveira-Costa, A., Tegmark, M., Gaensler, B. M., et al. 2008, *MNRAS*, 388, 247
 —, 2010, *Global Sky Model (GSM): A Model of Diffuse Galactic Radio Emission from 10 MHz to 100 GHz*, *Astrophysics Source Code Library*, ascl:1011.010
 DeCesar, M. E., Ransom, S. M., & Ray, P. S. 2011, *ArXiv e-prints*, arXiv:1111.0365
 DuPlain, R., Ransom, S., Demorest, P., et al. 2008, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 7019, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*
 Fregeau, J. M., Cheung, P., Portegies Zwart, S. F., & Rasio, F. A. 2004, *MNRAS*, 352, 1
 Freire, P. C., Kramer, M., & Lyne, A. G. 2001, *MNRAS*, 322, 885
 Freire, P. C. C., Ransom, S. M., Bégin, S., et al. 2008, *ApJ*, 675, 670
 Freire, P. C. C., Ransom, S. M., & Gupta, Y. 2007, *ApJ*, 662, 1177
 Giacomazzo, B., & Perna, R. 2012, *ApJ*, 758, L8
 Grindlay, J., Portegies Zwart, S., & McMillan, S. 2006, *Nature Physics*, 2, 116
 Harris, W. E. 1996, *AJ*, 112, 1487
 King, A., Olsson, E., & Davies, M. B. 2007, *MNRAS*, 374, L34
 Leigh, N., & Sills, A. 2011, *MNRAS*, 410, 2370
 Lorimer, D., & Kramer, M. 2005, *Cambridge Observing Handbooks for Research Astronomers*, Vol. 4, *Handbook of Pulsar Astronomy*, 1st edn. (Cambridge, U.K.; New York, U.S.A.: Cambridge University Press)
 Lynch, R. S., Freire, P. C. C., Ransom, S. M., & Jacoby, B. A. 2012, *ApJ*, 745, 109
 Maxwell, J. E., Lugger, P. M., Cohn, H. N., et al. 2012, *ApJ*, 756, 147
 McLaughlin, D. E., & van der Marel, R. P. 2005, *ApJS*, 161, 304
 Moretti, A., Piotto, G., Arcidiacono, C., et al. 2009, *A&A*, 493, 539
 Noyola, E., & Gebhardt, K. 2006, *AJ*, 132, 447
 Ortolani, S., Bica, E., & Barbuy, B. 1994, *A&A*, 286, 444
 Pallanca, C., Dalessandro, E., Ferraro, F. R., et al. 2010, *ApJ*, 725, 1165
 Papitto, A., Ferrigno, C., Bozzo, E., et al. 2013, *Nature*, 501, 517
 Paxton, B., Bildsten, L., Dotter, A., et al. 2011, *ApJS*, 192, 3
 Peters, P. C. 1964, *Phys. Rev.*, 136, B1224
 Phinney, E. S. 1992, *Royal Society of London Philosophical Transactions Series A*, 341, 39
 Phinney, E. S., & Sigurdsson, S. 1991, *Nature*, 349, 220
 Predehl, P., & Schmitt, J. H. M. M. 1995, *A&A*, 293, 889
 Ransom, S. M. 2001, PhD thesis, Harvard University
 Ransom, S. M. 2008, in *American Institute of Physics Conference Series*, Vol. 983, *40 Years of Pulsars: Millisecond Pulsars, Magnetars and More*, ed. C. Bassa, Z. Wang, A. Cumming, & V. M. Kaspi, 415–423
 Ransom, S. M., Greenhill, L. J., Herrnstein, J. R., et al. 2001, *ApJ*, 546, L25
 Rasio, F. A., & Heggie, D. C. 1995, *ApJ*, 445, L133
 Socrates, A., Katz, B., & Dong, S. 2012, *ArXiv e-prints*, arXiv:1209.5724
 Stacey, W. S., Heinke, C. O., Cohn, H. N., Lugger, P. M., & Bahramian, A. 2012, *ApJ*, 751, 62
 Tam, P. H. T., Kong, A. K. H., Hui, C. Y., et al. 2011, *ApJ*, 729, 90
 Tauris, T. M. 2011, in *Astronomical Society of the Pacific Conference Series*, Vol. 447, *Evolution of Compact Binaries*, ed. L. Schmidtobreick, M. R. Schreiber, & C. Tappert, 285
 Tauris, T. M., & Savonije, G. J. 1999, *A&A*, 350, 928
 Thorsett, S. E., Arzoumanian, Z., Camilo, F., & Lyne, A. G. 1999, *ApJ*, 523, 763
 Venter, C., De Jager, O. C., & Clapson, A.-C. 2009, *ApJ*, 696, L52
 Verbunt, F., & Freire, P. C. C. 2014, *A&A*, 561, A11

TABLE 1
TARGETED GLOBULAR CLUSTERS^a

Cluster Name	α	δ	l (degrees)	b (degrees)	Distance (pc)	Predicted DM (cm^{-3} pc)
NGC 6388	17 ^h 36 ^m 17 ^s .88	-44°44'0".24	345.56	-6.74	11.6 ± 2.0 ^b	340 ^c
NGC 6652	18 ^h 35 ^m 44 ^s .86	-32°59'25".10	1.53	-11.37	9 ± 1 ^d	190 ^c

^a The cluster positions were set to the optically determined positions of the cluster centers.

^b Moretti et al. (2009)

^c Cordes & Lazio (2002)

^d Chaboyer et al. (2000)

TABLE 2
OBSERVATION LOG^a

Date	t_{int} (s)	S_{min}^b (μJy)	S/N	S_2^c (μJy)	N_{TOA}	t_{TOA} (s)	$\langle\sigma_{\text{TOA}}\rangle$ (μs)	Barycentric P (ms)	Barycentric \dot{P} (10^{-12} s s ⁻¹)	z (Fourier bins)
NGC 6652										
2010 Oct 19	9470	5.8	25.2	22.5	8	1200	20	3.88937447(3)	-1.223 ± 0.027	8.50
2010 Oct 21	10062	5.6	14.1	12.2	8	1250	22	3.88915225(3)	-0.888 ± 0.023	7.00
2010 Oct 22	8878	6.0	5.8	5.4	7	1100	17	3.88904917(3)	-0.916 ± 0.028	6.00
2010 Oct 23 ^d	7504	6.5	17.4	17	4	940	23	3.888933(1)	-1.33 ± 2.09	... ^d
2010 Oct 24	6701	6.8	14.2	15.1	8	840	21	3.88878963(5)	-1.883 ± 0.060	6.25
2010 Oct 29	3086	10.1	12.6	19.7	7	380	32	3.8891541(1)	-0.934 ± 0.286	0.50
NGC 6388										
2010 Oct 21	6278	7.3
2010 Oct 24	3403	9.9
2010 Oct 29	4080	9.0
2011 Feb 05	5412	7.8
2011 Apr 08	6130	7.4
2011 May 06	4338	8.7

^a All observations were taken at 2 GHz with ~700 MHz effective bandwidth and 40.96 μs time resolution. The beamsize was ~6'.

^b S_{min} was calculated with the radiometer equation for pulsed signals (Appendix A1.4 of Lorimer & Kramer 2005), using $S/N_{\text{min}} = 5$ and pulse width $W = 0.1P$. For observations at S band, the correction factor $\beta = 1.05$, gain $G = 1.9 \text{ K J}^{-1}$, and $T_{\text{rec}} \approx 22 \text{ K}$; for these observations, the number of polarizations $n_p = 2$, and effective bandwidth $\Delta f = 700 \text{ MHz}$. The sky temperature $T_{\text{sky}} = 0.8 \text{ K}$ for NGC 6652 and 1.4 K for NGC 6388 (de Oliveira-Costa et al. 2008, 2010), giving $T_{\text{sys}} = T_{\text{rec}} + T_{\text{sky}} = 22.8 \text{ K}$ and 23.4 K, respectively.

^c The 2 GHz flux density S_2 was calculated with the same radiometer equation parameters as for S_{min} , but using the measured S/N rather than $S/N_{\text{min}} = 5$ and the measured $W = 0.0625P$ rather than $W = 0.1P$. The smaller measured W explains why we find $S_2 < S_{\text{min}}$ on 2010 October 22.

^d The DM used for coherent dedispersion of individual channels was accidentally set to 9.0 cm^{-3} pc for this observation. As a result, the MSP was not found in a straightforward acceleration search.

TABLE 3
NGC6652A TIMING SOLUTION^d

Timing Parameter	With JUMP	Without JUMP
Right Ascension ^b (J2000.0)	18 ^h 35 ^m 44 ^s .856	18 ^h 35 ^m 44 ^s .856
Declination ^b (J2000.0)	-32° 59' 25".08	-32° 59' 25".08
Dispersion Measure ^c (cm ⁻³ pc)	63.35	63.35
Spin period, P (ms)	3.888824(1)	3.8888289774(4)
Spin period epoch (MJD)	55488.931354	55488.931354
Spindown rate ^d , \dot{P} (s s ⁻¹)	0	0
Orbital period, P_b (days)	9.2460(5)	9.2459(5)
Projected semimajor axis, x (s)	19.6(3)	19.09(5)
Eccentricity, e	0.968(5)	0.950(1)
Epoch of periastron passage, T_0 (MJD)	55477.061(5)	55477.0401(6)
Longitude of periastron, ω (degrees)	291(1)	289.2(2)
Minimum companion mass, $m_{c,\min}$ (M_\odot)	0.765(14)	0.736(3)
Fit χ^2 per degrees of freedom	347.9/35	369.37/36

^a The solution uses the DD model (Damour & Deruelle 1985, 1986) and the TDB time system. The error on the last digit(s) of each parameter value is denoted in parentheses. The true solution is one in a family of solutions represented by the parameters listed here. The middle column gives the timing parameters obtained when allowing an arbitrary number of pulsar rotations between the fifth and sixth observations. The right column gives the parameters obtained with “forced” phase connection. The unique solution will be determined with further timing observations that include a periastron passage.

^b The position was fixed at the cluster’s center.

^c The DM was fixed to this best value from the discovery observation (2010 October 19).

^d The spindown rate was fixed at zero; a phase-connected timing solution spanning \sim one year will measure this parameter.




FIG. 1.— PRESTO discovery plot. *Top left*: Two cycles of the summed pulse profile. *Left*: Two pulse cycles, with the signal in 64 sub-integrations split evenly over the full integration time. The χ^2 plot on the right shows the steadily increasing S/N over time. *Middle*: Two pulse cycles with the signal in 32 frequency sub-bands split evenly over the full bandwidth. *Bottom middle*: The χ^2 increases dramatically near the pulsar DM. *Right*: The maximum χ^2 determines the best P (top) and \dot{P} (middle). *Bottom right*: Covariance between P and \dot{P} .




FIG. 2.— *Top:* Predicted spin periods for a circular orbit (dashed line) and the DD orbital model with a JUMP between observations 5 and 6 ($e = 0.967$; solid line). The incoherently measured P and \dot{P} , including uncertainties, are overlaid in black. *Middle:* Residuals from subtracting P predicted by the $e = 0.967$ model from the measured P . *Bottom:* Timing residuals from TOAs fit with the $e = 0.967$ model. We attribute the systematics in the TOAs to phase uncertainties which will be resolved with a fully phase-connected timing solution.