

# ON THE GREATEST AND LEAST ELEMENTS IN THE SET OF SEMISTANDARD TABLEAUX OF GIVEN SHAPE AND WEIGHT

AKIHIRO MUNEMASA AND MINWON NA

ABSTRACT. We give three algorithms to construct a semistandard tableau of given shape and weight, where the weight is a composition which is not necessarily a partition. With respect to a natural partial order on the set of semistandard tableaux, we show that the set of semistandard tableaux of given shape and weight has a unique greatest element and a unique least element. Two of our algorithms give each of these elements.

## 1. INTRODUCTION

Kostka numbers give the number of semistandard tableaux of given shape and weight, and they play a fundamental role in representation theory of symmetric groups (see [4]). Much work have been done on the problem of computing Kostka numbers, which is known to be  $\#P$  complete (see [3]). In this paper, we study the *set*, rather than the number, of semistandard tableaux of given shape and weight, in the hope that our study could shed some light on the computation of the cardinality of the set in question. We do not assume the weight is a partition, rather, it is an arbitrary composition.

Throughout this paper,  $n$  will denote a positive integer. Let  $\mu = (\mu_1, \mu_2, \dots, \mu_k)$  and  $a = (a_1, a_2, \dots, a_h)$  be a partition and composition of  $n$ , respectively. We say that  $h$  is the height  $h(a)$  of  $a$ . We denote by  $D_\mu$  the Young diagram of  $\mu$ , and by  $\text{STab}(\mu, a)$  the set of all semistandard tableaux of shape  $\mu$  and weight  $a$ . More precisely,

$$D_\mu = \{(i, j) \in \mathbb{Z}^2 \mid 1 \leq i \leq k, 1 \leq j \leq \mu_i\},$$

$$\text{STab}(\mu, a) = \{T \mid T : D_\mu \rightarrow \{1, \dots, h\},$$

$$T(i, j) \leq T(i, j + 1), T(i, j) < T(i + 1, j), (|T^{-1}(\{i\})|)_{i \geq 1} = a\}.$$

For compositions  $a = (a_1, a_2, \dots, a_h)$  and  $b = (b_1, b_2, \dots, b_k)$  of  $n$ , we say  $a$  dominates  $b$ , denoted  $a \trianglerighteq b$ , if  $k \geq h$  and

$$\sum_{i=1}^j a_i \geq \sum_{i=1}^j b_i$$

for  $j = 1, 2, \dots, h$ . The following is well known.

**Theorem 1** ([2, p. 26, Exercise 2]). *Let  $\mu$  and  $\lambda$  be partitions of  $n$ . Then  $\text{STab}(\mu, \lambda)$  is nonempty if and only if  $\mu \trianglerighteq \lambda$ .*

---

*Date:* June 29, 2015.

*2010 Mathematics Subject Classification.* 05A15, 05A17, 05A19, 05E10, 20C30.

*Key words and phrases.* Young diagram, semistandard tableau, Kostka number, partition, dominance order.

Let  $\lambda(a)$  denote the partition of  $n$  associated with  $a$ , that is, the partition obtained from  $a$  by rearranging the parts of  $a$  in non-increasing order. Then one can strengthen Theorem 1 using [1, Lemma 3.7.1], as follows:

**Theorem 2** ([2, p. 50, Proposition 2]). *Let  $\mu$  and  $a$  be a partition and composition of  $n$ , respectively. Then  $\text{STab}(\mu, a)$  is nonempty if and only if  $\mu \supseteq \lambda(a)$ .*

This theorem is incorrectly stated in [1, Lemma 3.7.3], where  $\mu \supseteq \lambda(a)$  is replaced by  $\mu \supseteq a$ . For example, let  $\mu = (5, 3) \vdash 8$  and  $a = (2, 6) \vDash 8$ . Then  $\mu \supseteq a$ , but  $\text{STab}(\mu, a) = \emptyset$ .

The purpose of this note is to give explicit algorithms to produce an element of  $\text{STab}(\mu, a)$ , thereby giving a direct proofs of Theorem 2. We also introduce a natural partial order on  $\text{STab}(\mu, a)$  and show that it has unique greatest and least elements, by showing that the elements produced by two of the three algorithms have the respective property. Although the proof of Theorem 2 using [1, Lemma 3.7.1] or [2, p. 50, Proposition 2] gives, in principle, a bijection between  $\text{STab}(\mu, a)$  and  $\text{STab}(\mu, \lambda(a))$ , it does not give an efficient algorithm to describe an element of  $\text{STab}(\mu, a)$  unless the permutation required to transform  $a$  to  $\lambda(a)$  is a transposition. We show that, in our Proposition 19 below, a direct approach for proving Theorem 2 along the line of [1, Lemma 3.7.3] can be justified, and we describe an algorithm to produce an element of  $\text{STab}(\mu, a)$  in this way.

This paper is organized as follows. After giving preliminaries in Section 2, we describe the procedure of constructing the greatest and least elements of  $\text{STab}(\mu, a)$  in Section 3 and 5, respectively. In Section 4, we justify the proof of [1, Lemma 3.7.3] in Proposition 19, using the ideas from Section 3.

## 2. PRELIMINARIES

A composition of  $n$  is a sequence  $a = (a_1, a_2, \dots, a_h)$  of positive integers such that  $\sum_{i=1}^h a_i = n$ . We write  $a \vDash n$  if  $a$  is a composition of  $n$ . A partition of  $n$  is a non-increasing sequence  $\mu = (\mu_1, \mu_2, \dots, \mu_k)$  of positive integers such that  $\sum_{i=1}^k \mu_i = n$ . We write  $\mu \vdash n$  if  $\mu$  is a partition of  $n$ . By convention, we define  $\mu_i = 0$  if  $i$  exceeds the number of parts in a partition  $\mu$ .

For a composition  $a = (a_1, a_2, \dots, a_h) \vDash n$ , we define

$$a^{(i)} = (a_1, \dots, a_{i-1}, a_i - 1, a_{i+1}, \dots, a_h)$$

for each  $1 \leq i \leq h$ . Set

$$a' = (a_1, a_2, \dots, a_{h-1}) \vDash n - a_h,$$

$$\tilde{a} = \begin{cases} (a_1, a_2, \dots, a_h - 1) & \text{if } a_h \geq 2, \\ (a_1, a_2, \dots, a_{h-1}) & \text{if } a_h = 1. \end{cases}$$

Then  $\tilde{a} \vDash n - 1$ . Let

$$q(a) = \max\{i \mid 1 \leq i \leq h, \lambda(a)_i = a_h\}.$$

Then

$$(1) \quad \lambda(a)_{q(a)} = a_h > \lambda(a)_{q(a)+1}.$$

Set

$$\tilde{\lambda}(a) = \lambda(a)^{(q(a))} \vdash n - 1.$$

For a partition  $\mu = (\mu_1, \dots, \mu_k) \vdash n$ , we define

$$s(\mu, a) = \max\{i \mid 1 \leq i \leq k, \mu_i \geq a_h\}.$$

Clearly,

$$(2) \quad \mu_{s(\mu, a)} \geq a_h > \mu_{s(\mu, a)+1}.$$

For  $\rho = (\rho_1, \rho_2, \dots, \rho_h) \vdash m$  and  $\mu = (\mu_1, \mu_2, \dots, \mu_k) \vdash n$ , we write

$$\rho \preceq \mu$$

if  $m \leq n$ ,  $h \leq k$  and  $\rho_i \leq \mu_i$  for all  $i$  with  $1 \leq i \leq h$ . For such  $\rho$  and  $\mu$ , we say that  $\mu/\rho$  is a *skew shape*, and we denote  $D_\mu \setminus D_\rho$  by  $D_{\mu/\rho}$ . We say that the skew shape  $\mu/\rho$  is totally disconnected if  $\rho_i \geq \mu_{i+1}$  for all  $i$  with  $1 \leq i \leq h$ . Set

$$\mathcal{B}(\mu, a) = \{\rho \vdash n - a_h \mid \rho \supseteq \lambda(a'), \rho \preceq \mu, \mu/\rho : \text{totally disconnected}\}.$$

**Lemma 3.** For a composition  $a \vDash n$ ,  $\lambda(\tilde{a}) = \tilde{\lambda}(a)$ .

*Proof.* Immediate from the definition.  $\square$

**Lemma 4.** Let  $p$  and  $q$  be positive integers. Let  $\mu \vdash n$  and  $\lambda \vdash n$  satisfy  $\mu_p > \mu_{p+1}$ ,  $\lambda_q > \lambda_{q+1}$  and  $\mu \supseteq \lambda$ . Then the following are equivalent.

- (i)  $\mu^{(p)} \supseteq \lambda^{(q)}$ ,
- (ii) either  $p \geq q$ , or  $p < q$  and  $\sum_{i=1}^j \mu_i > \sum_{i=1}^j \lambda_i$  for all  $j$  with  $p \leq j < q$ .

*Proof.* Observe

$$\sum_{i=1}^j \mu_i^{(p)} = \begin{cases} \sum_{i=1}^j \mu_i & \text{if } 1 \leq j < p, \\ \sum_{i=1}^j \mu_i - 1 & \text{otherwise,} \end{cases}$$

$$\sum_{i=1}^j \lambda_i^{(q)} = \begin{cases} \sum_{i=1}^j \lambda_i & \text{if } 1 \leq j < q, \\ \sum_{i=1}^j \lambda_i - 1 & \text{otherwise.} \end{cases}$$

Thus

$$\sum_{i=1}^j \mu_i^{(p)} \geq \sum_{i=1}^j \lambda_i^{(q)} \iff \begin{cases} \sum_{i=1}^j \mu_i \geq \sum_{i=1}^j \lambda_i & \text{if } 1 \leq j < \min\{p, q\}, \\ \sum_{i=1}^j \mu_i \geq \sum_{i=1}^j \lambda_i - 1 & \text{if } q \leq j < p, \\ \sum_{i=1}^j \mu_i - 1 \geq \sum_{i=1}^j \lambda_i & \text{if } p \leq j < q, \\ \sum_{i=1}^j \mu_i - 1 \geq \sum_{i=1}^j \lambda_i - 1 & \text{if } \max\{p, q\} \leq j. \end{cases}$$

Since  $\mu \supseteq \lambda$ , we have

$$(i) \iff \sum_{i=1}^j \mu_i - 1 \geq \sum_{i=1}^j \lambda_i \quad \text{if } p \leq j < q,$$

$$\iff (ii).$$

$\square$

**Definition 5.** Let  $\mu \vdash n$  and  $a \vDash n$  with  $\mu \supseteq \lambda(a)$ . A box of coordinate  $(i, \mu_i)$  is *removable for the pair*  $(\mu, a)$  if  $\mu_i > \mu_{i+1}$  and  $\mu^{(i)} \supseteq \lambda(\tilde{a})$ . We denote by  $R(\mu, a)$  the set of all  $i$  such that  $(i, \mu_i)$  is removable for the pair  $(\mu, a)$ .

**Lemma 6.** *Let  $\mu \vdash n$  and  $a \vDash n$  satisfy  $\mu \supseteq \lambda(a)$ . Then  $s(\mu, a) \in R(\mu, a)$ .*

*Proof.* Write  $p = s(\mu, a)$ ,  $q = q(a)$ ,  $\lambda = \lambda(a)$ . Then  $\lambda_q > \lambda_{q+1}$  by (1) and  $\mu_p > \mu_{p+1}$  by (2). Moreover, by Lemma 3, we have  $\lambda(\tilde{a}) = \tilde{\lambda}(a) = \lambda^{(a)}$ . Thus, in view of Lemma 4, it suffices to show

$$\text{either } p \geq q, \text{ or } p < q \text{ and } \sum_{i=1}^j \mu_i > \sum_{i=1}^j \lambda_i \text{ for all } j \text{ with } p \leq j < q.$$

Suppose  $p < q$  and let  $p \leq j < q$ . Let  $a = (a_1, \dots, a_h)$ . If  $j+1 \leq i \leq q$ , then  $p < i \leq q$ , so  $\mu_i < a_h \leq \lambda_i$ . Thus

$$\begin{aligned} \sum_{i=1}^j \mu_i &= \sum_{i=1}^q \mu_i - \sum_{i=j+1}^q \mu_i \\ &\geq \sum_{i=1}^q \lambda_i - \sum_{i=j+1}^q \mu_i \\ &> \sum_{i=1}^q \lambda_i - \sum_{i=j+1}^q \lambda_i \\ &= \sum_{i=1}^j \lambda_i. \end{aligned}$$

□

From Lemma 6, we find  $R(\mu, a) \neq \emptyset$ . Set

$$l(\mu, a) = \min R(\mu, a).$$

**Lemma 7.** *Let  $\mu \vdash n$  and  $a \vDash n$  satisfy  $\mu \supseteq \lambda(a)$ . Then  $l(\mu, a) \leq s(\mu, a)$ .*

*Proof.* Immediate from Lemma 6. □

**Lemma 8.** *Let  $\mu = (\mu_1, \mu_2, \dots, \mu_k) \vdash n$  and  $a = (a_1, a_2, \dots, a_h) \vDash n$ . Let  $i$  be an integer with  $1 \leq i \leq k$ . If there exists a tableau  $T \in \text{STab}(\mu, a)$  such that  $T(i, \mu_i) = h$ , then  $i \in R(\mu, a)$ .*

*Proof.* Since  $T \in \text{STab}(\mu, a)$  and  $T(i, \mu_i) = h$ , we have  $(i+1, \mu_i) \notin D_\mu$ , so  $\mu_i > \mu_{i+1}$ . Also,  $\tilde{T} = T|_{D_\mu \setminus \{(i, \mu_i)\}} \in \text{STab}(\mu^{(i)}, \tilde{a})$ . By [1, Lemma 3.7.1], we obtain  $\text{STab}(\mu^{(i)}, \lambda(\tilde{a})) \neq \emptyset$ . Thus  $\mu^{(i)} \supseteq \lambda(\tilde{a})$  and  $i \in R(\mu, a)$ . □

In fact, the converse of Lemma 8 is also true. We will prove it in Section 4.

**Lemma 9.** *Let  $\mu \vdash n$  and  $a = (a_1, a_2, \dots, a_h) \vDash n$  satisfy  $\mu \supseteq \lambda(a)$ . For  $T \in \text{STab}(\mu, a)$ , we have*

$$l(\mu, a) \leq \min\{i \mid T(i, \mu_i) = h\} \leq s(\mu, a).$$

*Proof.* Write  $q = \min\{i \mid T(i, \mu_i) = h\}$ . By Lemma 8, we have  $l(\mu, a) \leq q$ . Since  $a_h = |\{(i, j) \in D_\mu \mid T(i, j) = h\}| \leq \mu_q$ , we have  $q \leq s(\mu, a)$ . □

We will show in Theorem 13, Lemma 17 and Theorem 25 that equality can be achieved in both of the inequalities above. In Section 3, we give an algorithm to construct  $T \in \text{STab}(\mu, a)$  such that  $\min\{i \mid T(i, \mu_i) = h\} = s(\mu, a)$ . In Sections 4 and 5, we give an algorithm to construct  $T \in \text{STab}(\mu, a)$  such that  $\min\{i \mid T(i, \mu_i) = h\} = l(\mu, a)$ .

Finally, we define a partial order on  $\text{STab}(\mu, a)$  and a partition  $\rho(\mu, a) \vdash n - a_h$  as follows. We write  $\mu \triangleright \lambda$  to mean  $\mu \supseteq \lambda$  and  $\mu \neq \lambda$ .

**Definition 10.** Let  $\mu = (\mu_1, \mu_2, \dots, \mu_k) \vdash n$  and  $a = (a_1, a_2, \dots, a_h) \vDash n$  satisfy  $\mu \supseteq \lambda(a)$ . For  $T, S \in \text{STab}(\mu, a)$ , let

$$(3) \quad \tau^{(p)} = (|\{j \mid T(i, j) \leq p\}|)_{i=1}^k,$$

$$(4) \quad \sigma^{(p)} = (|\{j \mid S(i, j) \leq p\}|)_{i=1}^k$$

for all  $p$  with  $1 \leq p \leq h$ . We define that  $S \leq T$  if, either  $T = S$  or,  $\tau^{(h)} = \sigma^{(h)}$ ,  $\tau^{(h-1)} = \sigma^{(h-1)}, \dots, \tau^{(p+1)} = \sigma^{(p+1)}$ ,  $\tau^{(p)} \triangleright \sigma^{(p)}$  for some  $1 \leq p \leq h$ .

Since the relation  $\supseteq$  is a partial order, we see that  $(\text{STab}(\mu, a), \leq)$  is a partially ordered set.

Alternatively the partial order  $\leq$  on  $\text{STab}(\mu, a)$  can be defined recursively as follows: for  $T, S \in \text{STab}(\mu, a)$ , define  $\tau$  and  $\sigma$  by

$$(5) \quad T^{-1}(\{1, \dots, h-1\}) = D_\tau,$$

$$(6) \quad S^{-1}(\{1, \dots, h-1\}) = D_\sigma,$$

respectively. We define  $S \leq T$  if, either  $\tau \triangleright \sigma$ , or  $\tau = \sigma$  and  $S|_{D_\sigma} \leq T|_{D_\sigma}$ .

**Definition 11.** Let  $\mu = (\mu_1, \dots, \mu_k) \vdash n$  and  $a = (a_1, \dots, a_h) \vDash n$  satisfy  $\mu \supseteq \lambda(a)$ . Define  $\rho(\mu, a) = (\rho_1, \dots, \rho_{k-1}) \vdash n - a_h$  by setting

$$\rho_i = \begin{cases} \mu_i & \text{if } 1 \leq i < s, \\ \mu_s - (a_h - \mu_{s+1}) & \text{if } i = s, \\ \mu_{i+1} & \text{if } s < i \leq k-1, \end{cases}$$

where  $s = s(\mu, a)$ .

### 3. THE GREATEST ELEMENT OF $\text{STab}(\mu, a)$

**Lemma 12.** Let  $\mu = (\mu_1, \dots, \mu_k) \vdash n$  and  $a = (a_1, \dots, a_h) \vDash n$  satisfy  $\mu \supseteq \lambda(a)$ . Then  $\rho(\mu, a)$  is the greatest element of  $\mathcal{B}(\mu, a)$ .

*Proof.* Write  $\rho = \rho(\mu, a)$  and  $s = s(\mu, a)$ . By (2), we have  $\mu_s \geq a_h > \mu_{s+1}$ . Thus  $\rho \vdash n - a_h$  and  $\mu_s > \rho_s \geq \mu_{s+1}$ . So  $\mu_i \geq \rho_i \geq \mu_{i+1}$  for all  $i$  with  $1 \leq i \leq k$ . This implies that  $\rho \preceq \mu$  and  $\mu/\rho$  is totally disconnected.

Next, we show that  $\rho \supseteq \lambda(a')$ . Write  $\lambda(a) = (\lambda_1, \dots, \lambda_h)$ ,  $\lambda' = (\lambda'_1, \dots, \lambda'_{h-1}) = \lambda(a')$  and  $q = q(a)$ . Then  $\lambda_q = a_h$  and

$$\lambda' = (\lambda_1, \lambda_2, \dots, \lambda_{q-1}, \lambda_{q+1}, \lambda_{q+2}, \dots, \lambda_h) \vdash n - \lambda_q.$$

Observe

$$\sum_{i=1}^j \rho_i = \begin{cases} \sum_{i=1}^j \mu_i & \text{if } 1 \leq j < s, \\ \sum_{i=1}^{j+1} \mu_i - \lambda_q & \text{if } s \leq j \leq k-1 \end{cases}$$

$$(7) \quad \geq \begin{cases} \sum_{i=1}^j \lambda_i & \text{if } 1 \leq j < s, \\ \sum_{i=1}^{j+1} \lambda_i - \lambda_q & \text{if } s \leq j \leq k-1 \end{cases}$$

since  $\mu \geq \lambda(a)$ .

**Case 1.**  $q < s$ . By (7), we have

$$\begin{aligned} \sum_{i=1}^j \rho_i &\geq \begin{cases} \sum_{i=1}^j \lambda_i & \text{if } 1 \leq j < q, \\ \sum_{i=1}^{q-1} \lambda_i + \sum_{i=q+1}^{j+1} \lambda_i + \lambda_q - \lambda_{j+1} & \text{if } q \leq j < s, \\ \sum_{i=1}^{j+1} \lambda_i - \lambda_q & \text{if } s \leq j \leq k-1 \end{cases} \\ &\geq \begin{cases} \sum_{i=1}^j \lambda_i & \text{if } 1 \leq j < q, \\ \sum_{i=1}^{q-1} \lambda_i + \sum_{i=q+1}^{j+1} \lambda_i & \text{if } q \leq j < s, \\ \sum_{i=1}^{j+1} \lambda_i - \lambda_q & \text{if } s \leq j \leq k-1 \end{cases} \\ &= \sum_{i=1}^j \lambda'_i. \end{aligned}$$

**Case 2.**  $s \leq q$ . By (7), we have

$$\begin{aligned} \sum_{i=1}^j \rho_i &\geq \begin{cases} \sum_{i=1}^j \lambda_i & \text{if } 1 \leq j < s, \\ \sum_{i=1}^j \lambda_i + \lambda_{j+1} - \lambda_q & \text{if } s \leq j < \min\{q, k\}, \\ \sum_{i=1}^{j+1} \lambda_i - \lambda_q & \text{if } \min\{q, k\} \leq j \leq k \end{cases} \\ &\geq \begin{cases} \sum_{i=1}^j \lambda_i & \text{if } 1 \leq j < \min\{q, k\}, \\ \sum_{i=1}^{j+1} \lambda_i - \lambda_q & \text{if } q \leq j \leq k \end{cases} \\ &\geq \sum_{i=1}^j \lambda'_i. \end{aligned}$$

Thus we have  $\rho \in \mathcal{B}(\mu, a)$ .

It remains to show that  $\rho \geq \tau$  for all  $\tau \in \mathcal{B}(\mu, a)$ . Let  $\tau \in \mathcal{B}(\mu, a)$  with  $\tau \neq \rho$  and

$$(8) \quad r = \min\{i \mid 1 \leq i \leq k, \tau_i < \mu_i\}.$$

Then

$$(9) \quad 1 \leq r \leq s$$

and

$$(10) \quad \sum_{i=r}^k (\mu_i - \tau_i) = a_h.$$

By the definition of  $\mathcal{B}(\mu, a)$ , we have

$$(11) \quad \mu_i \geq \tau_i \geq \mu_{i+1}$$

for all  $i$  with  $1 \leq i \leq k$ .

If  $1 \leq j < s$ , then

$$\sum_{i=1}^j \rho_i = \sum_{i=1}^j \mu_i \geq \sum_{i=1}^j \tau_i.$$

If  $s \leq j \leq k$ , then

$$\begin{aligned}
\sum_{i=1}^j \rho_i &= \sum_{i=1}^{j+1} \mu_i - a_h \\
&= \sum_{i=1}^{j+1} \mu_i - \sum_{i=r}^k (\mu_i - \tau_i) && \text{(by (10))} \\
&= \sum_{i=1}^{r-1} \mu_i + \sum_{i=r}^{j+1} \mu_i - \sum_{i=r}^k \mu_i + \sum_{i=r}^k \tau_i && \text{(by (9))} \\
&= \sum_{i=1}^j \tau_i + \sum_{i=j+1}^k \tau_i - \sum_{i=j+2}^k \mu_i && \text{(by (8))} \\
&= \sum_{i=1}^j \tau_i + \sum_{i=j+1}^k (\tau_i - \mu_{i+1}) \\
&\geq \sum_{i=1}^j \tau_i && \text{(by (11)).}
\end{aligned}$$

Therefore,  $\rho \supseteq \tau$ . □

**Theorem 13.** Given  $\mu \vdash n$  and  $a = (a_1, a_2, \dots, a_h) \vDash n$  such that  $\mu \supseteq \lambda(a)$ , define  $\rho^i$  and  $a^i$  inductively by setting  $\rho^0 = \mu$ ,  $a^0 = a$ , and for  $1 \leq i \leq h$ ,

$$\begin{aligned}
\rho^i &= \rho(\rho^{i-1}, a^{i-1}) \vdash \sum_{j=1}^{h-i} a_j, \\
a^i &= (a^{i-1})' \vDash \sum_{j=1}^{h-i} a_j.
\end{aligned}$$

Define a tableau  $T$  of shape  $\mu$  and weight  $a$  by

$$(12) \quad T(p, q) = h - i \text{ if } (p, q) \in D_{\rho^i/\rho^{i+1}}.$$

Then  $T$  is the greatest element of  $\text{STab}(\mu, a)$ .

*Proof.* We prove the assertion by induction on  $h$ . Suppose first  $h = 1$ . Then  $\text{STab}(\mu, a)$  consists of a single element  $T$ , so that the assertion trivially holds.

Next suppose  $h > 1$ . Assume that the assertion holds for  $h - 1$ . Set  $\nu = \rho^1$  and  $b = a^1$ . Since  $\rho^1 \in \mathcal{B}(\mu, a)$  by Lemma 12, we have  $\nu \supseteq \lambda(b)$ . Define  $\nu^i$  and  $b^i$  inductively by setting  $\nu^0 = \nu$ ,  $b^0 = b$ , and for  $1 \leq i < h$ ,

$$\begin{aligned}
\nu^i &= \rho(\nu^{i-1}, b^{i-1}) \vdash \sum_{j=1}^{h-1-i} b_j, \\
b^i &= (b^{i-1})' \vDash \sum_{j=1}^{h-1-i} b_j.
\end{aligned}$$

Define a tableau  $T'$  of shape  $\nu$  and weight  $b$  by

$$T'(p, q) = h - (i + 1) \text{ if } (p, q) \in D_{\nu^i/\nu^{i+1}}.$$

By the inductive hypothesis,

$$(13) \quad T' \in \text{STab}(\nu, b),$$

$$(14) \quad T' \geq S' \text{ for all } S' \in \text{STab}(\nu, b).$$

It is easy to show that  $b^i = a^{i+1}$  and  $\nu^i = \rho^{i+1}$  by induction on  $i$ , and the latter implies  $T|_{D_\nu} = T'$ . Then by (13) and the fact that  $\mu/\nu$  is totally disconnected, we obtain  $T \in \text{STab}(\mu, a)$ .

It remains to show that  $T \geq S$  for all  $S \in \text{STab}(\mu, a)$ . Let  $S \in \text{STab}(\mu, a)$ . Define a partition  $\sigma$  by (6). By Lemma 12, we have  $\nu \supseteq \sigma$ . If  $\nu \triangleright \sigma$ , then  $T \geq S$ . If  $\nu = \sigma$ , then  $T|_{D_\nu} = T' \geq S|_{D_\nu}$  by (14), hence  $T \geq S$ .  $\square$

From Theorem 13, we obtain a tableau  $T \in \text{STab}(\mu, a)$ .

**Algorithm 1. Input:**  $\mu \vdash n$  and  $a \vDash n$  such that  $\mu \supseteq \lambda(a)$ .

**Output:**  $T \in \text{STab}(\mu, a)$ .

**Initialization:**  $\nu := \mu$ ,  $b := a$ .

**while**  $h(b) > 1$  **do**

$$\left| \begin{array}{l} T(i, j) := h(b) \text{ where } (i, j) \in D_{\nu/\rho(\nu, b)}. \\ \nu \leftarrow \rho(\nu, b), b \leftarrow b'. \end{array} \right.$$

**end**

$$T(1, j) := 1 \text{ where } 1 \leq j \leq \nu_1.$$

Output  $T$ .

**Example 14.** Let  $\mu = (4, 4, 1, 1) \vdash 10$  and  $a = (1, 3, 2, 2, 2) \vDash 10$ . Then a tableau  $T \in \text{STab}(\mu, a)$  is obtained via Algorithm 1.

$\nu$	$b$	$\rho(\nu, b)$	$T$
(4, 4, 1, 1)	(1, 3, 2, 2, 2)	(4, 3, 1)	$T(2, 4) = T(4, 1) = 5$
(4, 3, 1)	(1, 3, 2, 2)	(4, 2)	$T(2, 3) = T(3, 1) = 4$
(4, 2)	(1, 3, 2)	(4)	$T(2, 1) = T(2, 2) = 3$
(4)	(1, 3)	(1)	$T(1, 2) = T(1, 3) = T(1, 4) = 2$
(1)	(1)		$T(1, 1) = 1$

Thus

$$T = \begin{array}{cccc} 1 & 2 & 2 & 2 \\ 3 & 3 & 4 & 5 \\ 4 & & & \\ 5 & & & \end{array}$$

is the greatest element of  $\text{STab}(\mu, a)$ .

#### 4. REMOVABLE BOXES

Throughout this section, let  $\mu = (\mu_1, \mu_2, \dots, \mu_k) \vdash n$  and  $a = (a_1, a_2, \dots, a_h) \vDash n$ . Let  $\lambda = \lambda(a) = (\lambda_1, \lambda_2, \dots, \lambda_h)$ ,  $q = q(a)$  and  $l = l(\mu, a)$ . We assume  $\mu \supseteq \lambda$ .

**Lemma 15.** *Assume  $a_h \geq 2$ . For  $i \in R(\mu, a)$  with  $i \leq s(\mu^{(i)}, \tilde{a})$ , we have  $\rho(\mu^{(i)}, \tilde{a}) \in \mathcal{B}(\mu, a)$ .*



*Proof.* Write  $\rho = \rho(\mu^{(i)}, \tilde{a})$ . Since  $i \in R(\mu, a)$ , we have  $\mu^{(i)} \supseteq \lambda(\tilde{a})$ . Then by Lemma 12,  $\rho \in \mathcal{B}(\mu^{(i)}, \tilde{a})$ . Since  $a_h \geq 2$ , this implies  $\rho \supseteq \lambda(\tilde{a}') = \lambda(a')$ . To prove  $\rho \in \mathcal{B}(\mu, a)$ , it remains to show that  $\mu/\rho$  is totally disconnected. Since  $\mu^{(i)}/\rho$  is totally disconnected, it is enough to show  $\rho_{i-1} \geq \mu_i$ . Since  $i-1 < s(\mu^{(i)}, \tilde{a})$ , we obtain  $\rho_{i-1} = \mu_{i-1}^{(i)} = \mu_{i-1} \geq \mu_i$ .  $\square$

**Lemma 16.** *Assume  $a_h \geq 2$ . Then  $r \leq s(\mu, a)$  if and only if  $r \leq s(\mu^{(r)}, \tilde{a})$ .*

*Proof.* Immediate from the definition.  $\square$

**Lemma 17.** *Define  $\rho \vdash n - a_h$  by*

$$\rho = \begin{cases} \mu^{(l)} & \text{if } a_h = 1, \\ \rho(\mu^{(l)}, \tilde{a}) & \text{if } a_h \geq 2. \end{cases}$$

*Then  $\mu/\rho$  is totally disconnected,  $\rho_l < \mu_l$  and  $\rho \supseteq \lambda(a')$ .*

*Proof.* If  $a_h = 1$ , then  $\rho = \mu^{(l)} \supseteq \lambda(\tilde{a}) = \lambda(a')$ , since  $l \in R(\mu, a)$ . Thus the assertion holds.

Suppose  $a_h \geq 2$ . Then  $l \leq s(\mu^{(l)}, \tilde{a})$  by Lemma 7, Lemma 16, and hence  $\rho \in \mathcal{B}(\mu, a)$  by Lemma 15. Thus it remains to show that  $\rho_l < \mu_l$ . This can be shown as follows:

$$\begin{aligned} \rho_l &= \begin{cases} \mu_l^{(l)} - (\tilde{a}_h - \mu_{l+1}^{(l)}) & \text{if } l = s(\mu^{(l)}, \tilde{a}), \\ \mu_l^{(l)} & \text{if } l < s(\mu^{(l)}, \tilde{a}) \end{cases} \\ &\leq \mu_l^{(l)} \\ &< \mu_l, \end{aligned}$$

where the second inequality follows from the definition of  $s(\mu^{(l)}, \tilde{a})$ .  $\square$

From Lemma 17, we obtain a tableau  $U \in \text{STab}(\mu, a)$ .

**Algorithm 2. Input:**  $\mu = (\mu_1, \mu_2, \dots, \mu_k) \vdash n$  and  $a = (a_1, a_2, \dots, a_h) \vDash n$  such that  $\mu \supseteq \lambda(a)$ .

**Output:**  $U \in \text{STab}(\mu, a)$ .

**Initialization:**  $\nu := \mu$ ,  $b := a$ .

**while**  $h(b) > 1$  **do**

$l := l(\nu, b)$ .

**if**  $b_{h(b)} = 1$  **then**  $\rho := \nu^{(l)}$ .

**else**

$\rho := \rho(\nu^{(l)}, \tilde{b})$ .

**end if**

$U(i, j) := h(b)$  **where**  $(i, j) \in D_{\nu/\rho}$ .

$\nu \leftarrow \rho$ ,  $b \leftarrow b'$ .

**end**

$U(1, j) := 1$  **where**  $1 \leq j \leq \nu_1$ .

Output  $U$ .

**Example 18.** Let  $\mu = (4, 4, 1, 1) \vdash 10$  and  $a = (1, 3, 2, 2, 2) \vDash 10$ . Then a tableau  $U \in \text{STab}(\mu, a)$  is obtained via Algorithm 2.

$\nu$	$b$	$\rho$	$U$
$(4, 4, 1, 1)$	$(1, 3, 2, 2, 2)$	$(4, 3, 1)$	$U(2, 4) = U(4, 1) = 5$
$(4, 3, 1)$	$(1, 3, 2, 2)$	$(3, 3)$	$U(1, 4) = U(3, 1) = 4$
$(3, 3)$	$(1, 3, 2)$	$(3, 1)$	$U(2, 2) = U(2, 3) = 3$
$(3, 1)$	$(1, 3)$	$(1)$	$U(1, 2) = U(1, 3) = U(2, 1) = 2$
$(1)$	$(1)$		$U(1, 1) = 1$

Thus

$$U = \begin{array}{cccc} 1 & 2 & 2 & 4 \\ 2 & 3 & 3 & 5 \\ 4 & & & \\ 5 & & & \end{array}.$$

We note that the least element of  $\text{STab}(\mu, a)$  is

$$S = \begin{array}{cccc} 1 & 2 & 2 & 4 \\ 2 & 3 & 5 & 5 \\ 3 & & & \\ 4 & & & \end{array}.$$

In Section 5, we will show that there exists a unique least element of  $\text{STab}(\mu, a)$  whenever  $\mu \trianglerighteq \lambda(a)$ , and give an algorithm to construct it.

**Proposition 19.** *Let  $\mu = (\mu_1, \mu_2, \dots, \mu_k) \vdash n$  and  $a = (a_1, a_2, \dots, a_h) \vDash n$ , and assume  $\mu \trianglerighteq \lambda(a)$ . Let  $r$  be an integer with  $1 \leq r \leq k$ . Then there exists a tableau  $T \in \text{STab}(\mu, a)$  such that  $T(r, \mu_r) = h$  if and only if  $r \in R(\mu, a)$ .*

*Proof.* The ‘‘only if’’ part has been proved in Lemma 8. We prove the ‘‘if’’ part by induction on  $n$ . If  $n = 1$  then it is obvious. Let  $r \in R(\mu, a)$ ,  $s = s(\mu, a)$  and  $s' = s(\mu^{(r)}, \tilde{a})$ .

If  $a_h \geq 2$  then define  $\rho \vdash n - a_h$  by

$$(15) \quad \rho = \begin{cases} \rho(\mu, a) & \text{if } r > s', \\ \rho(\mu^{(r)}, \tilde{a}) & \text{otherwise.} \end{cases}$$

From Lemma 12 and 15, we have  $\rho \in \mathcal{B}(\mu, a)$ , so

$$(16) \quad \rho \trianglerighteq \lambda(a').$$

If  $a_h = 1$ , then define by  $\rho = \mu^{(r)}$ . From the definition of  $R(\mu, a)$ , (16) holds in this case also.

Since (16) implies  $R(\rho, \lambda(a')) \neq \emptyset$ , the inductive hypothesis implies that there exists a tableau  $T' \in \text{STab}(\rho, a')$ . Define a tableau  $T$  of shape  $\mu$  and weight  $a$  by

$$T(i, j) = \begin{cases} T'(i, j) & \text{if } (i, j) \in D_\rho, \\ h & \text{if } (i, j) \in D_{\mu/\rho}. \end{cases}$$

It remains to show that  $T(r, \mu_r) = h$ . This will follow if we can show  $\rho_r < \mu_r$ . If  $a_h = 1$ , then  $\rho_r = \mu_r - 1 < \mu_r$ . Suppose  $a_h \geq 2$ . If  $r > s'$  then we have  $r > s' \geq s$  by

the definition of  $s'$  and  $s$ . Since  $r \in R(\mu, a)$ , we have  $\rho_r = \mu_{r+1} < \mu_r$ . If  $r \leq s'$ , then

$$\begin{aligned} \rho_r &= \begin{cases} \mu_r^{(r)} - (\tilde{a} - \mu_{r+1}^{(r)}) & \text{if } r = s', \\ \mu_r^{(r)} & \text{if } r < s' \end{cases} \\ &\leq \mu_r^{(r)} \\ &= \mu_r - 1 \\ &< \mu_r, \end{aligned}$$

where the second inequality follows from the definition of  $s'$ .  $\square$

Proposition 19 justifies the proof of [1, Lemma 3.7.3]. It also gives an alternative proof of the ‘‘if’’ part of Theorem 2.

### 5. THE LEAST ELEMENT OF $\text{STab}(\mu, a)$

Throughout this section, we let  $\mu = (\mu_1, \mu_2, \dots, \mu_k) \vdash n$  and  $a = (a_1, a_2, \dots, a_h) \vDash n$ . We assume  $\mu \supseteq \lambda(a)$ . For a sequence  $(i_1, i_2, \dots, i_j)$  of positive integers, we abbreviate the partition

$$(\dots((\mu^{(i_1)})^{(i_2)})\dots)^{(i_j)}$$

of  $n - j$ , as  $\mu^{(i_1, i_2, \dots, i_j)}$ .

**Lemma 20.** *We have*

$$R(\mu, a) = \{i \mid l(\mu, a) \leq i \leq k, \mu_i > \mu_{i+1}\}.$$

*Proof.* Write  $q = q(a)$ ,  $\lambda = \lambda(a)$  and  $l = l(\mu, a)$ . Then  $\lambda_q > \lambda_{q+1}$  by (1). Let  $l \leq i \leq k$  and  $\mu_i > \mu_{i+1}$ . From Lemma 4 and the definition of  $l$ , we have  $\mu^{(i)} \supseteq \mu^{(l)} \supseteq \lambda^{(a)}$ . Thus  $i \in R(\mu, a)$ .  $\square$

For each  $i$  with  $1 \leq i \leq k$ , set

$$R(\mu, a, i) = \{r \in R(\mu, a) \mid r \geq i\}.$$

From Lemma 20, we have  $k \in R(\mu, a, i)$  for each  $i$  with  $1 \leq i \leq k$ . Set

$$l(\mu, a, i) = \min R(\mu, a, i).$$

Clearly,  $l(\mu, a, 1) = l(\mu, a)$ .

**Lemma 21.** *Let  $\mu$  and  $\mu'$  be partitions of  $n$ . Suppose that  $i \in R(\mu, a)$  and  $i' \in R(\mu', a)$  satisfy  $i \leq i'$ ,  $\mu_{i'} \geq \mu'_{i'}$  and  $\mu_j = \mu'_j$  for all  $j$  with  $j > i'$ . Then*

$$R(\mu'^{(i')}, \tilde{a}, i') \subseteq R(\mu^{(i)}, \tilde{a}, i).$$

*Proof.* Let  $r \in R(\mu'^{(i')}, \tilde{a}, i')$ . Since  $i \leq i' \leq r$ , we have

$$(17) \quad \delta_{i,r} \leq \delta_{i',r},$$

and

$$(18) \quad \mu_{r+1}^{(i')} = \mu'_{r+1} = \mu_{r+1} = \mu_{r+1}^{(i)}.$$

Thus

$$\begin{aligned}
\mu_r^{(i)} &= \mu_r - \delta_{i,r} \\
&\geq \mu'_r - \delta_{i,r} \\
&\geq \mu'_r - \delta_{i',r} && \text{(by (17))} \\
&= \mu_r^{(i')} \\
&> \mu_{r+1}^{(i')} \\
&= \mu_{r+1}^{(i)} && \text{(by (18)).}
\end{aligned}$$

It remains show that  $\mu^{(i,r)} \supseteq \lambda(\tilde{a})$ . If  $1 \leq q < r$ , then

$$\sum_{j=1}^q \mu_j^{(i,r)} = \sum_{j=1}^q \mu_j^{(i)} \geq \sum_{j=1}^q \lambda(\tilde{a})_j \geq \sum_{j=1}^q \lambda(\tilde{a})_j,$$

since  $i \in R(\mu, a)$ . If  $r \leq q$ , then  $i' \leq q$ , and hence  $\sum_{j>q} \mu_j = \sum_{j>q} \mu'_j$ . Thus

$$\begin{aligned}
\sum_{j=1}^q \mu_j^{(i,r)} &= \sum_{j=1}^q \mu_j - 2 \\
&= \sum_{j=1}^q \mu'_j - 2 \\
&= \sum_{j=1}^q \mu_j^{(i',r)} \\
&\geq \sum_{j=1}^q \lambda(\tilde{a})_j,
\end{aligned}$$

by  $r \in R(\mu^{(i')}, \tilde{a})$ . □

**Lemma 22.** *Assume  $a_h \geq 2$ ,  $r \in R(\mu, a)$ , and  $r \leq s(\mu, a)$ . Then  $s(\mu, a) \leq s(\mu^{(r)}, \tilde{a})$ . In particular,  $R(\mu^{(r)}, \tilde{a}, r) \neq \emptyset$  and  $l(\mu^{(r)}, \tilde{a}, r) \leq s(\mu^{(r)}, \tilde{a})$ .*

*Proof.* Since  $a_h \geq 2$ , we have

$$\begin{aligned}
\tilde{a}_h &= a_h - 1 \\
&\leq a_h - \delta_{r, s(\mu, a)} \\
&\leq \mu_{s(\mu, a)} - \delta_{r, s(\mu, a)} \\
&= \mu_{s(\mu, a)}^{(r)}.
\end{aligned}$$

Thus  $s(\mu^{(r)}, \tilde{a}) \geq s(\mu, a) \geq r$ , and hence  $s(\mu^{(r)}, \tilde{a}) \in R(\mu^{(r)}, \tilde{a}, r)$  by Lemma 6. □

**Notation 23.** Let  $r \in R(\mu, a)$  and suppose  $r \leq s(\mu, a)$ . Define  $a^i$ ,  $l_i$  and  $\mu^i$  inductively by setting  $a^0 = a$ ,  $l_0 = r$ ,  $\mu^0 = \mu$  and for  $0 \leq i < n$ ,

$$\begin{aligned} a^{i+1} &= \widetilde{a}^i \vdash n - i - 1, \\ l_{i+1} &= \begin{cases} l(\mu^i, a^i, 1) & \text{if } i \in A, \\ l(\mu^i, a^i, l_i) & \text{otherwise,} \end{cases} \\ \mu^{i+1} &= (\mu^i)^{(l_{i+1})} \vdash n - i - 1, \end{aligned}$$

where  $A = \{a_h, a_h + a_{h-1}, \dots, a_h + \dots + a_2\}$ .

In order to check  $l_{i+1}$  and  $\mu^{i+1}$  are well-defined, we show

$$(19) \quad \mu^i \trianglerighteq \lambda(a^i) \quad (0 \leq i < n),$$

$$(20) \quad R(\mu^i, a^i, l_i) \neq \emptyset \quad (0 \leq i < n, i \notin A),$$

$$(21) \quad l_{i+1} \leq s(\mu^i, a^i) \quad (0 \leq i < n).$$

Indeed, (19)–(20) guarantee that  $l_{i+1}$  is defined as an element of  $R(\mu^i, a^i)$ , even when  $i \notin A$ , so  $\mu^{i+1}$  is also defined.

We prove (19)–(21) by induction on  $i$ . If  $i = 0$  then, as  $\mu \trianglerighteq \lambda(a)$ , (19) holds. Also, (20) holds since  $r \in R(\mu, a, r)$ . Since  $0 \notin A$ , we have  $l_1 = l(\mu, a, l_0) = r \leq s(\mu, a)$ . Thus (21) holds for  $i = 0$  as well.

Assume (19)–(21) hold for some  $i \in \{0, 1, \dots, n-2\}$ . Since  $l_{i+1} \in R(\mu^i, a^i)$ , (19) holds for  $i+1$ .

If  $i+1 \notin A$ , then  $a_{h(a^i)}^i \geq 2$ . Also  $l_{i+1} \leq s(\mu^i, a^i)$  by induction. Lemma 22 then implies  $R(\mu^{i+1}, a^{i+1}, l_{i+1}) \neq \emptyset$  and  $l_{i+2} \leq s(\mu^{i+1}, a^{i+1})$ , so (20)–(21) hold for  $i+1$  as well.

If  $i+1 \in A$ , then  $l_{i+2} = l(\mu^{i+1}, a^{i+1}, 1) = \min R(\mu^{i+1}, a^{i+1}) \leq s(\mu^{i+1}, a^{i+1})$  by Lemma 6. Thus (21) holds for  $i+1$  as well.

Clearly,  $\mu^i = \mu^{(l_1, \dots, l_i)}$ .

**Lemma 24.** Let  $T \in \text{STab}(\mu, a)$ . With reference to Notation 23, suppose  $T^{-1}(h) = \{(t_1, t'_1), (t_2, t'_2), \dots, (t_{a_h}, t'_{a_h})\}$  and  $r \leq t_1 \leq t_2 \leq \dots \leq t_{a_h}$ . Then  $l_i \leq t_i$  for  $1 \leq i \leq a_h$ . In particular,  $\mu^{(t_1, \dots, t_{a_h})} \trianglerighteq \mu^{a_h}$ .

*Proof.* We prove the assertion by induction on  $i$ . If  $i = 1$ , then  $l_1 = r \leq t_1$ .

Assume  $l_1 \leq t_1, \dots, l_i \leq t_i$  hold for some  $i$  with  $1 \leq i < a_h$ . We aim to show  $l_{i+1} \leq t_{i+1}$  by deriving

$$(22) \quad R(\mu^{(t_1, \dots, t_i)}, a^i, t_i) \subseteq R(\mu^i, a^i, l_i)$$

from Lemma 21. In order to do so, we need to verify the hypotheses of Lemma 21. By the definition of  $l_i$ , we have  $l_i \in R(\mu^{i-1}, a^{i-1})$ . Since the restriction of  $T$  to  $D_{\mu^{(t_1, \dots, t_{i-1})}}$  is an element of  $\text{STab}(\mu^{(t_1, \dots, t_{i-1})}, a^{i-1})$ , Lemma 8 implies  $t_i \in R(\mu^{(t_1, \dots, t_{i-1})}, a^{i-1})$ , and our inductive hypothesis shows  $l_i \leq t_i$ . Similarly, we have

$$(23) \quad t_{i+1} \in R(\mu^{(t_1, \dots, t_i)}, a^i, t_i).$$

Since  $l_p \leq t_p \leq t_i$  for  $1 \leq p \leq i-1$  by our inductive hypothesis,

$$\begin{aligned} \mu_{t_i}^{i-1} &= \mu_{t_i}^{(l_1, \dots, l_{i-1})} \\ &= \mu_{t_i} - |\{p \mid 1 \leq p \leq i-1, l_p = t_i\}| \end{aligned}$$

$$\begin{aligned} &\geq \mu_{t_i} - |\{p \mid 1 \leq p \leq i-1, t_p = t_i\}| \\ &= \mu_{t_i}^{(t_1, \dots, t_{i-1})}. \end{aligned}$$

Finally, for  $j > t_i$ , we have  $\mu_j^{i-1} = \mu_j^{(l_1, \dots, l_{i-1})} = \mu_j = \mu_j^{(t_1, \dots, t_{i-1})}$ , since  $l_p \leq t_p \leq t_i$  for  $1 \leq p \leq i-1$ . Therefore, we have verified all the hypotheses of Lemma 21, and we obtain (22).

Now

$$\begin{aligned} l_{i+1} &= l(\mu^i, a^i, l_i) \\ &= \min R(\mu^i, a^i, l_i) \\ &\leq \min R(\mu^{(t_1, \dots, t_i)}, a^i, t_i) && \text{(by (22))} \\ &\leq t_{i+1} && \text{(by (23)).} \end{aligned}$$

□

**Theorem 25.** *Let  $\mu = (\mu_1, \mu_2, \dots, \mu_k) \vdash n$  and  $a = (a_1, a_2, \dots, a_h) \vDash n$ , and suppose  $\mu \triangleright \lambda(a)$ . Let  $r \in R(\mu, a)$  and suppose  $r \leq s(\mu, a)$ . Define  $a^i$ ,  $l_i$  and  $\mu^i$  as in Notation 23. Define a tableau  $S$  of shape  $\mu$  and weight  $a$  by*

$$S(l_{i+1}, \mu_{l_{i+1}}^i) = t,$$

where  $0 \leq i < n$  and  $\sum_{j=1}^{t-1} a_j < n - i \leq \sum_{j=1}^t a_j$ . Then  $S$  is the least element of the subposet

$$(24) \quad \{T \in \text{STab}(\mu, a) \mid \min\{i \mid T(i, \mu_i) = h\} \geq r\}$$

In particular, if  $r = l(\mu, a)$ , then  $S$  is the least element of  $\text{STab}(\mu, a)$ .

*Proof.* Note that the tableau  $S$  is well-defined. Indeed, by the definition of  $\mu^i$ , we have

$$(25) \quad D_{\mu^i} = D_{\mu^{i+1}} \cup \{(l_{i+1}, \mu_{l_{i+1}}^i)\}.$$

So  $D_\mu = \{(l_{i+1}, \mu_{l_{i+1}}^i) \mid 0 \leq i < n\}$ .

Next, we prove the statement by induction on  $n$ . If  $n = 1$  then  $\mu = a = (1)$ , so it is obvious.

Assume that the statement holds for  $n - 1$ . We apply Notation 23 with  $r, \mu, a$  replaced by  $l_2, \nu = \mu^1, b = a^1$ , respectively. This is admissible since  $l_2 \in R(\mu^1, a^1) = R(\nu, b)$  and  $l_2 \leq s(\mu^1, a^1) = s(\nu, b)$  by (21). Define  $b^i$ ,  $l'_i$  and  $\nu^i$  inductively by setting  $b^0 = b$ ,  $l'_0 = l_2$ ,  $\nu^0 = \nu$  and for  $0 \leq i < n - 1$ ,

$$\begin{aligned} b^{i+1} &= \tilde{b}^i \vDash n - i - 2, \\ l'_{i+1} &= \begin{cases} l(\nu^i, b^i, 1) & \text{if } i \in B, \\ l(\nu^i, b^i, l'_i) & \text{otherwise,} \end{cases} \\ \nu^{i+1} &= (\nu^i)^{(l'_{i+1})} \vdash n - i - 2, \end{aligned}$$

where

$$B = \begin{cases} \{b_{h-1}, b_{h-1} + b_{h-2}, \dots, b_{h-1} + \dots + b_2\} & \text{if } a_h = 1, \\ \{b_h, b_h + b_{h-1}, \dots, b_h + \dots + b_2\} & \text{otherwise,} \end{cases}$$

$$b = \begin{cases} (b_1, \dots, b_{h-1}) & \text{if } a_h = 1, \\ (b_1, \dots, b_h) & \text{otherwise.} \end{cases}$$

Define a tableau  $\tilde{S}$  of shape  $\nu$  and weight  $b$  by

$$\tilde{S}(l'_{i+1}, \nu^i_{l'_{i+1}}) = t,$$

where  $\sum_{j=1}^{t-1} b_j < n - 1 - i \leq \sum_{j=1}^t b_j$ . By the inductive hypothesis,  $\tilde{S}$  is the least element of the set

$$(26) \quad \{\tilde{T} \in \text{STab}(\nu, b) \mid \min\{i \mid \tilde{T}(i, \nu_i) = h(b)\} \geq l_2\}.$$

It is easy to see that  $b^i = a^{i+1}$  for  $0 \leq i < n$ . We show that

$$(27) \quad l'_i = l_{i+1} \text{ and } \nu^i = \mu^{i+1} \quad (1 \leq i < n)$$

by induction on  $i$ . Since  $0 \notin B$ , we have

$$(28) \quad \begin{aligned} l'_1 &= l(\nu^0, b^0, l'_0) \\ &= l(\mu^1, a^1, l_2) \\ &= \begin{cases} l(\mu^1, a^1, l(\mu^1, a^1, 1)) & \text{if } 1 \in A, \\ l(\mu^1, a^1, l(\mu^1, a^1, l_1)) & \text{otherwise} \end{cases} \\ &= \begin{cases} l(\mu^1, a^1, 1) & \text{if } 1 \in A, \\ l(\mu^1, a^1, l_1) & \text{otherwise} \end{cases} \\ &= l_2. \end{aligned}$$

Then  $\nu^1 = (\nu^0)^{(l'_1)} = (\mu^1)^{(l_2)} = \mu^2$ .

Assume  $i \geq 2$  and  $l'_{i-1} = l_i$  and  $\nu^{i-1} = \mu^i$ . Since  $i-1 \in B$  if and only if  $i \in A$ , we have

$$\begin{aligned} l'_i &= \begin{cases} l(\nu^{i-1}, b^{i-1}, 1) & \text{if } i-1 \in B, \\ l(\nu^{i-1}, b^{i-1}, l'_{i-1}) & \text{otherwise} \end{cases} \\ &= \begin{cases} l(\mu^i, a^i, 1) & \text{if } i \in A, \\ l(\mu^i, a^i, l_i) & \text{otherwise} \end{cases} \\ &= l_{i+1}. \end{aligned}$$

Then  $\nu^i = (\nu^{i-1})^{(l'_i)} = (\mu^i)^{(l_{i+1})} = \mu^{i+1}$ .

Next we show

$$(29) \quad S|_{D_\nu} = \tilde{S}.$$

First, since  $b = a^1$ , we obtain

$$\sum_{i=1}^j b_i = \begin{cases} \sum_{i=1}^j a_i & \text{if } j < h, \\ \sum_{i=1}^h a_i - 1 & \text{if } j = h. \end{cases}$$

Suppose that  $\sum_{j=1}^{t-1} b_j < n - 1 - i \leq \sum_{j=1}^t b_j$ . Then

$$\sum_{j=1}^{t-1} a_j < n - (i + 1) \leq \sum_{j=1}^t a_j,$$

so  $\tilde{S}(l'_{i+1}, \nu'_{i+1}) = t = S(l_{i+2}, \mu_{i+2}^{i+1})$ . Thus, we have proved (29).

Next we show  $S \in \text{STab}(\mu, a)$ . If  $l_1 = 1$  then this is clear, since  $\tilde{S} \in \text{STab}(\nu, b)$ . Suppose  $l_1 \geq 2$ . Since  $(l_1 - 1, \mu_{l_1}) \in D_\nu$ , there exists an  $i \in \{1, 2, \dots, n - 1\}$  such that  $(l_1 - 1, \mu_{l_1}) = (l_{i+1}, \mu_{l_{i+1}}^i)$ . Since  $l_1 \leq l_2 \leq \dots \leq l_{a_h}$ , we have

$$i + 1 > a_h = n - \sum_{j=1}^{h-1} a_j = n - \sum_{j=1}^{h-1} b_j,$$

and hence

$$n - 1 - (i - 1) \leq \sum_{j=1}^{h-1} b_j.$$

This implies

$$(30) \quad \tilde{S}(l'_i, \nu'^{i-1}_i) \leq h - 1.$$

Now

$$\begin{aligned} S(l_1 - 1, \mu_{l_1}) &= S(l_{i+1}, \mu_{l_{i+1}}^i) \\ &= \tilde{S}(l'_i, \nu'^{i-1}_i) && \text{(by (27), (29))} \\ &< h && \text{(by (30))} \\ &= S(l_1, \mu_{l_1}). \end{aligned}$$

Since  $\tilde{S} \in \text{STab}(\nu, b)$ , this implies  $S \in \text{STab}(\mu, a)$ .

It remains to show that  $S \leq T$  for all  $T$  in the set (24). Define partitions  $\tau$  and  $\sigma$  by (5) and (6), respectively.

Suppose first that  $\min\{i \mid T(i, \mu_i) = h\} > l_1$ . Write  $T^{-1}(h) = \{(t_1, t'_1), \dots, (t_{a_h}, t'_{a_h})\}$  with  $l_1 < t_1 \leq t_2 \leq \dots \leq t_{a_h}$ . Then Lemma 24 implies  $\tau = \mu^{(t_1, \dots, t_{a_h})} \triangleright \mu^{a_h} = \sigma$ . Since  $l_1 < t_1$ , we have  $\tau \triangleright \sigma$ . Thus  $S \leq T$ .

Next suppose that  $\min\{i \mid T(i, \mu_i) = h\} = l_1$ . Set  $\tilde{T} = T|_{D_\nu}$  and observe  $\tilde{T} \in \text{STab}(\nu, b)$ . Set  $m = \min\{i \mid \tilde{T}(i, \nu_i) = h(b)\}$ . By Lemma 8, we have  $m \in R(\nu, b)$ , so  $m \geq l(\nu, b)$ . If  $a_h = 1$ , then  $l_2 = l(\nu, b)$ , so  $m \geq l_2$ . If  $a_h \geq 2$ , then  $h(b) = h$ , so  $m \geq l_1$ . Thus  $m \geq l(\nu, b, l_1) = l_2$ . Therefore,  $\tilde{T}$  belong to the set (26). This implies  $\tilde{S} \leq \tilde{T}$ , and hence either  $\tau \triangleright \sigma$ , or  $\tau = \sigma$  and  $\tilde{S}|_{D_\sigma} \leq \tilde{T}|_{D_\sigma}$ . Since  $\tilde{S}|_{D_\sigma} = S|_{D_\sigma}$  and  $\tilde{T}|_{D_\sigma} = T|_{D_\sigma}$ , the recursive definition of the partial order implies  $S \leq T$ .  $\square$

**Algorithm 3. Input:**  $\mu = (\mu_1, \mu_2, \dots, \mu_k) \vdash n$  and  $a = (a_1, a_2, \dots, a_h) \vDash n$  such that  $\mu \triangleright \lambda(a)$ .

**Output:**  $S \in \text{STab}(\mu, a)$ .

**Initialization:**  $\nu := \mu$ ,  $b := a$ ,  $m := n$  and  $l' := 1$ .

**while**  $m > 1$  **do**



$h := h(b)$  and  $l := l(\nu, b, l')$ .  
 $S(l, \nu_l) := h$ .  
**if**  $b_h = 1$ , **then**  $l' \leftarrow 1$ .  
**else**  $l' \leftarrow l$ .  
 $\nu \leftarrow \nu^{(l)}$ ,  $b \leftarrow \tilde{b}$  and  $m \leftarrow m - 1$ .  
**end**  
 $S(1, 1) := 1$ .  
Output  $S$ .

**Example 26.** Let  $\mu = (4, 4, 1, 1) \vdash 10$  and  $a = (1, 3, 2, 2, 2) \vDash 10$ . Then a tableau  $S \in \text{STab}(\mu, a)$  is obtained via Algorithm 3.

$\nu$	$b$	$m$	$l'$	$h$	$l$	$S$
(4, 4, 1, 1)	(1, 3, 2, 2, 2)	10	1	5	2	$S(2, 4) = 5$
(4, 3, 1, 1)	(1, 3, 2, 2, 1)	9	2	5	2	$S(2, 3) = 5$
(4, 2, 1, 1)	(1, 3, 2, 2)	8	1	4	1	$S(1, 4) = 4$
(3, 2, 1, 1)	(1, 3, 2, 1)	7	1	4	4	$S(4, 1) = 4$
(3, 2, 1)	(1, 3, 2)	6	1	3	2	$S(2, 2) = 3$
(3, 1, 1)	(1, 3, 1)	5	2	3	3	$S(3, 1) = 3$
(3, 1)	(1, 3)	4	1	2	1	$S(1, 3) = 2$
(2, 1)	(1, 2)	3	1	2	1	$S(1, 2) = 2$
(1, 1)	(1, 1)	2	1	2	2	$S(2, 1) = 2$
(1)	(1)	1	1			$S(1, 1) = 1$

Thus

$$S = \begin{array}{cccc} 1 & 2 & 2 & 4 \\ 2 & 3 & 5 & 5 \\ 3 & & & \\ 4 & & & \end{array}$$

and  $S$  is the least element of  $\text{STab}(\mu, a)$ .

**Remark 27.** Let  $a$  and  $b$  be compositions of  $n$  with  $\lambda(a) = \lambda(b)$ . Then there exists a bijection from  $\text{STab}(\mu, a)$  to  $\text{STab}(\mu, b)$  using [1, Lemma 3.7.1], but they are not isomorphic as partially ordered sets. For example, let  $\mu = (4, 4, 1, 1) \vdash 10$ ,  $a = (1, 3, 2, 2, 2) \vDash 10$  and  $b = (1, 2, 2, 2, 3) \vDash 10$ . Then  $(\text{STab}(\mu, b), \leq)$  is a totally ordered set, while  $(\text{STab}(\mu, a), \leq)$  contains two incomparable tableaux:

$$T = \begin{array}{cccc} 1 & 2 & 2 & 3 \\ 2 & 4 & 4 & 5 \\ 3 & & & \\ 5 & & & \end{array},$$

$$S = \begin{array}{cccc} 1 & 2 & 2 & 4 \\ 2 & 3 & 3 & 5 \\ 4 & & & \\ 5 & & & \end{array}.$$

Indeed, define  $\tau^{(p)}$  and  $\sigma^{(p)}$  by (3) and (4), respectively. Then  $\tau^{(3)} = (4, 1, 1)$  and  $\sigma^{(3)} = (3, 3)$  are incomparable. Thus  $(\text{STab}(\mu, a), \leq)$  is not totally ordered, hence it is not isomorphic to  $(\text{STab}(\mu, b), \leq)$ .

## REFERENCES

- [1] T. Ceccherini-Silverstein, F. Scarabotti and F. Tolli, Representation Theory of the Symmetric Groups, Cambridge University Press, 2010.
- [2] W. Fulton, Young Tableaux, London Mathematical Society Student Texts 35, 1997.
- [3] H. Narayanan, On the complexity of computing Kostka numbers and Littlewood-Richardson coefficients, J. Algebraic Combin. 24 (2006), 347–354.
- [4] B. Sagan, The Symmetric Group, Springer, 2001.

RESEARCH CENTER FOR PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCES, TOHOKU UNIVERSITY, SENDAI 980–8579, JAPAN  
*E-mail address:* `munemasa@math.is.tohoku.ac.jp`

RESEARCH CENTER FOR PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCES, TOHOKU UNIVERSITY, SENDAI 980–8579, JAPAN  
*E-mail address:* `minwon@ims.is.tohoku.ac.jp`