

MULTIWAVELENGTH EVIDENCE FOR QUASI-PERIODIC MODULATION IN THE GAMMA-RAY BLAZAR PG 1553+113

M. ACKERMANN¹, M. AJELLO², A. ALBERT³, W. B. ATWOOD⁴, L. BALDINI^{5,3}, J. BALLE⁶, G. BARBIELLINI^{7,8}, D. BASTIERI^{9,10}, J. BECERRA GONZALEZ^{11,12}, R. BELLAZZINI¹³, E. BISSALDI¹⁴, R. D. BLANDFORD³, E. D. BLOOM³, R. BONINO^{15,16}, E. BOTTACINI³, J. BREGEON¹⁷, P. BRUEL¹⁸, R. BUEHLER¹, S. BUSON^{9,10}, G. A. CALIANDRO^{3,19}, R. A. CAMERON³, R. CAPUTO⁴, M. CARAGIULO¹⁴, P. A. CARAVEO²⁰, E. CAVAZZUTI²¹, C. CECCHI^{22,23}, A. CHEKHTMAN²⁴, J. CHIANG³, G. CHIARO¹⁰, S. CIPRINI^{21,22,25,26*}, J. COHEN-TANUGI¹⁷, J. CONRAD^{27,28,29}, S. CUTINI^{21,25,22,30*}, F. D'AMMANDO^{31,32}, A. DE ANGELIS³³, F. DE PALMA^{14,34}, R. DESIANTE^{35,15}, L. DI VENERE³⁶, A. DOMÍNGUEZ², P. S. DRELL³, C. FAVUZZI^{36,14}, S. J. FEGAN¹⁸, E. C. FERRARA¹¹, W. B. FOCKE³, L. FUHRMANN³⁷, Y. FUKAZAWA³⁸, P. FUSCO^{36,14}, F. GARGANO¹⁴, D. GASPARRINI^{21,25,22}, N. GIGLIETTO^{36,14}, P. GIOMMI²¹, F. GIORDANO^{36,14}, M. GIROLETTI³¹, G. GODFREY³, D. GREEN^{12,11}, I. A. GRENIER⁶, J. E. GROVE³⁹, S. GUIRIEC^{11,40}, A. K. HARDING¹¹, E. HAYS¹¹, J. W. HEWITT⁴¹, A. B. HILL^{42,3}, D. HORAN¹⁸, T. JOGLER³, G. JÓHANNESSEN⁴³, A. S. JOHNSON³, T. KAMAE⁴⁴, M. KUSS¹³, S. LARSSON^{45,28,46*}, L. LATRONICO¹⁵, J. LI⁴⁷, L. LI^{45,28}, F. LONGO^{7,8}, F. LOPARCO^{36,14}, B. LOTT⁴⁸, M. N. LOVELLETTE³⁹, P. LUBRANO^{22,23}, J. MAGILL¹², S. MALDERA¹⁵, A. MANFREDA¹³, W. MAX-MOERBECK^{49,50}, M. MAYER¹, M. N. MAZZIOTTA¹⁴, J. E. MCENERY^{11,12}, P. F. MICHELSON³, T. MIZUNO⁵¹, M. E. MONZANI³, A. MORSELLI⁵², I. V. MOSKALENKO³, S. MURGIA⁵³, E. NUSS¹⁷, M. OHNO³⁸, T. OHSUGI⁵¹, R. OJHA¹¹, N. OMODEI³, E. ORLANDO³, J. F. ORMES⁵⁴, D. PANEQUE^{55,3}, T. J. PEARSON⁵⁰, J. S. PERKINS¹¹, M. PERRI²¹, M. PESCE-ROLLINS^{13,3}, V. PETROSIAN³, F. PIRON¹⁷, G. PIVATO¹³, T. A. PORTER³, S. RAINÒ^{36,14}, R. RANDO^{9,10}, M. RAZZANO^{13,56}, A. READHEAD⁵⁰, A. REIMER^{57,3}, O. REIMER^{57,3}, A. SCHULZ¹, C. SGRÒ¹³, E. J. SISKIND⁵⁸, F. SPADA¹³, G. SPANDRE¹³, P. SPINELLI^{36,14}, D. J. SUSON⁵⁹, H. TAKAHASHI³⁸, J. B. THAYER³, D. J. THOMPSON^{11,60*}, L. TIBALDO³, D. F. TORRES^{47,61}, G. TOSTI^{22,23}, E. TROJA^{11,12}, Y. UCHIYAMA⁶², G. VIANELLO³, K. S. WOOD³⁹, M. WOOD³, S. ZIMMER^{27,28}, A. BERDYUGIN⁶³, R. H. D. CORBET^{64,65}, T. HOVATTA⁶⁶, E. LINDFORS⁶³, K. NILSSON⁶⁷, R. REINTHAL⁶³, A. SILLANPÄÄ⁶³, A. STAMERRA^{68,69,70*}, L. O. TAKALO⁶³, M. J. VALTONEN⁶⁷

Accepted to The Astrophysical Journal Letters

ABSTRACT

We report for the first time a γ -ray and multiwavelength nearly-periodic oscillation in an active galactic nucleus. Using the *Fermi* Large Area Telescope (LAT) we have discovered an apparent quasi-periodicity in the γ -ray flux ($E > 100$ MeV) from the GeV/TeV BL Lac object PG 1553+113. The marginal significance of the 2.18 ± 0.08 year-period γ -ray cycle is strengthened by correlated oscillations observed in radio and optical fluxes, through data collected in the OVRO, Tuorla, KAIT, and CSS monitoring programs and *Swift* UVOT. The optical cycle appearing in ~ 10 years of data has a similar period, while the 15 GHz oscillation is less regular than seen in the other bands. Further long-term multi-wavelength monitoring of this blazar may discriminate among the possible explanations for this quasi-periodicity.

Subject headings: gamma rays: galaxies — gamma rays: general — BL Lacertae objects: general — BL Lacertae objects: individual (PG 1553+113) — galaxies: jets — accretion, accretion disks

¹ Deutsches Elektronen Synchrotron DESY, D-15738 Zeuthen, Germany

² Department of Physics and Astronomy, Clemson University, Kinard Lab of Physics, Clemson, SC 29634-0978, USA

³ W. W. Hansen Experimental Physics Laboratory, Kavli Institute for Particle Astrophysics and Cosmology, Department of Physics and SLAC National Accelerator Laboratory, Stanford University, Stanford, CA 94305, USA

⁴ Santa Cruz Institute for Particle Physics, Department of Physics and Department of Astronomy and Astrophysics, University of California at Santa Cruz, Santa Cruz, CA 95064, USA

⁵ Università di Pisa and Istituto Nazionale di Fisica Nucleare, Sezione di Pisa I-56127 Pisa, Italy

⁶ Laboratoire AIM, CEA-IRFU/CNRS/Université Paris Diderot, Service d'Astrophysique, CEA Saclay, F-91191 Gif sur Yvette, France

⁷ Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, I-34127 Trieste, Italy

⁸ Dipartimento di Fisica, Università di Trieste, I-34127 Trieste, Italy

⁹ Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-35131 Padova, Italy

¹⁰ Dipartimento di Fisica e Astronomia "G. Galilei", Università di Padova, I-35131 Padova, Italy

¹¹ NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

¹² Department of Physics and Department of Astronomy, University of Maryland, College Park, MD 20742, USA

¹³ Istituto Nazionale di Fisica Nucleare, Sezione di Pisa, I-56127 Pisa, Italy

¹⁴ Istituto Nazionale di Fisica Nucleare, Sezione di Bari, I-70126 Bari,

Italy

¹⁵ Istituto Nazionale di Fisica Nucleare, Sezione di Torino, I-10125 Torino, Italy

¹⁶ Dipartimento di Fisica Generale "Amadeo Avogadro", Università degli Studi di Torino, I-10125 Torino, Italy

¹⁷ Laboratoire Univers et Particules de Montpellier, Université Montpellier, CNRS/IN2P3, Montpellier, France

¹⁸ Laboratoire Leprince-Ringuet, École polytechnique, CNRS/IN2P3, Palaiseau, France

¹⁹ Consorzio Interuniversitario per la Fisica Spaziale (CIFS), I-10133 Torino, Italy

²⁰ INFN-Istituto di Astrofisica Spaziale e Fisica Cosmica, I-20133 Milano, Italy

²¹ Agenzia Spaziale Italiana (ASI) Science Data Center, I-00133 Roma, Italy

²² Istituto Nazionale di Fisica Nucleare, Sezione di Perugia, I-06123 Perugia, Italy

²³ Dipartimento di Fisica, Università degli Studi di Perugia, I-06123 Perugia, Italy

²⁴ College of Science, George Mason University, Fairfax, VA 22030, resident at Naval Research Laboratory, Washington, DC 20375, USA

²⁵ INFN Osservatorio Astronomico di Roma, I-00040 Monte Porzio Catone (Roma), Italy

²⁶ email: stefano.ciprini@asdc.asi.it

²⁷ Department of Physics, Stockholm University, AlbaNova, SE-106 91 Stockholm, Sweden

²⁸ The Oskar Klein Centre for Cosmoparticle Physics, AlbaNova, SE-106 91 Stockholm, Sweden

1. INTRODUCTION

Among active galactic nuclei (AGN), blazars are distinguished by erratic variability at all energies on a wide range of timescales. They are generally thought to be powered by supermassive black holes (SMBHs, 10^8 – $10^9 M_{\odot}$). PG 1553+113 (1ES 1553+113, $z \sim 0.49$, Danforth et al. 2010; Aliu et al. 2015; Abramowski et al. 2015) is an optically/X-ray selected BL Lac object (Falomo & Treves

1990) emitting variable GeV/TeV γ radiation (Aleksic et al. 2015; Abramowski et al. 2015). As typical in very-high energy (VHE) BL Lacs, the energetic non-thermal emission of PG 1553+113 originates in a relativistic jet and has a spectral energy distribution (SED) with two humps, overwhelming any other component from either the nucleus or the host galaxy.

The Large Area Telescope (LAT) on the *Fermi Gamma-ray Space Telescope* is providing continuous monitoring of the high-energy γ -ray sky. The apparent modulation noted in the γ -ray flux of PG 1553+113 stimulated the multi-frequency and long-term variability study described in this paper.

In §2 we describe the *Fermi* LAT data analysis and the sources of multiwavelength data; §3 details the multiple approaches used for lightcurves and cross-correlation analysis; §4 outlines preliminary scenarios to interpret these results.

2. FERMI LAT AND RADIO, OPTICAL, X-RAY DATA

The LAT is a pair conversion detector with a 2.4 sr field of view, sensitive to γ rays from ~ 20 MeV to > 300 GeV (Atwood et al. 2009). The present work uses the new Pass 8 LAT database (Atwood et al. 2013). The LAT operating mode allows it to cover the entire sky every two ~ 1.6 -hour spacecraft orbits, providing a regular and uniform view of γ -ray sources, sampling timescales from hours to years. This work uses observations of PG 1553+113 covering ~ 6.9 years (2008 August 4 to 2015 July 19, Modified Julian Day, MJD, 54682.65–57222.65). The LAT data analysis employed the standard `ScienceTools v10r0p5`⁷² package, selecting events from 100 MeV–300 GeV with `P8R2_SOURCE_V6` instrument response functions, in a circular Region of Interest of 10° radius centered on the position of PG 1553+113. It used files `gll_iem_v06` and `iso_P8R2_SOURCE_V6_v06` to model the Galactic and isotropic diffuse emission. Contamination due to the γ -ray-bright Earth limb is avoided by excluding events with zenith angle $> 90^{\circ}$. An unbinned maximum likelihood model fit technique is applied to each time bin with a power-law spectral model and photon index fixed to the 3FGL Catalog average value (1.604 ± 0.025 , Acero et al. 2015) for PG 1553+113. The resulting lightcurves are shown in Fig. 1.

Optical R-band data covering an interval of ~ 9.9 years (2005 April 19 to 2015 March 29, MJD 53479–57110) are reported in Fig. 2. Most unpublished observations were performed as part of the Tuorla blazar monitoring program (Takalo et al. 2008)⁷³. The data are reduced using a semi-automatic pipeline (Nilsson et al. in prep.). Public data from the Katzman Automatic Imaging Telescope (KAIT) and the Catalina Sky Survey (CSS) programs are also added. V-band magnitudes are scaled to the R-band values.

²⁹ The Royal Swedish Academy of Sciences, Box 50005, SE-104 05 Stockholm, Sweden

³⁰ email: sara.cutini@asdc.asi.it

³¹ INAF Istituto di Radioastronomia, I-40129 Bologna, Italy

³² Dipartimento di Astronomia, Università di Bologna, I-40127 Bologna, Italy

³³ Dipartimento di Fisica, Università di Udine and Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, Gruppo Collegato di Udine, I-33100 Udine

³⁴ Università Telematica Pegaso, Piazza Trieste e Trento, 48, I-80132 Napoli, Italy

³⁵ Università di Udine, I-33100 Udine, Italy

³⁶ Dipartimento di Fisica “M. Merlin” dell’Università e del Politecnico di Bari, I-70126 Bari, Italy

³⁷ Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

³⁸ Department of Physical Sciences, Hiroshima University, Higashi-Hiroshima, Hiroshima 739-8526, Japan

³⁹ Space Science Division, Naval Research Laboratory, Washington, DC 20375-5352, USA

⁴⁰ NASA Postdoctoral Program Fellow, USA

⁴¹ University of North Florida, Department of Physics, 1 UNF Drive, Jacksonville, FL 32224, USA

⁴² School of Physics and Astronomy, University of Southampton, Highfield, Southampton, SO17 1BJ, UK

⁴³ Science Institute, University of Iceland, IS-107 Reykjavik, Iceland

⁴⁴ Department of Physics, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

⁴⁵ Department of Physics, KTH Royal Institute of Technology, AlbaNova, SE-106 91 Stockholm, Sweden

⁴⁶ email: stefan@astro.su.se

⁴⁷ Institute of Space Sciences (IEEC-CSIC), Campus UAB, E-08193 Barcelona, Spain

⁴⁸ Centre d’Études Nucléaires de Bordeaux Gradignan, IN2P3/CNRS, Université Bordeaux 1, BP120, F-33175 Gradignan Cedex, France

⁴⁹ National Radio Astronomy Observatory (NRAO), Socorro, NM 87801, USA

⁵⁰ Cahill Center for Astronomy and Astrophysics, California Institute of Technology, Pasadena, CA 91125, USA

⁵¹ Hiroshima Astrophysical Science Center, Hiroshima University, Higashi-Hiroshima, Hiroshima 739-8526, Japan

⁵² Istituto Nazionale di Fisica Nucleare, Sezione di Roma “Tor Vergata”, I-00133 Roma, Italy

⁵³ Center for Cosmology, Physics and Astronomy Department, University of California, Irvine, CA 92697-2575, USA

⁵⁴ Department of Physics and Astronomy, University of Denver, Denver, CO 80208, USA

⁵⁵ Max-Planck-Institut für Physik, D-80805 München, Germany

⁵⁶ Funded by contract FIRB-2012-RBFR12PM1F from the Italian Ministry of Education, University and Research (MIUR)

⁵⁷ Institut für Astro- und Teilchenphysik and Institut für Theoretische Physik, Leopold-Franzens-Universität Innsbruck, A-6020 Innsbruck, Austria

⁵⁸ NYCB Real-Time Computing Inc., Lattingtown, NY 11560-1025, USA

⁵⁹ Department of Chemistry and Physics, Purdue University Calumet, Hammond, IN 46323-2094, USA

⁶⁰ email: David.J.Thompson@nasa.gov

⁶¹ Institutió Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain

⁶² 3-34-1 Nishi-Ikebukuro, Toshima-ku, Tokyo 171-8501, Japan

⁶³ Tuorla Observatory, University of Turku, FI-21500 Piikkiö, Finland

⁶⁴ Center for Research and Exploration in Space Science and Technology (CRESTT) and NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

⁶⁵ Department of Physics and Center for Space Sciences and Technology, University of Maryland Baltimore County, Baltimore, MD 21250, USA

⁶⁶ Aalto University, Metsähovi Radio Observatory, Metsähovintie 114, Kylmälä, Finland

⁶⁷ Finnish Centre for Astronomy with ESO (FINCA), University of Turku, FI-21500 Piikkiö, Finland

⁶⁸ INAF, Osservatorio Astronomico di Torino, I-10025 Pino Torinese (TO), Italy

⁶⁹ Scuola Normale Superiore, Piazza dei Cavalieri, 7, I-56126 Pisa, Italy

⁷⁰ email: stamerra@oato.inaf.it

* Correspondence: sara.cutini@asdc.asi.it; stefano.ciprini@asdc.asi.it; stefan@astro.su.se; stamerra@oato.inaf.it; David.J.Thompson@nasa.gov

⁷² <http://fermi.gsfc.nasa.gov/ssc/data/analysis/documentation/>

⁷³ users.utu.fi/kani/1m




FIG. 1.— *Fermi* LAT γ -ray lightcurves of PG 1553+113 over ~ 6.9 years, from 2008 August 4 to 2015 July 19. The lightcurve above 1 GeV is shown with a constant 45-day binning (top panel); two light curves above 100 MeV are shown, with 45- and 20-day binning (middle and bottom panels)

As part of an ongoing blazar monitoring program supporting *Fermi* (Richards et al. 2011), the Owens Valley Radio Observatory (OVRO) 40-m radio telescope has been observing PG 1553+113 continually (about every 1 to 23 days) since 2008 August. Figure 2 reports published 15 GHz lightcurves for the period from 2008 August 19 to 2014 May 18 (MJD 54697-56795). OVRO instrumentation, data calibration and reduction are described in Richards et al. (2011).

Swift observed PG 1553+113 110 times between 2005 April 20 and 2015 July 18 (unabsorbed 0.3–2 keV flux lightcurve in Figure 2). X-Ray Telescope (XRT) data were first calibrated and cleaned (xrtpipeline, XRTDAS v.3.0.0) and energy spectra extracted from a region of 20 pixel ($\sim 47''$) radius, with a nearby 20 pixel radius region for background. Individual XRT spectra are well fitted with a log-parabolic model, with column density fixed to the Galactic value of $3.6 \times 10^{20} \text{ cm}^{-2}$ (Kalberla et al. 2011). Aperture photometry ($5''$ radius) for the UVOT V-band filter was performed.

3. TEMPORAL VARIABILITY ANALYSIS AND CROSS CORRELATION ANALYSIS

We performed continuous wavelet transform (CWT) and Lomb-Scargle Periodogram (LSP) analyses on the lightcurves. Fig. 3 shows clear peaks at ~ 2 years for γ -ray and optical power spectra. We also made an epoch folding (pulse shape) analysis used to extract the period, shape, amplitude and phase, with uncertainties (Larsson 1996). The χ^2 for the folded pulse as a function of trial periods was fitted with a model containing 4 Fourier components, giving a period of 798 ± 30 days (2.18 ± 0.08 years), consistent with the CWT and LSP findings (Fig. 3). The value of the signal power peak does not change using regular 20-day and 45-day bins or an adaptive-bin technique (Lott et al. 2012) for construction of the LAT lightcurve.

A direct Power Density Spectrum (PDS) constructed from a LAT count-rate lightcurve using exposure-weighted aperture photometry (Corbet et al. 2007; Kerr 2011) above 100 MeV for a region with 3° radius with 600 second time bins (Fig.




FIG. 2.— Multifrequency lightcurves of PG 1553+113 at X-ray, optical and radio bands. Top panel: *Swift*-XRT integrated flux (0.3-2.0 keV). Central panel: optical flux density from Tuorla (R filter, black filled circle points), Catalina CSS (V filter rescaled, blue filled squared points), KAIT (V filter rescaled, red filled diamond points), *Swift*-UVOT (V filter rescaled, green filled circle points). Dotted line: LAT flux ($E > 100$ MeV) with time bins of 20 days, scaled and y-shifted for comparison. Bottom panel: 15 GHz flux density from OVRO 40m (black filled circle points) and parsec-scale 15 GHz flux density by VLBA (MOJAVE program, yellow diamond filled points).

4), confirms previous results with a peak at 2.16 ± 0.08 years, at $82\times$ the mean power level. The low-frequency modulation prevents an easy fit subtraction to the PDS continuum. The peak is ~ 5 times the mean level using a 4th order polynomial fit.

The significance of any apparent periodic variation depends on what assumption is made about spurious stochastic variability mimicking a periodic variation. The significance of the ~ 2 -year γ -ray periodicity is difficult to assess given the limited length of the γ -ray lightcurve. Red-noise, i.e. random and relatively enhanced low-frequency fluctuations over intervals comparable to the sample length, hinders the evaluation of periodicity significance (e.g. Hsieh et al. 2005; Lasky et al. 2015). We have approached the problem with two procedures:

1) The red-noise is assumed to be produced by similar amplitude flares (as seen in PG 1553+113 and some other LAT blazars), and the probability for these to line up in a regular pattern is estimated. The coherence of the periodic modulation was investigated by studying phase variations along the lightcurve. The local phase at each minimum and maximum

was estimated by correlating a one-period long data segment with the Fourier template of the full lightcurve. The rms variations relative to a perfectly coherent modulation was 27.4 days. The chance probabilities for 3, 4 and 5 random events to be distributed with at least this coherence, as estimated by Monte Carlo simulations, are 0.0535, 0.0105 and 0.0027 respectively, implying a chance probability of a few percent for the 3.5-peak γ -ray lightcurve of PG 1553+113.

2) We modeled the red-noise using Monte Carlo simulations with a first-order autoregressive process as the null hypothesis to assess whether the signal is consistent with a stochastic origin. Non-linear influence on the PDS is minimal thanks to the evenly spaced γ -ray lightcurve. The power peak in Fig. 3 is above the 99% confidence contour level, i.e. has $< 1\%$ chance of being a statistical fluctuation. The optical power peak has $< 5\%$ chance of being a statistical fluctuation.

Although the γ -ray periodicity signal alone is not compelling, the 9.9-years of optical data support the finding of a periodic oscillation in PG 1553+113. The optical data, although affected by seasonal gaps, were analyzed using the




FIG. 3.— Left top panel: pulse shape (epoch-folded) γ -ray ($E > 100$ MeV) flux lightcurve at the 2.18 year period (two cycles shown). Left bottom panels: 2D plane contour plot of the CWT power spectrum (scalogram) of the γ -ray lightcurve, using a Morlet mother function (filled color contour). The side panel to this is the 1D smoothed, all-epoch averaged, spectrum of the CWT scalogram showing a signal power peak in agreement with the 2.18-year value, also showing the LSP. Dashed lines depict increasing levels of confidence against red-noise calculated with Monte Carlo simulation. The γ -ray signal peak is above the 99% confidence contour level ($< 1\%$ chance probability of being spurious). Right top panel: pulse shape from epoch folding of the optical flux lightcurve at the 2.18 year period (two cycles shown). Right bottom panels: the same CWT and LSP diagrams for the optical lightcurve. The optical signal peak is above the 95% confidence contour level.

same techniques as for the γ -ray data. This analysis gives a period of 754 ± 20 days (2.06 ± 0.05 years), consistent within uncertainties with the γ -ray results (Fig. 3).




FIG. 4.— Power Density Spectrum (PDS) of the LAT 0.1 – 300 GeV count rate lightcurve of PG 1553+113 from a 3° exposure-weighted aperture photometry technique with 600-second time bins.

The less coherent 15 GHz lightcurve (5.7-years OVRO data) shows a signal power peak at 1.9 ± 0.1 year, with an

additional power component at a 1.2-year timescale. *Swift* XRT data show a factor of 5 variation linearly correlated with the γ -ray flux, while the synchrotron peak frequency shows a factor ~ 6 increase during high X-ray states, as suggested by Reimer et al. (2008).

The long-term X-ray count rate lightcurve from the *Rossini* XTE ASM instrument (1996 February 20 to 2010 September 11) and the *Swift*-BAT (from 2005 May 29) were also analyzed but do not show any signal above the low-frequency noise, because of insufficient statistics.

An important diagnostic for multi-frequency periodicity analysis is the discrete cross-correlation function (DCCF) used with two independent and complementary approaches.

In the first procedure, flux variations are modeled assuming a simple power law $\propto 1/f^\alpha$ (with $f = 1/t$) in the PDS as measured directly from the lightcurve data, allowing us to estimate the cross-correlations significance avoiding the assumption of equal variability in all sources at the cost of a model assumption (Max-Moerbeck et al. 2014). For the γ -ray lightcurve with 20-day binning we obtain a best fit $\alpha = 0.8$, but the error is unconstrained, indicating that the length of the data set is too short (i.e. below five cycles), relative to the suspected periodic modulation, to enable a reliable data characterization. The 45-day bin lightcurve yields a best fit $\alpha = 0.1$ with unconstrained error. The optical PSD is constrained: the best fit value is $\alpha = 1.85$, with 1σ limits at $[1.75, 2.00]$. The 15 GHz flux light curve a slope of $\alpha = 1.4$, with unconstrained limits on the α values as for the γ -ray data.




FIG. 5.— Discrete cross-correlation plots from the approach with PDS model measured from the lightcurve data (Max-Moerbeck et al. 2014). In each plot the black dots are the DCCF estimates, and the red, orange and green lines are the 1σ , 2σ and 3σ significance levels respectively. Top panel: DCCF between the radio 15GHz and γ -ray (20-day time bins) lightcurve. Central panel: DCCF between the unbinned optical lightcurve and γ -ray (20-day time bins) lightcurve. Bottom panel: DCCF between the 20-day rebinned optical lightcurve and γ -ray (20-day time bins) lightcurve. The oscillating shape of the significance contours for this case is due to the number of samples in each bin.

The DCCF between the unbinned radio lightcurve and the 20-day bin γ -ray lightcurve results in a most probable time lag for radio-flux lagging the γ -ray flux by 50 ± 20 days, with a 98.14% significance for the best PSD fit with a range of [89.56%-99.99%] when fit errors are taken into account (Fig. 5), using the fitting procedure of Max-Moerbeck et al. (2014). The DCCF between the unbinned optical lightcurve and the 20-day bin γ -ray lightcurve results in a most probable time lag for γ -ray flux lagging the optical flux by 130 ± 14 days, with a 99.14% significance for the best PSD fit and [96.09%-99.97%] when fit errors are taken into account (Fig. 5). The DCCF peak is broad, however, and consistent with no lag. This is also seen when the optical data are rebinned into 20-day intervals, as shown in the bottom panel, where the most probable lag is 10 ± 51 days.

In the second procedure, the significance of the γ -ray – radio correlation was estimated to be 95% by a mixed

source correlation procedure (Fuhrmann et al. 2014), cross-correlating the PG 1553+113 lightcurve with those of 132 comparison sources in that work, and evaluating the average DCCF level for time lags -100 to $+100$ days. The γ -ray – optical correlation is significant at the 99% level, even though partly limited by the number of comparison sources and optical lightcurve gaps. With only 132 comparison lightcurves we can measure a minimum probability-value of 0.0075, therefore in principle a 99% level of significance, but in this approach the error in that estimate is hard to determine. With the mixed source methods there are two limitations: 1) the assumption that all the sources can be described with the same model for the variability, and 2) the sample variance due to the limited number of lightcurves must be assessed. The optical flux is found to lead the γ -ray variations by 75 ± 27 days and the radio by 158 ± 10 days (γ -ray variations lead the 15 GHz flux variations by 83 ± 27 days). The possible reverse γ -ray-optical time lag decreases to 28 ± 27 days when the optical lightcurve is binned.

The possible optical- γ -ray lag was already pointed out by Cohen et al. (2014), using KAIT unbinned optical lightcurves and LAT data. The high degree of γ -ray-radio correlation in PG 1553+113 is not typically found in other individual blazars/AGN (see Max-Moerbeck et al. 2014). Significant cross-correlations are, nevertheless, found when stacking blazar samples (radio lagging γ rays; Fuhrmann et al. 2014).

4. DISCUSSION AND CONCLUSIONS

Factors that led to the indication of a possible ~ 2 -year periodic modulation in PG 1553+113 are: the continuous all-sky survey of *Fermi*; the increased capability of the new *Fermi* LAT Pass 8 data; and the long-term radio/optical monitoring of γ -ray blazars. Although the statistical significance of periodicity is marginal in each band, the consistent positive cross-correlation between bands strengthens the case, making PG 1553+113 the first possible quasi-periodic GeV γ -ray blazar and a prime candidate for further studies. Hints of possible γ -ray periodicities are rare in literature (for example Sandrinelli et al. 2014). The similarity of the low- and high-energy modulation in PG 1553+113 is also a novel behavior for AGN (Rieger 2004, 2007). Any periodic driving scenario should be related to the relativistic jet itself or to the process feeding the jet for this VHE BL Lac object. We outline, as examples, four possibilities:

1. Pulsational accretion flow instabilities, approximating periodic behavior, are able to explain modulations in the energy outflow efficiency. Magnetically-arrested and magnetically-dominated accretion flows (MDAF) could be suitable regimes for radiatively inefficient BL Lacs (Fragile & Meier 2009), characterized by advection-dominated accretion flows and subluminal, turbulent and peculiar radio kinematics (Karouzos et al. 2012; Piner & Edwards 2014). Such kinematics are sometimes explained as a precessing or helical jet (Conway & Murphy 1993). MDAF in a inner disc portion can be able to efficiently impart energy to particles in the jets of VHE BL Lacs (Tchekhovskoy et al. 2011). Periodic instabilities are believed to have short periods, $\sim 10^5 \text{ s} \cdot (M_{SMBH}/10^8 M_{\odot})$ (Honma et al. 1992), but MHD simulations of magnetically choked accretion flows are seen to produce longer periods for slow-spinning SMBH (McKinney et al. 2012).
2. Jet precession (e.g., Romero et al. 2000; Stirling et

al. 2003; Caproni et al. 2013), rotation (Camenzind & Krockenberger 1992; Vlahakis & Tsinganos 1998; Hardee & Rosen 1999) or helical structure (e.g., Conway & Murphy 1993; Roland et al. 1994; Villata & Raiteri 1999; Nakamura & Meier 2004; Ostorero et al. 2004), i.e. geometrical models (Rieger 2004), in the presence of a jet wrapped by a sufficiently strong magnetic field, could have a net apparent periodicity from the change of the viewing angle. Correspondingly the resulting Doppler magnification factor changes periodically without the need for intrinsic variation in outflows and efficiency. Non-ballistic hydrodynamical jet precession may explain variations with periods > 1 year (Rieger 2004). A differential Doppler factor $\Delta\mathcal{D}(t) = \Gamma^{-1}(1 - \beta(t) \cos \theta(t))^{-1} \lesssim 40\%$ variation (precession angle $\sim 1^\circ$) might be sufficient to support the ~ 2.8 amplitude flux modulation seen in γ rays. A homogeneous curved helical jet scenario for PG 1553+113 was proposed in Raiteri et al. (2015).

3. A mechanism analogous to low-frequency QPO from Galactic high-mass binaries/microquasars could produce an accretion-outflow coupling mechanism as the basis of the periodicity (Fender & Belloni 2004). King et al. (2013) ascribed the radio QPO in the FSRQ CGRaBS J1359+4011 to this mechanism. However BL Lac objects like PG 1553+113 are thought to possess a lower accretion rate. The microquasar QPO mechanism of Lense-Thirring precession (Wilkins 1972) requires that the inner accretion flow forms a geometrically thick torus rather than a standard thin disc as the latter warps (Bardeen-Petterson effect, Bardeen & Petterson 1975) rather than precesses (Ingram et al. 2009). A low mass accretion rate means that the accretion process probably forms an Advection-Dominated Accretion Flow (ADAF), so it can precess (Fragile & Meier 2009). The X-ray emission in PG 1553+113 is probably from the jet rather than from the flow, making it unlikely that the changing inclination of the hot flow causes the QPO. However, Lense-Thirring precession of the flow could affect the jet direction, giving the QPO as in (2) above.
4. The presence of a gravitationally bound binary SMBH system (Begelman et al. 1980; Barnes & Hernquist 1992) with a total mass $\sim 10^8 M_\odot$, and a milli-pc separation in the early inspiral gravitational-wave driven regime, might be another hypothesis. Keplerian binary orbital motion, would induce periodic accretion perturbations (Valtonen et al. 2008; Pihajoki et al. 2013; Liu et al. 2015) or jet nutation expected from the misalignment of the rotating SMBH spins, or the gravitational torque on the disc exerted by the companion (Katz 1997; Romero et al. 2000; Caproni et al. 2013; Graham et al. 2015). Significant acceleration of the disc evolution and accretion onto a binary SMBH system is depicted by modeling (Nixon et al. 2013; Doğan et al. 2015).

Binary SMBH induced periodicities have timescales

ranging from ~ 1 to ~ 25 years (Komossa 2006; Rieger 2007). The SMBH total mass in PG 1553+113, estimated utilizing the putative link between inflow/accretion (disc luminosity) and outflow/jet (jet power) in blazars (Ghisellini et al. 2014), is $\simeq 1.6 \times 10^8 M_\odot$, using a $0.1 M_{Edd}$ rate and Doppler factor $\mathcal{D} = 30$, in agreement with estimates for VHE BL Lacs (Woo et al. 2005).

The observed 2.18-year period is equivalent to an intrinsic orbital time $T'_{Kep} \leq T_{obs}/(1+z) \simeq 1.5$ years, and the binary system size would be 0.005 pc (~ 100 Schwarzschild radii). The probability to be observing such milli-pc system, estimated from the binary mass ratios $\sim 0.1 - 0.01$ and the GW-driven regime lifetime (Peters 1964), $t_{GW} \simeq 10^5 - 10^6$ years might be too small.

Periodicities claimed for AGN are often controversial; however PG 1553+113 may potentially represent a key γ -ray/multimessenger laboratory in the hypothesis of low-frequency gravitational wave emission and may have associated PeV neutrino emission (Padovani & Resconi 2014). VLBI structure observations, radio/optical polarization data, and a prolonged multifrequency monitoring campaign will shed light on the situation. If the periodic modulation is real and coherent, as would be expected for a binary scenario, then subsequent maxima would be expected in 2017 and 2019, well within the possible lifetime of the *Fermi* mission.

We thank the anonymous referee for useful and constructive comments. We extend special thanks to Prof. C. Done of Durham University, UK, and Prof. R. W. Romani of Stanford University, USA, for useful comments during the course of this work. The *Fermi*-LAT Collaboration acknowledges support for LAT development, operation and data analysis from NASA and DOE (United States), CEA/Irfu and IN2P3/CNRS (France), ASI and INFN (Italy), MEXT, KEK, and JAXA (Japan), and the K.A. Wallenberg Foundation, the Swedish Research Council and the National Space Board (Sweden). Science analysis support in the operations phase from INAF (Italy) and CNES (France) is also gratefully acknowledged. Tuorla blazar monitoring program has been partially supported by Academy of Finland grant 127740. KAIT telescope program is supported by Katzman Foundation and the National Science Foundation. The CSS survey is funded by the National Aeronautics and Space Administration under Grant No. NNG05GF22G issued through the Science Mission Directorate Near-Earth Objects Observations Program. The CRTS survey is supported by the U.S. National Science Foundation under grants AST-0909182. The OVRO 40-m program is supported in part by NASA grants NNX08AW31G and NNX11A043G and NSF grants AST-0808050 and AST-1109911. The MOJAVE program is supported under NASA-Fermi grant NNX12A087G. The National Radio Astronomy Observatory (NRAO) is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. The NASA *Swift* γ -ray burst explorer is a MIDEX Gamma Ray Burst mission led by NASA with participation of Italy and the UK. This research has made use of the Smithsonian/NASA's ADS bibliographic database. This research has made use of the NASA/IPAC NED database (JPL CalTech and NASA, USA). This research has made use of the archives and services of the ASI Science Data Center (ASDC), a facility of the Italian Space Agency (ASI Headquarters, Rome, Italy). This research has made use of the XRT Data Analysis Software (XRTDAS) developed under the responsibility of the ASDC. This work is a product of the ASDC Fermi team developed in the frame of the INAF Senior Scientists project and the foreign visiting scientists program of ASDC.

Facilities: *Fermi* Gamma-ray Space Telescope — *Swift* — *OVRO* — *Tuorla* — *KVA* — *KAIT* — *CSS* — *CRTS*

REFERENCES

Abramowski, A., Aharonian, F., Ait Benkhali, F., et al. 2015, *ApJ*, 802, 65
 Acero, F., Ackermann, M., Ajello, M., et al. 2015, *ApJS*, 218, 23
 Aleksić, J., Ansoldi, S., Antonelli, L. A., et al. 2015, *MNRAS*, 450, 4399

Aliu, E., Archambault, S., Archer, A., et al. 2015, *ApJ*, 800, 61
 Atwood, W. B., Abdo, A. A., Ackermann, M., et al. 2009, *ApJ*, 697, 1071

- Atwood, W., Albert, A., Baldini, L., et al. 2013, in eConf C121028, p.8, (arXiv:1303.3514)
- Bardeen, J. M., & Petterson, J. A. 1975, *ApJ*, 195, L65
- Barnes, J. E., Hernquist, L. 1992, *ARA&A*, 30, 705
- Begelman, M. C., Blandford, R. D., Rees, M. J., 1980, *Nature*, 287, 307
- Camenzind, M., Krockenberger, M., 1992, *A&A*, 255, 59
- Caproni, A., Abraham, Z., & Monteiro, H. 2013, *MNRAS*, 428, 280
- Cohen, D. P., Romani, R. W., Filippenko, A. V., et al. 2014, *ApJ*, 797, 137
- Conway, J. E., & Murphy, D. W. 1993, *ApJ*, 411, 89
- Corbet, R. H. D., Markwardt, C. B., Tueller, J. 2007, *ApJ*, 655, 458
- Danforth, C. W., Keeney, B. A., Stocke, J. T., Shull, J. M., & Yao, Y. 2010, *ApJ*, 720, 976
- Doğan, S., Nixon, C., King, A., & Price, D. J. 2015, *MNRAS*, 449, 1251
- Falomo, R., & Treves, A. 1990, *PASP*, 102, 1120
- Fender, R., & Belloni, T. 2004, *ARA&A*, 42, 317
- Fragile, P. C., & Meier, D. L. 2009, *ApJ*, 693, 771
- Fuhrmann, L., Larsson, S., Chiang, J., et al. 2014, *MNRAS*, 441, 1899
- Ghisellini, G., Tavecchio, F., Maraschi, L., Celotti, A., & Sbarrato, T. 2014, *Nature*, 515, 376
- Graham, M. J., Djorgovski, S. G., Stern, D., et al. 2015, *Nature*, 518, 74
- Hardee, P. E., & Rosen, A. 1999, *ApJ*, 524, 650
- Honma, F., Matsumoto, R., & Kato, S. 1992, *PASJ*, 44, 529
- Hsieh, C.-H., Glaser, S. M., Lucas, A. J., & Sugihara, G. 2005, *Nature*, 435, 336
- Kalberla, P. M. W., Burton, W. B., Hartmann, D., et al. 2005, *A&A*, 440, 775
- Karouzos, M., Britzen, S., Witzel, A., Zensus, J. A., & Eckart, A. 2012, *A&A*, 537, AA112
- Katz, J. I. 1997, *ApJ*, 478, 527
- Kerr, M. 2011, *ApJ*, 732, 38
- Kharb, P., Gabuzda, D., & Shastri, P. 2008, *MNRAS*, 384, 230
- King, O. G., Hovatta, T., Max-Moerbeck, W., et al. 2013, *MNRAS*, 436, L114
- Komossa, S. 2006, *Mem. Soc. Astron. Italiana*, 77, 733
- Ingram, A., Done, C., & Fragile, P. C. 2009, *MNRAS*, 397, L101
- Larsson, S. 1996, *A&AS*, 117, 197
- Lasky, P. D., Melatos, A., Ravi, V., & Hobbs, G. 2015, *MNRAS*, 449, 3293
- Liu, T., Gezari, S., Heinis, S., et al. 2015, *ApJ*, 803, L16
- Lott, B., Escande, L., Larsson, S., Ballet, J. 2012, *A&A* 544, A6
- Max-Moerbeck, W., Hovatta, T., Richards, J. L., et al. 2014, *MNRAS*, 445, 428
- McKinney, J. C., Tchekhovskoy, A., & Blandford, R. D. 2012, *MNRAS*, 423, 3083
- Mohan, P., & Mangalam, A. 2015, *ApJ*, 805, 91
- Nakamura, M., & Meier, D. L. 2004, *ApJ*, 617, 123
- Nixon, C., King, A., & Price, D. 2013, *MNRAS*, 434, 1946
- Ostorero, L., Villata, M., & Raiteri, C. M. 2004, *A&A*, 419, 913
- Padovani, P., & Resconi, E. 2014, *MNRAS*, 443, 474
- Peters, P. C. 1964, *PhRv*, 136, 1224
- Pihajoki, P., Valtonen, M. & Ciprini, S. 2013, *MNRAS*, 434, 3122
- Piner, B. G., & Edwards, P. G. 2014, *ApJ*, 797, 25
- Raiteri, C. M., Stamerra, A., Villata, M., et al. 2015, *MNRAS*, accepted, (arXiv:1509.02706)
- Reimer, A., Costamante, L., Madejski, G., Reimer, O., Dorner, D. 2008, *ApJ*, 682, 775
- Richards, J. L., Max-Moerbeck, W., Pavlidou, V., et al. 2011, *ApJS*, 194, 29
- Rieger, F. M. 2004, *ApJ*, 615, L5
- Rieger, F. M. 2007, *Ap&SS*, 309, 271
- Roland, J., Teyssier, R., & Roos, N. 1994, *A&A*, 290, 357
- Romero, G. E., Chajet, L., Abraham, Z., & Fan, J. H. 2000, *A&A*, 360, 57
- Sandrinelli, A., Covino, S., & Treves, A. 2014, *ApJ*, 793, LL1
- Stirling, A. M., Cawthorne, T. V., Stevens, J. A., et al. 2003, *MNRAS*, 341, 405
- Takalo, L. O., Nilsson, K., Lindfors, E., et al. 2008, *AIP Conf. Ser.* 1085, 705
- Tchekhovskoy, A., Narayan, R., McKinney, J. C. 2011, *MNRAS*, 418, L79
- Valtonen, M. J., Lehto, H. J., Nilsson, K., et al. 2008, *Nature*, 452, 851
- Villata, M., & Raiteri, C. M. 1999, *A&A*, 347, 30
- Vlahakis, N., & Tsinganos, K. 1998, *MNRAS*, 298, 777
- Wilkins, D. C. 1972, *Phys. Rev. D*, 5, 814
- Woo, J.-H., Urry, C. M., van der Marel, R. P., Lira, P., & Maza, J. 2005, *ApJ*, 631, 762