LENGTH OF AN INTERSECTION

CHRISTIAN DELHOMME AND MAURICE POUZET ´

Abstract. A poset p is well-partially ordered (WPO) if all its linear extensions are well orders ; the supremum of ordered types of these linear extensions is the *length*, $\ell(\mathbf{p})$ of p. We prove that if the vertex set X of **p** is infinite, of cardinality κ , and the ordering \leq is the intersection of finitely many partial orderings \leq_i on X, $1 \leq i \leq n$, then, letting $\ell(X, \leq_i) = \kappa \cdot q_i + r_i$, with $r_i < \kappa$, denote the euclidian division by κ (seen as an initial ordinal) of the length of the corresponding poset :

$$
\ell(\mathbf{p}) < \kappa \cdot \bigotimes_{1 \le i \le n} q_i + \bigg| \sum_{1 \le i \le n} r_i \bigg|^{+}
$$

where $|\sum r_i|^+$ denotes the least initial ordinal greater than the ordinal $\sum r_i$. This inequality is optimal (for $n \geq 2$).

1. INTRODUCTION

1.1. **Presentation of the result.** Let $\mathbf{p} := (X, \leq)$ be an ordered set(poset). This poset is *well founded* if every non-empty subset of the vertex set X contains some minimal element. It is *well partially ordered* (WPO for short) if, in addition, it has no infinite antichain *i.e.* if every infinite set of vertices has comparable elements. Since its introduction by Erdös and Rado and by Higman [\[12\]](#page-12-0) the notion of WPO has attracted considerable interest in various areas of mathematics and computer science (*e.g.* [\[21\]](#page-12-1), [\[15\]](#page-12-2)). It was observed by Wolk [\[23\]](#page-12-3) that a poset p is WPO if and only *if and only if* its linear extensions are all well orders. It has been proved by de Jongh and Parikh [\[8\]](#page-12-4) that there is a largest ordinal type of these linear extensions, that we denote $\ell(\mathbf{p})$ and call the *length of* \mathbf{p} (or of its ordering \leq). The length behaves nicely *w.r.t.* to some poset and ordinal operations. For an example, in [\[8\]](#page-12-4), de Jongh and Parikh extended Carruth formulas [\[2\]](#page-12-5) for direct sum and cartesian product of well ordered chains, showing that if p and q are two WPO's then the lengths of their direct sum $\mathbf{p} \uplus \mathbf{q}$ and of their cartesian product $\mathbf{p} \times \mathbf{q}$ (which are WPO) satisfy:

(1.1)
$$
\ell(\mathbf{p} \uplus \mathbf{q}) = \ell(\mathbf{p}) \oplus \ell(\mathbf{q}) \text{ and } \ell(\mathbf{p} \times \mathbf{q}) = \ell(\mathbf{p}) \otimes \ell(\mathbf{q})
$$

where ⊕ and ⊗ denote the Hessenberg addition and multiplication, also called natural operations. Since then, the ordinal length of various WPO has be computed ([\[22,](#page-12-6) [14,](#page-12-7) [19,](#page-12-8) [20\]](#page-12-9)).

In this note we consider posets **p** of which the ordering \leq is the intersection of finitely many orderings $\leq_i, 1 \leq i \leq n$, such that the corresponding posets \mathbf{p}_i are WPO. Since such a poset can be embedded into the direct product $\prod_{1 \leq i \leq n} p_i$, it is WPO and it follows from de Jongh-Parikh formula [\(1.1\)](#page-0-0) above that $\ell(\mathbf{p}) \leq \bigotimes_{1 \leq i \leq n} \ell(\mathbf{p}_i)$. This upper bound is crude. The best upper bound is given in the next theorem:

Theorem 1.1. *Given finitely many equipotent ordinals* α_i , $1 \leq i \leq n$, *of cardinality* κ , *consider their euclidian division by* κ : $\alpha_i = \kappa \cdot q_i + r_i$ *with* $r_i < \kappa$, and the least initial ordinal $\sum_{1 \leq i \leq n} r_i |$ ⁺ greater than $\sum_{1 \leq i \leq n} r_i$ *. Then :*

$$
\kappa \cdot \bigotimes_{1 \le i \le n} q_i + \big| \sum_{1 \le i \le n} r_i \big|^+
$$

is the least strict upper bound of the lengths of the posets $\mathbf{p} = (X, \leq)$ *of which the ordering* \leq *is the intersection of* n *orderings* \leq_i , $1 \leq i \leq n$, such that each (X, \leq_i) is a WPO of length α_i . This strict least upper bound is *unchanged if each ordering* \leq_i *is required to be linear.*

In particular :

Date: September 1, 2018.

Key words and phrases. Ordered sets, ordinal length, well quasi order, well partial ordering.

Proposition 1.1. If an ordering \leq on a set X of cardinality κ is the intersection of n orderings \leq _{*i*}, $1 \leq i \leq n$, then the length of $\mathbf{p} := (X, \leq)$ is at most $\kappa \cdot (\otimes_{1 \leq i \leq n} q_i)$ provided that each corresponding poset \mathbf{p}_i *have length at most* $\kappa \cdot q_i$ *. This bound is attained for a family of n linear orderings of types* $\kappa \cdot q_i$ *.*

As a consequence :

Corollary 1.1. *The length of a WPO* p *is an initial ordinal* κ *whenever the ordering of* p *is the intersection of finitely many orderings of length* κ*.*

Indeed, the length of a poset \bf{p} is trivially at least its cardinality (viewed as an initial ordinal), hence $\ell(\mathbf{p}) \geq \kappa$. Since in the theorem above, $q_i = 1$ and $r_i = 0, \kappa + 1$ is a strict upper bound of the length of p. Hence the length of **p** is κ .

As another consequence :

Corollary 1.2. Let **p** *be a poset and* α *be a countably infinite ordinal. Then* $\ell(\mathbf{p}) = \alpha$ *provided that the ordering is the intersection of an ordering of length* ω *and an ordering of length* α *.*

Note that, according to Theorem [1.1,](#page-0-1) this result does not hold with ω_1 in place of ω and $\alpha = \omega_1 + \delta$ with $\omega \leq \delta < \omega_1$. Indeed, for each β satisfying $\alpha \leq \beta \leq \omega_1$, there a poset of length β which is the intersection of a linear order of type ω_1 and a linear order of type α .

Theorem [1.1](#page-0-1) answers a question raised by Forster in [\[10\]](#page-12-10) in terms of the height of the tree of bad sequences of a poset **p** (see page 46). We recall that a sequence x_0, x_1, \ldots (finite or not) of vertices of a poset **p** is called *good* if there are indices i, j such that $i < j$ and $x_i \leq x_j$ and is called *bad* otherwise. The set of all bad sequences of **p** is denoted by $Bad(p)$. If we compare sequences by extension, this set becomes a tree with root r the empty sequence. As it is easy to see, this tree has no infinite chain if and only if p is WPO, hence in this case r has an height in $Bad(p)$. It turns out that this height is equal to $\ell(p)$ ([\[14\]](#page-12-7)).

Instead of the length of orderings which are intersections of orderings, the height of orderings which are unions of non necessarily transitive relations has been computed and an analogous formula obtained, see [\[1,](#page-12-11) [6,](#page-12-12) [7\]](#page-12-13).

Corollary [1.2](#page-1-0) was conjectured by Forster [\[10\]](#page-12-10) page 47.

1.2. Organisation of the paper. In the preliminary Section [2,](#page-2-0) we fix our notation on ordinals and WPO and we recall basic facts on those. The proof of Theorem [1.1](#page-0-1) is handled in Section [3.](#page-5-0) We first introduce specific notation regarding our particular purpose, and we restate our main result in this framework. We reduce this proof to the case $n = 2$ (Section [3.1\)](#page-5-1), and then the proof of this case $n = 2$ is split into a minoration in Section [3.2,](#page-7-0) and a majoration in Section [3.3.](#page-9-0)

Let us close this introductory section with some consequences of Corollary [1.1](#page-1-1) above, that were our initial motivation for the present work.

1.3. Sierpinskisation. Let α be a countable order type and ω be the order type of non-negative integers. A *sierpinskisation* of α and ω , or simply of α , is any poset $\mathbf{s} := (X, \leq)$ such that the ordering on X is the intersection of two linear orderings on X, one of type α , the other of type ω . Such a sierpinskisation can be obtained from a bijective map $\varphi : \omega \to \alpha$, setting $X := \mathbb{N}$ and $x \leq y$ if $x \leq y$ w.r.t. the natural ordering on N and $\varphi(x) \leq \varphi(y)$ w.r.t. the ordering of type α .

A consequence of Corollary [1.2](#page-1-0) is this:

Lemma 1.1. If α and α' are two countable ordinals with $\alpha < \alpha'$ then no sierpinskization of α' can be *embedded in any sierpinskization of* α *.*

Proof. If a sierpinskization s' of α' is embeddable into a sierpinskization s of α then by Corollary [1.2](#page-1-0) and [\(2.2\)](#page-3-0) of Section [2.1.3,](#page-3-1) $\alpha' = \ell(\mathbf{s}') \le \ell(\mathbf{s}) = \alpha$.

In contrast, we recall the following result (Lemma 3.4.1 of [\[17\]](#page-12-14)).

Lemma 1.2. For every countable order type α , if s and s' are a sierpinskizations of α and $\omega.\alpha$ respectively, *then* s *is embeddable into* s ′ *.*

As a special case, if α and $\omega.\alpha$ are equimorphic then any two sierpinskizations of α are equimorphic. If α is an ordinal, the equimorphy of α and $ω$.α amounts to $ω$ *ind*($α$) = $α$ (where *ind*($γ$) := 0 if $γ = 0$ and otherwise $ind(\gamma) := \delta$, where δ is the least non-zero ordinal such that $\gamma = \gamma' + \delta$ for some γ'). If $\beta \geq \omega$ then $\alpha := \omega^{\beta}$ satisfies this condition. Hence:

Corollary 1.3. *The* ω_1 -sequence $(\mathbf{s}_{\omega^\beta})_{\omega \leq \beta \leq \omega_1}$ of sierpinskizations of ω^β is strictly increasing w.r.t. embed*dability.*

To a poset $\mathbf{p} := (X, \leq),$ there are two natural posets associated with p, namely, the poset $I(\mathbf{p})$ of initial segments of **p** and the poset $I_{< \omega}(\mathbf{p})$ of finitely generated initial segments of **p**, that is the finite unions of principal initial segment of **p**, sets of the form $\downarrow x := \{y \in X : y \leq x\}$ for $x \in X$. If $s := (X, \leq)$ is a sierpinskization of α then each principal initial segment of **s** is finite, hence $I_{\leq \omega}(\mathbf{s})$ is a distributive lattice which is embeddable as a sublattice into $[\omega]^{<\omega}$, the lattice of finite subsets of ω , ordered by inclusion. If α is not an ordinal then s contains an infinite antichain. This implies that $[\omega]^{<\omega}$ is also embeddable in $I_{<\omega}(\mathbf{s})$ as a join-semilattice. For ordinals the situation is different: since s is WPO, $I_{\leq w}(\mathbf{s})$ is WPO via Higman's result [\[12\]](#page-12-0) on finite sequences, hence $[\omega]^{<\omega}$ is not embeddable in $I_{<\omega}(\mathbf{s})$.

Similarly to Corollary [1.3,](#page-2-1) the lattices $I_{\leq\omega}(\mathbf{s}_{\omega}\beta)$ form a ω_1 -chain w.r.t. join-semilattices embeddings and no member of this chain is embeddable in a previous member w.r.t. posets embeddings. Indeed:

Lemma 1.3. Let α , α' be two countably infinite ordinals and **s**, **s'** be two sierpinskizations of α and α' . If **s** *is order-embeddable into* s' *then* $I_{< w}(s)$ *is embeddable in* $I_{< w}(s')$ *as a join-semilattice, and this in turn implies that* $I(s)$ *is embeddable in* $I(s')$ *as a join-semilattice.* If $I(s)$ *is embeddable in* $I(s')$ *as a join-semilattice then* $\alpha \leq \alpha'$. The converse of these implications holds if $\omega \cdot \alpha \leq \alpha'$.

Proof. An order-embedding of **s** into **s'** yields a join-semilattice embedding of $I_{\leq \omega}(\mathbf{s})$ into $I_{\leq \omega}(\mathbf{s}')$. A joinsemilattice embedding of $I_{\leq \omega}(\mathbf{s})$ into $I_{\leq \omega}(\mathbf{s}')$ yields too a join-semilattice embedding of $I(\mathbf{s})$ into $I(\mathbf{s}')$. According to Corollary [1.2,](#page-1-0) $\ell(s) = \alpha$, hence the maximum length of chains in $I(s)$ is $\ell(s) + 1$, thus if $I(s)$ is embeddable into $I(s')$ as a join-semilattice, it is embeddable as a poset and in this case $\ell(s) + 1 \leq \ell(s') + 1$, thus $\alpha \leq \alpha'$. If $\omega \cdot \alpha \leq \alpha'$, then since s' induces a sierpinskization of $\omega \cdot \alpha$ and s embeds into this sierpinskization by Lemma [1.2,](#page-1-2) it embeds into s' .

Let β be an order type and $\alpha := \omega \beta$. A sierpinskization of α is called *monotonic* if for every $\gamma \in \alpha$, the map φ_{-1} , once restricted to $\omega \times {\gamma}$ is monotonic. According to Lemma 3.4.3. of [\[17\]](#page-12-14) two monotonic sierpinskization of α are equimorphic. Hence, we extend the conclusion of Corollary [1.3](#page-2-1) to monotonic sierpinskizations of ω .β for every $\beta < \omega$ as well as to the distributive lattices they generate.

For a detailed study of possible lengths of chains in algebraic lattices and sierpinskizations, see [\[3\]](#page-12-15) and [\[5,](#page-12-16) [4\]](#page-12-17).

Acknowledgement : Results of this paper have been presented at the International Conference on Discrete Mathematics and Computer Science (DIMACOS'11) organized by A. Boussaïri, M. Kabil, and A. Taik in Mohammedia (Morocco) May, 5-8, 2011. We are pleased to thank the organizers for their warmful invitation.

2. Preliminaries

2.1. Ordinals and WPO. Notation and basic properties. See [\[9\]](#page-12-18).

2.1.1. *Ordinals*. Given ordinals α and β :

- |α| denotes the cardinality of α , considered as the least ordinal that is equipotent with α . If $|\alpha| = \alpha$, then the ordinal α is *initial*.
- α^+ denotes the least initial ordinal greater than α (its so called Hartog).
- If $\alpha \leq \beta$, then $(-\alpha) + \beta$ denotes the only ordinal γ such that $\beta = \alpha + \gamma$. Thus for a non zero ordinal α , $(-1) + \alpha$ is equal to $n - 1$ if α is an integer n, and it is equal to α itself if α is infinite.
- For each non-zero ordinal γ , $\alpha = \delta \cdot q_{\delta}(\alpha) + r_{\delta}(\alpha)$ denote the euclidian division of α by δ , that is characterized by the remainder being less than δ : $r_{\delta}(\alpha) < \delta$.

⊕ and ⊗ denote the natural addition and multiplication on ordinals (also called Hessenberg operations).

For every set A of ordinals, let sup A denote its *supremum*, *i.e.* the least ordinal greater than or equal to every element of A , and let sup⁺ A denote its least strict upper-bound, *i.e.* least ordinal greater than every element of A. In particular, $\sup^+ \varnothing = \sup \varnothing = 0$ and $\sup^+ A = \sup \{\alpha + 1 : \alpha \in A\}.$

2.1.2. *Posets.* We view a *poset* (or *ordered set*) **p** as a pair (X, \leq) , where \leq is the *ordering* of the poset and X the *vertex set*. Thus an ordering of X is a set of ordered pairs of elements of X. If needed, we may denote by \leq_{p} the ordering of a poset **p**.

Given two posets $\mathbf{p} := (X, \leq_{\mathbf{p}})$ and $\mathbf{q} := (X, \leq_{\mathbf{q}})$ with the same vertex set X:

- $p \sqsubseteq q$ means that q is an (edge)-*extension* of p, *i.e.* that $x \leq_{p} y \Rightarrow x \leq_{q} y$.
- let $\mathbf{p} \sqcap \mathbf{q} := (X, \leq_{\mathbf{p}} \sqcap \leq_{\mathbf{q}})$ denote the intersection poset on X. Likewise is defined the intersection of any number of posets on a same set.

An *initial segment* Y of a poset $\mathbf{p} := (X, \leq)$ is any set $Y \subseteq X$ of vertices such that $x \leq x' \in X' \Rightarrow x \in X'$. Final segments are defined likewise.

An *order type* is an isomorphy type of posets. The *order type* of a well order p, is identified with the unique ordinal it is isomorphic to, and will be denoted $\tau(\mathbf{p})$.

When we write that an application f between ordered sets is \leq -increasing, we mean that $x \leq y \Rightarrow f(x) \leq$ $f(y)$.

Notation 2.1. If \prec is a binary relation on a set X, *e.g.* an ordering ≤, or the corresponding strict ordering $\langle \cdot \rangle$, *etc.*, then for each $x \in X$ and $Y \subseteq X$, we let :

$$
\{Y \prec x\} := \{y \in Y : y \prec x\}.
$$

If in addition **r** is a relational structure with vertex set X, *e.g.* of the form (X, \leq) , or $(X, >)$, or (X, \geq) , *etc.*, then we let :

$$
\{\mathbf r \prec x\} := \mathbf r \restriction \{y \in X : y \prec x\}
$$

denote the corresponding induced substructure.

2.1.3. *WPO.* Basics on WPO can be found in [\[16\]](#page-12-19). Recall that, if a poset **p** is a WPO, then $\ell(\mathbf{p})$ denotes its length. Notice that if **p** is a well order then $\ell(\mathbf{p}) = \tau(\mathbf{p})$.

WPO also admit the following characterization : a poset p is a WPO *if and only if* the collection of its initial segments is well founded under inclusion, and in this case the length of \bf{p} is equal to the height of its vertex set.

We shall use the following observations. Given a WPO $\mathbf{p} := (X, \leq)$:

• The length may be inductively computed $[8]$:

(2.1)
$$
\ell(\mathbf{p}) = \sup_{x \in X} \ell(\{\mathbf{p} \ngeq x\}).
$$

• If $X' \subseteq X$ then :

 (2.2)

$$
\ell(\mathbf{p} \restriction X') \le \ell(\mathbf{p}) \le \ell((\mathbf{p} \restriction X') \uplus (\mathbf{p} \restriction X \setminus X')) = \ell(\mathbf{p} \restriction X') \oplus \ell(\mathbf{p} \restriction X \setminus X').
$$

The middle inequality follows from **p** being an edge-extension of $(\mathbf{p} \restriction X') \oplus (\mathbf{p} \restriction X \setminus X')$, while the right-hand one is (1.1) . Incidentally, notice that if, in addition, X' is an initial segment, then $\ell(\mathbf{p} \restriction X') + \ell(\mathbf{p} \restriction X \setminus X') \leq \ell(\mathbf{p}).$

We shall also need the following lemmas :

Lemma 2.1. *Given a WPO* **p** *of length* α *, consider a decomposition* $\alpha = \alpha' + \alpha''$ *of this length. There is a partition of the domain* X *of* p *into an initial segment* X′ *and a final segment* X′′ *of* p *such that* $\ell(\mathbf{a} \restriction X') = \alpha'$ and $\ell(\mathbf{a} \restriction X'') = \alpha''$.

Proof. Given a linear extension $\mathbf{a} := (X, \preccurlyeq)$ of type α of \mathbf{p} , consider the initial segment Y' of \mathbf{a} of type α' , and let Y'' denote the complementary final segment, of type α'' . Thus $\ell(p \restriction Y') \geq \alpha'$ and $\ell(p \restriction Y'') \geq \alpha''$. So consider an initial segment X' of $p \restriction Y'$ such that $\ell(p \restriction X') = \alpha'$ and let $X'' := X \setminus X' \supseteq Y''$. In particular $\ell(\mathbf{p} \restriction X'') \geq \ell(\mathbf{p} \restriction Y'') = \alpha''$. Then, from :

$$
\alpha = \ell(\mathbf{p}) \ge \ell(\mathbf{p} \restriction X') + \ell(\mathbf{p} \restriction X'') = \alpha' + \ell(\mathbf{p} \restriction X'') \ge \alpha' + \alpha'' = \alpha
$$

it follows that $\alpha' + \ell(p \restriction X'') = \alpha' + \alpha''$, and therefore $\ell(p \restriction X'') = \alpha''$.

Lemma 2.2. *Consider a WPO* $p = (X, \leq)$ *of cardinality* κ *. The length of* p *is equal to* κ if and only if *every proper initial segment of* **p** *has cardinality less than* κ , if and only if *for every vertex* $x \in X$, the cardinality *of* $\{X \not\geq x\}$ *is less than* κ *.*

Proof. The second assertion is equivalent to the first one since the collection of $\{X \geq x\}$'s is coifinal in the collection of proper initial segments. As for this second equivalence, recall from [\(2.1\)](#page-3-2) that :

$$
\ell(\mathbf{p}) = \sup_{x \in X} \ell(\{X \not\geq x\}) \geq \sup_{x \in X_i} |\{X_i \not\geq_i x\}|
$$

given that the length of a poset is equipotent with its vertex set. \Box

Lemma 2.3. *Given a set* X, *of infinite cardinaliy* κ , *let us consider finitely many WQO* $\mathbf{p}_1, \ldots, \mathbf{p}_n$ *on* X *of respective lengths* $\alpha_1, \ldots, \alpha_n$. *If* β_1, \ldots, β_n *are ordinals of cardinality* κ *such that* $\alpha_1 \leq \beta_1, \ldots, \alpha_n \leq \beta_n$, *then there are a supserset* Y *of* X *and* n $WQO \mathbf{q}_1, \ldots, \mathbf{q}_n$ *on* Y *such that :*

$$
\mathbf{q}_1\restriction X=\mathbf{p}_1,\ldots,\mathbf{q}_n\restriction X=\mathbf{p}_n
$$

and :

$$
\ell(\mathbf{q}_1)=\beta_1,\ldots,\ell(\mathbf{q}_n)=\beta_n.
$$

In particular :

$$
\ell(\mathbf{p}_1\sqcap\cdots\sqcap\mathbf{p}_n)\leq\ell(\mathbf{q}_1\sqcap\cdots\sqcap\mathbf{q}_n).
$$

Furthermore, if the \mathbf{p}_i *'s are well orders one can choose the* \mathbf{q}_i *'s to be well orders too.*

Proof. For $i \in \{1, \ldots, n\}$, let \leq_i denote the ordering of \mathbf{p}_i . First observe that it can be assumed that there is a j such that $\beta_i = \alpha_i$ for each $i \neq j$. Indeed the general case follows from the succession of n applications of this particular case. And, without loss of generality, j can be assumed to be equal to 1. So let us assume that $\beta_i = \alpha_i$ for every $i \neq 1$.

Let $\gamma_1 := (-\alpha_1 + \beta_1)$, so that $\beta_1 = \alpha_1 + \gamma_1$. Let Z be a set of cardinality $|\gamma_1|$ disjoint from X and let $Y := X \cup Z$. Given a well ordering \preccurlyeq of type γ_1 on Z, consider the poset $\mathbf{q}_1 := \mathbf{p}_1 + (Z, \preccurlyeq)$. Thus $\mathbf{q}_1 \restriction X = \mathbf{p}_1$ and, clearly, $\ell(\mathbf{q}_1) = \ell(\mathbf{q}_1 \restriction X) + \gamma_1 = \alpha_1 + \gamma_1 = \beta_1$. If $n = 1$, then the proof is complete. So assume that $n \geq 2$.

Consider $i \in \{2,\ldots,n\}$. Letting $\delta_i := (-|\gamma_1|) + \alpha_i$, so that $|\gamma_1| + \delta_i = \alpha_i$, consider, with Lemma [2.3,](#page-4-0) an initial segment X_i of \mathbf{p}_i such that $\ell(\mathbf{p}_i \upharpoonright X_i) = |\gamma_1|$ and $\ell(\mathbf{p}_i \upharpoonright X \setminus X_i) = \delta_i$. In particular $\{X_i \not\geq_i x\}$ has cardinality less than $|\gamma_1|$ for each $x \in X_i$ (Lemma [2.2\)](#page-3-3). Then consider the lexicographical product poset $\mathbf{r}_i := (\{0,1\}, \leq) \cdot (\mathbf{p}_i \restriction X_i)$ on $\{0,1\} \times X_i$. Observe that $\ell(\mathbf{r}_i) = |\gamma_1|$ because of Lemma [2.2.](#page-3-3) Indeed for each $(\varepsilon, x) \in \{0, 1\} \times X_i, \{0, 1\} \times X_i \not\geq_{\mathbf{r}_i} (\varepsilon, x) \} \subseteq \{0, 1\} \times \{X_i \not\geq_i x\} \cup \{(0, x)\}\$ has cardinality less than $|\gamma_1|$.

Now given a bijection $f: X_i \to Z$, let us consider a poset $\mathbf{q}_i = (Y, \leq'_i)$, on $Y = X \dot{\cup} Z = X_i \dot{\cup} (X \setminus X_i) \dot{\cup} Z$ such that :

(1) $\mathbf{q}_i \restriction X = \mathbf{p}_i;$

(2) $\mathbf{q}_i \restriction X_i \dot{\cup} Z$ be isomorphic with \mathbf{r}_i through $(0, x) \mapsto f(x)$ and $(1, x) \mapsto x$;

(3) $X_i \dot{\cup} Z$ be an initial segment of \mathbf{q}_i .

Such a poset can be obtained, starting from p_i , by substituting a two vertex linear order for each element of X_i . We claim that $\ell(\mathbf{q}_i) = \beta_i$. Indeed :

$$
\alpha_i = \ell(\mathbf{p}_i) = \ell(\mathbf{q}_i \upharpoonright X) \le \ell(\mathbf{q}_i) \le \ell(\mathbf{q}_i \upharpoonright (X_i \cup Z)) + \ell(\mathbf{q}_i \upharpoonright (X \setminus X_i)) = |\gamma_1| + \delta_i = \alpha_i = \beta_i.
$$

2.2. Conventions regarding ordinal operations. For each finitary operation ϕ on ordinals, we may consider terms of the form $\phi_{\#}(\cdots)$ of which the arguments are ordinals and underlined ordinals. Such a term denotes the least ordinal strictly greater than the evaluation of the expression obtained by replacing $\phi_{\#}$ by ϕ and each argument by a non-greater ordinal, by a lesser one if this argument is underlined; *e.g.* :

$$
\phi_{\#}(\alpha_1,\ldots,\alpha_m,\underline{\beta_1},\ldots,\underline{\beta_n}) := \sup^+ \{ \phi(\alpha'_1,\ldots,\alpha'_m,\beta'_1,\ldots,\beta'_n) : \newline \alpha'_1 \leq \alpha_1,\ldots,\alpha'_m \leq \alpha_m,\beta'_1 < \beta_1,\ldots,\beta'_n < \beta_n \}.
$$

Notice that if no argument is underlined and ϕ is \leq -increasing in each variable, then $\phi_{\#}(\cdots) = \phi(\cdots) + 1$.

Now we introduce a finitary operation ϕ on ordinals with the same arity as ϕ :

$$
\underline{\phi}(\alpha_1, \dots, \alpha_n) := \phi_{\#}(\underline{\alpha_1}, \dots, \underline{\alpha_n})
$$

= $\sup^+ \{ \phi(\alpha'_1, \dots, \alpha'_n) : \alpha'_1 < \alpha_1, \dots, \alpha'_n < \alpha_n \}$
:= $\sup \{ \phi(\alpha'_1, \dots, \alpha'_n) + 1 : \alpha'_1 < \alpha_1, \dots, \alpha'_n < \alpha_n \}.$

The following observations are easily checked :

- If ϕ is an associative binary operation, then ϕ is associative.
- If ϕ_n is a n-ary operation obtained from an associative binary operation ϕ_2 , then ϕ_2 is associative and ϕ_n is obtained from ϕ_2 .

In particular the expression $\alpha_1 \underline{\oplus} \cdots \underline{\oplus} \alpha_n$ is not ambiguous; namely :

$$
\alpha_1 \underline{\oplus} \cdots \underline{\oplus} \alpha_n = \sup^{\dagger} {\alpha'_1 \oplus \cdots \oplus \alpha_n : \alpha'_1 \leq \alpha_1, \ldots, \alpha'_m \leq \alpha_m}.
$$

Observe that an ordinal α is *indecomposable*, *i.e.*, it is not the sum of two lesser ordinals *if and only if* $\alpha \oplus \alpha = \alpha$. In particular every initial ordinal is indecomposable. We shall need the following distributivity property :

(2.3)
$$
\alpha \oplus \alpha = \alpha \Rightarrow \alpha \cdot (\beta \oplus \gamma) = (\alpha \cdot \beta) \oplus (\alpha \cdot \gamma).
$$

3. Proof of Theorem [1.1](#page-0-1)

Let us intoduce the following finitary operations on ordinals :

$$
\varphi^+_{P}(\alpha_1,\ldots,\alpha_n):=\sup^{\ast}\{\ell(\mathbf{p}_1\sqcap\cdots\sqcap\mathbf{a}_n):\mathbf{a}_1,\ldots,\mathbf{a}_n\text{ WPO},\ \ell(\mathbf{a}_1)=\alpha_1,\ldots,\ell(\mathbf{a}_n)=\alpha_n\}.
$$

Relativizing this operation to WPO's that are linear, *i.e.* to WLO's, we also let :

$$
\varphi_L^+(\alpha_1,\ldots,\alpha_n):=\sup\{\ell(\mathbf{a}_1\sqcap\cdots\sqcap\mathbf{a}_n):\mathbf{a}_1,\ldots,\mathbf{a}_n\text{ WLO},\,\tau(\mathbf{a}_1)=\alpha_1,\ldots,\tau(\mathbf{a}_n)=\alpha_n\}.
$$

Obviously, $\varphi_P^+(\alpha_1,\ldots,\alpha_n) = \varphi_L^+(\alpha_1,\ldots,\alpha_n) = 0$ whenever α_1,\ldots,α_n do not have the same cardinality. Also, $\varphi_L^+(\alpha) = \varphi_P^+(\alpha) = \alpha + 1$ for every ordinal α , and $\varphi_L^+(k,\ldots,k) = \varphi_P^+(k,\ldots,k) = k+1$ if $k < \omega$, thus we will not need to consider the case $n = 1$ nor the case of finite ordinals.

Besides, $\varphi_L^+ \leq \varphi_P^+$; in fact, we shall see that equality holds. Indeed, in terms of these two operations, Theorem [1.1](#page-0-1) can be rephrased as follows :

Theorem 3.1. $\varphi_P^+ = \varphi_L^+$ and for every $n \geq 2$:

(3.1)
$$
\varphi_P^+(\alpha_1,\ldots,\alpha_n)=\kappa\cdot (q_\kappa(\alpha_1)\otimes\cdots\otimes q_\kappa(\alpha_n))+|r_\kappa(\alpha_1)+\cdots+r_\kappa(\alpha_n)|^+
$$

provided that the arguments be equipotent, of common infinite cardinality κ .

In Section [3.1](#page-5-1) below we reduce the general case to the particular case $n = 2$. The minoration of φ_P^+ is proved in Section [3.2](#page-7-0) (Proposition [3.2\)](#page-9-1) and the majoration in Section [3.3](#page-9-0) (Proposition [3.3\)](#page-11-0).

3.1. The derivation of the general case from the case $n = 2$.

3.1.1. *Auxiliary operations.* For this reduction, we introduce two auxiliary operations $\tilde{\varphi}_P$ and $\tilde{\varphi}_L$. Let :

$$
\widetilde{\varphi}_P(\alpha_1,\ldots,\alpha_n):=\operatorname{sup} \{\ell(\mathbf{a}_1\sqcap\cdots\sqcap\mathbf{a}_n):\mathbf{a}_1,\ldots,\mathbf{a}_n\text{ WPO},\ell(\mathbf{a}_1)<\alpha_1,\ldots,\ell(\mathbf{a}_n)<\alpha_n\}
$$

and :

$$
\widetilde{\varphi}_L(\alpha_1,\ldots,\alpha_n):=\sup^+\{\ell(\mathbf{a}_1\sqcap\cdots\sqcap\mathbf{a}_n):\mathbf{a}_1,\ldots,\mathbf{a}_n\text{ WLO},\,\tau(\mathbf{a}_1)<\alpha_1,\ldots,\tau(\mathbf{a}_n)<\alpha_n\}.
$$

Obviously $\widetilde{\varphi}_L(\alpha) = \widetilde{\varphi}_P(\alpha) = \alpha$ and in general $\widetilde{\varphi}_L \leq \widetilde{\varphi}_P$.

3.1.2. The strategy of reduction to the case $n = 2$. The reduction will rely on the following observations. Letting $\widetilde{\varphi}_{P,n}$, $\widetilde{\varphi}_{L,n}$ denote the restrictions of these operations to *n* variables :

- $\tilde{\varphi}_{P,2}$ and $\tilde{\varphi}_{L,2}$ are associative and $\tilde{\varphi}_{P,n}, \tilde{\varphi}_{L,n}$ are their extensions by associativity.
- $\widetilde{\varphi}_{P,n}$ and $\varphi_{P,n}^+$ are recoverable from one another (and likewise for $\widetilde{\varphi}_{L,n}$ and $\varphi_{L,n}^+$).
- Letting θ^+ denote the operation corresponding to the right-hand member of [\(3.1\)](#page-5-2), consider the operation $\widetilde{\theta}$ defined from θ^+ as $\widetilde{\varphi}_{P,n}$ is definable $\varphi_{P,n}^+$; then $\widetilde{\theta}_2$ is associative and $\widetilde{\theta}$ is its extension by associativity.
- The theorem in the case $n = 2$ precisely says that $\varphi_{L,2}^+ = \theta_2^+ = \varphi_{P,2}^+$.

Note that $\hat{\theta}$ will not have to be explicited.

3.1.3. *The reduction.* So let us proceed to this reduction. First observe that Lemma [2.3](#page-4-0) yields :

Lemma 3.1. *Given two finite tuples* $(\alpha_1, \ldots, \alpha_n)$, and $(\alpha'_1, \ldots, \alpha'_n)$ *of equipotent ordinals with respective cardinalities* κ *and* κ' *and such that* $\alpha'_1 \leq \alpha_1, \ldots, \alpha'_n \leq \alpha_n$:

$$
\varphi_P^+(\alpha'_1,\ldots,\alpha'_n) \leq \varphi_P^+(\alpha_1,\ldots,\alpha_n) \text{ and } \varphi_L^+(\alpha'_1,\ldots,\alpha'_n) \leq \varphi_L^+(\alpha_1,\ldots,\alpha_n).
$$

Notation 3.1. Given a finitary operation ϕ on ordinals, let $[\phi]^+$ and $[\phi]^\sim$ denote the following two operations :

$$
[\phi]^+(\alpha_1,\ldots,\alpha_n) = \begin{cases} \phi(\alpha_1+1,\ldots,\alpha_n+1) & \text{if the } \alpha_i\text{'s are equipotent} \\ 0 & \text{otherwise} \end{cases}
$$

and :

$$
[\phi]^\sim(\alpha_1,\ldots,\alpha_n) = \sup^+_{\alpha_1} \{ \alpha : \exists \alpha'_1 < \alpha_1,\ldots,\alpha'_n < \alpha_n \ \alpha < \phi(\alpha'_1,\ldots,\alpha'_n) \}
$$
\n
$$
= \sup\{ \phi(\alpha'_1,\ldots,\alpha'_n) : \alpha'_1 < \alpha_1,\ldots,\alpha'_n < \alpha_n \}.
$$

Corollary 3.1. Each operation φ_P^+ and $\widetilde{\varphi}_P$, resp. φ_L^+ and $\widetilde{\varphi}_L$, can be recovered from the other one as *follows :* $\varphi_P^+ = [\tilde{\varphi}_P]^+, \varphi_L^+ = [\tilde{\varphi}_L]^+, \tilde{\varphi}_P = [\varphi_P^+]^\sim$ and $\tilde{\varphi}_L^- = [\varphi_L^+]^\sim$.

Corollary 3.2. For each n, $\varphi_{P,n}^+ = \varphi_{L,n}^+$ is equivalent to $\widetilde{\varphi}_{P,n} = \widetilde{\varphi}_{L,n}$.

Let us now come to the associtivity relations.

Lemma 3.2. *For any positive integers* m *and* n *:*

(3.2)
$$
\widetilde{\varphi}_P(\alpha_1,\ldots,\alpha_m,\beta_1,\ldots,\beta_n) \leq \widetilde{\varphi}_P(\widetilde{\varphi}_P(\alpha_1,\ldots,\alpha_m),\widetilde{\varphi}_P(\beta_1,\ldots,\beta_n))
$$

and, dually :

(3.3)
$$
\widetilde{\varphi}_L(\alpha_1,\ldots,\alpha_m,\beta_1,\ldots,\beta_n) \geq \widetilde{\varphi}_L(\widetilde{\varphi}_L(\alpha_1,\ldots,\alpha_m),\widetilde{\varphi}_L(\beta_1,\ldots,\beta_n)).
$$

Proof. For Inequality [\(3.2\)](#page-6-0), just write each $\mathbf{p}_1 \sqcap \cdots \sqcap \mathbf{p}_m \sqcap \mathbf{q}_1 \sqcap \cdots \sqcap \mathbf{q}_n$ as $(\mathbf{p}_1 \sqcap \cdots \sqcap \mathbf{p}_m) \sqcap (\mathbf{q}_1 \sqcap \cdots \sqcap \mathbf{q}_n)$. Let us then handle Inequality [\(3.3\)](#page-6-1). Consider the case of equipotent α_i 's and β_j 's. Observe that, according to Lemma [3.3](#page-6-2) below :

$$
\ell(\mathbf{a} \sqcap \mathbf{b}) \leq \ell(\underbrace{\mathbf{a}_1 \sqcap \cdots \sqcap \mathbf{a}_m \sqcap \mathbf{b}_1 \sqcap \cdots \sqcap \mathbf{b}_n}_{\sqsubseteq \mathbf{a} \sqcap \mathbf{b}}) < \widetilde{\varphi}_L(\alpha_1, \ldots, \alpha_m, \beta_1, \ldots, \beta_n).
$$

Thus $\tilde{\varphi}_L(\alpha_1,\ldots,\alpha_m,\beta_1,\ldots,\beta_n)$ is a strict upper bound of the lengths of the intersections of two well orderings of types less than $\tilde{\varphi}_L(\alpha_1, \ldots, \alpha_m)$ and $\tilde{\varphi}_L(\beta_1, \ldots, \beta_n)$ respectively, and therefore it is greater than or equal to $\tilde{\varphi}_L(\tilde{\varphi}_L(\alpha_1, \ldots, \alpha_m), \tilde{\varphi}_L(\beta_1, \ldots, \beta_n))$, that is the least of these stric equal to $\tilde{\varphi}_L(\tilde{\varphi}_L(\alpha_1,\ldots,\alpha_m),\tilde{\varphi}_L(\beta_1,\ldots,\beta_n))$, that is the least of these strict upper bounds.

Lemma 3.3. *For any two equipotent well orders* a *and* b *with the same vertex set* X *and of respective types :*

$$
\tau(\mathbf{a}) = \alpha < \widetilde{\varphi}_L(\alpha_1, \dots, \alpha_m) \text{ and } \tau(\mathbf{b}) = \beta < \widetilde{\varphi}_L(\beta_1, \dots, \beta_n)
$$

there are well orders \mathbf{a}_i 's and \mathbf{b}_j 's on X such that :

$$
\tau(\mathbf{a}_i) < \alpha_i \ \text{and} \ \tau(\mathbf{b}_j) < \beta_j
$$

and

$$
\mathbf{a}_1 \sqcap \cdots \sqcap \mathbf{a}_m \sqsubseteq \mathbf{a} \text{ and } \mathbf{b}_1 \sqcap \cdots \sqcap \mathbf{b}_n \sqsubseteq \mathbf{b}.
$$

Proof. Since α is not a strict upper bound of the set of which $\tilde{\varphi}_L(\alpha_1, \ldots, \alpha_m)$ is defined to be the least strict upper bound, it is less than or equal to an element of that set. So there are well orders a_i 's on a common vertex set X' with $\tau(\mathbf{a}'_i) < \alpha_i$ and such that $\ell(\mathbf{a}'_1 \cap \cdots \cap \mathbf{a}'_m) \ge \alpha$. Then consider a linear extension \mathbf{a}' of $\mathbf{a}'_1 \sqcap \cdots \sqcap \mathbf{a}'_m$ of type at least α , and consider $X'' \subseteq X'$ such that $\tau(\mathbf{a}' \restriction X'') = \alpha$. Then consider the bijection $f: X'' \to X$ that is an isomophism from $\mathbf{a}' \restriction X''$ to \mathbf{a} , and let each \mathbf{a}_i be the well order on X onto which f is an isomorphism from $\mathbf{a}'_i \restriction X''$. Incidentally note that indeed $\mathbf{a}_1 \cap \cdots \cap \mathbf{a}_m = \mathbf{a}$.

Perform likewise *w.r.t.* b.

From Lemma [3.2](#page-6-3) a straightforward induction yields :

Corollary 3.3. *If* $\widetilde{\varphi}_{P,2} = \widetilde{\varphi}_{L,2}$ *then* $\widetilde{\varphi}_{P,n} = \widetilde{\varphi}_{L,n}$ *for every integer* $n \geq 2$ *.*

In order to complete our derivation of the general case from the case $n = 2$, it remains to check :

Lemma 3.4. *If* [\(3.1\)](#page-5-2) *in Theorem [3.1](#page-5-3) holds for* $n = 2$ *, then it holds for any* n.

Proof. Consider the finitary operation $\tilde{\theta} := [\theta^+]^{\sim}$ on ordinals defined from :

$$
\theta^+ : (\alpha_1, \dots, \alpha_n) \mapsto \begin{cases} \kappa \cdot (q_\kappa(\alpha_1) \otimes \dots \otimes q_\kappa(\alpha_n)) + |r_\kappa(\alpha_1) + \dots + r_\kappa(\alpha_n)|^+ & \text{if } |\alpha_1| = \dots = |\alpha_n|, \text{ let } \kappa \\ 0 & \text{otherwise} \end{cases}
$$

 $(n \geq 2)$. As above $\theta^+ = [\theta]^+$, because θ^+ satisfies the monotonicity condition of Lemma [3.1.](#page-6-4)

To infer $\varphi^+ = \theta^+$ from $\varphi_2^+ = \theta_2^+$ (by φ^+ here, we mean φ_R^+ or φ_L^+), it suffices to check that $\tilde{\theta}_2$ is associative with $\hat{\theta}_n$ equal to its extension by associativity to n variables. Given that θ^+ is commutative, it suffices to check that $[\widetilde{\theta}_2(\widetilde{\theta}_n, \text{id}_1)]^+ = \theta_{n+1}^+$, which we do now. Below we assume that the arguments are equipotent (written $\alpha_i \sim \beta$) of cardinal κ , and we just write q and r for q_{κ} and r_{κ} .

$$
\begin{aligned}\n&\left[\tilde{\theta}_{2}(\tilde{\theta}_{n}, id_{1})\right]^{+}(\alpha_{1}, \ldots, \alpha_{n}, \beta) \\
&= \tilde{\theta}_{2}(\tilde{\theta}_{n}(\alpha_{1} + 1, \ldots, \alpha_{n} + 1), \beta + 1) \\
&= \tilde{\theta}_{2}(\theta_{n}^{+}(\alpha_{1}, \ldots, \alpha_{n}), \beta + 1) \\
&= \sup\{\theta_{2}^{+}(\alpha, \beta) : \alpha < \theta_{n}^{+}(\alpha_{1}, \ldots, \alpha_{n})\} \\
&= \sup\{\theta_{2}^{+}(\kappa \cdot (q(\alpha_{1}) \otimes \cdots \otimes q(\alpha_{n})) + \gamma, \kappa \cdot q(\beta) + r(\beta)) : \gamma \sim r(\alpha_{1}) + \cdots + r(\alpha_{n})\} \\
&= \sup\{\kappa \cdot ((q(\alpha_{1}) \otimes \cdots \otimes q(\alpha_{n})) \otimes q(\beta)) + |\gamma + r(\beta)|^{+} : \gamma \sim r(\alpha_{1}) + \cdots + r(\alpha_{n})\} \\
&= \kappa \cdot (q(\alpha_{1}) \otimes \cdots \otimes q(\alpha_{n}) \otimes q(\beta)) + |r(\alpha_{1}) + \cdots + r(\alpha_{n}) + r(\beta)|^{+} \\
&= \theta_{n+1}^{+}(\alpha_{1}, \ldots, \alpha_{n}, \beta).\n\end{aligned}
$$

3.1.4. *Strategy for the case* $n = 2$. We shall prove :

Proposition 3.1. *Given two equipotent ordinals* α *and* β *, of cardinality* κ :

(3.4)
$$
\varphi_P^+(\alpha,\beta) \leq \kappa \cdot (q_\kappa(\alpha) \otimes q_\kappa(\beta)) + |r_\kappa(\alpha) + r_\kappa(\beta)|^+ \leq \varphi_L^+(\alpha,\beta).
$$

As a consequence, we shall get :

Corollary 3.4.

and

$$
\varphi^+_{P,2}=\varphi^+_{L,2}
$$

$$
\varphi_P^+(\alpha,\beta) = \kappa \cdot (q_\kappa(\alpha) \otimes q_\kappa(\beta)) + |r_\kappa(\alpha) + r_\kappa(\beta)|^+
$$

if the arguments are equipotent, of common cardinality κ*.*

The left-hand inequality in Proposition [3.1](#page-7-1) is Proposition [3.3](#page-11-0) and the right-hand one is Proposition [3.2.](#page-9-1)

3.2. Minoration. The proof of the minoration of φ^+ is performed in two steps. We first consider the case of two arguments that are multiples of their common cardinality, and then we reduce the general case to this one.

 \Box

3.2.1. *Minoration for arguments multiples of their cardinality.* Given an infinite cardinal κ viewed as an initial ordinal, and two order types α and β of cardinality at most κ , let us consider three posets **k**, a and **b** of respective order types α , β and κ , and vertex sets K, A and B. Let $\mathbf{p} = (P, \leq)$ denote the cartesian product $(\mathbf{k} \cdot \mathbf{a}) \times (\mathbf{k} \cdot \mathbf{b})$. Thus $P = (K \times A) \times (K \times B)$.

We are interested in orders of the form $p \restriction R$, and in particular in those for wich $R \subseteq (K \times A) \times (K \times B)$ is the graph of a bijection between $K \times A$ and $K \times B$. Indeed we shall see (Corollary [3.5\)](#page-8-0) that if R is mixing (see below) and α and β are types of well orders, then **p** \restriction R has length at least $\kappa \cdot (\alpha \otimes \beta)$ and is an intersection of two well orders of respective types $\kappa \cdot \alpha$ and $\kappa \cdot \beta$.

Given $R \subseteq P$, let us say that R is *functional*, resp. *partial functional*, if it is the graph of a function, resp. partial function of $K \times A$ to $K \times B$, in other words if each vertical section $R \cap (\{(k, a)\} \times (K \times B))$ has exactly one element, resp. at most one element. Say that R is *co-functional*, resp. *partial co-functional* if each horizontal section $R \cap ((K \times A) \times \{(k, b)\})$ has exactly one element, resp. at most one element. Then say that R is *bi-functional*, resp. *partial bi-functional* if it is both fonctional and co-functional, resp. both partial functional and partial co-functional. Observe that R is bi-functional *if and only if* this is the graph of a bijection between $K \times A$ and $K \times B$, and that it is partial bi-functional *if and only if* this is the graph of a one-to-one partial function.

For $R \subseteq P$, and $(a, b) \in A \times B$, we let :

$$
R_a^b := \{(k_1, k_2) : ((k_1, a), (k_2, b)) \in R\} \subseteq K \times K.
$$

Note that, if R is partial functional, resp. partial co-functional, then so is each R_a^b (with the obvious extensions of the definitions).

Say that R is *mixing* if each R_a^b has cardinality κ .

Lemma 3.5. *Consider* $R \subseteq (K \times A) \times (K \times B)$.

- (1) If R is partial bi-functional then the poset $p \restriction R$ is the intersection of two orders of types at most $\kappa \cdot \alpha$ *and* $\kappa \cdot \beta$ *.*
- (2) If R is partial functional and mixing then the poset $p \restriction R$ has an (edge)-extension of type $\kappa \cdot (\alpha \times \beta)$.

If R is partial bi-functional and mixing then the poset $p \restriction R$ is the intersection of two orders of types $\kappa \cdot \alpha$ *and* $\kappa \cdot \beta$ *and has an (edge)-extension of type* $\kappa \cdot (\alpha \times \beta)$ *.*

Proof.

- (1) The ordering of **p** is the intersection of the two lexicographical products of $\mathbf{k} \cdot \mathbf{a}$ and $\mathbf{k} \cdot \mathbf{b}$ (the left one and the right one). If R is partial bi-functional then the restrictions to R of these two lexicographical products have types at most $\kappa \cdot \alpha$ and $\kappa \cdot \beta$. They have types exactly $\kappa \cdot \alpha$ and $\kappa \cdot \beta$ if in addition R is mixing.
- (2) Consider the mapping $((k_1, a), (k_2, b)) \rightarrow (k_1, (a, b))$. Its restriction to any partial functional R is increasing from $p \restriction R$ to $k \cdot (a \times b)$. If, in addition, R is mixing, then the image of R has type $\kappa \cdot (\alpha \times \beta).$

Lemma 3.6. *There exists a mixing bi-functional* $R \subseteq P$ *.*

Proof. Consider two partitions $(K_a : a \in A)$ and $(K^b : b \in B)$ of K into classes of size κ each. For each $(a, b) \in A \times B$, consider the graph $G_a^b \subseteq K \times K$ of a bijection between K^b and K_a . Then let :

$$
R := \{ ((k_1, a), (k_2, b)) : (k_1, k_2) \in G_a^b : a \in A, b \in B \} \subseteq P.
$$

Then observe that R is bi-functional and mixing (with $R_a^b = G_a^b$). $\qquad \qquad \Box$

In the case α and β are types of well orders, these lemma yield :

Corollary 3.5. *Consider an infinite cardinal* κ *and two ordinals* α *and* β *of cardinality at most* κ*. Then there is a WPO* r *of length at least* $\kappa \cdot (\alpha \otimes \beta)$ *and that is the intersection of two well orders of types* $\kappa \cdot \alpha$ *and* $\kappa \cdot \beta$ *respectively. In particular :*

$$
\varphi_L^+(\kappa \cdot \alpha, \kappa \cdot \beta) > \kappa \cdot (\alpha \otimes \beta).
$$

 \Box

Proof. Consider $\mathbf{r} := \mathbf{p} \restriction R$ for R bi-functional and mixing, as in Lemma [3.5.](#page-8-1) Thus, on the one hand r is the intersection of two well orders of types $\kappa \cdot \alpha$ and $\kappa \cdot \beta$ respectively, and on the other hand it has an edge-extension of isomorphy type $\kappa \cdot (\alpha \times \beta)$, so that $\ell(\mathbf{r}) > \ell(\kappa \cdot (\alpha \times \beta))$. Besides :

$$
\ell(\kappa \cdot (\alpha \times \beta)) \geq \kappa \cdot \ell(\alpha \times \beta) = \kappa \cdot (\alpha \otimes \beta).
$$

The last equality is Carruth's result, *cf.* [\(1.1\)](#page-0-0), while the inequality is easy to check : an extension of $\kappa \cdot (\alpha \times \beta)$ can be obtained as the ordinal product of κ by any extension of $\alpha \times \beta$.

3.2.2. *Minoration. General case.*

Lemma 3.7. *Consider ordinals* $\alpha_1, \ldots, \alpha_n$ *and* β_1, \ldots, β_n *such that* $|\alpha_1| = |\beta_1|, \ldots, |\alpha_n| = |\beta_n|$. *Then (observe the ordering of the* β_i *'s)*:

(3.5)
$$
\varphi_L^+(\alpha_1 + \cdots + \alpha_n, \beta_n + \cdots + \beta_1) \geq \varphi_L^+(\alpha_1, \beta_1) \underline{\oplus} \cdots \underline{\oplus} \varphi_L^+(\alpha_n, \beta_n).
$$

Proof. For each $i \in \{1, ..., n\}$, let X_i , be n pairwise disjoint sets of cardinalities $\kappa_i := |\alpha_i| = |\beta_i|$; let \mathbf{a}_i and \mathbf{b}_i be two well ordered sets of type α_i and β_i with vertex set X_i and let $\mathbf{p}_i := \mathbf{a}_i \sqcap \mathbf{b}_i$. The lexicographical sums $\mathbf{a} := \mathbf{a}_1 + \cdots + \mathbf{a}_n$ and $\mathbf{b} := \mathbf{b}_n + \cdots + \mathbf{b}_1$ on $X := \cup_{1 \leq i \leq n} X_i$ have order type $\alpha := \alpha_1 + \cdots + \alpha_n$ and $\beta := \beta_n + \cdots + \beta_1$ and the ordered set $\mathbf{p} := \mathbf{a} \cap \mathbf{b}$ is the direct sum $\mathbf{p}_1 \oplus \cdots \oplus \mathbf{p}_n$. Hence, according to de Jongh-Parikh formula, $\ell(\mathbf{p}) = \ell(\mathbf{p}_1) \oplus \cdots \oplus \ell(\mathbf{p}_n)$.

Summing up, the typical member, namely $\ell(p_1) \oplus \cdots \oplus \ell(p_n)$, of the set of ordinals of which the right-hand side of [\(3.5\)](#page-9-2) is the least strict upper bound belongs to the set of ordinals $\ell(p)$ of which the left-hand side of [\(3.5\)](#page-9-2) is the least strict upper bound. Inequality (3.5) between these least strict upper bounds follows. \Box

Proposition 3.2. *Given two equipotent infinite ordinals* α *and* β *, of cardinality* κ *:*

(3.6)
$$
\varphi_L^+(\alpha,\beta) \geq \kappa \cdot (q_\kappa(\alpha) \otimes q_\kappa(\beta)) + |r_\kappa(\alpha) + r_\kappa(\beta)|^+.
$$

Proof. Observe that, for any ordinals γ equipotent to $r_{\kappa}(\alpha)$ and δ equipotent to $r_{\kappa}(\beta)$:

- $\alpha = \delta + \kappa \cdot q_{\kappa}(\alpha) + r_{\kappa}(\alpha)$, and
- $\beta = \gamma + \kappa \cdot q_{\kappa}(\beta) + r_{\kappa}(\beta),$

and then invoke Lemma [3.7](#page-9-3) with these two three-block decompositions to get :

$$
\varphi_L^+(\alpha,\beta) \ge \varphi_L^+(\delta,r_\kappa(\beta)) \oplus \varphi_L^+(\kappa \cdot q_\kappa(\alpha),\kappa \cdot q_\kappa(\beta)) \oplus \varphi_L^+(r_\kappa(\alpha),\gamma)
$$

which, given Corollary [3.5](#page-8-0) (and the definition of \oplus), yields :

$$
\varphi_L^+(\alpha,\beta) > \delta \underline{\oplus} \kappa \cdot (q_\kappa(\alpha) \otimes q_\kappa(\beta)) \underline{\oplus} \gamma.
$$

Then observe that the the least strict upper bound of the right-hand member of this last inequality is equal to right-hand member of (3.6) .

3.3. Majoration. For any ordinals α and β , let $\varphi(\alpha, \beta)$ denote the supremum of the length $\ell(\mathbf{a} \sqcap \mathbf{b})$ where a and **b** are two WQO on the same set and $\ell(\mathbf{a}) = \alpha$ and $\ell(\mathbf{b}) = \beta$. We shall just write $\varphi^+(\alpha, \beta)$ for $\varphi^+_P(\alpha, \beta)$. Note that $\varphi^+(\alpha,\beta) = \varphi(\alpha,\beta) + 1$ if $\varphi(\alpha,\beta)$ is realized by some pair of WQO, and that $\varphi^+(\alpha,\beta) = \varphi(\alpha,\beta)$ otherwise (Item [\(5\)](#page-9-5) of Lemma [3.8](#page-9-6) below).

Lemma 3.8. *Consider two ordinals* α *and* β *.*

(1) $\varphi^+(\alpha, \beta) = \varphi^+(\beta, \alpha)$.

- (2) $\varphi^+(\alpha, \beta) = 0$ if and only if α *and* β *fail to be equipotent.*
- (3) For every integer n, $\varphi(n,n) = n$ and $\varphi^+(n,n) = n+1$.
- (4) $|\alpha| = |\beta| \Rightarrow \varphi^+(\alpha, \beta) > \max(\alpha, \beta)$.
- (5) $\varphi(\alpha, \beta) \leq \varphi^+(\alpha, \beta) \leq \varphi(\alpha, \beta) + 1$, and $\varphi^+(\alpha, \beta) = \varphi(\alpha, \beta) + 1$ if and only if the corresponding *supremum is attained,* if and only if $\varphi^+(\alpha, \beta)$ *is a successor ordinal.*
- (6) $\varphi(\alpha, \beta) \leq \alpha \otimes \beta$.
- (7) *Given a third ordinal* α' : $|\alpha| \leq \alpha' \leq \alpha \Rightarrow \varphi(\alpha', \beta) \leq \varphi(\alpha, \beta)$ *.*
- (8) $|\alpha| = |\beta| \Longrightarrow \varphi^+(\alpha, \beta) = \varphi_{\#}(\alpha, \beta).$

Proof. Items [\(1\)](#page-9-7), [\(2\)](#page-9-8), [\(3\)](#page-9-9) and [\(5\)](#page-9-5) are obvious. Below a and b denote well orders of type α and β with the same vertex set.

- [\(4\)](#page-9-10) Each of **a** and **b** is an extension of $\mathbf{a} \sqcap \mathbf{b}$.
- [\(6\)](#page-9-11) Note that, as mentionned in the introduction, the poset a ⊓ b is embeddable into the direct product $\mathbf{a} \otimes \mathbf{b}$. Thus $\ell(\mathbf{a} \cap \mathbf{b}) \leq \ell(\mathbf{a} \otimes \mathbf{b}) = \ell(\mathbf{a}) \otimes \ell(\mathbf{b}) = \alpha \otimes \beta$ by [\(2.2\)](#page-3-0) and de Jongh-Parikh formula [\(1.1\)](#page-0-0).
- [\(7\)](#page-9-12) Invoke Lemma [2.3.](#page-4-0)
- [\(8\)](#page-9-13) Consider Item [\(7\)](#page-9-12).

Lemma 3.9. *For any ordinals* α *and* β *:*

$$
\varphi(\alpha,\beta) \leq \varphi_{\#}(\underline{\alpha},\beta) \oplus \varphi_{\#}(\alpha,\beta).
$$

Proof of Lemma [3.9.](#page-10-0) If $|\alpha| \neq |\beta|$ then the left-hand member of the inequality is 0. So assume that $|\alpha| = |\beta|$ and consider two WPO a and b on a set X of respective lenghts α and β , and let p denote the poset intersection $\mathbf{a} \sqcap \mathbf{b}$. Recall (2.1) :

$$
\ell \mathbf{p} = \sup_{x \in X} \ell \{ \mathbf{p} \not\geq_{\mathbf{p}} x \}
$$

while :

$$
\{X \not\geq_{\mathbf{p}} x\} = \{X \not\geq_{\mathbf{a}} x\} \cup \{X \not\geq_{\mathbf{b}} x\}
$$

so that :

$$
\ell\{\mathbf{p} \not\geq_{\mathbf{p}} x\} \leq \ell\{\mathbf{p} \not\geq_{\mathbf{a}} x\} \oplus \ell\{\mathbf{p} \not\geq_{\mathbf{b}} x\}
$$

and then :

$$
\ell \mathbf{p} \leq \sup_{x \in X} \ell \{ \mathbf{p} \not\geq_{\mathbf{a}} x \} \oplus \ell \{ \mathbf{p} \not\geq_{\mathbf{b}} x \}.
$$

Now observe that the poset $\{p \nleq_\mathbf{a} x\} := \mathbf{p} \restriction \{X \nleq_\mathbf{a} x\}$ is the intersection of the posets $\{\mathbf{a} \nleq_\mathbf{a} x\} := \mathbf{a} \restriction \{X \nleq_\mathbf{a} x\}$ $\{X \nleq_{\mathbf{a}} x\}$ and $\{\mathbf{b} \nleq_{\mathbf{a}} x\} := \mathbf{b} \restriction \{X \nleq_{\mathbf{a}} x\}$, the first of which has length less than α and the second of which has length at most β . Therefore :

$$
\ell \{\mathbf{p} \not\geq_\mathbf{a} x\} < \varphi_\#(\underline{\alpha},\beta)
$$

and likewise :

so that :

$$
\ell\{\mathbf{p} \not\geq_{\mathbf{a}} x\} \oplus \ell\{\mathbf{p} \not\geq_{\mathbf{b}} x\} < \varphi_{\#}(\underline{\alpha}, \beta) \oplus \varphi_{\#}(\alpha, \underline{\beta}).
$$

 $\ell{\bf p}\ngeq_{\bf b} x\lt\varphi_{\#}(\alpha,\beta)$

Henceforth :

$$
\ell \mathbf{p} \le \varphi_{\#}(\underline{\alpha}, \beta) \oplus \varphi_{\#}(\alpha, \beta)
$$

and finally :

$$
\varphi(\alpha,\beta) \leq \varphi_{\#}(\underline{\alpha},\beta) \oplus \varphi_{\#}(\alpha,\beta).
$$

 \Box

Corollary 3.6. *For every cardinal* κ *(*i.e. *initial ordinal) :*

$$
\varphi^+(\kappa,\kappa)=\kappa+1.
$$

Proof. According to Lemma [3.9](#page-10-0) :

$$
\varphi(\kappa,\kappa) \leq \varphi_{\#}(\underline{\kappa},\kappa) \oplus \varphi_{\#}(\kappa,\underline{\kappa}) = \kappa \oplus \kappa = \kappa
$$

so that $\varphi^+(\kappa,\kappa) \leq \varphi(\kappa,\kappa) + 1 \leq \kappa + 1$. The reverse inequality is straightforward.

Lemma 3.10. *Given any ordinals* α' , α'' *and* β *:*

(3.7)
$$
\varphi^+(\alpha'+\alpha'',\beta) \leq \varphi_{\#}(\alpha',\beta) \oplus \varphi_{\#}(\alpha'',\beta) \leq \varphi_{\#}(\alpha',\beta) \oplus |\alpha''|^+.
$$

 \Box

Proof. Observe that if $\alpha := \alpha' + \alpha''$ fails to be equipotent with β , then the left-hand member of [\(3.7\)](#page-10-1) is 0, in which case this inequality is trivially satisfied. So assume that α and β are equipotent.

Given two posets **a** and **b** with the same vertex set X and respective lengths α and β , let $\mathbf{p} := \mathbf{a} \square \mathbf{b}$. Consider a partition of X into subsets X' and X'' of respective lengths α' and α'' w.r.t. **a** (*cf.* Lemma [2.1\)](#page-3-4). Letting β' and β'' denote the respective lengths of X' and X'' *w.r.t.* **b**:

$$
\ell(\mathbf{p}) \leq \ell(\mathbf{p} \upharpoonright X') \oplus \ell(\mathbf{p} \upharpoonright X'') \leq \varphi(\alpha', \beta') \oplus \varphi(\alpha'', \beta'') < \varphi_{\#}(\alpha', \beta) \oplus \varphi_{\#}(\alpha'', \beta) \leq \varphi_{\#}(\alpha', \beta) \oplus |\alpha''|^{+}.
$$

For the left-hand inequality, recall [\(2.2\)](#page-3-0) ; the second one holds by definition of φ ; the penultimate one holds by definition of $\varphi_{\#}$, given that, obviously, $\beta' \leq \beta$ and $\beta'' \leq \beta$; the right-hand one follows from $\varphi_{\#}(\alpha'', \beta) \leq |\alpha''|^+$. Then [\(3.7\)](#page-10-1) holds by definition of φ **+.** □ □

Proposition 3.3. *Given two equipotent infinite ordinals* α *and* β *, of cardinality* κ *:*

(3.8)
$$
\varphi_R^+(\alpha, \beta) \leq \kappa \cdot (q_\kappa(\alpha) \otimes q_\kappa(\beta)) + |r_\kappa(\alpha) + r_\kappa(\beta)|^+.
$$

Proof of Proposition [3.3.](#page-11-0) Given κ , we prove [\(3.8\)](#page-11-1) by induction on (α, β) . First note that, according to Corollary [3.6,](#page-10-2) [\(3.8\)](#page-11-1) holds for $(\alpha, \beta) = (\kappa, \kappa)$. So let us assume that $(\alpha, \beta) > (\kappa, \kappa)$, *i.e.* that at least one of the inequalities $\alpha \geq \kappa$ and $\beta \geq \kappa$ is strict; let us assume that [\(3.8\)](#page-11-1) holds for every pair (α_1, β_1) such that $(\kappa, \kappa) \leq (\alpha_1, \beta_1) \leq (\alpha, \beta)$; and let us prove that it holds also for (α, β) . We are led to distinguish three cases. We shall first assume that α or β is not a multiple of κ ; then we shall assume that they are both multiples of κ but that at least one of them fails to be indecomposable; and it will remain to consider the case of both being indecomposable (and in particular multiples of κ). Let $\alpha = \kappa \cdot \alpha' + \gamma$ and $\beta = \kappa \cdot \beta' + \delta$, with $\gamma < \kappa$ and $\delta < \kappa$, denote the euclidian divisions of α and β by κ .

(1) Assume that α or β is not a multiple of κ , *e.g.* that $\gamma > 0$. In this case:

$$
\varphi^+(\alpha, \beta) \leq \varphi_{\#}(\kappa \cdot \alpha', \kappa \cdot \beta' + \delta) \oplus |\gamma|^{+}
$$
 by (3.7)
\n
$$
= \varphi^+(\kappa \cdot \alpha', \kappa \cdot \beta' + \delta) \oplus |\gamma|^{+}
$$
 Lemma 3.8(8)
\n
$$
\leq (\kappa \cdot (\alpha' \otimes \beta') + |\delta|^{+}) \oplus |\gamma|^{+}
$$
 Induction assumption
\n
$$
\leq \kappa \cdot (\alpha' \otimes \beta') + |\delta + \gamma|^{+}
$$
 easily checked.

(2) Let us assume now that α and β are both multiples of κ and that at least one fails to be indecomposable, *e.g.* α . Observe that α being a multiple of $|\alpha|$, its being decomposable is equivalent to $\alpha' := q_{|\alpha|}(\alpha)$ being decomposable. Thus we assume that $\alpha' = \alpha'_1 + \alpha'_2$ with α'_1 and α'_2 both less than α' , and we can even assume that, indeed, $\alpha' = \alpha'_1 + \alpha'_2 = \alpha'_1 \oplus \alpha'_2$. Then :

$$
\varphi^+(\alpha,\beta) = \varphi^+(\kappa \cdot \alpha_1' + \kappa \cdot \alpha_2', \kappa \cdot \beta')
$$

\n
$$
\leq \varphi_{\#}(\kappa \cdot \alpha_1', \kappa \cdot \beta') \oplus \varphi_{\#}(\kappa \cdot \alpha_2', \kappa \cdot \beta')
$$

\n
$$
= \varphi^+(\kappa \cdot \alpha_1', \kappa \cdot \beta') \oplus \varphi^+(\kappa \cdot \alpha_2', \kappa \cdot \beta')
$$

\n
$$
\leq (\kappa \cdot (\alpha_1' \otimes \beta') + 1) \oplus (\kappa \cdot (\alpha_2' \otimes \beta') + 1)
$$

\n
$$
= \kappa \cdot (\alpha_1' \otimes \beta') \oplus \kappa \cdot (\alpha_2' \otimes \beta') + 1
$$

\n
$$
= \kappa \cdot ((\alpha_1' \oplus \alpha_2') \otimes \beta') + 1
$$

\nby (2.3)

which is indeed [\(3.8\)](#page-11-1), given that the two remainders are 0.

(3) Let us now assume that α and β are both indecomposable ; in particular $\gamma = 0$ and $\delta = 0$. Recall from Lemma [3.9](#page-10-0) that :

$$
\varphi(\alpha,\beta) \leq \varphi_{\#}(\underline{\alpha},\beta) \oplus \varphi_{\#}(\alpha,\underline{\beta}).
$$

We claim that $\lambda := \kappa \cdot (\alpha' \otimes \beta')$ is a strict upper bound of $\{\varphi(\alpha,\beta) : \alpha_1 < \alpha, \beta_1 \leq \beta\}$. With this claim $\varphi_{\#}(\underline{\alpha}, \beta) \leq \lambda$, and likewise $\varphi_{\#}(\alpha, \beta) \leq \lambda$. Hence, given that λ is indecomposable :

$$
\varphi(\alpha,\beta) \leq \varphi_{\#}(\underline{\alpha},\beta) \oplus \varphi_{\#}(\alpha,\underline{\beta}) \leq \lambda \oplus \lambda = \lambda = \kappa \cdot (\alpha' \otimes \beta')
$$

which implies [\(3.8\)](#page-11-1).

Thus it remains to check the claim. To this end, consider $\alpha_1 < \alpha = \kappa \cdot \alpha'$ and $\beta_1 < \beta = \kappa \cdot \beta'$.

• If $\alpha' = 1$, then $\alpha_1 < \kappa$, so that :

$$
\varphi(\alpha_1,\beta_1) < |\alpha_1|^+ \leq \kappa \leq \kappa \cdot \beta' = \kappa \cdot (\alpha' \otimes \beta').
$$

• If $\alpha' > 1$, then this ordinal, which is assumed to be indecomposable, is an infinite limit ordinal. In this case $\alpha_1 \leq \kappa \cdot \alpha'' < \alpha$ for some $\alpha'' < \alpha'$. Then, invoking the induction assumption for the middle inequality :

$$
\varphi(\alpha_1,\beta_1) \leq \varphi(\kappa \cdot \alpha'',\kappa \cdot \beta') \leq \kappa \cdot (\alpha'' \otimes \beta') + 1 < \kappa \cdot (\alpha' \otimes \beta').
$$

 \Box

REFERENCES

- [1] A. Blass, Y. Gurevich, Program termination and well partial orderings. ACM Trans. Comput. Log. 9 (2008), no. 3, Art. 18, 26 pp.
- [2] P.W. Carruth, Arithmetic of ordinals with applications to the theory of ordered Abelian groups. Bull. Amer. Math. Soc. 48, (1942). 262-271.
- [3] I. Chakir, Conditions de chaînes dans les treillis algébriques, Thèse de doctorat d'Etat, Université de Rabat, 11 Mai 2009.
- [4] I. Chakir, M. Pouzet, The length of chains in algebraic lattices, Proceedings ISOR'08, International Symposium on Operational Research,Algiers , Algeria , Nov 2-6, 2008, H.Ait Haddadene, I.Bouchemakh, M.Boudar, S.Bouroubi (Eds)LAID3, pp.379-390. [arXiv:0812.2193v](http://arxiv.org/abs/0812.2193)1 [math.CO]
- [5] I. Chakir, and M. Pouzet, A characterization of well-founded algebraic lattices, preprint, [arXiv:0812.2300](http://arxiv.org/abs/0812.2300) [math.CO].
- [6] C. Delhommé, Height of a superposition. Order 23 (2006), no. 2-3, 221–233.
- [7] C. Delhommé, Decomposition of tree-automatic structures. Submitted.
- [8] D. H. J. de Jongh and R. Parikh. Well-partial orderings and hierarchies. Nederl. Akad. Wetensch. Proc. Ser. A 80=Indag. Math., 39(3):195–207, 1977.
- [9] R. Fraïssé. Theory of relations. North-Holland Publishing Co., Amsterdam, 2000.
- [10] T. Forster, An Introduction to WQO and BQO Theory, 145p, 2000.
- [11] J. H. Gallier. What's so special about Kruskal's theorem and the ordinal Γ_0 ? A survey of some results in proof theory. Ann. Pure Appl. Logic, 53(3):199–260, 1991.
- [12] G. Higman. Ordering by divisibility in abstract algebras. Proc. London Math. Soc. (3), 2:326-336, 1952.
- [13] T. Jech, Set theory. The third millennium edition, revised and expanded. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. xiv+769 pp.
- [14] I. Křiž, R. Thomas, Ordinal types in Ramsey theory and well-partial-ordering theory. Mathematics of Ramsey theory, 5795, Algorithms Combin., 5, Springer, Berlin, 1990,
- [15] R. Lalement. Computation as logic. Prentice Hall International, Englewood Cliffs, NJ, 1993. With a foreword by Michel Demazure, Translated from the 1990 French original by John Plaice.
- [16] E. C. Milner. Basic wqo- and bqo-theory. In Graphs and order (Banff, Alta., 1984), pages 487–502. Reidel, Dordrecht, 1985.
- [17] M. Pouzet, N. Zaguia, Ordered sets with no chains of ideals of a given type. Order 1 (1984), 159-172.
- [18] M. Pouzet, N. Zaguia, Dimension de Krull des ensembles ordonnés. Discrete Math. 53 (1985), 173-192.
- [19] M. Pouzet, M. Sobrani, The order type of the collection of finite series-parallel posets. Discrete Math. 265 (2003), n 1-3, 189211.
- [20] M. Rathjen and A. Weiermann. Proof-theoretic investigations on Kruskal's theorem. Ann. Pure Appl. Logic, 60(1):49–88, 1993.
- [21] N. Robertson and P. D. Seymour. Graph minors—a survey. In Surveys in combinatorics 1985 (Glasgow, 1985), pages 153–171. Cambridge Univ. Press, Cambridge, 1985.
- [22] D. Schmidt. Well-partial orderings and their maximal order types. Habilitation, University of Heidelberg, Heidelberg, 1978. [23] E.S. Wolk, Partially well ordered sets and partial ordinals. Fund. Math. 60 (1967) 175-186.
-

LIM-ERMIT, UNIVERSITÉ DE LA RÉUNION, FACULTÉ DES SCIENCES ET TECHNOLOGIES - PTU, 2, RUE JOSEPH WETZEL, 97490 Sainte-Clotilde, France

E-mail address: delhomme@univ-reunion.fr

ICJ, MATHÉMATIQUES, UNIVERSITÉ CLAUDE-BERNARD LYON1, 43 BD. 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX, France and Mathematics & Statistics Department, University of Calgary, Calgary, Alberta, Canada T2N 1N4 E-mail address: pouzet@univ-lyon1.fr, mpouzet@ucalgary.ca