

Discrete photon statistics from continuous microwave measurements

Stéphane Virally,^{1,*} Jean Olivier Simoneau,¹ Christian Lupien,¹ and Bertrand Reulet¹

¹*Département de Physique, Université de Sherbrooke, Sherbrooke, Québec J1K 2R1, Canada*

(Dated: May 25, 2022)

Photocount statistics are an important tool for the characterization of electromagnetic fields, especially for fields with an irrelevant phase. In the microwave domain, continuous rather than discrete measurements are the norm. Using a novel approach, we recover discrete photon statistics from the cumulants of a continuous distribution of field quadrature measurements. The use of cumulants allows the separation between the signal of interest and experimental noise. Using a parametric amplifier as the first stage of the amplification chain, we extract useful data from up to the sixth cumulant of the continuous distribution of a coherent field, hence recovering up to the third moment of the discrete statistics associated with a signal with much less than one average photon.

Introduction. Photon statistics measurements provide a wealth of information on the state of the electromagnetic field. For instance, Glauber's theory of optical coherence [1] is solely based on correlations between multiple photon measurements. As the discrete nature of the interaction between light and matter is essentially a quantum feature, statistical distributions can also characterize the classicality of photonic states. For instance, single photon states [2, 3] exhibit sub-Poisson photocount distributions that are not predicted by classical theories [4]. These states are not just of theoretical interest, as they feature prominently in proposals for the development of quantum computation [5] and quantum communication networks [6].

With the advent of circuit QED [7], there is currently a great interest in quantum states of the electromagnetic field in the microwave domain. From early on, predictions have been made on the specific type of photon statistics that can be expected in mesoscopic conductors [8–10]. Recently, purely quantum photonic states have been demonstrated in the microwave domain, using superconducting devices [11, 12]. Entanglement has also been demonstrated in the GHz range using normal conductors [13, 14]. The detection of single photons in the microwave domain remains a challenge, but schemes have been proposed for the extraction of photocounts from continuous measurements with linear detectors [15]. Experiments have already been performed to specifically extract discrete statistics from continuous measurements [16–18], although they focused on the calculation of centered moments of the continuous distributions, rather than the cumulants, as is the case herein.

In this Letter, we derive simple formulas linking the cumulants [19] of the continuous variable (CV) distribution of field quadrature measurements to the centered moments of the photocount statistics. The latter fully characterize the state of the electromagnetic field when its phase is either not well defined or irrelevant. The additive nature of cumulants is especially important in the quantum regime, when we reconstruct photon statistics for signals with much less than one photon per mea-

surement on average and noise contributions cannot be discarded. We also use a quantum-limited parametric amplifier as the initial stage of the amplification chain, in order to lower the noise background to a few photons per measurement. With both techniques, we can characterize quantum sources with much less than one average photon in just a few seconds, without the need for excessive averaging.

This Letter is organized as follows. We first derive formulas linking the cumulants of the CV distribution to the discrete moments of the photocount distribution. We then discuss semi-classical limits and show how quantum states can be characterized from their statistics. Finally, in the experimental section, we recover the first three moments of the photocount distribution associated with a weak coherent state (featuring much less than one photon on average per measurement) by measuring the first six cumulants of the CV distribution.

Theory. We measure quadratures of a bosonic input field by mixing the signal with a local oscillator (heterodyne interferometric measurement). If the input field is associated with the ladder operator \hat{a} , observables are of the form

$$\hat{X}_{\hat{a},\theta} = \frac{e^{i\theta} \hat{a}^\dagger + e^{-i\theta} \hat{a}}{\sqrt{2}}, \quad (1)$$

with θ the phase between the input field and the local oscillator.

We focus on the case of a measurement where the phase difference θ is effectively averaged during the full time τ of statistics accumulation. This is the case when the phase varies randomly during the detection, but also in the narrow-band limit, when the pulsations ω_0 of the local oscillator and ω of the measured signal differ by $\delta\omega$, with $\delta\omega \tau \gg 1$. As the phase θ is averaged, expectations of all odd centered moments of the measured signal are zero, while expectations of the $2k^{\text{th}}$ centered moments are

$$\langle X_{\hat{a},\theta}^{2k} \rangle_\theta = \left(\frac{1}{2}\right)^k \left\langle \sum_{\text{c.s.}} \hat{a}^k \hat{a}^{\dagger k} \right\rangle, \quad (2)$$

with ‘‘c.s.’’ standing for completely symmetric (i.e., the sum is taken on all possible permutations of the non-commuting operators). Here, $\langle \bullet \rangle$ represents averaging over the quantum ensemble, while $\langle \bullet \rangle_\theta$ represents averaging over both the quantum ensemble and the phase θ . To simplify notations, we drop the θ indices in the calculations below.

The completely symmetric sum of Eq. 2 can be explicitly evaluated and yields [20]

$$\begin{aligned} \sum_{\text{c.s.}} \hat{a}^k \hat{a}^{\dagger k} &= \sum_{i=0}^k \left(\frac{1}{2}\right)^{k-i} \frac{(2k)!}{(i!)^2(k-i)!} \hat{a}^{\dagger i} \hat{a}^i \\ &= \sum_{i=0}^k \left(\frac{1}{2}\right)^{k-i} \frac{(2k)!}{(i!)^2(k-i)!} \prod_{j=0}^{i-1} (\hat{n} - j), \end{aligned} \quad (3)$$

with $\hat{n} = \hat{a}^\dagger \hat{a}$ the usual number operator. We have explicitly linked the even moments $\langle X_a^{2k} \rangle$ of the measured CV distribution to the moments $\langle n^\ell \rangle$ of the discrete photocount distribution.

As explicited before, we actually want to work with the cumulants $C_k = \langle X_a^k \rangle$ of the CV distribution rather than its moments. Cumulants characterize the distribution in the exact same way as its moments, but they also have the advantage of being additive for independant distributions. This is crucial when unavoidable noise contributions, including vacuum fluctuations and the weak thermal contributions of the amplification process, are taken into account. With the use of cumulants, these independant contributions can be measured separately and removed from the final measurement, leaving only the signal, however weak.

Simple algebra leads from Eq. 3 to the first three moments of the photocount distribution in the form

$$\langle n \rangle = C_2 - \frac{1}{2}; \quad (4)$$

$$\langle \delta n^2 \rangle = \frac{2}{3} C_4 + C_2^2 - \frac{1}{4}; \quad (5)$$

$$\langle \delta n^3 \rangle = \frac{2}{5} C_6 + 4 C_4 C_2 + 2 C_2^3 - \frac{1}{2} C_2, \quad (6)$$

where $\langle \delta n^\ell \rangle = \langle (n - \langle n \rangle)^\ell \rangle$ is the ℓ^{th} centered moment of the photocount distribution.

Cumulants can also be obtained through the use of the second cumulant-generating function [19], which is the natural logarithm of the characteristic function, or

$$\Gamma(\lambda) = \ln[\langle \exp(i\lambda X_{\hat{a},\theta}) \rangle_\theta], \quad (7)$$

as

$$C_k = (-i)^k \frac{d^k \Gamma}{d\lambda^k}(0). \quad (8)$$

The second cumulant-generating function can be computed for some usual states [21]. For a thermal state,

$\Gamma(\lambda) = -(\langle n \rangle + 1/2) \lambda^2/2$. This is the second cumulant-generating function of a gaussian distribution, leading to $C_2 = \langle n \rangle + 1/2$ and $C_k = 0, \forall k \geq 3$. In other words, there is an equivalence between gaussian distribution of quadratures and thermal (chaotic) statistics. When the fourth-order cumulant is not zero, the statistics is not chaotic. This is in contradiction with a claim made in ref. [18] for the noise emitted by a tunnel junction. Such noise is known to be caused by Poissonian charge tunneling statistics [22] and thus features a non-vanishing fourth-order current cumulant ($C_4 \propto e^3 B^3 \bar{I}$ where e is the unit charge, B the bandwidth and \bar{I} the average current). For this reason, the associated photocount statistics is not chaotic. However, the ratio $C_4/C_2^2 \propto e B/\bar{I}$ of the current statistics is typically small (10^{-4} to 10^{-3} in [18]). It can then be difficult to discern between chaotic and non-chaotic statistics, especially in the presence of detection noise.

For a coherent state, $\Gamma(\lambda) = -\lambda^2/4 + \ln[I_0(i\sqrt{2}|\alpha|\lambda)]$, where I_0 is the modified Bessel function of the first kind. This leads to $C_2 = |\alpha|^2 + 1/2$, $C_4 = -3|\alpha|^4/2$ and $C_6 = 10|\alpha|^6$. Using Eqs. 4, 5, 6 we get, as expected, $\langle n \rangle = \langle \delta n^2 \rangle = \langle \delta n^3 \rangle = |\alpha|^2$.

Semi-classical narrow-band model. We wish to extract useful limits for semi-classical narrow-band signals. We first model the amplitude of a fully classical narrow-band signal as

$$s(t) = a(t) \cos(\omega t) + b(t) \sin(\omega t), \quad (9)$$

where a and b are real functions of time with slow variations (in the sense that the time scale of their variations is much smaller than $1/\omega$). We consider a measurement process that cannot resolve the fast oscillations at ω . The intensity of a such a signal is then defined by

$$i(t) = \frac{1}{2} [a^2(t) + b^2(t)]. \quad (10)$$

A semi-classical prolongation of the model requires adding an ersatz for the vacuum, in the form

$$s_{\text{vac}}(t) = \xi_a(t) \cos(\omega t) + \xi_b(t) \sin(\omega t), \quad (11)$$

where ξ_a and ξ_b are two independant gaussian random processes with null average and $1/2$ variance.

By identifying $X_{\hat{a},\omega t}$ with $s(t) + s_{\text{vac}}(t)$, and applying Eqs. 4, 5, 6 we get (after averaging ωt over many cycles)

$$\langle n \rangle = \langle i \rangle; \quad (12)$$

$$\langle \delta n^2 \rangle = \langle i \rangle + \langle \delta i^2 \rangle, \quad (13)$$

$$\langle \delta n^3 \rangle = \langle i \rangle + 3 \langle \delta i^2 \rangle + \langle \delta i^3 \rangle, \quad (14)$$

where $\langle \delta i^k \rangle$ is the k^{th} centered moment of the intensity distribution.

Eqs. ?? provide interesting limits for classical signals. There exists inequalities verified by continuous distributions. For instance $\langle \delta i^2 \rangle \geq 0$ implies

$$\langle \delta n^2 \rangle \geq \langle n \rangle. \quad (15)$$

Thus, we recover the fact that classical signals cannot obey sub-Poissonian statistics [23]. When the intensity is constant, all moments $\langle \delta i^k \rangle$ vanish and we recognize the Poisson statistics of coherent signals. Super-Poissonian statistics exists as soon as $i(t)$ is not constant. This is the case, for instance, when the intensity is modulated, as shown in the experimental section.

Additional inequalities stem from Stieltjes moments theorem [24]. For instance, $\langle \delta i^3 \rangle \geq -3 \langle i \rangle \langle \delta i^2 \rangle$ leads to

$$\langle \delta n^3 \rangle \geq \langle n \rangle + 3 (\langle \delta n^2 \rangle - \langle n \rangle) (1 - \langle n \rangle). \quad (16)$$

In particular, if $\langle n \rangle \leq 1$, $\langle \delta n^3 \rangle \geq \langle n \rangle$.

In addition to the “hard” inequalities of Eqs. 15, 16, we can derive relations in the vanishing signal limit. For a classical signal, a reduction in intensity is equivalent to the addition of an attenuator in the line. It is possible to write $\langle i \rangle = \eta \langle I \rangle$ and $\langle \delta i^k \rangle = \eta^k \langle \delta I^k \rangle$ with a fixed intensity I and a varying attenuation factor η . Accordingly, the vanishing signal limit $\langle i \rangle \rightarrow 0$ is equivalent to the limit $\eta \rightarrow 0$. Thus, Eqs. 12, 13, 14 yield $\langle \delta n^2 \rangle, \langle \delta n^3 \rangle \sim \langle n \rangle$ as $\langle n \rangle \rightarrow 0$. Hence, the Fano factor [25] $\mathcal{F} = \langle \delta n^2 \rangle / \langle n \rangle$ of a narrow-band classical signal verifies

$$\mathcal{F} \rightarrow 1 \text{ as } \langle n \rangle \rightarrow 0. \quad (17)$$

In fact, the Fano factor is a great tool for characterizing distributions in the low $\langle n \rangle$ limit. It should always be close to 1 for classical states and only deviate reasonably from this value for non-classical states such as Fock states or squeezed vacua. Although Fano factors of a few hundreds have been reported in previous experiments [16], they only reflect the fact that moments were used, rather than cumulants, thus mixing signal and noise. They do not reflect the underlying statistics of coherent states.

Fig. 1 identifies classical and non-classical regions in the $(\langle n \rangle, \langle \delta n^2 \rangle)$ plane. For a given photon average, coherent states feature the lowest variance of all classical states. As shown above, there also exists an upper limit for narrow-band classical signals as the photon average vanishes, with a Fano factor approaching unity. There are thus two distinct non-classical zones in the plane. Quantum states below the $\langle \delta n^2 \rangle = \langle n \rangle$ line (i.e. states with sub-Poissonian statistics) are well-known and include Fock states (on the x-axis). On the other side, squeezed vacuum states exhibit super-Poissonian statistics even as $\langle n \rangle \rightarrow 0$ (with a limit $\langle \delta n^2 \rangle \sim 2 \langle n \rangle$ that reflects the production of photon pairs in these states). Although the two non-classical regions seem segregated, it is possible in some cases to move a state from one to the other

using a symplectic operation, such as a displacement. For instance, when a squeezed vacuum is displaced in the quadrature plane in the direction of squeezing, it can cross the classical region and end up as an amplitude squeezed state in the sub-Poisson region [26, 27]. Indeed, under application of the Glauber displacement operator [23] $\hat{D}(r e^{i\theta})$, the average photon number and variance of the photocount distribution become

$$\langle n \rangle \rightarrow \langle n \rangle + \sqrt{2} r \langle X_{\hat{a},\theta} \rangle + r^2 \quad (18)$$

and

$$\langle \delta n^2 \rangle \rightarrow \langle \delta n^2 \rangle + 2 r^2 \langle \delta \hat{X}_{\hat{a},\theta}^2 \rangle + 2 \sqrt{2} r \text{cov}(\hat{n}, \hat{X}_{\hat{a},\theta}), \quad (19)$$

where the covariance between two operators is defined as

$$\text{cov}(\hat{p}, \hat{q}) = \frac{1}{2} (\langle \hat{p} \hat{q} \rangle + \langle \hat{q} \hat{p} \rangle) - \langle \hat{p} \rangle \langle \hat{q} \rangle. \quad (20)$$

The migration from one non-classical region to the other is due to the presence of the covariance term that reflects correlations between \hat{n} and $\hat{X}_{\hat{a},\theta}$.




FIG. 1 (color online). States of a narrow-band semi-classical electromagnetic field are restricted in the $(\langle n \rangle, \langle \delta n^2 \rangle)$ plane. They can only exist in the shaded region. The lower limit (thick blue line) represents coherent states and Poissonian statistics. The shaded region is not bounded by a “hard” upper curve as classical states are only constrained in the limit $\langle n \rangle \rightarrow 0$. Squeezed vacua (solid red line) can exist in the non-classical upper part. They can cross from the upper to the lower non-classical parts when appropriately displaced (green arrow).

Experiment. We tested the results of Eqs. 5, 6 on a weak coherent state. The full experimental setup is represented in Fig. 2. The ideal signal is well approximated by a phase and amplitude-controlled sine wave at 6.01 GHz generated by an analog signal generator. It is attenuated down to much less than a photon per measurement bin before going through a Josephson parametric amplifier (paramp) [28] placed in a dilution refrigerator and thermalized at 7 mK. The parametric amplifier is pumped in




FIG. 4 (color online). Variance of photon statistics for different modulations schemes. Although all distributions are super-poissonian, in the small signal limit all variances move towards $\langle n \rangle$, as expected for classical states.

factor of 2 in the limit of vanishing $\langle n \rangle$.

We acknowledge fruitful discussions with J. Gabelli. We thank G. Laliberté for technical help. This work was supported by the Canada Excellence Research Chairs program, Canada NSERC, Québec MEIE, Québec FRQNT via INTRIQ, Université de Sherbrooke via EPIQ, and Canada Foundation for Innovation.

* Stephane.Virally@USherbrooke.ca

- [1] R. J. Glauber, *Physical Review* **130**, 2529 (1963); *Physical Review Letters* **10**, 84 (1963).
- [2] C. K. Hong and L. Mandel, *Physical Review Letters* **56**, 58 (1986).
- [3] M. Eisaman, J. Fan, A. Migdall, and S. Polyakov, *Review of Scientific Instruments* **82**, 071101 (2011).
- [4] L. Mandel and E. Wolf, *Optical Coherence and Quantum Optics* (Cambridge University Press, 1995).
- [5] E. Knill, R. Laflamme, and G. J. Milburn, *Nature* **409**, 46 (2001).
- [6] N. Sangouard, C. Simon, H. De Riedmatten, and N. Gisin, *Reviews of Modern Physics* **83**, 33 (2011).
- [7] S. M. Girvin, M. H. Devoret, and R. J. Schoelkopf, *Physica Scripta* **T137**, 014012 (2009).
- [8] C. W. J. Beenakker and H. Schomerus, *Physical Review Letters* **86**, 700 (2001); **93**, 096801 (2004).
- [9] A. V. Lebedev, G. B. Lesovik, and G. Blatter, *Physical Review B* **81**, 155421 (2010).
- [10] C. Padurariu, F. Hassler, and Y. V. Nazarov, *Physical Review B* **86**, 054514 (2012).
- [11] D. Bozyigit, C. Lang, L. Steffen, J. M. Fink, M. Baur, R. Bianchetti, P. J. Leek, S. Filipp, M. P. da Silva, A. Blais, and A. Wallraff, *Nature Physics* **7**, 154 (2011).
- [12] C. Lang, C. Eichler, L. Steffen, J. M. Fink, M. J. Woolley, A. Blais, and A. Wallraff, *Nature Physics* **9**, 345 (2013).
- [13] E. P. Menzel, R. Di Candia, F. Deppe, P. Eder, L. Zhong, M. Ihmig, M. Haeberlein, A. Baust, E. Hoffmann, D. Ballester, K. Inomata, T. Yamamoto, Y. Nakamura, E. Solano, A. Marx, and R. Gross, *Physical Review Letters* **109**, 250502 (2012).
- [14] J.-C. Fergues, C. Lupien, and B. Reulet, *Physical Review Letters* **113**, 043602 (2014); **114**, 130403 (2015).
- [15] M. P. da Silva, D. Bozyigit, A. Wallraff, and A. Blais, *Physical Review A* **82**, 043804 (2010).
- [16] J. Gabelli, L.-H. Reydellet, G. Fève, J.-M. Berroir, B. Plaçais, P. Roche, and D. C. Glattli, *Physical Review Letters* **93**, 056801 (2004).
- [17] E. P. Menzel, F. Deppe, M. Mariani, M. A. Araque Caballero, A. Baust, T. Niemczyk, E. Hoffmann, A. Marx, E. Solano, and R. Gross, *Physical Review Letters* **105**, 100401 (2010).
- [18] E. Zakka-Bajjani, J. Dufouleur, N. Coulombel, P. Roche, D. C. Glattli, and F. Portier, *Physical Review Letters* **104**, 206802 (2010).
- [19] J. F. Kenney and E. S. Keeping, *Mathematics of Statistics, Part Two*, 2nd ed. (Van Nostrand, New York, 1951).
- [20] K. Cahill and R. J. Glauber, *Physical Review* **177**, 1857 (1969).
- [21] C. W. Gardiner and P. Zoller, *Quantum Noise*, 3rd ed. (Springer Berlin / Heidelberg, 2004).
- [22] Y. Blanter and M. Büttiker, *Physics Reports* **336**, 1 (2000).
- [23] R. Loudon, *The Quantum Theory of Light*, 3rd ed. (Oxford University Press, 2000).
- [24] D. V. Widder, *The Laplace Transform* (Princeton University Press, 1946).
- [25] Alternatively, it is common in quantum optical experiments to measure related quantities such as Mandel's Q parameter ($Q = \mathcal{F} - 1$) or the degree of second-order coherence ($g^{(2)}(0) = 1 + Q/\langle n \rangle$).
- [26] N. B. Grosse, T. Symul, M. Stobiska, T. C. Ralph, and P. K. Lam, *Physical Review Letters* **98**, 2 (2007).
- [27] M.-A. Lemonde, N. Didier, and A. A. Clerk, *Physical Review A* **90**, 063824 (2014).
- [28] M. Hatridge, R. Vijay, D. H. Slichter, J. Clarke, and I. Siddiqi, *Physical Review B* **83**, 134501 (2011).