Global dynamics above the ground state energy for the combined power-type nonlinear Schrödinger equations with energy-critical growth at low frequencies

Takafumi Akahori, Slim Ibrahim, Hiroaki Kikuchi and Hayato Nawa

Abstract

We consider the combined power-type nonlinear Schrödinger equations with energycritical growth, and study the solutions slightly above the ground state threshold at low frequencies, so that we obtain a so-called nine-set theory developed by Nakanishi and Schlag [24, 25].

1 Introduction

We consider the following nonlinear Schrödinger equation:

$$i\frac{\partial\psi}{\partial t} + \Delta\psi + |\psi|^{p-1}\psi + |\psi|^{\frac{4}{d-2}}\psi = 0, \qquad (\text{NLS})$$

where $\psi = \psi(x,t)$ is a complex-valued function on $\mathbb{R}^d \times \mathbb{R}$ $(d \ge 3)$, Δ is the Laplace operator on \mathbb{R}^d and p satisfies that

$$2 + \frac{4}{d}
(1.1)$$

We denote the mass and the Hamiltonian of (NLS) by \mathcal{M} and \mathcal{H} , respectively, that is,

$$\mathcal{M}(u) := \frac{1}{2} \|u\|_{L^2}^2, \qquad (1.2)$$

$$\mathcal{H}(u) := \frac{1}{2} \|\nabla u\|_{L^2}^2 - \frac{1}{p+1} \|u\|_{L^{p+1}}^{p+1} - \frac{1}{2^*} \|u\|_{L^{2^*}}^{2^*}.$$
(1.3)

The Cauchy problem for (NLS) is locally well-posed in H^1 (see, e.g., Proposition 3.1 in [29]), and the mass and the Hamiltonian are conserved quantities for the flow defined by (NLS). Furthermore, for any solution ψ of finite variance, we have the "virial identity":

$$\frac{d^2}{dt^2} \int_{\mathbb{R}^d} |x|^2 |\psi(x,t)|^2 \, dx = 8\mathcal{K}(\psi(t)), \tag{1.4}$$

where

$$\mathcal{K}(u) := \|\nabla u\|_{L^2}^2 - \frac{d(p-1)}{2(p+1)} \|u\|_{L^{p+1}}^{p+1} - \|u\|_{L^{2^*}}^{2^*}.$$
(1.5)

It is easy to see that

$$\mathcal{H}(u) > \frac{1}{2}\mathcal{K}(u). \tag{1.6}$$

A standing wave of (NLS) with frequency ω is a solution to (NLS) of the form $e^{it\omega}u$, so that u solves the following semilinear elliptic equation:

$$\omega u - \Delta u - |u|^{p-1}u - |u|^{\frac{4}{d-2}}u = 0, \qquad u \in H^1(\mathbb{R}^d) \setminus \{0\}.$$
(1.7)

Moreover, a ground state of (1.7) is a solution to (1.7) of the minimal action, where by the action, we mean the functional S_{ω} defined by

$$\mathcal{S}_{\omega} := \omega \mathcal{M} + \mathcal{H}. \tag{1.8}$$

In [2], the same authors showed the existence of a ground state via the variational problem

$$m_{\omega} := \inf \left\{ \mathcal{S}_{\omega}(u) \colon u \in H^1(\mathbb{R}^d) \setminus \{0\}, \ \mathcal{K}(u) = 0 \right\}.$$
(1.9)

More precisely, we proved that for any $d \ge 4$, any p with $2 + \frac{4}{d} and any <math>\omega > 0$, the variational problem associated with m_{ω} has a minimizer, and any minimizer for m_{ω} becomes a ground state of (1.7). Moreover, we see from Proposition 1.2 in [2] that for any $d \ge 3$, any p with $2 + \frac{4}{d} and any <math>\omega > 0$,

$$m_{\omega} = \inf \left\{ \mathcal{I}_{\omega}(u) \colon u \in H^{1}(\mathbb{R}^{d}) \setminus \{0\}, \ \mathcal{K}(u) \leq 0 \right\}$$

$$= \inf \left\{ \mathcal{J}_{\omega}(u) \colon u \in H^{1}(\mathbb{R}^{d}) \setminus \{0\}, \ \mathcal{K}(u) \leq 0 \right\},$$
(1.10)

where

$$\mathcal{I}_{\omega}(u) := \mathcal{S}_{\omega}(u) - \frac{2}{d(p-1)}\mathcal{K}(u), \qquad (1.11)$$

$$\mathcal{J}_{\omega}(u) := \mathcal{S}_{\omega}(u) - \frac{1}{2}\mathcal{K}(u).$$
(1.12)

It is worthwhile noting that \mathcal{I}_{ω} and \mathcal{J}_{ω} do not contain the L^{p+1} -norm and \dot{H}^1 -norm, respectively: Precisely,

$$\mathcal{I}_{\omega}(u) = \frac{\omega}{2} \|u\|_{L^{2}}^{2} + \frac{p-1-\frac{4}{d}}{2(p-1)} \|\nabla u\|_{L^{2}}^{2} + \frac{2^{*}-(p+1)}{2^{*}(p-1)} \|u\|_{L^{2^{*}}}^{2^{*}},$$
(1.13)

$$\mathcal{J}_{\omega}(u) = \frac{\omega}{2} \|u\|_{L^{2}}^{2} + \frac{d(p-1-\frac{4}{d})}{4(p+1)} \|u\|_{L^{p+1}}^{p+1} + \frac{1}{d} \|u\|_{L^{2^{*}}}^{2^{*}}.$$
(1.14)

Remark 1.1. The authors have to say that Theorem 1.2 in [2] contains a mistake. Actually, if d = 3 and $3 , then for any <math>\omega > 0$, we can prove the existence of ground state of (1.7) (cf. Theorem 1 in [32]). Moreover, we find from Proposition A.1 in Section A that if d = 3 and $1 , then there exists <math>\omega_3 > 0$ such that for any $0 < \omega < \omega_3$, the equation (1.7) admits at least one ground state. Our aim in this paper is to study the behavior of solutions to (NLS) slightly above the "ground state threshold at low frequencies", in the spirit of Nakanishi and Schlag [24, 25]. To this end, we need detailed information about ground states of (1.7). We will see that for any $d \geq 3$ and $2 + \frac{4}{d} , there exists <math>\omega_1 > 0$ such that for any $0 < \omega < \omega_1$, a positive radial ground state Φ_{ω} of (1.7) exists and unique (see Proposition 2.1 and (i) of Proposition 2.4). Furthermore, the ground state Φ_{ω} satisfies that $\frac{d}{d\omega}\mathcal{M}(\Phi_{\omega}) < 0$ on $(0, \omega_1)$ (see (iii) of Proposition 2.4), so that the inverse of the mapping $\alpha \in (0, \omega_1) \rightarrow \mathcal{M}(\Phi_{\alpha}) \in (\mathcal{M}(\Phi_{\omega_1}), \infty)$ exists (we will see that $\lim_{\omega \to 0} \mathcal{M}(\Phi_{\omega}) = \infty$ in (2.9)).

Throughout this paper, we use the symbol Φ_{ω} to denote the positive radial ground state of (1.7). Moreover, $\mathcal{O}(\Phi_{\omega})$ denotes the orbit

$$\mathcal{O}(\Phi_{\omega}) := \left\{ e^{i\theta} \Phi_{\omega} \colon \theta \in \mathbb{R} \right\}.$$
(1.15)

Now, we state our main result.

Theorem 1.1. Assume either d = 3 and $3 \le p < 5$, or $d \ge 4$ and $1 + \frac{4}{d-1} .$ $Then, there exists <math>\omega_* > 0$ such that for any $\omega \in (0, \omega_*)$, there exists a positive function $\varepsilon_{\omega}: [0, \infty) \to (0, \infty)$ with the following property: Set

$$\widetilde{PW}_{\omega} := \left\{ u \in H^1(\mathbb{R}^d) \colon \mathcal{S}_{\omega}(u) < m_{\omega} + \varepsilon_{\omega}(\mathcal{M}(u)) \right\}.$$
(1.16)

Then, any radial solution ψ starting from \widehat{PW}_{ω} exhibits one of the following scenarios: In the statement below, $\alpha > 0$ denotes the constant such that $\mathcal{M}(\psi) = \mathcal{M}(\Phi_{\alpha})$.

(i) Scattering both forward and backward in time;

- (ii) Finite time blowup both forward and backward in time;
- (iii) Scattering forward in time, and finite time blowup backward in time;
- (iv) Finite time blowup forward in time, and scattering backward in time;
- (v) Trapped by $\mathcal{O}(\Phi_{\alpha})$ forward in time, and scattering backward in time;
- (vi) Scattering forward in time, and trapped by $\mathcal{O}(\Phi_{\alpha})$ backward in time;
- (vii) Trapped by $\mathcal{O}(\Phi_{\alpha})$ forward in time, and finite time blowup backward in time;
- (viii) Finite time blowup forward in time, and trapped by $\mathcal{O}(\Phi_{\alpha})$ backward in time;

(ix) Trapped by $\mathcal{O}(\Phi_{\alpha})$ both forward and backward in time.

Here, "scattering forward in time" means that the maximal lifespan of a solution ψ is infinity and there exists $\phi \in H^1(\mathbb{R}^d)$ such that

$$\lim_{t \to \infty} \left\| \psi(t) - e^{it\Delta} \phi \right\|_{H^1} = 0; \tag{1.17}$$

"blowup forward in time" means that the maximal lifespan of a solution is finite; and "trapped by $\mathcal{O}(\Phi_{\alpha})$ forward in time" means that the maximal lifespan of a solution is infinity and the solution stays in some neighborhood of $\mathcal{O}(\Phi_{\alpha})$ in $H^1(\mathbb{R}^d)$ after some time. The terms corresponding to "backward in time" are used in a similar manner.

Remark 1.2. We continue our study in anticipation of validity of Theorem 1.1 for $d \ge 3$ and $1 + \frac{4}{d} \le p < 2^* - 1$. We describe main obstructions to the extension: (1) The double critical case (namely, $p = 1 + \frac{4}{d}$) has a special difficulty that the same proof as (iii) of Proposition 2.4 cannot apply to derive the property (2.11). In fact, the property (2.11) fails for the equation (1.22) with $p = 1 + \frac{4}{d}$. Moreover, in the double critical case $p = 1 + \frac{4}{d}$, the scaling-exponent s_p (see (1.42)) vanishes $(s_{1+\frac{4}{d}} = 0)$, which requires some modifications in our argument, especially the proof of existence of unstable mode (Proposition 4.1).

(2) The restriction on the exponent p in Theorem 1.1 comes from the "one-pass theorem" (see Theorem 7.1 and Remark 7.1).

We will undertake the extension of Theorem 1.1 including the double critical case in a forthcoming paper as well as the high-frequency case $\omega \gg 1$.

We give a proof of Theorem 1.1 in Section 3 below. In particular, we show how to construct the function ε_{ω} (see (3.8)).

The equation (NLS) reminds us of the following ones

$$i\frac{\partial\psi}{\partial t} + \Delta\psi + |\psi|^{p-1}\psi = 0, \qquad (1.18)$$

$$i\frac{\partial\psi}{\partial t} + \Delta\psi + |\psi|^{\frac{4}{d-2}}\psi = 0.$$
(1.19)

These equations are invariant under the following scalings, respectively:

$$\psi(x,t) \mapsto \lambda^{-\frac{2}{p-1}} \psi(\frac{x}{\lambda}, \frac{t}{\lambda^2}),$$
(1.20)

$$\psi(x,t) \mapsto \lambda^{-\frac{d-2}{2}} \psi(\frac{x}{\lambda}, \frac{t}{\lambda^2}).$$
 (1.21)

On the other hand, there is no scaling which leaves (NLS) invariant.

The stationary problems corresponding to (1.18) and (1.19) are respectively

$$\omega v - \Delta v - |v|^{p-1} v = 0, (1.22)$$

$$\Delta v + |v|^{\frac{4}{d-2}}v = 0. \tag{1.23}$$

Any solution v to (1.22) gives rise to a standing wave $e^{i\omega t}v$ to (1.18), and any solution to (1.23) solves the equation (1.19). Here, we introduce the "scaling-operator" T_{ω} to be that for any function f on \mathbb{R}^d ,

$$T_{\omega}f(x) := \omega^{-\frac{1}{p-1}} f(\frac{x}{\sqrt{\omega}}).$$
(1.24)

Then, putting $u := T_{\omega} v$, we see that u solves

$$u - \Delta u - |u|^{p-1}u = 0. \tag{1.25}$$

It is well known that the equation (1.25) has a unique positive solution up to translations which is radially symmetric with respect to some point. We denote the positive solution symmetric about the origin by U. On the other hand, the equation (1.23) possesses a solution of the form

$$W(x) := \left(\frac{\sqrt{d(d-2)}}{1+|x|^2}\right)^{\frac{d-2}{2}}.$$
(1.26)

Here, a positive C^2 -solution of (1.23) is unique up to scalings and translations (see Corollary 8.2 in [8]). Following Brézis-Nirenberg [7], we introduce the variational value

$$\sigma := \inf \left\{ \left\| \nabla u \right\|_{L^2}^2 : u \in \dot{H}^1(\mathbb{R}^d), \ \left\| u \right\|_{L^{2^*}} = 1 \right\}.$$
(1.27)

Then, it is known that $u = W/||W||_{L^{2^*}}$ is a minimizer of the variational problem associated with σ and

$$\sigma^{\frac{d}{2}} = \|\nabla W\|_{L^2}^2 = \|W\|_{L^{2^*}}^{2^*}.$$
(1.28)

We also introduce the variational value

$$m^{\ddagger} := \inf \left\{ \mathcal{I}^{\ddagger}(u) \colon u \in \dot{H}^{1}(\mathbb{R}^{d}) \setminus \{0\}, \ \mathcal{K}^{\ddagger}(u) \le 0 \right\},$$
(1.29)

where

$$\mathcal{I}^{\ddagger}(u) := \mathcal{H}^{\ddagger}(u) - \frac{2}{d(p-1)} \mathcal{K}^{\ddagger}(u)$$
(1.30)

with

$$\mathcal{H}^{\ddagger}(u) := \frac{1}{2} \|\nabla u\|_{L^2}^2 - \frac{1}{2^*} \|u\|_{L^{2^*}}^{2^*}, \qquad (1.31)$$

$$\mathcal{K}^{\ddagger}(u) := \|\nabla u\|_{L^{2}}^{2} - \|u\|_{L^{2^{*}}}^{2^{*}}.$$
(1.32)

Note that

$$\mathcal{I}^{\dagger}(u) = \frac{d(p-1) - 4}{2d(p-1)} \|\nabla u\|_{L^2}^2 + \frac{4 - (d-2)(p-1)}{2d(p-1)} \|u\|_{L^{2*}}^{2*}.$$
 (1.33)

In particular, $\mathcal{I}^{\ddagger}(u) \ge 0$ for $1 + \frac{4}{d} \le p \le 2^* - 1$.

We can verify (see Appendix A) that

$$m^{\ddagger} = \mathcal{I}^{\ddagger}(W) = \mathcal{H}^{\ddagger}(W) = \frac{1}{d}\sigma^{\frac{d}{2}}.$$
(1.34)

Furthermore, we proved in [2] that for any $d \ge 4$, any $1 + \frac{4}{d} and any <math>\omega > 0$,

$$0 < m_{\omega} < \frac{1}{d}\sigma^{\frac{d}{2}}.\tag{1.35}$$

We can also prove (1.35) for d = 3 if $\omega > 0$ is sufficiently small (see Lemma A.4).

Now, we go back to the equation (1.7). If u is a solution to (1.7), then $v := T_{\omega}u$ satisfies that

$$v - \Delta v - |v|^{p-1}v - \omega^{\frac{2^* - (p+1)}{p-1}} |v|^{\frac{4}{d-2}}v = 0.$$
(1.36)

We should mention that if ω is small enough, it is expected that (1.7) has properties similar to (1.25). Indeed, this is the heart of our analysis.

We briefly review results for (1.18) and (1.19) which are used in this paper.

In [18], Kenig and Merle studied the equation (1.19) in the dimensions d = 3, 4, 5, and proved the scattering of radial solutions starting from the set

$$PW_{+}^{\ddagger} := \left\{ u \in H^{1}(\mathbb{R}^{d}) \colon \mathcal{H}^{\ddagger}(u) < \mathcal{H}^{\ddagger}(W), \ \|\nabla u\|_{L^{2}}^{2} < \|\nabla W\|_{L^{2}}^{2} \right\}.$$
(1.37)

Their result is extended to the higher dimensional cases by Killip and Visan [21]. We summarize these results:

Theorem 1.2 ([18, 21]). Assume $d \ge 3$. Then, the set PW_+^{\ddagger} is invariant under the flow defined by (1.19). Furthermore, any non-trivial radial solution ψ to (1.19) starting from PW_+^{\ddagger} exists globally in time, and satisfies that

$$\|\nabla\psi\|_{St(\mathbb{R})} < \infty, \tag{1.38}$$

$$\mathcal{H}^{\ddagger}(\psi) \ge \frac{1}{2} \inf_{t \in \mathbb{R}} \mathcal{K}^{\ddagger}(\psi(t)) > 0, \qquad (1.39)$$

where $St(\mathbb{R}) := L^{\infty}(\mathbb{R}, L^2(\mathbb{R}^d)) \cap L^2(\mathbb{R}, L^{2^*}(\mathbb{R}^d)).$

The result for (1.18) corresponding to Theorem 1.2 is derived by Duyckaerts, Holmer and Roudenko [11, 14] for d = p = 3. Furthermore, the first and the fourth authors extended their result to the general dimensions $d \ge 1$ and the powers p satisfying (1.1) in [1]; the result says that we have either the scattering or the blowup, if the solutions start from the set

$$PW^{\dagger} := \left\{ u \in H^1(\mathbb{R}^d) \setminus \{0\} \colon \mathcal{H}^{\dagger}(u) < \left(\frac{\mathcal{M}(U)}{\mathcal{M}(u)}\right)^{\frac{1-s_p}{s_p}} \mathcal{H}^{\dagger}(U) \right\},$$
(1.40)

where

$$\mathcal{H}^{\dagger}(u) := \frac{1}{2} \|\nabla u\|_{L^2}^2 - \frac{1}{p+1} \|u\|_{L^{p+1}}^{p+1}, \qquad (1.41)$$

and s_p is the "scaling-exponent" given by

$$s_p := \frac{d}{2} - \frac{2}{p-1}.$$
(1.42)

The condition (1.1) on p implies that $0 < s_p < 1$. Furthermore, we can classify the behavior of the solutions by the sign of the functional \mathcal{K}^{\dagger} defined by

$$\mathcal{K}^{\dagger}(u) := \|\nabla u\|_{L^{2}}^{2} - \frac{d(p-1)}{2(p+1)} \|u\|_{L^{p+1}}^{p+1}.$$
(1.43)

Note here that for any $\omega > 0$, the replacement of U with $T_{\omega}U$ in (1.40) leaves PW^{\dagger} unchanged. Moreover, introducing the set PW_{ω}^{\dagger} as

$$PW_{\omega}^{\dagger} := \left\{ u \in H^1(\mathbb{R}^d) \setminus \{0\} \colon \mathcal{S}_{\omega}^{\dagger}(u) < \mathcal{S}_{\omega}^{\dagger}(U) \right\},$$
(1.44)

we can express PW^{\dagger} as the union of PW_{ω}^{\dagger} over all $\omega > 0$:

$$PW^{\dagger} = \bigcup_{\omega > 0} PW_{\omega}^{\dagger}.$$
 (1.45)

Here, $\mathcal{S}^{\dagger}_{\omega}$ denotes the action for (1.22), namely,

$$S^{\dagger}_{\omega}(u) := \omega \mathcal{M}(u) + \mathcal{H}^{\dagger}(u). \tag{1.46}$$

In [25], Nakanishi and Schlag considered the equation (1.18) in the case d = p = 3. They developed a method to analyze the behavior of (radial) solutions starting from the set

$$PW^{\dagger,\varepsilon} := \left\{ u \in H^1(\mathbb{R}^3) \setminus \{0\} \colon \mathcal{H}^{\dagger}(u) < \left(\frac{\mathcal{M}(U)}{\mathcal{M}(u)}\right) (\mathcal{H}^{\dagger}(U) + \varepsilon) \right\}$$
(1.47)

for some $\varepsilon > 0$. Clearly, the set $PW^{\dagger,\varepsilon}$ is an enlargement of PW^{\dagger} . In particular, they proved that there exists $\varepsilon > 0$ such that all solutions starting from $PW^{\dagger,\varepsilon}$ exhibit one of the same nine scenarios as Theorem 1.1 above. This result motivated our study.

We note that the way to define PW^{\dagger} (see (1.40)) is based on the scaling-invariant nature of U (cf. [1]). Due to lack of such a scaling property, it is not appropriate to define the corresponding "potential well" for our equation (NLS) by simply replacing \mathcal{H}^{\dagger} and U with \mathcal{H} and Φ_{ω} , respectively in PW^{\dagger} . Instead, from the viewpoint of (1.45), we consider the frequency-wise "potential well" PW_{ω} defined by

$$PW_{\omega} := \left\{ u \in H^1(\mathbb{R}^d) \colon \mathcal{S}_{\omega}(u) < m_{\omega} \right\}.$$
(1.48)

This set is closely related to a variational nature of Φ_{ω} . Indeed, the definition of m_{ω} (see (1.9)) implies that if $u \in H^1(\mathbb{R}^d) \setminus \{0\}$ and $\mathcal{K}(u) = 0$, then $\mathcal{S}_{\omega}(u) \ge m_{\omega}$, so that we can split PW_{ω} into three parts according to the sign of the functional \mathcal{K} :

$$PW_{\omega} = PW_{\omega,+} \cup \{0\} \cup PW_{\omega,-}, \tag{1.49}$$

where

$$PW_{\omega,+} := \left\{ u \in H^1(\mathbb{R}^d) \colon \mathcal{S}_{\omega}(u) < m_{\omega}, \ \mathcal{K}(u) > 0 \right\},$$
(1.50)

$$PW_{\omega,-} := \left\{ u \in H^1(\mathbb{R}^d) \colon \mathcal{S}_{\omega}(u) < m_{\omega}, \ \mathcal{K}(u) < 0 \right\}.$$
(1.51)

The following theorem follows from the results in [2, 3] together with the existence theorem of ground state in \mathbb{R}^3 (see Proposition A.1 and Proposition A.4, and [32]).¹

Theorem 1.3. Let $\omega > 0$. Assume $d \ge 3$ and $1 + \frac{4}{d} . Furthermore, assume <math>\omega \in (0, \omega_3)$ if d = 3, where ω_3 is the frequency given in Proposition A.1. Then, the following hold: $PW_{\omega,+}$ and $PW_{\omega,-}$ are invariant under the flow defined by (NLS); and (i) Any radial solution ψ to (NLS) starting from $PW_{\omega,+}$ satisfies

$$\|\langle \nabla \rangle \psi\|_{St(\mathbb{R})} < \infty, \tag{1.52}$$

$$\mathcal{H}(\psi) \ge \frac{1}{2} \inf_{t \in \mathbb{R}} \mathcal{K}(\psi(t)) > 0, \qquad (1.53)$$

¹The papers [2, 3] dealt with the case where $d \ge 4$ and $2 + \frac{4}{d} only. However, we can easily verify that the arguments in [2, 3] work well for the case where <math>d = 3$ and $1 + \frac{4}{3} as long as there exists a ground state <math>Q_{\omega}$ with $m_{\omega} = S_{\omega}(Q_{\omega})$.

where $St(\mathbb{R}) := L^{\infty}(\mathbb{R}, L^2(\mathbb{R}^d)) \cap L^2(\mathbb{R}, L^{2^*}(\mathbb{R}^d))$. In particular, ψ scatters both forward and backward in time.

(ii) Any radial solution ψ to (NLS) starting from $PW_{\omega,-}$ satisfies

$$\sup_{t \in I_{\max}(\psi)} \mathcal{K}(\psi(t)) < 0, \tag{1.54}$$

and blows up in a finite time both forward and backward in time, where $I_{\max}(\psi)$ denotes the maximal existence-interval of ψ .

Note that Theorem 1.1 is an extension of Theorem 1.3.

This paper is organized as follows. In Section 2, we give fundamental properties of ground state of (1.7). In Section 3, we prove Theorem 1.1. The main ingredient of the proof is Theorem 3.1. We assign preliminaries for the proof of Theorem 3.1 to Section 4 through Section 7: In Section 4, we think about the decomposition of solution around the ground state in the same viewpoint of Nakanishi and Schlag [25]. In Section 5, we prove the "ejection lemma" for the equation (NLS). In Section 6, we introduce the distance function used in [25], and derive some variational lemmas. In Section 7, we prove the "one-pass theorem". In Section 8, we finally prove Theorem 3.1. This paper also has appended sections: In Section A, we prove the existence of ground state of (1.7) in three dimensions. In Section B, we state fundamental properties of linearized operator around a ground state of (1.7). In Section C, we give two well-known inequalities for radial functions. In Section D, we record a small-data theory, and in Section E we give a long-time perturbation theory in a general setting.

Notation. Besides the notation introduced so far, we use the following notation (see also the table of notation in Section F):

(i) We use $\partial_{\omega} \Phi_{\omega}$ to denote the derivative of the positive radial grand state Φ_{ω} of (1.7) with respect to ω (see Proposition 2.4 below for the differentiability).

(ii) The daggered symbols are related to (1.25):

$$L_{+}^{\dagger} := 1 - \Delta - pU^{p-1}, \qquad (1.55)$$

$$L_{-}^{\dagger} := 1 - \Delta - U^{p-1}. \tag{1.56}$$

(iii) Functionals with tilde are related to the rescaled equation (1.36):

$$\widetilde{\mathcal{K}}_{\omega}(u) := \|\nabla u\|_{L^{2}}^{2} - \frac{d(p-1)}{2(p+1)} \|u\|_{L^{p+1}}^{p+1} - \omega^{\frac{2^{*}-(p+1)}{p-1}} \|u\|_{L^{2^{*}}}^{2^{*}},$$
(1.57)

$$\widetilde{\mathcal{I}}_{\omega}(u) := \frac{1}{2} \|u\|_{L^{2}}^{2} + \frac{s_{p}}{d} \|\nabla u\|_{L^{2}}^{2} + \omega^{\frac{2^{*} - (p+1)}{p-1}} \frac{1 - s_{p}}{d} \|u\|_{L^{2^{*}}}^{2^{*}},$$
(1.58)

$$\widetilde{L}_{\omega,+} := 1 - \Delta - p \left(T_{\omega} \Phi_{\omega} \right)^{p-1} - \omega^{\frac{2^* - (p+1)}{p-1}} (2^* - 1) \left(T_{\omega} \Phi_{\omega} \right)^{\frac{4}{d-2}}, \tag{1.59}$$

$$\widetilde{L}_{\omega,-} := 1 - \Delta - \left(T_{\omega}\Phi_{\omega}\right)^{p-1} - \omega^{\frac{2^* - (p+1)}{p-1}} \left(T_{\omega}\Phi_{\omega}\right)^{\frac{4}{d-2}}.$$
(1.60)

(iv) We use $(\cdot, \cdot)_{L^2}$ to denote the inner product in $L^2(\mathbb{R}^d)$:

$$(u,v)_{L^2} := \int_{\mathbb{R}^d} u(x)\overline{v(x)} \, dx.$$

We also use $L^2_{real}(\mathbb{R}^d)$ to denote the real Hilbert space of complex-valued functions in $L^2(\mathbb{R}^d)$ which is equipped with the inner product

$$(u,v)_{L^2_{real}} := \Re \int_{\mathbb{R}^d} u(x) \overline{v(x)} \, dx.$$

Furthermore, $H^1_{real}(\mathbb{R}^d)$ denotes the real Hilbert space of functions in $H^1(\mathbb{R}^d)$ equipped with the inner product

$$(u,v)_{H^1_{real}} := (u,v)_{L^2_{real}} + (\nabla u, \nabla v)_{L^2_{real}}$$

(v) We use $\langle v, u \rangle_{H^{-1}, H^1}$ to denote the duality pair of $u \in H^1_{real}(\mathbb{R}^d)$ and $v \in H^{-1}_{real}(\mathbb{R}^d)$:

$$\langle v, u \rangle_{H^{-1}, H^1} := ((1 - \Delta)^{-\frac{1}{2}} v, (1 - \Delta)^{\frac{1}{2}} u)_{L^2_{real}}$$

(vi) Let I be an interval, and let $1 + \frac{4}{d} \le q \le 2^* - 1$. Then, we introduce Strichartz-type spaces on I as

$$St(I) := L_t^{\infty} L_x^2(I) \cap L_t^2 L_x^{2^*}(I),$$
(1.61)

$$V_{q+1}(I) := L_t^{\frac{(d+2)(q-1)}{2}} L_x^{\frac{2d(d+2)(q-1)}{d(d+2)(q-1)-8}}(I),$$
(1.62)

$$W_{q+1}(I) := L_{t,x}^{\frac{(d+2)(q-1)}{2}}(I).$$
(1.63)

Note here that Sobolev's embedding shows $|\nabla|^{-s_q}V_{q+1}(I) \hookrightarrow W_{q+1}(I)$. Moreover, by Strichartz' estimate, we mean the following estimate: for any appropriate space-time function u, any $t_0 \in I$ and any pair $(q, r) \in [2, 2^*] \times [2, \infty]$ with $\frac{2}{r} = d(\frac{1}{2} - \frac{1}{q})$,

$$\|u\|_{St(I)} \lesssim \|u(t_0)\|_{L^2} + \|i\frac{\partial u}{\partial t} + \Delta u\|_{L^{r'}_t L^{q'}_x(I)},$$
(1.64)

where q' and r' denote the Hölder conjugates of q and r respectively.



(vii) For a space-time function u, we define

$$F[u] := |u|^{p-1}u + |u|^{\frac{4}{d-2}}u, \quad F^{\dagger}[u] := |u|^{p-1}u, \quad F^{\dagger}[u] := |u|^{\frac{4}{d-2}}u, \tag{1.65}$$

$$e[u] := i\frac{\partial u}{\partial t} + \Delta u + F[u], \quad e^{\ddagger}[u] := i\frac{\partial u}{\partial t} + \Delta u + F^{\ddagger}[u].$$
(1.66)

2 Properties of Ground State

In this section, we give several fundamental properties of ground state of (1.7). As mentioned in Section 1, for any $d \ge 4$, any $1 + \frac{4}{d} and any <math>\omega > 0$, the equation (1.7) admits at least one ground state Q_{ω} satisfying $m_{\omega} = S_{\omega}(Q_{\omega})$ (see Proposition 1.1 and Theorem 1.1 in [2]). When d = 3, it follows from Proposition A.1 and Proposition A.4 in Appendix A that for any $1 + \frac{4}{3} , there exists <math>\omega_3 > 0$ such that for any $0 < \omega < \omega_3$, the equation (1.7) admits a ground state Q_{ω} satisfying $m_{\omega} = S_{\omega}(Q_{\omega})$.

The following proposition tells us that any ground state of (1.7) is essentially positive and radial:

Proposition 2.1. Assume $d \geq 3$ and $1 + \frac{4}{d} . Let <math>\omega > 0$, and let Q_{ω} be a ground state of (1.7). Then, there exist $\theta \in \mathbb{R}$, $y \in \mathbb{R}^d$ and a positive radial ground state Φ_{ω} of (1.7) such that $\Phi_{\omega} \in C^2(\mathbb{R}^d) \cap H^1(\mathbb{R}^d)$, $\frac{x}{|x|} \cdot \nabla \Phi_{\omega} < 0$ and $Q_{\omega} = e^{i\theta} \Phi_{\omega}(\cdot - y)$.

Proof of Proposition 2.1. Let Q_{ω} be a ground state of (1.7). Then, using (1.10), we can verify that $|Q_{\omega}|$ is also a ground state of (1.7). We also see that $|Q_{\omega}| \in W^{2,q}_{loc}(\mathbb{R}^d)$ for any $2 \leq q < \infty$ (see Theorem 2.3 in [6]). Furthermore, the Schauder theory shows that $Q_{\omega} \in C^2(\mathbb{R}^d)$ and $\lim_{|x|\to\infty} |Q_{\omega}(x)| = 0$. On the other hand, it follows from Theorem 9.10 in [23] that for any compact set K in \mathbb{R}^d , there exists a constant C depending only on K and ω such that for any $x \in K$,

$$|Q_{\omega}(x)| \ge C \int_{K} |Q_{\omega}(y)| \, dy.$$
(2.1)

Hence, $|Q_{\omega}|$ is a positive ground state of (1.7). We see from the result of Gidas, Ni and Nirenberg [12] that there exist a positive radial function Φ_{ω} with $\frac{x}{|x|} \cdot \nabla \Phi_{\omega}(x) < 0$ for any $x \in \mathbb{R}^d \setminus \{0\}$, and $y \in \mathbb{R}^d$, such that $|Q_{\omega}| = \Phi_{\omega}(\cdot - y)$. We introduce the sign of a complex-valued function u, denoted by sgn u, as

$$\operatorname{sgn} u := \frac{\overline{u}}{|u|}.$$
(2.2)

Then, we have

$$\operatorname{sgn}\overline{Q_{\omega}} := \frac{Q_{\omega}}{|Q_{\omega}|},\tag{2.3}$$

so that $Q_{\omega} = |Q_{\omega}| \operatorname{sgn} \overline{Q_{\omega}}$. We shall show that $\operatorname{sgn} \overline{Q_{\omega}}$ is constant, which together with $|\operatorname{sgn} \overline{Q_{\omega}}| \equiv 1$ completes the proof. Since

$$\Re\left[\operatorname{sgn}\overline{Q_{\omega}}\nabla\operatorname{sgn}Q_{\omega}(x)\right] = \Re\left[\frac{Q_{\omega}}{|Q_{\omega}|}\nabla\left(\frac{\overline{Q_{\omega}}}{|Q_{\omega}|}\right)\right] = \frac{\Re\left[Q_{\omega}\overline{\nabla Q_{\omega}}\right] - |Q_{\omega}|\nabla|Q_{\omega}|}{|Q_{\omega}|^{2}} = 0, \quad (2.4)$$

we have

$$\begin{aligned} \|\nabla Q_{\omega}\|_{L^{2}}^{2} &= \left\|\nabla \left(\left|Q_{\omega}|\operatorname{sgn} \overline{Q_{\omega}}\right.\right)\right\|_{L^{2}}^{2} \\ &= \int_{\mathbb{R}^{d}} \left|\nabla |Q_{\omega}(x)|\right|^{2} dx + \int_{\mathbb{R}^{d}} \left|Q_{\omega}(x)\right|^{2} \left|\nabla \operatorname{sgn} \overline{Q_{\omega}}(x)\right|^{2} dx \\ &+ \int_{\mathbb{R}^{d}} \nabla |Q_{\omega}(x)|^{2} \Re \left[\operatorname{sgn} \overline{Q_{\omega}} \nabla \operatorname{sgn} Q_{\omega}(x)\right] dx \\ &= \int_{\mathbb{R}^{d}} \left|\nabla |Q_{\omega}(x)|\right|^{2} dx + \int_{\mathbb{R}^{d}} \left|Q_{\omega}(x)\right|^{2} \left|\nabla \operatorname{sgn} \overline{Q_{\omega}}(x)\right|^{2} dx. \end{aligned}$$
(2.5)

Hence, if $\nabla \operatorname{sgn} \overline{Q_{\omega}} \neq 0$, then we would have

$$m_{\omega} = \mathcal{S}_{\omega}(Q_{\omega}) > \mathcal{S}_{\omega}(|Q_{\omega}|) = \mathcal{S}_{\omega}(\Phi_{\omega}) = m_{\omega}.$$
 (2.6)

However, this is a contradiction. Thus, $\operatorname{sgn} \overline{Q_{\omega}}$ must be constant.

Next, we give a decay property of a positive ground state.

Proposition 2.2. Assume $d \geq 3$ and $1 + \frac{4}{d} . Let <math>\omega > 0$, and let $\Phi_{\omega} \in C^2(\mathbb{R}^d) \cap H^1(\mathbb{R}^d)$ be a positive radial ground state of (1.7). Then, there exist positive constants $C(\omega) > 0$ and $\delta(\omega) > 0$ depending on ω such that

$$|\Phi_{\omega}(x)| + |\nabla \Phi_{\omega}(x)| + |\Delta \Phi_{\omega}(x)| \le C(\omega)e^{-\delta(\omega)|x|}$$
(2.7)

for any $x \in \mathbb{R}^d$. In particular, $\Phi_\omega \in H^2(\mathbb{R}^d)$.

Proof of Proposition 2.2. We can prove the proposition in a way similar to [5].

The following convergence result justifies the intuition that the equation (1.7) looks like (1.25) for a sufficiently small ω .

Proposition 2.3. Assume $d \ge 3$ and $1 + \frac{4}{d} . Let <math>\omega > 0$, and let Φ_{ω} be a positive radial ground state of (1.7). Moreover, let U be the unique positive radial ground state of (1.25). Then, we have

$$\lim_{\omega \downarrow 0} \|T_{\omega} \Phi_{\omega} - U\|_{H^1} = 0.$$
(2.8)

Proof of Proposition 2.3. We can prove the proposition in a way similar to [16]. \Box

Remark 2.1. We see from Proposition 2.3 that for any $1 \le q \le 2^* - 1$,

$$\lim_{\omega \downarrow 0} \omega^{s_p - \frac{q-1}{p-1}} \|\Phi_{\omega}\|_{L^{q+1}}^{q+1} = \|U\|_{L^{q+1}}^{q+1}$$
(2.9)

and

$$\lim_{\omega \downarrow 0} \omega^{s_p - 1} \| \nabla \Phi_{\omega} \|_{L^2}^2 = \| \nabla U \|_{L^2}^2.$$
(2.10)

Proposition 2.4. Assume that $d \ge 3$ and $1 + \frac{4}{d} . Then, there exists <math>\omega_1 > 0$ with the following properties:

(i) For any $\omega \in (0, \omega_1)$, a positive radial ground state of (1.7) uniquely exists.

(ii) The mapping $\omega \in (0, \omega_1) \mapsto \Phi_\omega \in H^1(\mathbb{R}^d)$ is continuously differentiable.

(iii) For any $\omega \in (0, \omega_1)$,

$$\frac{d}{d\omega}\mathcal{M}(\Phi_{\omega}) = \left(\Phi_{\omega}, \partial_{\omega}\Phi_{\omega}\right)_{L^{2}_{real}} < 0.$$
(2.11)

Here, Φ_{ω} denotes the unique positive radial ground state of (1.7).

In order to prove the claim (i) in Proposition 2.4, we need the following result (see, e.g., Theorem 0.2 in [17], [26] and Proposition 2.8 in [31]):

Lemma 2.5. Assume either d = 1, 2 and $1 , or <math>d \ge 3$ and 1 .Then, we have

$$\operatorname{Ker} L_{+}^{\dagger} = \operatorname{span} \{ \partial_{1} U, \dots, \partial_{d} U \}, \qquad (2.12)$$

where L_{+}^{\dagger} is the operator defined by (1.55), and $\partial_{1}U, \ldots, \partial_{d}U$ denote the partial derivatives of U.

Now, we give a proof of the claim (i):

Proof of (i) in Proposition 2.4. It suffices to prove the uniqueness of the ground state. Suppose for contradiction that for any $k \in \mathbb{N}$, there exists $\omega_k \in (0, \frac{1}{k})$ such that the equation (1.7) has two different positive radial ground states, say U_{ω_k} and V_{ω_k} . Put $\widetilde{U}_k := T_{\omega_k} U_{\omega_k}$, $\widetilde{V}_k := T_{\omega_k} V_{\omega_k}$ and

$$\widetilde{u}_k := \frac{\widetilde{U}_k - \widetilde{V}_k}{\|\widetilde{U}_k - \widetilde{V}_k\|_{H^1}}.$$
(2.13)

Then, \widetilde{u}_k satisfies the equation

$$\widetilde{u}_{k} - \Delta \widetilde{u}_{k} - \frac{\widetilde{U}_{k}^{p} - \widetilde{V}_{k}^{p}}{\|\widetilde{U}_{k} - \widetilde{V}_{k}\|_{H^{1}}} - \omega_{k}^{\frac{2^{*} - (p+1)}{p-1}} \frac{\widetilde{U}_{k}^{2^{*} - 1} - \widetilde{V}_{k}^{2^{*} - 1}}{\|\widetilde{U}_{k} - \widetilde{V}_{k}\|_{H^{1}}} = 0.$$
(2.14)

Moreover, it follows from $\widetilde{u}_k \in H^1_{rad}(\mathbb{R}^d)$ and $\|\widetilde{u}_k\|_{H^1} = 1$ for any $k \in \mathbb{N}$ that there exists a radial function $\widetilde{u} \in H^1(\mathbb{R}^d)$ such that, passing to a subsequence, we have

$$\lim_{k \to \infty} \widetilde{u}_k = \widetilde{u} \qquad \text{weakly in } H^1(\mathbb{R}^d), \tag{2.15}$$

and for any $2 < q < 2^*$,

$$\lim_{k \to \infty} \widetilde{u}_k = \widetilde{u} \qquad \text{strongly in } L^q(\mathbb{R}^d).$$
(2.16)

We shall show that $\tilde{u} \equiv 0$. First note that the fundamental theorem of calculus together with Proposition 2.3 and (2.16) implies that for any $\phi \in C_c^{\infty}(\mathbb{R}^d)$,

$$\lim_{k \to \infty} \left\langle \frac{U_k^p - V_k^p}{\|\widetilde{U}_k - \widetilde{V}_k\|_{H^1}}, \phi \right\rangle_{H^{-1}, H^1}$$

$$= \lim_{k \to \infty} p \int_0^1 \left\langle \left\{ \widetilde{V}_k + \theta(\widetilde{U}_k - \widetilde{V}_k) \right\}^{p-1} \widetilde{u}_k, \phi \right\rangle_{H^{-1}, H^1} d\theta \qquad (2.17)$$

$$= p \int_{\mathbb{R}^d} U^{p-1} \widetilde{u} \phi \, dx.$$

We can also verify that for a given function $\phi \in C_c^{\infty}(\mathbb{R}^d)$,

$$\lim_{k \to \infty} \omega_k^{\frac{2^* - (p+1)}{p-1}} \left\langle \frac{\widetilde{U}_k^{2^* - 1} - \widetilde{V}_k^{2^* - 1}}{\|\widetilde{U}_k - \widetilde{V}_k\|_{H^1}}, \phi \right\rangle_{H^{-1}, H^1} = 0.$$
(2.18)

Combining (2.14) with (2.15), (2.17) and (2.18), we find that for any $\phi \in C_c^{\infty}(\mathbb{R}^d)$,

$$\left\langle L_{+}^{\dagger}\widetilde{u}, \phi \right\rangle_{H^{-1},H^{1}} = \left\langle \widetilde{u} - \Delta \widetilde{u} - pU^{p-1}\widetilde{u}, \phi \right\rangle_{H^{-1},H^{1}} = 0.$$
 (2.19)

Furthermore, since \tilde{u} is a radial H^1 -solution of $L^{\dagger}_{+}\tilde{u} = 0$, we find that $\tilde{u} \in H^2_{rad}(\mathbb{R}^d)$. Hence, it follows from Lemma 2.5 (there is no radial function in the kernel of L^{\dagger}_{+}) that $\tilde{u} \equiv 0$.

Now, multiplying the equation (2.14) by \tilde{u}_k and then integrating the resulting equation over \mathbb{R}^d , we obtain

$$\|\widetilde{u}_{k}\|_{H^{1}}^{2} = \int_{\mathbb{R}^{d}} \frac{\widetilde{U}_{k}^{p} - \widetilde{V}_{k}^{p}}{\|\widetilde{U}_{k} - \widetilde{V}_{k}\|_{H^{1}}} \widetilde{u}_{k} \, dx + \omega_{k}^{\frac{2^{*} - (p+1)}{p-1}} \int_{\mathbb{R}^{d}} \frac{\widetilde{U}_{k}^{2^{*} - 1} - \widetilde{V}_{k}^{2^{*} - 1}}{\|\widetilde{U}_{k} - \widetilde{V}_{k}\|_{H^{1}}} \widetilde{u}_{k} \, dx.$$
(2.20)

Then, it follows from (2.16) and $\tilde{u} \equiv 0$ that the right-hand side of (2.20) tends to 0 as $k \to \infty$, whereas the left-hand side is identically 1 (see (2.13)). This is a contradiction. Thus, we have completed the proof of the claim (i) in Proposition 2.4.

Next, we mention how to prove the claim (ii) in Proposition 2.4.

Proof of (ii) in Proposition 2.4. The claim (ii) follows from the result by Shatah and Strauss [27] (see also [20]). \Box

Finally, we move on to the proof of the claim (iii) in Proposition 2.4. Let us notice that the claim (iii) is an analogy to that

$$\frac{d}{d\omega}\mathcal{M}(T_{\omega^{-1}}U) = \frac{1}{2}\frac{d}{d\omega}\left(\omega^{-s_p}\|U\|_{L^2}^2\right) = -\frac{s_p}{2}\omega^{-s_p-1}\|U\|_{L^2}^2 < 0.$$
(2.21)

Hence, it is convenient to introduce a function ΛU defined by

$$\Lambda U(x) := \partial_{\omega} \{ T_{\omega^{-1}} U(x) \} \Big|_{\omega = 1} = \frac{1}{p - 1} U(x) + \frac{1}{2} x \cdot \nabla U(x).$$
(2.22)

Then, we can verify that ΛU obeys

$$L_{+}^{\dagger}\Lambda U = -U. \tag{2.23}$$

Differentiation of the both sides of the equation (1.7) with respect to ω yields

$$\omega \partial_{\omega} \Phi_{\omega} - \Delta \partial_{\omega} \Phi_{\omega} - p \Phi_{\omega}^{p-1} \partial_{\omega} \Phi_{\omega} - (2^* - 1) \Phi_{\omega}^{\overline{d-2}} \partial_{\omega} \Phi_{\omega} = -\Phi_{\omega}.$$
(2.24)

Furthermore, we see from (2.24) that

$$\widetilde{L}_{\omega,+}(\omega T_{\omega}\partial_{\omega}\Phi_{\omega}) = -T_{\omega}\Phi_{\omega}, \qquad (2.25)$$

where $\widetilde{L}_{\omega,+}$ is the operator defined by (1.59). We state a property of the operator $\widetilde{L}_{\omega,+}$.

Lemma 2.6. Assume $d \ge 3$ and $1 + \frac{4}{d} . Then, there exist <math>\omega_0 > 0$ and C > 0 such that for any $\omega \in (0, \omega_0)$ and any $f \in H^2_{rad}(\mathbb{R}^d)$,

$$\|\widetilde{L}_{\omega,+}f\|_{L^2} \ge C \|f\|_{H^1}.$$
(2.26)

Proof of Lemma 2.6. Suppose for contradiction that for any $k \in \mathbb{N}$, there exist $\omega_k \in (0, \frac{1}{k})$ and $f_k \in H^2_{rad}(\mathbb{R}^d)$ with $||f_k||_{H^1} = 1$ such that

$$\|\widetilde{L}_{\omega_k,+}f_k\|_{L^2} < \frac{1}{k}.$$
(2.27)

Then, we can take $f_{\infty} \in H^1_{rad}(\mathbb{R}^d)$ such that, passing to some subsequence,

$$\lim_{k \to \infty} f_k = f_{\infty} \qquad \text{weakly in } H^1(\mathbb{R}^d), \tag{2.28}$$

and for any $2 < q < 2^*$,

$$\lim_{k \to \infty} f_k = f_{\infty} \qquad \text{strongly in } L^q(\mathbb{R}^d).$$
(2.29)

Furthermore, for any $\phi \in C_c^{\infty}(\mathbb{R}^d)$, we have

$$\begin{aligned} \left| \langle L_{+}^{\dagger} f_{\infty}, \phi \rangle_{H^{-1}, H^{1}} \right| \\ &\leq \left| \langle \widetilde{L}_{\omega_{k}, +} f_{k}, \phi \rangle_{H^{-1}, H^{1}} \right| + \left| \langle \widetilde{L}_{\omega_{k}, +} (f_{\infty} - f_{k}), \phi \rangle_{H^{-1}, H^{1}} \right| \\ &+ p \left| \langle \{ (T_{\omega_{k}} \Phi_{\omega_{k}})^{p-1} - U^{p-1} \} f_{\infty}, \phi \rangle_{H^{-1}, H^{1}} \right| \\ &+ \omega_{k}^{\frac{2^{*} - (p+1)}{p-1}} (2^{*} - 1) \left| \langle (T_{\omega_{k}} \Phi_{\omega_{k}})^{\frac{4}{d-2}} f_{\infty}, \phi \rangle_{H^{-1}, H^{1}} \right|. \end{aligned}$$
(2.30)

This together with the hypothesis (2.27), (2.28), (2.29), Proposition 2.3 and $\lim_{k\to\infty} \omega_k = 0$ shows that $L^{\dagger}_+ f_{\infty} = 0$ in the distribution sense. Since $(\operatorname{Ker} L^{\dagger}_+)|_{H^1_{rad}} = \{0\}$ (see Lemma 2.5), we conclude that $f_{\infty} = 0$. Furthermore, this fact $f_{\infty} = 0$ together with $||f_k||_{H^1} = 1$, (2.29) and $\lim_{k\to\infty} \omega_k = 0$ gives us that

$$\lim_{k \to \infty} (\tilde{L}_{\omega_k, +} f_k, f_k)_{L^2} = 1.$$
(2.31)

However, it follows from the hypothesis (2.27) that

$$\lim_{k \to \infty} \left| (\widetilde{L}_{\omega_k, +} f_k, f_k)_{L^2} \right| \le \lim_{k \to \infty} \| \widetilde{L}_{\omega_k, +} f_k \|_{L^2} = 0.$$
(2.32)

This is a contradiction. Thus, we have proved that (2.26) holds.

Lemma 2.7. Assume $d \ge 3$ and $1 + \frac{4}{d} . Then, we have$

$$\lim_{\omega \downarrow 0} \|\omega T_{\omega} \partial_{\omega} \Phi_{\omega} - \Lambda U\|_{H^1} = 0.$$
(2.33)

Proof of Lemma 2.7. Let $\omega_0 > 0$ and C > 0 be constants given in Lemma 2.6. Furthermore, let $\{\omega_n\}$ be a sequence in $(0, \omega_0)$ with $\lim_{n\to\infty} \omega_n = 0$, and set

$$\Lambda \Phi_{\omega_n} := \partial_\omega \Phi_\omega|_{\omega = \omega_n}. \tag{2.34}$$

Then, we find from Proposition 2.6, (2.25) and Proposition 2.3 that there exists a number N such that for any $n \ge N$,

$$C\|\omega_n T_{\omega_n} \Lambda \Phi_{\omega_n}\|_{H^1} \le \left\|\widetilde{L}_{\omega_n,+}(\omega_n T_{\omega_n} \Lambda \Phi_{\omega_n})\right\|_{L^2} = \|T_{\omega_n} \Phi_{\omega_n}\|_{H^1} \le 2\|U\|_{H^1}.$$
 (2.35)

Thus, $\{\omega_n T_{\omega_n} \Lambda \Phi_{\omega_n}\}$ is bounded in $H^1(\mathbb{R}^d)$ and therefore we can take $g_{\infty} \in H^1(\mathbb{R}^d)$ such that, passing to some subsequence,

$$\lim_{n \to \infty} \omega_n T_{\omega_n} \Lambda \Phi_{\omega_n} = g_{\infty} \qquad \text{weakly in } H^1(\mathbb{R}^d).$$
(2.36)

This together with Proposition 2.3 also shows that for any $\phi \in C_0^{\infty}(\mathbb{R}^d)$,

$$\lim_{n \to \infty} \langle \widetilde{L}_{\omega_n, +}(\omega_n T_{\omega_n} \Lambda \Phi_{\omega_n}), \phi \rangle_{H^{-1}, H^1} = \langle L_+^{\dagger} g_{\infty}, \phi \rangle_{H^{-1}, H^1}.$$
(2.37)

On the other hand, we see from (2.25) and Proposition 2.3 that

$$\lim_{n \to \infty} \widetilde{L}_{\omega_n, +}(\omega_n T_{\omega_n} \Lambda \Phi_{\omega_n}) = -U \quad \text{strongly in } H^1(\mathbb{R}^d).$$
(2.38)

Putting (2.37) and (2.38) together, we find that $L^{\dagger}_{+}g_{\infty} = -U$. Furthermore, since $(\operatorname{Ker} L^{\dagger}_{+})|_{H^{1}_{rad}} = \{0\}$ (see Lemma 2.5), this identity together with (2.23) shows that $g_{\infty} = \Lambda U$. It remains to show that

$$\lim_{n \to \infty} \|\omega_n T_{\omega_n} \Lambda \Phi_{\omega_n}\|_{H^1} = \|\Lambda U\|_{H^1}.$$
(2.39)

We see from (2.25), Proposition 2.2, Proposition 2.3, (2.36) with $g_{\infty} = \Lambda U$ and (2.23) that

$$\begin{split} \lim_{n \to \infty} \|\omega_n T_{\omega_n} \Lambda \Phi_{\omega_n}\|_{H^1}^2 \\ &= -\lim_{n \to \infty} (T_{\omega_n} \Phi_{\omega_n}, \, \omega_n T_{\omega_n} \Lambda \Phi_{\omega_n})_{L^2_{real}} \\ &+ \lim_{n \to \infty} p((T_{\omega_n} \Phi_{\omega_n})^{p-1} \omega_n T_{\omega_n} \Lambda \Phi_{\omega_n}, \omega_n T_{\omega_n} \Lambda \Phi_{\omega_n})_{L^2_{real}} \\ &+ \lim_{n \to \infty} \omega_n^{\frac{2^* - (p+1)}{p-1}} (2^* - 1)((T_{\omega_n} \Phi_{\omega_n})^{\frac{4}{d}} \omega_n T_{\omega_n} \Lambda \Phi_{\omega_n}, \omega_n T_{\omega_n} \Lambda \Phi_{\omega_n})_{L^2_{real}} \\ &= -(U, \Lambda U)_{L^2_{real}} + p(U^{p-1} \Lambda U, \Lambda U)_{L^2_{real}} \\ &= (L^{\dagger}_+ \Lambda U, \Lambda U)_{L^2_{real}} + p(U^{p-1} \Lambda U, \Lambda U)_{L^2_{real}} = \|\Lambda U\|_{H^1}^2. \end{split}$$

Thus, we have completed the proof.

Now, we are in a position to prove the claim (iii) in Proposition 2.4.

Proof of (iii) in Proposition 2.4. We see from Proposition 2.3, (2.7) and (2.22) that

$$\lim_{\omega \to 0} \omega^{s_p + 1} \frac{d}{d\omega} \mathcal{M}(\Phi_{\omega}) = \lim_{\omega \to 0} \omega^{s_p + 1} (\Phi_{\omega}, \Phi'_{\omega})_{L^2_{real}} = \lim_{\omega \to 0} (T_{\omega} \Phi_{\omega}, \omega T_{\omega} \partial_{\omega} \Phi_{\omega})_{L^2_{real}}$$
(2.41)
$$= (U, \Lambda U)_{L^2_{real}} = -\frac{s_p}{2} \|U\|_{L^2}^2 < 0,$$

which gives us the desired result.

3 Proof of Theorem 1.1

In order to prove Theorem 1.1, for a given $\varepsilon \geq 0$, we consider the set

$$A^{\varepsilon}_{\omega} := \left\{ u \in H^1(\mathbb{R}^d) \colon \mathcal{S}_{\omega}(u) < m_{\omega} + \varepsilon, \ \mathcal{M}(u) = \mathcal{M}(\Phi_{\omega}) \right\}.$$
(3.1)

Then, a key fact to prove Theorem 1.1 is the following:

L		I
L		I

Theorem 3.1. Assume either d = 3 and $3 \le p < 5$, or $d \ge 4$ and $1 + \frac{4}{d-1} .$ $Then, there exists <math>\omega_* > 0$ with the following property: for any $\omega \in (0, \omega_*)$, there exists a positive constant $\varepsilon(\omega)$ such that all radial solutions starting from $A_{\omega}^{\varepsilon(\omega)}$ exhibit one of the following scenarios:

(i) Scattering both forward and backward in time;

- (ii) Finite time blowup both forward and backward in time;
- (iii) Scattering forward in time, and finite time blowup backward in time;
- (iv) Finite time blowup forward in time, and scattering backward in time;
- (v) Trapped by $\mathcal{O}(\Phi_{\omega})$ forward in time, and scattering backward in time;
- (vi) Scattering forward in time, and trapped by $\mathcal{O}(\Phi_{\omega})$ backward in time;
- (vii) Trapped by $\mathcal{O}(\Phi_{\omega})$ forward in time, and finite time blowup backward in time;
- (viii) Finite time blowup forward in time, and trapped by $O(\Phi_{\omega})$ backward in time;
- (ix) Trapped by $O(\Phi_{\omega})$ both forward and backward in time.

We give a proof of this theorem in Section 8.

In order to prove Theorem 1.1, we also use the following fact:

Lemma 3.1. Assume $d \ge 3$ and $1 + \frac{4}{d} . Let <math>\omega_1$ be the frequency given by Proposition 2.4. Then, we have the following:

(i) m_{ω} is differentiable on $(0, \omega_1)$, and

$$\frac{dm_{\omega}}{d\omega} = \mathcal{M}(\Phi_{\omega}) \tag{3.2}$$

for all $\omega \in (0, \omega_1)$. In particular, m_{ω} is strictly increasing on $(0, \omega_1)$. (ii) $\frac{m_{\omega}}{\omega}$ is differentiable and strictly decreasing on $(0, \omega_1)$.

(iii) Let $0 < \alpha < \beta < \omega_1$. Then, we have that

$$\mathcal{M}(\Phi_{\beta}) < \frac{m_{\beta} - m_{\alpha}}{\beta - \alpha} < \mathcal{M}(\Phi_{\alpha}).$$
(3.3)

Proof of Lemma 3.1. We shall prove the claim (i). Since $\mathcal{S}'(\Phi_{\omega}) = 0$ and Φ_{ω} is differentiable with respect to ω on $(0, \omega_1)$ (see (ii) of Proposition 2.4), we see that for any $\omega \in (0, \omega_1)$,

$$\frac{dm_{\omega}}{d\omega} = \frac{d}{d\omega} \mathcal{S}_{\omega}(\Phi_{\omega}) = \mathcal{M}(\Phi_{\omega}) + \mathcal{S}'_{\omega}(\Phi_{\omega})\partial_{\omega}\Phi_{\omega} = \mathcal{M}(\Phi_{\omega}).$$
(3.4)

Thus, we have proved the first claim.

Next, we shall prove the second claim (ii). The differentiability of $\frac{m_{\omega}}{\omega}$ follows from the first claim. Furthermore, we conclude from (ii) of Proposition 2.4 that for any $\omega \in (0, \omega_1)$,

$$\frac{d}{d\omega}\left(\frac{m_{\omega}}{\omega}\right) = \frac{d}{d\omega}\left(\frac{\mathcal{S}_{\omega}(\Phi_{\omega})}{\omega}\right) = \frac{\mathcal{S}_{\omega}'(\Phi_{\omega})\partial_{\omega}\Phi_{\omega}\omega - \mathcal{S}_{\omega}(\Phi_{\omega})}{\omega^2} = -\frac{\mathcal{S}_{\omega}(\Phi_{\omega})}{\omega^2} < 0.$$
(3.5)

Thus, we find that $\frac{m_{\omega}}{\omega}$ is strictly decreasing on $(0, \omega_1)$.

Finally, we shall prove the last claim (iii). It follows from the mean value theorem and (3.2) that for any $0 < \alpha < \beta < \omega_1$, there exists $\theta \in (0, 1)$ such that

$$\frac{m_{\beta} - m_{\alpha}}{\beta - \alpha} = \mathcal{M}(\Phi_{\alpha + \theta(\beta - \alpha)}).$$
(3.6)

On the other hand we see from (iii) of Proposition 2.4 that for any $\theta \in (0, 1)$,

$$\mathcal{M}(\Phi_{\beta}) < \mathcal{M}(\Phi_{\alpha+\theta(\beta-\alpha)}) < \mathcal{M}(\Phi_{\alpha}).$$
(3.7)

Putting (3.6) and (3.7) together, we obtain the desired result (3.3).

We see from (2.9) with q = 1 and (iii) of Proposition 2.4 that there is a strictly decreasing function $\alpha: (\mathcal{M}(\Phi_{\omega_1}), \infty) \to (0, \omega_1)$ such that $M = \mathcal{M}(\Phi_{\alpha(M)})$ for any $M \in (\mathcal{M}(\Phi_{\omega_1}), \infty)$; α is the inverse of the function $\omega \mapsto \mathcal{M}(\Phi_{\omega})$. Let ω_* be the frequency given by Theorem 3.1. Then, for each $\omega \in (0, \omega_*)$, we define a positive function $\varepsilon_{\omega}: [0, \infty) \to (0, \infty)$ to be that for any $M \ge 0$,

$$\varepsilon_{\omega}(M) := \begin{cases} m_{\omega_*} - m_{\omega} - (\omega_* - \omega)\mathcal{M}(\Phi_{\omega_*}) & \text{if } M \leq \mathcal{M}(\Phi_{\omega_*}), \\ \varepsilon(\alpha(M)) + m_{\alpha(M)} - m_{\omega} - (\alpha(M) - \omega)M & \text{if } \mathcal{M}(\Phi_{\omega_*}) < M < \mathcal{M}(\Phi_{\omega}), \\ \varepsilon(\omega) & \text{if } M = \mathcal{M}(\Phi_{\omega}), \\ \varepsilon(\alpha(M)) + (\omega - \alpha(M))M - (m_{\omega} - m_{\alpha(M)}) & \text{if } M > \mathcal{M}(\Phi_{\omega}), \end{cases}$$

$$(3.8)$$

where $\varepsilon(\alpha(M))$ and $\varepsilon(\omega)$ are the positive constants given by Theorem 3.1.

Remark 3.1. It follows from (iii) of Lemma 3.1 that $\varepsilon_{\omega}(M)$ is positive for all $M \ge 0$. Furthermore, the continuity of $\varepsilon(\omega)$ with respect to ω implies that

$$\inf_{M \ge 0} \varepsilon_{\omega}(M) > 0. \tag{3.9}$$

Now, we are in a position to prove Theorem 1.1.

Proof of Theorem 1.1. Let ω_* be the frequency found in Theorem 3.1, and let $\omega \in (0, \omega_*)$. Furthermore, let ε_{ω} be the positive function define by (3.8), and let \widetilde{PW}_{ω} be the set defined by (1.16).



Figure 2: How to extend the potential well PW_{ω}

We consider a solution ψ to (NLS) starting from \widetilde{PW}_{ω} . If $\mathcal{M}(\psi) \leq \mathcal{M}(\Phi_{\omega_*})$, then we can verify that $\psi \in PW_{\omega_*}$ (see Figure 2 above). Hence, it follows from Theorem 1.3 that ψ exhibits the scenario (i) or (ii) in Theorem 1.1. On the other hand, if $\mathcal{M}(\Phi_{\omega_*}) < \mathcal{M}(\psi) < \mathcal{M}(\Phi_{\omega})$, then we can verify that $\psi \in A_{\alpha}^{\varepsilon(\alpha)}$, where $\alpha := \alpha(\mathcal{M}(\psi))$. Similarly, if $\mathcal{M}(\psi) \geq \mathcal{M}(\Phi_{\omega})$, then $\psi \in A_{\alpha}^{\varepsilon(\alpha)}$. Hence, it follows from Theorem 3.1 that ψ exhibits one of the nine scenarios in Theorem 1.1.

4 Decomposition around ground state

In this section, for a positive radial ground state Φ_{ω} to (1.7) and a solution ψ to (NLS) with the maximal existence-interval I_{max} , we consider the decomposition of the form

$$\psi(x,t) = e^{i\theta(t)} \left(\Phi_{\omega}(x) + \eta(x,t) \right), \tag{4.1}$$

where $\theta(t)$ is some function of $t \in I_{\text{max}}$ to be chosen later, and η is the remainder. In this decomposition, we do not assume the orthogonality between Φ_{ω} and $\eta(t)$ in L^2_{real} (see (4.57) below).

4.1 Linearized operator

The decomposition (4.1) leads us to the linearized operator \mathcal{L}_{ω} around Φ_{ω} which is defined by

$$\mathcal{L}_{\omega}u := \omega u - \Delta u - \frac{p+1}{2}\Phi_{\omega}^{p-1}u - \frac{p-1}{2}\Phi_{\omega}^{p-1}\overline{u} - \frac{2^{*}}{2}\Phi_{\omega}^{\frac{4}{d-2}}u - \frac{2^{*}-2}{2}\Phi_{\omega}^{\frac{4}{d-2}}\overline{u}$$

$$= \left(\omega - \Delta - p\Phi_{\omega}^{p-1} - (2^{*}-1)\Phi_{\omega}^{\frac{4}{d-2}}\right)\Re[u] + i\left(\omega - \Delta - \Phi_{\omega}^{p-1} - \Phi_{\omega}^{\frac{4}{d-2}}\right)\Im[u].$$
(4.2)

The operator \mathcal{L}_{ω} is self-adjoint in $L^2_{real}(\mathbb{R}^d)$, and for any $u, v \in H^1(\mathbb{R}^d)$,

$$\left[\mathcal{S}_{\omega}^{\prime\prime}(\Phi_{\omega})u\right]v = \langle \mathcal{L}_{\omega}u, v \rangle_{H^{-1}, H^{1}}.$$
(4.3)

It is convenient to introduce the operators $L_{\omega,+}$ and $L_{\omega,-}$:

$$L_{\omega,+} := \omega - \Delta - p\Phi_{\omega}^{p-1} - (2^* - 1)\Phi_{\omega}^{\frac{4}{d-2}}, \qquad (4.4)$$

$$L_{\omega,-} := \omega - \Delta - \Phi_{\omega}^{p-1} - \Phi_{\omega}^{\frac{4}{d-2}}.$$
(4.5)

Then, we have

$$\mathcal{L}_{\omega}u = L_{\omega,+}\Re[u] + iL_{\omega,-}\Im[u]. \tag{4.6}$$

Moreover, since Φ_{ω} is a solution to (1.7), we can verify that

$$\mathcal{L}_{\omega}\Phi_{\omega} = L_{\omega,+}\Phi_{\omega} = -(p-1)\Phi_{\omega}^{p} - (2^{*}-2)\Phi_{\omega}^{2^{*}-1}, \qquad (4.7)$$

$$\mathcal{L}_{\omega}(i\Phi_{\omega}) = L_{\omega,-}\Phi_{\omega} = 0, \tag{4.8}$$

$$\mathcal{L}_{\omega}\partial_{\omega}\Phi_{\omega} = L_{\omega,+}\partial_{\omega}\Phi_{\omega} = -\Phi_{\omega}.$$
(4.9)

Inserting the decomposition $\psi(t) = e^{i\theta(t)}(\Phi_{\omega} + \eta(t))$ into the equation (NLS), we obtain the equation for η :

$$\frac{\partial \eta}{\partial t}(t) = -i\mathcal{L}_{\omega}\eta(t) - i\Big\{\frac{d\theta}{dt}(t) - \omega\Big\}(\Phi_{\omega} + \eta(t)) + iN_{\omega}(\eta(t)), \tag{4.10}$$

where $N_{\omega}(\eta)$ denotes the higher order term of η , i.e.,

$$N_{\omega}(\eta) := |\Phi_{\omega} + \eta|^{p-1} (\Phi_{\omega} + \eta) - |\Phi_{\omega}|^{p-1} \Phi_{\omega} - \frac{p+1}{2} \Phi_{\omega}^{p-1} \eta - \frac{p-1}{2} \Phi_{\omega}^{p-1} \overline{\eta} + |\Phi_{\omega} + \eta|^{\frac{4}{d-2}} (\Phi_{\omega} + \eta) - |\Phi_{\omega}|^{\frac{4}{d-2}} \Phi_{\omega} - \frac{2^{*}}{2} \Phi_{\omega}^{\frac{4}{d-2}} \eta - \frac{2^{*}-2}{2} \Phi_{\omega}^{\frac{4}{d-2}} \overline{\eta}$$
(4.11)
$$= O(|\eta|^{\min\{2,p\}}).$$

We see from (4.10) that the operator $-i\mathcal{L}_{\omega}$ relates to the behavior of the remainder $\eta(t)$. Unfortunately, $-i\mathcal{L}_{\omega}$ is not symmetric in $L^2_{real}(\mathbb{R}^d)$, and therefore we do not have the orthogonality of eigenfunctions in this space. Thus, we need to work in the symplectic space $(L^2(\mathbb{R}^d), \Omega)$ instead of $L^2_{real}(\mathbb{R}^2)$, where Ω is the symplectic form defined by

$$\Omega(f,g) := \Im \int_{\mathbb{R}^d} f(x)\overline{g(x)} \, dx = \left(f, ig\right)_{L^2_{real}}.$$
(4.12)

Proposition 4.1. Assume $d \geq 3$ and $1 + \frac{4}{d} . Let <math>\omega_1$ be the frequency given by Proposition 2.4. Then, there exists $\omega_2 \in (0, \omega_1)$ such that for any $\omega \in (0, \omega_2)$, $-i\mathcal{L}_{\omega}$ has a positive eigenvalue μ , as an operator in $L^2_{real}(\mathbb{R}^d)$. Furthermore, the eigenvalue μ satisfies that

$$-\mu^{2} = \inf \left\{ \frac{\langle L_{\omega,+}u, u \rangle_{H^{-1}, H^{1}}}{((L_{\omega,-})^{-1}u, u)_{L^{2}_{real}}} \colon u \in H^{1}(\mathbb{R}^{d}), \ (u, \Phi_{\omega})_{L^{2}_{real}} = 0 \right\}.$$
(4.13)

Proof of Proposition 4.1. We prove the proposition following the exposition in Section 2 of [9]. What we need to prove is that there exist a function $u \in H^1(\mathbb{R}^d)$ and $\nu < 0$ such that

$$L_{\omega,-}L_{\omega,+}u = \nu u. \tag{4.14}$$

Indeed, putting $f_1 = -\Re[u]$ and $f_2 = -\sqrt{-\nu}(L_{\omega,-})^{-1}\Re[u]$, we see from (4.6) and (4.14) that

$$-i\mathcal{L}_{\omega}(f_{1}+if_{2}) = -iL_{\omega,+}f_{1} + L_{\omega,-}f_{2}$$

$$= i(L_{\omega,-})^{-1}\nu\Re[u] - L_{\omega,-}\sqrt{-\nu}(L_{\omega,-})^{-1}\Re[u] = \sqrt{-\nu}(f_{1}+if_{2}).$$
(4.15)

Furthermore, we can verify that the problem (4.14) is equivalent to that there exist $u \in H^1(\mathbb{R}^d)$, $\nu < 0$ and $\alpha \in \mathbb{R}$ such that

$$\begin{cases} L_{\omega,+}u = \nu (L_{\omega,-})^{-1}u + \alpha \Phi_{\omega}, \\ (u, \Phi_{\omega})_{L^2_{real}} = 0. \end{cases}$$

$$(4.16)$$

Note here that Ker $L_{\omega,-} = \text{span} \{ \Phi_{\omega} \}$ (see Lemma B.1) and $-\Phi_{\omega} = L_{\omega,+} \partial_{\omega} \Phi_{\omega}$ (see (4.9)). The problem (4.16) leads us to the following minimizing problem

$$\nu_{\omega} := \inf \left\{ \frac{\langle L_{\omega,+}u, u \rangle_{H^{-1}, H^{1}}}{((L_{\omega,-})^{-1}u, u)_{L^{2}_{real}}} : u \in H^{1}(\mathbb{R}^{d}), \ (u, \Phi_{\omega})_{L^{2}_{real}} = 0 \right\}.$$
(4.17)

We can verify that any minimizer of (4.17) satisfies the equation (4.16) with $\nu = \nu_{\omega}$ for some $\alpha \in \mathbb{R}$. Thus, it suffices to show that $-\infty < \nu_{\omega} < 0$ and the existence of a minimizer.

First, we shall show that $-\infty < \nu_{\omega}$. We take any $u \in H^1(\mathbb{R}^d)$ with $(u, \Phi_{\omega})_{L^2_{real}} = 0$. Then, we see from Proposition 2.2 that

$$\frac{\langle L_{\omega,+}u,u\rangle_{H^{-1},H^{1}}}{((L_{\omega,-})^{-1}u,u)_{L_{real}^{2}}} = \frac{\langle L_{\omega,-}u,u\rangle_{H^{-1},H^{1}}}{\|(L_{\omega,-})^{-\frac{1}{2}}u\|_{L^{2}}^{2}} - \frac{\langle \{(p-1)\Phi_{\omega}^{p-1} + (2^{*}-2)\Phi_{\omega}^{\frac{4}{d-2}}\}u,u\rangle_{H^{-1},H^{1}}}{\|(L_{\omega,-})^{-\frac{1}{2}}u\|_{L^{2}}^{2}} = \frac{\|(L_{\omega,-})^{\frac{1}{2}}u\|_{L^{2}}^{2}}{\|(L_{\omega,-})^{-\frac{1}{2}}u\|_{L^{2}}^{2}} - C(\omega)\frac{\|u\|_{L^{2}}^{2}}{\|(L_{\omega,-})^{-\frac{1}{2}}u\|_{L^{2}}^{2}},$$
(4.18)

where $C(\omega)$ is some positive constant depending on ω . Moreover, it is easy to see that

$$\|u\|_{L^{2}}^{2} = ((L_{\omega,-})^{\frac{1}{2}}u, (L_{\omega,-})^{-\frac{1}{2}}u)_{L^{2}_{real}} \leq \|(L_{\omega,-})^{\frac{1}{2}}u\|_{L^{2}}\|(L_{\omega,-})^{-\frac{1}{2}}u\|_{L^{2}}$$

$$\leq \frac{1}{2C(\omega)}\|(L_{\omega,-})^{\frac{1}{2}}u\|_{L^{2}}^{2} + \frac{C(\omega)}{2}\|(L_{\omega,-})^{-\frac{1}{2}}u\|_{L^{2}}^{2}.$$
(4.19)

Putting the estimates (4.18) and (4.19) together, we find that $\nu_{\omega} > -C(\omega)^2 > -\infty$.

Next, we shall show that ν_{ω} is negative for any sufficiently small $\omega > 0$. To this end, we introduce projections Π_{ω} and Π :

$$\Pi_{\omega} u := u - \frac{(u, \Phi_{\omega})_{L^{2}_{real}}}{\|\Phi_{\omega}\|_{L^{2}}^{2}} \Phi_{\omega}, \quad \Pi u := u - \frac{(u, U)_{L^{2}_{real}}}{\|U\|_{L^{2}}^{2}} U.$$
(4.20)

Then, we see from substitution of variables, Proposition 2.3 and Lemma 2.7 that

$$\omega^{-2}\nu_{\omega} = \omega^{-2} \inf \left\{ \frac{\langle L_{\omega,+}u, u \rangle_{H^{-1},H^{1}}}{((L_{\omega,-})^{-1}u, u)_{L^{2}_{real}}} : u \in H^{1}(\mathbb{R}^{d}), (u, \Phi_{\omega})_{L^{2}_{real}} = 0 \right\}$$

$$\leq \omega^{-2} \frac{\langle L_{\omega,+}\Pi_{\omega}\partial_{\omega}\Phi_{\omega}, \Pi_{\omega}\partial_{\omega}\Phi_{\omega}\rangle_{H^{-1},H^{1}}}{((L_{\omega,-})^{-1}\Pi_{\omega}\partial_{\omega}\Phi_{\omega}, \Pi_{\omega}\partial_{\omega}\Phi_{\omega})_{L^{2}_{real}}} = \frac{\langle \widetilde{L}_{\omega,+}(\omega T_{\omega}\Pi_{\omega}\partial_{\omega}\Phi_{\omega}), \omega T_{\omega}\Pi_{\omega}\partial_{\omega}\Phi_{\omega}\rangle_{H^{-1},H^{1}}}{((\widetilde{L}_{\omega,-})^{-1}(\omega T_{\omega}\Pi_{\omega}\partial_{\omega}\Phi_{\omega}), \omega T_{\omega}\Pi_{\omega}\partial_{\omega}\Phi_{\omega})_{L^{2}_{real}}}$$

$$\rightarrow \frac{\langle L^{\dagger}_{+}\Pi\Lambda U, \Pi\Lambda U \rangle_{H^{-1},H^{1}}}{\langle (L^{\dagger}_{-})^{-1}\Pi\Lambda U, \Pi\Lambda U \rangle_{L^{2}_{real}}} \quad \text{as } \omega \downarrow 0,$$

$$(4.21)$$

where $\widetilde{L}_{\omega,+}$, $\widetilde{L}_{\omega,-}$, L^{\dagger}_{+} and L^{\dagger}_{-} are operators defined by (1.59), (1.60), (1.55) and (1.56), respectively. Thus, it suffices to show that

$$\frac{\langle L^{\dagger}_{+}\Pi\Lambda U,\Pi\Lambda U\rangle_{H^{-1},H^{1}}}{\langle (L^{\dagger})^{-1}_{-}\Pi\Lambda U,\Pi\Lambda U\rangle_{L^{2}_{real}}} < 0.$$

$$(4.22)$$

Note here that we see from (2.22) and integration by parts that

$$\Pi \Lambda U = \Lambda U + \frac{s_p}{2}U. \tag{4.23}$$

Moreover, it follows from (2.22) and (2.23) that

$$L_{+}^{\dagger}\Pi\Lambda U = L_{+}^{\dagger}\Lambda U - \frac{(\Lambda U, U)_{L_{real}^{2}}}{\|U\|_{L^{2}}^{2}}L_{+}^{\dagger}U = -U - \frac{(p-1)s_{p}}{2}U^{p}.$$
 (4.24)

Using (4.23), (4.24), (2.22) and integration by parts, we obtain that

$$\langle L_{+}^{\dagger}\Pi\Lambda U,\Pi\Lambda U\rangle_{H^{-1},H^{1}} = \langle -U - \frac{(p-1)s_{p}}{2}U^{p},\Lambda U + \frac{s_{p}}{2}U\rangle_{H^{-1},H^{1}}$$

$$= -(U,\Lambda U)_{L_{real}^{2}} - \frac{(p-1)s_{p}}{2}(U^{p},\Lambda U)_{L_{real}^{2}} - \frac{s_{p}}{2}\|U\|_{L^{2}}^{2} - \frac{(p-1)s_{p}^{2}}{4}\|U\|_{L^{p+1}}^{p+1}$$

$$= \frac{d(p-1)s_{p}}{4}\left(\frac{1}{2} - \frac{1}{p+1}\right)\|U\|_{L^{p}}^{p} < 0.$$

$$(4.25)$$

Since Lemma B.1 implies that $(\Pi \Lambda U, (L_{-}^{\dagger})^{-1} \Pi \Lambda U)_{L_{real}^{2}} > 0$, we find from (4.25) that (4.22) holds.

Finally, we shall show the existence of a minimizer u for ν_{ω} . We can take a sequence $\{u_n\}$ in $H^1(\mathbb{R}^d)$ such that

$$(u_n, \Phi_\omega)_{L^2_{real}} \equiv 0, \tag{4.26}$$

$$((L_{\omega,-})^{-1}u_n, u_n)_{L^2_{real}} \equiv 1,$$
 (4.27)

$$\lim_{n \to \infty} \langle L_{\omega,+} u_n, u_n \rangle_{H^{-1}, H^1} = \nu_{\omega}.$$
(4.28)

Note here that Lemma B.1 shows that the square root of $L_{\omega,-}$ is well-defined. We see from (4.28), Proposition 2.2, the Cauchy-Schwarz inequality, (B.1) and (4.27) that for any sufficiently large number n,

$$\begin{aligned} \|u_n\|_{H^1}^2 &\leq (1+\omega^{-1})\langle L_{\omega,+}u_n, u_n\rangle_{H^{-1},H^1} \\ &+ \int_{\mathbb{R}^d} \left\{ p(\Phi_{\omega}(x))^{p-1} + (2^*-1)(\Phi_{\omega}(x))^{\frac{4}{d-2}} \right\} |u_n(x)|^2 \, dx \\ &\lesssim (1+\omega^{-1})\nu_{\omega} + C_1(\omega)((L_{\omega,-})^{\frac{1}{2}}u_n, (L_{\omega,-})^{-\frac{1}{2}}u_n)_{L^2_{real}} \\ &\leq (1+\omega^{-1})\nu_{\omega} + C_1(\omega)\|(L_{\omega,-})^{\frac{1}{2}}u_n\|_{L^2} \|(L_{\omega,-})^{-\frac{1}{2}}u_n\|_{L^2} \\ &= (1+\omega^{-1})\nu_{\omega} + C_1(\omega)(L_{\omega,-}u_n, u_n)_{L^2_{real}}^{\frac{1}{2}} ((L_{\omega,-})^{-1}u_n, u_n)_{L^2_{real}}^{\frac{1}{2}} \\ &\leq (1+\omega^{-1})\nu_{\omega} + C_2(\omega)\|u_n\|_{H^1}, \end{aligned}$$

$$(4.29)$$

where $C_1(\omega)$ and $C_2(\omega)$ are some constants depending only on d, p and ω . This implies that $\{u_n\}$ is bounded in $H^1(\mathbb{R}^d)$ and therefore we can extract a subsequence of $\{u_n\}$ (still denoted by the same symbol) and a function $u_{\infty} \in H^1(\mathbb{R}^d)$ such that

$$\lim_{n \to \infty} u_n = u_{\infty} \quad \text{weakly in } H^1(\mathbb{R}^d), \tag{4.30}$$

$$(u_{\infty}, \Phi_{\omega})_{L^2_{real}} = 0, \tag{4.31}$$

$$\lim_{n \to \infty} \int_{\mathbb{R}^d} \Phi_{\omega}(x)^q |u_n(x)|^2 \, dx = \int_{\mathbb{R}^d} \Phi_{\omega}(x)^q |u_{\infty}(x)|^2 \, dx, \tag{4.32}$$

where q indicates p - 1 or $\frac{4}{d-2}$: Here, (4.31) follows from (4.26) and (4.30), and (4.32) follows from (4.30) and Propositionre 2.2. Furthermore, we can verify that

$$\langle L_{\omega,+}u_{\infty}, u_{\infty} \rangle_{H^{-1},H^{1}} \leq \liminf_{n \to \infty} \langle L_{\omega,+}u_{n}, u_{n} \rangle_{H^{-1},H^{1}} = \nu_{\omega} < 0.$$
(4.33)

In particular, the limit u_{∞} is non-trivial. We also see from Lemma B.1, (4.8), (4.26) and (4.27) that for any number n,

$$\|(L_{\omega,-})^{-1}u_n\|_{H^1}^2 \lesssim (u_n, (L_{\omega,-})^{-1}u_n)_{L^2_{real}} \le 1,$$
(4.34)

where the implicit constant depends on ω . Hence, there exists a subsequence of $\{u_n\}$ (still denoted by the same symbol) and a function $v_{\infty} \in H^1(\mathbb{R}^d)$ such that

$$\lim_{n \to \infty} (L_{\omega,-})^{-1} u_n = v_{\infty} \quad \text{weakly in } H^1(\mathbb{R}^d).$$
(4.35)

Furthermore, we see from (4.8) that for any test function $\phi \in C_c^{\infty}(\mathbb{R}^d)$,

$$\langle L_{\omega,-}v_{\infty},\phi\rangle_{H^{-1},H^{1}} = \langle L_{\omega,-}\phi,v_{\infty}\rangle_{H^{-1},H^{1}}$$

$$= \lim_{n \to \infty} ((L_{\omega,-})^{-1}u_{n},L_{\omega,-}\phi)_{L^{2}_{real}} = (u_{\infty},\phi)_{L^{2}_{real}}.$$

$$(4.36)$$

This together with (4.31) shows that

$$v_{\infty} = (L_{\omega,-})^{-1} u_{\infty}.$$
(4.37)

We also see from (4.37) and Lemma B.1 that

$$((L_{\omega,-})^{-1}u_{\infty}, u_{\infty})_{L^{2}_{real}} = (v_{\infty}, L_{\omega,-}v_{\infty})_{L^{2}_{real}} > 0.$$
(4.38)

Furthermore, it follows from (4.35), (4.37), (4.26) and (4.27) that

$$((L_{\omega,-})^{-1}u_{\infty}, u_{\infty})_{L^{2}_{real}} \le \liminf_{n \to \infty} ((L_{\omega,-})^{-1}u_{n}, u_{n})_{L^{2}_{real}} = 1,$$
(4.39)

so that we can take $\rho_1 \ge 1$ such that $((L_{\omega,-})^{-1}\rho_1 u_{\infty}, \rho_1 u_{\infty})_{L^2_{real}} = 1$. Suppose here that $\rho_1 > 1$. Then, we see from (4.31) and (4.33) that

$$\nu_{\omega} \le \langle L_{\omega,+}\rho_1 u_{\infty}, \rho_1 u_{\infty} \rangle_{H^{-1},H^1} < \langle L_{\omega,+} u_{\infty}, u_{\infty} \rangle_{H^{-1},H^1} \le \nu_{\omega}.$$

$$(4.40)$$

This is a contradiction. Thus, we have shown that

$$((L_{\omega,-})^{-1}u_{\infty}, u_{\infty})_{L^{2}_{real}} = 1.$$
(4.41)

Then, the same argument as (4.40) shows that

$$\nu_{\omega} = \langle L_{\omega,+} u_{\infty}, u_{\infty} \rangle_{H^{-1}, H^1}.$$
(4.42)

This together with (4.31) and (4.41) shows that u_{∞} is a minimizer for the problem (4.17).

4.2 Symplectic decomposition

In this subsection, we assume that $d \ge 3$, $1 + \frac{4}{d} and <math>0 < \omega < \omega_2$, where ω_2 is the frequency given in Proposition 4.1. Note that $-i\mathcal{L}_{\omega}$ has a positive eigenvalue $\mu > 0$ as an operator in $L^2_{real}(\mathbb{R}^d)$.

We will apply the "symplectic decomposition" corresponding to the discrete modes of $-i\mathcal{L}_{\omega}$ to the remainder η in (4.1) (see (4.48) below). Moreover, we determine the function $\theta(t)$ in (4.1).

Let \mathcal{U}_+ be an eigenfunction corresponding to the positive eigenvalue μ , and put

$$\mathcal{U}_{-} := \overline{\mathcal{U}_{+}}.\tag{4.43}$$

Then, we have

$$-i\mathcal{L}_{\omega}\mathcal{U}_{-} = \overline{i\mathcal{L}_{\omega}\mathcal{U}_{+}} = -\overline{\mu}\mathcal{U}_{+} = -\mu\mathcal{U}_{-}.$$
(4.44)

Hence, \mathcal{U}_{-} is an eigenfunction of $-i\mathcal{L}_{\omega}$ corresponding to $-\mu$. We assume that \mathcal{U}_{+} and \mathcal{U}_{-} are normalized in the following sense:

$$\Omega(\mathcal{U}_+, \mathcal{U}_-) = 1, \quad \Omega(\mathcal{U}_-, \mathcal{U}_+) = -1.$$
(4.45)

It is obvious that

$$\Omega(\mathcal{U}_+, \mathcal{U}_+) = \Omega(\mathcal{U}_-, \mathcal{U}_-) = 0.$$
(4.46)

Furthermore, it follows from (4.8), (4.9) and $\mathcal{L}_{\omega}\mathcal{U}_{\pm} = \pm i\mu\mathcal{U}_{\pm}$ that

$$\Omega(i\Phi_{\omega}, \mathfrak{U}_{\pm}) = \Omega(\partial_{\omega}\Phi_{\omega}, \mathfrak{U}_{\pm}) = 0.$$
(4.47)

Now, we expand the remainder $\eta(t)$ in the decomposition (4.1) by the discrete modes of $-i\mathcal{L}_{\omega}$:

$$\eta(t) = \lambda_{+}(t)\mathcal{U}_{+} + \lambda_{-}(t)\mathcal{U}_{-} + a(t)i\Phi_{\omega} + b(t)\partial_{\omega}\Phi_{\omega} + \gamma(t), \qquad (4.48)$$

where

$$\Omega(\gamma(t), \mathcal{U}_{+}) = \Omega(\gamma(t), \mathcal{U}_{-}) = \Omega(\gamma(t), i\Phi_{\omega}) = \Omega(\gamma(t), \partial_{\omega}\Phi_{\omega}) = 0.$$
(4.49)

We see from (4.45), (4.46) and (4.47) that the coefficients are as follows:

$$\lambda_{+}(t) = \Omega(\eta(t), \mathfrak{U}_{-}), \qquad \lambda_{-}(t) = -\Omega(\eta(t), \mathfrak{U}_{+}), \qquad (4.50)$$

$$a(t) = \frac{\Omega(\eta(t), \partial_{\omega} \Phi_{\omega})}{(\Phi_{\omega}, \partial_{\omega} \Phi_{\omega})_{L^{2}_{real}}}, \qquad b(t) = -\frac{\Omega(\eta(t), i\Phi_{\omega})}{(\Phi_{\omega}, \partial_{\omega} \Phi_{\omega})_{L^{2}_{real}}}.$$
(4.51)

Note here that the denominators in (4.51) are non-zero (see (iii) of Proposition 2.4). Moreover, it follows from Hölder's inequality that

$$\lambda_{\pm}^{2}(t) = \Omega(\eta(t), \mathfrak{U}_{\mp})^{2} \le \|\mathfrak{U}_{+}\|_{L^{2}}^{2} \|\eta(t)\|_{L^{2}}^{2}.$$
(4.52)

We require that the mode $i\Phi_{\omega}$ does not appear in the decomposition (4.48), that is, $a(t) \equiv 0$. To this end, we choose the function $\theta(t)$ in (4.1) so that

$$\Omega(e^{-i\theta(t)}\psi(t),\partial_{\omega}\Phi_{\omega}) \equiv 0.$$
(4.53)

Then, it follows from $\Omega(\Phi_{\omega}, \partial_{\omega}\Phi_{\omega}) = 0$ that $\Omega(\eta(t), \partial_{\omega}\Phi_{\omega}) \equiv 0$, and therefore $a(t) \equiv 0$. Furthermore, the choice of $\theta(t)$ has room of an integer multiple of π . Hence, in addition to (4.53), we can choose $\theta(t)$ so that

$$(e^{-i\theta(t)}\psi(t),\partial_{\omega}\Phi_{\omega})_{L^{2}_{real}} < 0.$$
(4.54)

This choice of $\theta(t)$ plays an important role in the argument below (see (6.10)).

4.3 Modulation equations

We continue the discussion about the decompositions (4.1) and (4.48) for a solution ψ to (NLS). Throughout this subsection, as well as Section 4.2, we assume that $d \geq 3$, $1 + \frac{4}{d} and <math>0 < \omega < \omega_2$. Furthermore, we assume that ψ satisfies (4.53), (4.54) and

$$\mathcal{M}(\psi) = \mathcal{M}(\Phi_{\omega}). \tag{4.55}$$

Since

$$\mathcal{M}(\psi) = \mathcal{M}(\Phi_{\omega}) + \mathcal{M}(\eta(t)) + \left(\Phi_{\omega}, \eta(t)\right)_{L^{2}_{real}},\tag{4.56}$$

the condition (4.55) implies that for any $t \in I_{\max}$,

$$\left(\Phi_{\omega},\eta(t)\right)_{L^2_{real}} = -\mathcal{M}(\eta(t)). \tag{4.57}$$

In particular, there is no orthogonality between Φ_{ω} and $\eta(t)$ in $L^2_{real}(\mathbb{R}^d)$.

Our aim here is to derive ordinary differential equations for θ , λ_{+} and λ_{-} under the condition (4.55).

First, we shall derive an equation for θ . We see from (4.53) that

$$0 = \frac{d}{dt}\Omega(\Phi_{\omega} + \eta(t), \partial_{\omega}\Phi_{\omega}) = \left(\frac{d\eta}{dt}(t), i\partial_{\omega}\Phi_{\omega}\right)_{L^{2}_{real}}.$$
(4.58)

Putting (4.10) and (4.58) together, we obtain

$$0 = \left(-\mathcal{L}_{\omega}\eta(t) - \left\{\frac{d\theta}{dt}(t) - \omega\right\} \left(\Phi_{\omega} + \eta(t)\right) + N_{\omega}(\eta(t)), \ \partial_{\omega}\Phi_{\omega}\right)_{L^{2}_{real}}$$
$$= -\left(\eta(t), \mathcal{L}_{\omega}\partial_{\omega}\Phi_{\omega}\right)_{L^{2}_{real}} - \left\{\frac{d\theta}{dt}(t) - \omega\right\} \left(\Phi_{\omega} + \eta(t), \partial_{\omega}\Phi_{\omega}\right)_{L^{2}_{real}}$$
$$+ \left(N_{\omega}(\eta(t)), \partial_{\omega}\Phi_{\omega}\right)_{L^{2}_{real}}.$$
(4.59)

Furthermore, using (4.9) and (4.57), we obtain the equation for θ :

$$\left\{\frac{d\theta}{dt}(t) - \omega\right\} \left(\Phi_{\omega} + \eta(t), \partial_{\omega}\Phi_{\omega}\right)_{L^{2}_{real}} = -\mathcal{M}(\eta(t)) + \left(N_{\omega}(\eta(t)), \partial_{\omega}\Phi_{\omega}\right)_{L^{2}_{real}}.$$
 (4.60)

Next, we shall derive equations for λ_+ and λ_- . It follows from (4.10), (4.47), (4.50) and $\mathcal{L}_{\omega}\mathcal{U}_{\pm} = \pm i\mu\mathcal{U}_{\pm}$ that

$$\frac{d\lambda_{\pm}}{dt}(t) = \pm \Omega(\frac{d\eta}{dt}(t), \mathfrak{U}_{\mp})$$

$$= \pm \Omega(-i\mathcal{L}_{\omega}\eta(t) - i\left\{\frac{d\theta}{dt}(t) - \omega\right\}\eta(t) + iN_{\omega}(\eta(t)), \mathfrak{U}_{\mp})$$

$$= \pm (\eta(t), i\mu\mathfrak{U}_{\mp})_{L^{2}_{real}} \mp \left(\left\{\frac{d\theta}{dt}(t) - \omega\right\}\eta(t) - N_{\omega}(\eta(t)), \mathfrak{U}_{\mp}\right)_{L^{2}_{real}}$$

$$= \pm \mu\lambda_{\pm}(t) \mp \left(\left\{\frac{d\theta}{dt}(t) - \omega\right\}\eta(t) - N_{\omega}(\eta(t)), \mathfrak{U}_{\mp}\right)_{L^{2}_{real}}.$$
(4.61)

Thus, we have obtained equations for λ_+ and λ_- :

$$\frac{d\lambda_{+}}{dt}(t) = \mu\lambda_{+}(t) - \left(\left\{\frac{d\theta}{dt}(t) - \omega\right\}\eta(t) - N_{\omega}(\eta(t)), \mathcal{U}_{-}\right)_{L^{2}_{real}},$$
(4.62)

$$\frac{d\lambda_{-}}{dt}(t) = -\mu\lambda_{-}(t) + \left(\left\{\frac{d\theta}{dt}(t) - \omega\right\}\eta(t) - N_{\omega}(\eta(t)), \,\mathcal{U}_{+}\right)_{L^{2}_{real}}.$$
(4.63)

Note here that it follows from $i\mathcal{L}_{\omega}\mathcal{U}_{\pm} = \pm \mu \mathcal{U}_{\pm}$, (4.62), (4.63) and (4.45) that

$$\langle \mathcal{L}_{\omega} \left(\lambda_{+}(t) \mathcal{U}_{+} + \lambda_{-}(t) \mathcal{U}_{-} \right), \frac{d\lambda_{+}}{dt}(t) \mathcal{U}_{+} + \frac{d\lambda_{-}}{dt}(t) \mathcal{U}_{-} \rangle_{H^{-1}, H^{1}}$$

$$= -\mu \lambda_{-}(t) \left(\left\{ \frac{d\theta}{dt}(t) - \omega \right\} \eta(t) - N_{\omega}(\eta(t)), \mathcal{U}_{-} \right)_{L^{2}_{real}}$$

$$+ \mu \lambda_{+}(t) \left(\left\{ \frac{d\theta}{dt}(t) - \omega \right\} \eta(t) - N_{\omega}(\eta(t)), \mathcal{U}_{+} \right)_{L^{2}_{real}}.$$

$$(4.64)$$

4.4 Linearized energy norm

Our aim here it to introduce the "linearized energy norm" for the remainder η in the decomposition (4.1). Throughout this subsection, as well as Section 4.3, we assume that $d \geq 3$, $1 + \frac{4}{d} , <math>0 < \omega < \omega_2$ and ψ is a solution to (NLS) satisfying (4.53) through (4.55).

Put

$$\Gamma(t) := b(t)\partial_{\omega}\Phi_{\omega} + \gamma(t). \tag{4.65}$$

Then, since $a(t) \equiv 0$ (see (4.53)), the decomposition (4.48) is rewritten by

$$\eta(t) = \lambda_{+}(t)\mathcal{U}_{+} + \lambda_{-}(t)\mathcal{U}_{-} + \Gamma(t).$$
(4.66)

We see from (4.47) and (4.49) that

$$\Omega(\Gamma(t), \mathcal{U}_{+}) = \Omega(\Gamma(t), \mathcal{U}_{-}) = \Omega(\Gamma(t), \partial_{\omega} \Phi_{\omega}) = 0.$$
(4.67)

As a consequence of these orthogonalities, we have the following relationship:

Lemma 4.2. The function Γ in the decomposition (4.66) satisfies

$$\langle \mathcal{L}_{\omega} \Gamma(t), \Gamma(t) \rangle_{H^{-1}, H^1} \sim \| \Gamma(t) \|_{H^1}^2$$
(4.68)

for all $t \in I_{max}$, where the implicit constant depends on ω as well as d and p.

Proof of Lemma 4.2. Lemma B.5 together with (4.67) gives the desired result.

Now, we see from (4.55), Taylor's expansion around Φ_{ω} , $S'_{\omega}(\Phi_{\omega}) = 0$, (4.3), $\mathcal{L}_{\omega}\mathcal{U}_{\pm} = \pm i\mu\mathcal{U}_{\pm}$ and (4.45) that

$$\begin{aligned} \mathcal{H}(\psi) - \mathcal{H}(\Phi_{\omega}) &= \mathcal{S}_{\omega}(\psi) - \mathcal{S}_{\omega}(\Phi_{\omega}) \\ &= \mathcal{S}_{\omega}(e^{-i\theta(t)}\psi) - \mathcal{S}_{\omega}(\Phi_{\omega}) = \mathcal{S}_{\omega}(\Phi_{\omega} + \eta(t)) - \mathcal{S}_{\omega}(\Phi_{\omega}) \\ &= \frac{1}{2} \langle \mathcal{L}_{\omega}\eta(t), \eta(t) \rangle_{H^{-1}, H^{1}} + O(\|\eta(t)\|_{H^{1}}^{\min\{3, p+1\}}) \\ &= \frac{1}{2} \lambda_{+}(t)\lambda_{-}(t)(-i\mu\mathcal{U}_{+}, \mathcal{U}_{-})_{L^{2}_{real}} + \frac{1}{2} \lambda_{+}(t)\lambda_{-}(t)(i\mu\mathcal{U}_{-}, \mathcal{U}_{+})_{L^{2}_{real}} \\ &+ \frac{1}{2} \langle \mathcal{L}_{\omega}\Gamma(t), \Gamma(t) \rangle_{H^{-1}, H^{1}} + O(\|\eta(t)\|_{H^{1}}^{\min\{3, p+1\}}) \\ &= -\mu\lambda_{+}(t)\lambda_{-}(t) + \frac{1}{2} \langle \mathcal{L}_{\omega}\Gamma(t), \Gamma(t) \rangle_{H^{-1}, H^{1}} + O(\|\eta(t)\|_{H^{1}}^{\min\{3, p+1\}}). \end{aligned}$$

$$(4.69)$$

We define the linearized energy norm $\|\eta(t)\|_E$ by

$$\|\eta(t)\|_{E}^{2} := \frac{\mu}{2} \left(\lambda_{+}^{2}(t) + \lambda_{-}^{2}(t)\right) + \frac{1}{2} \langle \mathcal{L}_{\omega} \Gamma(t), \Gamma(t) \rangle_{H^{-1}, H^{1}}.$$
(4.70)

Then, we see from (4.69) that

$$\mathcal{H}(\psi) - \mathcal{H}(\Phi_{\omega}) + \frac{\mu}{2} \left(\lambda_{+}(t) + \lambda_{-}(t) \right)^{2} - \|\eta(t)\|_{E}^{2} = O(\|\eta(t)\|_{H^{1}}^{\min\{3,p+1\}}).$$
(4.71)

Lemma 4.3. The function Γ in the decomposition (4.66) satisfies

$$\|\Gamma(t)\|_{H^1} \lesssim \|\eta(t)\|_{H^1} \tag{4.72}$$

for all $t \in I_{max}$. Moreover, we have

$$\|\eta(t)\|_{H^1} \sim \|\eta(t)\|_E. \tag{4.73}$$

Here, the implicit constants in (4.72) and (4.73) depend on ω as well as d and p.

Proof of Lemma 4.3. We see from (4.52) that

$$\|\Gamma(t)\|_{H^{1}}^{2} = \|\eta(t) - \lambda_{+}(t)\mathcal{U}_{+} - \lambda_{-}(t)\mathcal{U}_{-}\|_{H^{1}}^{2}$$

$$\lesssim \|\eta(t)\|_{H^{1}}^{2} + \lambda_{+}^{2}(t)\|\mathcal{U}_{+}\|_{H^{1}}^{2} + \lambda_{-}^{2}(t)\|\mathcal{U}_{-}\|_{H^{1}}^{2}$$

$$\lesssim (1 + 2\|\mathcal{U}_{+}\|_{H^{1}}^{4})\|\eta(t)\|_{H^{1}}^{2},$$
(4.74)

which proves (4.72).

Next, we shall prove (4.73). We see from (4.66), $\mathcal{U}_{-} = \overline{\mathcal{U}_{+}}$ (see (4.43)), Lemma 4.2 and the definition (4.70) that

$$\begin{aligned} \|\eta(t)\|_{H^{1}}^{2} &= \|\lambda_{+}(t)\mathcal{U}_{+} + \lambda_{-}(t)\mathcal{U}_{-} + \Gamma(t)\|_{H^{1}}^{2} \\ &\lesssim \lambda_{+}^{2}(t)\|\mathcal{U}_{+}\|_{H^{1}}^{2} + \lambda_{-}^{2}(t)\|\mathcal{U}_{-}\|_{H^{1}}^{2} + \|\Gamma(t)\|_{H^{1}}^{2} \\ &\lesssim \|\mathcal{U}_{+}\|_{H^{1}}^{2} \left(\lambda_{+}^{2}(t) + \lambda_{-}^{2}(t)\right) + \langle \mathcal{L}_{\omega}\Gamma(t), \Gamma(t) \rangle_{H^{-1}, H^{1}} \\ &\lesssim \max\{1, \mu^{-1}\}(1 + \|\mathcal{U}_{+}\|_{H^{1}}^{2})\|\eta(t)\|_{E}^{2}. \end{aligned}$$

$$(4.75)$$

Moreover, it follows from (4.52), Lemma 4.2 and (4.74) that

$$\begin{aligned} \|\eta(t)\|_{E}^{2} &\lesssim \mu \|\mathcal{U}_{+}\|_{L^{2}}^{2} \|\eta(t)\|_{L^{2}}^{2} + \|\Gamma(t)\|_{H^{1}}^{2} \\ &\lesssim \mu \|\mathcal{U}_{+}\|_{L^{2}}^{2} \|\eta(t)\|_{H^{1}}^{2} + \left(1 + 2\|\mathcal{U}_{+}\|_{H^{1}}^{4}\right) \|\eta(t)\|_{H^{1}}^{2}. \end{aligned}$$

$$(4.76)$$

Putting (4.75) and (4.76) together, we find that (4.73) holds.

4.5 Distance function from the ground state

Our aim here is to introduce a distance function from the ground state Φ_{ω} by using the linearized energy norm (4.70) (cf. [25]). Throughout this subsection, we assume that $d \geq 3$, $1 + \frac{4}{d} and <math>0 < \omega < \omega_2$, where ω_2 is the frequency given by Proposition 4.1.

We see from (4.71), and (4.73) in Lemma 4.3 that there exists a constant $\delta_E(\omega) > 0$ with the following property: for any solution ψ to (NLS) satisfying $\mathcal{M}(\psi) = \mathcal{M}(\Phi_{\omega})$ and any $t \in I_{\max}(\psi)$ for which $\|\eta(t)\|_E \leq 4\delta_E(\omega)$,

$$\left| \mathcal{H}(\psi(t)) - \mathcal{H}(\Phi_{\omega}) + \frac{\mu}{2} \left(\lambda_{+}(t) + \lambda_{-}(t) \right)^{2} - \|\eta(t)\|_{E}^{2} \right| \leq \frac{\|\eta(t)\|_{E}^{2}}{10}, \qquad (4.77)$$

where $\eta(t)$, $\lambda_+(t)$ and $\lambda_-(t)$ are the functions appearing in the decomposition for ψ of the form (4.1) with (4.66) (see also (4.48), (4.53) and (4.54)). Regarding the initial data of a solution to (NLS) as a general function in $H^1(\mathbb{R}^d)$, we find that the following fact holds:

Proposition 4.4. There exists a constant $\delta_E(\omega) > 0$ with the following property: let u be a function in $H^1(\mathbb{R}^d)$ satisfying $\mathcal{M}(u) = \mathcal{M}(\Phi_\omega)$. Consider the decomposition of the

form

$$u = e^{i\theta[u]}(\Phi_{\omega} + \eta[u]), \quad \Omega(e^{-i\theta[u]}u, \partial_{\omega}\Phi_{\omega}) \equiv 0, \quad (e^{-i\theta[u]}u, \partial_{\omega}\Phi_{\omega})_{L^{2}_{real}} < 0, \quad (4.78)$$

$$\eta[u] = \lambda_{+}[u]\mathcal{U}_{+} + \lambda_{-}[u]\mathcal{U}_{-} + \Gamma[u].$$
(4.79)

Define $\|\eta[u]\|_E$ by

$$\|\eta[u]\|_{E}^{2} := \frac{\mu}{2} \left(\lambda_{+}^{2}[u] + \lambda_{-}^{2}[u]\right) + \frac{1}{2} \langle \mathcal{L}_{\omega} \Gamma[u], \Gamma[u] \rangle_{H^{-1}, H^{1}}.$$
(4.80)

Furthermore, assume that

$$\|\eta[u]\|_E \le 4\delta_E(\omega). \tag{4.81}$$

Then,

$$\left| \mathcal{H}(u) - \mathcal{H}(\Phi_{\omega}) + \frac{\mu}{2} \left(\lambda_{+}[u] + \lambda_{-}[u] \right)^{2} - \|\eta[u]\|_{E}^{2} \right| \leq \frac{\|\eta[u]\|_{E}^{2}}{10}.$$
(4.82)

Now, we introduce a distance function d_{ω} . To this end, fix a non-increasing smooth function χ on $[0, \infty)$ such that

$$\chi(r) = \begin{cases} 1 & \text{if } 0 \le r \le 1, \\ 0 & \text{if } r \ge 2. \end{cases}$$
(4.83)

Moreover, we define

$$H^1_{\omega}(\mathbb{R}^d) := \left\{ u \in H^1(\mathbb{R}^d) \colon \mathcal{M}(u) = \mathcal{M}(\Phi_{\omega}) \right\}.$$
(4.84)

Then, we define a function $d_\omega \colon H^1_\omega(\mathbb{R}^d) \to [0,\infty)$ by

$$d_{\omega}(u)^{2} := \|\eta[u]\|_{E}^{2} + \chi \left(\frac{\|\eta[u]\|_{E}}{2\delta_{E}(\omega)}\right) C_{\omega}(u), \qquad (4.85)$$

where $\delta_E(\omega)$ is the constant given by Proposition 4.4, and

$$C_{\omega}(u) := \mathcal{H}(u) - \mathcal{H}(\Phi_{\omega}) + \frac{\mu}{2} (\lambda_{+}[u] + \lambda_{-}[u])^{2} - \|\eta[u]\|_{E}^{2}.$$
(4.86)

We rephrase (4.82) as follows: if $\|\eta[u]\|_E \leq 4\delta_E(\omega)$, then

$$|C_{\omega}(u)| \le \frac{\|\eta[u]\|_E^2}{10}.$$
(4.87)

Now, we consider a solution ψ to (NLS) satisfying $\mathcal{M}(\psi) = \mathcal{M}(\Phi_{\omega})$. In the decomposition (4.1) with (4.66), it is convenient to introduce new parameters $\lambda_1(t)$ and $\lambda_2(t)$ defined by

$$\lambda_1(t) := \frac{\lambda_+(t) + \lambda_-(t)}{2}, \qquad \lambda_2(t) := \frac{\lambda_+(t) - \lambda_-(t)}{2}. \tag{4.88}$$

We see from (4.62) and (4.63) that

$$\frac{d\lambda_1}{dt}(t) = \mu\lambda_2(t) + \frac{1}{2}\left(\left\{\frac{d\theta}{dt}(t) - \omega\right\}\eta(t) - N_\omega(\eta(t)), \ \mathcal{U}_+ - \mathcal{U}_-\right)_{L^2_{real}},\tag{4.89}$$

$$\frac{d\lambda_2}{dt}(t) = \mu\lambda_1(t) - \frac{1}{2}\left(\left\{\frac{d\theta}{dt}(t) - \omega\right\}\eta(t) - N_\omega(\eta(t)), \,\mathcal{U}_+ + \mathcal{U}_-\right)_{L^2_{real}}.$$
(4.90)

An important property of the distance function $d_{\omega}(\psi(t))$ is the following:

Lemma 4.5. Assume that there exists an interval I on which

$$\sup_{t \in I} d_{\omega}(\psi(t)) \le \delta_E(\omega). \tag{4.91}$$

Then, all of the following hold for all $t \in I$:

$$\frac{1}{2} \|\eta(t)\|_E^2 \le d_\omega(\psi(t))^2 \le \frac{3}{2} \|\eta(t)\|_E^2, \tag{4.92}$$

$$d_{\omega}(\psi(t))^2 = \mathcal{H}(\psi) - \mathcal{H}(\Phi_{\omega}) + 2\mu\lambda_1^2(t), \qquad (4.93)$$

$$\frac{d}{dt}d_{\omega}(\psi(t))^2 = 4\mu^2\lambda_1(t)\lambda_2(t) + 4\mu\lambda_1(t)\Omega\left(\left\{\frac{d\theta}{dt}(t) - \omega\right\}\eta(t) - N_{\omega}(\eta(t)), f_2\right).$$
(4.94)

Furthermore, if

$$\mathcal{S}_{\omega}(\psi) < m_{\omega} + \frac{1}{2} d_{\omega}(\psi(t))^2 \tag{4.95}$$

holds for all $t \in I$, then

$$d_{\omega}(\psi(t)) \sim |\lambda_1(t)| \tag{4.96}$$

for all $t \in I$, where the implicit constant depends on ω as well as d and p.

Proof of Lemma 4.5. First, we shall show that for any $t \in I$,

$$\|\eta(t)\|_E \le 4\delta_E(\omega). \tag{4.97}$$

Suppose for contradiction that $\|\eta(t_0)\|_E > 4\delta_E(\omega)$ for some $t_0 \in I$. Then, it follows from the definition (4.85) that $d_{\omega}(\psi(t_0)) = \|\eta(t_0)\|_E \ge 4\delta_E(\omega)$. However, this contradicts the assumption (4.91). Thus, we have proved (4.97). Using (4.87) and (4.97), we can verify (4.92). Indeed,

$$\frac{1}{2} \|\eta(t)\|_{E}^{2} \leq \|\eta(t)\|_{E}^{2} - \left|C_{\omega}(\psi(t))\right| \\
\leq d_{\omega}(\psi(t))^{2} \leq \|\eta(t)\|_{E}^{2} + \left|C_{\omega}(\psi(t))\right| \leq \frac{3}{2} \|\eta(t)\|_{E}^{2}.$$
(4.98)

Next, we shall derive the equation (4.93). We see from (4.91) and (4.92) that

$$\|\eta(t)\|_{E}^{2} \leq 2d_{\omega}(\psi(t))^{2} \leq 2\delta_{E}(\omega)^{2}.$$
(4.99)

Furthermore, it follows from the definition of $d_{\omega}(\psi(t))$ (see (4.85) and (4.86)) that

$$d_{\omega}(\psi(t))^{2} = \|\eta(t)\|_{E}^{2} + C_{\omega}(\psi(t)) = \mathcal{H}(\psi) - \mathcal{H}(\Phi_{\omega}) + \frac{\mu}{2} (\lambda_{+}(t) + \lambda_{-}(t))^{2}, \quad (4.100)$$

so that (4.93) holds.

The equation (4.94) follows from (4.93) and (4.89).

Finally, we shall prove (4.96). We see from the definition of $C_{\omega}(\psi(t))$ (see (4.86)), (4.97) and (4.87) that

$$\begin{aligned} \|\eta(t)\|_{E}^{2} &= \mathcal{H}(\psi) - \mathcal{H}(\Phi_{\omega}) + \frac{\mu}{2} \big(\lambda_{+}(t) + \lambda_{-}(t)\big)^{2} - C_{\omega}(\psi(t)) \\ &\leq \mathcal{H}(\psi) - \mathcal{H}(\Phi_{\omega}) + \frac{\mu}{2} \big(\lambda_{+}(t) + \lambda_{-}(t)\big)^{2} + \frac{\|\eta(t)\|_{E}^{2}}{10}, \end{aligned}$$

$$(4.101)$$

so that

$$\frac{9}{10} \|\eta(t)\|_{E}^{2} \leq \mathcal{H}(\psi) - \mathcal{H}(\Phi_{\omega}) + 2\mu\lambda_{1}(t)^{2}.$$
(4.102)

Moreover, it follows from the assumptions (4.55) and (4.95), and $m_{\omega} = \mathcal{S}_{\omega}(\Phi_{\omega})$ that

$$\mathcal{H}(\psi) - \mathcal{H}(\Phi_{\omega}) = \mathcal{S}_{\omega}(\psi) - \mathcal{S}_{\omega}(\Phi_{\omega}) < \frac{1}{2}d_{\omega}(\psi(t))^{2}.$$
(4.103)

Putting (4.92), (4.102) and (4.103) together, we obtain that

$$d_{\omega}(\psi(t))^{2} \leq \frac{3}{2} \|\eta(t)\|_{E}^{2} < \frac{5}{6} d_{\omega}(\psi(t))^{2} + \frac{10}{3} \mu \lambda_{1}(t)^{2}.$$
(4.104)

Hence, we have

$$d_{\omega}(\psi(t))^2 < 200\mu\lambda_1(t)^2.$$
(4.105)

On the other hand, we see from (4.52), Lemma 4.3 and (4.92) that

$$\mu\lambda_1(t)^2 \le 2\mu \left(\lambda_+^2(t) + \lambda_-^2(t)\right) \le 4\mu \|\mathcal{U}_+\|_{L^2}^2 \|\eta(t)\|_{L^2}^2 \lesssim d_\omega(\psi(t))^2, \tag{4.106}$$

where the implicit constant depends on μ and $\|\mathcal{U}_+\|_{L^2}^2$. Combining (4.105) and (4.106), we obtain (4.96).

4.6 Fundamental properties of the eigenfunctions

In connection with λ_1 and λ_2 (see (4.88)), we introduce the real-valued functions f_1 and f_2 :

$$f_1 := \frac{\mathcal{U}_+ + \mathcal{U}_-}{2} = \Re[\mathcal{U}_+], \qquad f_2 := \frac{\mathcal{U}_+ - \mathcal{U}_-}{2i} = \Im[\mathcal{U}_+].$$
(4.107)

We shall observe the properties of f_1 and f_2 . Throughout this subsection, we assume that $d \ge 3$, $1 + \frac{4}{d} , <math>0 < \omega < \omega_2$ and (4.55), where ω_2 is the constant given by Proposition 4.1.

First, we note that the decomposition (4.66) of $\eta(t)$ is expressed as follows in terms of the functions f_1 and f_2 defined by (4.107):

$$\eta(t) = 2\lambda_1(t)f_1 + 2i\lambda_2(t)f_2 + \Gamma(t).$$
(4.108)

We see from (4.6) and $\mathcal{L}_{\omega}\mathcal{U}_{+} = i\mu\mathcal{U}_{+}$ that

$$L_{\omega,+}f_1 = -\mu f_2, \qquad L_{\omega,-}f_2 = \mu f_1.$$
 (4.109)

Furthermore, it follows from the elliptic regularity for (4.109) that

$$\|f_1\|_{L^{\infty}} + \|f_2\|_{L^{\infty}} < \infty.$$
(4.110)

Lemma 4.6. We have that $f_2 \notin \text{span} \{\Phi_\omega\} = \text{Ker} L_{\omega,-}$.

Proof of Lemma 4.6. Suppose for a contradiction that $f_2 = k\Phi_{\omega}$ for some $k \neq 0$. Then, it follows from (4.8) and (4.107) that

$$0 = k\mathcal{L}_{\omega}(i\Phi_{\omega}) = \mathcal{L}_{\omega}(if_2) = \frac{1}{2}\mathcal{L}_{\omega}(\mathcal{U}_+ - \mathcal{U}_-) = \frac{1}{2}i\mu\mathcal{U}_+ + \frac{1}{2}i\mu\mathcal{U}_- = i\mu f_1.$$
(4.111)

Furthermore, we see from (4.109) and (4.111) that

$$-\mu f_2 = L_{\omega,+} f_1 = 0. \tag{4.112}$$

Thus, $f_1 = f_2 \equiv 0$. This is a contradiction.

Lemma 4.7. We have the following orthogonalities:

$$\left(\Phi_{\omega}, f_1\right)_{L^2} = \left(\partial_{\omega}\Phi_{\omega}, f_2\right)_{L^2} = 0. \tag{4.113}$$

Furthermore, we have

$$(f_1, f_2)_{L^2} > 0 \tag{4.114}$$

and

$$\left(\Phi_{\omega}, f_2\right)_{L^2} \neq 0. \tag{4.115}$$

Proof of Lemma 4.7. Since $L_{\omega,-}\Phi_{\omega} = 0$ and $L_{\omega,-}$ is self-adjoint in $L^2(\mathbb{R}^d)$, we find from (4.109) that

$$\left(\Phi_{\omega}, f_{1}\right)_{L^{2}} = \mu^{-1} \langle L_{\omega, -} f_{2}, \Phi_{\omega} \rangle_{H^{-1}, H^{1}} = \mu^{-1} \left(L_{\omega, -} \Phi_{\omega}, f_{2}\right)_{L^{2}} = 0.$$
(4.116)

On the other hand, it follows from (4.109), (4.9) and the self-adjointness of $L_{\omega,+}$ in $L^2(\mathbb{R}^d)$ that

$$\left(\partial_{\omega}\Phi_{\omega}, f_{2}\right)_{L^{2}} = -\mu^{-1} \langle L_{\omega,+}f_{1}, \partial_{\omega}\Phi_{\omega} \rangle_{H^{-1},H^{1}} = -\mu^{-1} \left(L_{\omega,+}\partial_{\omega}\Phi_{\omega}, f_{1}\right)_{L^{2}} = \mu^{-1} \left(\Phi_{\omega}, f_{1}\right)_{L^{2}},$$

$$(4.117)$$

which together with (4.116) proves (4.113).

Next, we shall prove (4.114). It follows from (4.109) that

$$(f_1, f_2)_{L^2} = \mu^{-1} \langle L_{\omega, -} f_2, f_2 \rangle_{H^{-1}, H^1}.$$
(4.118)

Since $f_2 \notin \text{Ker } L_{\omega,-}$ (see Lemma 4.6), Lemma B.1 together with (4.118) shows the desired result.

Finally, we prove (4.115). Suppose for contradiction that $(\Phi_{\omega}, f_2)_{L^2} = 0$. Then, we see from (4.7), the self-adjointness of $L_{\omega,+}$ and (4.109) that

$$\left((p-1)\Phi_{\omega}^{p} + (2^{*}-2)\Phi_{\omega}^{2^{*}-1}, f_{1}\right)_{L^{2}} = -\left(L_{\omega,+}\Phi_{\omega}, f_{1}\right)_{L^{2}} = \mu\left(\Phi_{\omega}, f_{2}\right)_{L^{2}} = 0.$$
(4.119)

Hence, it follows from Lemma B.3 that $\langle L_{\omega,+}f_1, f_1 \rangle_{H^{-1},H^1} \geq 0$. However, it must follow from (4.109) and (4.114) that

$$\langle L_{\omega,+}f_1, f_1 \rangle_{H^{-1},H^1} = -\mu (f_2, f_1)_{L^2} < 0.$$
 (4.120)

This is a contradiction. Thus, we find that (4.115) holds.

The relation (4.115) in Lemma 4.7 allows us to choose f_2 so that

$$(\Phi_{\omega}, f_2)_{L^2} < 0. \tag{4.121}$$

In our analysis, the frequency ω varies. In particular, we need to take $\omega \to 0$. Hence, we have to pay attention to the dependence of ω . Such a bother does not appear in the scale invariant cases such as (1.18) (see [25]). The following lemma plays an important role to prove the ejection lemma (Lemma 5.2):

Lemma 4.8. For any constant C > 0, there exists $\omega(C) > 0$ such that for any $\omega \in (0, \omega(C))$,

$$\mu |(\Phi_{\omega}, f_2)_{L^2}| \ge C |(\Phi_{\omega}^{2^*-1}, f_1)_{L^2}|.$$
(4.122)

Proof of Lemma 4.8. We use the notation μ_{ω} , $f_{1,\omega}$ and $f_{2,\omega}$ instead of μ , f_1 and f_2 in order to emphasise the dependence on ω .

Suppose for contradiction that there exists a constant $C_0 > 0$ with the following property: for any number n, there exists $\omega_n \in (0, \frac{1}{n})$ such that

$$\mu_{\omega_n} \left| (\Phi_{\omega_n}, f_{2,\omega_n})_{L^2} \right| < C_0 \left| (\Phi_{\omega_n}^{2^*-1}, f_{1,\omega_n})_{L^2} \right|, \tag{4.123}$$

where f_{1,ω_n} and f_{2,ω_n} are functions given by (4.107) for ω_n . We consider the functions $f_{1,n}$ and $f_{2,n}$ defined by

$$f_{1,n} := \omega_n^{-s_p} T_{\omega_n} f_{1,\omega_n}, \qquad f_{2,n} := \omega_n^{-s_p} T_{\omega_n} f_{2,\omega_n}.$$
(4.124)

It follows from (4.109) that

$$\widetilde{L}_{\omega_n,+}f_{1,n} = -\mu_{\omega_n}\omega_n^{-1}f_{2,n}, \quad \widetilde{L}_{\omega_n,-}f_{2,n} = \mu_{\omega_n}\omega_n^{-1}f_{1,n}, \quad (4.125)$$

where $\widetilde{L}_{\omega,+}$ and $\widetilde{L}_{\omega,-}$ are the operators defined by (1.59) and (1.60), respectively. We put

$$g_n := \frac{f_{1,n}}{\|f_{1,n}\|_{H^1}}, \quad h_n := \frac{f_{2,n}}{\|f_{1,n}\|_{H^1}}.$$
(4.126)

Then, we can take a real-valued function g in $H^1(\mathbb{R}^d)$ such that

$$\lim_{n \to \infty} g_n = g \qquad \text{weakly in } H^1(\mathbb{R}^d). \tag{4.127}$$

We shall show that g is non-trivial. Suppose for contradiction that g was trivial. Then, it follows from Lemma 4.7, (4.125), Proposition 2.3 and (4.127) that

$$0 \ge \lim_{n \to \infty} \frac{-\omega_n^{-s_p - 1} \mu_{\omega_n}}{\|f_{1,n}\|_{H^1}^2} (f_{1,\omega_n}, f_{2,\omega_n})_{L^2} = \lim_{n \to \infty} \frac{-\omega_n^{-1} \mu_{\omega_n}}{\|f_{1,n}\|_{H^1}^2} (f_{1,n}, f_{2,n})_{L^2}$$

$$= \lim_{n \to \infty} \frac{1}{\|f_{1,n}\|_{H^1}^2} \langle \widetilde{L}_{\omega_n, +} f_{1,n}, f_{1,n} \rangle_{H^{-1}, H^1}$$

$$= 1 - \lim_{n \to \infty} \int_{\mathbb{R}^d} \left\{ p(T_{\omega_n} \Phi_{\omega_n})^{p-1} |g_n|^2 + \omega_n^{\frac{2^* - (p+1)}{p-1}} (2^* - 1)(T_{\omega_n} \Phi_{\omega_n})^{\frac{4}{d-2}} |g_n|^2 \right\} dx$$

$$= 1.$$
(4.128)

This is a contradiction, and therefore g is non-trivial.

Next, we shall show that there exists a constant $C_0 > 0$ such that for any number nand any real-valued function $f \in H^1(\mathbb{R}^d)$ with $(f, T_{\omega_n} \Phi_{\omega_n})_{L^2} = 0$,

$$\langle \widetilde{L}_{\omega_n,-}f, f \rangle_{H^{-1},H^1} \ge C_0 \|f\|_{H^1}^2.$$
 (4.129)

Suppose for contradiction that we could take a subsequence of $\{\omega_n\}$ (still denoted by the same symbol) and a sequence $\{f_n\}$ in $H^1(\mathbb{R}^d)$ such that

$$\lim_{n \to \infty} \langle \widetilde{L}_{\omega_n, -} f_n, f_n \rangle_{H^{-1}, H^1} = 0, \qquad \|f_n\|_{H^1} = 1, \qquad (f_n, T_{\omega_n} \Phi_{\omega_n})_{L^2} = 0.$$
(4.130)

Furthermore, we can take $f_0 \in H^1(\mathbb{R}^d)$ such that $\lim_{n\to\infty} f_n = f_0$ weakly in $H^1(\mathbb{R}^d)$. Then, we see from the weak lower semicontinuity and Proposition 2.3 that

$$0 = \lim_{n \to \infty} \langle \widetilde{L}_{\omega_n, -} f_n, f_n \rangle_{H^{-1}, H^1}$$

=
$$\lim_{n \to \infty} \left\{ \|f_n\|_{H^1}^2 - \int (T_{\omega_n} \Phi_{\omega_n})^{p-1} |f_n|^2 dx - \omega_n^{\frac{2^* - (p+1)}{p-1}} \int (T_{\omega_n} \Phi_{\omega_n})^{\frac{4}{d-2}} |f_n|^2 dx \right\}$$

$$\geq \langle L_{-}^{\dagger} f_0, f_0 \rangle_{H^{-1}, H^1}.$$

(4.131)

Moreover, it follows from Proposition 2.3 and the hypothesis (4.130) that

$$(f_0, U)_{L^2} = \lim_{n \to \infty} (f_n, T_{\omega_n} \Phi_{\omega_n})_{L^2} = 0.$$
 (4.132)

Hence, we conclude from (4.131) and the positivity of L_{-}^{\dagger} together with (4.132) that

$$0 \ge \langle L_{-}^{\dagger} f_{0}, f_{0} \rangle \gtrsim \| f_{0} \|_{H^{1}}^{2}, \qquad (4.133)$$

so that f_0 is trivial. However, the same argument as (4.128) yields that f_0 is non-trivial. Thus, we arrive at a contradiction and therefore (4.129) holds.

We shall show that $\{h_n\}$ is bounded in $H^1(\mathbb{R}^d)$. We see from (4.129) and (4.125) that

$$C_{0} \| f_{2,n} \|_{H^{1}}^{2} \leq \langle \widetilde{L}_{\omega_{n},-} f_{2,n}, f_{2,n} \rangle_{H^{-1},H^{1}} = \mu_{\omega_{n}} \omega_{n}^{-1} (f_{1,n}, f_{2,n})_{L^{2}}$$

$$= -\langle \widetilde{L}_{\omega_{n},+} f_{1,n}, f_{1,n} \rangle_{H^{-1},H^{1}} \leq C \| f_{1,n} \|_{H^{1}}^{2}.$$
(4.134)

Dividing the both sides above by $||f_{1,n}||^2_{H^1}$, we find that $\{h_n\}$ is bounded in $H^1(\mathbb{R}^d)$.

We shall show that the sequence $\{\mu_{\omega_n}\omega_n^{-1}\}$ is bounded. It follows from (4.125) that

$$|\mu_{\omega_n}\omega_n^{-1}| = \frac{|\langle \widetilde{L}_{\omega_n,-}f_{2,n}, f_{1,n} \rangle_{H^{-1},H^1}|}{\|f_{1,n}\|_{L^2}^2}.$$
(4.135)

Furthermore, it follows from (4.127) and the boundedness of $\{g_n\}$ and $\{h_n\}$ in $H^1(\mathbb{R}^d)$ that

$$\left|\mu_{\omega_{n}}\omega_{n}^{-1}\right| \leq \frac{\sup_{n\in\mathbb{N}}\left|\langle \tilde{L}_{\omega_{n},-}h_{n},g_{n}\rangle_{H^{-1},H^{1}}\right|}{\lim\inf_{n\to\infty}\|g_{n}\|_{L^{2}}^{2}} \lesssim \frac{\sup_{n\in\mathbb{N}}\left(\|g_{n}\|_{H^{1}}^{2}+\|h_{n}\|_{H^{1}}^{2}\right)}{\|g\|_{L^{2}}^{2}} \lesssim 1.$$
(4.136)

Thus, we find that $\{\mu_{\omega_n}\omega_n^{-1}\}$ is bounded.

Since $\{h_n\}$ is bounded in $H^1(\mathbb{R}^d)$, there exists $h \in H^1(\mathbb{R}^d)$ such that

$$\lim_{n \to \infty} h_n = h \qquad \text{weakly in } H^1(\mathbb{R}^d).$$
(4.137)

Moreover, we can take a subsequence of $\{\mu_{\omega_n}\omega_n^{-1}\}$ (still denoted by the same symbol) and $\nu_* \in [0,\infty)$ such that $\lim_{n\to\infty} \mu_{\omega_n}\omega_n^{-1} = \nu_*$. We shall show that $\nu_* \neq 0$ and h is non-trivial. Recall here that μ_{ω_n} is a positive eigenvalue of $-i\mathcal{L}_{\omega_n}$. We find from the proof of Proposition 4.1 (see (4.15) and (4.17)) that

$$-\mu_{\omega_n}^2 = \inf\left\{\frac{\langle L_{\omega_n,+}u, u \rangle_{H^{-1}, H^1}}{((L_{\omega_n,-})^{-1}u, u)_{L_{real}^2}} : u \in H^1(\mathbb{R}^d), (u, \Phi_{\omega_n})_{L_{real}^2} = 0\right\}.$$
 (4.138)

Furthermore, it follows from the estimates (4.21) and (4.22) in the proof of Proposition 4.1 shows that

$$-\nu_*^2 = \lim_{n \to \infty} \omega_n^{-2} (-\mu_{\omega_n}^2) = \frac{\langle L_+^{\dagger} \Pi \Lambda U, \Pi \Lambda U \rangle_{H^{-1}, H^1}}{((L_-^{\dagger})^{-1} \Pi \Lambda U, \Pi \Lambda U)_{L_{real}^2}} < 0, \qquad (4.139)$$

where Π is the projection given by (4.20). Thus, we find that $\nu_* \neq 0$. Next, suppose for contradiction that h was trivial. Then, we see from (4.125) and (4.127) that for any $\varphi \in C_0^{\infty}(\mathbb{R}^d)$,

$$0 = \lim_{n \to \infty} \langle \widetilde{L}_{\omega_n, -} h_n, \varphi \rangle_{H^{-1}, H^1} = \lim_{n \to \infty} (\mu_{\omega_n} \omega_n^{-1} g_n, \varphi)_{L^2_{real}} = (\nu_* g, \varphi)_{L^2_{real}}.$$
 (4.140)

However, this contradicts that $\nu_* \neq 0$ and g is non-trivial. Thus, we find that h is non-trivial.

Note that g and h satisfies that

$$L^{\dagger}_{+}g = -\nu_{*}h, \qquad L^{\dagger}_{-}h = \nu_{*}g.$$
 (4.141)

Then, applying the same proof of (4.115) (use Lemma 2.2 in [9] instead of Lemma B.3), we find that

$$\langle L^{\dagger}_{+}U,h\rangle_{H^{-1},H^{1}} = (p-1)(U^{p},h)_{L^{2}} \neq 0.$$
 (4.142)

We see from (4.109) and the scaling that

$$-\omega_n^{-1}\mu_{\omega_n}(\Phi_{\omega_n}, f_{2,\omega_n})_{L^2} = \langle \widetilde{L}_{\omega_n, +} T_{\omega_n} \Phi_{\omega_n}, f_{1,n} \rangle_{H^{-1}, H^1}.$$
(4.143)

Moreover, it follows from Proposition 2.3 that

$$\lim_{n \to \infty} \widetilde{L}_{\omega_n, +} T_{\omega_n} \Phi_{\omega_n} = L_+^{\dagger} U \qquad \text{strongly in } H^{-1}(\mathbb{R}^d).$$
(4.144)

This together with (4.137) yields

$$\lim_{n \to \infty} \langle \widetilde{L}_{\omega_n, +} T_{\omega_n} \Phi_{\omega_n}, h_n \rangle_{H^{-1}, H^1} = \langle L_+^{\dagger} U, h \rangle_{H^{-1}, H^1}.$$

$$(4.145)$$

We find from (4.143), (4.145) and (4.142) that

$$\lim_{n \to \infty} \|f_{1,n}\|_{H^1}^{-1} \omega_n^{-1} \mu_{\omega_n} |(\Phi_{\omega_n}, f_{2,\omega_n})_{L^2}| = \lim_{n \to \infty} |\langle \widetilde{L}_{\omega_n,+} T_{\omega_n} \Phi_{\omega_n}, h_n \rangle_{H^{-1},H^1}|
= |\langle L_+^{\dagger} U, h \rangle_{H^{-1},H^1}| \gtrsim 1,$$
(4.146)

On the other hand, it follows from Proposition 2.3 and (4.127) that

$$\lim_{n \to 0} \|f_{1,n}\|_{H^1}^{-1} \omega_n^{-\frac{2^*-2}{p-1}} \left| (\Phi_{\omega_n}^{2^*-1}, f_{1,\omega_n})_{L^2} \right| = \lim_{n \to 0} \left| ((T_{\omega_n} \Phi_{\omega_n})^{2^*-1}, g_n)_{L^2} \right|
= \left| (U^{2^*-1}, g)_{L^2} \right| \lesssim 1.$$
(4.147)

Since $\frac{2^*-2}{p-1} > 1$, we conclude from (4.146) and (4.147) that for any sufficiently large number n,

$$\mu_{\omega_n} \left| (\Phi_{\omega_n}, f_{2,,\omega_n})_{L^2} \right| \gtrsim \omega_n^{-\frac{2^*-2}{p-1}+1} \left| (\Phi_{\omega_n}^{2^*-1}, f_{1,,\omega_n})_{L^2} \right| \gg C_0 \left| (\Phi_{\omega_n}^{2^*-1}, f_{1,\omega_n})_{L^2} \right|, \quad (4.148)$$

where C_0 is the constant given in the hypothesis (4.123). However, this contradicts (4.123). Thus, the desired result (4.122) holds.

Now, we see from (4.47) and (4.57) that

$$(\Phi_{\omega}, \Gamma(t))_{L^2_{real}} = -\mathcal{M}(\eta(t)). \tag{4.149}$$

Moreover, we see from (4.7) and (4.8) that

$$2\omega\Phi_{\omega} - 2\Delta\Phi_{\omega} - \frac{d(p-1)}{2}\Phi_{\omega}^{p} - 2^{*}\Phi_{\omega}^{2^{*}-1}$$

$$= (2-s_{p})L_{\omega,-}\Phi_{\omega} + s_{p}L_{\omega,+}\Phi_{\omega} - (1-s_{p})(2^{*}-2)\Phi_{\omega}^{2^{*}-1}$$

$$= s_{p}L_{\omega,+}\Phi_{\omega} - (1-s_{p})(2^{*}-2)\Phi_{\omega}^{2^{*}-1} = -s_{p}(p-1)\Phi_{\omega}^{p} - (2^{*}-2)\Phi_{\omega}^{2^{*}-1}.$$
(4.150)

Furthermore, this together with (4.108), (4.109) and (4.149) shows that

$$\begin{aligned} \mathcal{K}'(\Phi_{\omega})\eta(t) \\ &= 2\langle\omega\Phi_{\omega} - \Delta\Phi_{\omega} - \frac{d(p-1)}{4}\Phi_{\omega}^{p} - \frac{2^{*}}{2}\Phi_{\omega}^{2^{*}-1}, \eta(t)\rangle_{H^{-1},H^{1}} - 2\omega(\Phi_{\omega},\eta(t))_{L^{2}_{real}} \\ &= 2s_{p}\lambda_{1}(t)(L_{\omega,+}\Phi_{\omega},f_{1})_{L^{2}} - 2(1-s_{p})(2^{*}-2)\lambda_{1}(t)(\Phi_{\omega}^{2^{*}-1},f_{1})_{L^{2}} \\ &- s_{p}((p-1)\Phi_{\omega}^{p} + (2^{*}-2)\Phi_{\omega}^{2^{*}-1},\Gamma(t))_{L^{2}_{real}} \\ &- (1-s_{p})(2^{*}-2)(\Phi_{\omega}^{2^{*}-1},\Gamma(t))_{L^{2}_{real}} + \omega\mathcal{M}(\eta(t)) \\ &= -2\mu s_{p}\lambda_{1}(t)(\Phi_{\omega},f_{2})_{L^{2}} - 2(1-s_{p})(2^{*}-2)\lambda_{1}(t)(\Phi_{\omega}^{2^{*}-1},f_{1})_{L^{2}} \\ &- s_{p}(p-1)(\Phi_{\omega}^{p},\Gamma(t))_{L^{2}_{real}} - (2^{*}-2)(\Phi_{\omega}^{2^{*}-1},\Gamma(t))_{L^{2}_{real}} + \omega\mathcal{M}(\eta(t)). \end{aligned}$$
5 Ejection lemma

Let ω_2 denote the frequency given by Proposition 4.1 throughout this section. Furthermore, for a given $\omega \in (0, \omega_2)$, $\delta_E(\omega)$ denotes the constant given by Proposition 4.4.

We see from the argument in the previous section that any solution ψ satisfying $\mathcal{M}(\psi) = \mathcal{M}(\Phi_{\omega})$ has the decomposition of the form (4.1) with (4.48), (4.53) and (4.54). Recall that $d_{\omega} \colon H^1_{\omega}(\mathbb{R}^d) \to [0, \infty)$ denotes the distance function defined by (4.85).

The following lemma is a criterion of continuation for the solutions to (NLS) in terms of the ground state:

Lemma 5.1. Assume $d \ge 3$ and $1 + \frac{4}{d} . Then, for any <math>\omega \in (0, \omega_2)$, there exists $\delta_0(\omega) \in (0, \delta_E(\omega))$ such that if ψ is a solution to (NLS) satisfying

$$\mathcal{M}(\psi) = \mathcal{M}(\Phi_{\omega}),\tag{5.1}$$

then ψ extends as long as $d_{\omega}(\psi(t)) \leq \delta_0(\omega)$.

Proof of Lemma 5.1. We prove the claim by contradiction. Hence, suppose to the contrary that there exists $\omega \in (0, \omega_2)$ with the following property: for any $\delta_0 \in (0, \delta_E(\omega))$, there exists a solution ψ to (NLS) such that: $\mathcal{M}(\psi) = \mathcal{M}(\Phi_\omega)$; $T_{\max} := \sup I_{\max}(\psi) < \infty$; and $\sup_{t \in [t_0, T_{\max})} d_{\omega}(\psi(t)) \leq \delta_0$ for some $t_0 \in I_{\max}$. Then, it follows from Lemma 4.3 and Lemma 4.5 that

$$\sup_{t \in [t_0, T_{\max})} \|\eta(t)\|_{H^1} \lesssim \sup_{t \in [t_0, T_{\max})} d_\omega(\psi(t)) < \delta_0,$$
(5.2)

where the implicit constant depends only on d, p and ω . In particular, we have

$$\sup_{t \in [t_0, T_{\max})} \|\psi(t)\|_{H^1} \lesssim \|\Phi_\omega\|_{H^1} + \delta_0.$$
(5.3)

Let $t_1 \in (t_0, T_{\text{max}})$. Then, Strichartz' estimate together with (5.2) shows that

$$\begin{aligned} \|\langle \nabla \rangle e^{i(t-t_1)\Delta} \psi(t_1)\|_{St([t_1,T_{\max}])} &\lesssim \|\langle \nabla \rangle e^{i(t-t_1)\Delta} \Phi_\omega\|_{St([t_1,T_{\max}])} + \|\eta(t_1)\|_{H^1} \\ &\lesssim \|\langle \nabla \rangle e^{i(t-t_1)\Delta} \Phi_\omega\|_{St([t_1,T_{\max}])} + \delta_0. \end{aligned}$$
(5.4)

Furthermore, the small-data theory (Lemma D.1) together with (5.3) shows that there exists $\delta(\omega) > 0$ such that if

$$\|\langle \nabla \rangle e^{i(t-t_1)\Delta} \psi(t_1)\|_{St([t_1,T_{\max}])} \le \delta(\omega), \tag{5.5}$$

then the solution ψ exists on $[t_1, T_{\text{max}}]$. Since

$$\lim_{t_1 \uparrow T_{\max}} \|\langle \nabla \rangle e^{i(t-t_1)\Delta} \Phi_\omega\|_{St([t_1, T_{\max}])} = 0,$$
(5.6)

we see from (5.4) that if $\delta_0 \ll \delta(\omega)$, then ψ extends beyond T_{max} . However, this is a contradiction. Thus, we have proved the lemma.

Note here that if follows from Lemma 4.8 that there exists a frequency $\omega(2^*) \in (0, \omega_2)$ such that for any $\omega \in (0, \omega(2^*))$,

$$\mu s_p |(\Phi_{\omega}, f_2)_{L^2}| \ge 2(1 - s_p)(2^* - 2) |(\Phi_{\omega}^{2^* - 1}, f_1)_{L^2}|.$$
(5.7)

We use this fact in the proof of the ejection lemma below.

Lemma 5.2 (Ejection lemma). Assume $d \ge 3$ and $1 + \frac{4}{d} . Then, for any <math>\omega \in (0, \omega(2^*))$, there exist constants $\delta_X \in (0, \delta_0(\omega))$ ($\delta_0(\omega)$ denotes the constant given by Lemma 5.1), $A_* > 0$, $B_* > 0$, $C_* > 0$ and $T_* > 0$ with the following properties: for any $t_0 \in \mathbb{R}$ and any solution ψ to (NLS) defined around t_0 satisfying

$$\mathcal{M}(\psi) = \mathcal{M}(\Phi_{\omega}),\tag{5.8}$$

$$0 < R_0 := d_\omega(\psi(t_0)) < \delta_X, \tag{5.9}$$

$$\mathcal{S}_{\omega}(\psi) < m_{\omega} + \frac{R_0^2}{2},\tag{5.10}$$

we can extend ψ as long as $d_{\omega}(\psi(t)) \leq \delta_X$. Furthermore, assume that there exists $T > t_0$ such that

$$R_0 \le \min_{t \in [t_0, T]} d_\omega(\psi(t)), \tag{5.11}$$

and define

$$T_X := \inf \{ t \in [t_0, T] : d_{\omega}(\psi(t)) = \delta_X \},$$
(5.12)

where we interpret $T_X = T$ if $d_{\omega}(\psi) < \delta_X$ on $[t_0, T]$. Then, for any $t \in [t_0, T_X]$,

$$A_* e^{\mu(t-t_0)} R_0 \le d_\omega(\psi(t)) \le B_* e^{\mu(t-t_0)} R_0, \tag{5.13}$$

$$\|\eta(t)\|_{H^1} \sim \mathfrak{s}\lambda_1(t) \sim \mathfrak{s}\lambda_+(t) \sim e^{\mu(t-t_0)}R_0, \qquad (5.14)$$

$$|\lambda_{-}(t)| + \|\Gamma(t)\|_{H^{1}} \lesssim R_{0} + \left(e^{\mu(t-t_{0})}R_{0}\right)^{\frac{\min\{3,p+1\}}{2}},\tag{5.15}$$

$$\mathfrak{s}\mathcal{K}(\psi(t)) \gtrsim \left(e^{\mu(t-t_0)} - C_*\right) R_0,\tag{5.16}$$

where \mathfrak{s} is either $\mathfrak{s} = 1$ or $\mathfrak{s} = -1$. Moreover, $d_{\omega}(\psi(t))$ is increasing on the region $\{t \in [t_0, T_X] : t_0 + T_* R_0^{\min\{1, p-1\}} \leq t\};$ and

$$\left| d_{\omega}(\psi(t)) - R_0 \right| \lesssim R_0^{\min\{2,p\}} \tag{5.17}$$

on the region $\{t \in [t_0, T_X] : t_0 \le t \le t_0 + T_* R_0^{\min\{1, p-1\}} \}.$

Proof of Lemma 5.2. Let $\delta_X \in (0, \delta_0(\omega))$ be a small constant to be chosen later, and let ψ be a solution satisfying (5.8), (5.9) and (5.10). Since the equation (NLS) is invariant under the time translations, it suffices to consider the case where $t_0 = 0$.

First, we find from Lemma 5.1 that ψ extends as long as $d_{\omega}(\psi(t)) \leq \delta_X$.

Next, we assume (5.11) as well as (5.8), (5.9) and (5.10). Then, it follows from the definition of T_X and $t_0 = 0$ that for any $t \in [0, T_X]$,

$$d_{\omega}(\psi(t)) \le \delta_X < \delta_E(\omega). \tag{5.18}$$

Hence, we see from Lemma 4.3 and Lemma 4.5 that for any $t \in [0, T_X]$,

$$\|\Gamma(t)\|_{H^1} \lesssim \|\eta(t)\|_{H^1} \sim \|\eta(t)\|_E \sim d_\omega(\psi(t)) \sim |\lambda_1(t)|,$$
(5.19)

$$\frac{d}{dt}d_{\omega}(\psi(t))^{2} = 4\mu^{2}\lambda_{1}(t)\lambda_{2}(t) + 4\mu\lambda_{1}(t)\Omega\left(\left\{\frac{d\theta}{dt}(t) - \omega\right\}\eta(t) - N_{\omega}(\eta(t)), f_{2}\right).$$
 (5.20)

In particular, we deduce from (4.52), (5.18) and (5.19) that for any $t \in [0, T_X]$,

$$|\lambda_{-}(t)| + \|\Gamma(t)\|_{H^{1}} \lesssim \|\eta(t)\|_{H^{1}} \sim |\lambda_{1}(t)| \lesssim \delta_{X} \ll 1.$$
(5.21)

Furthermore, we find from (5.11) and (5.19) that for any $t \in [0, T_X]$.

$$0 < R_0 \lesssim |\lambda_1(t)|, \tag{5.22}$$

which together with the continuity of $\lambda_1(t)$ shows that for any $t \in [0, T_X]$

$$\mathfrak{s} := \operatorname{sgn}[\lambda_1(t)] \equiv \operatorname{sgn}[\lambda_1(0)]. \tag{5.23}$$

Now, we further choose δ_X so small that

$$\delta_X \ll \frac{|(\Phi_\omega, \partial_\omega \Phi_\omega)_{L^2}|}{2\|\partial_\omega \Phi_\omega\|_{L^2}}.$$
(5.24)

Then, we find from (5.21) that for any $t \in [0, T_X]$,

$$\|\eta(t)\|_{L^{2}} \leq \|\eta(t)\|_{H^{1}} \ll \min\left\{1, \frac{|(\Phi_{\omega}, \partial_{\omega}\Phi_{\omega})_{L^{2}}|}{2\|\partial_{\omega}\Phi_{\omega}\|_{L^{2}}}\right\}.$$
(5.25)

Furthermore, (5.25) implies that for any $t \in [0, T_X]$,

$$\left| (\Phi_{\omega} + \eta(t), \partial_{\omega} \Phi_{\omega})_{L^2} \right| \ge \left| (\Phi_{\omega}, \partial_{\omega} \Phi_{\omega})_{L^2} \right| - \|\eta(t)\|_{L^2} \|\partial_{\omega} \Phi_{\omega}\|_{L^2} \ge \frac{1}{2} \left| (\Phi_{\omega}, \partial_{\omega} \Phi_{\omega})_{L^2} \right|.$$
(5.26)

We see from Hölder's inequality and $N_{\omega}(\eta) = O(|\eta|^{\min\{2,p\}})$ (see (4.11)) that for any function $u \in L^1(\mathbb{R}^d) \cap L^{\infty}(\mathbb{R}^d)$,

$$\left| (N_{\omega}(\eta(t)), u)_{L^{2}_{real}} \right| \leq \left\| N_{\omega}(\eta(t)) \right\|_{L^{\frac{p+1}{\min\{2,p\}}}} \left\| u \right\|_{L^{\frac{p+1}{\max\{1,p-1\}}}}$$

$$\lesssim \left\| \eta(t) \right\|_{L^{p+1}}^{\min\{2,p\}} \left\| u \right\|_{L^{1} \cap L^{\infty}}.$$
(5.27)

We deduce from the equation (4.60) for $\theta(t)$, (5.26), (5.27) and (5.25) that for any $t \in [0, T_X]$,

$$\left|\frac{d\theta}{dt}(t) - \omega\right| \lesssim \frac{\|\eta(t)\|_{L^2}^2 + \|\eta(t)\|_{L^{p+1}}^{\min\{2,p\}} \|\partial_\omega \Phi_\omega\|_{L^1 \cap L^\infty}}{|(\Phi_\omega, \partial_\omega \Phi_\omega)_{L^2}|} \lesssim \|\eta(t)\|_{H^1}^{\min\{2,p\}}, \tag{5.28}$$

where the implicit constant depends on ω . Furthermore, it follows from (5.28), (5.27) and (5.25) that for any $t \in [0, T_X]$,

$$\left| \Omega\left(\left\{ \frac{d\theta}{dt}(t) - \omega \right\} \eta(t) - N_{\omega}(\eta(t)), f_{2}\right) \right| \\
\leq \left| \frac{d\theta}{dt}(t) - \omega \right| \|\eta(t)\|_{L^{2}} \|f_{2}\|_{L^{2}} + \|\eta(t)\|_{L^{p+1}}^{\min\{2,p\}} \|f_{2}\|_{L^{2} \cap L^{\infty}} \\
\lesssim \|\eta(t)\|_{H^{1}}^{\min\{2,p\}+1} + \|\eta(t)\|_{H^{1}}^{\min\{2,p\}} \lesssim \|\eta(t)\|_{H^{1}}^{\min\{2,p\}},$$
(5.29)

where the implicit constant depends on ω . Hence, we find form (5.29) and (5.19) that for any $t \in [0, T_X]$,

$$\left|\Omega\left(\left\{\frac{d\theta}{dt}(t) - \omega\right\}\eta(t) - N_{\omega}(\eta(t)), f_2\right)\right| \lesssim \left|\lambda_1(t)\right|^{\min\{2, p\}},\tag{5.30}$$

where the implicit constant depends on ω . Similarly, we can verify that for any $t \in [0, T_X]$,

$$\left| \left(\left\{ \frac{d\theta}{dt}(t) - \omega \right\} \eta(t) - N_{\omega}(\eta(t)), \mathfrak{U}_{\pm} \right)_{L^{2}_{real}} \right| \lesssim \left| \lambda_{1}(t) \right|^{\min\{2, p\}},$$
(5.31)

where the implicit constant depends on ω .

We shall show that

$$\lambda_1(0) \sim \lambda_+(0) \sim \mathfrak{s}R_0, \tag{5.32}$$

where the implicit constants depend on ω . We see from (5.9) and (5.11) that

$$\left. \frac{d}{dt} d_{\omega} \left(\psi(t) \right)^2 \right|_{t=t_0=0} = 2R_0 \frac{d}{dt} d_{\omega} \left(\psi(t) \right) \right|_{t=0} \ge 0, \tag{5.33}$$

which together with (5.20) yields

$$0 \le \mu^2 \lambda_1(0) \lambda_2(0) + \mu \lambda_1(0) \left| \Omega\left(\left\{ \frac{d\theta}{dt}(0) - \omega \right\} \eta(0) - N_\omega(\eta(0)), f_2 \right) \right|.$$
 (5.34)

Combining (5.34) with (5.30), we obtain that

$$0 \le \mu^2 \operatorname{sgn}[\lambda_1(0)] |\lambda_1(0)| \lambda_2(0) + C\mu |\lambda_1(0)|^{\min\{3,p+1\}}$$
(5.35)

for some constant C > 0 depending on ω . Furthermore, it follows from (5.35) that

$$-C|\lambda_1(0)|^{\min\{2,p\}} \le \mu \mathfrak{s}\lambda_2(0).$$
(5.36)

Suppose here that $\mathfrak{s} = \operatorname{sgn}[\lambda_1(0)] = 1$. Then, we see from (5.22), (5.36) and (5.21) that

$$0 < \lambda_1(0) \lesssim \lambda_1(0) - \frac{C}{\mu} |\lambda_1(0)|^{\min\{2,p\}} \lesssim \lambda_1(0) + \lambda_2(0) = \lambda_+(0),$$
 (5.37)

so that $\operatorname{sgn}[\lambda_+(0)] = 1$. Suppose next that $\operatorname{sgn}[\lambda_1(0)] = -1$. Then, (5.36) becomes $\mu\lambda_2(0) \leq C|\lambda_1(0)|^{\min\{2,p\}}$. This together with (5.21) shows that

$$\lambda_{+}(0) = \lambda_{1}(0) + \lambda_{2}(0) \le \lambda_{1}(0) + \frac{C}{\mu} |\lambda_{1}(0)|^{\min\{2,p\}} \sim \lambda_{1}(0).$$
(5.38)

Thus, we conclude that

$$\operatorname{sgn}[\lambda_1(0)] = \operatorname{sgn}[\lambda_+(0)]. \tag{5.39}$$

Since $\lambda_1(0)$ and $\lambda_+(0)$ are independent of t, (5.39) together with (5.19) and $d_{\omega}(\psi(0)) = R_0$ implies (5.32).

Next, we shall prove (5.14) and (5.13). Since we have (5.19), it suffices for (5.14) to show that for any $t \in [0, T_X]$,

$$\lambda_1(t) \sim \lambda_+(t) \sim \mathfrak{s} R_0 e^{\mu t}. \tag{5.40}$$

Let $\alpha > 1$ be a constant satisfying $|\lambda_1(0)| \leq \frac{1}{2}\alpha R_0$, and consider

$$T_{\alpha} := \sup \left\{ T \in [0, T_X] : |\lambda_1(t)| \le \alpha R_0 e^{\mu t} \text{ for all } t \in [0, T] \right\}.$$

$$(5.41)$$

Then, we have $T_{\alpha} > 0$. We shall show that $T_{\alpha} = T_X$. Suppose for contradiction that $T_{\alpha} < T_X$. Then,

$$\alpha R_0 e^{\mu T_\alpha} = |\lambda_1(T_\alpha)| \lesssim \delta_X, \tag{5.42}$$

where we have used (5.21) to obtain the inequality. Using the equation (4.62) for λ_+ , we have that

$$\frac{d}{dt} \left(e^{-\mu t} \lambda_{+}(t) \right) = e^{-\mu t} \left(\frac{d\lambda_{+}}{dt}(t) - \mu \lambda_{+}(t) \right)
= -e^{-\mu t} \left(\left\{ \frac{d\theta}{dt}(t) - \omega \right\} \eta(t) - N_{\omega} \left(\eta(t) \right), \, \mathfrak{U}_{-} \right)_{L^{2}_{real}}.$$
(5.43)

Furthermore, integrating the equation (5.43), and then using (5.31) and (5.21), we find that for any $t \in [0, T_X]$,

$$\begin{aligned} \left|\lambda_{+}(t) - e^{\mu t}\lambda_{+}(0)\right| &= \left|e^{\mu t}\int_{0}^{t} e^{-\mu s}\left(\left\{\frac{d\theta}{dt}(s) - \omega\right\}\eta(s) - N_{\omega}(\eta(s)), \mathcal{U}_{-}\right)_{L^{2}_{real}}ds\right| \\ &\lesssim \int_{0}^{t} e^{\mu(t-s)}\left|\lambda_{1}(s)\right|^{\min\{2,p\}}ds, \end{aligned}$$
(5.44)

where the implicit constant depends on ω . Similarly, we have

$$\left|\lambda_{-}(t) - e^{-\mu t}\lambda_{-}(0)\right| \lesssim \int_{0}^{t} e^{-\mu(t-s)} \left|\lambda_{1}(s)\right|^{\min\{2,p\}} ds.$$
 (5.45)

Then, we find from (5.44), (5.45), (5.32) and (5.21) that for any $t \in [0, T_X]$,

$$\begin{aligned} |\lambda_{1}(t)| &\leq |\lambda_{+}(t)| + |\lambda_{-}(t)| \\ &\lesssim e^{\mu t} |\lambda_{+}(0)| + e^{-\mu t} |\lambda_{-}(0)| + \int_{0}^{t} e^{\mu (t-s)} |\lambda_{1}(s)|^{\min\{2,p\}} ds \\ &\lesssim e^{\mu t} R_{0} + R_{0} + e^{\mu t} \int_{0}^{t} e^{-\mu s} |\lambda_{1}(s)|^{\min\{2,p\}} ds. \end{aligned}$$
(5.46)

This together with the definition of T_{α} (see (5.41)) and (5.42) shows that

$$\begin{aligned} |\lambda_1(T_{\alpha})| &\lesssim e^{\mu T_{\alpha}} R_0 + \frac{1}{\mu \min\{1, p-1\}} (\alpha R_0 e^{\mu T_{\alpha}})^{\min\{2, p\}} \\ &\leq e^{\mu T_{\alpha}} R_0 + \frac{1}{\mu \min\{1, p-1\}} \delta_X^{\min\{1, p-1\}} \alpha R_0 e^{\mu T_{\alpha}}. \end{aligned}$$
(5.47)

Hence, if $\frac{1}{\mu \min\{1,p-1\}} \delta_X^{\min\{1,p-1\}} \ll 1$, then (5.47) implies that $|\lambda_1(T_\alpha)| \leq \frac{1}{2} \alpha R_0 e^{\mu T_\alpha}$. However, this contradicts (5.42). Thus, we have proved that $T_\alpha = T_X$, and therefore for any $t \in [0, T_X]$,

$$|\lambda_1(t)| \lesssim R_0 e^{\mu t}.\tag{5.48}$$

We also see from (5.44), (5.45) and (5.48) that for any $t \in [0, T_X]$,

$$\left|\lambda_{+}(t) - e^{\mu t}\lambda_{+}(0)\right| \lesssim (R_{0}e^{\mu t})^{\min\{2,p\}},$$
(5.49)

$$\left|\lambda_{-}(t) - e^{-\mu t}\lambda_{-}(0)\right| \lesssim (R_0 e^{\mu t})^{\min\{2,p\}}.$$
 (5.50)

Furthermore, we find from (5.32), (5.49) and (5.50) that (5.40) holds. Combining (5.19) and (5.40), we also obtain (5.13).

We shall prove (5.15). It follows from (5.40) that for any $t \in [0, T_X]$,

$$|\lambda_{+}(t)| + |\lambda_{-}(t)| \lesssim |\lambda_{+}(t)| + |\lambda_{1}(t)| \sim R_{0}e^{\mu t}.$$
 (5.51)

Furthermore, (5.50) together with (5.51) shows that

$$\left|\lambda_{-}(t)\right| \lesssim e^{-\mu t} \left|\lambda_{-}(0)\right| + (R_{0}e^{\mu t})^{\min\{2,p\}} \lesssim R_{0} + (R_{0}e^{\mu t})^{\min\{2,p\}}.$$
 (5.52)

In order to complete the proof of (5.15), we employ the "nonlinear energy projected onto \mathcal{U}_{\pm} plane":

$$E_{\{\mathcal{U}_+,\mathcal{U}_-\}}(t) := \mathcal{S}_{\omega} \big(\Phi_{\omega} + \lambda_+(t)\mathcal{U}_+ + \lambda_-(t)\mathcal{U}_- \big) - \mathcal{S}_{\omega} \big(\Phi_{\omega} \big).$$
(5.53)

The second order Taylor's expansion around Φ_{ω} together with $\mathcal{S}'_{\omega}(\Phi_{\omega}) = 0, (4.3), \mathcal{L}_{\omega}\mathcal{U}_{\pm} = \pm i\mu\mathcal{U}_{\pm}$ and (4.45) shows that

$$E_{\{\mathfrak{U}_{+},\mathfrak{U}_{-}\}}(t) = \frac{1}{2} \left[S_{\omega}''(\Phi_{\omega}) \{\lambda_{+}(t)\mathfrak{U}_{+} + \lambda_{-}(t)\mathfrak{U}_{-}\} \right] \{\lambda_{+}(t)\mathfrak{U}_{+} + \lambda_{-}(t)\mathfrak{U}_{-}\} + O(\|\lambda_{+}(t)\mathfrak{U}_{+} + \lambda_{-}(t)\mathfrak{U}_{-}\|_{H^{1}}^{\min\{3,p+1\}}) = \frac{1}{2} \langle \mathcal{L}_{\omega} (\lambda_{+}(t)\mathfrak{U}_{+} + \lambda_{-}(t)\mathfrak{U}_{-}), \lambda_{+}(t)\mathfrak{U}_{+} + \lambda_{-}(t)\mathfrak{U}_{-}\rangle_{H^{-1},H^{1}} + O(\|\lambda_{+}(t)\mathfrak{U}_{+} + \lambda_{-}(t)\mathfrak{U}_{-}\|_{H^{1}}^{\min\{3,p+1\}}) = \frac{1}{2} (i\mu\lambda_{+}(t)\mathfrak{U}_{+} - i\mu\lambda_{-}(t)\mathfrak{U}_{-}, \lambda_{+}(t)\mathfrak{U}_{+} + \lambda_{-}(t)\mathfrak{U}_{-})_{L^{2}_{real}} + O(\|\lambda_{+}(t)\mathfrak{U}_{+} + \lambda_{-}(t)\mathfrak{U}_{-}\|_{H^{1}}^{\min\{3,p+1\}}) = -\mu\lambda_{+}(t)\lambda_{-}(t) + O(\|\lambda_{+}(t)\mathfrak{U}_{+} + \lambda_{-}(t)\mathfrak{U}_{-}\|_{H^{1}}^{\min\{3,p+1\}}).$$
(5.54)

We find from (4.69), (5.8), (5.54) and Lemma 4.2 that

$$S_{\omega}(\psi) - S_{\omega}(\Phi_{\omega}) - E_{\{\mathfrak{U}_{+},\mathfrak{U}_{-}\}}(t)$$

$$= \frac{1}{2} \langle \mathcal{L}_{\omega} \Gamma(t), \Gamma(t) \rangle_{H^{-1}, H^{1}}$$

$$+ O(\|\eta(t)\|_{H^{1}}^{\min\{3, p+1\}}) + O(\|\lambda_{+}(t)\mathfrak{U}_{+} + \lambda_{-}(t)\mathfrak{U}_{-}\|_{H^{1}}^{\min\{3, p+1\}})$$

$$\sim \|\Gamma(t)\|_{H^{1}}^{2} + O(\|\eta(t)\|_{H^{1}}^{\min\{3, p+1\}}) + O(\|\lambda_{+}(t)\mathfrak{U}_{+} + \lambda_{-}(t)\mathfrak{U}_{-}\|_{H^{1}}^{\min\{3, p+1\}}).$$
(5.55)

Moreover, it follows from $\mathcal{S}'_{\omega}(\Phi_{\omega}) = 0$, (4.3), (4.64) and (5.31) that

$$\begin{aligned} \left| \frac{d}{dt} E_{\{\mathcal{U}_{+},\mathcal{U}_{-}\}}(t) \right| \\ &= \left| \mathcal{S}_{\omega}'(\Phi_{\omega} + \lambda_{+}(t)\mathcal{U}_{+} + \lambda_{-}(t)\mathcal{U}_{-}) \right\{ \frac{d\lambda_{+}}{dt}(t)\mathcal{U}_{+} + \frac{d\lambda_{-}}{dt}(t)\mathcal{U}_{-} \right\} \right| \\ &\lesssim \left| \left[\mathcal{S}_{\omega}''(\Phi_{\omega}) \left(\lambda_{+}(t)\mathcal{U}_{+} + \lambda_{-}(t)\mathcal{U}_{-} \right) \right] \left\{ \frac{d\lambda_{+}}{dt}(t)\mathcal{U}_{+} + \frac{d\lambda_{-}}{dt}(t)\mathcal{U}_{-} \right\} \right| \\ &+ \left\| \lambda_{+}(t)\mathcal{U}_{+} + \lambda_{-}(t)\mathcal{U}_{-} \right\|_{H^{1}}^{\min\{2,p\}} \left\| \frac{d\lambda_{+}}{dt}(t)\mathcal{U}_{+} + \frac{d\lambda_{-}}{dt}(t)\mathcal{U}_{-} \right\|_{H^{1}} \\ &= \left| \left\langle \mathcal{L}_{\omega} \left(\lambda_{+}(t)\mathcal{U}_{+} + \lambda_{-}(t)\mathcal{U}_{-} \right), \frac{d\lambda_{+}}{dt}(t)\mathcal{U}_{+} + \frac{d\lambda_{-}}{dt}(t)\mathcal{U}_{-} \right\rangle_{H^{-1},H^{1}} \right| \\ &+ \left\| \lambda_{+}(t)\mathcal{U}_{+} + \lambda_{-}(t)\mathcal{U}_{-} \right\|_{H^{1}}^{\min\{2,p\}} \left\| \frac{d\lambda_{+}}{dt}(t)\mathcal{U}_{+} + \frac{d\lambda_{-}}{dt}(t)\mathcal{U}_{-} \right\|_{H^{1}} \\ &\lesssim \mu(|\lambda_{-}(t)| + |\lambda_{+}(t)|)^{\min\{3,p+1\}} \\ &+ \left(|\lambda_{+}(t)| + |\lambda_{-}(t)| \right)^{\min\{2,p\}} \left(\left| \frac{d\lambda_{+}}{dt}(t) \right| + \left| \frac{d\lambda_{-}}{dt}(t) \right| \right). \end{aligned}$$
(5.56)

Here, the equations (4.62) and (4.63) together with (5.28) and (5.27) show that for any $t \in [0, T_X]$,

$$\left| \frac{d\lambda_{\pm}}{dt}(t) \right| \leq \mu |\lambda_{\pm}(t)| + \left| \frac{d\theta}{dt}(t) - \omega \right| \|\eta(t)\|_{L^{2}} \|\mathcal{U}_{\pm}\|_{L^{2}} + \left| \left(N_{\omega} \left(\eta(t) \right), \mathcal{U}_{\mp} \right)_{L^{2}_{real}} \right| \\ \lesssim |\lambda_{+}(t)| + |\lambda_{-}(t)| + \|\eta(t)\|_{H^{1}}^{\min\{3,p+1\}} + \|\eta(t)\|_{H^{1}}^{\min\{2,p\}} \\ \lesssim |\lambda_{+}(t)| + |\lambda_{-}(t)| + \|\lambda_{+}(t)\mathcal{U}_{+} + \lambda_{-}(t)\mathcal{U}_{-} + \Gamma(t)\|_{H^{1}}^{\min\{2,p\}} \\ \lesssim |\lambda_{+}(t)| + |\lambda_{-}(t)| + \|\Gamma(t)\|_{H^{1}}^{\min\{2,p\}},$$
(5.57)

where the implicit constant depends on ω . Putting (5.56) and (5.57) together, we find that for any $t \in [0, T_X]$,

$$\left| \frac{d}{dt} E_{\{u_+, u_-\}}(t) \right|
\lesssim \left\| \Gamma(t) \right\|_{H^1}^{\min\{2, p\}} \left(|\lambda_+(t)| + |\lambda_-(t)| \right) + \left(|\lambda_+(t)| + |\lambda_-(t)| \right)^{\min\{3, p+1\}},$$
(5.58)

where the implicit constant depends on ω . Furthermore, we find from (5.55), the decomposition (4.66) and (5.58) that for any $t \in [0, T_X]$,

$$\begin{split} \|\Gamma(t)\|_{H^{1}}^{2} \\ \lesssim S_{\omega}(\psi) - S_{\omega}(\Phi_{\omega}) - E_{\{\mathcal{U}_{+},\mathcal{U}_{-}\}}(0) - \left(E_{\{\mathcal{U}_{+},\mathcal{U}_{-}\}}(t) - E_{\{\mathcal{U}_{+},\mathcal{U}_{-}\}}(0)\right) \\ + O(\|\eta(t)\|_{H^{1}}^{\min\{3,p+1\}}) + O(\|\lambda_{+}(t)\mathcal{U}_{+} + \lambda_{-}(t)\mathcal{U}_{-}\|_{H^{1}}^{\min\{3,p\}}) \\ \lesssim \|\Gamma(0)\|_{H^{1}}^{2} + \int_{0}^{t} \left|\frac{d}{dt}E_{\{\mathcal{U}_{+},\mathcal{U}_{-}\}}(t')\right| dt' \\ + \sup_{0 \leq t' \leq t} \|\Gamma(t') + \lambda_{+}(t')\mathcal{U}_{+} + \lambda_{-}(t')\mathcal{U}_{-}\|_{H^{1}}^{\min\{3,p+1\}} \\ + \sup_{0 \leq t' \leq t} \left(|\lambda_{+}(t')| + |\lambda_{-}(t')|\right)^{\min\{3,p+1\}} \\ \lesssim \|\Gamma(0)\|_{H^{1}}^{2} + \int_{0}^{t} \|\Gamma(t')\|_{H^{1}}^{\min\{2,p\}} \left(|\lambda_{+}(t')| + |\lambda_{-}(t')|\right) dt' \\ + \int_{0}^{t} \left(|\lambda_{+}(t')| + |\lambda_{-}(t')|\right)^{\min\{3,p+1\}} dt' \\ + \sup_{0 \leq t' \leq t} \|\Gamma(t')\|_{H^{1}}^{\min\{3,p+1\}} + \sup_{0 \leq t' \leq t} \left(|\lambda_{+}(t')| + |\lambda_{-}(t')|\right)^{\min\{3,p+1\}}. \end{split}$$

Suppose here that p < 2, so that $\min\{3, p+1\} = p+1$. Note that $p+1 < \frac{2}{2-p}$. Then, it follows from (5.59), Young's inequality, (5.19), (5.51) and (5.40) that for any $t \in [0, T_X]$,

$$\sup_{0 \le t' \le t} \|\Gamma(t')\|_{H^{1}}^{2} \lesssim \|\Gamma(0)\|_{H^{1}}^{2} + \int_{0}^{t} \left(|\lambda_{+}(t')| + |\lambda_{-}(t')|\right)^{\frac{2}{2-p}} dt' + \int_{0}^{t} \left(|\lambda_{+}(t')| + |\lambda_{-}(t')|\right)^{p+1} dt' + \sup_{0 \le t' \le t} \left\{|\lambda_{+}(t')| + |\lambda_{-}(t')|\right\}^{p+1} \lesssim |\lambda_{1}(0)|^{2} + \int_{0}^{t} |\lambda_{1}(t')|^{p+1} dt' + \sup_{0 \le t' \le t} |\lambda_{1}(t')|^{p+1} \lesssim R_{0}^{2} + \int_{0}^{t} \left(R_{0}e^{\mu t'}\right)^{p+1} dt + \sup_{0 \le t' \le t} \left(R_{0}e^{\mu t'}\right)^{p+1} \lesssim R_{0}^{2} + (R_{0}e^{\mu t})^{p+1}.$$
(5.60)

This together with (5.52) gives us the desired result (5.15) for p < 2. Similarly, we can prove (5.15) for $p \ge 2$.

We shall prove that there exists $T_* > 0$ such that $d_{\omega}(\psi(t))$ is strictly increasing on

 $[T_*R_0^{\min\{1,p-1\}}, T_X]$. It follows from (5.19), (5.20) and (5.30) that for any $t \in [0, T_X]$,

$$\begin{aligned} |\lambda_1(t)| \frac{d}{dt} d_\omega(\psi(t)) &\sim d_\omega(\psi(t)) \frac{d}{dt} d_\omega(\psi(t)) = \frac{1}{2} \frac{d}{dt} d_\omega(\psi(t))^2 \\ &\geq 2\mu^2 |\lambda_1(t)| \mathfrak{s} \lambda_2(t) - C_1 \mu |\lambda_1(t)|^{\min\{3,p+1\}}, \end{aligned}$$
(5.61)

so that

$$\frac{d}{dt}d_{\omega}(\psi(t)) \gtrsim \mu^2 \mathfrak{s}\lambda_2(t) - C_1 \mu |\lambda_1(t)|^{\min\{2,p\}},\tag{5.62}$$

where $C_1 > 0$ is constant depending only on d, p and ω . Here, multiplying the equation (4.90) by \mathfrak{s} , integrating the resulting equation, and using (5.31) and (5.40), we obtain that

$$\mathfrak{s}\lambda_{2}(t) = \mathfrak{s}\lambda_{2}(0) + \mu \int_{0}^{t} \mathfrak{s}\lambda_{1}(t') dt' -\mathfrak{s}\frac{1}{2} \int_{0}^{t} \left(\left\{\frac{d\theta}{dt}(t') - \omega\right\} \eta(t') - N_{\omega}(\eta(t')), \mathcal{U}_{+} + \mathcal{U}_{-}\right)_{L^{2}_{real}} dt'$$
(5.63)
$$\geq \mathfrak{s}\lambda_{2}(0) + C_{2}R_{0}(e^{\mu t} - 1) - C_{3} \int_{0}^{t} (R_{0}e^{\mu t'})^{\min\{2,p\}} dt',$$

where C_2 and C_3 are some positive constants depending only on d, p and ω . Here, we see from (5.36) and (5.32) that

$$\mu \mathfrak{s} \lambda_2(0) \gtrsim -|\lambda_1(0)|^{\min\{2,p\}} \sim -R_0^{\min\{2,p\}},\tag{5.64}$$

where the implicit constants depend on ω . Moreover, it follows from (5.40) and (5.21) that for any $t \in [0, T_X]$,

$$|\lambda_1(t)| \sim R_0 e^{\mu t} \lesssim \delta_X, \tag{5.65}$$

so that, taking δ_X sufficiently small dependently on ω , we have

$$C_3 \int_0^t (R_0 e^{\mu t'})^{\min\{2,p\}} dt' \le \frac{C_2}{2} \mu \int_0^t R_0 e^{\mu t'} dt' = \frac{C_2}{2} R_0 (e^{\mu t} - 1).$$
(5.66)

Putting the estimates (5.63), (5.64) and (5.66) together, we find that for any $t \in [0, T_X]$,

$$\mathfrak{s}\lambda_2(t) \ge \frac{C_2}{2} R_0(e^{\mu t} - 1) - C_4 R_0^{\min\{2,p\}} \ge \frac{C_2}{2} R_0 \mu t - C_4 R_0^{\min\{2,p\}}$$
(5.67)

for some constants $C_4 > 0$ depending only on d, p and ω . Hence, if we choose $T_* \ge \frac{4C_4}{\mu C_2}$, then for any $t \in [T_*R_0^{\min\{1,p-1\}}, T_X]$,

$$\mathfrak{s}\lambda_2(t) \ge T_*C_2\mu R_0^{\min\{2,p\}}.$$
(5.68)

Furthermore, it follows from (5.50), (5.40) and (5.51) that

$$\begin{aligned} |\lambda_{2}(t)| &\geq |\lambda_{1}(t)| - |\lambda_{-}(t)| \\ &\geq |\lambda_{1}(t)| - e^{-\mu t} \lambda_{-}(0) - C_{5}(e^{\mu t}R_{0})^{\min\{2,p\}} \\ &\geq c_{1}R_{0}e^{\mu t} - C_{6}R_{0}e^{-\mu t} - C_{5}(R_{0}e^{\mu t})^{\min\{2,p\}} \end{aligned}$$
(5.69)

for some constants $c_1 \in (0,1)$, $C_5 > 0$ and $C_6 > 0$ depending only on d, p and ω . Hence, if $t \geq \frac{1}{2\mu} \log(\frac{4C_6}{c_1})$, then we find from (5.69) and (5.40) that

$$|\lambda_2(t)| \ge \frac{c_1}{10} R_0 e^{\mu t} \sim |\lambda_1(t)|.$$
(5.70)

Furthermore, we see from (5.62), (5.70) and (5.65) that for any $t \ge \frac{1}{2\mu} \log \left(\frac{4C_6}{c_1}\right)$,

$$\frac{d}{dt}d_{\omega}(\psi(t)) > 0.$$
(5.71)

On the other hand, if $t \leq \frac{1}{2\mu} \log(\frac{4C_6}{c_1})$, then we have

$$|\lambda_1(t)| \sim R_0 e^{\mu t} \lesssim R_0. \tag{5.72}$$

Hence, choosing T_* suitably, and using (5.62), (5.68) and (5.72), we conclude that for any $t \in [T_*R_0^{\min\{1,p-1\}}, T_X]$ with $t \leq \frac{1}{2\mu} \log(\frac{4C_6}{c_1})$,

$$\frac{d}{dt}d_{\omega}(\psi(t)) \ge \frac{\mu^3 C_2 T_*}{2} R_0^{\min\{2,p\}} - C_1 |\lambda_1(t)|^{\min\{2,p\}} > 0.$$
(5.73)

Putting (5.71) and (5.73) together, we obtain the desired result.

We shall prove (5.17). It follows from $d_{\omega}(\psi(0)) = R_0$ and the fundamental theorem of calculus that

$$\left|d_{\omega}(\psi(t)) - R_0\right| \le \int_0^t \left|\frac{d}{ds}d_{\omega}(\psi(s))\right| ds.$$
(5.74)

Here, it follows from (5.19), (5.20), (5.30) and (5.40) that

$$\begin{aligned} |\lambda_1(t)| \Big| \frac{d}{dt} d_\omega(\psi(t)) \Big| &\sim \Big| 2d_\omega(\psi(t)) \frac{d}{dt} d_\omega(\psi(t)) \Big| = \Big| \frac{d}{dt} d_\omega(\psi(t))^2 \Big| \\ &\lesssim |\lambda_1(t)| |\lambda_2(t)| + |\lambda_1(t)|^{\min\{3,p+1\}} \\ &= |\lambda_1(t)| |\lambda_+(t) - \lambda_1(t)| + |\lambda_1(t)|^{\min\{3,p+1\}} \lesssim |\lambda_1(t)|^2, \end{aligned}$$

$$(5.75)$$

where the implicit constants depend on ω . Furthermore, this together with (5.40) gives us that for any $t \in [0, T_*R_0^{\min\{1, p-1\}}]$

$$\left|\frac{d}{dt}d_{\omega}(\psi(t))\right| \lesssim R_0 e^{\mu t}.$$
(5.76)

Putting (5.74) and (5.76) together, we find that for any $t \in [0, T_*R_0^{\min\{1, p-1\}}]$,

$$\left| d_{\omega}(\psi(t)) - R_0 \right| \lesssim \int_0^t R_0 e^{\mu s} \, ds \lesssim T_* R_0^{\min\{2,p\}} e^{\mu T_* R_0^{\min\{1,p-1\}}} \lesssim R_0^{\min\{2,p\}}, \qquad (5.77)$$

where the implicit constants depend on ω . Thus, we have proved (5.17).

Finally, we shall prove (5.16). Taylor's expansion of \mathcal{K} around Φ_{ω} together with $\mathcal{K}(\Phi_{\omega}) = 0$ and (4.151) shows that

$$\mathcal{K}(\psi(t)) = \mathcal{K}(\Phi_{\omega} + \eta(t)) = \mathcal{K}'(\Phi_{\omega})\eta(t) + O(\|\eta(t)\|_{H^{1}}^{2})$$

$$= -2\mu s_{p}\lambda_{1}(t)(\Phi_{\omega}, f_{2})_{L^{2}} - 2(1 - s_{p})(2^{*} - 2)\lambda_{1}(t)(\Phi_{\omega}^{2^{*}-1}, f_{1})_{L^{2}}$$

$$- s_{p}(p - 1)(\Phi_{\omega}^{p}, \Gamma(t))_{L^{2}_{real}} - (2^{*} - 2)(\Phi_{\omega}^{2^{*}-1}, \Gamma(t))_{L^{2}_{real}}$$

$$+ \omega \mathcal{M}(\eta(t)) + O(\|\eta(t)\|_{H^{1}}^{2}).$$
(5.78)

Furthermore, multiplying the both sides above by $\mathfrak{s} = \operatorname{sgn}[\lambda_1]$, and using $(\Phi_{\omega}, f_2)_{L^2} < 0$ (see (4.121)), (5.7) and (5.19), we find that for any $t \in [0, T_X]$,

$$\mathfrak{sK}(\psi(t)) \geq 2s_{p}\mu |(\Phi_{\omega}, f_{2})_{L^{2}}||\lambda_{1}(t)| - 2(1 - s_{p})(2^{*} - 2)|(\Phi_{\omega}^{2^{*}-1}, f_{1})_{L^{2}}||\lambda_{1}(t)| - (p - 1)||\Phi_{\omega}||_{L^{p+1}}^{p}||\Gamma(t)||_{L^{p+1}} - (2^{*} - 2)||\Phi_{\omega}||_{L^{2^{*}}}^{2^{*}-1}||\Gamma(t)||_{L^{2^{*}}} - O(|\lambda_{1}(t)|^{2}) \geq s_{p}\mu |(\Phi_{\omega}, f_{2})_{L^{2}}||\lambda_{1}(t)| - (p - 1)||\Phi_{\omega}||_{L^{p+1}}^{p}||\Gamma(t)||_{L^{p+1}} - (2^{*} - 2)||\Phi_{\omega}||_{L^{2^{*}}}^{2^{*}-1}||\Gamma(t)||_{L^{2^{*}}} - O(|\lambda_{1}(t)|^{2}).$$
(5.79)

Recall here that $|\lambda_1(t)| \leq \delta_X \ll 1$ for $t \in [0, T_X]$ (see (5.19)). Then, we conclude from (5.79), (5.40) and the proved result (5.15) that there exists a constant $C_* > 0$ depending only on d, p and ω such that

$$\mathfrak{s}\mathcal{K}(\psi(t)) \gtrsim R_0 e^{\mu t} - C_* R_0, \tag{5.80}$$

where the implicit constant depends on ω . Thus, we have completed the proof.

6 Modified distance function

In this section, we continue to study the decomposition of the form (4.1) with (4.48), (4.53) and (4.54). Our aim here is to introduce the "modified distance function" \tilde{d}_{ω} (see Proposition 6.2). To this end, we introduce two distances between a function $u \in H^1(\mathbb{R}^d)$ and the orbit of Φ_{ω} :

$$\operatorname{dist}_{H^1}(u, \mathcal{O}(\Phi_\omega)) := \inf_{\theta \in \mathbb{R}} \|u - e^{i\theta} \Phi_\omega\|_{H^1},$$
(6.1)

$$\operatorname{dist}_{L^2}(u, \mathcal{O}(\Phi_\omega)) := \inf_{\theta \in \mathbb{R}} \|u - e^{i\theta} \Phi_\omega\|_{L^2}.$$
(6.2)

Lemma 6.1. Assume $d \geq 3$, $1 + \frac{4}{d} , and let <math>\omega_2$ be the frequency given by Proposition 4.1. Then, for any $\omega \in (0, \omega_2)$, there exists $\gamma_1(\omega) > 0$ with the following property: let ψ be a solution to (NLS) satisfying $\mathcal{M}(\psi) = \mathcal{M}(\Phi_\omega)$. If $\operatorname{dist}_{H^1}(\psi(t), \mathcal{O}(\Phi_\omega)) \leq \gamma_1(\omega)$, then

$$\|\eta(t)\|_E \lesssim \operatorname{dist}_{H^1}(\psi(t), \mathcal{O}(\Phi_\omega)) \le \|\eta(t)\|_E,$$
(6.3)

and if $\operatorname{dist}_{L^2}(\psi(t), \mathcal{O}(\Phi_\omega)) \leq \gamma_1(\omega)$, then

$$\|\eta(t)\|_{L^2} \sim \operatorname{dist}_{L^2}(\psi(t), \mathcal{O}(\Phi_\omega)).$$
(6.4)

Here, η is the function appearing in the decomposition for ψ of the form (4.1) with (4.48), (4.53) and (4.54). Moreover, the implicit constants in (6.3) and (6.4) depend on ω as well as d and p. Proof of Lemma 6.1. Note first that Lemma 4.3 shows

$$\operatorname{dist}_{H^{1}}(\psi(t), \mathcal{O}(\Phi_{\omega})) \leq \|\psi(t) - e^{i\theta(t)}\Phi_{\omega}\|_{H^{1}} = \|\eta(t)\|_{H^{1}} \lesssim \|\eta(t)\|_{E}.$$
(6.5)

Moreover, we can take a continuous function $\theta_0(t)$ of t such that

$$\|\psi(t) - e^{i\theta_0(t)}\Phi_\omega\|_{H^1} = \operatorname{dist}_{H^1}(\psi(t), \mathcal{O}(\Phi_\omega)).$$
(6.6)

Put $\eta_0(t) := \psi(t) - e^{i\theta_0(t)} \Phi_\omega$, so that $\|\eta_0(t)\|_{H^1} = \operatorname{dist}_{H^1}(\psi(t), \mathcal{O}(\Phi_\omega))$. Then, it follows from $\Omega(\Phi_\omega, \partial_\omega \Phi_\omega) = (\Phi_\omega, i\partial_\omega \Phi_\omega)_{L^2_{real}} = 0$ and (4.53) that

$$\Omega(e^{-i\theta(t)}\eta_0(t),\partial_{\omega}\Phi_{\omega}) = \Omega(e^{-i\theta(t)}\psi(t),\partial_{\omega}\Phi_{\omega}) - \Omega(e^{i(\theta_0(t)-\theta(t))}\Phi_{\omega},\partial_{\omega}\Phi_{\omega})$$

$$= -\left(ie^{i(\theta_0(t)-\theta(t))}\Phi_{\omega},\partial_{\omega}\Phi_{\omega}\right)_{L^2_{real}}$$

$$= \sin\left(\theta_0(t) - \theta(t)\right)\left(\Phi_{\omega},\partial_{\omega}\Phi_{\omega}\right)_{L^2_{real}}.$$
(6.7)

Hence, we have

$$\begin{aligned} \left| \sin \left(\theta_0(t) - \theta(t) \right) \right| \left| \left(\Phi_\omega, \partial_\omega \Phi_\omega \right)_{L^2_{real}} \right| &\leq \| \eta_0(t) \|_{L^2} \| \partial_\omega \Phi_\omega \|_{L^2} \\ &\leq \operatorname{dist}_{H^1} \left(\psi(t), \mathcal{O}(\Phi_\omega) \right) \| \partial_\omega \Phi_\omega \|_{L^2}. \end{aligned}$$

$$\tag{6.8}$$

We see from (6.8) and $\operatorname{dist}_{H^1}(\psi(t), \mathcal{O}(\Phi_\omega)) \ll 1$ that

$$\inf_{k\in\mathbb{Z}} \left| \theta_0(t) - \theta(t) + k\pi \right| \lesssim \operatorname{dist}_{H^1}(\psi(t), \mathcal{O}(\Phi_\omega)),$$
(6.9)

where the implicit constant depends on ω . Here, we can eliminate the case where k is odd in the above infimum. Indeed, it follows from the choice of $\theta(t)$ (see (4.54)) that

$$0 > (e^{-i\theta(t)}\psi(t), \partial_{\omega}\Phi_{\omega})_{L^{2}_{real}}$$

$$= \cos\left(\theta_{0}(t) - \theta(t)\right)\left(\Phi_{\omega}, \partial_{\omega}\Phi_{\omega}\right)_{L^{2}_{real}} + \left(e^{-i\theta(t)}\eta_{0}(t), \partial_{\omega}\Phi_{\omega}\right)_{L^{2}_{real}},$$
(6.10)

so that

$$-\cos\left(\theta_0(t) - \theta(t)\right)\left(\Phi_\omega, \partial_\omega \Phi_\omega\right)_{L^2_{real}} > \left(e^{-i\theta(t)}\eta_0(t), \partial_\omega \Phi_\omega\right)_{L^2_{real}}.$$
(6.11)

Thus, supposing for contradiction that $\theta_0(t) - \theta(t)$ lay near an odd multiple of π , we have

$$\frac{1}{2}(\Phi_{\omega},\partial_{\omega}\Phi_{\omega})_{L^{2}_{real}} > (e^{-i\theta(t)}\eta_{0}(t),\partial_{\omega}\Phi_{\omega})_{L^{2}_{real}}.$$
(6.12)

However, this together with (iii) of Proposition 2.4 shows

$$1 \lesssim \frac{1}{2} \left| (\Phi_{\omega}, \partial_{\omega} \Phi_{\omega})_{L^{2}_{real}} \right| < \|\eta_{0}(t)\|_{L^{2}} \|\partial_{\omega} \Phi_{\omega}\|_{L^{2}} \le \operatorname{dist}_{H^{1}} \left(\psi(t), \mathcal{O}(\Phi_{\omega}) \right) \|\partial_{\omega} \Phi_{\omega}\|_{L^{2}}, \quad (6.13)$$

which contradicts that $\operatorname{dist}_{H^1}(\psi(t), \mathcal{O}(\Phi_\omega)) \ll 1$. Since $\operatorname{dist}_{H^1}(\psi(t), \mathcal{O}(\Phi_\omega)) \ll 1$ implies that $\theta_0(t) - \theta(t)$ lies near an even multiple of π , we see from Lemma 4.3 and (6.9) that

$$\begin{aligned} \|\eta(t)\|_{E} &\lesssim \|\eta(t)\|_{H^{1}} = \|e^{-i\theta(t)}\psi(t) - \Phi_{\omega}\|_{H^{1}} \\ &\leq |e^{i(\theta_{0}(t) - \theta(t))} - 1| \|\Phi_{\omega}\|_{H^{1}} + \|\eta_{0}(t)\|_{H^{1}} \\ &\lesssim |\cos\left(\theta_{0}(t) - \theta(t)\right) - 1| + |\sin\left(\theta_{0}(t) - \theta(t)\right)| + \|\eta_{0}(t)\|_{H^{1}} \\ &\lesssim \inf_{k \in \mathbb{Z}} |\theta_{0}(t) - \theta(t) - 2k\pi| + \|\eta_{0}(t)\|_{H^{1}} \\ &\lesssim \operatorname{dist}_{H^{1}}(\psi(t), \mathcal{O}(\Phi_{\omega})). \end{aligned}$$
(6.14)

Putting (6.5) and (6.14) together, we obtain the desired result (6.3). The same argument as the above shows (6.4). \Box

Now, we recall that $H^1_{\omega}(\mathbb{R}^d)$ denotes the set $\{u \in H^1(\mathbb{R}^d) : \mathcal{M}(u) = \mathcal{M}(\Phi_{\omega})\}$ (see (4.84)). The following proposition is an immediate consequence of Lemma 6.1:

Proposition 6.2. Assume $d \geq 3$, $1 + \frac{4}{d} , and let <math>\omega_2$ be the frequency given by Proposition 4.1. Then, for any $\omega \in (0, \omega_2)$, there exist a constant $\tilde{\gamma}(\omega) \in (0, \delta_E(\omega))$ ($\delta_E(\omega)$ denotes the constant given by Proposition 4.4) and a continuous function $\tilde{d}_{\omega}: H^1_{\omega}(\mathbb{R}^d) \to [0, \infty)$ such that:

$$\widetilde{d}_{\omega}(u) \sim \operatorname{dist}_{H^1}(u, \mathcal{O}(\Phi_{\omega})),$$
(6.15)

where the implicit constant depends only on d, p and ω ; and if $\widetilde{d}_{\omega}(u) \leq \widetilde{\gamma}(\omega)$, then

$$\widetilde{d}_{\omega}(u) = d_{\omega}(u), \tag{6.16}$$

where d_{ω} is the distance function defined by (4.85).

Remark 6.1. We can take the constant δ_X given by the ejection lemma (Lemma 5.2) so small that $\delta_X \leq \tilde{\gamma}(\omega)$. Hence, we may assume that $\delta_X < \tilde{\gamma}(\omega)$ in what follows.

Lemma 6.3. Assume $d \ge 3$ and $1 + \frac{4}{d} , and let <math>\omega_2$ be the frequency given by Proposition 4.1. Then, for any $\omega \in (0, \omega_2)$ and any $\delta > 0$, there exist $\varepsilon_0(\delta) > 0$ and $\kappa_1(\delta) > 0$ with the following properties: $\varepsilon_0(\delta)$ and $\kappa_1(\delta)$ are non-decreasing with respect to δ ; and for any radial function $u \in H^1(\mathbb{R}^d)$ satisfying

$$\mathcal{M}(u) = \mathcal{M}(\Phi_{\omega}),\tag{6.17}$$

$$\mathcal{S}_{\omega}(u) < m_{\omega} + \varepsilon_0(\delta), \tag{6.18}$$

$$\widetilde{d}_{\omega}(u) \ge \delta, \tag{6.19}$$

we have either

$$\mathcal{K}(u) \le -\kappa_1(\delta) \tag{6.20}$$

or

$$\mathcal{K}(u) \ge \min\left\{\kappa_1(\delta), \frac{1}{2} \|\nabla u\|_{L^2}^2\right\}.$$
 (6.21)

Here, \tilde{d}_{ω} denotes the function given by Proposition 6.2.

Proof of Lemma 6.3. We see from Hölder's inequality that if $u \in H^1(\mathbb{R}^d)$ satisfies $\mathcal{M}(u) = \mathcal{M}(\Phi_\omega)$, then

$$\|u\|_{L^{p+1}}^{p+1} \le \|u\|_{L^2}^{p+1-\frac{d(p-1)}{2}} \|u\|_{L^{2^*}}^{\frac{d(p-1)}{2}} = \mathcal{M}(\Phi_\omega)^{\frac{p+1}{2}-\frac{d(p-1)}{4}} \|u\|_{L^{2^*}}^{\frac{d(p-1)}{2}}.$$
 (6.22)

Since $2 < \frac{d(p-1)}{2} < 2^*$, we see from (6.22) and Sobolev's embedding that there exists $c_0 > 0$ depending only on d, p and ω such that if $u \in H^1(\mathbb{R}^d)$ satisfies $\mathcal{M}(u) = \mathcal{M}(\Phi_\omega)$ and $\|\nabla u\|_{L^2}^2 \leq c_0$, then $\mathcal{K}(u) \geq \frac{1}{2} \|\nabla u\|_{L^2}^2$. Thus, for the desired result, it is sufficient to prove that for any $\delta > 0$, there exist $\varepsilon_0(\delta) > 0$ and $\kappa_0(\delta) > 0$ with the following property: for any radial function $u \in H^1(\mathbb{R}^d)$ satisfying (6.17) through (6.19) and $\|\nabla u\|_{L^2}^2 \geq c_0$, we have $|\mathcal{K}(u)| > \kappa_1(\delta)$. We prove this by contradiction argument. Hence, we suppose to the contrary that there exists $\delta_0 > 0$ with the following property: for any number $n \geq 1$, there exists a radial function $u_n \in H^1(\mathbb{R}^d)$ satisfying

$$\mathcal{M}(u_n) = \mathcal{M}(\Phi_\omega), \tag{6.23}$$

$$\mathcal{S}_{\omega}(u_n) < m_{\omega} + \frac{1}{n},\tag{6.24}$$

$$\widetilde{d}_{\omega}(u_n) \ge \delta_0, \tag{6.25}$$

$$\inf_{n \ge 1} \|\nabla u_n\|_{L^2} \ge c_0, \tag{6.26}$$

$$|\mathcal{K}(u_n)| \le \frac{1}{n}.\tag{6.27}$$

Then, we see from (6.24) and (6.27) that

$$\frac{s_p}{d} \|\nabla u_n\|_{L^2}^2 \le \mathcal{I}_{\omega}(u_n) = \mathcal{S}_{\omega}(u_n) - \frac{2}{d(p-1)} \mathcal{K}(u_n) \le m_{\omega} + 2, \tag{6.28}$$

so that

$$||u_n||_{H^1}^2 \le 2\mathcal{M}(\Phi_{\omega}) + \frac{d}{s_p}(m_{\omega} + 2).$$
(6.29)

Hence, we can take a radial function $u_{\infty} \in H^1(\mathbb{R}^d)$ and a subsequence of $\{u_n\}$ (still denoted by the same symbol $\{u_n\}$) such that

$$\lim_{n \to \infty} u_n = u_{\infty} \qquad \text{weakly in } H^1(\mathbb{R}^d) \tag{6.30}$$

and for any $2 < q < 2^*$,

$$\lim_{n \to \infty} u_n = u_{\infty} \qquad \text{strongly in } L^q(\mathbb{R}^d).$$
(6.31)

We shall show that u_{∞} is non-trivial by using the contradiction argument. Hence, suppose to the contrary that $u_{\infty} \equiv 0$. Then, we see from (6.31) that

$$\lim_{n \to \infty} \|u_n\|_{L^{p+1}} = 0. \tag{6.32}$$

Moreover, (6.27) together with (6.32) shows that, passing to some subsequence,

$$\lim_{n \to \infty} \|\nabla u_n\|_{L^2}^2 = \lim_{n \to \infty} \|u_n\|_{L^{2^*}}^{2^*}.$$
(6.33)

We see from the definition of σ (see (1.27)) and (6.33) that

$$\lim_{n \to \infty} \|\nabla u_n\|_{L^2}^2 \ge \sigma \lim_{n \to \infty} \|u_n\|_{L^{2^*}}^2 = \sigma \lim_{n \to \infty} \|\nabla u_n\|_{L^2}^{\frac{4}{2^*}},$$
(6.34)

which together with (6.33) and (6.26) yields

$$\lim_{n \to \infty} \|u_n\|_{L^{2^*}}^2 \ge \sigma^{\frac{d}{2}}.$$
(6.35)

Furthermore, we see from the definition of \mathcal{J}_{ω} (see (1.12) and (1.14)), (6.24), (6.27) and (6.35), we find that

$$\frac{1}{d}\sigma^{\frac{d}{2}} \le \lim_{n \to \infty} \frac{1}{d} \|u_n\|_{L^{2^*}}^{2^*} \le \lim_{n \to \infty} \mathcal{J}_{\omega}(u_n) \le \lim_{n \to \infty} \left\{ \mathcal{S}_{\omega}(u_n) + \frac{1}{2} |\mathcal{K}(u_n)| \right\} \le m_{\omega}.$$
(6.36)

However, this contradicts (1.35) and therefore u_{∞} is non-trivial.

Next, we shall prove $u_{\infty} = e^{i\theta_0} \Phi_{\omega}$ for some θ_0 . We see from (6.24) and (6.27) that

$$\mathcal{I}_{\omega}(u_{\infty}) \leq \lim_{n \to \infty} \mathcal{I}_{\omega}(u_n) \leq \lim_{n \to \infty} \left\{ \mathcal{S}_{\omega}(u_n) + \frac{2}{d(p-1)} |\mathcal{K}(u_n)| \right\} \leq m_{\omega}.$$
(6.37)

Hence, from the point of view of (1.10), what we need to prove is that $\mathcal{K}(u_{\infty}) \leq 0$. We prove this by contradiction. Hence, suppose to the contrary that $\mathcal{K}(u_{\infty}) > 0$. Then, the Brezis-Lieb lemma together with (6.27) shows

$$\lim_{n \to \infty} \mathcal{K}(u_n - u_\infty) = -\lim_{n \to \infty} \left\{ \mathcal{K}(u_n) - \mathcal{K}(u_n - u_\infty) \right\} = -\mathcal{K}(u_\infty) < 0, \tag{6.38}$$

so that for any sufficiently large $n \in \mathbb{N}$,

$$\mathcal{K}(u_n - u_\infty) < 0. \tag{6.39}$$

Hence, we can take $\lambda_n < 1$ such that $\mathcal{K}(\lambda_n^{\frac{d}{2}} \{ u_n(\lambda_n \cdot) - u_\infty(\lambda_n \cdot) \}) = 0$ (cf. Lemma 2.1 in [3]). Moreover, this implies that

$$m_{\omega} \leq \mathcal{I}_{\omega} \left(\lambda_n^{\frac{u}{2}} \left\{ u_n(\lambda_n \cdot) - u_{\infty}(\lambda_n \cdot) \right\} \right) \leq \mathcal{I}_{\omega}(u_n - u_{\infty})$$

$$= \mathcal{I}_{\omega}(u_n) - \mathcal{I}_{\omega}(u_{\infty}) + o_n(1) = \mathcal{S}_{\omega}(u_n) - \frac{2}{d(p-1)} \mathcal{K}(u_n) - \mathcal{I}_{\omega}(u_{\infty}) + o_n(1),$$
(6.40)

which together with (6.24) and (6.27) shows

$$m_{\omega} \le m_{\omega} - \mathcal{I}_{\omega}(u_{\infty}). \tag{6.41}$$

Since $\mathcal{I}_{\omega}(u_{\infty}) > 0$, this is a contradiction. Hence, $\mathcal{K}(u_{\infty}) \leq 0$. In particular, we find from (6.37) that u_{∞} is a minimizer of the variational problem (1.10). Since any minimizer of \mathcal{I}_{ω} is also a ground state (see Proposition 1.2 in [2]), we conclude from the radial symmetry and non-triviality of u_{∞} , and Proposition 2.1 that $u_{\infty} = e^{i\theta_0}\Phi_{\omega}$ for some $\theta_0 \in \mathbb{R}$. In particular, $\mathcal{K}(u_{\infty}) = 0$ and $\mathcal{I}(u_{\infty}) = m_{\omega}$. However, the condition (6.25) prevents u_{∞} from existing. Indeed, it follows from the weak convergence (6.30) and $\lim_{n\to\infty} \mathcal{I}_{\omega}(u_n) \leq m_{\omega} = \mathcal{I}(u_{\infty})$ that

$$\lim_{n \to \infty} \|u_n\|_{L^2} = \|u_\infty\|_{L^2}, \qquad \lim_{n \to \infty} \|\nabla u_n\|_{L^2} = \|\nabla u_\infty\|_{L^2}.$$
(6.42)

Hence, we find that

$$\lim_{n \to \infty} u_n = u_{\infty} \qquad \text{strongly in } H^1(\mathbb{R}^d). \tag{6.43}$$

Using the strong convergence (6.43), we have

$$\delta_0 < \inf_{\theta} \|u_n - e^{i\theta} \Phi_{\omega}\|_{H^1} \le \|u_n - e^{i\theta_0} \Phi_{\omega}\|_{H^1} = \|u_n - u_\infty\|_{H^1} = o_n(1).$$
(6.44)

However, this is a contradiction. Thus, we cannot take a sequence $\{u_n\}$ satisfying (6.23) through (6.27) and therefore the lemma holds.

Lemma 6.4. Assume $d \ge 3$ and $1 + \frac{4}{d} , and let <math>\omega_2$ be the frequency given by Proposition 4.1. Then, for any $\omega \in (0, \omega_2)$ and any $\delta > 0$, there exists $\kappa_2(\delta) > 0$ such that for any radial function $u \in H^1(\mathbb{R}^d)$ satisfying $\mathcal{J}_{\omega}(u) \le m_{\omega} - \delta$, the following holds:

$$\mathcal{K}(u) \ge \min\left\{\kappa_2(\delta), \frac{1}{2} \|\nabla u\|_{L^2}^2\right\}.$$
 (6.45)

Proof of Lemma 6.4. Let u be a function satisfying $\mathcal{J}_{\omega}(u) \leq m_{\omega} - \delta$. Then, we see from (1.14) and (1.10) that

$$\|u\|_{L^2}^2 \le \frac{2}{\omega} \mathcal{J}_{\omega}(u) \le \frac{2}{\omega} m_{\omega}, \tag{6.46}$$

$$\mathcal{K}(u) > 0. \tag{6.47}$$

Furthermore, it follows from Hölder's inequality and (6.46) that

$$\|u\|_{L^{p+1}}^{p+1} \le \|u\|_{L^2}^{p+1-\frac{d(p-1)}{2}} \|u\|_{L^{2^*}}^{\frac{d(p-1)}{2}} \le \left(\frac{2}{\omega^2}m_\omega\right)^{\frac{p+1}{2}-\frac{d(p-1)}{4}} \|u\|_{L^{2^*}}^{\frac{d(p-1)}{2}}.$$
 (6.48)

Hence, we find from (6.48), $2 < \frac{d(p-1)}{2} < 2^*$ and Sobolev's embedding that there exists $c_0 > 0$ depending only on d, p and ω such that if $u \in H^1(\mathbb{R}^d)$ satisfies $\mathcal{J}_{\omega}(u) \leq m_{\omega} - \delta$ and $\|\nabla u\|_{L^2}^2 \leq c_0$, then $\mathcal{K}(u) \geq \frac{1}{2} \|\nabla u\|_{L^2}^2$. Thus, for the desired result (6.45), it is sufficient to prove that for any $\delta > 0$, there exists $\kappa_2(\delta) > 0$ with the following property: if $u \in H^1(\mathbb{R}^d)$ is radial and satisfies that $\mathcal{J}_{\omega}(u) \leq m_{\omega} - \delta$ and $\|\nabla u\|_{L^2}^2 \geq c_0$, then $\mathcal{K}(u) > \kappa_2(\delta)$. We prove this by contradiction. Hence, suppose to the contrary that there exists $\delta_0 > 0$ with

the following property: for any $n \in \mathbb{N}$, there exists a radial function $u_n \in H^1(\mathbb{R}^d)$ such that

$$\mathcal{J}_{\omega}(u_n) \le m_{\omega} - \delta_0, \tag{6.49}$$

$$\inf_{n\ge 1} \|\nabla u_n\|_{L^2} \ge c_0,\tag{6.50}$$

$$0 < \mathcal{K}(u_n) \le \frac{1}{n}.\tag{6.51}$$

Furthermore, it follows from (1.12), (6.49) and (6.51) that

$$\mathcal{S}_{\omega}(u_n) \le \mathcal{J}_{\omega}(u_n) + \frac{2}{n} \le m_{\omega} + \frac{2}{n} - \delta_0.$$
(6.52)

Now, using (1.14), (6.49) and (6.51), we find that for any $n \ge 1$,

$$\begin{aligned} \|u\|_{L^{2}}^{2} + \|\nabla u_{n}\|_{L^{2}}^{2} &\leq \mathcal{K}(u_{n}) + \|u\|_{L^{2}}^{2} + \|u\|_{L^{p+1}}^{p+1} + \|u\|_{L^{2^{*}}}^{2^{*}} \\ &\lesssim 1 + \mathcal{J}_{\omega}(u_{n}) \leq 1 + m_{\omega}. \end{aligned}$$
(6.53)

Hence, we can extract a subsequence of $\{u_n\}$ (still denoted by the same symbol $\{u_n\}$) and a radial function $u_{\infty} \in H^1(\mathbb{R}^d)$ such that

$$\lim_{n \to \infty} u_n = u_{\infty} \qquad \text{weakly in } H^1(\mathbb{R}^d) \tag{6.54}$$

and for any $2 < q < 2^*$,

$$\lim_{n \to \infty} u_n = u_{\infty} \qquad \text{strongly in } L^q(\mathbb{R}^d). \tag{6.55}$$

Then, we see from the same argument as the proof of Lemma 6.3 that u_{∞} is non-trivial. Furthermore, the lower semi-continuity of the weak limit together with (6.54), (6.55) and (6.49) implies that

$$\mathcal{J}_{\omega}(u_{\infty}) \le \liminf_{n \to \infty} \mathcal{J}_{\omega}(u_n) < m_{\omega}.$$
(6.56)

Hence, we find from (1.10) and $u_{\infty} \neq 0$ that $\mathcal{K}(u_{\infty}) > 0$. However, the same argument as the proof of Lemma 6.3 shows that $\mathcal{K}(u_{\infty}) \leq 0$. This is a contradiction. Thus, we have completed the proof.

7 One-pass theorem

In this section, we derive the one-pass theorem for our equation (NLS) (cf. Theorem 4.1 in [25]). To this end, we need the following inequalities for radial functions:

Lemma 7.1. The following hold for all M > 0 and all radial functions g in $H^1(\mathbb{R}^d)$: if d = 3 and $3 \le q \le 5 = 2^* - 1$, then

$$\int_{\mathbb{R}^3} \frac{M|x|}{(M+|x|)^2} |g(x)|^{q+1} \, dx \lesssim \frac{1}{M} \|g\|_{H^1}^{q-1} \left\| \nabla \left(\frac{M}{M+|x|}g\right) \right\|_{L^2}^2,\tag{7.1}$$

and if $d \ge 4$ and $1 + \frac{4}{d-1} \le q \le 2^* - 1$, then

$$\int_{\mathbb{R}^{d}} \frac{M|x|}{(M+|x|)^{2}} |g(x)|^{q+1} dx \lesssim M^{-\frac{(d-2)q-d}{4}} \|\nabla g\|_{L^{2}}^{q-1} \left\|\nabla \left(\frac{M}{M+|x|}g\right)\right\|_{L^{2}}^{2} + M^{-\frac{(d-1)(q-1)-4}{4}} \|g\|_{H^{1}}^{q-1} \int_{\mathbb{R}^{d}} \frac{M}{|x|(M+|x|)^{2}} |g(x)|^{2} dx.$$
(7.2)

Here, the implicit constants depend only on d and q.

Remark 7.1. We rely on the radial Sobolev inequalities to prove Lemma 7.1, which causes the restriction of p in the one-pass theorem (Theorem 7.1) and Theorem 1.1.

Proof of Lemma 7.1. First, we prove the inequality (7.1). We see from (C.1) and Sobolev's embedding that for any $3 \le q \le 5$,

$$\int_{\mathbb{R}^{3}} \frac{M|x|}{(M+|x|)^{2}} |g(x)|^{q+1} dx \leq \frac{1}{M} \sup_{x \in \mathbb{R}^{3}} |x| \left| \frac{M}{M+|x|} g(x) \right|^{2} \int_{\mathbb{R}^{3}} |g(x)|^{q-1} dx \\
\lesssim \frac{1}{M} \|g\|_{H^{1}}^{q-1} \left\| \nabla \left(\frac{M}{M+|x|} g \right) \right\|_{L^{2}}^{2}.$$
(7.3)

Next, we prove (7.2). We see from (C.1), Hölder's inequality and Hardy's inequality that for any $d \ge 4$ and $1 + \frac{4}{d} \le q \le 2^* - 1$ (hence $\frac{d}{d-2} \le q \le 3$),

$$\begin{split} &\int_{|x| \le \sqrt{M}} \frac{M|x|}{(M+|x|)^2} |g(x)|^{q+1} dx \\ \le \frac{1}{M} \left\| |x|^{\frac{d-2}{2}} \frac{M}{M+|x|} g \right\|_{L^{\infty}}^2 \int_{|x| \le \sqrt{M}} \frac{|g(x)|^{q-1}}{|x|^{d-3}} dx \\ \lesssim \frac{1}{M} \left\| \nabla \left(\frac{M}{M+|x|} g \right) \right\|_{L^2}^2 \left(\int_{|x| \le \sqrt{M}} |x|^{-\frac{2(d-2-q)}{3-q}} dx \right)^{\frac{3-q}{2}} \left(\int_{\mathbb{R}^d} \frac{|g|^2}{|x|^2} dx \right)^{\frac{q-1}{2}} \\ \lesssim M^{-\frac{(d-2)q-d}{4}} \left\| \nabla g \right\|_{L^2}^{q-1} \left\| \nabla \left(\frac{M}{M+|x|} g \right) \right\|_{L^2}^2. \end{split}$$
(7.4)

On the other hand, we see from (C.2) that for any $1 + \frac{4}{d-1} \le q \le 2^* - 1$,

$$\int_{\sqrt{M} \le |x|} \frac{M|x|}{(M+|x|)^2} |g(x)|^{q+1} dx
\le \left\| |x|^{d-1} |g|^2 \right\|_{L^{\infty}}^{\frac{q-1}{2}} \int_{\sqrt{M} \le |x|} \frac{|x|^{-\frac{(d-1)(q-1)}{2}+2}M}{|x|(M+|x|)^2} |g(x)|^2 dx
\lesssim M^{-\frac{(d-1)(q-1)-4}{4}} \|g\|_{H^1}^{q-1} \int_{\mathbb{R}^d} \frac{M}{|x|(M+|x|)^2} |g(x)|^2 dx.$$
(7.5)

Putting (7.4) and (7.5) together, we obtain the desired estimate (7.2).

Next, we recall that: ω_2 denotes the frequency given by Proposition 4.1; $\omega(2^*) \in (0, \omega_2)$ is the frequency satisfying (5.7); and for a given $\omega \in (0, \omega(2^*))$, δ_X , A_* , B_* and C_* denote the constants given by the ejection lemma (Lemma 5.2). Moreover, for given $\omega \in (0, \omega_2)$ and $\delta > 0$, $\varepsilon_0(\delta)$ denotes the constant determined by Lemma 6.3. Here, we may assume that

$$\delta_X \ll \mu |(\Phi_\omega, f_2)_{L^2}|. \tag{7.6}$$

Furthermore, we define $\delta_S > 0$ as a constant satisfying

$$\delta_S \le \frac{A_* \delta_X}{2B_* C_*}$$
 and $\delta_S < \delta_X$. (7.7)

Note that $\delta_S < \delta_X < \delta_0(\omega) < \delta_E(\omega)$ and $\delta_X < \tilde{\gamma}(\omega)$, where $\delta_E(\omega)$, $\delta_0(\omega)$ and $\tilde{\gamma}(\omega)$ are the constants given by Proposition 4.4, Lemma 5.1 and Proposition 6.2, respectively.

In the proof of the one-pass theorem (Theorem 7.1) below, we use the following sign function:

$$\operatorname{sign}[\alpha] = \begin{cases} 1 & \text{if } \alpha \ge 0, \\ -1 & \text{if } \alpha < 0. \end{cases}$$
(7.8)

In particular, sign[0] = 1.

Theorem 7.1 (One-pass theorem). Assume that either d = 3 and $3 \le p < 5$, or $d \ge 4$ and $1 + \frac{4}{d-1} . Then, there exists <math>\omega_* > 0$ such that for any $\omega \in (0, \omega_*)$, there exist positive constants ε_* , δ_* and R_* with the following properties:

$$\sqrt{\varepsilon_*} \ll R_* \ll \delta_* \ll \delta_S, \quad \varepsilon_* \le \varepsilon_0(\delta_*), \tag{7.9}$$

and for any $\varepsilon \in (0, \varepsilon_*]$, any $R \in (\sqrt{2\varepsilon}, R_*)$ and any radial solution ψ to (NLS) satisfying

$$\mathcal{M}(\psi) = \mathcal{M}(\Phi_{\omega}), \tag{7.10}$$

$$\mathcal{S}_{\omega}(\psi) < m_{\omega} + \varepsilon, \tag{7.11}$$

$$d_{\omega}(\psi(0)) < R, \tag{7.12}$$

we have either

(i)
$$\widetilde{d}_{\omega}(\psi(t)) < R + R^{\frac{\min\{3,p+1\}}{2}}$$
 for all $t \in [0, T_{\max}]$; or
(ii) there exists $t_* > 0$ such that $\widetilde{d}_{\omega}(\psi(t)) \ge R + R^{\frac{\min\{3,p+1\}}{2}}$ for all $t \in [t_*, T_{\max}]$.

Here, T_{max} denotes the maximal lifespan of ψ .

Proof of Theorem 7.1. We prove the claim by contradiction. Hence, we suppose to the contrary that for any $0 < \omega_* \ll 1$, there exists $0 < \omega < \omega_*$ such that for any $\varepsilon_*, \delta_*, R_* > 0$ with $\sqrt{\varepsilon_*} \ll R_* \ll \delta_* \ll \delta_S$ and $\varepsilon_* < \varepsilon_0(\delta_*)$, there exist $\varepsilon \in (0, \varepsilon_*]$, $R \in (\sqrt{2\varepsilon}, R_*)$ and a radial solution ψ such that for these ω , ε , R and ψ , the conditions (7.10), (7.11) and (7.12) hold, while both (i) and (ii) fail. Here, we may assume that δ_* is so small that

$$\log\left(\frac{\delta_X}{\delta_*}\right) \le \frac{1}{M_0} \frac{\delta_X}{\delta_*} \tag{7.13}$$

for some positive constant M_0 to be specified later dependently only on d, p and ω .

The failure of (i) together with (7.9) and (7.12) shows that there exist times $t_2 > t_1 > 0$ such that

$$\widetilde{d}_{\omega}(\psi(t_1)) = R < \delta_*, \tag{7.14}$$

$$\widetilde{d}_{\omega}(\psi(t_2)) = R + R^{\frac{\min\{3,p+1\}}{2}} < \delta_* \ll \delta_S < \delta_X,$$
(7.15)

$$R < \widetilde{d}_{\omega}(\psi(t)) < R + R^{\frac{\min\{3, p+1\}}{2}} < \delta_* \ll \delta_S < \delta_X \quad \text{for all } t \in (t_1, t_2).$$

$$(7.16)$$

Moreover, the failure of (ii) shows that there exists $t_3 > t_2$ such that

$$R < \widetilde{d}_{\omega}(\psi(t_3)) < R + R^{\frac{\min\{3, p+1\}}{2}},$$
(7.17)

$$\widetilde{d}_{\omega}(\psi(t_3)) < \widetilde{d}_{\omega}(\psi(t)) \quad \text{for all } t \in (t_2, t_3).$$
 (7.18)

We see from (7.14) through (7.18) that

$$R = \widetilde{d}_{\omega}(\psi(t_1)) = \min_{t \in [t_1, t_3]} \widetilde{d}_{\omega}(\psi(t)).$$
(7.19)

Note here that the ejection lemma (Lemma 5.2) together with (7.15) shows that

$$t_1 + T_* R^{\min\{1, p-1\}} < t_2, \tag{7.20}$$

where T_* is the constant given by the ejection lemma. Put

$$t'_{2} := \inf\{t \in [t_{1}, t_{3}] : \widetilde{d}_{\omega}(\psi(t)) = \delta_{X}\}.$$
(7.21)

Then, (7.16) shows $t_2 < t'_2$. Moreover, we see from (6.16) and the ejection lemma that

$$A_* e^{\mu(t-t_1)} R \le d_\omega(\psi(t)) \le B_* e^{\mu(t-t_1)} R, \tag{7.22}$$

$$\|\eta(t)\|_{H^1} \sim |\lambda_1(t)| \sim e^{\mu(t-t_1)} R \sim d_\omega(\psi(t)), \tag{7.23}$$

$$\|\Gamma(t)\|_{H^1} \lesssim R + \{e^{\mu(t-t_1)}R\}^{\frac{\min\{3,p+1\}}{2}} \sim \widetilde{d}_{\omega}(\psi(t_1)) + |\lambda_1(t)|^{\frac{\min\{3,p+1\}}{2}},$$
(7.24)

$$\operatorname{sign}[\lambda_1(t)]\mathcal{K}(\psi(t)) \gtrsim (e^{\mu(t-t_1)} - C_*)R \tag{7.25}$$

for all $t \in [t_1, t'_2]$, and $\tilde{d}_{\omega}(\psi(t))$ increases on $[t_1 + T_*R^{\min\{1, p-1\}}, t'_2]$. In particular, $\tilde{d}_{\omega}(\psi(t))$ must reach δ_X before t_3 and therefore

$$t_2 < t_2' < t_3. \tag{7.26}$$

Furthermore, we see from (7.16) that there exists $t_S \in (t_2, t'_2)$ such that $\widetilde{d}_{\omega}(\psi(t_S)) = \delta_S$. Now, note that (7.11) together with $\sqrt{2\varepsilon} < R$ and (7.19) shows that

$$\mathcal{S}_{\omega}(\psi) < m_{\omega} + \frac{1}{2}\widetilde{d}_{\omega}(\psi(t))^2 \quad \text{for all } t \in [t_1, t_3].$$
(7.27)

Hence, we find from Lemma 4.5, (6.16) and (7.19) that

$$R \le d_{\omega}(\psi(t)) \sim |\lambda_1(t)| \quad \text{for all } t \in [t_1, t_3] \text{ with } \widetilde{d}_{\omega}(\psi(t)) \le \delta_E(\omega).$$
(7.28)

In particular, sign[$\lambda_1(t)$] is constant on the set { $t \in [t_1, t_3]$: $\tilde{d}_{\omega}(\psi(t)) \leq \delta_E(\omega)$ }. Moreover, Lemma 6.3 together with (7.11) and $\varepsilon < \varepsilon_* < \varepsilon_0(\delta_*)$ shows that sign[$\mathcal{K}(\psi(t))$] is constant on the set { $t \in [t_1, t_3]$: $\tilde{d}_{\omega}(\psi(t)) \geq \delta_*$ }.

We shall show that for any $t \in [t_1, t_3]$ with $\delta_* \leq \tilde{d}_{\omega}(\psi(t)) \leq \delta_E(\omega)$,

$$\operatorname{sign}[\lambda_1(t)] = \operatorname{sign}[\mathcal{K}(\psi(t))].$$
(7.29)

Since both $\operatorname{sign}[\lambda_1(t)]$ and $\operatorname{sign}[\mathcal{K}(\psi(t))]$ are constant on the set $\{t \in [t_1, t_3]: \delta_* \leq \tilde{d}_{\omega}(\psi(t)) \leq \delta_E(\omega)\}$, it suffices to show that $\operatorname{sign}[\lambda_1(t'_2)] = \operatorname{sign}[\mathcal{K}(\psi(t'_2))]$. We see from (7.21), (7.22), (6.16) and (7.7) that

$$\delta_X \le B_* e^{\mu(t'_2 - t_1)} R \le B_* e^{\mu(t'_2 - t_1)} \frac{e^{-\mu(t_S - t_1)}}{A_*} \widetilde{d}_{\omega}(\psi(t_S))$$

$$= \frac{B_*}{A_*} e^{\mu(t'_2 - t_S)} \delta_S \le \frac{\delta_X}{2C_*} e^{\mu(t'_2 - t_S)}.$$
(7.30)

Dividing the both sides of (7.30) by δ_X , and taking the logarithm, we obtain

$$\frac{1}{\mu} \log C_* < t_2' - t_S. \tag{7.31}$$

Since $t_1 < t_S$, we find from (7.25) and (7.31) that $\operatorname{sign}[\lambda_1(t'_2)] = \operatorname{sign}[\mathcal{K}(\psi(t'_2))]$. Hence, (7.29) holds.

We find from (7.29) that the following function $\mathfrak{S}: [t_1, t_3] \to \{1, -1\}$ is well-defined:

$$\mathfrak{S}(t) := \begin{cases} \operatorname{sign}[\lambda_1(t)] & \text{if } \widetilde{d}_{\omega}(\psi(t)) \leq \delta_E(\omega), \\ \operatorname{sign}[\mathcal{K}(\psi(t))] & \text{if } \widetilde{d}_{\omega}(\psi(t)) \geq \delta_*. \end{cases}$$
(7.32)

Note that the function $\mathfrak{S}(t)$ is constant on $[t_1, t_3]$. Put $\mathfrak{s} := \mathfrak{S}(t) \in \{1, -1\}$. When $\mathfrak{s} = 1$, we see that there exists $C(\omega) > 0$ such that for any $t \in [t_1, t_3]$,

$$\sup_{t \in [t_1, t_3]} \|\psi(t)\|_{H^1} \le C(\omega).$$
(7.33)

Indeed, if $\widetilde{d}_{\omega}(\psi(\tau)) \geq \delta_S$ for some $\tau \in [t_1, t_3]$, then (7.32) shows $\inf_{t \in [t_1, t_3]} \mathcal{K}(\psi(t)) \geq 0$. Hence, we see from (1.11), (1.13) and the assumption (7.11) that for any $t \in [t_1, t_3]$,

$$\frac{\omega}{2} \|\psi(t)\|_{L^2}^2 + \frac{s_p}{d} \|\nabla\psi(t)\|_{L^2}^2 \le \mathcal{I}_{\omega}(\psi(t)) \le \mathcal{S}_{\omega}(\psi) \le m_{\omega} + 1.$$
(7.34)

On the other hand, if $\tilde{d}_{\omega}(\psi(t)) \leq \delta_S$ for all $t \in [t_1, t_3]$, then we see from Lemma 4.3, Lemma 4.5 and (6.16) that for any $t \in [t_1, t_3]$,

$$\|\psi(t)\|_{H^1}^2 \lesssim \|\Phi_{\omega}\|_{H^1}^2 + \|\eta(t)\|_{H^1}^2 \lesssim \|\Phi_{\omega}\|_{H^1}^2 + d_{\omega}(\psi(t))^2 \lesssim \|\Phi_{\omega}\|_{H^1}^2 + \delta_S.$$
(7.35)

Thus, we have verified that (7.33) holds.

Let us summarize information obtained so far: there exist $t'_1(=t_1+T_*R^{\min\{1,p-1\}}) \in (t_1,t_2)$ and $t'_2 \in (t_2,t_3)$ such that: $\tilde{d}_{\omega}(\psi(t))$ increases on $[t'_1,t'_2]$; $\tilde{d}_{\omega}(\psi(t'_2)) = \delta_X$; and for any $t \in (t_1,t'_2)$,

$$\delta_X \ge \widetilde{d}_{\omega}(\psi(t)) \sim \|\eta(t)\|_{H^1} \sim |\lambda_1(t)| \sim e^{\mu(t-t_1)}R, \tag{7.36}$$

$$\|\Gamma(t)\|_{H^1} \lesssim \widetilde{d}_{\omega}(\psi(t_1)) + |\lambda_1(t)|^{\frac{\min\{3,p+1\}}{2}},$$
(7.37)

$$\mathfrak{s}\mathcal{K}(\psi(t)) \gtrsim (e^{\mu(t-t_1)} - C_*)R. \tag{7.38}$$

Arguing from t_3 backward in time, we are also able to obtain a time interval $(t''_2, t_3) \subset (t'_2, t_3)$ such that: $\tilde{d}_{\omega}(\psi(t''_2)) = \delta_X$; for any $t \in (t''_2, t_3)$,

$$\delta_X \ge \widetilde{d}_{\omega}(\psi(t)) \sim \|\eta(t)\|_{H^1} \sim |\lambda_1(t)| \sim e^{\mu(t_3 - t)} \widetilde{d}_{\omega}(\psi(t_3)), \tag{7.39}$$

$$\|\Gamma(t)\|_{H^1} \lesssim \tilde{d}_{\omega}(\psi(t_3)) + |\lambda_1(t)|^{\frac{\min\{3,p+1\}}{2}},$$
(7.40)

$$\mathfrak{s}\mathcal{K}(\psi(t)) \gtrsim (e^{\mu(t_3-t)} - C_*)\widetilde{d}_{\omega}(\psi(t_3)), \tag{7.41}$$

and $\widetilde{d}_{\omega}(\psi(t))$ decreases at least in the region $\{t \in [t_2'', t_3]: \widetilde{d}_{\omega}(\psi(t)) \geq 2\widetilde{d}_{\omega}(\psi(t_3))\}$ (cf. (5.17) in the ejection lemma (Lemma 5.2)).

Suppose here that there exists a time $\tau \in (t'_2, t''_2)$ such that $\tilde{d}_{\omega}(\psi(\tau))$ is a local minimum and $\tilde{d}_{\omega}(\psi(\tau)) < \delta_*$. Then, we can apply the ejection lemma (Lemma 5.2) from τ both forward and backward in time to obtain an open interval $I_{\tau} \subset (t'_2, t''_2)$ such that

$$\widetilde{d}_{\omega}(\psi(\inf I_{\tau})) = \widetilde{d}_{\omega}(\psi(\sup I_{\tau})) = \delta_X, \qquad (7.42)$$

and for any $t \in I_{\tau}$,

$$\delta_X \ge \widetilde{d}_{\omega}(\psi(t)) \sim \|\eta(t)\|_{H^1} \sim |\lambda_1(t)| \sim e^{\mu|t-\tau|} \widetilde{d}_{\omega}(\psi(\tau)), \tag{7.43}$$

$$\|\Gamma(t)\|_{H^1} \lesssim \tilde{d}_{\omega}(\psi(\tau)) + |\lambda_1(t)|^{\frac{\min\{3,p+1\}}{2}},\tag{7.44}$$

$$\mathfrak{s}\mathcal{K}(\psi(t)) \gtrsim (e^{\mu|t-\tau|} - C_*)\widetilde{d}_{\omega}(\psi(\tau)), \tag{7.45}$$

and $\tilde{d}_{\omega}(\psi(t))$ is monotone in the region $\tilde{d}_{\omega}(\psi(t)) \geq 2\tilde{d}_{\omega}(\psi(\tau))$. Note that for any distinct local minimum points τ_1 and τ_2 of $\tilde{d}_{\omega}(\psi(t))$ in (t'_2, t''_2) , the monotonicity away from them implies that the intervals I_{τ_1} and I_{τ_2} are either disjoint or identical.



Figure 3: Behavior of $\tilde{d}_{\omega}(\psi(t))$ in the case where there exists a local minimum point $\tau \in (t'_2, t''_2)$ such that $\tilde{d}_{\omega}(\psi(\tau)) < \delta_*$.

We find from the above observation that there exist a number $n \ge 2$ and disjoint open subintervals I_1, \ldots, I_n of $[t_1, t_3]$ with the following properties²: • inf $I_1 = t_1$, sup $I_n = t_3$, $\tilde{d}_{\omega}(\psi(\sup I_1)) = \tilde{d}_{\omega}(\psi(\inf I_n)) = \delta_X$, and

$$\widetilde{d}_{\omega}(\psi(\inf I_j)) = \widetilde{d}_{\omega}(\psi(\sup I_j)) = \delta_X \quad \text{except for } j = 1, n.$$
(7.46)

• For each $1 \leq j \leq n$, there exists $\tau_j \in I_j$ such that

$$\widetilde{d}_{\omega}(\psi(\tau_j)) < \delta_*, \tag{7.47}$$

and for any $t \in I_j$,

$$\delta_X \ge \widetilde{d}_{\omega}(\psi(t)) \sim \|\eta(t)\|_{H^1} \sim |\lambda_1(t)| \sim e^{\mu|t-\tau_j|} \widetilde{d}_{\omega}(\psi(\tau_j)), \tag{7.48}$$

$$\|\Gamma(t)\|_{H^1} \lesssim \widetilde{d}_{\omega}(\psi(\tau_j)) + |\lambda_1(t)|^{\frac{\min\{3,p+1\}}{2}},\tag{7.49}$$

$$\mathfrak{s}\mathcal{K}(\psi(t)) \gtrsim (e^{\mu|t-\tau_j|} - C_*)\widetilde{d}_{\omega}(\psi(\tau_j)).$$
(7.50)

• If t satisfies $\sup I_{j-1} < t < \inf I_j$ for some $2 \le j \le n$, then

$$\widetilde{d}_{\omega}(\psi(t)) > \delta_*. \tag{7.51}$$

²Note that $I_1 = (t_1, t'_2)$. Moreover, if $\widetilde{d}_{\omega}(\psi(t)) \ge \delta_*$ on $[t'_2, t''_2]$, then n = 2 and $I_2 = (t''_2, t_3)$.

Put $I' := [t_1, t_3] \setminus \bigcup_{j=1}^n I_j$. Then, for any $t \in I'$, we have $\widetilde{d}_{\omega}(\psi(t)) > \delta_*$ (see (7.51)), which together with Lemma 6.3 gives us that for any $t \in I'$,

$$\begin{cases} \mathcal{K}(\psi(t)) \ge \min\left\{\kappa_1(\delta_*), \frac{1}{2} \|\nabla\psi(t)\|_{L^2}^2\right\} & \text{if} \quad \mathfrak{s} = 1, \\ \mathcal{K}(\psi(t)) \le -\kappa_1(\delta_*) & \text{if} \quad \mathfrak{s} = -1. \end{cases}$$
(7.52)

We can find from elementary calculations, (7.48), $\tilde{d}_{\omega}(\psi(\tau_j)) \leq \delta_*$ and (7.13) that for any $1 \leq j \leq n$,

$$\mu |I_j| \widetilde{d}_{\omega}(\psi(\tau_j)) = \left\{ \log e^{\mu |\sup I_j - \tau_j|} + \log e^{\mu |\inf I_j - \tau_j|} \right\} \widetilde{d}_{\omega}(\psi(\tau_j))$$

$$\lesssim \log \left(\frac{\delta_X}{\widetilde{d}_{\omega}(\psi(\tau_j))} \right) \widetilde{d}_{\omega}(\psi(\tau_j)) \le \frac{1}{M_0} \delta_X.$$
(7.53)

We shall derive a contradiction in the case where $\mathfrak{s} = 1$. To this end, we use the following identity: for any $t \in I_{\max}(\psi)$ and any M > 0,

$$\frac{d}{dt} \Im \int_{\mathbb{R}^{d}} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla \psi(x,t) \overline{\psi(x,t)} \, dx$$

$$= 2\mathcal{K} \left(\frac{M}{M+|x|} \psi(t) \right)$$

$$- \frac{p-1}{p+1} \int_{\mathbb{R}^{d}} \left\{ \frac{M^{2}}{(M+|x|)^{2}} + \frac{(d-1)M}{M+|x|} - \frac{dM^{p+1}}{(M+|x|)^{p+1}} \right\} |\psi(t)|^{p+1} \, dx$$

$$- \frac{2}{d} \int_{\mathbb{R}^{d}} \left\{ \frac{M^{2}}{(M+|x|)^{2}} + \frac{(d-1)M}{M+|x|} - \frac{dM^{2^{*}}}{(M+|x|)^{2^{*}}} \right\} |\psi(t)|^{2^{*}} \, dx$$

$$+ \frac{1}{2} \int_{\mathbb{R}^{d}} \left\{ \frac{2M^{2}}{(M+|x|)^{4}} + \frac{(d-3)(d-1)M}{|x|(M+|x|)^{2}} \right\} |\psi(t)|^{2} \, dx.$$
(7.54)

Here, note that an elementary computation shows that for any $d \ge 3$ and any $1 \le q \le 2^* - 1$,

$$\int_{\mathbb{R}^{d}} \left\{ \frac{M^{2}}{(M+|x|)^{2}} + \frac{(d-1)M}{M+|x|} - \frac{dM^{q+1}}{(M+|x|)^{q+1}} \right\} |\psi(t)|^{q+1} dx \\
= \int_{\mathbb{R}^{d}} \left\{ \frac{dM}{M+|x|} \left(\frac{M}{M+|x|} - \frac{M^{q}}{(M+|x|)^{q}} \right) + \frac{(d-1)M|x|}{(M+|x|)^{2}} \right\} |\psi(t)|^{q+1} dx \quad (7.55) \\
\lesssim \int_{\mathbb{R}^{d}} \frac{M|x|}{(M+|x|)^{2}} |\psi(t)|^{q+1} dx.$$

We shall derive a contradiction in three steps:

Step 1. We shall derive an estimate for (7.54) on the region $\bigcup_{j=1}^{n} I_j$.

We see from Sobolev's embedding that for any $d \ge 3$, any $1 + \frac{4}{d} \le q \le 2^* - 1$, any M > 0 and any $t \in I_{\max}(\psi)$,

$$\int_{\mathbb{R}^{d}} \frac{M|x|}{(M+|x|)^{2}} |\psi(t)|^{q+1} dx \lesssim \frac{1}{M} \int_{\mathbb{R}^{d}} |x| |\Phi_{\omega}(x)|^{q+1} dx + \|\eta(t)\|_{L^{q+1}}^{q+1} \\ \lesssim \frac{C(\omega)}{M} + \|\eta(t)\|_{H^{1}}^{q+1},$$
(7.56)

where $C(\omega)$ is some positive constant depending only on d, p and ω . Furthermore, it follows from (7.54), (7.55) and (7.56) that for any $d \geq 3$, any $1 + \frac{4}{d} \leq p < 2^* - 1$, any M > 0 and any $t \in I_{\max}(\psi)$, then

$$\frac{d}{dt} \Im \int_{\mathbb{R}^d} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla \psi(x,t) \overline{\psi(x,t)} \, dx$$

$$\geq 2\mathcal{K} \left(\frac{M}{M+|x|} \psi(t)\right) - \frac{C(\omega)}{M} - C\left\{ \|\eta(t)\|_{H^1}^{p+1} + \|\eta(t)\|_{H^1}^{2^*} \right\}$$
(7.57)

where $C(\omega)$ is some positive constant depending only on d, p and ω , and C is some positive constant depending only on d and p.

We consider the first two terms on the right-hand side of (7.57). Note first that it follows from (4.150) that for any $u \in H^1(\mathbb{R}^d)$,

$$\mathcal{K}'(\Phi_{\omega})u = -s_p \langle (p-1)\Phi_{\omega}^p, u \rangle_{H^{-1}, H^1} - (2^* - 2) \langle \Phi_{\omega}^{2^* - 1}, u \rangle_{H^{-1}, H^1} - 2\omega \langle \Phi_{\omega}, u \rangle_{H^{-1}, H^1}.$$
(7.58)

Furthermore, the first order Taylor's expansion of \mathcal{K} around Φ_{ω} together with (7.58) shows that

$$\begin{aligned} \mathcal{K}\Big(\frac{M}{M+|x|}\psi(t)\Big) &= \mathcal{K}\Big(\Phi_{\omega} + \eta(t) - \frac{|x|}{M+|x|} \{\Phi_{\omega} + \eta(t)\}\Big) \\ &= -s_p \langle (p-1)\Phi_{\omega}^p, \eta(t) \rangle_{H^{-1},H^1} + s_p \langle (p-1)\Phi_{\omega}^p, \frac{|x|}{M+|x|} \{\Phi_{\omega} + \eta(t)\} \rangle_{H^{-1},H^1} \\ &- (2^*-2) \langle \Phi_{\omega}^{2^*-1}, \eta(t) \rangle_{H^{-1},H^1} + (2^*-2) \langle \Phi_{\omega}^{2^*-1}, \frac{|x|}{M+|x|} \{\Phi_{\omega} + \eta(t)\} \rangle_{H^{-1},H^1} \quad (7.59) \\ &- 2\omega \langle \Phi_{\omega}, \eta(t) \rangle_{H^{-1},H^1} + 2\omega \langle \Phi_{\omega}, \frac{|x|}{M+|x|} \{\Phi_{\omega} + \eta(t)\} \rangle_{H^{-1},H^1} \\ &+ O\left(\left\|\frac{M}{M+|x|}\eta(t) - \frac{|x|}{M+|x|}\Phi_{\omega}\right\|_{H^1}^2\right). \end{aligned}$$

Here, we see from the decomposition (4.108), (4.7), (4.109), (4.121) and $\mathfrak{s} = 1$ (hence $\lambda_1(t) \ge 0$ by (7.32)) that the first and third terms on the right-hand side of (7.59) are rewritten as follows:

$$- s_{p} \langle (p-1)\Phi_{\omega}^{p}, \eta(t) \rangle_{H^{-1}, H^{1}} - (2^{*}-2) \langle \Phi_{\omega}^{2^{*}-1}, \eta(t) \rangle_{H^{-1}, H^{1}} \\ = -s_{p} \langle (p-1)\Phi_{\omega}^{p}, 2\lambda_{1}(t)f_{1} \rangle_{H^{-1}, H^{1}} - s_{p}(p-1) \langle \Phi_{\omega}^{p}, \Gamma(t) \rangle_{H^{-1}, H^{1}} \\ - (2^{*}-2) \langle \Phi_{\omega}^{2^{*}-1}, 2\lambda_{1}(t)f_{1} \rangle_{H^{-1}, H^{1}} - (2^{*}-2) \langle \Phi_{\omega}^{2^{*}-1}, \Gamma(t) \rangle_{H^{-1}, H^{1}} \\ = 2s_{p}\lambda_{1}(t) \langle L_{\omega, +}\Phi_{\omega}, f_{1} \rangle_{H^{-1}, H^{1}} - s_{p}(p-1) \langle \Phi_{\omega}^{p}, \Gamma(t) \rangle_{H^{-1}, H^{1}} \\ - 2(1-s_{p})(2^{*}-2)\lambda_{1}(t) \langle \Phi_{\omega}^{2^{*}-1}, f_{1} \rangle_{H^{-1}, H^{1}} - (2^{*}-2) \langle \Phi_{\omega}^{2^{*}-1}, \Gamma(t) \rangle_{H^{-1}, H^{1}} \\ = 2s_{p}\mu |\lambda_{1}(t)| |(\Phi_{\omega}, f_{2})_{L^{2}}| - s_{p}(p-1) \langle \Phi_{\omega}^{p}, \Gamma(t) \rangle_{H^{-1}, H^{1}} \\ - 2(1-s_{p})(2^{*}-2) |\lambda_{1}(t)| \langle \Phi_{\omega}^{2^{*}-1}, f_{1} \rangle_{H^{-1}, H^{1}} - (2^{*}-2) \langle \Phi_{\omega}^{2^{*}-1}, \Gamma(t) \rangle_{H^{-1}, H^{1}}. \end{cases}$$
(7.60)

We also see from the decomposition (4.108), Lemma 4.7 and (4.149) that

$$-2\omega \langle \Phi_{\omega}, \eta(t) \rangle_{H^{-1}, H^{1}} = -2\omega \langle \Phi_{\omega}, \Gamma(t) \rangle_{H^{-1}, H^{1}} = \omega \| \eta(t) \|_{L^{2}}^{2}.$$
(7.61)

Moreover, it follows from Hölder's inequality, Sobolev's embedding and (2.9) that for any $p \leq q \leq 2^* - 1$ and any sufficiently small $\omega > 0$,

$$\left| (\Phi_{\omega}^{q}, \Gamma(t))_{L^{2}} \right| \leq \|\Phi_{\omega}\|_{L^{q+1}}^{q} \|\Gamma(t)\|_{L^{q+1}} \lesssim \omega^{\frac{q}{q+1}(\frac{q-1}{p-1}-s_{p})} \|\Gamma(t)\|_{H^{1}} \leq \omega^{\frac{1-s_{p}}{2}} \|\Gamma(t)\|_{H^{1}}, \quad (7.62)$$

where the implicit constant depends only on d and q; it follows from Hölder's inequality and Sobolev's embedding that for any $1 \le q \le 2^* - 1$ and any sufficiently small $\omega > 0$,

$$\langle \Phi_{\omega}^{q}, \frac{|x|}{M+|x|} \{ \Phi_{\omega} + \eta(t) \} \rangle_{H^{-1},H^{1}} \geq -\int_{\mathbb{R}^{d}} \frac{|x|}{M+|x|} \Phi_{\omega}^{q} |\eta(t)| \, dx$$

$$\geq -\frac{1}{M} \bigg(\int_{\mathbb{R}^{d}} |x|^{\frac{q+1}{q}} \Phi_{\omega}^{q+1} \, dx \bigg)^{\frac{q}{q+1}} \|\eta(t)\|_{L^{q+1}} \geq -\frac{1}{M} G(\omega) \|\eta(t)\|_{H^{1}},$$

$$(7.63)$$

where $G(\omega)$ is some positive constant depending only on d, q and ω ; and it follows from Sobolev's embedding that

$$\left\|\frac{M}{M+|x|}\eta(t) - \frac{|x|}{M+|x|}\Phi_{\omega}\right\|_{H^{1}}^{2} \lesssim \|\eta(t)\|_{H^{1}}^{2} + \frac{1}{M}G(\omega),$$
(7.64)

where $G(\omega)$ is some positive constant depending only on d, q and ω .

Plugging (7.60) and (7.61) into (7.59), and then using Lemma 4.8 with $C = s_p - 2(1-s_p)(2^*-2)$, (7.62), (7.63) and (7.64), we find that if ω_* is sufficiently small (hence so is ω), then

$$\begin{aligned} &\mathcal{K}\Big(\frac{M}{M+|x|}\psi(t)\Big) - \frac{C(\omega)}{M} \\ &= 2s_p\mu|\lambda_1(t)||(\Phi_{\omega}, f_2)_{L^2}| - s_p(p-1)\langle\Phi_{\omega}^p, \Gamma(t)\rangle_{H^{-1}, H^1} \\ &- 2(1-s_p)(2^*-2)|\lambda_1(t)|(\Phi_{\omega}^{2^*-1}, f_1)_{L^2} - (2^*-2)\langle\Phi_{\omega}^{2^*-1}, \Gamma(t)\rangle_{H^{-1}, H^1} \\ &+ s_p\langle(p-1)\Phi_{\omega}^p, \frac{|x|}{M+|x|} \{\Phi_{\omega} + \eta(t)\}\rangle_{H^{-1}, H^1} \\ &+ (2^*-2)\langle\Phi_{\omega}^{2^*-1}, \frac{|x|}{M+|x|} \{\Phi_{\omega} + \eta(t)\}\rangle_{H^{-1}, H^1} \\ &+ (2^*-2)\langle\Phi_{\omega}^{2^*-1}, \frac{|x|}{M+|x|} \{\Phi_{\omega} + \eta(t)\}\rangle_{H^{-1}, H^1} \\ &+ O\left(\left\|\frac{M}{M+|x|}\eta(t) - \frac{|x|}{M+|x|}\Phi_{\omega}\right\|_{H^1}^2\right) - \frac{C(\omega)}{M} \\ &\geq s_p\mu|\lambda_1(t)||(\Phi_{\omega}, f_2)_{L^2}| - C_1\omega^{\frac{1-s_p}{2}}\|\Gamma(t)\|_{H^1} - \frac{1}{M}G_1(\omega)\|\eta(t)\|_{H^1} \\ &- C_2\|\eta(t)\|_{H^1}^2 - \frac{1}{M}G_2(\omega), \end{aligned}$$

where C_1 and C_2 are some positive constants depending only on d and p, and $G_1(\omega)$ and $G_2(\omega)$ are some positive constants depending only on d, q and ω . Recall here that (7.48), (7.49) and (7.6) give us that for any $t \in I_j$,

$$\|\eta(t)\|_{H^1} \sim |\lambda_1(t)| \sim e^{\mu|t-\tau_j|} \widetilde{d}_{\omega}(\psi(\tau_j)), \qquad (7.66)$$

$$\|\eta(t)\|_{H^1}^2 \lesssim \delta_X |\lambda_1(t)| \ll \mu |(\Phi_\omega, f_2)_{L^2}||\lambda_1(t)|,$$
(7.67)

$$\|\Gamma(t)\|_{H^1} \lesssim \widetilde{d}_{\omega}(\psi(\tau_j)). \tag{7.68}$$

Moreover, it follows from (7.19) that if $M \ge G_2(\omega)/R$, then

$$\frac{G_2(\omega)}{M} \le R \le \tilde{d}_{\omega}(\psi(\tau_j)).$$
(7.69)

Hence, (7.65) together with (7.66) through (7.69) shows that for any sufficiently small $\omega > 0$, any $M \ge G_2(\omega)/R$ and any $t \in I_j$,

$$\mathcal{K}\left(\frac{M}{M+|x|}\psi(t)\right) - \frac{C(\omega)}{M} \ge 2C_0^*(\omega)\left\{e^{\mu|t-\tau_j|} - C_1^*(\omega)\right\}\widetilde{d}_{\omega}(\psi(\tau_j)),\tag{7.70}$$

where $C_0^*(\omega)$ and $C_1^*(\omega)$ are some positive constant depending only on d, p and ω .

We find from (7.57), (7.70), $\|\eta(t)\|_{H^1} \lesssim \delta_X$ (see (7.48)) and (7.66) that if we take δ_X sufficiently small dependently only on d, p and ω in advance, then for any $M \ge G_2(\omega)/R$ and any $t \in I_j$,

$$\frac{d}{dt} \Im \int_{\mathbb{R}^d} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla \psi(x,t) \overline{\psi(x,t)} \, dx$$

$$\geq 2C_0^*(\omega) \{ e^{\mu|t-\tau_j|} - C_1^*(\omega) \} \widetilde{d}_{\omega}(\psi(\tau_j)) - C \delta_X^{p-1} e^{\mu|t-\tau_j|} \widetilde{d}_{\omega}(\psi(\tau_j)) \qquad (7.71)$$

$$\geq C_0^*(\omega) \{ e^{\mu|t-\tau_j|} - C_1^*(\omega) \} \widetilde{d}_{\omega}(\psi(\tau_j)).$$

Step 2. Next, we consider the "variational region" I'. We have $\tilde{d}_{\omega}(\psi) \geq \delta_*$ on I'. Moreover, it follows from (7.54), (7.55), Lemma 7.1 and (7.33) that if $d \geq 3$ and $3 \leq p < 5$, or $d \geq 4$ and $1 + \frac{4}{d-1} , then for any sufficiently large <math>M > 0$ depending only on d, p and ω , we have

$$\frac{d}{dt} \Im \int_{\mathbb{R}^d} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla \psi(x,t) \overline{\psi(x,t)} \, dx$$

$$\geq 2\mathcal{K} \Big(\frac{M}{M+|x|} \psi(t) \Big) - o_M(1) \Big\| \nabla \Big(\frac{M}{M+|x|} \psi(t) \Big) \Big\|_{L^2}^2.$$
(7.72)

Suppose here that $\|\nabla \psi(t)\|_{L^2} \leq \alpha$ for some $\alpha > 0$. Then, it follows from Hölder's inequality, $\frac{d(p-1)}{2} > 2$, Sobolev's embedding and the assumption (7.10) that

$$\begin{aligned} \left\| \frac{M}{M+|x|} \psi(t) \right\|_{L^{p+1}}^{p+1} &\leq \left\| \frac{M}{M+|x|} \psi(t) \right\|_{L^{2}}^{p+1-\frac{d(p-1)}{2}} \left\| \frac{M}{M+|x|} \psi(t) \right\|_{L^{2^{*}}}^{\frac{d(p-1)}{2}} \\ &\leq \left\| \psi(t) \right\|_{L^{2}}^{p+1-\frac{d(p-1)}{2}} \left\| \psi(t) \right\|_{L^{2^{*}}}^{\frac{d(p-1)}{2}-2} \left\| \frac{M}{M+|x|} \psi(t) \right\|_{L^{2^{*}}}^{2} \\ &\lesssim \mathcal{M}(\Phi_{\omega})^{\frac{p+1}{2}-\frac{d(p-1)}{4}} \alpha^{\frac{d(p-1)}{2}-2} \left\| \nabla \left(\frac{M}{M+|x|} \psi(t) \right) \right\|_{L^{2}}^{2}. \end{aligned}$$
(7.73)

Similarly, we have

$$\left\|\frac{M}{M+|x|}\psi(t)\right\|_{L^{2^{*}}}^{2^{*}} \lesssim \alpha^{2^{*}-2} \left\|\nabla\left(\frac{M}{M+|x|}\psi(t)\right)\right\|_{L^{2}}^{2}.$$
(7.74)

Furthermore, we find from (7.73) and (7.74) that for a given $\omega > 0$ there exist $\alpha_0 > 0$ and $\kappa_0 > 0$ such that for any $t \in I_{\max}(\psi)$ with $\|\nabla \psi(t)\|_{L^2} \leq \alpha_0$,

$$\mathcal{K}\left(\frac{M}{M+|x|}\psi(t)\right) \ge \kappa_0 \left\|\nabla\left(\frac{M}{M+|x|}\psi(t)\right)\right\|_{L^2}^2.$$
(7.75)

On the other hand, we see from (7.52) that for any $t \in I'$ with $\|\nabla \psi(t)\|_{L^2} > \alpha_0$,

$$\mathcal{K}(\psi(t)) \ge \min\left\{\kappa_1(\delta_*), \, \frac{1}{2}\alpha_0^2\right\}.$$
(7.76)

Choosing $\varepsilon_* > 0$ so small that $\varepsilon_*^2 < \frac{1}{4} \min\{\kappa_1(\delta_*), \frac{1}{2}\alpha_0^2\}$, and using (7.11) and (7.76), we obtain that for any $t \in I'$ with $\|\nabla \psi(t)\|_{L^2} > \alpha_0$,

$$\mathcal{J}_{\omega}\left(\frac{M}{M+|x|}\psi(t)\right) \le \mathcal{J}_{\omega}(\psi(t)) = \mathcal{S}_{\omega}(\psi) - \frac{1}{2}\mathcal{K}(\psi(t)) < m_{\omega} - \varepsilon_*^2, \qquad (7.77)$$

which together with Lemma 6.4 gives us that

$$\mathcal{K}\left(\frac{M}{M+|x|}\psi(t)\right) \ge \min\{\kappa_2(\varepsilon_*^2), \ \frac{1}{2} \left\|\nabla\left(\frac{M}{M+|x|}\psi(t)\right)\right\|_{L^2}^2\}.$$
(7.78)

Putting (7.75) and (7.78) together, we find that for any $t \in I'$,

$$\mathcal{K}\left(\frac{M}{M+|x|}\psi(t)\right) \ge \min\{\kappa_2(\varepsilon_*^2), \ \kappa_0 \left\|\nabla\left(\frac{M}{M+|x|}\psi(t)\right)\right\|_{L^2}^2\}$$
(7.79)

for some constant $\kappa_0 > 0$ depending only on d, p and ω . Note here that (7.33) shows

$$\left\|\nabla\left(\frac{M}{M+|x|}\psi(t)\right)\right\|_{L^{2}} \le \left\|\frac{M}{(M+|x|)^{2}}\psi(t)\right\|_{L^{2}} + \left\|\frac{M}{M+|x|}\nabla\psi(t)\right\|_{L^{2}} \lesssim C(\omega)$$
(7.80)

for all $t \in [t_1, t_3]$. Furthermore, we find from (7.72), (7.79) and (7.80) that if M satisfies $o_M(1) \ll \min\{\kappa_2(\varepsilon_*^2)/C(\omega)^2, \kappa_0\}$ and $t \in I'$, then

$$\frac{d}{dt}\Im \int_{\mathbb{R}^d} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla \psi(x,t) \overline{\psi(x,t)} \, dx \ge 0.$$
(7.81)

Step 3. We finish the proof by deriving a contradiction.

We see from (7.71) and (7.81) that for any sufficiently small $\omega > 0$, there exists

 $L(\omega) \gg 1$ such that if $M \ge L(\omega)/R$, then

$$\left[\Im\int_{\mathbb{R}^{d}} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla\psi(x,t)\overline{\psi(x,t)} \, dx\right]_{t_{1}}^{t_{3}}$$

$$= \int_{t_{1}}^{t_{3}} \frac{d}{dt}\Im\int_{\mathbb{R}^{d}} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla\psi(x,t)\overline{\psi(x,t)} \, dx \, dt$$

$$\geq C_{0}^{*}(\omega) \sum_{j=1}^{n} \int_{I_{j}} \left\{ e^{\mu|t-\tau_{j}|} - C_{1}^{*}(\omega) \right\} \widetilde{d}_{\omega}(\psi(\tau_{j})) \, dt \qquad (7.82)$$

$$= C_{0}^{*}(\omega) \sum_{j=1}^{n} \frac{1}{\mu} \left\{ e^{\mu|\sup I_{j}-\tau_{j}|} + e^{\mu|\inf I_{j}-\tau_{j}|} - 2 \right\} \widetilde{d}_{\omega}(\psi(\tau_{j}))$$

$$- C_{0}^{*}(\omega) C_{1}^{*}(\omega) \sum_{j=1}^{n} |I_{j}| \widetilde{d}_{\omega}(\psi(\tau_{j})).$$

Here, it follows from (7.46) and (7.48) that for any $1 \le j \le n$,

$$\delta_X = \widetilde{d}_{\omega}(\psi(\sup I_j)) \sim e^{\mu |\sup I_j - \tau_j|} \widetilde{d}_{\omega}(\psi(\tau_j)) \quad \text{except for } j = n,$$
(7.83)

$$\delta_X = \widetilde{d}_{\omega}(\psi(\inf I_j)) \sim e^{\mu |\inf I_j - \tau_j|} \widetilde{d}_{\omega}(\psi(\tau_j)) \quad \text{except for } j = 1.$$
(7.84)

Moreover, we see from (7.53) and $\tilde{d}_{\omega}(\psi(\tau_j)) \geq R$ (see (7.19)) that if M_0 is sufficiently large dependently only on d, p and ω ,

$$\mu C_1^*(\omega) |I_j| \widetilde{d}_{\omega}(\psi(\tau_j)) \lesssim C_1^*(\omega) \frac{1}{M_0} \delta_X \ll \delta_X.$$
(7.85)

Thus, (7.82) together with (7.83), (7.84), $\tilde{d}_{\omega}(\psi(\tau_j)) \leq \delta_* \ll \delta_X$ and (7.85) shows that for any sufficiently large M > 0 depending only on d, p, ω and R,

$$\left[\Im \int_{\mathbb{R}^d} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla \psi(x,t) \overline{\psi(x,t)}\right]_{t_1}^{t_3}$$

$$\geq 2C^{**}(\omega) \sum_{j=1}^n \left\{\delta_X - \delta_*\right\} - C^{**}(\omega) \sum_{j=1}^n \delta_X \ge \frac{1}{2} C^{**}(\omega) n \delta_X \ge C^{**}(\omega) \delta_X$$
(7.86)

for some positive constant $C^{**}(\omega)$ depending only on d, p and ω . Note here that

$$\left[\Im \int_{\mathbb{R}^d} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla \Phi_\omega(x) \Phi_\omega(x) dx\right]_{t_1}^{t_3} = 0.$$
(7.87)

Moreover, we have

$$\left|\Im \int_{\mathbb{R}^d} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \Phi_{\omega}(x) \overline{\nabla \eta(x,t)} \, dx\right| \le \||x|\Phi_{\omega}\|_{L^2} \|\nabla \eta(t)\|_{L^2}. \tag{7.88}$$

Here, since Φ_{ω} decays exponentially, $||x|\Phi_{\omega}||_{L^2}$ is finite.

We see from the decomposition (4.1), (7.87), (7.88), Lemma 4.3, Lemma 4.5, $\|\eta(t_1)\|_{H^1} \sim \widetilde{d}_{\omega}(t_1) = R$ and $\|\eta(t_3)\|_{H^1} \sim \widetilde{d}_{\omega}(t_3) \lesssim R$ that if R_* is sufficiently small dependently on ω ,

and $M = L(\omega)/R$ for some sufficiently large constant $L(\omega)$ depending only on d, p and ω , then

$$\begin{split} & \left| \left[\Im \int_{\mathbb{R}^{d}} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla \psi(x,t) \overline{\psi(x,t)} \, dx \right]_{t_{1}}^{t_{3}} \right| \\ \leq & \left| \left[\Im \int_{\mathbb{R}^{d}} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \left(\nabla \eta(x,t) \Phi_{\omega}(x) + \nabla \Phi_{\omega}(x,t) \overline{\eta(x,t)} \right) \, dx \right]_{t_{1}}^{t_{3}} \right| \\ & + \left| \left[\Im \int_{\mathbb{R}^{d}} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla \eta(x,t) \overline{\eta(x,t)} \, dx \right]_{t_{1}}^{t_{3}} \right| \\ \leq & \left| \left[\Im \int_{\mathbb{R}^{d}} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla \eta(x,t) \Phi_{\omega}(x) \, dx \right]_{t_{1}}^{t_{3}} \right| \\ & + \left| \left[\Im \int_{\mathbb{R}^{d}} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla \eta(x,t) \Phi_{\omega}(x) \, dx \right]_{t_{1}}^{t_{3}} \right| \\ & + \left| \left[\Im \int_{\mathbb{R}^{d}} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \Phi_{\omega}(x,t) \overline{\nabla \eta(x,t)} \, dx \right]_{t_{1}}^{t_{3}} \right| \\ & + \left| \left[\Im \int_{\mathbb{R}^{d}} \frac{M|x|}{(M+|x|)^{2}} + \frac{(d-1)M|x|}{(M+|x|)^{2}} \right] \Phi_{\omega}(x,t) \overline{\eta(x,t)} \, dx \right]_{t_{1}}^{t_{3}} \right| \\ & + \left| \left[\Im \int_{\mathbb{R}^{d}} \frac{M|x|}{M+|x|} \frac{x}{|x|} \cdot \nabla \eta(x,t) \overline{\eta(x,t)} \, dx \right]_{t_{1}}^{t_{3}} \right| \\ & \lesssim \|\eta(t_{1})\|_{H^{1}} + \|\eta(t_{3})\|_{H^{1}} + M\|\eta(t_{1})\|_{H^{1}}^{2} + M\|\eta(t_{3})\|_{H^{1}}^{2} \ll C^{**}(\omega) \delta_{X}. \end{split}$$

This contradicts (7.86). Hence, the case $\mathfrak{s} = 1$ never happens.

We can deal with the case where $\mathfrak{s} = -1$ in a way similar to Section 4.1 of [25]. Hence, we have completed the proof.

8 Proof of Theorem 3.1

Our aim in this section is to prove Theorem 3.1. By the time reversality, it suffices to show that there are only three possibilities (scattering, blowup, trapping) forward in time. Throughout this section, we fix $\omega \in (0, \omega_*)$, where $\omega_* > 0$ is the frequency given by the one-pass theorem (Theorem 7.1).

We introduce several notation used in this section:

• We use $I_{\max}(\psi)$ to denote the maximal existence-interval of a solution ψ to (NLS), and set $T_{\max}(\psi) := \sup I_{\max}(\psi)$ and $T_{\min}(\psi) := \inf I_{\min}(\psi)$.

• Since the set A_{ω}^{ε} is invariant under the flow defined by (NLS) for all $\varepsilon > 0$ (see (3.1)), we use the convention $\psi \in A_{\omega}^{\varepsilon}$ to indicate that $\psi(t) \in A_{\omega}^{\varepsilon}$ for all $t \in I_{\max}(\psi)$.

• For given R > 0 and $\varepsilon > 0$, we define $S_{\omega,R}^{\varepsilon}$ by

$$S_{\omega,R}^{\varepsilon} := \left\{ \psi : \begin{array}{l} \psi \text{ is a radial solution to (NLS) such that } 0 \in I_{\max}(\psi), \\ \psi \in A_{\omega}^{\varepsilon}, \text{ and } \widetilde{d}_{\omega}(\psi(t)) \geq R \text{ for all } t \in [0, T_{\max}(\psi)) \end{array} \right\}.$$
(8.1)

Next, we give a fundamental fact in A_{ω}^{ε} :

Lemma 8.1. Assume $d \ge 3$ and $1 + \frac{4}{d} . Let <math>\psi$ be a solution to (NLS) and $\varepsilon > 0$. Furthermore, assume that $\psi \in A_{\omega}^{\varepsilon}$ and

$$\inf_{t \in I_{\max}(\psi)} \|\nabla \psi(t)\|_{L^2}^2 \le \mathcal{H}(\Phi_\omega).$$
(8.2)

Then, we have either

$$\sup_{t \in I_{\max}(\psi)} \mathcal{K}(\psi(t)) < 0, \tag{8.3}$$

or

$$\inf_{t \in I_{\max}(\psi)} \mathcal{K}(\psi(t)) > 0.$$
(8.4)

Proof of Lemma 8.1. Note first that $\mathcal{M}(\psi) = \mathcal{M}(\Phi_{\omega})$ and therefore ψ is non-trivial. We also see from the assumption (8.2) that there exists $t_0 \in I_{\max}(\psi)$ such that

$$\|\nabla\psi(t_0)\|_{L^2}^2 \le \frac{3}{2}\mathcal{H}(\Phi_\omega).$$
 (8.5)

Furthermore, we see from $\mathcal{M}(\psi) = \mathcal{M}(\Phi_{\omega})$ and (8.5) that

$$S_{\omega}(\psi) = \omega \mathcal{M}(\psi) + \frac{1}{2} \|\nabla \psi(t_0)\|_{L^2}^2 - \frac{1}{p+1} \|\psi(t_0)\|_{L^{p+1}}^{p+1} - \frac{1}{2^*} \|\psi(t_0)\|_{L^{2*}}^{2*}$$

$$\leq \omega \mathcal{M}(\Phi_{\omega}) + \frac{3}{4} \mathcal{H}(\Phi_{\omega}) = S_{\omega}(\Phi_{\omega}) - \frac{1}{4} \mathcal{H}(\Phi_{\omega}) < m_{\omega}.$$
(8.6)

Hence, the claim follows from (1.49) and Theorem 1.3.

Now, let us recall that δ_X and δ_S denote the constants given by the ejection lemma (Lemma 5.2) and (7.7), respectively (see also Remark 6.1). Furthermore, the one-pass theorem shows that there exist constants $0 < \varepsilon_* \ll R_* \ll \delta_* (\ll \delta_S < \delta_X)$ such that: $\varepsilon_* \leq \varepsilon_0(\delta_*)$ (see Lemma 6.3 for the definition of $\varepsilon_0(\delta_*)$); and for any $\varepsilon \in (0, \varepsilon_*)$, any $R \in (\sqrt{2\varepsilon}, R_*)$ and any solution ψ to (NLS) with $\psi \in A_{\omega}^{\varepsilon}$, the following alternative holds: **Case 1.** There exists $t_0 \in I_{\max}(\psi)$ such that $\tilde{d}_{\omega}(\psi(t_0)) < R$ and $\tilde{d}_{\omega}(\psi(t)) < R + R^{\frac{\min\{3,p+1\}}{2}}$ for all $t \in [t_0, T_{\max}(\psi))$; or

Case 2. There exists $t_1 \in I_{\max}(\psi)$ such that for all $t_1 \leq t < T_{\max}(\psi)$,

$$\widetilde{d}_{\omega}(\psi(t)) \ge R. \tag{8.7}$$

In Case 1, it follows from $R + R^{\frac{\min\{3,p+1\}}{2}} < \delta_S < \delta_X$, (6.16) in Proposition 6.2, Remark 6.1 and Lemma 5.1 that $T_{\max}(\psi) = \infty$. Hence, ψ is trapped by $\mathcal{O}(\Phi_{\omega})$ forward in time.

Next, we consider Case 2. In this case, we have that $\psi(\cdot - t_1) \in S_{\omega,R}^{\varepsilon}$ and

$$\varepsilon \le \min\left\{\frac{\widetilde{d}_{\omega}(\psi(t))}{2}, \varepsilon_0(\delta_*)\right\}$$
(8.8)

for any $t \in [t_1, T_{\max}(\psi))$. In Section 8.1 and Section 8.2, we will prove that the solution ψ blows up in a finite time or scatters. To this end, we need some preparations. Let us begin with the following:

Lemma 8.2. Let $\varepsilon \in (0, \varepsilon_*)$, $R \in (\sqrt{2\varepsilon}, R_*)$, and let $\psi \in S_{\omega,R}^{\varepsilon}$. Then, for any $T \in (0, T_{\max}(\psi))$, there exists $\tau \in [T, T_{\max}(\psi))$ such that

$$\widetilde{d}_{\omega}(\psi(\tau)) \ge \delta_X. \tag{8.9}$$

Proof of Lemma 8.2. Suppose for contradiction that there exists $T_0 \in (0, T_{\max}(\psi))$ such that for any $t \in [T_0, T_{\max}(\psi))$,

$$d_{\omega}(\psi(t)) = \widetilde{d}_{\omega}(\psi(t)) < \delta_X, \qquad (8.10)$$

where the equality follows from (6.16) in Proposition 6.2 and Remark 6.1. Then, we see from Lemma 5.1 that $T_{\max}(\psi) = \infty$. Let $L \gg \frac{1}{\mu}$, and consider a time $t_L \in [T_0, T_0 + L \log \frac{\delta_X}{\varepsilon_*}]$ such that $R_L := d_{\omega}(\psi(t_L))$ is the minimum of $d_{\omega}(\psi(t))$ over the interval $[T_0, T_0 + L \log \frac{\delta_X}{\varepsilon_*}]$. Note that $\delta_X > R_L \ge R > \varepsilon_*$. If $T_0 \le t_L \le T_0 + \frac{L}{2} \log \frac{\delta_X}{\varepsilon_*}$, then we see from the ejection lemma (Lemma 5.2) that

$$d_{\omega}\left(\psi\left(t_{L}+\frac{L}{2}\log\frac{\delta_{X}}{\varepsilon_{*}}\right)\right) \sim e^{\mu\left(\frac{L}{2}\log\frac{\delta_{X}}{\varepsilon_{*}}\right)}R_{L} > \left(\frac{\delta_{X}}{\varepsilon_{*}}\right)^{\frac{\mu L}{2}}\varepsilon_{*} \gg \delta_{X}.$$
(8.11)

However, this contradicts (8.10). Similarly, when $t_L \ge T_0 + \frac{L}{2} \log \frac{\delta_X}{\varepsilon_*}$, applying the ejection lemma backward in time, we reach a contradiction. Thus, we have proved the claim.

Next, we show that the solution ψ stays away from Φ_{ω} after some time.

Lemma 8.3. Let $\varepsilon \in (0, \varepsilon_*)$, $R \in (\sqrt{2\varepsilon}, R_*)$, and let $\psi \in S_{\omega,R}^{\varepsilon}$. Then, there exists a time $T_0 \in [0, T_{\max}(\psi))$ such that for any $t \in [T_0, T_{\max}(\psi))$,

$$\widetilde{d}_{\omega}(\psi(t)) \ge R_*. \tag{8.12}$$

Proof of Lemma 8.3. It suffices to consider the case where there exists a time $\tau_0 \in [0, T_{\max}(\psi))$ such that

$$R \le \widetilde{R}_* := \widetilde{d}_{\omega} \left(\psi(\tau_0) \right) < R_*.$$
(8.13)

Here, it follows from Lemma 8.2 that we may assume that

$$\frac{\sqrt{1+4R_*}-1}{2} \le \tilde{R}_* < R_*.$$
(8.14)

Then, we see from the one-pass theorem (Theorem 7.1) that either

$$\sup_{t \ge \tau_0} \widetilde{d}_{\omega}(\psi(t)) \le \widetilde{R}_* + \widetilde{R}_*^{\frac{\min\{3,p+1\}}{2}}, \tag{8.15}$$

or there exists $T_0 \ge \tau_0$ such that

$$\inf_{t \ge T_0} \widetilde{d}_{\omega}(\psi(t)) \ge \widetilde{R}_* + \widetilde{R}_*^{\frac{\min\{3,p+1\}}{2}} \ge \widetilde{R}_* + \widetilde{R}_*^2 \ge R_*.$$
(8.16)

We find from Lemma 8.2 and $R_* \ll \delta_X$ that the former case (8.15) never happens. Thus, the latter case (8.16) only happens, and the proof is completed.

Lastly, we introduce a sign function which will determine the scattering or blowup.

Lemma 8.4. Let $\varepsilon \in (0, \varepsilon_*)$ and $R \in (\sqrt{2\varepsilon}, R_*)$. Then, there exists a function $\mathfrak{S} \colon S_{\omega,R}^{\varepsilon} \to \{1, -1\}$ such that

$$\mathfrak{S}(\psi) = \begin{cases} \operatorname{sign}[\lambda_1(t)] & \text{if } t \in [0, T_{\max}(\psi)) \text{ with } \widetilde{d}_{\omega}(\psi(t)) \leq \delta_E(\omega), \\ \operatorname{sign}[\mathcal{K}(\psi(t))] & \text{if } t \in [0, T_{\max}(\psi)) \text{ with } \widetilde{d}_{\omega}(\psi(t)) \geq \delta_*. \end{cases}$$

$$(8.17)$$

In addition, there exists a positive constant $C(\omega)$ which is independent of ε and R and satisfies that

$$\sup\left\{\|\psi\|_{L^{\infty}_{t}H^{1}_{x}([0,T_{\max}(\psi)))} \colon \psi \in S^{\varepsilon}_{\omega,R}, \ \mathfrak{S}(\psi) = 1\right\} \le C(\omega).$$

$$(8.18)$$

Proof of Lemma 8.4. We introduced a sign function (7.32) in the proof of one-pass theorem (Theorem 7.1). Employing the same argument as (7.32) and Lemma 8.2, we can prove the existence of the desired sign function. Furthermore, we can prove (8.18) in the same way as (7.33).

We divide $S_{\omega,R}^{\varepsilon}$ into two parts according to the sign of \mathfrak{S} :

$$S_{\omega,R,\pm}^{\varepsilon} := \left\{ \psi \in S_{\omega,R}^{\varepsilon} \colon \mathfrak{S}(\psi) = \pm 1 \right\}.$$
(8.19)

8.1 Analysis on $S_{\omega,R,-}^{\varepsilon}$

In this section, we shall prove that any radial solution $\psi \in S_{\omega,R,-}^{\varepsilon}$ blows up forward in time.

Proposition 8.5. Assume that $0 < \varepsilon < \min\{\varepsilon_0(R_*), \varepsilon_*\}$, where $\varepsilon_0(R_*)$ is a constant given by Lemma 6.3. Let $R \in (\sqrt{2\varepsilon}, R_*)$, and let $\psi \in S_{\omega,R,-}^{\varepsilon}$. Then, the maximal lifespan $T_{\max}(\psi)$ is finite: the finite time blowup forward in time.

Proof of Proposition 8.5. We see from Lemma 8.3, Lemma 6.3 and Lemma 8.1 that there exist a time $T_0 \in [0, T_{\max}(\psi))$ and $\kappa_1(R_*) > 0$ such that

$$\inf_{t \in [T_0, T_{\max}(\psi))} |\mathcal{K}(\psi(t))| \ge \kappa_1(R_*).$$
(8.20)

Moreover, it follows from Lemma 8.2 that there exists a time $\tau \in [T_0, T_{\max}(\psi))$ such that

$$\widetilde{d}_{\omega}(\psi(\tau)) \ge \delta_X \ge \delta_*. \tag{8.21}$$

Hence, Lemma 8.4 together with $\mathfrak{S}(\psi) = -1$ shows that

$$\mathcal{K}(\psi(\tau)) < 0. \tag{8.22}$$

Putting (8.20) and (8.22) together, we find that

$$\sup_{t \in [T_0, T_{\max}(\psi))} \mathcal{K}(\psi(t)) \le -\kappa_1(R_*).$$
(8.23)

Then, the same argument as the proof of Theorem 1.3 in [2] is available. Thus, we find that $T_{\max}(\psi) < \infty$.

8.2 Analysis on $S_{\omega,R,+}^{\varepsilon}$

In this section, we shall prove that any solution $\psi \in S_{\omega,R,+}^{\varepsilon}$ scatters forward in time. Lemma 8.3 together with time-translation allows us to restrict ourselves to the solutions in $S_{\omega,R_*,+}^{\varepsilon}$. Then, Lemma 6.3 together with Lemma 8.1 determines constants $\varepsilon_0(R_*) > 0$ and $\kappa_1(R_*) > 0$ such that if $\varepsilon \in (0, \varepsilon_0(R_*))$, then any solution $\psi \in S_{\omega,R_*,+}^{\varepsilon}$ obeys

$$\inf_{t \in [0, T_{\max}(\psi))} |\mathcal{K}(\psi(t))| \ge \kappa_1(R_*).$$
(8.24)

Furthermore, this together with Lemma 8.2 and Lemma 8.4 shows that for any $\varepsilon \in (0, \varepsilon_0(R_*))$ and any $\psi \in S_{\omega,R_*,+}^{\varepsilon}$,

$$\inf_{[0,T_{\max}(\psi))} \mathcal{K}(\psi(t)) \ge \kappa_1(R_*).$$
(8.25)

It is well known (see, e.g, the claim (v) of Theorem 4.1 in [3]) that the boundedness of a solution ψ in the Strichartz-type space $W_{p+1}([0, T_{\max}(\psi)) \cap W_{2^*}([0, T_{\max}(\psi)))$ implies the scattering forward in time. Thus, our aim is to prove the following:

Proposition 8.6. For any $\omega \in (0, \omega_*)$, there exists $\varepsilon \in (0, \varepsilon_*)$ such that for any $\psi \in S^{\varepsilon}_{\omega, R_*, +}$,

$$\|\psi\|_{W_{p+1}([0,T_{\max}(\psi))\cap W^{2^*}([0,T_{\max}(\psi)))} < \infty.$$
(8.26)

In particular, the solution ψ scatters forward in time.

Proof of Proposition 8.6. We divide the proof into several parts: In section 8.2.1, we suppose for contradiction that the claim was false, and extract some sequence of non-scattering solutions in $S_{\omega,R_{*,+}}^{\varepsilon_{*}}$. In Section 8.2.2, we apply a profile decomposition to this sequence, and obtain "linear profiles". In Section 8.2.3, we introduce the "nonlinear profiles", and investigate their fundamental properties. Furthermore, in Section 8.2.4, we show the existence of a nonlinear profile whose Strichartz norm diverges. Finally, in Section 8.2.5, we derive a contradiction by showing the existence of the "critical element", and complete the proof.

8.2.1 Setup

For any E > 0, we define $\nu(E)$ by

$$\nu(E) := \sup \left\{ \|\psi\|_{W_{p+1}([0,T_{\max}(\psi))) \cap W_{2^*}([0,T_{\max}(\psi)))} \colon \psi \in S_{\omega,R_*,+}^{\varepsilon_*}, \ \mathcal{H}(\psi) \le E \right\}.$$
(8.27)

Furthermore, we put

$$E_* := \sup\{E > 0 \colon \nu(E) < \infty\}.$$
(8.28)

If $\psi \in S_{\omega,R_{*},+}^{\varepsilon_{*}}$ and $\mathcal{H}(\psi) < \mathcal{H}(\Phi_{\omega})$, then we see from (8.25) that $\psi \in PW_{\omega,+}$. Hence, it follows from Theorem 1.3 that

$$E_* \ge \mathcal{H}(\Phi_\omega). \tag{8.29}$$

Thus, what we want to prove is that $E_* > \mathcal{H}(\Phi_{\omega})$. We prove this by contradiction, and therefore suppose that

$$E_* = \mathcal{H}(\Phi_\omega). \tag{8.30}$$

Then, we can take a sequence $\{\varepsilon_n\}$ of constants in $(0, \varepsilon_*)$ and a sequence $\{\psi_n\}$ of solutions such that

$$\lim_{n \to \infty} \varepsilon_n = 0, \tag{8.31}$$

$$\varepsilon_n < \min\left\{\varepsilon_0(R_*), \ \frac{\kappa_1(R_*)}{10d(p-1)+10}\right\},\tag{8.32}$$

$$\psi_n \in S^{\varepsilon_n}_{\omega, R_*, +},\tag{8.33}$$

$$\|\psi_n\|_{W_{p+1}([0,T_{\max}(\psi_n)))\cap W_{2^*}([0,T_{\max}(\psi_n)))} = \infty.$$
(8.34)

In particular, we have that

$$\inf_{n \ge 1} \inf_{t \in [0, T_{\max}(\psi_n))} \widetilde{d}_{\omega}(\psi_n(t)) \ge R_*,$$
(8.35)

$$\lim_{n \to \infty} \mathcal{H}(\psi_n) = \mathcal{H}(\Phi_\omega) = E_*.$$
(8.36)

Furthermore, it follows from (8.25) and Lemma 8.4 that

$$\inf_{n \ge 1} \inf_{t \in [0, T_{\max}(\psi_n))} \mathcal{K}(\psi_n(t)) \ge \kappa_1(R_*), \tag{8.37}$$

$$\sup_{n \ge 1} \sup_{t \in [0, T_{\max}(\psi_n))} \|\psi_n(t)\|_{H^1} < \infty.$$
(8.38)

8.2.2 Linear profiles

Let $\{\psi_n\}$ be the sequence of solutions obtained in Section 8.2.1. We employ the profile decomposition for a sequence in the homogeneous space $\dot{H}^1(\mathbb{R}^d)$ (see Theorem 1.6 in [19] and Lemma 2.10 in [21]). We apply it to the sequence $\{|\nabla|^{-1}\langle \nabla \rangle \psi_n(0)\}$ in $\dot{H}^1(\mathbb{R}^d)$, instead of $\{\psi_n\}$ itself. Then, we find³ that there exists a subsequence of $\{\psi_n\}$ (denoted by the same symbol $\{\psi_n\}$) with the following properties: there exist a family $\{\{(x_n^1, t_n^1, \lambda_n^1)\}, \{(x_n^2, t_n^2, \lambda_n^2)\}, \ldots\}$ of sequences in $\mathbb{R}^d \times \mathbb{R} \times (0, \infty)$, a family $\{\tilde{u}^1, \tilde{u}^2, \ldots\}$ of functions in $H^1(\mathbb{R}^d)$, and a family $\{\{w_n^1\}, \{w_n^2\}, \ldots\}$ of sequences in $H^1(\mathbb{R}^d)$ such that: • For any $j \geq 1$,

$$\lim_{n \to \infty} t_n^j = t_\infty^j \in \mathbb{R} \cup \{\pm \infty\},\tag{8.39}$$

$$\lim_{n \to \infty} \lambda_n^j = \lambda_\infty^j \in \{0, 1, \infty\}, \qquad \begin{cases} \lambda_n^j \equiv 1 & \text{if } \lambda_\infty^j = 1, \\ \lambda_n^j \leq 1 & \text{if } \lambda_\infty^j = 0. \end{cases}$$
(8.40)

³We do not use the assumption that every ψ_n is radial in this subsection (Section 8.2.2) and therefore translation parameters appear in the linear profile decomposition. In Section 8.2.4 and 8.2.5, we need the radial symmetry of the solutions so that we can take every translation $x_n^j = 0$.

• For any $j' \neq j$,

$$\lim_{n \to \infty} \left\{ \frac{\lambda_n^{j'}}{\lambda_n^j} + \frac{\lambda_n^j}{\lambda_n^{j'}} + \frac{|x_n^j - x_n^{j'}|}{\lambda_n^j} + \frac{|t_n^j - t_n^{j'}|}{(\lambda_n^j)^2} \right\} = \infty.$$
(8.41)

• For any \dot{H}^1 -admissible pair⁴ (q, r),

$$\lim_{j \to \infty} \lim_{n \to \infty} \left\| |\nabla|^{-1} \langle \nabla \rangle e^{it\Delta} w_n^j \right\|_{L^r_t L^q_x(\mathbb{R})} = 0.$$
(8.42)

• For any $n \ge 1$ and $k \ge 1$,

$$e^{it\Delta}\psi_n(0) = \sum_{j=1}^k \langle \nabla \rangle^{-1} |\nabla| G_n^j \left(e^{it\Delta} |\nabla|^{-1} \langle \nabla \rangle \widetilde{u}^j \right) + e^{it\Delta} w_n^k$$

$$= \sum_{j=1}^k \frac{\langle \nabla \rangle^{-1} \langle \lambda_n^j \nabla \rangle}{\lambda_n^j} e^{i(t-t_n^j)\Delta} g_n^j \widetilde{u}^j + e^{it\Delta} w_n^k,$$
(8.43)

where G_n^j and g_n^j are the operators defined by

$$G_{n}^{j}v(x,t) := \frac{1}{(\lambda_{n}^{j})^{\frac{d-2}{2}}} v\Big(\frac{x - x_{n}^{j}}{\lambda_{n}^{j}}, \frac{t - t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\Big),$$
(8.44)

$$g_n^j u(x) := \frac{1}{(\lambda_n^j)^{\frac{d-2}{2}}} u\Big(\frac{x - x_n^j}{\lambda_n^j}\Big).$$
(8.45)

Note that

$$\frac{\langle \nabla \rangle^{-1} \langle \lambda_n^j \nabla \rangle}{\lambda_n^j} e^{i(t-t_n^j)\Delta} g_n^j \widetilde{u}^j = g_n^j \sigma_n^j e^{i\frac{t-t_n^j}{(\lambda_n^j)^2}\Delta} \widetilde{u}^j = G_n^j \sigma_n^j e^{it\Delta} \widetilde{u}^j,$$
(8.46)

where

$$\sigma_n^j := \frac{\langle (\lambda_n^j)^{-1} \nabla \rangle^{-1} \langle \nabla \rangle}{\lambda_n^j}.$$
(8.47)

• For any $k \ge 1$ and any $1 \le j \le k$,

$$\lim_{n \to \infty} \langle \nabla \rangle^{-1} |\nabla| e^{-i \frac{t_n^j}{(\lambda_n^j)} \Delta} (g_n^j)^{-1} |\nabla|^{-1} \langle \nabla \rangle w_n^k = 0 \quad \text{weakly in } H^1(\mathbb{R}^d).$$
(8.48)

• For any $k \ge 1$,

$$\lim_{n \to \infty} \left\{ \| \langle \nabla \rangle \psi_n(0) \|_{L^2}^2 - \sum_{j=1}^k \| \langle \nabla \rangle g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \|_{L^2}^2 - \| \langle \nabla \rangle w_n^k \|_{L^2}^2 \right\} = 0.$$
(8.49)

⁴A pair (q,r) is said to be \dot{H}^1 -admissible if $\frac{1}{r} = \frac{d}{2}(\frac{1}{2} - \frac{1}{q} - \frac{1}{d})$ and $(q,r) \in [2,\infty] \times [2,\infty]$.
Furthermore, the following hold for all $k \ge 1$ (see Lemma 2.2. in [3]):

$$\lim_{n \to \infty} \left\{ \mathcal{M}(\psi_n) - \sum_{j=1}^k \mathcal{M}\left(g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j\right) - \mathcal{M}(w_n^k) \right\} = 0,$$
(8.50)

$$\lim_{n \to \infty} \left\{ \mathcal{H}(\psi_n) - \sum_{j=1}^k \mathcal{H}\left(g_n^j \sigma_n^j e^{-i\frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j\right) - \mathcal{H}(w_n^k) \right\} = 0,$$
(8.51)

$$\lim_{n \to \infty} \left\{ \mathcal{S}_{\omega}(\psi_n) - \sum_{j=1}^k \mathcal{S}_{\omega} \left(g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right) - \mathcal{S}_{\omega}(w_n^k) \right\} = 0,$$
(8.52)

$$\lim_{n \to \infty} \left\{ \mathcal{I}_{\omega}(\psi_n(0)) - \sum_{j=1}^k \mathcal{I}_{\omega}\left(g_n^j \sigma_n^j e^{-i\frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j\right) - \mathcal{I}_{\omega}(w_n^k) \right\} = 0.$$
(8.53)

Note that it follows from Strichartz' estimate, (8.38) and (8.49) that for any $k \ge 1$, there exists a number N(k) such that for any $n \ge N(k)$,

$$\left\| \langle \nabla \rangle e^{it\Delta} w_n^k \right\|_{St(\mathbb{R})} \lesssim \|w_n^k\|_{H^1} \lesssim 1.$$
(8.54)

Moreover, it follows from (8.42) and (8.54) that for any $1 + \frac{4}{d} < q \le 2^* - 1$,

$$\lim_{k \to \infty} \lim_{n \to \infty} \|e^{it\Delta} w_n^k\|_{W_{q+1}(\mathbb{R})} \leq \lim_{k \to \infty} \lim_{n \to \infty} \|e^{it\Delta} w_n^k\|_{W_{2+\frac{4}{4}}(\mathbb{R})}^{1-s_q} \|e^{it\Delta} w_n^k\|_{W_{2^*}(\mathbb{R})}^{s_q} \\
\lesssim \lim_{k \to \infty} \lim_{n \to \infty} \||\nabla|^{-1} \langle \nabla \rangle e^{it\Delta} w_n^k\|_{W_{2^*}(\mathbb{R})}^{s_q} = 0.$$
(8.55)

We shall discuss fundamental properties of operators g_n^j (see (8.45)) and σ_n^j (see (8.47)). To this end, for each $j \ge 1$, we introduce an operator σ_{∞}^j as

$$\sigma_{\infty}^{j} := \begin{cases} |\nabla|^{-1} \langle \nabla \rangle & \text{if } \lambda_{\infty}^{j} = 0, \\ 1 & \text{if } \lambda_{\infty}^{j} = 1, \\ 0 & \text{if } \lambda_{\infty}^{j} = \infty. \end{cases}$$

$$(8.56)$$

Lemma 8.7. The following hold for all function $f \in H^1(\mathbb{R}^d)$ and all $j \ge 1$: (i)

$$\lim_{n \to \infty} \left\| \nabla \left\{ \sigma_n^j f - \sigma_\infty^j f \right\} \right\|_{L^2} = 0; \tag{8.57}$$

(ii)

$$\lim_{n \to \infty} \left\| g_n^j \sigma_n^j f \right\|_{L^2} = \lim_{n \to \infty} \left\| \lambda_n^j \sigma_n^j f \right\|_{L^2} = \begin{cases} 0 & \text{if } \lambda_\infty^j = 0, \\ \|f\|_{L^2} & \text{if } \lambda_\infty^j = 1, \\ \|\langle \nabla \rangle f\|_{L^2} & \text{if } \lambda_\infty^j = \infty. \end{cases}$$
(8.58)

Proof of Lemma 8.7. We shall prove (8.57). If $\lambda_{\infty}^{j} = 1$, then $\sigma_{n}^{j} \equiv 1$ and therefore the claim is trivial. Assume that $\lambda_{\infty}^{j} = 0$. Then, we see from Parseval's identity that

$$\lim_{n \to \infty} \left\| |\nabla| \left\{ \sigma_n^j f - \sigma_\infty^j f \right\} \right\|_{L^2} = \lim_{n \to \infty} \left\| |\nabla| \left\{ \frac{\langle (\lambda_n^j)^{-1} \nabla \rangle^{-1}}{\lambda_n^j} - |\nabla|^{-1} \right\} \langle \nabla \rangle f \right\|_{L^2}$$

$$= \lim_{n \to \infty} \left\| \left\{ \frac{|\xi|}{\sqrt{(\lambda_n^j)^2 + |\xi|^2}} - 1 \right\} \langle \xi \rangle \mathcal{F}[f] \right\|_{L^2_{\xi}}.$$
(8.59)

Moreover, since $\lim_{n\to\infty} \lambda_n^j = \lambda_{\infty}^j = 0$, we see that

$$\lim_{n \to \infty} \left\{ \frac{|\xi|}{\sqrt{(\lambda_n^j)^2 + |\xi|^2}} - 1 \right\} \langle \xi \rangle \mathcal{F}[f](\xi) = 0 \quad \text{for almost all } \xi \in \mathbb{R}^d.$$
(8.60)

Hence, we find from Lebesgue's convergence theorem that (8.57) holds. Similarly, we can prove the case where $\lambda_{\infty}^{j} = \infty$.

Next, we give a proof of (8.58). The case where $\lambda_{\infty}^{j} = 1$ is trivial. Assume that $\lambda_{\infty}^{j} = 0$. Then, we see from the substitution of variables and Parseval's identity that

$$\lim_{n \to \infty} \|g_n^j \sigma_n^j f\|_{L^2} = \lim_{n \to \infty} \|\lambda_n^j \sigma_n^j f\|_{L^2} = \lim_{n \to \infty} \left\| \frac{1}{\sqrt{1 + \frac{|\xi|^2}{(\lambda_n^j)^2}}} \langle \xi \rangle \mathcal{F}[f] \right\|_{L^2_{\xi}}.$$
 (8.61)

Since $\lim_{n\to\infty} \lambda_n^j = \lambda_{\infty}^j = 0$, we find from (8.61) and Lebesgue's convergence theorem that (8.58) holds. Similarly, we can prove the case where $\lambda_{\infty}^j = \infty$.

Lemma 8.8. Let j be a number such that $\lambda_{\infty}^{j} \neq \infty$. Then, for any $n \geq 1$, any $q \in [2, \infty)$ and any function f with $\langle \nabla \rangle f \in L^{q}(\mathbb{R}^{d})$,

$$\left\|\lambda_n^j \sigma_n^j f\right\|_{L^q} + \left\|\nabla \sigma_n^j f\right\|_{L^q} \lesssim \left\|\langle \nabla \rangle f\right\|_{L^q},\tag{8.62}$$

where the implicit constant depends only on d and q.

Proof of Lemma 8.8. If $\lambda_{\infty}^{j} = 1$, then the claim is trivial. When $\lambda_{\infty}^{j} = 0$, we find from the Mihlin multiplier theorem that

$$\begin{aligned} \left\|\lambda_{n}^{j}\sigma_{n}^{j}f\right\|_{L^{q}} + \left\|\nabla\sigma_{n}^{j}f\right\|_{L^{q}} \lesssim \left\|\langle\nabla\rangle f\right\|_{L^{q}} + \left\|\sigma_{n}^{j}(\sigma_{\infty}^{j})^{-1}|\nabla|\sigma_{\infty}^{j}f\right\|_{L^{q}} \\ \lesssim \left\|\langle\nabla\rangle f\right\|_{L^{q}} + \left\||\nabla|\sigma_{\infty}^{j}f\right\|_{L^{q}} \lesssim \left\|\langle\nabla\rangle f\right\|_{L^{q}}. \end{aligned}$$

$$(8.63)$$

Thus, we have proved the lemma.

Lemma 8.9. Let $1 + \frac{4}{d} < q < 2^* - 1$, and let $j \ge 1$ be a number for which $\lambda_{\infty}^j \in \{0, \infty\}$. Then, the linear profile \tilde{u}^j satisfies

$$\lim_{n \to \infty} \|G_n^j \sigma_n^j e^{it\Delta} \widetilde{u}^j |_{t=0} \|_{L^{q+1}} = \lim_{n \to \infty} \|g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \|_{L^{q+1}} = 0.$$
(8.64)

Proof of Lemma 8.9. Assume that $\lambda_{\infty}^{j} \in \{0,\infty\}$. Then, we see from the Gagliardo-Nirenberg inequality that

$$\left\|g_{n}^{j}\sigma_{n}^{j}e^{-i\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\Delta}\widetilde{u}^{j}\right\|_{L^{q+1}}^{q+1} \lesssim \left\|\lambda_{n}^{j}\sigma_{n}^{j}\widetilde{u}^{j}\right\|_{L^{2}}^{q+1-\frac{d(q-1)}{2}} \left\|\nabla\sigma_{n}^{j}\widetilde{u}^{j}\right\|_{L^{2}}^{\frac{d(q-1)}{2}}.$$
(8.65)

If $\lambda_{\infty}^{j} = 0$, then it follows from (8.65) and Lemma 8.7 that

$$\lim_{n \to \infty} \left\| g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right\|_{L^{q+1}}^{q+1} \lesssim \lim_{n \to \infty} \left\| \lambda_n^j \sigma_n^j \widetilde{u}^j \right\|_{L^2}^{q+1 - \frac{d(q-1)}{2}} \left\| \langle \nabla \rangle \widetilde{u}^j \right\|_{L^2}^{\frac{d(q-1)}{2}} = 0.$$
(8.66)

On the other hand, if $\lambda_{\infty}^{j} = \infty$, then it follows from (8.65) and Lemma 8.7 that

$$\lim_{n \to \infty} \left\| g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right\|_{L^{q+1}}^{q+1} \lesssim \left\| \langle \nabla \rangle \widetilde{u}^j \right\|_{L^2}^{q+1 - \frac{d(q-1)}{2}} \lim_{n \to \infty} \left\| \nabla \sigma_n^j \widetilde{u}^j \right\|_{L^2}^{\frac{d(q-1)}{2}} = 0.$$
(8.67)

Thus, we have completed the proof.

8.2.3 Nonlinear profiles

In this subsection, we introduce the "nonlinear profile" associated with $(\tilde{u}^j, \{(x_n^j, t_n^j, \lambda_n^j)\})$ for each $j \ge 1$. From the point of view of (8.43), we first introduce a function U_n^j as the solution to (NLS) satisfying

$$U_n^j(0) = \frac{\langle \nabla \rangle^{-1} \langle \lambda_n^j \nabla \rangle}{\lambda_n^j} e^{-it_n^j \Delta} g_n^j \widetilde{u}^j.$$
(8.68)

Then, we find from (8.46) that

$$e^{it\Delta}U_n^j(0) = G_n^j \sigma_n^j e^{it\Delta} \widetilde{u}^j.$$
(8.69)

Furthermore, Duhamel's formula for U_n^j together with (8.69) shows that U_n^j satisfies

$$U_{n}^{j}(t) - G_{n}^{j}\sigma_{n}^{j}e^{it\Delta}\widetilde{u}^{j} = i\int_{0}^{t}e^{i(t-s)\Delta} \left\{ F^{\dagger}[U_{n}^{j}](s) + F^{\dagger}[U_{n}^{j}](s) \right\} ds,$$
(8.70)

where see (1.65) for the definitions of F^{\dagger} and F^{\ddagger} . Applying $(\sigma_n^j)^{-1} (G_n^j)^{-1}$ to the both sides in (8.70), we find that

$$\begin{split} (\sigma_n^j)^{-1} (G_n^j)^{-1} U_n^j(t) &- e^{it\Delta} \widetilde{u}^j \\ &= i(\sigma_n^j)^{-1} (g_n^j)^{-1} \int_{-\frac{t_n^j}{(\lambda_n^j)^2}}^t e^{i(\lambda_n^j)^2(t-s)\Delta} \left\{ (\lambda_n^j)^{-\frac{(d-2)(p-1)}{2}} g_n^j F^{\dagger}[(G_n^j)^{-1} U_n^j](s) \right. \\ &+ (\lambda_n^j)^{-2} g_n^j F^{\dagger}[(G_n^j)^{-1} U_n^j](s) \left. \right\} (\lambda_n^j)^2 \, ds \end{split}$$

$$\end{split}$$

$$\begin{split} (8.71) \\ &= i(\sigma_n^j)^{-1} \int_{-\frac{t_n^j}{(\lambda_n^j)^2}}^t e^{i(t-s)\Delta} \left\{ (\lambda_n^j)^{\frac{4-(d-2)(p-1)}{2}} F^{\dagger}[\sigma_n^j(\sigma_n^j)^{-1} (G_n^j)^{-1} U_n^j](s) \right. \\ &+ F^{\dagger}[\sigma_n^j(\sigma_n^j)^{-1} (G_n^j)^{-1} U_n^j](s) \left. \right\} \, ds. \end{split}$$

From the point of view of (8.71), for each number $j \ge 1$ with $\lambda_{\infty}^{j} \in \{0, 1\}$, we define the nonlinear profile $\tilde{\psi}^{j}$ associated with $(\tilde{u}^{j}, \{x_{n}^{j}, t_{n}^{j}, \lambda_{n}^{j}\})$ as the solution to

$$\begin{split} \widetilde{\psi}^{j}(t) &= e^{it\Delta} \widetilde{u}^{j} + i \int_{-\frac{t_{\infty}^{j}}{(\lambda_{\infty}^{j})^{2}}}^{t} e^{i(t-s)\Delta} (\sigma_{\infty}^{j})^{-1} \Big\{ (\lambda_{\infty}^{j})^{\frac{4-(d-2)(p-1)}{2}} F^{\dagger}[\sigma_{\infty}^{j} \widetilde{\psi}^{j}](s) \\ &+ F^{\ddagger}[\sigma_{\infty}^{j} \widetilde{\psi}^{j}](s) \Big\} \, ds. \end{split}$$

$$(8.72)$$

When $-\frac{t_{\infty}^{j}}{(\lambda_{\infty}^{j})^{2}} \in \{\pm\infty\}$, the equation (8.72) is interpreted as the final value problem at $\pm\infty$. Moreover, if $j \ge 1$ is a number for which $\lambda_{\infty}^{j} = \infty$, then we define $\tilde{\psi}^{j}(t) := e^{it\Delta}\tilde{u}^{j}$ (we define $\tilde{\psi}^{j}$ as a free solution to (NLS) so that (8.76) holds; in particular, if $\tilde{u}^{j} \equiv 0$, then $\tilde{\psi}^{j} \equiv 0$).

We simply write $I_{\max}^j := I_{\max}(\widetilde{\psi}^j)$, $T_{\max}^j := T_{\max}(\widetilde{\psi}^j)$ and $T_{\min}^j := T_{\min}(\widetilde{\psi}^j)$. Moreover, for an interval I and a number $j \ge 1$, we define

$$\mathcal{W}^{j}(I) := \begin{cases} W_{p+1}(I) \cap W_{2^{*}}(I) & \text{if } \lambda_{\infty}^{j} \in \{1, \infty\}, \\ W_{2^{*}}(I) & \text{if } \lambda_{\infty}^{j} = 0. \end{cases}$$
(8.73)

We find from the construction of the nonlinear profiles (see (8.72)) that the following hold for all numbers $j \ge 1$:

- $\widetilde{\psi}^j \in C(I^j_{\max}, H^1(\mathbb{R}^d)).$
- $\widetilde{\psi}^j$ satisfies

$$\begin{cases}
i\frac{\partial\sigma_{\infty}^{j}\widetilde{\psi}^{j}}{\partial t} + \Delta\sigma_{\infty}^{j}\widetilde{\psi}^{j} + F[\sigma_{\infty}^{j}\widetilde{\psi}^{j}] = 0 \quad \text{if} \quad \lambda_{\infty}^{j} = 1, \\
i\frac{\partial\sigma_{\infty}^{j}\widetilde{\psi}^{j}}{\partial t} + \Delta\sigma_{\infty}^{j}\widetilde{\psi}^{j} + F^{\ddagger}[\sigma_{\infty}^{j}\widetilde{\psi}^{j}] = 0 \quad \text{if} \quad \lambda_{\infty}^{j} = 0, \\
i\frac{\partial\widetilde{\psi}^{j}}{\partial t} + \Delta\widetilde{\psi}^{j} = 0 \quad \text{if} \quad \lambda_{\infty}^{j} = \infty.
\end{cases}$$
(8.74)

• There exists a number N(j) such that for any $n \ge N(j)$,

$$-\frac{t_n^j}{(\lambda_n^j)^2} \in I_{\max}^j \tag{8.75}$$

and

$$\lim_{n \to \infty} \left\| \widetilde{\psi}^j \left(-\frac{t_n^j}{(\lambda_n^j)^2} \right) - e^{-i\frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right\|_{H^1} = 0.$$
(8.76)

• If $-\frac{t_{\infty}^{j}}{(\lambda_{\infty}^{j})^{2}} = \infty$, then $T_{\max}^{j} = \infty$ and for any $T > T_{\min}^{j}$,

$$\left\|\sigma_{\infty}^{j}\widetilde{\psi}^{j}\right\|_{\mathcal{W}^{j}\left([T,\infty)\right)} < \infty; \tag{8.77}$$

• If $-\frac{t_{\infty}^j}{(\lambda_{\infty}^j)^2} = -\infty$, then $T_{\min}^j = -\infty$ and for any $T < T_{\max}^j$,

$$\left\|\sigma_{\infty}^{j}\widetilde{\psi}^{j}\right\|_{\mathcal{W}^{j}((-\infty,T])} < \infty.$$
(8.78)

When $\lambda_{\infty}^{j} \in \{0, \infty\}$, it is convenient rewriting (8.76) in the form

$$\begin{cases} \lim_{n \to \infty} \left\| \widetilde{\psi}^{j} \left(\cdot -x_{n}^{j}, -\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \right) - g_{n}^{j} \sigma_{n}^{j} e^{-i \frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \Delta} \widetilde{u}^{j} \right\|_{H^{1}} = 0 \quad \text{if} \quad \lambda_{\infty}^{j} = 1, \\ \lim_{n \to \infty} \left\| \nabla g_{n}^{j} \left\{ \sigma_{\infty}^{j} \widetilde{\psi}^{j} \left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \right) - \sigma_{n}^{j} e^{-i \frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \Delta} \widetilde{u}^{j} \right\} \right\|_{L^{2}} = 0 \quad \text{if} \quad \lambda_{\infty}^{j} = 0. \end{cases}$$

$$(8.79)$$

Now, for each number $j \ge 1$, we define

$$\psi_n^j := G_n^j \sigma_n^j \widetilde{\psi}^j, \tag{8.80}$$

and let $I_{\max,n}^{j}$ be the maximal existence-interval of ψ_{n}^{j} . Then, we see from (8.74) that

$$\begin{cases}
i\frac{\partial\psi_n^j}{\partial t} + \Delta\psi_n^j + F[\psi_n^j] = 0 & \text{if } \lambda_{\infty}^j = 1, \\
i\frac{\partial\psi_n^j}{\partial t} + \Delta\psi_n^j + (\sigma_{\infty}^j)^{-1}F^{\ddagger}[G_n^j\sigma_{\infty}^j\widetilde{\psi}^j] = 0 & \text{if } \lambda_{\infty}^j = 0, \\
i\frac{\partial\psi_n^j}{\partial t} + \Delta\psi_n^j = 0 & \text{if } \lambda_{\infty}^j = \infty,
\end{cases}$$
(8.81)

and

$$I_{\max,n}^{j} = \begin{cases} ((\lambda_{n}^{j})^{2} T_{\min}^{j} + t_{n}^{j}, \ (\lambda_{n}^{j})^{2} T_{\max}^{j} + t_{n}^{j}) & \text{if } \lambda_{\infty}^{j} \in \{0,1\}, \\ (-\infty,\infty) & \text{if } \lambda_{\infty}^{j} = \infty. \end{cases}$$
(8.82)

Furthermore, (8.75) shows that $0 \in I^{j}_{\max,n}$ for any $n \ge N(j)$.

In what follows, we discuss fundamental properties of nonlinear profiles in a series of lemmas. In particular, we show that there is at most one "bad nonlinear profile"; the other profiles are "good" in the sense of (8.135) and (8.136).

Lemma 8.10. Let j be a number for which $\lambda_{\infty}^{j} = 0$, and let I be an interval on which

$$\left\| \langle \nabla \rangle \widetilde{\psi}^{j} \right\|_{St(I)} < \infty.$$
(8.83)

Then, the following hold:

$$\lim_{n \to \infty} \|\sigma_{\infty}^{j} \widetilde{\psi}^{j} - \sigma_{n}^{j} \widetilde{\psi}^{j}\|_{W_{2^{*}}(I)} \lesssim \lim_{n \to \infty} \|\nabla\{\sigma_{\infty}^{j} \widetilde{\psi}^{j} - \sigma_{n}^{j} \widetilde{\psi}^{j}\}\|_{V_{2^{*}}(I)} = 0,$$
(8.84)

$$\lim_{n \to \infty} \left\| \nabla \left\{ \sigma_{\infty}^{j} \widetilde{\psi}^{j} - \sigma_{n}^{j} \widetilde{\psi}^{j} \right\} \right\|_{V_{2+\frac{4}{d}}(I)} = 0, \tag{8.85}$$

$$\lim_{n \to \infty} \left\| \lambda_n^j \sigma_n^j \widetilde{\psi}^j \right\|_{V_{2+\frac{4}{d}}(I)} = 0.$$
(8.86)

Proof of Lemma 8.10. The inequality in (8.84) follows from Sobolev's embedding. We see from Hölder's inequality, Lemma 8.7, Lemma 8.8 and the assumption (8.83) that for almost all $t \in I$,

$$\lim_{n \to \infty} \|\nabla \{\sigma_{\infty}^{j} \widetilde{\psi}^{j}(t) - \sigma_{n}^{j} \widetilde{\psi}^{j}(t)\}\|_{L^{2d(d+2)}}^{\frac{2(d+2)}{d-2}} \leq \lim_{n \to \infty} \|\nabla \{\sigma_{\infty}^{j} \widetilde{\psi}^{j}(t) - \sigma_{n}^{j} \widetilde{\psi}^{j}(t)\}\|_{L^{2}(I)}^{\frac{8}{d-2}} \|\langle \nabla \rangle \widetilde{\psi}^{j}(t)\|_{L^{2^{*}}(I)}^{2} = 0.$$
(8.87)

Furthermore, we see from Lemma 8.8 and the assumption (8.83) that

$$\|\nabla\{\sigma_{\infty}^{j}\widetilde{\psi}^{j}(t) - \sigma_{n}^{j}\widetilde{\psi}^{j}(t)\}\|_{L^{\frac{2d(d+2)}{d-2}}}^{\frac{2(d+2)}{d-2}} \lesssim \|\langle\nabla\rangle\widetilde{\psi}^{j}(t)\|_{L^{\frac{2d(d+2)}{d-2}}}^{\frac{2(d+2)}{d-2}} \in L^{1}(I).$$
(8.88)

Thus, Lebesgue's convergence theorem shows that the equality in (8.84) holds. Similarly, we can prove (8.85). It remains to prove (8.86). We see from Hölder's inequality, Lemma 8.7, Lemma 8.8 and the assumption (8.83) that for almost all $t \in I$,

$$\lim_{n \to \infty} \left\| \lambda_n^j \sigma_n^j \widetilde{\psi}^j(t) \right\|_{L_{\frac{2(d+2)}{d}}}^{\frac{2(d+2)}{d}} \le \lim_{n \to \infty} \left\| \lambda_n^j \sigma_n^j \widetilde{\psi}^j(t) \right\|_{L^2(I)}^{\frac{4}{d}} \| \langle \nabla \rangle \widetilde{\psi}^j(t) \|_{L^{2^*}(I)}^2 = 0.$$
(8.89)

Furthermore, we see from Lemma 8.8 and the assumption (8.83) that

$$\|\lambda_{n}^{j}\sigma_{n}^{j}\widetilde{\psi}^{j}(t)\|_{L^{\frac{2(d+2)}{d}}}^{\frac{2(d+2)}{d}} \lesssim \|\langle\nabla\rangle\widetilde{\psi}^{j}(t)\|_{L^{\frac{2(d+2)}{d}}}^{\frac{2(d+2)}{d}} \in L^{1}(I).$$
(8.90)

Thus, Lebesgue's convergence theorem shows that (8.86) holds.

Lemma 8.11. Let j be a number for which $\lambda_{\infty}^{j} = 0$, and let I be an interval on which

$$\left\| \langle \nabla \rangle \widetilde{\psi}^{j} \right\|_{St(I)} < \infty. \tag{8.91}$$

Then, the nonlinear term in (8.81) satisfies

$$\lim_{n \to \infty} \left\| \langle \nabla \rangle \left\{ (\sigma_{\infty}^{j})^{-1} F^{\dagger}[G_{n}^{j} \sigma_{\infty}^{j} \widetilde{\psi}^{j}] - F[\psi_{n}^{j}] \right\} \right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_{n}^{j})} = 0,$$
(8.92)

where $I_n^j := ((\lambda_n^j)^2 \inf I + t_n^j, \ (\lambda_n^j)^2 \sup I + t_n^j).$

Proof of Lemma 8.11. Note first that

$$F[\psi_n^j] = F^{\ddagger}[G_n^j \sigma_n^j \widetilde{\psi}^j] + F^{\dagger}[G_n^j \sigma_n^j \widetilde{\psi}^j].$$
(8.93)

Next, observe that

$$\begin{split} &\lim_{n\to\infty} \left\| \langle \nabla \rangle \left\{ (\sigma_{\infty}^{j})^{-1} F^{\dagger}[G_{n}^{j} \sigma_{\infty}^{j} \widetilde{\psi}^{j}] - F^{\dagger}[G_{n}^{j} \sigma_{n}^{j} \widetilde{\psi}^{j}] \right\} \right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}} (I_{n}^{j}) \\ &= \lim_{n\to\infty} (\lambda_{n}^{j})^{-2} \left\| G_{n}^{j} (\lambda_{n}^{j})^{-1} |\nabla| F^{\dagger}[\sigma_{\infty}^{j} \widetilde{\psi}^{j}] - G_{n}^{j} \langle (\lambda_{n}^{j})^{-1} \nabla \rangle F^{\dagger}[\sigma_{n}^{j} \widetilde{\psi}^{j}] \right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}} (I_{n}^{j}) \\ &= \lim_{n\to\infty} \left\| |\nabla| F^{\dagger}[\sigma_{\infty}^{j} \widetilde{\psi}^{j}] - \lambda_{n}^{j} \langle (\lambda_{n}^{j})^{-1} \nabla \rangle F^{\dagger}[\sigma_{n}^{j} \widetilde{\psi}^{j}] \right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}} (I) \\ &\lesssim \lim_{n\to\infty} \left\| \nabla \left\{ F^{\dagger}[\sigma_{\infty}^{j} \widetilde{\psi}^{j}] - F^{\dagger}[\sigma_{n}^{j} \widetilde{\psi}^{j}] \right\} \right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}} (I) \\ &+ \lim_{n\to\infty} \lambda_{n}^{j} \left\| \left\{ |(\lambda_{n}^{j})^{-1} \nabla| - \langle (\lambda_{n}^{j})^{-1} \nabla \rangle \right\} F^{\dagger}[\sigma_{n}^{j} \widetilde{\psi}^{j}] \right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}} (I) \end{split}$$

We consider the first term on the right-hand side of (8.94). We see from elementary computations and Hölder's inequality that

$$\begin{split} &\lim_{n\to\infty} \left\| \nabla \left\{ F^{\ddagger}[\sigma_{\infty}^{j}\widetilde{\psi}^{j}] - F^{\ddagger}[\sigma_{n}^{j}\widetilde{\psi}^{j}] \right\} \right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I)} \\ &\leq \lim_{n\to\infty} \left\| |\sigma_{\infty}^{j}\widetilde{\psi}^{j}|^{\frac{4}{d-2}} \nabla \sigma_{\infty}^{j}\widetilde{\psi}^{j} - |\sigma_{n}^{j}\widetilde{\psi}^{j}|^{\frac{4}{d-2}} \nabla \sigma_{n}^{j}\widetilde{\psi}^{j} \right\} \right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I)} \\ &+ \lim_{n\to\infty} \left\| |\sigma_{\infty}^{j}\widetilde{\psi}^{j}|^{\frac{8-2d}{d-2}} (\sigma_{\infty}^{j}\widetilde{\psi}^{j})^{2} \overline{\nabla \sigma_{\infty}^{j}\widetilde{\psi}^{j}} - |\sigma_{n}^{j}\widetilde{\psi}^{j}|^{\frac{8-2d}{d-2}} (\sigma_{n}^{j}\widetilde{\psi}^{j})^{2} \overline{\nabla \sigma_{n}^{j}\widetilde{\psi}^{j}} \right\} \right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I)}. \end{split}$$
(8.95)

Furthermore, we see from the triangle inequality, Hölder's inequality, Sobolev's embedding, Lemma 8.8, Lemma 8.10 and the assumption (8.91) that if $d \ge 6$ (hence $\frac{4}{d-2} \le 1$), then

$$\begin{split} \lim_{n \to \infty} \left\| |\sigma_{\infty}^{j} \widetilde{\psi}^{j}|^{\frac{4}{d-2}} \nabla \sigma_{\infty}^{j} \widetilde{\psi}^{j} - |\sigma_{n}^{j} \widetilde{\psi}^{j}|^{\frac{4}{d-2}} \nabla \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|_{L^{\frac{2(d+2)}{d+4}}(I)} \\ &+ \lim_{n \to \infty} \left\| |\sigma_{\infty}^{j} \widetilde{\psi}^{j}|^{\frac{8-2d}{d-2}} (\sigma_{\infty}^{j} \widetilde{\psi}^{j})^{2} \nabla \overline{\sigma_{\infty}^{j}} \widetilde{\psi}^{j} - |\sigma_{n}^{j} \widetilde{\psi}^{j}|^{\frac{8-2d}{d-2}} (\sigma_{n}^{j} \widetilde{\psi}^{j})^{2} \nabla \overline{\sigma_{n}^{j}} \widetilde{\psi}^{j} \right\|_{L^{\frac{2(d+2)}{d+4}}(I)} \\ &\lesssim \lim_{n \to \infty} \left\| |\sigma_{\infty}^{j} \widetilde{\psi}^{j}|^{\frac{4-2}{d-2}} \nabla \sigma_{\infty}^{j} \widetilde{\psi}^{j} - |\sigma_{n}^{j} \widetilde{\psi}^{j}|^{\frac{4}{d-2}} \nabla \sigma_{\infty}^{j} \widetilde{\psi}^{j} \right\|_{L^{\frac{2(d+2)}{d+4}}(I)} \\ &+ \lim_{n \to \infty} \left\| |\sigma_{\infty}^{j} \widetilde{\psi}^{j}|^{\frac{8-2d}{d-2}} (\sigma_{n}^{j} \widetilde{\psi}^{j})^{2} \nabla \overline{\sigma_{\infty}^{j}} \widetilde{\psi}^{j} - |\sigma_{n}^{j} \widetilde{\psi}^{j}|^{\frac{4}{d-2}} \nabla \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|_{L^{\frac{2(d+2)}{d+4}}(I)} \\ &+ \lim_{n \to \infty} \left\| |\sigma_{n}^{j} \widetilde{\psi}^{j}|^{\frac{4-2}{d-2}} \nabla \sigma_{\infty}^{j} \widetilde{\psi}^{j} - |\sigma_{n}^{j} \widetilde{\psi}^{j}|^{\frac{4}{d-2}} \nabla \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|_{L^{\frac{2(d+2)}{d+4}}(I)} \\ &+ \lim_{n \to \infty} \left\| |\sigma_{n}^{j} \widetilde{\psi}^{j}|^{\frac{8-2d}{d-2}} (\sigma_{n}^{j} \widetilde{\psi}^{j})^{2} \nabla \overline{\sigma_{\infty}^{j}} \widetilde{\psi}^{j} - |\sigma_{n}^{j} \widetilde{\psi}^{j}|^{\frac{8-2d}{d-2}} (\sigma_{n}^{j} \widetilde{\psi}^{j})^{2} \nabla \overline{\sigma_{n}^{j}} \widetilde{\psi}^{j} \right\|_{L^{\frac{2(d+2)}{d+4}}(I)} \\ &\leq \lim_{n \to \infty} \left\| \sigma_{n}^{j} \widetilde{\psi}^{j}|^{\frac{4-2}{d-2}} (\sigma_{n}^{j} \widetilde{\psi}^{j}) - \sigma_{\infty}^{j} \widetilde{\psi}^{j} \right\|_{V_{2}+\frac{4}{d}}(I) \\ &\leq \lim_{n \to \infty} \left\| \sigma_{n}^{j} \widetilde{\psi}^{j}|^{\frac{4-2}{d-2}} (\sigma_{n}^{j} \widetilde{\psi}^{j}) - \sigma_{\infty}^{j} \widetilde{\psi}^{j} \right\|_{V_{2}+\frac{4}{d}}(I) \\ &\leq \lim_{n \to \infty} \left\| \sigma_{n}^{j} \widetilde{\psi}^{j}|^{\frac{4-2}{d-2}} (\sigma_{n}^{j} \widetilde{\psi}^{j}) - \sigma_{\infty}^{j} \widetilde{\psi}^{j} \right\|_{V_{2}+\frac{4}{d}}(I) \\ &= 0. \end{split}$$

Similarly, we can verify that if $3 \le d \le 5$ (hence $1 < \frac{4}{d-2} = 1 + \frac{6-d}{d-2}$), then

$$\begin{split} \lim_{n \to \infty} \left\| \left\| \sigma_{\infty}^{j} \widetilde{\psi}^{j} \right\|^{\frac{4}{d-2}} \nabla \sigma_{\infty}^{j} \widetilde{\psi}^{j} - \left\| \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|^{\frac{4}{d-2}} \nabla \sigma_{n}^{j} \widetilde{\psi}^{j} \right\} \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(I)} \\ &+ \lim_{n \to \infty} \left\| \left\| \sigma_{\infty}^{j} \widetilde{\psi}^{j} \right\|^{\frac{8-2d}{d-2}} (\sigma_{\infty}^{j} \widetilde{\psi}^{j})^{2} \nabla \sigma_{\infty}^{j} \widetilde{\psi}^{j} - \left\| \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|^{\frac{8-2d}{d-2}} (\sigma_{n}^{j} \widetilde{\psi}^{j})^{2} \nabla \sigma_{n}^{j} \widetilde{\psi}^{j} \right\} \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(I)} \\ &\lesssim \lim_{n \to \infty} \left\{ \left\| \sigma_{\infty}^{j} \widetilde{\psi}^{j} \right\|^{\frac{6-d}{d-2}}_{W_{2^{*}}(I)} + \left\| \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|^{\frac{6-d}{d-2}}_{W_{2^{*}}(I)} \right\} \right\| \sigma_{\infty}^{j} \widetilde{\psi}^{j} - \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|_{W_{2^{*}}(I)} \left\| \langle \nabla \rangle \widetilde{\psi}^{j} \right\|_{V_{2+\frac{4}{d}}(I)} \\ &+ \lim_{n \to \infty} \left\| \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|^{\frac{4}{d-2}}_{W_{2^{*}}(I)} \left\| \nabla \left\{ \sigma_{\infty}^{j} \widetilde{\psi}^{j} - \sigma_{\infty}^{j} \widetilde{\psi}^{j} \right\} \right\|_{V_{2+\frac{4}{d}}(I)} \\ &= 0. \end{split}$$

Next, we consider the second term on the right-hand side of (8.94). We see from the Mihlin multiplier theorem, the assumption (8.91) and Lemma 8.10 that

$$\begin{split} &\lim_{n\to\infty}\lambda_n^j \left\|\left\{|(\lambda_n^j)^{-1}\nabla|-\langle(\lambda_n^j)^{-1}\nabla\rangle\right\}F^{\ddagger}[\sigma_n^j\widetilde{\psi}^j]\right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I)} \\ &\lesssim &\lim_{n\to\infty}\lambda_n^j \left\|F^{\ddagger}[\sigma_n^j\widetilde{\psi}^j]\right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I)} \lesssim &\lim_{n\to\infty} \left\|\sigma_n^j\widetilde{\psi}^j\right\|_{W_{2^*}(I)}^{\frac{4}{d-2}} \|\lambda_n^j\sigma_n^j\widetilde{\psi}^j\|_{V_{2+\frac{4}{d}}(I)} = 0. \end{split}$$

$$(8.98)$$

Putting (8.94) through (8.98) together, we find that

$$\lim_{n \to \infty} \left\| \langle \nabla \rangle (\sigma_{\infty}^j)^{-1} F^{\ddagger}[G_n^j \sigma_{\infty}^j \widetilde{\psi}^j] - \langle \nabla \rangle F^{\ddagger}[G_n^j \sigma_n^j \widetilde{\psi}^j] \right\|_{L^{\frac{2(d+2)}{d+4}}(I_n^j)} = 0.$$
(8.99)

We can also verify in a way similar to the estimates (8.94) and (8.98) that

$$\begin{split} \lim_{n \to \infty} \left\| \langle \nabla \rangle F^{\dagger}[G_{n}^{j} \sigma_{n}^{j} \widetilde{\psi}^{j}] \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(l_{n}^{j})} \\ &= \lim_{n \to \infty} (\lambda_{n}^{j})^{\frac{4-(d-2)(p-1)}{2}} \left\| \lambda_{n}^{j} \langle (\lambda_{n}^{j})^{-1} \nabla \rangle F^{\dagger}[\sigma_{n}^{j} \widetilde{\psi}^{j}] \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(l_{n}^{j})} \\ &\lesssim \lim_{n \to \infty} (\lambda_{n}^{j})^{\frac{4-(d-2)(p-1)}{2}} \lambda_{n}^{j} \left\| \{ |(\lambda_{n}^{j})^{-1} \nabla | - \langle (\lambda_{n}^{j})^{-1} \nabla \rangle \} F^{\dagger}[\sigma_{n}^{j} \widetilde{\psi}^{j}] \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(l_{n}^{j})} \\ &+ \lim_{n \to \infty} (\lambda_{n}^{j})^{\frac{4-(d-2)(p-1)}{2}} \left\| \nabla F^{\dagger}[\sigma_{n}^{j} \widetilde{\psi}^{j}] \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(l_{n}^{j})} \\ &\lesssim \lim_{n \to \infty} (\lambda_{n}^{j})^{\frac{4-(d-2)(p-1)}{2}} \left\| \nabla F^{\dagger}[\sigma_{n}^{j} \widetilde{\psi}^{j}] \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(l_{n}^{j})} \\ &\lesssim \lim_{n \to \infty} (\lambda_{n}^{j})^{\frac{4-(d-2)(p-1)}{2}} \left\| \nabla F^{\dagger}[\sigma_{n}^{j} \widetilde{\psi}^{j}] \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(l_{n}^{j})} \\ &\lesssim \lim_{n \to \infty} (\lambda_{n}^{j})^{\frac{4-(d-2)(p-1)}{2}} \left\| \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|_{W_{p+1}(l)}^{p-1} \left\| \lambda_{n}^{j} \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|_{V_{2}+\frac{4}{d}}(l)} \\ &+ \lim_{n \to \infty} (\lambda_{n}^{j})^{\frac{4-(d-2)(p-1)}{2}} \left\| \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|_{W_{p+1}(l)}^{p-1} \left\| \nabla \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|_{V_{2}+\frac{4}{d}}(l)} \\ &\leq \lim_{n \to \infty} \left\| \lambda_{n}^{j} \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|_{V_{2}+\frac{4}{d}}(l)}^{(p-1)(1-s_{p})} \left\| \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|_{W_{2}*(l)}^{(p-1)s_{p}} \left\| \nabla \sigma_{n}^{j} \widetilde{\psi}^{j} \right\|_{V_{2}+\frac{4}{d}}(l)} \\ &= 0. \end{split}$$

Then, the desired result (8.92) follows from (8.99) and (8.100).

Lemma 8.12. For any $\delta > 0$ and any $k \ge 1$, there exists a number $N(\delta, k)$ such that for any $n \ge N(\delta, k)$,

$$\sum_{\substack{1 \le j \le k \\ \lambda_{\infty}^j = \infty}} \| \langle \nabla \rangle F[\psi_n^j] \|_{L^{2(d+2)}_{t,x}(\mathbb{R})} \le \delta,$$
(8.101)

where the sum is taken over all integers j satisfying $1 \leq j \leq k$ and $\lambda_{\infty}^{j} = \infty$.

Proof of Lemma 8.12. Let q denote p or $2^* - 1$, and let $j \ge 1$ be a number for which $\lambda_{\infty}^j = \infty$, so that $\psi_n^j = G_n^j \sigma_n^j e^{it\Delta} \tilde{u}^j$. Then, we see from Hölder's inequality, Sobolev's

embedding, Strichartz' estimate, Lemma 8.7 with $\sigma_{\infty}^{j} = 0$ (see (8.56)) that

$$\lim_{n \to \infty} \| |\psi_{n}^{j}|^{q-1} \psi_{n}^{j} \|_{L^{2(d+2)}_{t,x}}^{2(d+2)}(\mathbb{R}) \leq \lim_{n \to \infty} \| \psi_{n}^{j} \|_{W_{q+1}(I_{n})}^{q-1} \| \psi_{n}^{j} \|_{V_{2+\frac{4}{d}}}(\mathbb{R}) \leq \lim_{n \to \infty} (\lambda_{n}^{j})^{(q-1)(1-s_{q})} \| |\nabla|^{s_{q}} \sigma_{n}^{j} e^{it\Delta} \widetilde{u}^{j} \|_{V_{q+1}(I_{n})}^{q-1} \lambda_{n}^{j} \| \sigma_{n}^{j} e^{it\Delta} \widetilde{u}^{j} \|_{V_{2+\frac{4}{d}}}(\mathbb{R}) \leq \lim_{n \to \infty} (\lambda_{n}^{j})^{(q-1)(1-s_{q})} \| |\nabla|^{s_{q}} \sigma_{n}^{j} \widetilde{u}^{j} \|_{L^{2}}^{q-1} \| \lambda_{n}^{j} \sigma_{n}^{j} \widetilde{u}^{j} \|_{L^{2}} \leq \lim_{n \to \infty} \| \lambda_{n}^{j} \sigma_{n}^{j} \widetilde{u}^{j} \|_{L^{2}}^{(q-1)(1-s_{q})} \| \nabla \sigma_{n}^{j} \widetilde{u}^{j} \|_{L^{2}}^{(q-1)s_{q}} \| \lambda_{n}^{j} \sigma_{n}^{j} \widetilde{u}^{j} \|_{L^{2}} = 0.$$
(8.102)

Similarly, we can verify that

$$\lim_{n \to \infty} \|\nabla\{|\psi_n^j|^{q-1}\psi_n^j\}\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(\mathbb{R})} \lesssim \lim_{n \to \infty} \||\psi_n^j|^{q-1} |\nabla\psi_n^j|\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(\mathbb{R})} \leq \lim_{n \to \infty} \|\psi_n^j\|_{W_{q+1}(I_n)}^{q-1} \|\nabla\psi_n^j\|_{V_{2+\frac{4}{d}}(\mathbb{R})} = 0.$$
(8.103)

Thus, we find from (8.102) and (8.103) that for any $\delta > 0$ and any $k \ge 1$, there exists $N(\delta, k) \ge 1$ such that for any $n \ge N(\delta, k)$,

$$\sum_{\substack{1 \le j \le k \\ j_{\infty}^{j} = \infty}} \| \langle \nabla \rangle \big\{ |\psi_{n}^{j}|^{q-2} \psi_{n}^{j} \big\} \|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(\mathbb{R})} \le \delta,$$
(8.104)

which gives us the desired result.

Lemma 8.13. Let $j \ge 1$, and let I be an interval on which $\tilde{\psi}^j$ exists. Then, we have

$$\sup_{n\geq 1} \|\langle \nabla \rangle \psi_n^j \|_{St(I_n^j)} \lesssim \|\langle \nabla \rangle \widetilde{\psi}^j \|_{St(I)}, \tag{8.105}$$

where $I_n^j := ((\lambda_n^j)^2 \inf I + t_n^j, \ (\lambda_n^j)^2 \sup I + t_n^j).$

Proof of Lemma 8.13. We can easily verify that

$$\|\langle \nabla \rangle \psi_n^j \|_{L^\infty_t L^2_x(I^j_n)} \le \|\lambda_n^j \sigma_n^j \widetilde{\psi}^j \|_{L^\infty_t L^2_x(I)} + \|\nabla \sigma_n^j \widetilde{\psi}^j \|_{L^\infty_t L^2_x(I)}, \tag{8.106}$$

$$\|\langle \nabla \rangle \psi_n^j \|_{L^2_t L^{2^*}_x(I_n^j)} \le \|\lambda_n^j \sigma_n^j \widetilde{\psi}^j \|_{L^2_t L^{2^*}_x(I)} + \|\nabla \sigma_n^j \widetilde{\psi}^j \|_{L^2_t L^{2^*}_x(I)}.$$
(8.107)

When $\lambda_{\infty}^{j} \in \{0, 1\}$, these estimates (8.106) and (8.107) together with Lemma 8.8 prove the lemma. Assume $\lambda_{\infty}^{j} = \infty$, so that $\tilde{\psi}^{j}(t) = e^{it\Delta}\tilde{u}^{j}$. Then, we see from (8.106), (8.107) and Strichartz' estimate that

$$\begin{aligned} \|\langle \nabla \rangle \psi_n^j \|_{L_t^\infty L_x^2(I_n^j)} + \|\langle \nabla \rangle \psi_n^j \|_{L_t^2 L_x^{2^*}(I_n^j)} &\lesssim \|\lambda_n^j \sigma_n^j \widetilde{u}^j \|_{L^2} + \|\nabla \sigma_n^j \widetilde{u}^j \|_{L^2} \\ &\lesssim \|\langle \nabla \rangle \widetilde{u}^j \|_{L^2} = \|\langle \nabla \rangle \widetilde{\psi}^j \|_{L_t^\infty L_x^2(I)}. \end{aligned}$$

$$(8.108)$$

Thus, we have completed the proof.

Lemma 8.14 (cf. Lemma 6.8 in [3]). There exists a number J_0 such that $I_{\max}^j = \mathbb{R}$ for any $j > J_0$; and for any $r \ge 2$,

$$\sum_{j>J_0} \|\langle \nabla \rangle \widetilde{\psi}^j \|_{St(\mathbb{R})}^r \lesssim \sum_{j>J_0} \|\widetilde{u}^j\|_{H^1}^r < \infty.$$
(8.109)

Proof of Lemma 8.14. It follows from (8.49) and the uniform boundedness (8.38) that

$$\lim_{n \to \infty} \sum_{j=1}^{k} \left\| g_{n}^{j} \sigma_{n}^{j} e^{-i \frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \Delta} \widetilde{u}^{j} \right\|_{H^{1}}^{2} \lesssim 1$$
(8.110)

for all $k \geq 1$. Furthermore, we see from Lemma 8.7 that: if $\lambda_{\infty}^{j} = 0$, then

$$\lim_{n \to \infty} \left\| g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right\|_{L^2} = \lim_{n \to \infty} \left\| \lambda_n^j \sigma_n^j \widetilde{u}^j \right\|_{L^2} = 0,$$
(8.111)

$$\lim_{n \to \infty} \left\| \nabla g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right\|_{L^2} = \lim_{n \to \infty} \left\| \nabla \sigma_n^j \widetilde{u}^j \right\|_{L^2} = \| \widetilde{u}^j \|_{H^1}; \tag{8.112}$$

and if $\lambda_{\infty}^{j} \in \{1, \infty\}$, then

$$\lim_{n \to \infty} \left\| g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right\|_{H^1} = \| \widetilde{u}^j \|_{H^1}.$$
(8.113)

Putting (8.110) through (8.113) together, we obtain that

$$\sum_{j=1}^{\infty} \|\widetilde{u}^{j}\|_{H^{1}}^{2} = \lim_{k \to \infty} \sum_{j=1}^{k} \|\widetilde{u}^{j}\|_{H^{1}}^{2} \lesssim \lim_{k \to \infty} \sum_{j=1}^{k} \lim_{n \to \infty} \left\| g_{n}^{j} \sigma_{n}^{j} e^{-i\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \Delta} \widetilde{u}^{j} \right\|_{H^{1}}^{2} \lesssim 1.$$
(8.114)

In particular, for any $\delta > 0$, we can take $J(\delta) \ge 1$ such that for any $j \ge J(\delta)$,

$$\|\widetilde{u}^j\|_{H^1} < \delta. \tag{8.115}$$

Note here that if $\lambda_{\infty}^{j} = 0$, then (8.115) is rewritten as $\|\nabla \sigma_{\infty}^{j} \widetilde{u}^{j}\|_{L^{2}}^{2} < \delta$. Since $\widetilde{\psi}^{j}$ satisfies (8.74) and (8.76), the small-data theory (Lemma D.1) together with (8.115) implies that there exists a number J_{0} such that for any $j \geq J_{0}$,

$$\|\langle \nabla \rangle \widetilde{\psi}^{j}\|_{St(\mathbb{R})} \lesssim \|\widetilde{u}^{j}\|_{H^{1}}.$$
(8.116)

Then, the claim (8.109) follows from (8.114), (8.116) and Jensen's inequality.

Lemma 8.15. For any number $k \ge 1$, there exists a number N(k) such that for any $n \ge N(k)$,

$$\sum_{j=1}^{k} \mathcal{I}_{\omega} \left(g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right) + \mathcal{I}_{\omega} (w_n^k) \le m_{\omega} - \frac{\kappa_1(R_*)}{10d(p-1)}.$$
(8.117)

Proof of Lemma 8.15. It follows from (8.32), (8.37) and $S_{\omega}(\psi_n) \leq m_{\omega} + \varepsilon_n$ that for any numbers n and k,

$$m_{\omega} - \frac{\kappa_1(R_*)}{2d(p-1)} \ge m_{\omega} + \varepsilon_n - \frac{2}{d(p-1)} \mathcal{K}(\psi_n(0))$$

$$\ge \mathcal{S}_{\omega}(\psi_n) - \frac{2}{d(p-1)} \mathcal{K}(\psi_n(0)) = \mathcal{I}_{\omega}(\psi_n(0)).$$
(8.118)

This together with (8.53) gives us the desired result (8.117).

Lemma 8.16. For any number $k \ge 1$, there exists a number N(k) such that for any $n \ge N(k)$ and any $j \le k$ for which the linear profile \tilde{u}^j is non-trivial,

$$\mathcal{H}\left(g_{n}^{j}\sigma_{n}^{j}e^{-i\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\Delta}\widetilde{u}^{j}\right) > \frac{1}{2}\mathcal{K}\left(g_{n}^{j}\sigma_{n}^{j}e^{-i\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\Delta}\widetilde{u}^{j}\right) > 0,$$
(8.119)

$$\mathcal{H}(w_n^k) \ge \frac{1}{2}\mathcal{K}(w_n^k) \ge 0.$$
(8.120)

Proof of Lemma 8.16. Since \mathcal{I}_{ω} is non-negative, Lemma 8.15 together with (1.10) shows that for any $k \geq 1$, there exists a number N(k) such that for any $n \geq N(k)$ and any $j \leq k$ for which \tilde{u}^{j} is non-trivial,

$$\mathcal{K}\left(g_{n}^{j}\sigma_{n}^{j}e^{-i\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\Delta}\widetilde{u}^{j}\right) > 0, \quad \mathcal{K}(w_{n}^{k}) \ge 0.$$
(8.121)

This together with (1.6) gives us the desired result.

Lemma 8.17. There is at most one number j_0 such that $\lambda_{\infty}^{j_0} \in \{0,1\}$ and

$$\begin{cases} \mathcal{S}_{\omega}(\sigma_{\infty}^{j_{0}}\widetilde{\psi}^{j_{0}}) \geq m_{\omega} & \text{if } \lambda_{\infty}^{j_{0}} = 1, \\ \mathcal{H}^{\ddagger}(\sigma_{\infty}^{j_{0}}\widetilde{\psi}^{j_{0}}) \geq \mathcal{H}^{\ddagger}(W) & \text{if } \lambda_{\infty}^{j_{0}} = 0. \end{cases}$$

$$(8.122)$$

Proof of Lemma 8.17. First, we consider a number k such that

$$\mathcal{S}_{\omega}\left(g_{n}^{k}\sigma_{n}^{k}e^{-i\frac{t_{n}^{k}}{(\lambda_{n}^{k})^{2}}\Delta}\widetilde{u}^{k}\right) \leq \frac{2}{3}m_{\omega}.$$
(8.123)

Then, it follows from Lemma 8.7, Lemma 8.9, (8.74), (8.79), (1.34) and (1.35) that for any sufficiently large number n,

$$\begin{cases} \mathcal{S}_{\omega}(\sigma_{\infty}^{k}\widetilde{\psi}^{k}) = \mathcal{S}_{\omega}\left(\sigma_{\infty}^{k}\widetilde{\psi}^{k}\left(-\frac{t_{n}^{k}}{(\lambda_{n}^{k})^{2}}\right)\right) < m_{\omega} & \text{if } \lambda_{\infty}^{k} = 1, \\ \mathcal{H}^{\ddagger}(\sigma_{\infty}^{k}\widetilde{\psi}^{k}) = \mathcal{H}^{\ddagger}\left(\sigma_{\infty}^{k}\widetilde{\psi}^{k}\left(-\frac{t_{n}^{k}}{(\lambda_{n}^{k})^{2}}\right)\right) < \mathcal{H}^{\ddagger}(W) & \text{if } \lambda_{\infty}^{k} = 0. \end{cases}$$

$$(8.124)$$

Thus, it suffices for the desired result to show that there is at most one number j_0 such that for any sufficiently large n,

$$\mathcal{S}_{\omega}\left(g_{n}^{j_{0}}\sigma_{n}^{j_{0}}e^{-i\frac{t_{n}^{j_{0}}}{(\lambda_{n}^{j_{0}})^{2}}\Delta}\widetilde{u}^{j_{0}}\right) \geq \frac{2}{3}m_{\omega}.$$
(8.125)

We see from Lemma 8.16 that for any $j \ge 1$ and any sufficiently large n,

$$\mathcal{S}_{\omega}\left(g_{n}^{j}\sigma_{n}^{j}e^{-i\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\Delta}\widetilde{u}^{j}\right) \geq 0.$$
(8.126)

Hence, it follows from (8.52) and $S_{\omega}(\psi_n(0)) < m_{\omega} + \varepsilon_n$ that there are at most one linear profile satisfying (8.125).

Now, using Lemma 8.17 and reordering the indices, we may assume that for any $j \ge 2$,

$$\begin{cases} \mathcal{S}_{\omega}(\sigma_{\omega}^{j}\widetilde{\psi}^{j}) < m_{\omega} & \text{if } \lambda_{\infty}^{j} = 1, \\ \mathcal{H}^{\ddagger}(\sigma_{\infty}^{j}\widetilde{\psi}^{j}) < \mathcal{H}^{\ddagger}(W) & \text{if } \lambda_{\infty}^{j} = 0. \end{cases}$$

$$(8.127)$$

Lemma 8.18. Assume (8.127). Then, for any $j \ge 1$ for which the nonlinear profile $\tilde{\psi}^{j}$ is non-trivial and $\lambda_{\infty}^{j} \in \{0,1\}$, there exists a number N(j) such that for any $n \ge N(j)$,

$$\begin{cases} \mathcal{K}\big(\widetilde{\psi}^{j}\Big(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\Big)\big) > 0 & \text{if } \lambda_{\infty}^{j} = 1, \\ \mathcal{K}^{\ddagger}\big(\sigma_{\infty}^{j}\widetilde{\psi}^{j}\Big(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\Big) > 0 & \text{if } \lambda_{\infty}^{j} = 0. \end{cases}$$

$$(8.128)$$

Furthermore, if $\lambda_{\infty}^{j} = 0$ and $\mathcal{H}^{\ddagger}(\sigma_{\infty}^{j}\widetilde{\psi}^{j}) < \mathcal{H}^{\ddagger}(W)$, then for any $n \geq N(j)$,

$$\left\|\nabla \sigma_{\infty}^{j} \widetilde{\psi}^{j} \left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\right)\right\|_{L^{2}}^{2} < \|\nabla W\|_{L^{2}}^{2}.$$
(8.129)

Proof of Lemma 8.18. We see from Lemma 8.15 that there exists a number N(j) such that for any $n \ge N(j)$,

$$\mathcal{I}_{\omega}\left(g_{n}^{j}\sigma_{n}^{j}e^{-i\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\Delta}\widetilde{u}^{j}\right) \leq m_{\omega} - \frac{\kappa_{1}(R_{*})}{100d(p-1)}.$$
(8.130)

When $\lambda_{\infty}^{j} = 1$, the desired result (8.128) follows from (8.79), (8.130) and (1.10). In order to prove (8.128) in the case $\lambda_{\infty}^{j} = 0$, note that it follows from (1.30) and (8.130) that for any $n \geq N(j)$,

$$\mathcal{I}^{\ddagger}_{\omega} \left(g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right) = \mathcal{H}^{\ddagger} \left(g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right) - \frac{2}{d(p-1)} \mathcal{K}^{\ddagger} \left(g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right) \\
\leq \mathcal{S}_{\omega} \left(g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right) - \frac{2}{d(p-1)} \mathcal{K} \left(g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right) \\
= \mathcal{I}_{\omega} \left(g_n^j \sigma_n^j e^{-i \frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j \right) \leq m_{\omega} - \frac{\kappa_1(R_*)}{100d(p-1)}.$$
(8.131)

Furthermore, we see from (8.131) and (8.79) that for any $n \ge N(j)$,

$$\mathcal{I}^{\ddagger}_{\omega}\left(\sigma^{j}_{\infty}\widetilde{\psi}^{j}\left(-\frac{t^{j}_{n}}{(\lambda^{j}_{n})^{2}}\right)\right) = \lim_{n \to \infty} \mathcal{I}^{\ddagger}\left(g^{j}_{n}\sigma^{j}_{n}e^{-i\frac{t^{j}_{n}}{(\lambda^{j}_{n})^{2}}}\widetilde{u}^{j}\right) \le m_{\omega} - \frac{\kappa_{1}(R_{*})}{100d(p-1)}.$$
(8.132)

Thus, when $\lambda_{\infty}^{j} = 0$, (1.29) together with (1.34), (1.35) and (8.132) shows that for any $n \geq N(j)$,

$$\mathcal{K}^{\ddagger}\left(\sigma_{\infty}^{j}\widetilde{\psi}^{j}\left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\right)\right) = \left\|\nabla\sigma_{\infty}^{j}\widetilde{\psi}^{j}\left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\right)\right\|_{L^{2}}^{2} - \left\|\sigma_{\infty}^{j}\widetilde{\psi}^{j}\left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\right)\right\|_{L^{2^{*}}}^{2^{*}} > 0, \quad (8.133)$$

which completes the proof of (8.128).

Next, we shall show (8.129). Assume $\lambda_{\infty}^{j} = 0$ and $\mathcal{H}^{\ddagger}(\sigma_{\infty}^{j}\widetilde{\psi}^{j}) < \mathcal{H}^{\ddagger}(W)$. Then, we see from (8.133), (1.34) and (1.28) that for any $n \geq N(j)$,

$$\frac{1}{d} \| \nabla \sigma_{\infty}^{j} \widetilde{\psi}^{j} \left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \right) \|_{L^{2}}^{2}
= \frac{1}{2} \| \nabla \sigma_{\infty}^{j} \widetilde{\psi}^{j} \left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \right) \|_{L^{2}}^{2} - \frac{1}{2^{*}} \| \nabla \sigma_{\infty}^{j} \widetilde{\psi}^{j} \left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \right) \|_{L^{2}}^{2}
\leq \frac{1}{2} \| \nabla \sigma_{\infty}^{j} \widetilde{\psi}^{j} \left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \right) \|_{L^{2}}^{2} - \frac{1}{2^{*}} \| \sigma_{\infty}^{j} \widetilde{\psi}^{j} \left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \right) \|_{L^{2^{*}}}^{2^{*}}
= \mathcal{H}^{\ddagger} \left(\sigma_{\infty}^{j} \widetilde{\psi}^{j} \right) < \mathcal{H}^{\ddagger}(W) = \frac{1}{d} \| \nabla W \|_{L^{2}}^{2}.$$
(8.134)

Thus, we have completed the proof.

We see from Lemma 8.18 and (8.127) that for any $j \ge 2$ for which $\tilde{\psi}^j$ is non-trivial and $\lambda_{\infty}^j \neq \infty$,

$$\begin{cases} \sigma_{\infty}^{j} \widetilde{\psi}^{j} \in PW_{\omega,+} & \text{if } \lambda_{\infty}^{j} = 1, \\ \sigma_{\infty}^{j} \widetilde{\psi}^{j} \in PW_{+}^{\ddagger} & \text{if } \lambda_{\infty}^{j} = 0. \end{cases}$$

$$(8.135)$$

Furthermore, since $\tilde{\psi}^j(t) = e^{it\Delta}\tilde{u}^j$ for $j \ge 1$ with $\lambda_{\infty}^j = \infty$, it follows from Theorem 1.2, Theorem 1.3 and Lemma 8.14 that $I_{\max}^j = \mathbb{R}$ for any $j \ge 2$, and

$$\sup_{j\geq 2} \|\langle \nabla \rangle \widetilde{\psi}^j \|_{St(\mathbb{R})} < \infty.$$
(8.136)

8.2.4 Existence of bad profile

In the previous section, we showed that there are at most one bad profile (see Lemma 8.17), and the candidate is $\tilde{\psi}^1$ (see (8.136)). Our aim in this section is to show that for any $T \in (T_{\min}^1, T_{\max}^1)$,

$$\|\sigma_{\infty}^{1}\widetilde{\psi}^{1}\|_{\mathcal{W}^{1}([T,T_{\max}^{1}))} = \infty, \qquad (8.137)$$

where \mathcal{W}^1 is the function space defined by (8.73) (see Proposition 8.25 below). To this end, we observe properties of $\tilde{\psi}^1$ on an interval where

$$\|\sigma_{\infty}^{1}\widetilde{\psi}^{1}\|_{\mathcal{W}^{1}(I)} < \infty.$$
(8.138)

Lemma 8.19. For any interval I satisfying (8.138), we can take a constant A(I) > 0 such that

$$\|\langle \nabla \rangle \widetilde{\psi}^1\|_{St(I)} \le A(I). \tag{8.139}$$

Proof of Lemma 8.19. When $\lambda_{\infty}^1 = \infty$, $\tilde{\psi}^1(t) = e^{it\Delta}\tilde{u}^1$ and therefore the claim follows from Strichartz' estimate. Moreover, when I is a compact interval in I_{\max}^1 , the claim follows from the well-posedness theory. Thus, we may assume that $\lambda_{\infty}^j \in \{0, 1\}$, and $\sup I = T_{\max}^1$ or $\inf I = T_{\min}^1$. We only consider the case where $\sup I = T_{\max}^1$ and $T_{\min}^1 < \inf I$. The same proof is applicable for the other cases. Since $\sup I = T_{\max}^1$, the blowup criterion (cf. Theorem 4.1 in [3] and Lemma 2.11 in [18]) together with (8.138) shows that $T_{\max}^1 = \infty$ and

$$\|\sigma_{\infty}^{1}\widetilde{\psi}^{1}\|_{\mathcal{W}^{1}([\inf I,\infty))} < \infty.$$
(8.140)

We shall show that

$$\sup_{\mathbf{t}\in[\inf I,\infty)} \|\widetilde{\psi}^1(t)\|_{H^1} < \infty.$$
(8.141)

Suppose for contradiction that (8.141) was false. Then, we could take a sequence $\{t_n\}$ in $[\inf I, \infty)$ such that $\lim_{n\to\infty} t_n = \infty$ and

$$\lim_{n \to \infty} \left\| \widetilde{\psi}^1(t_n) \right\|_{H^1} = \infty.$$
(8.142)

Assume $\lambda_{\infty}^1 = 1$. Then, $\tilde{\psi}^1$ is a solution to (NLS), and we see from Strichartz' estimate and Hölder's inequality that for any $t_m < t_n$,

$$\begin{split} \|e^{-it_{m}\Delta}\widetilde{\psi}^{1}(t_{m}) - e^{-it_{n}\Delta}\widetilde{\psi}^{1}(t_{n})\|_{H^{1}} &\leq \sup_{T \in [t_{m}, t_{n}]} \left\| \int_{t_{m}}^{T} e^{-it\Delta}F[\widetilde{\psi}^{1}(t)] dt \right\|_{H^{1}} \\ &\lesssim \|\widetilde{\psi}^{1}\|_{W_{p+1}([t_{m}, \infty))}^{p-1} \|\langle \nabla \rangle \widetilde{\psi}^{1}\|_{V_{p+1}([t_{m}, \infty))} + \|\widetilde{\psi}^{1}\|_{W_{2^{*}}([t_{m}, \infty))}^{\frac{4}{d-2}} \|\langle \nabla \rangle \widetilde{\psi}^{1}\|_{V_{2^{*}}([t_{m}, \infty))}. \end{split}$$

$$(8.143)$$

Moreover, we see from (8.140) and $\sigma_{\infty}^1 = 1$ that

$$\lim_{T \to \infty} \| \widetilde{\psi}^1 \|_{W_{p+1}([T,\infty)) \cap W_{2^*}([T,\infty))} = 0.$$
(8.144)

In particular, for any $\delta > 0$, there exists $T(\delta) > 0$ such that

$$\|\widetilde{\psi}^1\|_{W_{p+1}([T(\delta),\infty))\cap W_{2^*}([T(\delta),\infty))} \le \delta.$$
(8.145)

An estimate similar to (8.143) together with (8.145) also yields that

$$\|\langle \nabla \rangle \psi^{1} \|_{V_{p+1}([T(\delta),\infty)) \cap V_{2^{*}}([T(\delta),\infty))}$$

$$\lesssim \|\widetilde{\psi}^{1}(T(\delta))\|_{H^{1}} + (\delta^{p-1} + \delta^{\frac{4}{d-2}}) \|\langle \nabla \rangle \widetilde{\psi}^{1} \|_{V_{p+1}([T(\delta),\infty)) \cap V_{2^{*}}([T(\delta),\infty))}.$$

$$(8.146)$$

Thus, we can take $\delta_0 > 0$ such that

$$\|\langle \nabla \rangle \widetilde{\psi}^{1}\|_{V_{p+1}([T(\delta_{0}),\infty)) \cap V_{2^{*}}([T(\delta_{0}),\infty))} \lesssim \|\widetilde{\psi}^{1}(T(\delta_{0}))\|_{H^{1}}.$$
(8.147)

Combining (8.143) with (8.144) and (8.147), we find that $\{e^{-it_n\Delta}\widetilde{\psi}^1(t_n)\}$ is a Cauchy sequence in $H^1(\mathbb{R}^d)$. Similarly, when $\lambda_{\infty}^1 = 0$, we can verify that $\{e^{-it_n\Delta}\sigma_{\infty}^1\widetilde{\psi}^1(t_n)\}$ is

a Cauchy sequence in $\dot{H}^1(\mathbb{R}^d)$. Thus, in both cases $\lambda_{\infty}^1 = 1$ and $\lambda_{\infty}^1 = 0$, we can take $\phi_+ \in H^1(\mathbb{R}^d)$ such that

$$\lim_{n \to \infty} \| \tilde{\psi}^1(t_n) \|_{H^1} = \| \phi_+ \|_{H^1} < \infty.$$
(8.148)

However, this contradicts (8.142). Hence, we have proved (8.141).

Now, we are able to show (8.139). If $\lambda_{\infty}^1 = 1$, then $e[\tilde{\psi}^1] \equiv 0$ (see (1.66) for the definition of $e[\tilde{\psi}^1]$), and Lemma D.2 together with (8.140) and (8.141) gives us the desired estimate (8.139). If $\lambda_{\infty}^1 = 0$, then $e^{\ddagger}[\sigma_{\infty}^1 \tilde{\psi}^1] \equiv 0$. Moreover, it follows from (8.73), (8.140) and (8.141) that

$$\|\sigma_{\infty}^{1}\widetilde{\psi}^{1}\|_{W_{2^{*}}([\inf I,\infty))} < \infty, \quad \sup_{t \in [\inf I,\infty)} \|\nabla\sigma_{\infty}^{1}\widetilde{\psi}^{1}\|_{L^{2}} < \infty.$$

$$(8.149)$$

Hence, we see from Lemma D.2 that

$$\|\langle \nabla \rangle \widetilde{\psi}^1\|_{St([\inf I,\infty))} = \|\nabla \sigma^1_{\infty} \widetilde{\psi}^1\|_{St([\inf I,\infty))} < \infty,$$
(8.150)

which completes the proof.

Since the estimate (8.42) is insufficient to control the remainder $e^{it\Delta}w_n^j$ in the Strichartz space $\langle \nabla \rangle^{-1}St(\mathbb{R})$, we need the following estimate:

Lemma 8.20. Assume (8.127), and let q denote p or $2^* - 1$. Then, for any $\delta > 0$, any interval I satisfying (8.138), and any number $j \ge 1$, there exists a number $K(\delta, I, j)$ with the following property: for any $k \ge K(\delta, I, j)$, there exists a number $N(\delta, I, j, k)$ such that for any $n \ge N(\delta, I, j, k)$ and $s \in \{0, 1\}$,

$$\left\|\psi_{n}^{j}|\nabla|^{s}e^{it\Delta}w_{n}^{k}\right\|_{L^{\frac{2(d+2)(q-1)}{d(q-1)+4}}_{t,x}(I_{n})} \leq \delta,$$
(8.151)

where $I_n := ((\lambda_n^1)^2 \inf I + t_n^1, \ (\lambda_n^1)^2 \sup I + t_n^1).$

Proof of Lemma 8.20. Let I be an interval on which (8.138) holds. Put $I^1 := I$ and $I^j := \mathbb{R}$ for $j \geq 2$. Then, it follows from Lemma 8.19 and (8.136) that there exists a constant A(I) > 0 such that

$$\sup_{j\geq 1} \|\langle \nabla \rangle \widetilde{\psi}^j\|_{St(I^j)} \le A(I).$$
(8.152)

Moreover, we see from (8.80) and elementary computations that for any numbers $j, k, n \ge 1$ and $s \in \{0, 1\}$,

$$\begin{aligned} \|\psi_{n}^{j}|\nabla|^{s}e^{it\Delta}w_{n}^{k}\|_{L^{\frac{2(d+2)(q-1)}{d(q-1)+4}}_{t,x}(I_{n})} \\ &\leq (\lambda_{n}^{j})^{2-s_{q}}\|\sigma_{n}^{j}\widetilde{\psi}^{j}(G_{n}^{j})^{-1}|\nabla|^{s}e^{it\Delta}w_{n}^{k}\|_{L^{\frac{2(d+2)(q-1)}{d(q-1)+4}}_{t,x}(I^{j})}. \end{aligned}$$

$$\tag{8.153}$$

We consider the right-hand side of (8.153) according to s and λ_{∞}^{j} :

Case 1.1. Assume s = 0 and $\lambda_{\infty}^{j} \in \{0, 1\}$ in (8.153). Then, we see from Hölder's inequality, Lemma 8.8, (8.152) and (8.55) that there exists a number $K_1(\delta, I)$ with the following property: for any $k \ge K_1(\delta, I)$, there exists a number $N_1(\delta, I, k)$ such that for any $j \ge 1$ and any $n \ge N_1(\delta, I, k)$,

$$\begin{aligned} &(\lambda_{n}^{j})^{2-s_{q}} \|\sigma_{n}^{j} \widetilde{\psi}^{j} (G_{n}^{j})^{-1} e^{it\Delta} w_{n}^{k} \|_{L^{\frac{2(d+2)(q-1)}{d(q-1)+4}}_{t,x}(I^{j})} \\ &\leq \lambda_{n}^{j} \|\sigma_{n}^{j} \widetilde{\psi}^{j} \|_{V_{2+\frac{4}{d}}(I^{j})} \| e^{it\Delta} w_{n}^{k} \|_{W_{q+1}(\mathbb{R})} \lesssim \|\langle \nabla \rangle \widetilde{\psi}^{j} \|_{V_{2+\frac{4}{d}}(I^{j})} \| e^{it\Delta} w_{n}^{k} \|_{W_{q+1}(\mathbb{R})} \leq \delta. \end{aligned}$$

$$(8.154)$$

Case 1.2. Assume s = 0 and $\lambda_{\infty}^{j} = \infty$. Then, $\tilde{\psi}^{j}(t) = e^{it\Delta}\tilde{u}^{j}$ and we see from an estimate similar to (8.154), Strichartz' estimate, Lemma 8.7 and Lemma 8.14 that for any $k \geq K_{1}(\delta, I)$, there exists a number $N_{1}(\delta, I, k)$ such that for any $j \geq 1$ and any $n \geq N_{1}(\delta, I, k)$,

$$\begin{aligned} &(\lambda_{n}^{j})^{2-s_{q}} \|\sigma_{n}^{j} \widetilde{\psi}^{j}(G_{n}^{j})^{-1} e^{it\Delta} w_{n}^{k} \|_{L^{\frac{2(d+2)(q-1)}{d(q-1)+4}}_{t,x}(I^{j})} \\ &\leq \lambda_{n}^{j} \|e^{it\Delta} \sigma_{n}^{j} \widetilde{u}^{j} \|_{V_{2+\frac{4}{d}}(I^{j})} \|e^{it\Delta} w_{n}^{k} \|_{W_{q+1}(\mathbb{R})} \lesssim \|\langle \nabla \rangle \widetilde{u}^{j} \|_{L^{2}} \|e^{it\Delta} w_{n}^{k} \|_{W_{q+1}(\mathbb{R})} \leq \delta. \end{aligned}$$

$$(8.155)$$

Case 2.1. Assume s = 1 and $\lambda_{\infty}^j \in \{0, 1\}$ in (8.153). Take a sequence $\{v_m^j\}_{m \ge 1}$ of smooth functions on $\mathbb{R}^d \times \mathbb{R}$ with the following properties:

$$\lim_{m \to \infty} \left\| \langle \nabla \rangle \left(\widetilde{\psi}^j - v_m^j \right) \right\|_{V_{2+\frac{4}{d}}(I^j) \cap V_{2^*}(I^j)} = 0, \tag{8.156}$$

and for each $m \ge 1$, there exist $R_m^j > 0$, $T_m^j > 0$ and $a_m^j \in \mathbb{R}$ such that

$$E_m^j := \left\{ (x, t) \colon |x| \le R_m^j, \ |t - a_m^j| \le T_m^j \right\} \subset \mathbb{R}^d \times I^j$$
(8.157)

and the support of v_m^j is contained in E_m^j .

We see from (8.152) and (8.156) that for each $j \ge 1$, there exists a number $M_1(I, j)$ such that for any $m \ge M_1(I, j)$,

$$\|\langle \nabla \rangle v_m^j \|_{V_{2+\frac{4}{d}}(I^j) \cap V_{2^*}(I^j)} \le A(I) + 1.$$
(8.158)

Moreover, it follows from (8.156) that for any $\delta > 0$ and any $j \ge 1$, there exists $M_2(\delta, I, j) \ge 1$ such that for any $m \ge M_2(\delta, I, j)$,

$$\|\langle \nabla \rangle \big(\widetilde{\psi}^{j} - v_{m}^{j} \big) \|_{V_{2+\frac{4}{d}}(I^{j}) \cap V_{2^{*}}(I^{j})} \le \delta.$$
(8.159)

Put $m_0 := \max\{M_1(I, j), M_2(\delta, I, j)\}$. We see from the triangle inequality, Hölder's inequality and Sobolev's embedding that the right-hand side of (8.153) with s = 1 is

estimated as follows:

$$\begin{aligned} &(\lambda_{n}^{j})^{2-s_{q}} \|\sigma_{n}^{j} \widetilde{\psi}^{j} (G_{n}^{j})^{-1} |\nabla| e^{it\Delta} w_{n}^{k} \|_{L^{\frac{2(d+2)(q-1)}{d(q-1)+4}}_{t,x}(I^{j})} \\ &\leq (\lambda_{n}^{j})^{1-s_{q}} \|\sigma_{n}^{j} \widetilde{\psi}^{j} - \sigma_{n}^{j} v_{m_{0}}^{j} \|_{W_{q+1}(I^{j})} \||\nabla| e^{it\Delta} w_{n}^{k} \|_{V_{2+\frac{4}{d}}(I^{j})} \\ &+ (\lambda_{n}^{j})^{2-s_{q}} \|\sigma_{n}^{j} v_{m_{0}}^{j} (G_{n}^{j})^{-1} |\nabla| e^{it\Delta} w_{n}^{k} \|_{L^{\frac{2(d+2)(q-1)}{d(q-1)+4}}_{t,x}(I^{j})}. \end{aligned}$$
(8.160)

We consider the first term on the right-hand side of (8.160). Using (8.54), Hölder's inequality, Lemma 8.8 and (8.159) that if n is sufficiently large dependently on k, then

$$\begin{aligned} &(\lambda_{n}^{j})^{1-s_{q}} \| \sigma_{n}^{j} \widetilde{\psi}^{j} - \sigma_{n}^{j} v_{m_{0}}^{j} \|_{W_{q+1}(I^{j})} \| |\nabla| e^{it\Delta} w_{n}^{k} \|_{V_{2+\frac{4}{d}}(I^{j})} \\ &\lesssim (\lambda_{n}^{j})^{1-s_{q}} \| \sigma_{n}^{j} \widetilde{\psi}^{j} - \sigma_{n}^{j} v_{m_{0}}^{j} \|_{V_{2+\frac{4}{d}}(I^{j})}^{1-s_{q}} \| \sigma_{n}^{j} \widetilde{\psi}^{j} - \sigma_{n}^{j} v_{m_{0}}^{j} \|_{W_{2^{*}}(I^{j})}^{s_{q}} \\ &\lesssim \| \langle \nabla \rangle \{ \widetilde{\psi}^{j} - v_{m_{0}}^{j} \} \|_{V_{2+\frac{4}{d}}(I)}^{1-s_{q}} \| \langle \nabla \rangle \{ \widetilde{\psi}^{j} - v_{m_{0}}^{j} \} \|_{V_{2^{*}}(I)}^{1-s_{q}} \leq \delta. \end{aligned}$$

$$(8.161)$$

Next, we consider the second term on the right-hand side of (8.160). Assume first that $\lambda_{\infty}^{j} = 1$, so that $\lambda_{n}^{j} \equiv 1$ and $\sigma_{n}^{j} \equiv 1$. Then, we see from Hölder's inequality, Sobolev's embedding, (8.158), (8.54), Lemma 2.5 of [21] and (8.55) that there exists a number $K_{2}(\delta, I, j)$ with the following property: for any $k \geq K_{2}(\delta, I, j)$, there exists $N_{2}(\delta, I, j, k) \geq 1$ such that for any $n \geq N_{2}(\delta, I, j, k)$,

$$\begin{aligned} &(\lambda_{n}^{j})^{2-s_{q}} \| \sigma_{n}^{j} v_{m_{0}}^{j}(G_{n}^{j})^{-1} | \nabla | e^{it\Delta} w_{n}^{k} \|_{L^{\frac{2(d+2)(q-1)}{d(q-1)+4}}(I^{j})} \\ &= \| v_{m_{0}}^{j} | \nabla | e^{it\Delta} w_{n}^{k} \|_{L^{\frac{2(d+2)(q-1)}{d(q-1)+4}}(E_{m_{0}}^{j})} \\ &\leq \| v_{m_{0}}^{j} \|_{W_{2^{*}}(I^{j})} \| | \nabla | e^{it\Delta} w_{n}^{k} \|_{L^{\frac{(d+2)(q-1)}{q+1}}(E_{m_{0}}^{j})} \\ &\lesssim \| \nabla v_{m_{0}}^{j} \|_{V_{2^{*}}(I^{j})} \| | \nabla | e^{it\Delta} w_{n}^{k} \|_{L^{2}(E_{m_{0}}^{j-1})} \\ &\leq \{A(I)+1\} \| | \nabla | e^{it\Delta} w_{n}^{k} \|_{L^{2}(E_{m_{0}}^{j})} \| | \nabla | e^{it\Delta} w_{n}^{k} \|_{V_{2^{+}\frac{4}{d}}}^{s_{q}} (\mathbb{R}) \\ &\lesssim \{A(I)+1\} (T_{m_{0}}^{j})^{\frac{2(1-s_{q})}{3(d+2)}} (R_{m_{0}}^{j})^{\frac{(3d+2)(1-s_{q})}{6(d+2)}} \| e^{it\Delta} w_{n}^{k} \|_{W_{2^{*}}^{\frac{1-s_{q}}{3}}} \| \nabla w_{n}^{k} \|_{L^{2}}^{\frac{2(1-s_{q})}{3}} \\ &\lesssim \{A(I)+1\} (T_{m_{0}}^{j})^{\frac{2(1-s_{q})}{3(d+2)}} (R_{m_{0}}^{j})^{\frac{(3d+2)(1-s_{q})}{6(d+2)}} \| e^{it\Delta} w_{n}^{k} \|_{W_{2^{*}}^{\frac{1-s_{q}}{3}}} \\ &\lesssim \{A(I)+1\} (T_{m_{0}}^{j})^{\frac{2(1-s_{q})}{3(d+2)}} (R_{m_{0}}^{j})^{\frac{(3d+2)(1-s_{q})}{6(d+2)}} \| e^{it\Delta} w_{n}^{k} \|_{W_{2^{*}}^{\frac{1-s_{q}}{3}}} \\ &\lesssim \delta. \end{aligned}$$

Next, we assume that $\lambda_{\infty}^{j} = 0$. We see from Lemma 8.8 and the support property of $v_{m_{0}}^{j}$ that

$$\|\lambda_n^j \sigma_n^j v_{m_0}^j(t)\|_{L^2}^2 \lesssim \|\langle \nabla \rangle v_{m_0}^j(t)\|_{L^2}^2 \in L^1(\mathbb{R}).$$
(8.163)

We also see from Lemma 8.7 that for any $t \in \mathbb{R}$,

$$\lim_{n \to \infty} \|\lambda_n^j \sigma_n^j v_{m_0}^j(t)\|_{L^2} = 0.$$
(8.164)

Hence, Lebesgue's convergence theorem shows that

$$\lim_{n \to \infty} \|\lambda_n^j \sigma_n^j v_{m_0}^j\|_{L^2_{t,x}(I^j)}^2 = \lim_{n \to \infty} \int_{\mathbb{R}} \|\lambda_n^j \sigma_n^j v_{m_0}^j(t)\|_{L^2}^2 dt = 0.$$
(8.165)

Furthermore, it follows from Hölder's inequality, (8.54), Sobolev's embedding, Lemma 8.8, (8.158), $\lambda_n^j \leq 1$ and (8.165) that for any $k \geq 1$, there exists $N_3(\delta, I, j, k) \geq 1$ such that for any $n \geq N_3(\delta, I, j, k)$,

$$\begin{aligned} &(\lambda_{n}^{j})^{2-s_{q}} \| \sigma_{n}^{j} v_{m_{0}}^{j} (G_{n}^{j})^{-1} | \nabla | e^{it\Delta} w_{n}^{k} \|_{L_{t,x}^{\frac{2(d+2)(q-1)}{d(q-1)+4}} (I^{j})} \\ &\leq (\lambda_{n}^{j})^{1-s_{q}} \| \sigma_{n}^{j} v_{m_{0}}^{j} \|_{V_{2+\frac{4}{d}}(I^{j})}^{1-s_{q}} \| \sigma_{n}^{j} v_{m_{0}}^{j} \|_{W_{2^{*}}(I^{j})}^{s_{q}} \| \| \nabla | e^{it\Delta} w_{n}^{k} \|_{V_{2+\frac{4}{d}}} (\mathbb{R}) \\ &\lesssim \| \lambda_{n}^{j} \sigma_{n}^{j} v_{m_{0}}^{j} \|_{L_{t}^{2}L_{x}^{2}(I^{j})}^{1-s_{q}} \{A(I)+1\}^{s_{q}} \\ &\lesssim \| \lambda_{n}^{j} \sigma_{n}^{j} v_{m_{0}}^{j} \|_{L_{t}^{2}L_{x}^{2}(I^{j})}^{1-s_{q}} \| \nabla \sigma_{n}^{j} v_{m_{0}}^{j} \|_{V_{2^{*}}(I^{j})}^{1-s_{q}} \{A(I)+1\}^{s_{q}} \\ &\lesssim \| \lambda_{n}^{j} \sigma_{n}^{j} v_{m_{0}}^{j} \|_{L_{t}^{2}L_{x}^{2}(I^{j})}^{1-s_{q}} \| \nabla \sigma_{n}^{j} v_{m_{0}}^{j} \|_{V_{2^{*}}(I^{j})}^{1-s_{q}} \{A(I)+1\}^{s_{q}} \\ &\lesssim \| \lambda_{n}^{j} \sigma_{n}^{j} v_{m_{0}}^{j} \|_{L_{t}^{2}L_{x}^{2}(I^{j})}^{1-s_{q}} \{A(I)+1\}^{\frac{1+s_{q}}{2}} \leq \delta. \end{aligned}$$

Case 2.2. Assume s = 1 and $\lambda_{\infty}^{j} = \infty$ in (8.153). We see from Hölder's inequality, (8.54), Sobolev's embedding, Strichartz' estimate and Lemma 8.7 that for any $k \ge 1$, there exists $N_4(\delta, j, k) \ge 1$ such that for any $n \ge N_4(\delta, j, k)$,

$$\begin{aligned} &(\lambda_{n}^{j})^{2-s_{q}} \| \sigma_{n}^{j} \widetilde{\psi}^{j} (G_{n}^{j})^{-1} | \nabla | e^{it\Delta} w_{n}^{k} \|_{L^{\frac{2(d+2)(q-1)}{d(q-1)+4}}(I^{j})} \\ &\leq (\lambda_{n}^{j})^{1-s_{q}} \| \sigma_{n}^{j} \widetilde{\psi}^{j} \|_{V_{2+\frac{4}{d}}(I^{j})}^{1-s_{q}} \| \sigma_{n}^{j} \widetilde{\psi}^{j} \|_{W_{2^{*}}(I^{j})}^{s_{q}} \| \| \nabla | e^{it\Delta} w_{n}^{k} \|_{V_{2+\frac{4}{d}}}(\mathbb{R}) \\ &\lesssim (\lambda_{n}^{j})^{1-s_{q}} \| e^{it\Delta} \sigma_{n}^{j} \widetilde{u}^{j} \|_{V_{2+\frac{4}{d}}(I^{j})}^{1-s_{q}} \| e^{it\Delta} \nabla \sigma_{n}^{j} \widetilde{u}^{j} \|_{V_{2^{*}}(I^{j})}^{s_{q}} \\ &\lesssim \| \lambda_{n}^{j} \sigma_{n}^{j} \widetilde{u}^{j} \|_{L^{2}}^{1-s_{q}} \| \nabla \sigma_{n}^{j} \widetilde{u}^{j} \|_{L^{2}}^{s_{q}} \leq \delta. \end{aligned}$$

$$(8.167)$$

Thus, putting the estimates (8.153), (8.154), (8.160), (8.161), (8.162), (8.166) and (8.167) together, we obtain the desired result (8.151).

Now, for a number $k \ge 1$, we define an approximate solution ψ_n^{k-app} of ψ_n by

$$\psi_n^{k-app}(t) := \sum_{j=1}^k \psi_n^j(t) + e^{it\Delta} w_n^k.$$
(8.168)

Assume (8.127). Then, we have $I_{\max}^j = \mathbb{R}$ for all $j \geq 2$, which together with (8.82) shows that $I_{\max,n}^j = \mathbb{R}$ for all $j \geq 2$. Hence, for any numbers k and n, the maximal existence-interval of ψ_n^{k-app} is $I_{\max,n}^1 := ((\lambda_n^1)^2 T_{\min}^1 + t_n^1, (\lambda_n^1)^2 T_{\max}^1 + t_n^1)$. Furthermore, we see from (8.75) that $0 \in I_{\max,n}^1$ for any sufficiently large number n.

Lemma 8.21. Assume (8.127), and let q denote one of the numbers $1 + \frac{4}{d}$, p and $2^* - 1$. Then, for any $j_0 \ge 2$, any $k > j_0$ and any $\delta > 0$, there exists a number $N(j_0, k, \delta)$ such that for any $n \ge N(j_0, k, \delta)$,

$$\left\|\sum_{j=j_{0}}^{k}\psi_{n}^{j}\right\|_{W_{q+1}(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} \leq \sum_{j=j_{0}}^{k}\|\psi_{n}^{j}\|_{W_{q+1}(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} + \delta,$$
(8.169)

$$\left\|\nabla\sum_{j=j_{0}}^{k}\psi_{n}^{j}\right\|_{V_{2+\frac{4}{d}}(\mathbb{R})}^{\frac{2(d+2)}{d}} \leq \sum_{j=j_{0}}^{k}\left\|\nabla\psi_{n}^{j}\right\|_{V_{2+\frac{4}{d}}(\mathbb{R})}^{\frac{2(d+2)}{d}} + \delta.$$
(8.170)

Proof of Lemma 8.21. Let j_0 and k be numbers with $2 \leq j_0 < k$. Then, we can verify that there exists a constant $C(j_0, k)$ such that for any number n,

$$\begin{split} \| \sum_{j=j_{0}}^{k} \psi_{n}^{j} \|_{W_{q+1}(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} \\ &\leq \sum_{j=j_{0}}^{k} \| \psi_{n}^{j} \|_{W_{q+1}(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} + C(j_{0},k) \sum_{j=2}^{k} \sum_{\substack{2 \leq j' \leq k; \\ j' \neq j}} \int_{\mathbb{R}} \int_{\mathbb{R}^{d}} |\psi_{n}^{j}|^{\frac{(d+2)(q-1)}{2}-1} |\psi_{n}^{j'}| \, dx dt. \end{split}$$

$$(8.171)$$

Furthermore, the orthogonality (8.41) shows that for any $\delta > 0$, we can take a number $N(j_0, k, \delta)$ such that for any distinct numbers $j, j' \in \{2, \ldots, k\}$ and any $n \ge N(j_0, k, \delta)$,

$$\int_{\mathbb{R}} \int_{\mathbb{R}^d} \left| \psi_n^j(x,t) \right|^{\frac{(d+2)(q-1)}{2} - 1} \left| \psi_n^{j'}(x,t) \right| dx dt \le \frac{\delta}{C(j_0,k)k^2}.$$
(8.172)

Putting (8.171) and (8.172) together, we obtain the desired result (8.169). Similarly, we can verify that (8.170) holds.

Lemma 8.22. Assume (8.127). Then, for any interval I satisfying (8.138), we can take a constant B(I) > 0 with the following property: for any number k, there exists a number N(k) such that for any $n \ge N(k)$,

$$\|\psi_{n}^{k\text{-}app}\|_{W_{p+1}(I_{n})\cap W_{2^{*}}(I_{n})} + \|\langle\nabla\rangle\psi_{n}^{k\text{-}app}\|_{V_{2+\frac{4}{d}}(I_{n})} \le B(I),$$
(8.173)

where $I_n := ((\lambda_n^1)^2 \inf I + t_n^1, (\lambda_n^1)^2 \sup I + t_n^1).$

Proof of Lemma 8.22. We consider the first term on the right-hand side of (8.173). Let q denote p or $2^* - 1$, and let J_0 be the number found in Lemma 8.14. Then, it follows from Lemma 8.21 with $\delta = 1$, Lemma 8.13, (8.54), Lemma 8.19, (8.136) and Lemma 8.14

that for any number k, there exists a number N(k) such that for any $n \ge N(k)$,

$$\begin{split} \|\psi_{n}^{k-app}\|_{W_{q+1}(I_{n})}^{\frac{(d+2)(q-1)}{2}} \\ \lesssim \|\psi_{n}^{1}\|_{W_{q+1}(I_{n})}^{\frac{(d+2)(q-1)}{2}} + \sum_{j=2}^{k} \|\psi_{n}^{j}\|_{W_{q+1}(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} + 1 + \|e^{it\Delta}w_{n}^{k}\|_{W_{q+1}(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} \\ \lesssim \|\langle\nabla\rangle\widetilde{\psi}^{1}\|_{St(I)}^{\frac{(d+2)(q-1)}{2}} + \sum_{j=2}^{J_{0}} \|\langle\nabla\rangle\widetilde{\psi}^{j}\|_{St(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} + \sum_{j=J_{0}}^{\infty} \|\langle\nabla\rangle\widetilde{\psi}^{j}\|_{St(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} + 1 \\ \lesssim A(I)^{\frac{(d+2)(q-1)}{2}} + J_{0} + 1. \end{split}$$

$$(8.174)$$

Thus, we have obtained the desired estimate for the first term. We can deal with the second term in a similar way. $\hfill \Box$

Lemma 8.23. Assume (8.127), and let q denote p or 2^*-1 . Then, for any $\delta > 0$ and any interval I satisfying (8.138), we can take a number $K(\delta, I)$ with the following property: for any $k \ge K(\delta, I)$, there exists a number $N(\delta, I, k)$ such that for any $n \ge N(\delta, I, k)$ and $s \in \{0, 1\}$,

$$\left\| |\psi_n^{k-app}|^{q-1} |\nabla|^s e^{it\Delta} w_n^k \right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_n)} \le \delta,$$
(8.175)

where $I_n := ((\lambda_n^1)^2 \inf I + t_n^1, \ (\lambda_n^1)^2 \sup I + t_n^1).$

Proof of Lemma 8.23. Let us begin with some preparation. We see from Sobolev's embedding, Lemma 8.13 and Lemma 8.14 that for any $\delta > 0$, there exists a number $K_1(\delta)$ such that

$$\sum_{j=K_1(\delta)}^{\infty} \left\|\psi_n^j\right\|_{W_{q+1}(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} \lesssim \sum_{j=K_1(\delta)}^{\infty} \left\|\langle\nabla\rangle\widetilde{\psi}^j\right\|_{St(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} \le \delta.$$
(8.176)

This together with Lemma 8.21 shows that for any $\delta > 0$ and any $k > K_1(\delta)$, there exists a number $N_1(\delta, k)$ such that for any $n \ge N_1(\delta, k)$,

$$\left\|\sum_{j=K_1(\delta)+1}^k \psi_n^j\right\|_{W_{q+1}(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} \lesssim \delta.$$
(8.177)

Furthermore, we see from Hölder's inequality, (8.54) and (8.55) that for any $\delta > 0$, there exists a number $K_2(\delta)$ with the following property: for any $k \ge K_2(\delta)$, there exists a number $N_2(\delta, k)$ such that for any $n \ge N_2(\delta, k)$ and $s \in \{0, 1\}$,

$$\begin{aligned} \left\| e^{it\Delta} w_{n}^{k} |\nabla|^{s} e^{it\Delta} w_{n}^{k} \right\|_{L_{t,x}^{\frac{2(d+2)(q-1)}{2}}(\mathbb{R})} &\leq \| e^{it\Delta} w_{n}^{k} \|_{W_{q+1}(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} \| |\nabla|^{s} e^{it\Delta} w_{n}^{k} \|_{V_{2+\frac{4}{d}}(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} \\ &\lesssim \| e^{it\Delta} w_{n}^{k} \|_{W_{q+1}(\mathbb{R})}^{\frac{(d+2)(q-1)}{2}} \leq \delta. \end{aligned}$$

$$(8.178)$$

Now, we shall prove (8.175). First, we consider the case where $q \leq 2$. Then, we see from Hölder's inequality, (8.54), (8.177) and (8.178) that for any $\delta > 0$, any $k > \max\{K_1(\delta), K_2(\delta)\}$, any $n \geq \max\{N_1(\delta, k), N_2(\delta, k)\}$ and $s \in \{0, 1\}$,

where $C(\delta)$ is some constant depending only on d, q and $K_1(\delta)$. Moreover, Lemma 8.20 shows that we can take a number $K_3(\delta, I)$ with the following property: for any $k \geq K_3(\delta, I)$, there exists a number $N_3(\delta, I, k)$ such that for any $1 \leq j \leq K_1(\delta)$, any $n \geq N_3(\delta, I, k)$ and $s \in \{0, 1\}$,

$$\left\|\psi_{n}^{j}|\nabla|^{s}e^{it\Delta}w_{n}^{k}\right\|_{L_{t,x}^{\frac{2(d+2)(q-1)}{2}}(I_{n})}^{\frac{(d+2)(q-1)}{2}} \leq \frac{\delta}{C(\delta)K_{1}(\delta)}.$$
(8.180)

Putting (8.179) and (8.180) together, we obtain the desired result (8.175) in the case $q \leq 2$.

Next, we consider the case where q > 2. In this case, we see from Hölder's inequality and Lemma 8.22 that there exists B(I) > 0 with the following property: for any number k, there exists a number $N_4(k)$ such that for any $n \ge N_4(k)$ and $s \in \{0, 1\}$,

$$\begin{aligned} \left\| |\psi_{n}^{k-app}|^{q-1} |\nabla|^{s} e^{it\Delta} w_{n}^{k} \right\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(I_{n})} \\ &\leq \|\psi_{n}^{k-app} |\nabla|^{s} e^{it\Delta} w_{n}^{k} \|_{L_{t,x}^{\frac{2(d+2)(q-1)}{d(q-1)+4}}(I_{n})} \|\psi_{n}^{k-app}\|_{W_{q+1}}^{q-2} \\ &\leq \|\psi_{n}^{k-app} |\nabla|^{s} e^{it\Delta} w_{n}^{k} \|_{L_{t,x}^{\frac{2(d+2)(q-1)}{d(q-1)+4}}(I_{n})} B(I)^{q-2}. \end{aligned}$$

$$(8.181)$$

Then, we can obtain the desired estimate (8.175) in a way similar to the case $q \leq 2$.

The following lemma enables us to control the error term of the approximate solution:

Lemma 8.24. Assume (8.127). Then, for any $\delta \in (0,1)$ and any interval I satisfying (8.138), there exist numbers k_0 (depending on δ and I) and $N(\delta, I)$ such that for any $n \geq N(\delta, I)$,

$$\|\langle \nabla \rangle e[\psi_n^{k_0 - app}]\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_n)} \le \delta,$$
(8.182)

where $I_n := ((\lambda_n^1)^2 \inf I + t_n^1, (\lambda_n^1)^2 \sup I + t_n^1).$

Proof of Lemma 8.24. Let q denote p or $2^* - 1$. Then, we see from (8.55), Lemma 8.22 and Lemma 8.23 that for any $\delta \in (0, 1)$ and any interval I satisfying (8.138), there exist a number $k_0 \ge 1$ (depending on δ and I), a constant B(I) > 0 (independent of δ) and a number $N_1(\delta, I) \ge 1$ such that if $n \ge N_1(\delta, I)$, then

$$\|e^{it\Delta}w_n^{k_0}\|_{W_{q+1}(\mathbb{R})} \le \frac{\delta}{(1+B(I))^{2^*}},\tag{8.183}$$

$$\|\psi_{n}^{k_{0}-app}\|_{W_{q+1}(I_{n})} + \|\langle\nabla\rangle\psi_{n}^{k_{0}-app}\|_{V_{2+\frac{4}{d}}(I_{n})} \le B(I),$$
(8.184)

$$\||\psi_{n}^{k_{0}-app}|^{q-1}e^{it\Delta}w_{n}^{k_{0}}\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_{n})} + \||\psi_{n}^{k_{0}-app}|^{q-1}|\nabla|e^{it\Delta}w_{n}^{k_{0}}\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_{n})} \leq \delta.$$
(8.185)

We rewrite the "error term" $e[\psi_n^{k_0-app}]$ as follows:

$$e[\psi_n^{k_0 - app}] = \sum_{j=1}^{k_0} \left(i \frac{\partial \psi_n^j}{\partial t} + \Delta \psi_n^j \right) + F\left[\sum_{j=1}^{k_0} \psi_n^j \right]$$

$$+ F\left[\psi_n^{k_0 - app} \right] - F\left[\psi_n^{k_0 - app} - e^{it\Delta} w_n^{k_0} \right].$$
(8.186)

Then, we have

$$\begin{aligned} \|\langle \nabla \rangle e[\psi_{n}^{k_{0}-app}]\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_{n})} \\ &\leq \|\langle \nabla \rangle \left\{ \sum_{j=1}^{k_{0}} \left(i \frac{\partial \psi_{n}^{j}}{\partial t} + \Delta \psi_{n}^{j} \right) + F\left[\sum_{j=1}^{k_{0}} \psi_{n}^{j} \right] \right\} \|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_{n})} \\ &+ \|\langle \nabla \rangle \left\{ F\left[\psi_{n}^{k_{0}-app} \right] - F\left[\psi_{n}^{k_{0}-app} - e^{it\Delta} w_{n}^{k_{0}} \right] \right\} \|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_{n})}. \end{aligned}$$
(8.187)

We consider the first term on the right-hand side of (8.187). It follows from (8.81), Lemma 8.11 and Lemma 8.12 that there exists a number $N_2(\delta, I)$ such that for any $n \ge N_2(\delta, I),$

$$\|\langle \nabla \rangle \left\{ \sum_{j=1}^{k_0} \left(i \frac{\partial \psi_n^j}{\partial t} + \Delta \psi_n^j \right) + F \left[\sum_{j=1}^{k_0} \psi_n^j \right] \right\} \|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_n)}$$

$$\leq \|\langle \nabla \rangle \left\{ \sum_{j=1}^{k_0} F \left[\psi_n^j \right] - F \left[\sum_{j=1}^{k_0} \psi_n^j \right] \right\} \|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_n)} + \delta.$$

$$(8.188)$$

Then, we see from elementary computations that for q = p or $q = 2^* - 1$,

$$\begin{aligned} \|\sum_{j=1}^{k_{0}} |\psi_{n}^{j}|^{q-1} \psi_{n}^{j} - |\sum_{j=1}^{k_{0}} \psi_{n}^{j}|^{q-1} \sum_{j=1}^{k_{0}} \psi_{n}^{j} \|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_{n})} \\ \lesssim \sum_{j=1}^{k_{0}} \sum_{\substack{1 \le k \le k_{0} \\ k \ne j}} \||\psi_{n}^{j}|^{q-1} \psi_{n}^{k}\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_{n})} \end{aligned}$$

$$(8.189)$$

and

$$\|\nabla \left\{ \sum_{j=1}^{k_0} |\psi_n^j|^{q-1} \psi_n^j - \left| \sum_{j=1}^{k_0} \psi_n^j \right|^{q-1} \sum_{j=1}^{k_0} \psi_n^j \right\} \|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_n)}$$

$$\lesssim \sum_{j=1}^{k_0} \sum_{\substack{1 \le k \le k_0 \\ k \ne j}} \||\psi_n^j|^{q-1} \nabla \psi_n^k\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_n)}.$$

$$(8.190)$$

Here, when q > 2, we must add the following term to the right-hand side of (8.190):

$$\sum_{i=1}^{k_0} \sum_{\substack{j=1\\k\neq j}}^{k_0} \sum_{\substack{1\le k\le k_0\\k\neq j}} \||\psi_n^i|^{q-2} |\nabla \psi_n^j| |\psi_n^k|\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_n)}.$$
(8.191)

We see from the orthogonality (8.41) that there exists a number $N_3(\delta, I)$ such that for any distinct numbers $j, k \in \{1, \ldots, k_0\}$ and any $n \ge N_3(\delta, I)$,

$$\||\psi_{n}^{j}|^{q-1}\psi_{n}^{k}\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_{n})} + \||\psi_{n}^{j}|^{q-1}\nabla\psi_{n}^{k}\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_{n})} \le \frac{\delta}{k_{0}^{2}}.$$
(8.192)

When q > 2, we also see that for any number $i \in \{1, \ldots, k_0\}$, any distinct numbers $j, k \in \{1, \ldots, k_0\}$ and any $n \ge N_3(\delta, I)$,

$$\||\psi_n^i|^{q-2}\nabla\psi_n^j\psi_n^k\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_n)} \le \frac{\delta}{k_0^3}.$$
(8.193)

Putting the estimates (8.188) through (8.193) together, we find that for any $n \ge \max\{N_2(\delta, I), N_3(\delta, I)\},\$

$$\|\langle \nabla \rangle \left\{ \sum_{j=1}^{k_0} \left(i \frac{\partial \psi_n^j}{\partial t} + \Delta \psi_n^j \right) + F\left[\sum_{j=1}^{k_0} \psi_n^j \right] \right\} \|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_n)} \lesssim \delta.$$
(8.194)

We move on to the second term on the right-hand side of (8.187). We can verify that for q = p or $q = 2^* - 1$, and any number $n \ge 1$,

$$\begin{split} \|\langle \nabla \rangle \Big\{ \|\psi_{n}^{k_{0}-app}\|^{q-1}\psi_{n}^{k_{0}-app} - |\psi_{n}^{k_{0}-app} - e^{it\Delta}w_{n}^{k_{0}}|^{q-1}(\psi_{n}^{k_{0}-app} - e^{it\Delta}w_{n}^{k_{0}}) \Big\} \|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(I_{n})} \\ &\lesssim \||\psi_{n}^{k_{0}-app}|^{q-1}e^{it\Delta}w_{n}^{k_{0}}\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(I_{n})} + \||e^{it\Delta}w_{n}^{k_{0}}|^{q-1}\psi_{n}^{k_{0}-app}\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(I_{n})} \\ &+ \||\psi_{n}^{k_{0}-app}|^{q-1}\nabla e^{it\Delta}w_{n}^{k_{0}}\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(I_{n})} + \||e^{it\Delta}w_{n}^{k_{0}}|^{q-1}\nabla \psi_{n}^{k_{0}-app}\|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(I_{n})} \\ &+ \|\langle \nabla \rangle \big\{ |e^{it\Delta}w_{n}^{k_{0}}|^{q-1}e^{it\Delta}w_{n}^{k_{0}}\big\} \|_{L_{t,x}^{\frac{2(d+2)}{d+4}}(I_{n})}. \end{split}$$

$$(8.195)$$

When q > 2, we must add the following terms to the right-hand side of (8.195):

$$\||\psi_{n}^{k_{0}-app}|^{q-2}e^{it\Delta}w_{n}^{k_{0}}\nabla\psi_{n}^{k_{0}-app}\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_{n})},$$
(8.196)

$$\||e^{it\Delta}w_n^{k_0}|^{q-2}\psi_n^{k_0-app}\nabla e^{it\Delta}w_n^{k_0}\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_n)}.$$
(8.197)

Furthermore, using (8.185), Hölder's inequality, (8.183), (8.184) and (8.54), we can verify that the right-hand side of (8.195) vanishes as n tends to the infinity (cf. the proof of (3.10) in [21]). Thus, we conclude that

$$\lim_{n \to \infty} \left\| \langle \nabla \rangle \left\{ F \left[\psi_n^{k_0 - app} \right] - F \left[\psi_n^{k_0 - app} - e^{it\Delta} w_n^{k_0} \right] \right\} \right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_n)} = 0.$$
(8.198)

Putting (8.187), (8.194) and (8.198) together, we find that the desired estimate (8.182) holds. $\hfill \Box$

Proposition 8.25. Assume (8.127). Then, $\lambda_{\infty}^1 \in \{0,1\}$, and for any $T \in I_{\max}^1 = (T_{\min}^1, T_{\max}^1)$,

$$\|\sigma_{\infty}^{1}\psi^{1}\|_{\mathcal{W}^{1}([T,T_{\max}^{1}))} = \infty.$$
(8.199)

Furthermore, the following holds:

$$\begin{cases} \mathcal{S}_{\omega}(\sigma_{\infty}^{1}\widetilde{\psi}^{1}) \geq m_{\omega} & \text{if } \lambda_{\infty}^{1} = 1, \\ \mathcal{H}^{\ddagger}(\sigma_{\infty}^{1}\widetilde{\psi}^{1}) \geq \frac{1}{d}\sigma^{\frac{d}{2}} > m_{\omega} & \text{if } \lambda_{\infty}^{1} = 0. \end{cases}$$

$$(8.200)$$

Proof of Proposition 8.25. First, we assume $\lambda_{\infty}^1 \in \{0, 1\}$ and prove (8.199). Suppose for contradiction that (8.199) failed for some $T \in I_{\max}^1$. Then, we see from the well-posedness theory that $T_{\max}^1 = \infty$ and for any $\tau \in (T_{\min}^1, \infty)$,

$$\|\sigma_{\infty}^{1}\psi^{1}\|_{\mathcal{W}^{1}([\tau,\infty))} < \infty.$$

$$(8.201)$$

In order to derive a contradiction, we consider the approximate solution ψ_n^{k-app} defined by (8.168). Note that for any numbers k and n, the maximal existence interval of ψ_n^{k-app} is identical to $I_{\max,n}^1 := ((\lambda_n^1)^2 T_{\min}^1 + t_n^1, \infty).$

We shall show that there exists a constant B > 0 with the following property: for any number k, there exists a number N(k) such that for any $n \ge N(k)$,

$$\|\psi_n^{k-app}\|_{V_{2+\frac{4}{d}}([0,\infty))\cap W_{2^*}([0,\infty))} \le B.$$
(8.202)

If $-\frac{t_{\infty}^1}{(\lambda_{\infty}^1)^2} = -\infty$, then we see from (8.201) and (8.78) that Lemma 8.22 is available on the whole interval \mathbb{R} . Thus, we can take a constant $B_1 > 0$ with the following property: for any number k, there exists a number $N_1(k)$ such that for any $n \ge N_1(k)$,

$$\|\psi_n^{k-app}\|_{V_{2+\frac{4}{d}}(\mathbb{R})\cap W_{2^*}(\mathbb{R})} \le B_1.$$
(8.203)

On the other hand, if $-\frac{t_{\infty}^1}{(\lambda_{\infty}^1)^2} \neq -\infty$, then it follows from the construction of $\tilde{\psi}^1$ (see (8.72)) that $-\frac{t_{\infty}^1}{(\lambda_{\infty}^1)^2} \in (T_{\min}^1, \infty) \cup \{\infty\}$. This implies that there exist $\tau_0 \in (T_{\min}^1, \infty)$ and a number N_0 such that $\tau_0 < -\frac{t_n^1}{(\lambda_n^1)^2}$ for any $n \geq N_0$, so that $[0, \infty) \subset ((\lambda_n^1)^2 \tau_0 + t_n^1, \infty) \subset I_{\max,n}^1$. Furthermore, we see from Lemma 8.22 and (8.201) with $\tau = \tau_0$ that there exists a constant $B_2 > 0$ with the following property: for any number k, there exists a number $N_2(k)$ such that for any $n \geq \max\{N_0, N_2(k)\}$,

$$\|\psi_n^{k-app}\|_{V_{2+\frac{4}{d}}([0,\infty))\cap W_{2^*}([0,\infty))} \le B_2.$$
(8.204)

Thus, (8.203) and (8.204) give the desired result (8.202).

Now, we shall finish the proof of (8.199) for $\lambda_{\infty}^1 \in \{0, 1\}$. Note that it follows from (8.43), (8.46) and Lemma 8.8 that

$$\|\psi_{n}(0) - \psi_{n}^{k-app}(0)\|_{H^{1}} = \left\|\sum_{j=1}^{k} g_{n}^{j} \sigma_{n}^{j} e^{-i \frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \Delta} \widetilde{u}^{j} - \sum_{j=1}^{k} g_{n}^{j} \sigma_{n}^{j} \widetilde{\psi}^{j} \left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\right)\right\|_{H^{1}}$$

$$\lesssim \sum_{j=1}^{k} \left\|e^{-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \Delta} \widetilde{u}^{j} - \widetilde{\psi}^{j} \left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\right)\right\|_{H^{1}}.$$
(8.205)

This estimate together with (8.76) shows that for any number k, there exists a number $N_3(k)$ such that

$$\sup_{n \ge N_3(k)} \|\psi_n(0) - \psi_n^{k-app}(0)\|_{H^1} \le 1.$$
(8.206)

We also see from (8.38) that

$$\sup_{n \ge 1} \|\psi_n\|_{L^{\infty}_t H^1_x([0,\infty))} \lesssim 1.$$
(8.207)

Let $\delta_0 > 0$ be a constant determined by the long-time perturbation theory (Lemma E.1) together with (8.202), (8.206) and (8.207). Then, we see from Sobolev's embedding,

Strichartz' estimate, (8.205) and (8.76) that for any number k, there exists a number $N_4(k)$ such that for any $n \ge N_4(k)$,

$$\begin{aligned} \|e^{it\Delta} \{\psi_n(0) - \psi_n^{k\text{-}app}(0)\}\|_{V_{2+\frac{4}{d}}([0,\infty)\cap W_{2^*}([0,\infty))} \\ \lesssim \|\langle \nabla \rangle e^{it\Delta} \{\psi_n(0) - \psi_n^{k\text{-}app}(0)\}\|_{St([0,\infty))} \lesssim \|\psi_n(0) - \psi_n^{k\text{-}app}(0)\|_{H^1} \ll \delta_0. \end{aligned}$$

$$(8.208)$$

Moreover, it follows from Lemma 8.24 that we can take numbers k_0 and N_5 such that for any $n \ge N_5$,

$$\|\langle \nabla \rangle e[\psi_n^{k_0 - app}]\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}([0,\infty))} \le \delta_0.$$
(8.209)

Thus, applying the long-time perturbation theory (Lemma E.1) to ψ_n and $\psi_n^{k_0-app}$ with $n \ge \max\{N(k_0), N_3(k_0), N_4(k_0), N_5\}$, we find that

$$\|\langle \nabla \rangle \psi_n\|_{St([0,\infty))} < \infty. \tag{8.210}$$

However, this contradicts (8.34). Thus, we have shown that (8.199) holds when $\lambda_{\infty}^1 \in \{0, 1\}$.

Next, we assume that $\lambda_{\infty}^1 = \infty$. Then, (8.202) holds without the hypothesis (8.201). Furthermore, the same argument as (8.205) through (8.210) shows that the case $\lambda_{\infty}^1 = \infty$ never happens.

It remains to prove (8.200). Suppose to the contrary that the claim (8.200) was false. Then, it follows from (8.74), Theorem 1.2, Theorem 1.3, Lemma 8.18 and (8.199) that $\|\sigma_{\infty}^{j}\tilde{\psi}^{j}\|_{St([0,\infty))} < \infty$. However, this contradicts (8.199). Thus, we have proved that the claim (8.200) is true.

8.2.5 Critical element and completion of the proof

We shall show the existence of "critical element" (see Lemma 8.29 below). Furthermore, we will derive a contradiction by using it under the hypothesis (8.30), which completes the proof of Proposition 8.6.

Let us begin by defining the functionals \mathcal{S}^j_{ω} and \mathcal{I}^j_{ω} . For each $j \geq 1$, we define

$$\mathcal{S}^{j}_{\omega} := \begin{cases} \mathcal{S}_{\omega} & \text{if } \lambda^{j}_{\infty} \in \{1, \infty\}, \\ \mathcal{H}^{\ddagger} & \text{if } \lambda^{j}_{\infty} = 0, \end{cases} \qquad \qquad \mathcal{I}^{j}_{\omega} := \begin{cases} \mathcal{I}_{\omega} & \text{if } \lambda^{j}_{\infty} \in \{1, \infty\}, \\ \mathcal{I}^{\ddagger} & \text{if } \lambda^{j}_{\infty} = 0. \end{cases}$$
(8.211)

Lemma 8.26. Assume (8.127). Then, for any $\delta > 0$ and any number $k \ge 2$, there exists a number $N(\delta, k)$ such that for any $t \in \mathbb{R}$ and any $n \ge N(\delta, k)$,

$$\sum_{j=2}^{k} \mathcal{I}_{\omega}^{j} \left(\sigma_{\infty}^{j} \widetilde{\psi}^{j}(t) \right) + \mathcal{I}_{\omega}(w_{n}^{k}) \leq \delta.$$
(8.212)

Proof of Lemma 8.26. We see from (8.31), (8.33) and (8.52) that for any $\delta > 0$ and any number k, there exists a number $N_1(\delta, k)$ such that for any $n \ge N_1(\delta, k)$,

$$\mathcal{S}_{\omega}\left(g_{n}^{1}\sigma_{n}^{1}e^{-i\frac{t_{n}^{1}}{(\lambda_{n}^{1})^{2}}\Delta}\widetilde{u}^{1}\right) + \sum_{j=2}^{k}\mathcal{S}_{\omega}\left(g_{n}^{j}\sigma_{n}^{j}e^{-i\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\Delta}\widetilde{u}^{j}\right) + \mathcal{S}_{\omega}(w_{n}^{k}) \leq m_{\omega} + \delta.$$
(8.213)

If $\lambda_{\infty}^1 = \lim_{n \to \infty} \lambda_n^1 = 0$, then Lemma 8.7 shows that for each $t \in I_{\max}^1$,

$$\lim_{n \to \infty} \|g_n^1 \sigma_n^1 \widetilde{\psi}^1(t)\|_{L^2} = 0.$$
(8.214)

We see from Lemma 8.18, (8.120) in Lemma 8.16, (8.79), (8.213), (8.214) and (8.200) in Proposition 8.25 that for any $\delta > 0$ and any number k, there exists $N_2(\delta, k)$ such that for any $t \in \mathbb{R}$ and any $n \ge N_2(\delta, k)$,

$$\sum_{j=2}^{k} \mathcal{I}_{\omega}^{j} \left(\sigma_{\infty}^{j} \widetilde{\psi}^{j}(t) \right) + \mathcal{I}_{\omega}(w_{n}^{k}) \leq \sum_{j=2}^{k} \mathcal{S}_{\omega}^{j} \left(\sigma_{\infty}^{j} \widetilde{\psi}^{j} \right) + \mathcal{S}_{\omega}(w_{n}^{k})$$

$$\leq \sum_{j=2}^{k} \mathcal{S}_{\omega}^{j} \left(g_{n}^{j} \sigma_{n}^{j} e^{-i \frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \Delta} \widetilde{u}^{j} \right) + \delta + \mathcal{S}_{\omega}(w_{n}^{k})$$

$$\leq m_{\omega} - \mathcal{S}_{\omega}^{1} (\sigma_{\infty}^{1} \widetilde{\psi}^{1}) + 2\delta \leq 2\delta.$$
(8.215)

Thus, we have completed the proof.

Put $\tau_n^1 := -\frac{t_n^j}{(\lambda_n^1)^2}$. Then, passing to a subsequence, we may assume that $\tau_\infty^1 := \lim_{n \to \infty} \tau_n^1 \in \mathbb{R} \cup \{\pm \infty\}.$

Lemma 8.27. Assume (8.127). Then, we have that $\lambda_{\infty}^1 = 1$. Furthermore, there exists a time $T \in I_{\max}^1$ such that

$$\inf_{\in [T,T_{\max}^1)} \widetilde{d}_{\omega} \left(\widetilde{\psi}^1(t) \right) \ge \frac{R_*}{2}$$
(8.217)

(8.216)

and

$$\inf_{t \in [T, T_{\max}^1)} \mathcal{K}\big(\widetilde{\psi}^1(t)\big) \ge \frac{\kappa_1(R_*)}{2},\tag{8.218}$$

where $\kappa_1(R_*)$ is the constant appearing in (8.37).

t

Proof of Lemma 8.27. Proposition 8.25 together with (8.77) shows that $\lambda_{\infty}^1 \neq \infty$ and $\tau_{\infty}^1 \neq \infty$. If $\tau_{\infty}^1 = -\infty$, then it follows from (8.78) that $T_{\min}^1 = -\infty$ and $\|\sigma_{\infty}^1 \widetilde{\psi}^1\|_{\mathcal{W}^1((-\infty,T])} < \infty$ for any $T < T_{\max}^1$. If $\tau_{\infty}^1 \in \mathbb{R}$, then we see from the construction of $\widetilde{\psi}^1$ (see (8.72)) that $\tau_{\infty}^1 \in I_{\max}^1$. Put

$$\tau_{\min}^{1} := \begin{cases} \frac{T_{\min}^{1} + \tau_{\infty}^{1}}{2} & \text{if } T_{\min}^{1} > -\infty, \\ \tau_{\infty}^{1} - 1 & \text{if } \tau_{\infty}^{1} \in \mathbb{R} \text{ and } T_{\min}^{1} = -\infty, \\ -\infty & \text{if } \tau_{\infty}^{1} = -\infty. \end{cases}$$
(8.219)

Then, $T_{\min}^1 \leq \tau_{\min}^1 \leq \tau_{\infty}^1$, and we can take a number N_1 such that for any $n \geq N_1$,

$$\tau_{\min}^1 < -\frac{t_n^1}{(\lambda_n^1)^2} < T_{\max}^1.$$
(8.220)

Note that $0 \in I_{\max,n}^1$ for all $n \ge N_1$. Furthermore, for each $T \in (\tau_{\min}^1, T_{\max}^1)$,

$$\|\sigma_{\infty}^{1}\widetilde{\psi}^{1}\|_{\mathcal{W}^{1}((\tau_{\min}^{1},T])} < \infty, \qquad (8.221)$$

which together with Lemma 8.19 shows that for any $T \in (\tau_{\min}^1, T_{\max}^1)$, there exists a constant A(T) > 0 such that

$$\|\langle \nabla \rangle \widetilde{\psi}^1\|_{St((\tau_{\min}^1, T])} \le A(T).$$
(8.222)

Sobolev's embedding also gives us that

$$\|\sigma_{\infty}^{1}\widetilde{\psi}^{1}\|_{\mathcal{W}^{1}((\tau_{\min}^{1},T])} \lesssim \|\langle \nabla \rangle \widetilde{\psi}^{1}\|_{St((\tau_{\min}^{1},T])} \le A(T).$$
(8.223)

Now, we consider the approximate solution ψ_n^{k-app} defined by (8.168). Lemma 8.22 together with (8.223) shows that for any $T \in (\tau_{\min}^1, T_{\max}^1)$, there exists B(T) > 0 with the following property: for any number k, there exists a number $N_2(k)$ such that for any $n \ge N_2(k)$,

$$\|\psi_{n}^{k-app}\|_{V_{2+\frac{4}{d}}(I_{n}(T))\cap W_{2^{*}}(I_{n}(T))} + \|\langle\nabla\rangle\psi_{n}^{k-app}\|_{L_{t}^{\infty}L_{x}^{2}(I_{n}(T))} \le B(T),$$
(8.224)

where

$$I_n(T) := \left((\lambda_n^1)^2 \tau_{\min}^1 + t_n^1, \ (\lambda_n^1)^2 T + t_n^1 \right].$$
(8.225)

Lemma 8.24 together with (8.223) shows that for any $\delta \in (0, 1)$ and any $T \in (\tau_{\min}^1, T_{\max}^1)$, there exists numbers k_0 (depending on δ and T) and $N_3(\delta, T)$ such that for any $n \geq N_3(\delta, T)$,

$$\|\langle \nabla \rangle e[\psi_n^{k_0 - app}]\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I_n(T))} \le \delta.$$
(8.226)

We also see from (8.43), Lemma 8.8 and (8.76) that for any number k and any $\gamma > 0$, there exists a number $N_4(k, \gamma)$ such that for any $n \ge N_4(k, \gamma)$,

$$\begin{aligned} \left\|\psi_{n}(0) - \psi_{n}^{k-app}(0)\right\|_{H^{1}} &= \left\|\sum_{j=1}^{k} g_{n}^{j} \sigma_{n}^{j} e^{-i \frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \Delta} \widetilde{u}^{j} - \sum_{j=1}^{k} g_{n}^{j} \sigma_{n}^{j} \widetilde{\psi}^{j} \left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\right)\right\|_{H^{1}} \\ &\lesssim \sum_{j=1}^{k} \left\|e^{-i \frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}} \Delta} \widetilde{u}^{j} - \widetilde{\psi}^{j} \left(-\frac{t_{n}^{j}}{(\lambda_{n}^{j})^{2}}\right)\right\|_{H^{1}} \leq \gamma. \end{aligned}$$
(8.227)

Furthermore, it follows from Strichartz' estimate and (8.227) that for any number k and any $\delta > 0$, there exists a number $N_5(k, \delta)$ such that for any $n \ge N_5(k, \delta)$,

$$\left\| \langle \nabla \rangle e^{it\Delta} \left\{ \psi_n(0) - \psi_n^{k\text{-}app}(0) \right\} \right\|_{V_{p+1}(\mathbb{R})} \lesssim \left\| \psi_n(0) - \psi_n^{k\text{-}app}(0) \right\|_{H^1} \le \delta.$$
(8.228)

The long-time perturbation theory (Lemma E.1) together with (8.224), (8.226), (8.227) and (8.228) shows that for any $T \in (\tau_{\min}^1, T_{\max}^1)$, there exist constants $\delta_0(T, \gamma) > 0$ and $C(T, \gamma) > 0$ with the following properties: for any $\delta \in (0, \delta_0(T, \gamma))$, we can take numbers k_0 (depending on δ and T) and $N_6(\delta, T)$ such that for any $n \ge N_6(\delta, T)$,

$$\left\| \langle \nabla \rangle \psi_n \right\|_{St(I_n(T))} \le C(T, \gamma), \tag{8.229}$$

$$\sup_{t \in I_n(T)} \left\| \langle \nabla \rangle \left\{ \psi_n(t) - \psi_n^{k_0 - app}(t) \right\} \right\|_{L^2} \le C(T, \gamma) \gamma.$$
(8.230)

Here, $C(T, \gamma)$ is non-decreasing with respect to γ . Furthermore, it follows from (8.230), Lemma 8.7 and Lemma 8.26 that for any $T \in (\tau_{\min}^1, T_{\max}^1)$, any $\delta \in (0, \delta_0(T, \gamma))$ and the number k_0 determined by T and δ , we can take a number $N_7(\delta, T, \omega)$ such that for any $n \geq N_7(\delta, T, \omega)$ and $s \in \{0, 1\}$,

$$\begin{split} \||\nabla|^{s} \left\{\psi_{n}\left((\lambda_{n}^{1})^{2}T+t_{n}^{1}\right)-\psi_{n}^{1}\left((\lambda_{n}^{1})^{2}T+t_{n}^{1}\right)\right\}\|_{L^{2}}^{2} \\ \lesssim \||\nabla|^{s} \left\{\psi_{n}\left((\lambda_{n}^{1})^{2}T+t_{n}^{1}\right)-\psi_{n}^{k_{0}-app}\left((\lambda_{n}^{1})^{2}T+t_{n}^{1}\right)\right\}\|_{L^{2}}^{2} \\ &+ \left\||\nabla|^{s} \sum_{j=2}^{k_{0}}\psi_{n}^{j}\left((\lambda_{n}^{1})^{2}T+t_{n}^{1}\right)\right\|_{L^{2}}^{2} + \left\||\nabla|^{s}w_{n}^{k_{0}}\right\|_{L^{2}}^{2} \\ \lesssim C(T,\gamma)^{2}\gamma^{2}+k_{0} \sum_{j=2}^{k_{0}}\left\||\nabla|^{s}\psi_{n}^{j}\left((\lambda_{n}^{1})^{2}T+t_{n}^{1}\right)\right\|_{L^{2}}^{2} + \left\||\nabla|^{s}w_{n}^{k_{0}}\right\|_{L^{2}}^{2} + \delta \\ &= C(T,\gamma)^{2}\gamma^{2}+k_{0} \sum_{j=2}^{k_{0}}\left\||\nabla|^{s}g_{n}^{j}\sigma_{n}^{j}\widetilde{\psi}^{j}(T)\right\|_{L^{2}}^{2} + \left\||\nabla|^{s}w_{n}^{k_{0}}\right\|_{L^{2}}^{2} + \delta \\ &\lesssim C(T,\gamma)^{2}\gamma^{2}+\delta. \end{split}$$
(8.231)

We shall show that $\lambda_{\infty}^1 = 1$. Suppose for contradiction that $\lambda_{\infty}^1 = 0$. Fix a time $T \in (\tau_{\min}^1, T_{\max}^1)$. Then, it follows from Lemma 8.7 that

$$\lim_{n \to \infty} \left\| \psi_n^1 \big((\lambda_n^1)^2 T + t_n^1 \big) \right\|_{L^2} = \lim_{n \to \infty} \left\| g_n^1 \sigma_n^1 \widetilde{\psi}^1(T) \right\|_{L^2} = 0.$$
(8.232)

Furthermore, we see from $\mathcal{M}(\psi_n) \equiv \mathcal{M}(\Phi_\omega)$, (8.231) and (8.232) that for any sufficiently small $\omega > 0$ and any $\delta \in (0, \delta_0(T, \gamma))$,

$$\|\Phi_{\omega}\|_{L^{2}}^{2} = \lim_{n \to \infty} \|\psi_{n}\left((\lambda_{n}^{1})^{2}T + t_{n}^{1}\right)\|_{L^{2}}^{2} \lesssim C(T,\gamma)^{2}\gamma^{2} + \delta.$$
(8.233)

Taking $\gamma \to 0$ and $\delta \to 0$, we deduce that $\Phi_{\omega} \equiv 0$. However, this is a contradiction. Thus, we find that $\lambda_{\infty}^1 = 1$, so that $\lambda_n^1 \equiv 1$, $\tau_{\infty}^1 = -t_{\infty}^1$ and $\sigma_n^1 \equiv 1$.

Now, we are in a position to prove (8.217) and (8.218). We first consider the case $\tau_{\infty}^1 = -\infty$. In this case, we have $t_{\infty}^1 = \infty$. Suppose for contradiction that (8.217) failed. Then, for any $T \in (\tau_{\min}^1, T_{\max}^1)$, we can take $T' \in [T, T_{\max}^1)$ such that

$$\widetilde{d}_{\omega}\left(\widetilde{\psi}^{1}(T')\right) \leq \frac{2}{3}R_{*}.$$
(8.234)

Since $\lim_{n\to\infty} t_n^1 = \infty$, we can take a number $N_8(T')$ such that $T' + t_n^1 \ge 0$ for any $n \ge N_8(T')$. Thus, we find from (8.35), (8.231) with $\lambda_n^1 \equiv 1$ and (8.234) that

$$R_* \leq \widetilde{d}_{\omega} \left(\psi_n \left(T' + t_n^1 \right) \right) \leq \widetilde{d}_{\omega} \left(\psi_n^1 \left(T' + t_n^1 \right) \right) + o_{\gamma}(1) + o_{\delta}(1)$$

$$= \widetilde{d}_{\omega} \left(\widetilde{\psi}^1(T') \right) + o_{\gamma}(1) + o_{\delta}(1) \leq \frac{2}{3} R_* + o_{\gamma}(1) + o_{\delta}(1).$$
(8.235)

However, this is impossible for sufficiently small γ and δ . Hence, the claim (8.217) is true. Suppose next that (8.218) failed. Then, for any $T \in (\tau_{\min}^1, T_{\max}^1)$, there exists $T' \in [T, T_{\max}^1)$ such that

$$\mathcal{K}\big(\widetilde{\psi}^1(T')\big) \le \frac{2}{3}\kappa_1(R_*). \tag{8.236}$$

Let $N_8(T')$ be the number obtained above. Then, we find from (8.37), (8.231) with $\lambda_n^1 \equiv 1$ and (8.236) that

$$\kappa_{1}(R_{*}) \leq \mathcal{K}(\psi_{n}(T'+t_{n}^{1}))$$

$$\leq \mathcal{K}(\psi_{n}^{1}(T'+t_{n}^{1})) + o_{\gamma}(1) + o_{\delta}(1) \leq \frac{2}{3}\kappa_{1}(R_{*}) + o_{\gamma}(1) + o_{\delta}(1).$$
(8.237)

However, for a sufficiently small γ and δ , this is a contradiction. Hence, the claim (8.218) is true.

It remains to consider the case $\tau_{\infty}^1 \in \mathbb{R}$. In this case, we have that $\tau_{\min}^1 < \tau_{\infty}^1 = -t_{\infty}^1 < T_{\max}^1$ (see (8.220)). Let $T' \in [-t_{\infty}^1, T_{\max}^1)$ be a time for which (8.234) or (8.236) holds. Then, we can derive a contradiction as well as the case $\tau_{\infty}^1 = -\infty$.

Lemma 8.28. Assume (8.127). Then, we have $\mathcal{M}(\widetilde{\psi}^1) = \mathcal{M}(\Phi_{\omega})$.

Proof of Lemma 8.28. We see from $\lambda_n^1 \equiv 1$ (see Lemma 8.27), (8.50) with k = 2, (8.76), Lemma 8.26 and Lemma 8.7 that for any $\delta > 0$, there exists a number $N(\delta)$ such that for any $n \geq N(\delta)$,

$$\omega \delta \geq \omega \left| \mathcal{M}(\psi_n(0)) - \mathcal{M}(e^{-it_n^1 \Delta} \widetilde{u}^1) \right| - \omega \mathcal{M}(g_n^2 e^{-it_n^2 \Delta} \widetilde{u}^2) - \omega \mathcal{M}(w_n^2)$$

$$\geq \omega \left| \mathcal{M}(\psi_n(0)) - \mathcal{M}(e^{-it_n^1 \Delta} \widetilde{u}^1) \right| - \delta$$

$$\geq \omega \left| \mathcal{M}(\Phi_\omega) - \mathcal{M}(\widetilde{\psi}^1) \right| - \omega \delta - \delta.$$
(8.238)

Since δ is an arbitrary constant, (8.238) implies the desired result.

Proposition 8.29. There exists a solution Ψ to (NLS) such that

$$\Psi \in S^{\varepsilon_*}_{\omega, R_*, +},\tag{8.239}$$

$$\inf_{t \in [0,T_{\max})} \mathcal{K}(\Psi(t)) \ge \frac{\kappa_1(R_*)}{2},\tag{8.240}$$

$$\mathcal{H}(\Psi) = E_*,\tag{8.241}$$

$$\|\Psi\|_{W_{p+1}([0,T_{\max}))\cap W_{2^*}([0,T_{\max}))} = \infty, \tag{8.242}$$

$$\left\{\Psi(t): t \in [0, T_{\max})\right\} \text{ is precompact in } H^1(\mathbb{R}^d), \tag{8.243}$$

where T_{max} denotes the maximal lifespan of Ψ .

Proof of Proposition 8.29. Without loss of generality, we may assume (8.127). Then, we see from Lemma 8.27 that $\lambda_{\infty}^1 = 1$ (hence $\sigma_{\infty}^1 = 1$ and $\tilde{\psi}^1$ is a solution to (NLS)), and there exists $T \in I_{\max}^1$ such that

$$\inf_{t \in [T, T_{\max}^1)} \widetilde{d}_{\omega} \big(\widetilde{\psi}^1(t) \big) \ge \frac{R_*}{2}, \quad \inf_{t \in [T, T_{\max}^1)} \mathcal{K} \big(\widetilde{\psi}^1(t) \big) \ge \frac{\kappa_1(R_*)}{2}.$$
(8.244)

We also see from (8.31), (8.33), (8.36), (8.52), (8.76) and Lemma 8.16 that

$$\mathcal{S}_{\omega}(\widetilde{\psi}^{1}) \leq \lim_{n \to \infty} \mathcal{S}_{\omega}(\psi_{n}) \leq \omega \mathcal{M}(\Phi_{\omega}) + E_{*} = m_{\omega}.$$
(8.245)

This together with Lemma 8.28 and (8.244) shows that $\tilde{\psi}^1(\cdot + T) \in S_{\omega, \frac{R_*}{2}}^{\frac{\varepsilon_*}{2}}$. Furthermore, we find from Lemma 8.2, Lemma 8.4, (8.244) and Lemma 8.3 that $\tilde{\psi}^1(\cdot + T_0) \in S_{\omega, R_*, +}^{\varepsilon_*}$ for some $T_0 \geq T$. Put $\Psi(t) := \tilde{\psi}^1(t+T_0)$. Then, the maximal lifespan of Ψ is $T_{\max}(\Psi) := T_{\max}^1 - T_0$, and $\Psi \in S_{\omega, R_*, +}^{\varepsilon_*}$. We also find from Proposition 8.25, (1.6) and (8.244) that

$$\|\Psi\|_{W_{p+1}([0,T_{\max}(\Psi)))\cap W_{2^*}([0,T_{\max}(\Psi)))} = \infty,$$
(8.246)

$$\mathcal{H}(\Psi) \ge \frac{1}{2} \inf_{t \in [0, T_{\max}(\Psi))} \mathcal{K}(\Psi(t)) \ge \frac{\kappa_1(R_*)}{4}.$$
(8.247)

Since $\mathcal{M}(\Psi) = \mathcal{M}(\tilde{\psi}^1) = \mathcal{M}(\Phi_{\omega})$, the definition of E_* (see (8.28)) together with (8.245) and (8.246) shows the property (8.241).

Now, suppose for contradiction that there existed a number $j \ge 2$ such that $\tilde{\psi}^j$ is non-trivial. Without loss of generality, we may assume that $\tilde{\psi}^2$ is non-trivial. Then, it follows from (8.135) and Theorem 1.2 that

$$\begin{cases} \mathcal{H}(\widetilde{\psi}^2) > 0 & \text{if } \lambda_{\infty}^2 = 1, \\ \mathcal{H}^{\ddagger}(\sigma_{\infty}^2 \widetilde{\psi}^2) > 0 & \text{if } \lambda_{\infty}^2 = 0. \end{cases}$$

$$(8.248)$$

We see from (8.79), Lemma 8.9 and (8.248) that

$$0 < h_2 := \begin{cases} \lim_{n \to \infty} \mathcal{H}\left(g_n^2 \sigma_n^2 e^{-i\frac{t_n^2}{(\lambda_n^2)^2} \Delta} \widetilde{u}^2\right) = \mathcal{H}\left(\widetilde{\psi}^2\right) & \text{if } \lambda_\infty^2 = 1, \\ \lim_{n \to \infty} \mathcal{H}\left(g_n^2 \sigma_n^2 e^{-i\frac{t_n^2}{(\lambda_n^2)^2} \Delta} \widetilde{u}^2\right) = \mathcal{H}^{\ddagger}\left(\sigma_\infty^2 \widetilde{\psi}^2\right) & \text{if } \lambda_\infty^2 = 0. \end{cases}$$

$$(8.249)$$

Furthermore, we see from (8.51), (8.36), (8.120) in Lemma 8.16 and (8.249) that

$$0 = \lim_{n \to \infty} \left\{ \mathcal{H}(\psi_n) - \sum_{j=1}^2 \mathcal{H}\left(g_n^j \sigma_n^j e^{-i\frac{t_n^j}{(\lambda_n^j)^2} \Delta} \widetilde{u}^j\right) - \mathcal{H}(w_n^2) \right\}$$

$$\leq E_* - \lim_{n \to \infty} \mathcal{H}\left(e^{-it_n^1 \Delta} \widetilde{u}^1\right) - h_2.$$
(8.250)

This together with (8.79) shows that

$$\mathcal{H}(\Psi) = \mathcal{H}(\widetilde{\psi}^1) = \lim_{n \to \infty} \mathcal{H}(e^{-it_n^1 \Delta} \widetilde{u}^1) \le E_* - h_2.$$
(8.251)

However, this contradicts the proved property (8.241). Thus, we have found that $\tilde{\psi}^j \equiv 0$ for all $j \geq 2$. Then, we also see that $\tilde{u}^j \equiv 0$ for all $j \geq 2$.

We return to the decomposition (8.43): $\psi_n(0) = e^{-it_n^1 \Delta} \widetilde{u}^1 + w_n^1$. It follows from (8.50), (8.79) and $\mathcal{M}(\widetilde{\psi}^1) = \mathcal{M}(\Psi) = \mathcal{M}(\Phi_\omega) = \mathcal{M}(\psi_n)$ that

$$\lim_{n \to \infty} \|w_n^1\|_{L^2} = 0.$$
(8.252)

Moreover, it follows from (8.36), (8.51), (8.79) and (8.241) that

$$\lim_{n \to \infty} \mathcal{H}\left(w_n^1\right) = 0,\tag{8.253}$$

which together with (8.120) in Lemma 8.16 shows

$$\lim_{n \to \infty} \|\nabla w_n^1\|_{L^2}^2 \le \frac{d}{s_p} \lim_{n \to \infty} \left\{ \mathcal{H}(w_n^1) - \frac{2}{d(p-1)} \mathcal{K}(w_n^k) \right\} \le \frac{d}{s_p} \lim_{n \to \infty} \mathcal{H}(w_n^1) = 0. \quad (8.254)$$

Thus, we find that

$$\lim_{n \to \infty} \|w_n^1\|_{H^1} = 0. \tag{8.255}$$

Finally, we shall prove the precompactness of $\{\Psi(t)\}$ in $H^1(\mathbb{R}^d)$. Take a sequence $\{\tau_n\}$ in $[0, T_{\max}(\Psi))$. By the continuity in time, it suffices to consider the case where $\lim_{n\to\infty} \tau_n = T_{\max}(\Psi)$. Applying the above argument to $\{\Psi(t+\tau_n)\}$, we can take a subsequence of $\{\tau_n\}$ (still denoted by the same symbol), a sequence $\{t_n\}$ in \mathbb{R} , a sequence $\{w_n\}$ in $H^1(\mathbb{R}^d)$ and a function $\tilde{u} \in H^1(\mathbb{R}^d)$ such that

$$\Psi(\tau_n) = e^{-it_n \Delta} \widetilde{u} + w_n, \qquad (8.256)$$

$$\lim_{n \to \infty} \|w_n\|_{H^1} = 0, \tag{8.257}$$

$$t_{\infty} := \lim_{n \to \infty} t_n \in \mathbb{R} \cup \{\infty\},\tag{8.258}$$

and if $t_{\infty} = \infty$, then $T_{\min}(\Psi) = -\infty$. We see from (8.256) and (8.257) that

$$\lim_{n \to \infty} \|\Psi(\tau_n) - e^{-it_n \Delta} \widetilde{u}\|_{H^1} = 0.$$
(8.259)

Hence, in order to prove the precompactness, it suffices to show that $t_{\infty} \in \mathbb{R}$. Suppose for contradiction that $t_{\infty} = \infty$. Then, we find from Strichartz' estimate and (8.259) that

$$\lim_{n \to \infty} \|e^{it\Delta} \Psi(\tau_n)\|_{V_{2+\frac{4}{d}}((-\infty,0]) \cap W_{2^*}((-\infty,0])} = \lim_{n \to \infty} \|e^{i(t+t_n)\Delta} \Psi(\tau_n)\|_{V_{2+\frac{4}{d}}((-\infty,-t_n]) \cap W_{2^*}((-\infty,-t_n])} \\
\leq \lim_{n \to \infty} \|e^{i(t+t_n)\Delta} e^{-it_n\Delta} \widetilde{u}\|_{V_{2+\frac{4}{d}}((-\infty,-t_n]) \cap W_{2^*}((-\infty,-t_n])} \\
+ \lim_{n \to \infty} \|e^{i(t+t_n)\Delta} \{\Psi(\tau_n) - e^{-it_n\Delta} \widetilde{u}\}\|_{V_{2+\frac{4}{d}}((-\infty,-t_n]) \cap W_{2^*}((-\infty,-t_n])} \\
\lesssim \lim_{n \to \infty} \|\langle \nabla \rangle e^{it\Delta} \widetilde{u}\|_{V_{2+\frac{4}{d}}((-\infty,-t_n]) \cap V_{2^*}((-\infty,-t_n])} + \lim_{n \to \infty} \|\Psi(\tau_n) - e^{-it_n\Delta} \widetilde{u}\|_{H^1} \\
= 0.$$
(8.260)

Thus, the small-data theory (see Lemma D.1 below) shows that there exists a number $N \ge 1$ such that for any $n \ge N$,

$$\|\Psi\|_{W_{p+1}((-\infty,\tau_n])\cap W_{2^*}((-\infty,\tau_n])} = \|\Psi(\cdot+\tau_n)\|_{W_{p+1}((-\infty,0])\cap W_{2^*}((-\infty,0])} \lesssim 1, \qquad (8.261)$$

which together with $\lim_{n\to\infty} \tau_n = T_{\max}(\Psi)$ shows

$$\|\Psi\|_{W_{p+1}((-\infty,T_{\max}(\Psi)))\cap W_{2^*}((-\infty,T_{\max}(\Psi)))} \lesssim 1.$$
(8.262)

However, this contradicts (8.242). Thus, we have $t_{\infty} \in \mathbb{R}$ and therefore the precompactness holds.

Now, we finish the proof of Proposition 8.6. Under the hypothesis (8.30), we have shown the existence of solution Ψ with the properties (8.239) through (8.243) in Proposition 8.29. However, the same argument as the proof of Proposition 5.3 in [18] shows that such a solution Ψ never exists, and therefore we have arrived at a contradiction. Thus, the hypothesis (8.30) is false, and we have proved Proposition 8.6.

A Existence of ground state

In this section, we show the existence of ground state of (1.7) in three dimensions. We also give a proof of (1.34) at the end of this section.

Proposition A.1. Assume d = 3 and $1 . Then, there exists <math>\omega_3 > 0$ such that for any $0 < \omega < \omega_3$, a ground state of (1.7) exists.

In order to prove Proposition A.1, we consider the following equation in \mathbb{R}^d :

$$v - \Delta v - \omega^{-\frac{5-p}{4}} |v|^{p-1} v - |v|^{\frac{4}{d-2}} v = 0, \qquad v \in H^1(\mathbb{R}^d) \setminus \{0\}.$$
 (A.1)

The action associated with the equation (A.1) is

$$\widetilde{\mathcal{S}}_{\omega}(v) = \frac{1}{2} \|v\|_{L^{2}}^{2} + \frac{1}{2} \|\nabla v\|_{L^{2}}^{2} - \frac{\omega^{-\frac{5-p}{4}}}{p+1} \|v\|_{L^{p+1}}^{p+1} - \frac{1}{2^{*}} \|v\|_{L^{2^{*}}}^{2^{*}}.$$
(A.2)

It is easy to see that: if d = 3 and v is a solution to (A.1), then the H^1 -rescaled function $\omega^{\frac{1}{4}}v(\sqrt{\omega})$ becomes a solution to (1.7) in \mathbb{R}^3 ; and if u is a solution to (1.7) in \mathbb{R}^3 , then $\omega^{-\frac{1}{4}}u(\sqrt{\omega})$ becomes a solution to (A.1). Furthermore, if d = 3 and $\widetilde{\Phi}_{\omega}$ is a ground state of (A.1) (namely, $\widetilde{\Phi}_{\omega}$ is a solution which minimizes the action \widetilde{S}_{ω} among the solutions to (A.1)), then for any solution u to (1.7) with d = 3,

$$\mathcal{S}_{\omega}(\omega^{\frac{1}{4}}\widetilde{\Phi}_{\omega}(\sqrt{\omega}\cdot)) = \widetilde{\mathcal{S}}_{\omega}(\widetilde{\Phi}_{\omega}) \le \widetilde{\mathcal{S}}_{\omega}(\omega^{-\frac{1}{4}}u(\cdot/\sqrt{\omega})) = \mathcal{S}_{\omega}(u).$$
(A.3)

Thus, we find that it is sufficient for Proposition A.1 to prove the existence of ground state of (A.1) in \mathbb{R}^3 . An advantage to consider the equation (A.2) is that we can take into account the smallness of ω easier than the original problem (1.7) in our variational argument below.

We define

$$\widetilde{m}_{\omega} := \inf \left\{ \widetilde{\mathcal{T}}_{\omega}(v) \colon v \in H^1(\mathbb{R}^3) \setminus \{0\}, \ \widetilde{\mathcal{N}}_{\omega}(v) \le 0 \right\},$$
(A.4)

where

$$\widetilde{\mathcal{N}}_{\omega}(v) := \|v\|_{L^{2}}^{2} + \|\nabla v\|_{L^{2}}^{2} - \omega^{-\frac{5-p}{4}} \|v\|_{L^{p+1}}^{p+1} - \|v\|_{L^{2^{*}}}^{2^{*}},$$
(A.5)

$$\widetilde{\mathcal{T}}_{\omega}(v) := \frac{p-1}{2(p+1)} \left\{ \|v\|_{L^2}^2 + \|\nabla v\|_{L^2}^2 \right\} + \frac{5-p}{6(p+1)} \|v\|_{L^{2^*}}^{2^*}.$$
(A.6)

It is easy to verify that

$$\widetilde{\mathcal{T}}_{\omega} = \widetilde{\mathcal{S}}_{\omega} - \frac{1}{p+1} \widetilde{\mathcal{N}}_{\omega}.$$
(A.7)

Moreover, we have the following:

Lemma A.2. Assume d = 3 and 1 . Then, we have the following:(i)

$$\widetilde{m}_{\omega} = \inf \left\{ \widetilde{\mathcal{S}}_{\omega}(v) \colon v \in H^1(\mathbb{R}^3) \setminus \{0\}, \ \widetilde{\mathcal{N}}_{\omega}(v) = 0 \right\}.$$
(A.8)

(ii) Any minimizer of the variational problem associated with \tilde{m}_{ω} becomes a ground state of (A.1).

Proof of Lemma A.2. First, we shall prove (i). Since $\widetilde{\mathcal{T}}_{\omega}(v) = \widetilde{\mathcal{S}}_{\omega}(v)$ for every $v \in H^1(\mathbb{R}^3)$ with $\widetilde{\mathcal{N}}_{\omega}(v) = 0$, we see that $\widetilde{m}_{\omega} \leq \inf \{\widetilde{\mathcal{S}}_{\omega}(v) : v \in H^1(\mathbb{R}^3) \setminus \{0\}, \ \widetilde{\mathcal{N}}_{\omega}(v) = 0\}$. In order to prove the opposite inequality, we note that for any $v_0 \in H^1(\mathbb{R}^3) \setminus \{0\}$ with $\widetilde{\mathcal{N}}_{\omega}(v_0) \leq 0$, there exists $0 < \lambda_0 \leq 1$ such that $\widetilde{\mathcal{N}}_{\omega}(\lambda_0 v_0) = 0$. Furthermore, this together with (A.7) shows that

$$\inf\left\{\widetilde{\mathcal{S}}_{\omega}(v)\colon v\in H^{1}(\mathbb{R}^{3})\setminus\{0\},\ \widetilde{\mathcal{N}}_{\omega}(v)=0\right\}\leq\widetilde{\mathcal{S}}_{\omega}(\lambda_{0}v_{0})=\widetilde{\mathcal{T}}_{\omega}(\lambda_{0}v_{0})\leq\widetilde{\mathcal{T}}_{\omega}(v_{0}).$$
 (A.9)

Since v_0 is an arbitrary function in $H^1(\mathbb{R}^3) \setminus \{0\}$ with $\widetilde{\mathcal{N}}_{\omega}(v_0) \leq 0$, we find that (A.8) holds.

Next, we shall prove the claim (ii). It is well known that if \widetilde{Q}_{ω} is a minimizer of $\inf \{\widetilde{S}_{\omega}(v) : v \in H^1(\mathbb{R}^3) \setminus \{0\}, \ \widetilde{\mathcal{N}}_{\omega}(v) = 0\}$, then \widetilde{Q}_{ω} is also a ground state of (A.1). Hence, we see from (A.7) and (A.8) that it suffices to prove that $\widetilde{\mathcal{N}}_{\omega}(\widetilde{Q}_{\omega}) = 0$ for all minimizer \widetilde{Q}_{ω} for \widetilde{m}_{ω} . Suppose to the contrary that there exists a minimizer \widetilde{Q}_{ω} for \widetilde{m}_{ω} such that $\widetilde{\mathcal{N}}_{\omega}(\widetilde{Q}_{\omega}) < 0$. Then, we could take $0 < \lambda_0 < 1$ such that $\widetilde{\mathcal{N}}_{\omega}(\lambda_0 \widetilde{Q}_{\omega}) = 0$. Furthermore, we see from the definition of \widetilde{m}_{ω} (see (A.4)) and $0 < \lambda_0 < 1$ that

$$\widetilde{m}_{\omega} \le \widetilde{\mathcal{T}}_{\omega}(\lambda_0 \widetilde{Q}_{\omega}) < \widetilde{\mathcal{T}}_{\omega}(\widetilde{Q}_{\omega}) = \widetilde{m}_{\omega}, \tag{A.10}$$

which is a contradiction. Thus, $\widetilde{\mathcal{N}}_{\omega}(\widetilde{Q}_{\omega}) = 0$.

A key lemma to prove the existence of the minimizer is the following:

Lemma A.3. Assume d = 3 and $1 . Then, there exists <math>\omega_3 > 0$ such that for any $0 < \omega < \omega_3$,

$$0 < \widetilde{m}_{\omega} < \frac{1}{3}\sigma^{\frac{3}{2}}.\tag{A.11}$$

Proof of Lemma A.3. First, we shall show that $\widetilde{m}_{\omega} > 0$. Let v be a nontrivial function in $H^1(\mathbb{R}^3)$ such that $\widetilde{\mathcal{N}}_{\omega}(v) \leq 0$. Then, it follows from Sobolev's embedding that

$$\|v\|_{H^{1}}^{2} = \|v\|_{L^{2}}^{2} + \|\nabla v\|_{L^{2}}^{2} \le \omega^{-\frac{5-p}{4}} \|v\|_{L^{p+1}}^{p+1} + \|v\|_{L^{2^{*}}}^{2^{*}} \lesssim \omega^{-\frac{5-p}{4}} \|v\|_{H^{1}}^{p+1} + \|v\|_{H^{1}}^{6}.$$
 (A.12)

In particular, (A.12) implies that

$$\omega^{\frac{5-p}{4}} \lesssim \|v\|_{H^1}^{p-1} + \omega^{\frac{5-p}{4}} \|v\|_{H^1}^4.$$
(A.13)

Since the implicit constant of the inequality (A.13) depends only on p, we find that for any $1 and any <math>\omega > 0$, there exists $C(\omega) > 0$ such that $C(\omega) \leq \tilde{\mathcal{T}}_{\omega}(v)$ for all $v \in H^1(\mathbb{R}^d) \setminus \{0\}$ with $\tilde{\mathcal{N}}_{\omega}(v) \leq 0$. Thus, we have proved that $\tilde{m}_{\omega} > 0$.

Next, we shall show that $\widetilde{m}_{\omega} < \frac{1}{3}\sigma^{\frac{3}{2}}$. We define

$$W_{\omega}(x) := \omega^{-\frac{1}{2}} W(x/\omega) = \omega^{\frac{1}{2}} \left(\omega^2 + \frac{|x|^2}{3}\right)^{-\frac{1}{2}}.$$
 (A.14)

Note that W_{ω} does not belong to $L^2(\mathbb{R}^3)$. Let χ be an even smooth function on \mathbb{R} such that $\chi(r) = 1$ for $0 \le r \le 1$, $\chi(r) = 0$ for $r \ge 2$, and χ is non-increasing on $[0, \infty)$. Then, we define

$$W_{\omega}(x) := \chi(|x|)W_{\omega}(x). \tag{A.15}$$

We see from (1.28) that

$$\|\nabla \widetilde{W}_{\omega}\|_{L^2}^2 = \sigma^{\frac{3}{2}} + O(\omega), \qquad (A.16)$$

$$\|\widetilde{W}_{\omega}\|_{L^{2^*}}^{2^*} = \sigma^{\frac{3}{2}} + O(\omega^3).$$
(A.17)
Moreover, we find that for any $2 \le q < 6$,

$$\|\widetilde{W}_{\omega}\|_{L^{q}}^{q} = \begin{cases} O(\omega^{\frac{6-q}{2}}) & \text{if } 3 < q < 6, \\ O(\omega^{\frac{3}{2}}|\log\omega|) & \text{if } q = 3, \\ O(\omega^{\frac{q}{2}}) & \text{if } 2 \le q < 3. \end{cases}$$
(A.18)

Note here that there exists a unique $\tau_{\omega,0} > 0$ such that

$$\widetilde{\mathcal{N}}_{\omega}(\tau_{\omega,0}\widetilde{W}_{\omega}) = 0. \tag{A.19}$$

Furthermore, it follows from (A.16) through (A.19) that: if 2 , then

$$0 = \tau_{\omega,0}^{2} \|\widetilde{W}_{\omega}\|_{L^{2}}^{2} + \tau_{\omega,0}^{2} \|\nabla\widetilde{W}_{\omega}\|_{L^{2}}^{2} - \omega^{-\frac{5-p}{4}} \tau_{\omega,0}^{p+1} \|\widetilde{W}_{\omega}\|_{L^{p+1}}^{p+1} - \tau_{\omega,0}^{2^{*}} \|\widetilde{W}_{\omega}\|_{L^{2^{*}}}^{2^{*}}$$

$$= \tau_{\omega,0}^{2} O(\omega) + \tau_{\omega,0}^{2} \{\sigma^{\frac{3}{2}} + O(\omega)\} - \tau_{\omega,\varepsilon}^{p+1} O(\omega^{\frac{5-p}{4}}) - \tau_{\omega,0}^{6} \{\sigma^{\frac{3}{2}} + O(\omega^{3})\};$$
(A.20)

if p = 2, then

$$0 = \tau_{\omega,0}^2 O(\omega) + \tau_{\omega,0}^2 \{ \sigma^{\frac{3}{2}} + O(\omega) \} - \tau_{\omega,0}^3 O(\omega^{\frac{3}{4}} |\log \omega|) - \tau_{\omega,0}^6 \{ \sigma^{\frac{3}{2}} + O(\omega^3) \};$$
(A.21)

and if 1 , then

$$0 = \tau_{\omega,0}^2 O(\omega) + \tau_{\omega,0}^2 \{ \sigma^{\frac{3}{2}} + O(\omega) \} - \tau_{\omega,0}^{p+1} O(\omega^{\frac{p+1}{2} - \frac{5-p}{4}}) - \tau_{\omega,0}^6 \{ \sigma^{\frac{3}{2}} + O(\omega^3) \}.$$
 (A.22)

Thus, we find from (A.20) through (A.22) that

$$\tau_{\omega,0}^4 = 1 + O(\omega) - \tau_{\omega,0}^{p-1} o_\omega(1), \tag{A.23}$$

where $o_{\omega}(1)$ means a certain function such that $\lim_{\omega \to 0} o_{\omega}(1) = 0$. Furthermore, since 0 , this implies that

$$\lim_{\omega \to +0} \tau_{\omega,0} \ge \frac{1}{2}.$$
(A.24)

We introduce a function $y_{\omega} \colon (0, \infty) \to \mathbb{R}$ as

$$y_{\omega}(t) := \frac{1}{2} t^2 \left\{ \|\widetilde{W}_{\omega}\|_{L^2}^2 + \|\nabla\widetilde{W}_{\omega}\|_{L^2}^2 \right\} - \frac{t^{2^*}}{2^*} \|\widetilde{W}_{\omega}\|_{L^{2^*}}^{2^*}.$$
(A.25)

It is easy to verify that the function y_{ω} attains its maximum only at the point

$$\tau_{\omega,\max} := \frac{\left\{ \|\widetilde{W}_{\omega}\|_{L^{2}}^{2} + \|\nabla\widetilde{W}_{\omega}\|_{L^{2}}^{2} \right\}^{\frac{1}{4}}}{\|\widetilde{W}_{\omega}\|_{L^{2^{*}}}^{\frac{3}{2}}}.$$
(A.26)

Note here that it follows from the definition of σ (see (1.27)), (A.16) and (A.17) that

$$\frac{\|\nabla \widetilde{W}_{\omega}\|_{L^{2}}^{2}}{\|\widetilde{W}_{\omega}\|_{L^{2^{*}}}^{2}} = \frac{\sigma^{\frac{3}{2}} + O(\omega)}{\sigma^{\frac{1}{2}} + O(\omega^{3})} = \sigma + O(\omega).$$
(A.27)

Moreover, we see from (A.16) and (A.18) that

$$\frac{\|\widetilde{W}_{\omega}\|_{L^{2}}^{2}}{\|\nabla\widetilde{W}_{\omega}\|_{L^{2}}^{2}} = O(\omega).$$
(A.28)

Hence, if ω is sufficiently small, then we have

$$y(\tau_{\omega,\max}) = \frac{1}{3} \left(\frac{\|\widetilde{W}_{\omega}\|_{L^{2}}^{2} + \|\nabla\widetilde{W}_{\omega}\|_{L^{2}}^{2}}{\|\widetilde{W}_{\omega}\|_{L^{2}^{*}}^{2}} \right)^{\frac{3}{2}} = \frac{1}{3} \frac{\|\nabla\widetilde{W}_{\omega}\|_{L^{2}}^{3}}{\|\widetilde{W}_{\omega}\|_{L^{2}^{*}}^{3}} \left(1 + \frac{\|\widetilde{W}_{\omega}\|_{L^{2}}^{2}}{\|\nabla\widetilde{W}_{\omega}\|_{L^{2}}^{2}} \right)^{\frac{3}{2}}$$

$$= \frac{1}{3} \frac{\|\nabla\widetilde{W}_{\omega}\|_{L^{2}}^{3}}{\|\widetilde{W}_{\omega}\|_{L^{2}}^{3}} \left\{1 + \frac{3}{2} \frac{\|\widetilde{W}_{\omega}\|_{L^{2}}^{2}}{\|\nabla\widetilde{W}_{\omega}\|_{L^{2}}^{2}} + O\left(\frac{\|\widetilde{W}_{\omega}\|_{L^{2}}^{4}}{\|\nabla\widetilde{W}_{\omega}\|_{L^{2}}^{4}}\right)\right\} = \frac{1}{3}\sigma^{\frac{3}{2}} + O(\omega).$$
(A.29)

Furthermore, it follows from (A.8), (A.18), (A.19), the same computations concerning $\|\widetilde{W}_{\omega}\|_{L^{p+1}}^{p+1}$ as (A.20) through (A.22), and (A.24) that

$$\widetilde{m}_{\omega} \leq \widetilde{\mathcal{S}}_{\omega}(\tau_{\omega,0}\widetilde{W}_{\omega}) = y_{\omega}(\tau_{\omega,0}) - \frac{\omega^{-\frac{5-p}{4}}}{p+1}\tau_{\omega,0}^{p+1} \|\widetilde{W}_{\omega}\|_{L^{p+1}}^{p+1}$$

$$\leq y_{\omega}(\tau_{\omega,\max}) - c\omega^{\theta} = \frac{1}{3}\sigma^{\frac{3}{2}} + O(\omega) - c\omega^{\theta}$$
(A.30)

for some $0 < \theta < 1$ and some c > 0 depending only on p. Thus, we find that if ω is sufficiently small depending only on p, then (A.11) holds.

Now, we give a brief proof of Proposition A.1:

Proof of Proposition A.1. As mentioned above, if d = 3 and $\tilde{\Phi}_{\omega}$ is a ground state of (A.1), then $\omega^{\frac{1}{4}} \tilde{\Phi}_{\omega}(\sqrt{\omega} \cdot)$ is one of (1.7). We also see from Lemma A.2 that any minimizer of the variational problem associated with \tilde{m}_{ω} becomes a ground state of (A.1). Thus, it suffices to prove that for the frequency $\omega_3 > 0$ given in Lemma A.3 and any $0 < \omega < \omega_3$, there exists a minimizer for \tilde{m}_{ω} . The proof is standard: using Lemma A.3 and the Schwarz symmetrization, we can prove the existence of minimizer for \tilde{m}_{ω} (see, e.g, Propositoin 2.1 in [2], and [7]).

At the end of this section, we discuss the variational value m_{ω} defined by (1.9) in three dimensions.

Proposition A.4. Assume d = 3 and $1 + \frac{4}{3} . Let <math>\omega_3$ be the frequency given in Proposition A.1. Then, for any $0 < \omega < \omega_3$,

$$0 < m_{\omega} < \frac{1}{3}\sigma^{\frac{3}{2}}.\tag{A.31}$$

Furthermore, any ground state Q_{ω} of (1.7) satisfies

$$\mathcal{S}_{\omega}(Q_{\omega}) = m_{\omega}.\tag{A.32}$$

Proof of Proposition A.4. First, we shall show that $m_{\omega} > 0$. To this end, we recall that (1.10) holds for d = 3 and $1 + \frac{4}{3} . Let <math>u$ be a nontrivial function in $H^1(\mathbb{R}^3)$ such that $\mathcal{K}(u) \leq 0$. Then, it follows from Sobolev's embedding that

$$\|\nabla u\|_{L^{2}}^{2} \lesssim \|u\|_{L^{2}}^{\frac{5-p}{2}} \|\nabla u\|_{L^{2}}^{\frac{3(p-1)}{2}} + \|\nabla u\|_{L^{2}}^{6}.$$
(A.33)

In particular, if $\|u\|_{L^2} \ll 1$, then (A.33) implies that $1 \lesssim \|\nabla u\|_{L^2}$. Hence, if $\|u\|_{L^2} \ll 1$, then $\mathcal{I}_{\omega}(u) \geq \frac{p-1-\frac{4}{3}}{p-1} \|\nabla u\|_{L^2}^2 \gtrsim 1$. On the other hand, if $\|u\|_{L^2} \gtrsim 1$, then $\mathcal{I}_{\omega}(u) \geq \frac{\omega}{2} \|u\|_{L^2}^2 \gtrsim \omega$. Thus, we find that $\mathcal{I}_{\omega}(u) \gtrsim \min\{1,\omega\}$ for all $u \in H^1(\mathbb{R}^3) \setminus \{0\}$ with $\mathcal{K}(u) \leq 0$. This together with (1.10) shows $m_{\omega} > 0$.

Next, we shall prove (A.31). Let Q_{ω} be a minimizer of the variational problem associated with \widetilde{m}_{ω} (see the proof of Proposition A.1 for the existence of the minimizer). Then, it follows from Lemma A.2 that \widetilde{Q}_{ω} is a ground state of (A.1). Furthermore, it is easy to verify that $\widetilde{\mathcal{N}}_{\omega}(\widetilde{Q}) = 0$. Put $Q_{\omega} := \omega^{\frac{1}{4}} \widetilde{Q}_{\omega}(\sqrt{\omega} \cdot)$. Then, Q_{ω} becomes a ground state of (1.7) and satisfies the following identity:

$$\mathcal{K}(Q_{\omega}) = 0. \tag{A.34}$$

Hence, we find from the definition of m_{ω} , (A.8) and Lemma A.3 that

$$m_{\omega} \leq \mathcal{S}_{\omega}(Q_{\omega}) = \widetilde{\mathcal{S}}_{\omega}(\widetilde{Q}_{\omega}) = \widetilde{m}_{\omega} < \frac{1}{3}\sigma^{\frac{3}{2}}.$$
 (A.35)

Then, the standard argument (see, e.g., Proposition 2.1 in [2]) proves the existence of minimizer of the variational problem associated with m_{ω} . Furthermore, it is well-known that any minimizer for m_{ω} becomes a ground state of (1.7) (see, e.g., Proposition 1.1 in [2]). Thus, (A.32) holds.

Next, we give a proof of (1.34).

Proof of (1.34). We find from (1.28) that the function W (see (1.26)) satisfies $\mathcal{K}^{\ddagger}(W) = 0$ and $\mathcal{I}^{\ddagger}(W) = \mathcal{H}^{\ddagger}(W) = \frac{1}{d}\sigma^{\frac{d}{2}}$. Hence, $m^{\ddagger} \leq \frac{1}{d}\sigma^{\frac{d}{2}}$ and therefore it remains to prove $\frac{1}{d}\sigma^{\frac{d}{2}} \leq m^{\ddagger}$. Let $u \in H^1(\mathbb{R}^d) \setminus \{0\}$ satisfy $\mathcal{K}^{\ddagger}(u) \leq 0$. Then, we can take $0 < \lambda_0 \leq 1$ such that $\mathcal{K}^{\ddagger}(\lambda_0 u) = 0$. Furthermore, it follows from the definition of σ (see (1.27)) and $\mathcal{K}^{\ddagger}(\lambda_0 u) = \lambda_0^2 \|\nabla u\|_{L^2}^2 - \lambda_0^{2^*} \|u\|_{L^{2^*}}^{2^*} = 0$ that

$$\sigma \le \frac{\|\nabla u\|_{L^2}^2}{\|u\|_{L^{2^*}}^2} \le \lambda_0^{\frac{4}{d-2}} \|u\|_{L^{2^*}}^{\frac{4}{d-2}}.$$
(A.36)

Raising the both sides of (A.36) to the power of $\frac{d}{2}$, and using $\mathcal{K}^{\ddagger}(\lambda_0 u) = 0$ and $\lambda_0 \leq 1$, we see that

$$\frac{1}{d}\sigma^{\frac{d}{2}} \leq \frac{1}{d} \|\lambda_0 u\|_{L^{2^*}}^{2^*} = \mathcal{H}^{\ddagger}(\lambda_0 u)
= \mathcal{H}^{\ddagger}(\lambda_0 u) - \frac{2}{d(p-1)} \mathcal{K}^{\ddagger}(\lambda_0 u) = \mathcal{I}^{\ddagger}(\lambda_0 u) \leq \mathcal{I}^{\ddagger}(u).$$
(A.37)

Since u is arbitrary, we find that $\frac{1}{d}\sigma^{\frac{d}{2}} \leq m^{\ddagger}$. Thus, we have completed the proof.

B Fundamental properties of the linearized operators

In this section, we discuss fundamental properties of the operators $L_{\omega,+}$ and $L_{\omega,-}$ defined by (4.4) and (4.5), respectively. Throughout this section, we assume that $d \ge 3$, $1 + \frac{4}{d} and <math>0 < \omega < \omega_1$, where ω_1 is the frequency given by Proposition 2.4.

First, we can easily verify that for any $f \in H^1(\mathbb{R}^d)$,

$$\left| \langle L_{\omega,+}f,f \rangle_{H^{-1},H^{1}} \right| + \left| \langle L_{\omega,-}f,f \rangle_{H^{-1},H^{1}} \right| \lesssim \left(\omega + 1 + \|\Phi_{\omega}\|_{L^{p+1}}^{p-1} + \|\Phi_{\omega}\|_{L^{2^{*}}}^{\frac{4}{d-2}} \right) \|f\|_{H^{1}}^{2},$$
(B.1)

where the implicit constant is independent of ω . Moreover, we see from (4.3) and (4.6) that for any $u, v \in H^1_{real}(\mathbb{R}^d)$,

$$\left[\mathcal{S}_{\omega}^{\prime\prime}(\Phi_{\omega})u\right]v = \langle L_{\omega,+}\Re[u] + iL_{\omega,-}\Im[u], v \rangle_{H^{-1},H^{1}}.$$
(B.2)

Lemma B.1. The operator $L_{\omega,-}$ is non-negative and Ker $L_{\omega,-} = \text{span}\{\Phi_{\omega}\}$; in particular, there exists $\delta_{\omega} > 0$ depending on ω such that for any $u \in H^1(\mathbb{R}^d)$ with $(u, \Phi_{\omega})_{L^2_{real}} = 0$,

$$\langle L_{\omega,-}u, u \rangle_{H^{-1},H^1} \ge \delta_{\omega} \|u\|_{H^1}^2.$$
 (B.3)

Proof of Lemma B.1. We can prove the claim in a way similar to [30].

Differentiating the equation (1.7) for Φ_{ω} with respect to x_j , $1 \leq j \leq d$, we have

$$0 = \partial_j(\omega\Phi_\omega - \Delta\Phi_\omega - \Phi_\omega^p - \Phi_\omega^{2^*-1}) = L_{\omega,+}\partial_j\Phi_\omega.$$
(B.4)

Lemma B.2. There exists $\omega_3 > 0$ such that for any $\omega \in (0, \omega_3)$, we have

$$\operatorname{Ker} L_{\omega,+} = \operatorname{span} \{ \partial_1 \Phi_\omega, \dots, \partial_d \Phi_\omega \}.$$
(B.5)

Proof of Lemma B.2. We can prove this lemma by regarding the operator $L_{\omega,+}$ as a perturbation of L_{ω}^{\dagger} (see [13] and Lemma 2.5).

Now, we introduce the functional \mathcal{N}_{ω} as

$$\mathcal{N}_{\omega}(u) := \omega \|u\|_{L^2}^2 + \|\nabla u\|_{L^2}^2 - \|u\|_{L^{p+1}}^{p+1} - \|u\|_{L^{2^*}}^{2^*}.$$
 (B.6)

Then, we can verify that

$$m_{\omega} = \inf \left\{ \mathcal{S}_{\omega}(u) \colon u \in H^1(\mathbb{R}^d) \setminus \{0\}, \ \mathcal{N}_{\omega}(u) = 0 \right\}.$$
(B.7)

Lemma B.3. For any $u \in H^1(\mathbb{R}^d)$ satisfying

$$((p-1)\Phi_{\omega}^{p} + (2^{*}-2)\Phi_{\omega}^{2^{*}-1}, u)_{L_{real}^{2}} = 0,$$
 (B.8)

we have

$$\left\langle L_{\omega,+}u, u \right\rangle_{H^{-1}, H^1} \ge 0. \tag{B.9}$$

Lemma B.4. As an operator in $L^2_{rad}(\mathbb{R}^d)$, $L_{\omega,+}$ has only one negative eigenvalue which is non-degenerate and 0 is not an eigenvalue.

Proof of Lemma B.4. We can prove this lemma in a way similar to [30] and [13]. \Box

Lemma B.5. Let ω_2 be the frequency given by Proposition 4.1. Then, there exists $\omega_0 \in (0, \omega_2)$ such that the following holds for all $0 < \omega < \omega_0$: Let $\mu > 0$ be a positive eigenvalue of $i\mathcal{L}_{\omega}$ as an operator in $L^2_{real}(\mathbb{R}^d)$, and let \mathcal{U}_+ be a corresponding eigenfunction. Then, we have:

(i) For any non-trivial, real-valued radial function $g \in H^1(\mathbb{R}^d)$ with $(g, f_2)_{L^2_{n-1}} = 0$,

$$\langle L_{\omega,+}g,g\rangle_{H^{-1},H^1} \sim ||g||_{H^1}^2,$$
 (B.10)

where the implicit constant may depend on ω .

(ii) For any non-trivial, real-valued radial function $g \in H^1(\mathbb{R}^d)$ with $(g, \partial_\omega \Phi_\omega)_{L^2} = 0$,

$$\langle L_{\omega,-}g,g\rangle_{H^{-1},H^1} \sim ||g||_{H^1}^2,$$
 (B.11)

where the implicit constant may depend on ω .

Proof of Lemma B.5. We prove the claim (i).

First, we shall show that for any $0 < \omega < \omega_2$ and any nontrivial, radial real-valued function $g \in H^1(\mathbb{R}^d)$ satisfying $(g, f_2)_{L^2_{real}} = 0$,

$$\langle L_{\omega,+}g,g \rangle_{H^{-1},H^1} > 0.$$
 (B.12)

Suppose for contradiction that there exists $\omega > 0$ and a nontrivial radial real-valued function $g_{-} \in H^{1}(\mathbb{R}^{d})$ such that

$$(g_{-}, f_2)_{L^2_{real}} = 0, (B.13)$$

$$\langle L_{\omega,+}g_{-},g_{-}\rangle_{H^{-1},H^{1}} \le 0.$$
 (B.14)

Since $L_{\omega,+}$ is self-adjoint in $L^2_{real}(\mathbb{R}^d)$, we see from (4.109) and (B.13) that

$$\langle L_{\omega,+}g_{-}, f_1 \rangle_{H^{-1},H^1} = (g_{-}, L_{\omega,+}f_1)_{L^2_{real}} = -\mu(g_{-}, f_2)_{L^2_{real}} = 0.$$
 (B.15)

Moreover, it follows from (4.109) and (4.114) that

$$\langle L_{\omega,+}f_1, f_1 \rangle_{H^{-1},H^1} = -(\mu f_2, f_1)_{L^2_{real}} = -\mu (f_1, f_2)_{L^2_{real}} < 0.$$
 (B.16)

Note here that (B.15) and (B.16) show that g_{-} and f_{1} are linearly independent in $L^{2}_{real}(\mathbb{R}^{d})$. We see from the hypothesis (B.14) and (B.15) that for any $a, b \in \mathbb{R}$,

$$\langle L_{\omega,+} \left(af_1 + b \frac{g_-}{\|g_-\|_{L^2}} \right), \left(af_1 + b \frac{g_-}{\|g_-\|_{L^2}} \right) \rangle_{H^{-1},H^1}$$

$$= a^2 (L_{\omega,+}f_1, f_1)_{L^2_{real}} + \frac{b^2}{\|g_-\|_{L^2}^2} \langle L_{\omega,+}g_-, g_- \rangle_{H^{-1},H^1} + \frac{2ab}{\|g_-\|_{L^2}} \langle L_{\omega,+}g_-, f_1 \rangle_{H^{-1},H^1}$$

$$\leq a^2 (L_{\omega,+}f_1, f_1)_{L^2_{real}} \leq 0.$$
(B.17)

Put

$$e_1 := \frac{g_-}{\|g_-\|_{L^2}}, \quad e_2 := \frac{f_1 - (f_1, e_1)_{L^2_{real}} e_1}{\|f_1 - (f_1, e_1)_{L^2_{real}} e_1\|_{L^2}}.$$
 (B.18)

Then, it is clear that $||e_1||_{L^2} = ||e_2||_{L^2} = 1$ and $(e_1, e_2)_{L^2_{real}} = 0$. Moreover, we find from (B.17) that for any $\alpha, \beta \in \mathbb{R}$,

$$\langle L_{\omega,+}(\alpha e_1 + \beta e_2), \, \alpha e_1 + \beta e_2 \rangle_{H^{-1},H^1} \le 0.$$
(B.19)

Since Weyl's essential spectrum theorem shows $\sigma_{ess}(L_{\omega,+}) = \sigma_{ess}(-\Delta + \omega^2) = [\omega^2, \infty)$, the min-max theorem (see Theorem 12.1 in [23]) together with (B.19) shows that the second eigenvalue of $L_{\omega,+}$ is non-positive. However, this contradicts the fact that $L_{\omega,+}$ has only one non-positive eigenvalue as an operator in $L^2_{rad}(\mathbb{R}^d)$ (see Lemma B.4). Thus, we have proved the claim (B.12).

Now, we are in a position to prove the claim (i). It follows from (B.1) and (2.9) that there exists $\omega_0 > 0$ such that for any $\omega \in (0, \omega_0)$ and any $g \in H^1(\mathbb{R}^d)$,

$$\langle L_{\omega,+}g,g\rangle_{H^{-1},H^{1}} \lesssim \left(\omega + 1 + \omega^{\frac{p-1}{p+1}(1-s_{p})} \|U\|_{L_{p+1}}^{p-1} + \omega^{\frac{4}{(d-2)(p-1)}-1} \|U\|_{L_{2^{*}}}^{2^{*}-2}\right) \|g\|_{H^{1}}^{2}.$$
(B.20)

Thus, it suffices to show that

$$\inf_{\substack{g \in H_{rad}^{1}(\mathbb{R}^{d},\mathbb{R}) \\ \|g\|_{H^{1}}=1 \\ (g,f_{2})_{L^{2}_{real}}=0}} \langle L_{\omega,+}g, g \rangle_{H^{-1},H^{1}} > 0, \tag{B.21}$$

where $H^1_{rad}(\mathbb{R}^d, \mathbb{R})$ denotes the space of radial real-valued functions in $H^1(\mathbb{R}^d)$. Suppose for a contradiction that (B.21) failed. Then, we could take a sequence $\{g_n\}$ of radial real-valued functions in $H^1(\mathbb{R}^d)$ such that

$$\|g_n\|_{H^1} \equiv 1, \tag{B.22}$$

$$(g_n, f_2)_{L^2_{real}} \equiv 0, \tag{B.23}$$

$$\lim_{n \to \infty} \langle L_{\omega,+} g_n, g_n \rangle_{H^{-1}, H^1} = 0.$$
(B.24)

Here, passing to some subsequence, we may assume that there exists a radial real-valued function $g_{\infty} \in H^1(\mathbb{R}^d)$ such that

$$\lim_{n \to \infty} g_n = g_{\infty} \quad \text{weakly in } H^1(\mathbb{R}^d).$$
(B.25)

If $g_{\infty} = 0$, then we see from (B.24) and (B.25) that for any $\omega \in (0, 1)$,

$$0 = \lim_{n \to \infty} \langle L_{\omega,+}g_n, g_n \rangle_{H^{-1}, H^1} = \lim_{n \to \infty} \left(L_{\omega,+}g_n, g_n \right)_{L^2_{real}}$$

$$= \lim_{n \to \infty} \left\{ \omega \int_{\mathbb{R}^d} |g_n|^2 + \int_{\mathbb{R}^d} |\nabla g_n|^2 - p \int_{\mathbb{R}^d} \Phi^{p-1}_{\omega} |g_n|^2 - (2^* - 1) \int_{\mathbb{R}^d} \Phi^{\frac{4}{d-2}}_{\omega} |g_n|^2 \right\}$$

$$\geq \omega - \lim_{n \to \infty} \left\{ p \int_{\mathbb{R}^d} \Phi^{p-1}_{\omega} |g_n|^2 + (2^* - 1) \int_{\mathbb{R}^d} \Phi^{\frac{4}{d-2}}_{\omega} |g_n|^2 \right\}$$

$$= \omega,$$

(B.26)

which is a contradiction. Thus, we have found that g_{∞} is nontrivial. Then, it follows from (B.24), the lower semicontinuity and (B.12) that

$$0 = \lim_{n \to \infty} \langle L_{\omega,+}g_n, g_n \rangle_{H^{-1}, H^1} \ge \langle L_{\omega,+}g_\infty, g_\infty \rangle_{H^{-1}, H^1} > 0.$$
(B.27)

This absurd conclusion comes from the hypothesis that (B.21) failed. Thus, we have prove the claim (i).

Next, we prove the claim (ii). Let $g \in H^1(\mathbb{R}^d)$ be a non-trivial, real-valued, radial function with $(g, \partial_\omega \Phi_\omega)_{L^2_{real}} = 0$. We write g in the form

$$g = a\Phi_{\omega} + h, \tag{B.28}$$

where h satisfies $(h, \Phi_{\omega})_{L^2_{real}} = 0$. Note that the condition $(g, \partial_{\omega} \Phi_{\omega})_{L^2_{real}} = 0$ implies that

$$a = -\frac{\left(h, \partial_{\omega} \Phi_{\omega}\right)_{L^{2}_{real}}}{\left(\Phi_{\omega}, \partial_{\omega} \Phi_{\omega}\right)_{L^{2}_{real}}}.$$
(B.29)

Hence, we find from $(h, \Phi_{\omega})_{L^2_{real}} = 0$ that

$$\begin{split} \|g\|_{H^{1}}^{2} &= (g,g)_{L^{2}_{real}} + (\nabla g,\nabla g)_{L^{2}_{real}} \\ &= a^{2} \|\Phi_{\omega}\|_{H^{1}}^{2} + \|h\|_{H^{1}}^{2} - 2a(\nabla\Phi_{\omega},\nabla h)_{L^{2}_{real}} \\ &\leq \frac{\|\partial_{\omega}\Phi_{\omega}\|_{L^{2}}^{2}\|h\|_{L^{2}}^{2}}{\left|(\Phi_{\omega},\partial_{\omega}\Phi_{\omega})_{L^{2}_{real}}\right|^{2}} \|\Phi_{\omega}\|_{H^{1}}^{2} + \|h\|_{H^{1}}^{2} + \frac{2\|\partial_{\omega}\Phi_{\omega}\|_{L^{2}}\|h\|_{L^{2}}}{\left|(\Phi_{\omega},\partial_{\omega}\Phi_{\omega})_{L^{2}_{real}}\right|} \|\nabla\Phi_{\omega}\|_{L^{2}} \|\nabla h\|_{L^{2}} \\ &\lesssim \|h\|_{H^{1}}^{2}. \end{split}$$
(B.30)

Moreover, we see from $L_{\omega,-}\Phi_{\omega} = 0$ and Lemma B.1, that

$$\langle L_{\omega,-}g,g \rangle_{H^{-1},H^1} = \langle L_{\omega,-}(a\Phi_\omega + h), a\Phi_\omega + h \rangle_{H^{-1},H^1}$$

$$= \langle L_{\omega,-}h,h \rangle_{H^{-1},H^1} \gtrsim \|h\|_{H^1}^2.$$
(B.31)

Putting (B.30) and (B.31) together, we obtain

$$||g||_{H^1}^2 \lesssim \langle L_{\omega,-}g,g \rangle_{H^{-1},H^1}.$$
 (B.32)

The opposite relation follows from (B.1). Hence, we have completed the proof.

C Inequalities for the radial functions

We see from the fundamental theorem of calculus and Hardy's inequality that for any $d \geq 3$ and any radial function $g \in H^1(\mathbb{R}^d)$,

$$\sup_{x \in \mathbb{R}^{d}} \{ |x|^{d-2} |g(x)|^{2} \} = \sup_{x \in \mathbb{R}^{d}} \int_{|x|}^{\infty} \frac{d}{dr} \left\{ -r^{d-2} |g(r)|^{2} \right\} dr$$

$$\lesssim \int_{0}^{\infty} r^{\frac{d-3}{2}} |g(r)| r^{\frac{d-1}{2}} |g'(r)| dr$$

$$\leq \left(\int_{0}^{\infty} |g(r)|^{2} r^{d-3} dr \right)^{\frac{1}{2}} \left(\int_{0}^{\infty} |g'(r)|^{2} r^{d-1} dr \right)^{\frac{1}{2}} \lesssim \|\nabla g\|_{L^{2}(\mathbb{R}^{d})}^{2}.$$
(C.1)

Similarly, we can verify that for any $d \geq 3$ and any radial function $g \in H^1(\mathbb{R}^d)$,

$$\sup_{x \in \mathbb{R}^d} \left\{ |x|^{d-1} |g(x)|^2 \right\} \lesssim \int_0^\infty r^{\frac{d-1}{2}} |g(r)| r^{\frac{d-1}{2}} |g'(r)| \, dr \lesssim \|g\|_{L^2(\mathbb{R}^d)} \|\nabla g\|_{L^2(\mathbb{R}^d)}.$$
(C.2)

D Small-data theory

We record standard small-data theories for (NLS) and (1.19) (see Lemma E.1 below and Corollary 3.9 in [22]):

Lemma D.1. (i) Assume $d \ge 3$ and $1 + \frac{4}{d} . Then, for any <math>A > 0$, there exists $\delta(A) > 0$ with the following property: for any $t_0 \in \mathbb{R}$ and any $\psi_0 \in H^1(\mathbb{R}^d)$ satisfying

$$\|\langle \nabla \rangle \psi_0\|_{L^2} \le A,\tag{D.1}$$

$$\|e^{i(t-t_0)\Delta}\psi_0\|_{W_{2+\frac{4}{d}}(\mathbb{R})\cap W_{2^*}(\mathbb{R})} \le \delta(A),$$
(D.2)

there exists a unique strong H^1 -solution ψ to (NLS) with $\psi(t_0) = \psi_0$. Furthermore, the maximal existence-interval of ψ is the whole of \mathbb{R} , and ψ satisfies

$$\|\langle \nabla \rangle \psi\|_{St(\mathbb{R})} \lesssim \|\langle \nabla \rangle \psi_0\|_{L^2}. \tag{D.3}$$

(ii) Assume $d \geq 3$. Then, there exists $\delta_0 > 0$ with the following property: for any $t_0 \in \mathbb{R}$ and any $\psi_0 \in \dot{H}^1(\mathbb{R}^d)$ satisfying

$$\|\nabla\psi_0\|_{L^2} \le \delta_0,\tag{D.4}$$

there exists a unique strong \dot{H}^1 -solution ψ to (1.19) with $\psi(t_0) = \psi_0$. Furthermore, the maximal existence-interval of ψ is the whole of \mathbb{R} and

$$\|\nabla\psi\|_{St(\mathbb{R})} \lesssim \|\nabla\psi_0\|_{L^2}.$$
 (D.5)

We can control the norm of the full Strichartz space by a few particular norms:

Lemma D.2. Assume $d \ge 3$, and let $A_1, A_2, A_3 > 0$. Then, there exists a constant $C(A_1, A_2, A_3) > 0$ with the following property:

(i) for any interval I and any space-time function u satisfying

$$|u||_{L^{\infty}(I,H^1)} \le A_1,$$
 (D.6)

$$\|u\|_{W_{p+1}(I)\cap W_{2^*}(I)} \le A_2,\tag{D.7}$$

$$\left\| \langle \nabla \rangle e[u] \right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I)} \le A_3, \tag{D.8}$$

we have

$$\|\langle \nabla \rangle u\|_{St(I)} \le C(A_1, A_2, A_3). \tag{D.9}$$

(ii) for any interval I and any space-time function u satisfying

$$\|\nabla u\|_{L^{\infty}(I,L^2)} \le A_1,$$
 (D.10)

$$\|u\|_{W_{2^*}(I)} \le A_2,\tag{D.11}$$

$$\left\|\nabla e^{\ddagger}[u]\right\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I)} \le A_3,$$
 (D.12)

we have

$$\|\nabla u\|_{St(I)} \le C(A_1, A_2, A_3).$$
 (D.13)

Proof of Lemma D.2. See the proof of Lemma 4.1 in [3].

E Long-time perturbation theory

In this section, we give the long-time perturbation theory for a general nonlinear Schrödinger equation including (NLS) with $1 + \frac{4}{d} \le p < 2^* - 1$:

$$i\frac{\partial\psi}{\partial t} + \Delta\psi + f(\psi) = 0, \qquad (E.1)$$

where $f : \mathbb{C} \to \mathbb{C}$ is a continuously differentiable function in the real-sense.

Fix an even, smooth cut-off function χ defined on \mathbb{R} such that $\chi(r) = 1$ if $0 \le r \le 1$ and $\chi(r) = 0$ if $r \ge 2$, and define

$$f_{\leq 1}(z) := \chi(|z|) f(z),$$
 (E.2)

$$f_{>1}(z) := f(z) - f_{\le 1}(z).$$
 (E.3)

Then, our assumptions about the nonlinearity f are the followings:

1.

$$f(0) = \frac{\partial f}{\partial z}(0) = \frac{\partial f}{\partial \bar{z}}(0) = 0.$$
(E.4)

2. There exists a constant $C_{\leq 1} > 0$ such that for any $z_1, z_2 \in \mathbb{C}$,

$$\left| \frac{\partial f_{\leq 1}}{\partial z}(z_1) - \frac{\partial f_{\leq 1}}{\partial z}(z_2) \right| + \left| \frac{\partial f_{\leq 1}}{\partial \bar{z}}(z_1) - \frac{\partial f_{\leq 1}}{\partial \bar{z}}(z_2) \right| \\
\leq C_{\leq 1} \begin{cases} |z_1 - z_2| \min\left\{ 10, (|z_1| + |z_2|)^{\frac{4}{d} - 1} \right\} & \text{if } d = 3, \\ |z_1 - z_2|^{\frac{4}{d}} & \text{if } d \geq 4. \end{cases}$$
(E.5)

3. There exists a constant $C_{>1} > 0$ such that for any $z_1, z_2 \in \mathbb{C}$,

$$\begin{aligned} \left| \frac{\partial f_{>1}}{\partial z}(z_1) - \frac{\partial f_{>1}}{\partial z}(z_2) \right| + \left| \frac{\partial f_{>1}}{\partial \bar{z}}(z_1) - \frac{\partial f_{>1}}{\partial \bar{z}}(z_2) \right| \\ \leq C_{>1} \begin{cases} |z_1 - z_2| (|z_1| + |z_2|)^{\frac{4}{d-2} - 1} & \text{if } 3 \le d \le 5, \\ |z_1 - z_2|^{\frac{4}{d-2}} & \text{if } d \ge 6. \end{cases} \end{aligned}$$
(E.6)

Throughout this section, we allow the implicit constants to depend on $C_{\leq 1}$ and $C_{>1}$, so that the assumptions (E.4) through (E.6) imply that

$$|f_{\leq 1}(z)| \lesssim |z|^{1+\frac{4}{d}},$$
 (E.7)

$$|f_{>1}(z)| \lesssim |z|^{1+\frac{4}{d-2}},$$
 (E.8)

$$\left|\frac{\partial f_{\leq 1}}{\partial z}(z)\right| + \left|\frac{\partial f_{\leq 1}}{\partial \bar{z}}(z)\right| \lesssim |z|^{\frac{4}{d}},\tag{E.9}$$

$$\left|\frac{\partial f_{>1}}{\partial z}(z)\right| + \left|\frac{\partial f_{>1}}{\partial \bar{z}}(z)\right| \lesssim |z|^{\frac{4}{d-2}}$$
(E.10)

for any $z \in \mathbb{C}$.

Now, we state the long-time perturbation theory for the equation (E.1):

Lemma E.1. Let $d \ge 3$, I an interval, $\psi \in C(I, H^1(\mathbb{R}^d))$ a solution to (E.1), and let u be a function in $C(I, H^1(\mathbb{R}^d))$. Assume that

$$\|u\|_{L^{\infty}_{t}H^{1}_{x}(I)} \le A, \tag{E.11}$$

$$\|u\|_{V_{2+\frac{4}{d}}(I)\cap W_{2^*}(I)} \le B,\tag{E.12}$$

$$\|\psi(t_0) - u(t_0)\|_{H^1} \le \gamma$$
(E.13)

for some $t_0 \in I$ and some constants A > 0, B > 0 and $\gamma > 0$. Then, there exist constants $\delta_0(A, B, \gamma) > 0$ and $C(A, B, \gamma) > 0$ such that if

$$\|e^{i(t-t_0)\Delta}\{\psi(t_0) - u(t_0)\}\|_{V_{2+\frac{4}{d}}(I) \cap W_{2^*}(I)} \le \delta,$$
(E.14)

$$\|\langle \nabla \rangle e(u)\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I)} \le \delta \tag{E.15}$$

for some $0 < \delta < \delta_0(A, B, \gamma)$, then we have

$$\|\psi - u\|_{V_{2+\frac{4}{d}}(I) \cap W_{2^*}(I)} \le C(A, B, \gamma) \delta^{\frac{1}{d(d+2)}},$$
(E.16)

$$\|\langle \nabla \rangle \{\psi - u\}\|_{St(I)} \le C(A, B, \gamma)\gamma, \tag{E.17}$$

$$|\langle \nabla \rangle \psi \|_{St(I)} \le C(A, B, \gamma). \tag{E.18}$$

Here, the constant $C(A, B, \gamma)$ is non-decreasing with respect to each of the parameters A, B and γ .

We can prove Lemma E.1 in a way similar to the proof of Theorem 3.8 in [22]. The main difference of the proof is that we need another exotic Strichartz estimate. For example, the following exotic Strichartz norms work well:

$$\|u\|_{ES(I)} := \|\langle \nabla \rangle^{\frac{4}{d+2}} u\|_{L_t^{\frac{d(d+2)}{2(d-2)}} L_x^{\frac{2d^2(d+2)}{d^3-4d+16}}(I)} + \|u\|_{L_t^{\frac{d+2}{2}} L_x^{\frac{2d(d+2)}{(d-2)(d+4)}}(I)},$$
(E.19)

$$\|u\|_{ES^*(I)} := \|\langle \nabla \rangle^{\frac{4}{d+2}} u\|_{L_t^{\frac{d}{2}} L_x^{\frac{2d^2(d+2)}{d^3+4d^2+4d-16}}(I)} + \|u\|_{L_t^{\frac{d(d+2)}{2(d+4)}} L_x^{\frac{2d^2(d+2)}{(d+4)^2(d-2)}}(I)},$$
(E.20)

where I is an interval. In particular, we have the following exotic Strichartz estimate.

Lemma E.2 (Exotic Strichartz estimate). Let $d \ge 3$. Then, we have

$$\|u - e^{i(t-t_0)\Delta}u(t_0)\|_{ES(I)} \lesssim \|i\frac{\partial u}{\partial t} + \Delta u - g\|_{ES^*(I)} + \|\langle \nabla \rangle g\|_{L^{\frac{2(d+2)}{d+4}}_{t,x}(I)}$$
(E.21)

for any interval I, any $t_0 \in I$, and any functions u and g defined on $\mathbb{R}^d \times I$.

Moreover, we can derive an analogue of Lemma 3.11 in [22]:

Lemma E.3. Let $d \geq 3$. Then, we have

$$\|u\|_{ES(I)} \lesssim \|u\|_{L^{\frac{1}{d+2}}_{t,x}(I)} \|\nabla u\|_{St(I)}^{\frac{d+1}{d+2}} + \|u\|_{L^{\frac{16(d-2)}{d^2(d+2)}}_{t,x}(I)} \|\langle \nabla \rangle u\|_{St(I)}^{\frac{d^3+2d^2-16d+32}{d^2(d+2)}}$$
(E.22)

for any interval I and any function u on $\mathbb{R}^d \times I$. In particular, we have

$$\|u\|_{ES(I)} \lesssim \|\langle \nabla \rangle u\|_{St(I)}.$$
(E.23)

Moreover, there exist constants $\theta_0, \theta_1 \in (0, 1]$ depending only on d such that

$$\|u\|_{L^{\frac{2(d+2)}{d}}_{t,x}(I)} \le \|u\|_{St(I)}^{1-\theta_0} \|u\|_{ES(I)}^{\theta_0},$$
(E.24)

$$\|u\|_{L^{\frac{2(d+2)}{d-2}}_{t,x}(I)} \le \|\nabla u\|_{St(I)}^{1-\theta_1} \|u\|_{ES(I)}^{\theta_1}$$
(E.25)

for any interval I and any function u on $\mathbb{R}^d \times I$.

Symbols	Description or equation number
\mathcal{M}	(1.2)
$\mathcal{H}, \mathcal{H}^{\dagger}, \mathcal{H}^{\ddagger},$	(1.3), (1.41), (1.32)
$\mathcal{S}_\omega, \mathcal{S}_\omega^\dagger, \mathcal{S}_\omega^j$	(1.8), (1.46), (8.211)
$\mathcal{K}, \widetilde{K}_\omega, \mathcal{K}^\dagger, \mathcal{K}^\ddagger$	(1.5), (1.57), (1.43), (1.31)
$\mathcal{I}_{\omega},\widetilde{\mathcal{I}}_{\omega},\mathcal{I}^{\ddagger},\mathcal{I}_{\omega}^{j}$	(1.11), (1.58), (1.30), (8.211)
\mathcal{J}_ω	(1.12)
Φ_ω, U	positive, radial ground states of (1.7) and (1.22)
W	(1.26)
$m_\omega,\sigma, u_\omega$	(1.9), (1.27), (4.17)
$\widetilde{PW}_{\omega}, PW_{\omega}, PW_{\omega,+}, PW_{\omega,-}$	(1.16), (1.48), (1.50), (1.51)
$PW^{\dagger}, PW^{\dagger}_{\omega}, PW^{\dagger, \varepsilon}$	(1.40), (1.44), (1.47)
PW^{\ddagger}_+	(1.37)
T_ω	(1.24)
s_p	(1.42)
$\mathcal{L}_{\omega}, L_{\omega,+}, L_{\omega,-}$	(4.2), (4.4), (4.5)
$L^{\dagger}_+, L^{\dagger}$	(1.55), (1.56)
$\widetilde{L}_{\omega,+},\widetilde{L}_{\omega,-}$	(1.59), (1.60)
μ	positive eigen-value of $-i\mathcal{L}_{\omega}$
\mathfrak{U}_{\pm}	eigen-function of $-i\mathcal{L}_{\omega}$ corresponding to $\pm\mu$
$\lambda_{\pm}(t)$	(4.50)
$\Gamma(t)$	(4.65)
$\lambda_1(t), \lambda_2(t)$	(4.88)
$d_\omega(\psi(t)), \widetilde{d}_\omega(\psi(t))$	(4.85), (6.15)
$\delta_E, \delta_X, \delta_S, \delta_*$	(4.77), (5.24), (7.7), (??)
T_X	(5.12)
$\kappa_1(\delta)$	(6.20)
$A_{\omega}^{\varepsilon}, S_{\omega,R}^{\varepsilon}, S_{\omega,R,\pm}^{\varepsilon}$	(3.1), (8.1), (8.19)
$St(I), V_q(I), V(I), W_q(I), W(I), W^j(I)$	(1.61), (1.62), (8.73)
G_n^j,g_n^j	(8.44), (8.45)
$\sigma_n^j, \sigma_\infty^j$	(8.47), (8.56)
$x_n^j, t_n^j, \lambda_n^j$	(8.39), (8.40), (8.41)
\widetilde{u}^j, w_n^k	(??), (8.42)
$\widetilde{\psi}^j,\psi^j_n$	(8.72), (8.80)
ψ_n^{k-app}	(8.168)

Acknowledgements

T.A. is partially supported by the Grant-in-Aid for Young Scientists (B) # 25800077 of JSPS.

S. I. is partially supported by NSERC Discovery grant # 371637-2014.

H. K. is partially supported by the Grant-in-Aid for Young Scientists (B) # 00612277 of JSPS.

H. N. is partially supported by the Grant-in-Aid for Scientific Research (B) 23340030 of JSPS and the Grant-in-Aid for Challenging Exploratory Research 23654052 of JSPS.

References

- Akahori. T. and Nawa. H., Blowup and Scattering problems for the Nonlinear Schrödinger equations, Kyoto J. Math (2013), 629–672.
- [2] Akahori T., Ibrahim S., Kikuchi H. and Nawa H., Existence of a ground state and blow-up for a nonlinear Schrödinger equation with critical growth. Differential and Integral Equations 25 (2012), 383–402.
- [3] Akahori T., Ibrahim S., Kikuchi H. and Nawa H., Existence of a ground state and scattering for a nonlinear Schrödinger equation with critical growth. Selecta Mathematica 19 (2013), 545–609.
- [4] Berestycki, H., Cazenave, T., Instabilité des états stationnarires dans les équations de Schroödinger et de Klein-Gordon non linéaires. C.R. Acad. Sci. Paris Sér. I Math. 293 (1981), 489–492.
- [5] Berestycki, H. and Lions, P.-L., Nonlinear scalar field equations. I. Existence of a ground state. Arch. Rational Mech. Anal. 82 (1983), 313–345.
- [6] Brezis, H. and Lieb, E. H., Minimum action solutions of some vector field equations. Comm. Math. Phys. 96 (1984), 97–113.
- [7] Brezis, H. and Nirenberg, L., Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents. Comm. Pure Appl. Math. 36 (1983), 437–477.
- [8] Caffarelli, L., Gidas, B. and Spruck, J., Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth. Comm. Pure Appl. Math. 42 (1989), 271–297.
- [9] Chang, S.M., Gustafson, S., Nakanishi, K. and Tsai, T.P., Spectra of linearized operators for NLS solitary waves. SIAM J. Math. Anal. 39 (2007/08), 1070–1111.

- [10] Cheng, X., Miao, C. and Zhao, L., Global well-posedness and scattering for nonlinear Schrödinger equations with combined nonlinearities in the radial case. J. Differential Equations 261 (2016), 2881–2934.
- [11] Duyckaerts, T., Holmer, J. and Roudenko, S., Scattering for the non-radial 3D cubic nonlinear Schrödinger equation. Math. Res. Lett. 15 (2008), 1233–1250.
- [12] Gidas, B., Ni, W.M. and Nirenberg, L., Symmetry of positive solutions of nonlinear elliptic equations in \mathbb{R}^N , Math. Anal. Appl. (1981), 369–402.
- [13] Grillakis, M., Linearized instability for nonlinear Schrödinger and Klein-Gordon equations. Comm. Pure Appl. Math. 41 (1988), 747–774.
- [14] Holmer, J. and Roudenko, S., A sharp condition for scattering of the radial 3D cubic nonlinear Schrödinger equation, Comm. Math. Phys. 282 (2008), 435–467.
- [15] Ibrahim, S., Masmoudi, N. and Nakanishi, K., Scattering threshold for the focusing nonlinear Klein-Gordon equation. Anal. PDE 4 (2011), 405–460.
- [16] Fukuizumi R., Remarks on the stable standing waves for nonlinear Schrödinger equations with double power nonlinearity. Adv. Math. Sci. Appl. 13 (2003), 549–564.
- [17] Kabeya, Y. and Tanaka, K., Uniqueness of positive radial solutions of semilinear elliptic equations in RN and Sere's non-degeneracy condition. Comm. Partial Differential Equations 24 (1999), 563–598.
- [18] Kenig, C.E. and Merle, F., Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case. Invent. Math. 166 (2006), 645–675.
- [19] Keraani, S., On the defect of compactness for the Strichartz estimates of the Schrödinger equations. J. Differential Equations 175 (2001), 353–392.
- [20] Killip R., Oh, T., Pocovnicu O. and Visan. M, Solitons and scattering for the cubicquintic nonlinear Schrödinger equation on ℝ³, arXiv:1409.6734.
- [21] Killip R. and Visan M., The focusing energy-critical nonlinear Schrödinger equation in dimensions five and higher. Amer. J. Math. 132 (2010), 361–424.
- [22] Killip R. and Visan M., Nonlinear Schrödinger equations at critical regularity, Clay Lecture Notes.
- [23] Lieb, E.H. and Loss, M., ANALYSIS, second edition, American Mathematical Society (2001).

- [24] Nakanishi, K. and Schlag, W., Global dynamics above the ground state energy for the focusing nonlinear Klein-Gordon equation. Journal of Differential Equations 250 (2011), 2299–2333.
- [25] Nakanishi, K. and Schlag, W., Global dynamics above the ground state energy for the cubic NLS equation in 3D. Calc. Var. and PDE 44 (2012), 1–45.
- [26] Ni, W.M. and Takagi, I., Locating the peaks of least-energy solutions to a semilinear Neumann problem. Duke Math. J. 70 (1993), 247–281.
- [27] Shatah, J. and Strauss, W., Instability of nonlinear bound states. Comm. Math. Phys. 100 (1985), 173–190.
- [28] Tao, T. and Visan, M., Stability of energy-critical nonlinear Schrödinger equations in high dimensions, Electron. J. Differential Equations 118 (2005), 1–28.
- [29] Tao, T., Visan, M. and Zhang, X., The nonlinear Schrödinger equation with combined power-type nonlinearities, Comm. Partial Differential Equations 32 (2007), 1281–1343.
- [30] Weinstein, M., Nonlinear Schrödinger equations and sharp interpolation estimates. Comm. Math. Phys. 87 (1982/83), 567–576.
- [31] Weinstein, M., Modulational stability of ground states of nonlinear Schrödinger equations. SIAM Math. ANAL. 16 (1985), 472–491.
- [32] Zhang, J. and Zou, W., The critical case for a Berestycki-Lions theorem, Science China Math. 57 (2014), 541–554.