A Statistical Model for Motifs Detection

Hamid Javadi[∗] and Andrea Montanari†

August 22, 2018

Abstract

We consider a statistical model for the problem of finding subgraphs with specified topology in an otherwise random graph. This task plays an important role in the analysis of social and biological networks. In these types of networks, small subgraphs with a specific structure have important functional roles, and they are referred to as 'motifs.'

Within this model, one or multiple copies of a subgraph is added ('planted') in an Erdős-Renyi random graph with n vertices and edge probability q_0 . We ask whether the resulting graph can be distinguished reliably from a pure Erd˝os-Renyi random graph, and we present two types of result. First we investigate the question from a purely statistical perspective, and ask whether there is any test that can distinguish between the two graph models. We provide necessary and sufficient conditions that are essentially tight for small enough subgraphs.

Next we study two polynomial-time algorithms for solving the same problem: a spectral algorithm, and a semidefinite programming (SDP) relaxation. For the spectral algorithm, we establish sufficient conditions under which it distinguishes the two graph models with high probability. Under the same conditions the spectral algorithm indeed identifies the hidden subgraph.

The spectral algorithm is substantially sub-optimal with respect to the optimal test. We show that a similar gap is present for the more sophisticated SDP approach.

1 Introduction

'Motifs' play a key role in the analysis of social and biological networks. Quoting from an influential paper in this area $[MSOI^+02]$ $[MSOI^+02]$, the term 'motif' broadly refers to

"patterns of interconnections occurring in complex networks at numbers that are significantly higher than those in randomized networks."

For instance, the authors of $[MSOI^+02]$ $[MSOI^+02]$ considered directed graph representations of various types of data: gene regulation networks, neural circuits, food webs, the world wide web, electronic circuits. They identified a number of small subgraphs that are found in atypically large numbers in such networks, and provided interpretations of their functional role. The analysis of motifs in large biological networks was pursued in a number of publications, see e.g. [\[KIMA04,](#page-14-0) [YLSK](#page-15-1)+04, [KA05,](#page-14-1) [SSR](#page-15-2)+05, [Alo07\]](#page-12-0).

[∗]Department of Electrical Engineering, Stanford University

[†]Department of Electrical Engineering and Statistics, Stanford University

The analysis of subgraph frequencies has an even longer history within sociology, in part because sociological theories are predictive of specific subgraph occurrences. We refer to [\[Gra73\]](#page-14-2) for early insights, and to [\[WF94,](#page-15-3) [EK10\]](#page-13-0) for recent reviews of this research area.

This paper studies a statistical model for the motif detection problem. In order to provide a formal statement of the problem, denote by \mathcal{G}_n the space of graphs over vertex set $V_n = [n]$.

Definition 1.1. We say that the two (sequences of) probability laws $\mathbb{P}_{0,n}$, $\mathbb{P}_{1,n}$ over \mathcal{G}_n are strongly distinguishable if there exists a sequence of functions (a 'test') $T : \mathcal{G}_n \to \{0,1\}$ such that

$$
\limsup_{n \to \infty} \mathbb{P}_{0,n}(T(G_n) = 1) = \limsup_{n \to \infty} \mathbb{P}_{1,n}(T(G_n) = 0) = 0.
$$

We say that they are weakly distinguishable if there exists $T: \mathcal{G}_n \to \{0,1\}$ such that

$$
\limsup_{n \to \infty} \left[\mathbb{P}_{0,n}(T(G_n) = 1) + \mathbb{P}_{1,n}(T(G_n) = 0) \right] < 1.
$$

We say that they are polynomial-time weakly (or strongly) distinguishable if a test exists that achieves the above and can be computed by a polynomial-time algorithm.

Throughout the paper, $\mathbb{P}_{0,n}$ will correspond to a standard Erdős-Renyi random graph, while $\mathbb{P}_{1,n}$ will be an Erdős-Renyi random graph with planted copies of a small graph H_n . Namely, fix $0 \le q_0 \le 1$, and let $H_n = (V(H_n), E(H_n))$ be a sequence of graphs, indexed by n. Let us emphasize that H_n is a non-random graph on $v(H_n) \equiv |V(H_n)| \ll n$ vertices. Given a graph F, we denote by $\mathcal{L}(F,n)$ the set of labelings of the vertices of F taking values in [n], i.e.

$$
\mathcal{L}(F,n) \equiv \left\{ \varphi : V(F) \to [n] \text{ s.t. } \varphi(i) \neq \varphi(j) \ \forall i \neq j \right\}. \tag{1.1}
$$

(In particular $|\mathcal{L}(H_n, n)| = n!/(n - v(H_n))!$) For an edge $e = (i, j) \in E(H_n)$, we let $\varphi(e)$ be the unordered pair $(\varphi(i), \varphi(j))$, and hence $\varphi(E(H_n)) = {\varphi(e) : e \in E(H_n)}$.

We then let $\mathbb{P}_{0,n}((i,j) \in E) = q_0$ independently for all pairs (i,j) . On the other hand

$$
\mathbb{P}_{1,n}(\cdot) = \frac{1}{|\mathcal{L}(H_n, n)|} \sum_{\varphi \in \mathcal{L}(H_n, n)} \mathbb{P}_{1,n}(\cdot | \varphi), \qquad (1.2)
$$

$$
\mathbb{P}_{1,n}((i,j)\in E|\varphi) = \begin{cases} 1 & \text{if } (i,j)\in\varphi(E(H_n)), \\ q_0 & \text{otherwise.} \end{cases}
$$
 (1.3)

Here it is understood that edges are independent conditional on the labeling φ . In Section [4](#page-10-0) we will generalize this defintion by considering the case in which a large number of identical copies of H_n is planted.

Whenever the two laws $\mathbb{P}_{0,n}$ and $\mathbb{P}_{1,n}$ are distinguishable, we will also say that the motif H_n is detectable. We will consider the following fundamental questions:

- Under which conditions on q_0 , $\{H_n\}_{n\in\mathbb{N}}$ the are two laws $\mathbb{P}_{0,n}$, $\mathbb{P}_{1,n}$ weakly distinguishable? Under which conditions are they strongly distinguishable?
- Assuming the conditions for distinguishability of $\mathbb{P}_{0,n}$, $\mathbb{P}_{1,n}$ are satisfied, under which conditions there exists a polynomial-time computable test $T(\cdot)$ that distinguishes $\mathbb{P}_{0,n}$ from $\mathbb{P}_{1,n}$?

• What features of the motif H_n controls it detectability?

The first two questions have attracted a substantial amount of work since the nineties for special cases such as planted cliques or dense subgraphs [\[Jer92,](#page-14-3) [FK00,](#page-13-1) [FPK01a\]](#page-13-2). A brief overview of this line of work is presented in Section [1.1.](#page-3-0) However, applied studies investigate a much broader collection of motifs than just cliques $[MSOI⁺02, Alo07, WF94, EK10]$ and $-in$ fact– cliques are rarely the most interesting motif from a scientific perspective.

It is a priori unclear whether intuitions developed on the planted clique problem apply to general motif detection. Quantitative implications, namely tight necessary and sufficient conditions for detectability of an arbitrary motif H_n are even less clear. The present paper builds upon some key insights that were developed in the earlier literature, to obtain a broader picture that is potentially applicable to motifs of scientific interest.

In the rest of the paper, we present the following results:

1. Section [2](#page-5-0) presents upper and lower bounds on the statistical threshold for motif detection. The bounds match for sufficiently small subgraphs (in particular for $v(H_n) = n^{o(1)}$) yielding a sharp characterization in that regime. The key feature controlling statistical detection turns out to be the maximum graph density $d(H_n)$ (see Section [2](#page-5-0) for a definition).

Determining sharp detectability conditions for larger motifs, or for background density q_0 depending on n , remains an open problem.

2. In order to explore the computational limits of motif detection, Section [3.1](#page-6-0) analyzes a simple spectral algorithm that computes the leading eigenvector of the centered adjacency matrix of G. The key feature controlling the behavior of this algorithm is the leading eigenvalue of the adjacency matrix of H_n . We prove that the spectral approach succeeds with high probability if this eigenvalue is larger than $C\sqrt{n}$.

We also show that, in the same regime, the spectral algorithms can be augmented with a combinatorial step that identifies the planted subgraph. We prove that this step is successful under a certain 'balancedness' condition on the motif H_n .

3. Section [3.2](#page-9-0) apply a semidefinite programming (SDP) relaxation of the quadratic assignment problem to motif detection. We prove that the SDP approach is not successful unless –again– problem to moth detection. We prove that the SDT approach is not successive unless $-\text{again}$ -
the leading eigenvalue of H_n is of order \sqrt{n} . In other words, a similarly large gap between statistical and algorithmic thresholds exists for SDP as for spectral methods.

As is often the case in proving negative results for SDP relaxation, our analysis relies on a careful construction of a primal witness. This construction is of potential interest for other applications of quadratic assignment.

4. Finally, Section [4](#page-10-0) considers the more general case in which m_n copies of the motif H_n are planted in the same graph. We obtain upper and lower bounds on the statistical threshold. These bounds match when the motif size is small enough or the number of copies is small enough.

These results suggest that motifs detection might be computationally hard in regimes in which it is statistically feasible. As discussed in the next section, this phenomenon has been already extensively investigated for the planted clique problem. More interestingly, our analysis suggests that the two thresholds are controlled by different features of the motif H_n . While the statistical threshold depends on the maximum density of H_n , the spectral threshold is related to its principal eigenvalue.

1.1 Related work

Statistical models for motif detection have been studied so far only for specific cases. An important line of work within theoretical computer science has focused on the planted clique problem, which corresponds to the case $H_n = K_{k(n)}$, the complete graph over $k(n)$ vertices. It is a classical result that the planted model is distinguishable from a pure Erdős-Renyi random graph, for $q_0 = 1/2$, provided $k(n) \geq 2(1+\varepsilon) \log_2 n$ (with ε arbitrarily small), while it is undistinguishable for $k(n) \leq$ $2(1 - \varepsilon) \log_2 n$, see e.g. [\[GM75\]](#page-14-4). On the other hand, approximating the size of the largest clique in a graph is hard in a worst case setting, even within a factor $n^{1-\varepsilon}$ [Hås99, [Kho01\]](#page-14-6). Starting with Jerrum's seminal work, a number of authors analyzed broad classes of algorithms for the statistical model. A short list includes Monte Carlo Markov Chain [\[Jer92\]](#page-14-3), spectral algorithms [\[AKS98\]](#page-12-1), SDP relaxations within the Lovasz-Schrijver hierarchy [\[FK00\]](#page-13-1), statistical query models [\[FGR](#page-13-3)+13], message passing algorithms [\[DM14\]](#page-13-4), and, most recently, SDP relaxations within the Sum-Of-Squares hierarchy [\[MPW15,](#page-15-4) [DM15,](#page-13-5) [RS15,](#page-15-5) [HKP15,](#page-14-7) [BHK](#page-13-6)⁺16].

While these works provide precious insights into the tradeoff between statistical and computational limits for graph estimation, they cannot be applied directly to general motif detection. In particular, they do not clarify what features of the motif H_n are relevant for detection.

The planted clique problem is arguably the most studied statistical problem presenting a large gap between optimal statistical estimation and computationally efficient algorithms. In fact, several works *assume* hardness of planted clique to prove hardness of other statistical estimation problems, see e.g. [\[BR13\]](#page-13-7). Similar in spirit is the recent work in [\[DLSS14\]](#page-13-8), that instead assumes hardness of refuting random satisfiability formulas, as well as [\[ADBL11\]](#page-12-2).

A generalization of the maximum-clique problem is provided by the densest k-subgraph problem. Given a graph G and an integer k, this requires to find a subset of vertices of size k that contains the largest number of edges [\[FPK01a\]](#page-13-2). The best polynomial-time algorithm guarantees an $O(n^{1/4+\epsilon})$ approximation ratio [\[BCC](#page-12-3)+10]. The recent paper [\[Man17\]](#page-14-8) proves that an approximation ratio $n^{1/(\log \log n)^{\Omega(1)}}$ cannot be achieved under the exponential time hypothesis (see also [\[Fei02,](#page-13-9) [Kho06,](#page-14-9) [AAM](#page-12-4)+11] for other inapproximability results). Semidefinite programming relaxations were studied, among others in [\[FPK01b,](#page-14-10) [BCV](#page-13-10)+12].

Once more, these results do not apply directly to the motif detection problem. In particular, it is not clear –in general– that searching for a dense subgraph is necessarily the best approach to detect a given motif H_n . Further, the focus of this line of work is on worst-case approximation guarantees, rather than on statistical thresholds.

Closer to the scope of the present paper is some of the recent work on 'community detection'. A random graph $G = (V, E)$ over $|V| = n$ vertices is generated by selecting a subset $S \subseteq V$ of $|S| = k$ vertices uniformly at random. Edges are conditionally independent given S. Two vertices i, j are connected by an edge with probability q_1 if $\{i, j\} \subseteq S$ and probability $q_0 < q_1$ otherwise. Statistical and computational thresholds for detection and estimation of the set S were studied by a number of authors [\[VAC](#page-15-6)+15, [ACV14,](#page-12-5) [HWX15,](#page-14-11) [Mon15,](#page-14-12) [HWX17\]](#page-14-13). Again, these works focuses uniquely on the density of the planted subgraph, and do not model its structure.

Following the original work on motifs in biological networks $[MSOI^+02]$ $[MSOI^+02]$, several papers developed algorithms to sample uniformly random networks with certain given characteristics, e.g. with a given degree sequence $[MKI^+03, TS07, BD11]$ $[MKI^+03, TS07, BD11]$ $[MKI^+03, TS07, BD11]$ $[MKI^+03, TS07, BD11]$. Uniform samples can be used to assess the significance level of specific subgraph counts in a real network under consideration. Let, for instance, $N_H(G)$ denote the number of copies of a certain small graph H in G. If in a real network of interest we find $N_H(G) = t$, the probability $\mathbb{P}_{0,n}(N_{H_n}(G_n) \geq t)$ is used as significance level for this discovery. Let us mention two key differences with respect to problem considered in the present paper. First, we focus on conditions under which the laws $\mathbb{P}_{0,n}$ and $\mathbb{P}_{1,n}$ are strongly distinguishable. For instance, in the case of a single planted subgraph, under the null model $N_{H_n}(G_n) = 0$ with very high probability. Hence, Monte Carlo is not effective in accurately computing p -values in this regime.

Second, the work of [\[MKI](#page-14-14)+03, [TS07,](#page-15-7) [BD11\]](#page-13-11) implicitly assumes that the subgraph H_n has bounded size so that $N_{H_n}(G_n)$ can be computed in time $n^{v(H_n)}$ by exhaustive search. Here we consider instead large subgraphs H_n , and address the computational challenge of testing for H_n in polynomial time. Let us emphasize that, while we typically assume H_n to have diverging size, in practice it is impossible to perform exhaustive search already for quite small subgraphs. For instance, if $n = 10^5$, and $v(H_n) = 6$, exhaustive search requires of the order of $10^{30}/6! \ge 10^{27}$ operations.

1.2 Notations

Given $n \in \mathbb{Z}$, we let $[n] = \{1, 2, \ldots, n\}$ denote the set of first n integers. We write $|S|$ for the cardinality of a set S. We denote by $(n)_k = n!/(n-k)!$ the incomplete factorial.

Throughout this paper, we will use lowercase boldface (e.g. $\mathbf{v} = (v_1, \ldots, v_n), \mathbf{x} = (x_1, \ldots, x_n),$ etc.) to denote vectors and uppercase boldface (e.g. $\mathbf{A} = (A_{i,j})_{i,j \in [n]}, \mathbf{Y} = (Y_{i,j})_{i,j \in [n]},$ etc.) to denote matrices. For a vector $\mathbf{v} \in \mathbb{R}^n$ and a set $A \subseteq [n]$, we define $\mathbf{v}_A \in \mathbb{R}^n$ as $(v_A)_i = (v)_i$ for $i \in A$ and $(v_A)_i = 0$, otherwise. For a matrix Y we use $||Y||_2$ to denote its spectral norm. Given a square matrix $\boldsymbol{X} \in \mathbb{R}^{n \times n}$, we denote its trace by $\text{Tr}(\boldsymbol{X}) = \sum_{i=1}^{n} X_{i,i}$. Given a symmetric matrix M, we denote by $\lambda_1(M) \geq \lambda_2(M) \geq \cdots \geq \lambda_n(M)$ its ordered eigenvalues.

We denote by $\mathbf{1}_n = (1, 1, \ldots, 1) \in \mathbb{R}^n$ the all-ones vector, and by \mathbf{I}_n , $\mathbf{J}_n = \mathbf{1}_n \mathbf{1}_n^{\mathsf{T}} \in \mathbb{R}^{n \times n}$ the identity and all-ones matrices, respectively. For a matrix $\mathbf{A} \in \mathbb{R}^{m \times n}$, $\text{vec}(\mathbf{A}) \in \mathbb{R}^{mn}$ is the vector whose l'th entry is A_{ij} where $i-1$ and $j-1$ are the quotient and the remainder in dividing l by n, respectively. Also, $e_i \in \mathbb{R}^n$ denotes the *i*'th standard unit vector.

A simple graph is a pair $G = (V, E)$, where V is a vertex set and E is a set of unordered pairs $(i, j), i, j \in E$. We will write $V(G), E(G)$ whenever necessary to specify which graph we are referring to. Throughout, we will be focusing on finite graphs. We let $v(G) = |V(G)|$, $e(G) = |E(G)|$. For $v \in V(G)$, degree of node v is shown by deg(v). For $H \subseteq G, v \in V(G)$, the number of nodes in H which are connected to v is denoted by $\deg_H(v)$. We let \mathcal{G}_n be the set of graphs over vertex set $V=[n].$

We follow the standard Big-Oh notation. Given functions $f(n)$, $g(n)$, we write $f(n) = O(g(n))$ if there exists a constant C such that $f(n) \leq C g(n)$, $f(n) = \Omega(g(n))$ if there exists a constant C such that $f(n) \geq g(n)/C$, and $f(n) = \Theta(g(n))$ if $f(n) = O(g(n))$ and $f(n) = \Omega(g(n))$. Further $f(n) = o(q(n))$ if $f(n) \leq C q(n)$ for all $C > 0$ and n large enough, and $f(n) = \omega(q(n))$ if $f(n) > 0$ $C g(n)$ for all $C > 0$ and n large enough.

2 Statistical limits on hypothesis testing

In this section we address the first question stated in Section [1:](#page-0-0) under which conditions on q_0 , ${H_n}_{n\in\mathbb{N}}$ are the two laws $\mathbb{P}_{0,n}$, $\mathbb{P}_{1,n}$ distinguishable? We focus on the case of a single planted subgraph, $m_n = 1$, deferring the generalization to $m_n > 1$ to Section [4.](#page-10-0) We note that strong and weak distinguishability are equivalent to $\liminf_{n\to\infty} \|P_{0,n} - P_{1,n}\|_{TV} = 1$, and $\liminf_{n\to\infty} \|P_{0,n} - P_{1,n}\|_{TV}$ $\mathbb{P}_{1,n}||_{TV} > 0$, respectively.

Our results depend on the graph sequence ${H_n}_{n\in\mathbb{N}}$ through its maximum density $d(H_n)$. For a graph H , we define

$$
d(H) \equiv \max_{F \subseteq H} \left(\frac{e(F)}{v(F)} \right). \tag{2.1}
$$

The following theorem provides a sufficient condition on the distinguishability of laws \mathbb{P}_0 , \mathbb{P}_1 .

Theorem 1. Let $\{H_n\}_{n>1}$ be a sequence of non-empty graphs such that $v(H_n) = o(n)$ and for $q_0 \in (0,1)$ let $\mathbb{P}_{0,n}$ be the null model with edge density q_0 , and $\mathbb{P}_{1,n}$ be planted model with parameters H_n , q_0 . If

$$
\liminf_{n \to \infty} \frac{d(H_n) \log(1/q_0)}{\log n} > 1,
$$
\n(2.2)

then the two laws $\mathbb{P}_{0,n}$, $\mathbb{P}_{1,n}$ are strongly distinguishable.

Remark 2.1. The proof of this theorem also provides an explicit test $T : \mathcal{G}_n \to \{0,1\}$ which has asymptotically vanishing error probability under the assumptions of the theorem. Let $k(n)$ $v(F_n)$, where $F_n \subseteq H_n$ is the subgraph of H_n with the smallest number of vertices, such that $e(F_n)/v(F_n) = d(H_n)$. Then, the test developed in the proof requires searching over all subsets of $k(n)$ vertices which, in most cases, is non-polynomial.

The next theorem provides condition under which the two laws are indistinguishable.

Theorem 2. Let $\{H_n\}_{n>1}$, $q_0 \in (0,1)$, $\mathbb{P}_{0,n}$, $\mathbb{P}_{1,n}$ be as in Theorem [1.](#page-5-1) Then the two models are not weakly distinguishable if

$$
\limsup_{n \to \infty} \frac{d(H_n) \log(1/q_0) + (5/2) \log v(H_n)}{\log n} < 1. \tag{2.3}
$$

Further, if $d(H_n) = o(\log v(H_n))$, then the laws \mathbb{P}_0 , \mathbb{P}_1 are not weakly distinguishable if

$$
\limsup_{n \to \infty} \frac{v(H_n)}{n^{1/2}} = 0.
$$
\n(2.4)

Note that, under the condition $v(H_n) = o(n^{\alpha})$ for any $\alpha > 0$ (i.e. when the hidden subgraph is 'not too large'), or when $d(H_n)/\log v(H_n) \to \infty$ as $n \to \infty$ (i.e. when the hidden subgraph is 'dense enough'), this bound matches the positive result of Theorem [1.](#page-5-1) We illustrate these results with a few examples in Appendix [A.](#page-15-8)

3 Computationally efficient tests

In this section we study two computationally plausible algorithms for detecting the planted subgraph in the setting described in Section [1.](#page-0-0) The first method leverages the spectral properties of the given graph for solving the problem. In this case, we establish sufficient conditions under which the algorithm succeeds with high probability. We then show that a modification of the spectral algorithm can be used to identify the hidden subgraph. The second approach uses an SDP relaxation of the problem, that is a priori more powerful than the spectral approach.

3.1 Spectral algorithm

For $p \in [0, 1)$ we denote by $\mathbf{A}_{\mathcal{C}}^p$ $_G^p$ the *shifted adjacency matrix* of the graph G , defined as follows:

$$
\left(\boldsymbol{A}_{G}^{p}\right)_{ij} = \begin{cases} 1 & \text{if } (i,j) \in E(G), \\ -p/(1-p) & \text{otherwise.} \end{cases}
$$
\n(3.1)

Further, we will denote by $A_G = A_G^0$ the $0-1$ adjacency matrix of G. Recall that $\lambda_1(A_G^p)$ $_{G}^{p})\geq$ $\lambda_2(\boldsymbol{A}_{\mathcal{C}}^p)$ $\binom{p}{G} \geq \cdots \geq \lambda_n(\boldsymbol{A}_{C}^p)$ $\binom{p}{G}$ denote the eigenvalues of \mathbf{A}_{G}^{p} C_G^p . The spectral test is simply based on the leading eigenvalue:

$$
T_{\rm spec}(G) = \begin{cases} 1 & \text{if } \lambda_1(\mathbf{A}_G^{q_0}) \ge 2.1 \,\sigma(q_0)\sqrt{n}, \\ 0 & \text{otherwise} \end{cases} \tag{3.2}
$$

$$
\sigma(q_0) \equiv \sqrt{\frac{q_0}{1 - q_0}}.
$$
\n(3.3)

This algorithm was first proposed for the planted clique problem in [\[AKS98\]](#page-12-1). Note that this test uses the knowledge of q_0 , but does not assume the knowledge of planted subgraph H_n .

Theorem 3. Let $\{H_n\}_{n>1}$ be a sequence of non-empty graphs such that $v(H_n) = o(n)$ and for $q_0 \in (0,1)$ let $\mathbb{P}_{0,n}$ be the null model with edge density q_0 , and $\mathbb{P}_{1,n}$ be planted model with parameters H_n , q₀. Define $\sigma(q_0)$ as per Eq. [\(3.3\)](#page-6-1). If

$$
\liminf_{n \to \infty} \frac{\lambda_1(A_{H_n})}{\sqrt{n}} > 3\sigma(q_0), \tag{3.4}
$$

then the two laws $\mathbb{P}_{0,n}$, $\mathbb{P}_{1,n}$ are strongly distinguishable.

Remark 3.1. The constant 2.1 in Eq. [\(3.2\)](#page-6-2) can be reduced to $2 + \varepsilon$ for any $\varepsilon > 0$. In addition, we expect that with further work the constant 3 in Eq. [\(3.4\)](#page-6-3) can be reduced to $1 + \varepsilon$ for any $\varepsilon > 0$. These improvements are not the focus of the present paper.

Can spectral methods be used to identify the hidden subgraph? We start by noting that, even if H_n can be detected, a subset of its node might remain un-identified. As an example, let H_n be a graph over $k(n)$ vertices, whereby vertices $\{1, \ldots, k(n) - 1\}$ are connected by a clique, and vertex $k(n)$ is connected to the others by a single edge, see figure below:

Algorithm 1 Spectral algorithm for identifying hidden subgraphs in G

Input: Graph G, edge probability q_0 , size of hidden subgraph $k = v(H)$ **Output:** Estimated support of the hidden subgraph $S \subseteq V(G)$ **Initialize:** $n = v(G), t = kq_0 + 3\sqrt{kq_0 \log k}, S = \emptyset$ 1: for $i \in V(G)$ do

2: Set $v^{(i)} \equiv$ principal eigenvector of $(A_G^{q_0})$ $\,{}^{q_0}_{G})_{-i,-i}.$

3: Order the entries of
$$
\mathbf{v}^{(i)}: |v_{j(1)}^{(i)}| \geq |v_{j(2)}^{(i)}| \geq \cdots \geq |v_{j(n)}^{(i)}|
$$

- 4: Set $S_i \equiv \{j(1), \ldots, j(k)\}\;$
- 5: Set $d^{(i)} \equiv \text{# of edges between vertex } i$ and vertices in S_i
- 6: if $d^{(i)} > t$ then

7:
$$
S = S \cup \{i\}
$$

Ensure: S

Then Example [A.1](#page-15-9) implies that H_n can be detected with high probability as soon as $k(n) \geq$ $(1+\varepsilon)\log n/\log(1/q_0)$. As we will see below, the spectral algorithm detects H_n with high probability if $k(n) \geq 3\sigma(q_0)\sqrt{n} = \Theta(\sqrt{n})$. However it is intuitively clear (and not hard to prove) that the degree-one vertex in H_n cannot be identified reliably.

With this caveat in mind, Algorithm 1 gives a spectral approach to identify a subset of the vertices of the hidden subgraph. In order to characterize the set of 'important' vertices of H_n , we introduce the following notion.

Definition 3.1. Given a graph $H = (V(H), E(H))$, and $c \in (0, 1)$, we define the c-significant set of H, $S_c(H) \subseteq V(H)$ as the following set of vertices

$$
S_c(H) := \{ i \in V(H) : \deg(i) > c v(H) \}.
$$
\n(3.5)

We also need to to assume that the leading eigenvector of H is sufficiently spread out.

Definition 3.2. Let $H = (V(H), E(H))$ be a graph with adjacency matrix $A_H \in \{0, 1\}^{n \times n}$. For $\varepsilon \in (0,1)$, we say that H has spectral expansion ε , if

$$
1 - \varepsilon \ge \frac{\max\left(\lambda_2(A_H); -\lambda_n(A_H)\right)}{\lambda_1(A_H)}
$$
\n(3.6)

Finally let v be the leading eigenvector of A_H . We say that H is (ε, μ) -balanced in spectrum if it

has spectral expansion ε and

$$
\min_{i \in V(H)} |v_i| \ge \frac{\mu}{\sqrt{v(H)}}.\tag{3.7}
$$

The following definition helps us present our result on Algorithm [1.](#page-7-0)

Definition 3.3. Let $H = (V(H), E(H))$ be a graph. For any $i \in V(H)$, the graph obtained by removing i from H is denoted by $H \setminus i$. Then:

- 1. We say that H is (ε, μ) -strictly balanced in spectrum if for all $i \in V(H)$, $H \setminus i$ is (ε, μ) balanced in spectrum.
- 2. We define $\lambda_-(H)$ as

$$
\lambda_{-}(H) \equiv \min_{i \in V(H)} \lambda_{1}(\mathbf{A}_{H \setminus i}). \tag{3.8}
$$

The next theorem states sufficient conditions under which Algorithm [1](#page-7-0) succeeds in identifying the significant set of the planted subgraph.

Theorem 4. Given ${H_n}_{n\in\mathbb{N}}$, $q_0 \in (0,1)$, let $\mathbb{P}_{1,n}$ be the law of the random graph with edge density q₀ and planted subgraph H_n , cf. Section [1,](#page-0-0) and assume $G_n \sim \mathbb{P}_{1,n}$. Assume $v(H_n) = o(n)$ and that, for each n, H_n is (ε,μ) -strictly balanced in spectrum for some $\mu > 0, \varepsilon \in (0,1)$. Let δ be such that

$$
\frac{2\delta}{\mu^2(1-\delta)} < 1.
$$

Finally, assume that

$$
\liminf_{n \to \infty} \frac{|\lambda_{-}(H_n)|}{\sqrt{n}} > \frac{3\sigma(q_0)}{\varepsilon \delta},\tag{3.9}
$$

where $\lambda_-(H_n)$ is defined as per Eq. [\(3.8\)](#page-8-0).

Let S be the output of Algorithm [1,](#page-7-0) and set $c > \alpha$, $\alpha \equiv 2\delta/(\mu^2(1-\delta))$. Then the following statements hold with high probability as $n, v(H_n) \to \infty$:

- 1. S contains all the vertices of G_n that correspond to the c-significant set $S_c(H_n)$ of planted subgraph H_n .
- 2. S does not contain any vertex that does not correspond to those of the planted subgraph H_n .

Remark 3.2. Note that if $\min_{i \in V(H_n)} \deg(i) > cv(H_n)$ where c is as in Theorem [4](#page-8-1) (the minimum degree of nodes in the hidden subgraph is 'sufficiently large'), $S_c(H_n) = V(H_n)$ and under the assumptions of Theorem [4,](#page-8-1) Algorithm [1](#page-7-0) will find all the nodes of the planted subgraph H_n .

In the opposite case, H_n contains some 'low degree' vertices, namely, for some $i \in V(H_n)$, $deg(i) \le cv(H_n)$, we have $S_c(H_n) \subset V(H_n)$ strictly. Then, in order to find all vertices of the planted subgraph H_n in G_n , after finding the output of Algorithm [1,](#page-7-0) S, we can select the nodes $i \in V(G_n)$ such that $\deg_S(i) > (1+\varepsilon)q_0|S|$, for some $\varepsilon > 0$. Note that if $|S_c(H_n)|$ is $\omega(\log n)$ then for any $i \notin \varphi_0(H_n)$, $\deg_S(i) \leq (1+\varepsilon)q_0|S|$ with high probability. Hence, this procedure will not choose any node i such that $i \notin \varphi_0(H_n)$. Moreover, this procedure will find the planted subgraph H_n if for all nodes $i \in V(H_n)$, $\deg_S(i) > (1 + \varepsilon)q_0|S|$.

Note that for any graph $H = (V(H), E(H))$, $\lambda_1(A_H) \leq v(H)$. Hence by definition

$$
\lambda_{-}(H) \le v(H \setminus i) \le \frac{v(H)}{q_0}.
$$
\n(3.10)

Hence, the assumptions of Theorem [4](#page-8-1) imply in particular

$$
\liminf_{n \to \infty} \frac{v(H_n)}{n^{1/2}} > 0. \tag{3.11}
$$

We can compare this condition with the one of Theorem [1.](#page-5-1) If H_n is a dense graph, we expect generically $d(H_n) = \Theta(v(H_n))$, and hence there is a large gap between the condition of Theorem [1](#page-5-1) (that guarantees distinguishability) and that of Theorem [4.](#page-8-1) We illustrate this with a few examples in appendix [B.](#page-16-0)

3.2 SDP relaxation

Since the spectral method is generally sub-optimal with respect to the statistical detection threshold, it is natural to look for more powerful algorithms. In this section we use an SDP relaxation of the quadratic assignment problem, first proposed in [\[ZKRW98\]](#page-15-10), for motif detection.

Recall that we denote by A_G the adjacency matrix of graph G

$$
\left(\mathbf{A}_{G}\right)_{ij} = \begin{cases} 1 & \text{if } (i,j) \in E(G), \\ 0 & \text{otherwise.} \end{cases}
$$
\n(3.12)

We want to find a planted copy of a given graph H in graph G. Let $v(H) = k$, $v(G) = n$. We consider therefore the problem

maximize
$$
\mathbf{Tr}(\mathbf{A}_H \mathbf{\Pi}^\mathsf{T} \mathbf{A}_G \mathbf{\Pi})
$$

subject to $\mathbf{\Pi}^\mathsf{T} \mathbf{\Pi} = \mathbf{I}_k$
 $\mathbf{\Pi} \in \{0, 1\}^{n \times k}.$ (3.13)

This is a non-convex optimization problem known as Quadratic Assignment Problem (QAP) and is well studied in the literature, for example see [\[Bur13\]](#page-13-12). We will denote the value of this problem as $\mathsf{OPT}(G;H)$.

Note indeed that, $\Pi \in \{0,1\}^{n \times k}$ is feasible if it contains exactly one non-zero entry per column and at most one per row. Call $\varphi(i) \in [n]$ the position of the non-zero-entry of column $i \in [k]$. Then $\varphi \in \mathcal{L}(H, n)$ is a labeling of the vertices of H, and the objective function can be rewritten as

$$
\operatorname{Tr}(\mathbf{A}_{H}\mathbf{\Pi}^{\mathsf{T}}\mathbf{A}_{G}\mathbf{\Pi}) = 2 \sum_{(i,j)\in E(H)} (\mathbf{A}_{G})_{\varphi(i),\varphi(j)} \tag{3.14}
$$

Hence, if G contains a planted copy of H (e.g. under model $G \sim \mathbb{P}_1$), we have $\mathsf{OPT}(G;H) \geq 2e(H)$. This suggests the following optimization-based test:

$$
T_{\mathsf{OPT}}(G) = \begin{cases} 1 & \text{if } \mathsf{OPT}(G; H) \ge 2\,e(H), \\ 0 & \text{otherwise.} \end{cases} \tag{3.15}
$$

The proof of Theorem [1](#page-5-1) suggests that this test is nearly optimal, provided $d(H) = e(H)/v(H)$, i.e. H has no subgraph denser than H itself^{[1](#page-10-1)}.

Unfortunately, in general, $\mathsf{OPT}(G;H)$ is NP-complete even to approximate within a constant factor [\[SG76\]](#page-15-11). We will then resort to an SDP relaxation of the same problem. The following Lemma provides a different formulation of [\(3.13\)](#page-9-1).

Lemma 3.4. Let Π^* be an optimal solution of problem [\(3.13\)](#page-9-1). Then $\text{vec}(\Pi^*) = y^*$ such that $\boldsymbol{y}^* \boldsymbol{y}^{*^{\sf T}} = \boldsymbol{Y}^*$ is an optimal solution of the following problem

maximize
$$
\text{Tr}((A_G \otimes A_H)Y)
$$

\nsubject to $Y \in \{0,1\}^{nk \times nk}$
\n $Y \succeq 0$
\n $\text{Tr}(Y\mathbf{J}_{nk}) = k^2$
\n $\text{Tr}(Y(\mathbf{I}_n \otimes (e_i e_i^T))) = 1$ for $i = 1,2,...,k$
\n $\text{Tr}(Y((e_j e_j^T) \otimes \mathbf{I}_k)) \le 1$ for $j = 1,2,...,n$
\n $\text{rank}(Y) = 1$. (3.16)

Now, we try the following SDP relaxation of problem [\(3.16\)](#page-10-2) which is proposed in [\[ZKRW98\]](#page-15-10)

maximize
$$
\text{Tr}((\mathbf{A}_G \otimes \mathbf{A}_H)\mathbf{Y})
$$

\nsubject to $\mathbf{Y} \succeq 0$
\n $0 \le \mathbf{Y} \le 1$
\n $\text{Tr}(\mathbf{Y}\mathbf{J}_{nk}) = k^2$
\n $\text{Tr}(\mathbf{Y}(\mathbf{I}_n \otimes (e_i e_i^{\mathsf{T}}))) = 1$ for $i = 1, 2, ..., k$
\n $\text{Tr}(\mathbf{Y}((e_j e_j^{\mathsf{T}}) \otimes \mathbf{I}_k)) \le 1$ for $j = 1, 2, ..., n$ (3.17)

The following theorem states an upper bound on the performance of the hypothesis testing method that rejects the null hypothesis if $SDP(G; H) \geq 2e(H)$.

Theorem 5. Let $\{H_n\}_{n>1}$, $\mathbb{P}_{0,n}$, $\mathbb{P}_{1,n}$ be as in Theorem [1.](#page-5-1) Consider the hypothesis testing problem in which under null G_n is generated according to $\mathbb{P}_{0,n}$ and under alternative it is generated according to $\mathbb{P}_{1,n}$. Define $\sigma(q_0)$ as per Eq. [\(3.3\)](#page-6-1). If

$$
\limsup_{n\to\infty}\frac{\lambda_1(\mathbf{A}_{H_n})}{\sqrt{n}}<\frac{1}{4}\sigma(q_0),
$$

then for the method that rejects the null hypothesis if $\text{SDP}(G_n; H_n) \geq 2e(H_n)$,

$$
\mathbb{P}_{0,n}\{T(G_n)=1\} \to 1
$$

as $n \to \infty$.

4 The case of multiple planted subgraphs

In this section we would like to generalize the results given in Section [2](#page-5-0) to the regime in which $m_n > 1$ atypical subgraphs are added to a random graph. Namely, fix H_n and let $\varphi_1, \varphi_2, \ldots, \varphi_{m_n} \in$ $\mathcal{L}(H_n, n)$ be independent uniformly random labelings of $V(H_n)$ in [n]. As before, we let $\mathbb{P}_{0,n}$ be

¹If this is the not case, the optimization problem (3.13) can be modified replacing H by its densest subgraph.

the law of an Erdős-Renyi random graph with edge probability q_0 . On the other hand under $\mathbb{P}_{1,n}$ edges are conditionally independent given $\varphi_1, \varphi_2, \ldots, \varphi_{m_n}$, with

$$
\mathbb{P}_{1,n}((i,j)\in E \mid (\varphi_l)_{l\leq m_n}) = \begin{cases} 1 & \text{if } (i,j)\in \varphi_l(E(H_n)) \text{ for some } l\in \{1,2,\ldots,m_n\}, \\ q_0 & \text{otherwise.} \end{cases}
$$
(4.1)

We would like to find the conditions on $m_n, q_0, \{H_n\}$ under which the two laws $\mathbb{P}_{0,n}, \mathbb{P}_{1,n}$ are strongly or weakly distinguishable, generalizing Theorem [1](#page-5-1) and Theorem [2](#page-5-2) to $m_n > 1$. The following theorem states the sufficient condition under which the two laws are indistinguishable.

Theorem 6. Let $\{H_n\}_{n\geq 1}$ be a sequence of non-empty graphs and for $q_0 \in (0,1)$ let $\mathbb{P}_{0,n}$ be the null model with edge density q_0 and $\mathbb{P}_{1,n}$ be the planted model as in [\(4.1\)](#page-11-0) with parameters H_n , q_0 , m_n . Then the two models are not weakly distinguishable if

$$
\limsup_{n \to \infty} \frac{(5/2) \log v(H_n) + \log m_n}{\log n} < 1,\tag{4.2}
$$

and

$$
\limsup_{n \to \infty} \frac{d(H_n) \log(1/q_0) + (5/2) \log v(H_n)}{\log n} < 1. \tag{4.3}
$$

(4.4)

The following Theorem states sufficient conditions under which the two models are strongly distinguishable.

Theorem 7. Let $\{H_n\}_{n\geq 1}$, $q_0 \in (0,1)$, m_n , $\mathbb{P}_{0,n}$, $\mathbb{P}_{1,n}$ be as in Theorem [6.](#page-11-1) Then the two laws $\mathbb{P}_{0,n}, \mathbb{P}_{1,n}$ are strongly distinguishable if

$$
\liminf_{n \to \infty} \frac{m_n e(H_n)}{n} = \infty, \tag{4.5}
$$

or if

$$
\liminf_{n \to \infty} \frac{d(H_n) \log(1/q_0)}{\log n} > 1.
$$
\n(4.6)

Remark 4.1. While the necessary and sufficient conditions in the above theorems do not match in general, they do match in specific regimes of interest. A first regime is the one of m_n bounded: in this case we recover the same asymptotics as for $m_n = 1$, cf. Section [2.](#page-5-0)

A second regime is obtained when the planted graphs $\{H_n\}$ have bounded size: $v(H_n) \leq C$ for all n. Theorem [6](#page-11-1) implies that the two models are not weakly distinguishable if $m_n \leq n^{1-\varepsilon}$ for some $\varepsilon > 0$ and all *n* large enough.

Vice versa, by Theorem [7,](#page-11-2) they are strongly distinguishable if $m_n/n \to \infty$. In other words, we have a characterization of the distinguishability threshold that is tight up to sub-polynomial factors.

Remark 4.2. The expected number of edges under the null model $\mathbb{P}_{0,n}$ is roughly $n^2q_0/2$, and its standard deviation is of order n. The sufficient condition (4.5) is therefore equivalent to requiring that the total number of edges in the planted graphs is much larger than this standard deviation.

The proof of Theorem [7](#page-11-2) constructs a simple test $T: \mathcal{G}_n \to \{0,1\}$, by letting $T(G_n) = 1$ if $e(G_n) \ge t_*$ and $T(G_n) = 0$ otherwise.

Notice that condition [\(4.6\)](#page-11-4) is instead the same as in Theorem [1.](#page-5-1) In this regime, it is sufficient to find a single copy of the highest density subgraph of H_n , and the multiplicity m_n does not seem to help.

Remark 4.3. It is possible to use the spectral algorithm in subsection [3.1](#page-6-0) to detect the m_n planted subgraphs by setting $k = v(H_n)m_n$ in Algorithm [1.](#page-7-0)

Note that if $m_n v(H_n)/n^{1/2} \to 0$ then the m_n planted subgraphs will have disjoint vertex sets with high probability as $n \to \infty$. Hence the law $\mathbb{P}_{1,n}$ studied in this section is the same as the one obtained by planting a graph consisting in the disjoint union of m_n copies of H_n . The top eigenvalue of the graph consisting of m_n disjoint copies of a graph H_n is the same as the top eigenvalue of H_n . Hence, in this regime, we expect the spectral method to succeed in detecting m_n motifs under the same conditions under which it succeeds in detecting one motif.

Acknowledgments

H.J. was supported by the William R. Hewlett Stanford Graduate Fellowship. A.M. was partially supported by NSF grants CCF-1319979 and DMS-1106627 and the AFOSR grant FA9550-13-1- 0036.

References

- [AAM+11] Noga Alon, Sanjeev Arora, Rajsekar Manokaran, Dana Moshkovitz, and Omri Weinstein, Inapproximability of densest κ -subgraph from average case hardness, Unpublished manuscript 1 (2011).
- [ACV14] Ery Arias-Castro and Nicolas Verzelen, Community detection in dense random networks, The Annals of Statistics 42 (2014), no. 3, 940–969.
- [ADBL11] Alekh Agarwal, John C Duchi, Peter L Bartlett, and Clement Levrard, Oracle inequalities for computationally budgeted model selection, Proceedings of the 24th Annual Conference on Learning Theory, 2011, pp. 69–86.
- [AKS98] Noga Alon, Michael Krivelevich, and Benny Sudakov, Finding a large hidden clique in a random graph, Proceedings of the ninth annual ACM-SIAM symposium on Discrete algorithms, Society for Industrial and Applied Mathematics, 1998, pp. 594–598.
- [Alo07] Uri Alon, Network motifs: theory and experimental approaches, Nature Reviews Genetics 8 (2007), no. 6, 450–461.
- [BCC+10] Aditya Bhaskara, Moses Charikar, Eden Chlamtac, Uriel Feige, and Aravindan Vijayaraghavan, Detecting high log-densities: an o (n 1/4) approximation for densest k-subgraph, Proceedings of the forty-second ACM symposium on Theory of computing, ACM, 2010, pp. 201–210.
- [BCV+12] Aditya Bhaskara, Moses Charikar, Aravindan Vijayaraghavan, Venkatesan Guruswami, and Yuan Zhou, Polynomial integrality gaps for strong sdp relaxations of densest ksubgraph, Proceedings of the twenty-third annual ACM-SIAM symposium on Discrete Algorithms, Society for Industrial and Applied Mathematics, 2012, pp. 388–405.
- [BD11] Joseph Blitzstein and Persi Diaconis, A sequential importance sampling algorithm for generating random graphs with prescribed degrees, Internet Mathematics $6(2011)$, no. 4, 489–522.
- [BHK+16] Boaz Barak, Samuel B Hopkins, Jonathan Kelner, Pravesh Kothari, Ankur Moitra, and Aaron Potechin, A nearly tight sum-of-squares lower bound for the planted clique problem, Foundations of Computer Science (FOCS), 2016 IEEE 57th Annual Symposium on, FOCS '16, IEEE, 2016, pp. 428–437.
- [BR13] Quentin Berthet and Philippe Rigollet, Complexity theoretic lower bounds for sparse principal component detection, Conference on Learning Theory, 2013, pp. 1046–1066.
- [Bur13] Rainer E Burkard, Quadratic assignment problems, Springer, 2013.
- [Chu97] Fan RK Chung, Spectral graph theory, no. 92, American Mathematical Soc., 1997.
- [DLSS14] Amit Daniely, Nati Linial, and Shai Shalev-Shwartz, From average case complexity to improper learning complexity, Proceedings of the forty-sixth annual ACM symposium on Theory of computing, ACM, 2014, pp. 441–448.
- [DM14] Yash Deshpande and Andrea Montanari, Finding hidden cliques of size $\sqrt{N/e}$ in nearly linear time, Foundations of Computational Mathematics (2014), 1–60.
- [DM15] Yash Deshpande and Andrea Montanari, Improved sum-of-squares lower bounds for hidden clique and hidden submatrix problems, Proceedings of The 28th Conference on Learning Theory (Paris, France) (Peter Grnwald, Elad Hazan, and Satyen Kale, eds.), Proceedings of Machine Learning Research, vol. 40, PMLR, 03–06 Jul 2015, pp. 523– 562.
- [EK10] David Easley and Jon Kleinberg, Networks, crowds, and markets: Reasoning about a highly connected world, Cambridge University Press, 2010.
- [Fei02] Uriel Feige, Relations between average case complexity and approximation complexity, Proceedings of the thiry-fourth annual ACM symposium on Theory of computing, ACM, 2002, pp. 534–543.
- [FGR+13] Vitaly Feldman, Elena Grigorescu, Lev Reyzin, Santosh Vempala, and Ying Xiao, Statistical algorithms and a lower bound for detecting planted cliques, Proceedings of the Forty-fifth Annual ACM Symposium on Theory of Computing (New York, NY, USA), STOC '13, ACM, 2013, pp. 655–664.
- [FK00] Uriel Feige and Robert Krauthgamer, Finding and certifying a large hidden clique in a semirandom graph, Random Structures and Algorithms 16 (2000), no. 2, 195–208.
- [FPK01a] Uriel Feige, David Peleg, and Guy Kortsarz, The dense k-subgraph problem, Algorithmica 29 (2001), no. 3, 410–421.
- $[FPK01b] \quad _____$, The dense k-subgraph problem, Algorithmica 29 (2001), no. 3, 410–421.
- [GM75] G. R. Grimmett and C. J. H. McDiarmid, On colouring random graphs, Mathematical Proceedings of the Cambridge Philosophical Society 77 (1975), no. 2, 313?324.
- [Gra73] Mark S Granovetter, The strength of weak ties, American journal of sociology (1973), 1360–1380.
- [Hås99] Johan Håstad, *Clique is hard to approximate withinn1- ε*, Acta Mathematica 182 (1999), no. 1, 105–142.
- [HKP15] Samuel B Hopkins, Pravesh K Kothari, and Aaron Potechin, Sos and planted clique: Tight analysis of mpw moments at all degrees and an optimal lower bound at degree $four,$ arXiv:1507.05230 $(2015).$
- [HWX15] Bruce Hajek, Yihong Wu, and Jiaming Xu, Computational lower bounds for community detection on random graphs, Proceedings of The 28th Conference on Learning Theory (Paris, France) (Peter Grnwald, Elad Hazan, and Satyen Kale, eds.), Proceedings of Machine Learning Research, vol. 40, PMLR, 03–06 Jul 2015, pp. 899–928.
- [HWX17] Bruce Hajek, Yihong Wu, and Jiaming Xu, Information limits for recovering a hidden community, IEEE Transactions on Information Theory 63 (2017), no. 8, 4729–4745.
- [Jer92] Mark Jerrum, Large cliques elude the metropolis process, Random Structures & Algorithms 3 (1992), no. 4, 347–359.
- [KA05] Nadav Kashtan and Uri Alon, Spontaneous evolution of modularity and network motifs, Proceedings of the National Academy of Sciences of the United States of America 102 (2005), no. 39, 13773–13778.
- [Kho01] Subhash Khot, *Improved inapproximability results for maxclique, chromatic number* and approximate graph coloring, Foundations of Computer Science, 2001. Proceedings. 42nd IEEE Symposium on, IEEE, 2001, pp. 600–609.
- [Kho06] , Ruling out ptas for graph min-bisection, dense k-subgraph, and bipartite clique, SIAM Journal on Computing 36 (2006), no. 4, 1025–1071.
- [KIMA04] Nadav Kashtan, Shalev Itzkovitz, Ron Milo, and Uri Alon, Efficient sampling algorithm for estimating subgraph concentrations and detecting network motifs, Bioinformatics 20 (2004), no. 11, 1746–1758.
- [Man17] Pasin Manurangsi, Almost-polynomial ratio eth-hardness of approximating densest ksubgraph, Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing, ACM, 2017, pp. 954–961.
- [MKI+03] Ron Milo, Nadav Kashtan, Shalev Itzkovitz, Mark EJ Newman, and Uri Alon, On the uniform generation of random graphs with prescribed degree sequences, arXiv preprint cond-mat/0312028 (2003).
- [Mon15] Andrea Montanari, Finding one community in a sparse graph, Journal of Statistical Physics 161 (2015), no. 2, 273–299.
- [MPW15] Raghu Meka, Aaron Potechin, and Avi Wigderson, Sum-of-squares lower bounds for planted clique, Proceedings of the Forty-seventh Annual ACM Symposium on Theory of Computing (New York, NY, USA), STOC '15, ACM, 2015, pp. 87–96.
- [MSOI+02] Ron Milo, Shai Shen-Orr, Shalev Itzkovitz, Nadav Kashtan, Dmitri Chklovskii, and Uri Alon, Network motifs: simple building blocks of complex networks, Science 298 (2002), no. 5594, 824–827.
- [RS15] Prasad Raghavendra and Tselil Schramm, Tight lower bounds for planted clique in the degree-4 sos program, arXiv:1507.05136 (2015).
- [SG76] Sartaj Sahni and Teofilo Gonzalez, P-complete approximation problems, Journal of the ACM (JACM) 23 (1976), no. 3, 555–565.
- [SSR+05] Sen Song, Per Jesper Sjöström, Markus Reigl, Sacha Nelson, and Dmitri B Chklovskii, Highly nonrandom features of synaptic connectivity in local cortical circuits, PLoS Biol 3 (2005), no. 3, e68.
- [Tao12] Terence Tao, Topics in random matrix theory, vol. 132, American Mathematical Soc., 2012.
- [TS07] Thomas Thorne and Michael PH Stumpf, Generating confidence intervals on biological networks, BMC bioinformatics 8 (2007), no. 1, 467.
- [VAC+15] Nicolas Verzelen, Ery Arias-Castro, et al., Community detection in sparse random networks, The Annals of Applied Probability 25 (2015), no. 6, 3465–3510.
- [WF94] Stanley Wasserman and Katherine Faust, Social network analysis: Methods and applications, vol. 8, Cambridge university press, 1994.
- [YLSK+04] Esti Yeger-Lotem, Shmuel Sattath, Nadav Kashtan, Shalev Itzkovitz, Ron Milo, Ron Y Pinter, Uri Alon, and Hanah Margalit, Network motifs in integrated cellular networks of transcription–regulation and protein–protein interaction, Proceedings of the National Academy of Sciences of the United States of America 101 (2004), no. 16, 5934–5939.
- [ZKRW98] Qing Zhao, Stefan E Karisch, Franz Rendl, and Henry Wolkowicz, Semidefinite programming relaxations for the quadratic assignment problem, Journal of Combinatorial Optimization 2 (1998), no. 1, 71–109.

A Examples: Statistical limits

Example A.1. Recall that K_m denotes the complete graph over m vertices (hence having degree $m-1$). Setting $H_n = K_{k(n)}$ we recover the hidden clique problem. In this case $d(H_n) = (k(n)-1)/2$. Hence, our theorems imply that the two laws are strongly distinguishable if $\liminf_{n\to\infty} k(n)/\log n$ $2/\log(1/q_0)$, and are not weakly distinguishable if $\limsup_{n\to\infty} k(n)/\log n < 2/\log(1/q_0)$.

Example A.2. Let Q_m be the hypercube graph over 2^m vertices (hence having degree m): this is the graph whose vertices are binary vectors of length m, connected by an edge whenever their Hamming distance is exactly equal to one. Set $H_n = Q_{\log_2 k(n)}$. In other words, H_n is an hypercube over $k(n)$ vertices. It is easy to see that $d(H_n) = (\log_2 k(n))/2$.

Let $\gamma(q_0) \equiv 2/\log_2(1/q_0)$. Theorem [1](#page-5-1) implies that this graph can be detected provided $k(n) \ge$ $n^{\gamma(q_0)+\varepsilon}$ for some $\varepsilon > 0$ and all n large enough. On the other hand, Theorem [2](#page-5-2) implies that it cannot be detected if $k(n) \leq n^{2\gamma(q_0)/(2+5\gamma(q_0))-\varepsilon}$ for some $\varepsilon > 0$ and all n large enough. Hence, we can see that the lower and upper bounds for distinguishability are close for small q_0 and as q_0 increases the gap between the bounds increases.

Example A.3. Let H_n be a regular tree with degree $d(n)$ and $r(n)$ generations (hence $v(H_n)$ = $1 + d(n)[(d(n) - 1)^{r(n)} - 1]/(d(n) - 2), e(H_n) = v(H_n) - 1).$ In this case for any $F_n \subseteq H_n$, $e(F_n) \le v(F_n) - 1$. Therefore, $d(H_n) = 1 - 1/v(H_n) < 1$ and Theorem [1,](#page-5-1) cannot guarantee the strong distinguishability of the hypotheses. Furthermore, $\limsup_{n\to\infty} d(H_n)/\log v(H_n) = 0$ and we are in the low density region. Hence, Theorem [2](#page-5-2) implies that the null and planted models are not weakly distinguishable if $\limsup_{n\to\infty} d(n)^{r(n)+1}/n^{1/2} = 0$.

Example A.4. Let C_k^m be the m-th power of the cycle over k vertices. This is the graph with vertex set $\{1, \ldots, k\}$, and two vertices i, j are connected if $|i - j - bk| \leq m$ for some $b \in \mathbb{N}$. Let $H_n = C_{k(n)}^{m(n)}$ $\binom{m(n)}{k(n)}$, for two functions $m(n)$, $k(n)$. In this case, $v(H_n) = k(n)$ and for all $i \in V(H_n)$, $deg(i) = 2m(n)$. Therefore, for any $F_n \subseteq H_n$, $e(F_n) \leq m(n)v(F_n)$. Since $e(H_n) = m(n)k(n)$, by definition, $d(H_n) = m(n)$. Using Theorem [1,](#page-5-1) two models are strongly distinguishable if $\liminf_{n\to\infty}m(n)/\log(n) > \log(1/q_0)$. In addition, depending on $k(n)$, $m(n)$, we can be in different graph density regimes. If $m(n) = \omega(\log k(n))$, the laws are not weakly distinguishable if $\limsup_{n\to\infty} m(n)/\log(n) < \log(1/q_0)$ and we get a tight characterization. If $m(n) = o(\log k(n)),$ two models cannot be weakly distinguished if $\limsup_{n\to\infty} k(n)/n^{1/2} = 0$. Finally, for the intermediate regime where $m(n) = \Theta(\log k(n))$, if $\limsup_{n\to\infty}((5/2)\log k(n) + m(n)\log(1/q_0))/\log n < 1$ the models are not weakly distinguishable.

B Examples: Spectral Algorithm

Example B.1. Let $H_n = K_{k(n)}$. Assume that, $v(H_n) = k(n)$ is $o(n)$. We have $\lambda_1(A_{H_n}) =$ $k(n) - 1$ and Theorem [3](#page-6-4) implies that the laws are strongly distinguishable using the spectral test if $\liminf_{n\to\infty} k(n)/n^{1/2} > 3\sigma(q_0)$. This shows a gap between the performance of the spectral test and the statistical bound of Theorem [1.](#page-5-1)

In order to express results on identifying the hidden subgraph in this case, first note that for all $i \in V(H_n)$, $\deg(i) = k(n) - 1$ and all nodes of H_n are in the c-significant set of H_n for $c < 1 - 1/k(n)$ as per Definition [3.1.](#page-7-1) Assuming that $k(n) \to \infty$ as $n \to \infty$, for any $c > 0$ all nodes of H_n are in the c-significant set of H_n for large enough n. Also, the leading eigenvector of A_{H_n} is $\mathbf{1}_{k(n)}$, its corresponding eigenvalue is $k(n) - 1$ and the rest of eigenvalues are -1. Setting $\varepsilon_0(k(n)) \equiv 1 - 1/(k(n-1))$, based on definition [3.2,](#page-7-2) for each n, H_n is $(\varepsilon, 1)$ balanced in spectrum for $\varepsilon < \varepsilon_0(k(n))$. Using the fact that for any $i \in V(H_n)$, $H_n \setminus i$ is $K_{k(n)-1}$, we deduce that for each n, H_n is $(\varepsilon, 1)$ - strictly balanced in spectrum for $\varepsilon < \varepsilon_0(k(n)-1)$ and $\lambda_-(H_n) = k(n)-2$. Note that, $\varepsilon_0(k(n)-1) \to 1$ as $n \to \infty$. Therefore, using Theorem [4,](#page-8-1) if $\liminf_{n \to \infty} k(n)/n^{1/2} > 9\sigma(q_0)$, Algorithm [1](#page-7-0) can find the planted clique with high probability as $n, v(H_n) \to \infty$.

Example B.2. Set $H_n = Q_{\log_2 k(n)}$ as in Example [A.2.](#page-16-1) Since the hypercube is a regular graph, $\lambda_1(A_{H_n}) = \log_2 k(n)$ and Theorem [3](#page-6-4) implies that two models can be strongly distinguished using the spectral test if $\liminf_{n\to\infty} \log_2 k(n)/n^{1/2} > 3\sigma(q_0)$. However, this never happens since $k \leq n$. Therefore, Theorem [3](#page-6-4) cannot guarantee the strong distinguishability of the hypotheses using the spectral test. Similarly, Theorem [4](#page-8-1) does not imply the success of Algorithm [1](#page-7-0) in finding the planted hypercube.

Example B.3. Let H_n be a regular tree with degree $d(n)$ and $r(n)$ generations as in Example [A.3.](#page-16-2) For a large regular tree, $\lambda_1(A_{H_n})$ is of order of $2\sqrt{d(n)-1}$ as $v(H_n) \to \infty$. Hence, based on Theorem [3,](#page-6-4) two laws are strongly distinguishable using the spectral test if $\liminf_{n\to\infty} d(n)/n$ $(9/4)\sigma^2(q_0)$. Therefore, $v(H_n)$ cannot be $o(n)$ and Theorem [3](#page-6-4) cannot guarantee the strong distinguishability of two models under any conditions. Recall that Theorem [1,](#page-5-1) also, could not guarantee the strong distinguishability for this example under any conditions.

As a side note, note that if q_0 is known a priori -which is not a practical assumption- and $\liminf_{n\to\infty} d(n)/(nq_0) > 0 = c$, the null and planted models can be distinguished only by looking at the maximum degree in the graph G_n . In fact, under the null model the maximum degree of at the maximum degree in the graph G_n . In fact, under the nun moder the maximum degree of graph G_n is less than or equal $nq_0 + \Theta(\sqrt{nq_0 \log n})$ with high probability. Therefore, the test that rejects the null iff the maximum degree of G_n is bigger than or equal $(1+\varepsilon)nq_0$ strongly distinguishes two models under this assumption. Subsequently, under this condition, the high degree nodes can be used to find the planted tree.

In addition, since $d(n)/v(H_n) \to 0$ as $v(H_n) \to \infty$ for the sequence of regular trees, Theorem [4](#page-8-1) cannot imply the success of Algorithm [1](#page-7-0) in finding the planted regular tree. In other words, Algorithm [1](#page-7-0) fails in identifying the planted regular tree because it does not contain *sufficiently high* degree vertices.

Example B.4. Set $H_n = C_{k(n)}^{m(n)}$ $\binom{m(n)}{k(n)}$ as in Example [A.4.](#page-16-3) As we had for previous examples, since H_n is a sequence of regular graphs, $\lambda_1(A_{H_n}) = 2m(n)$. Therefore, two models are strongly distinguishable using the spectral test if $\liminf_{n\to\infty} m(n)/n^{1/2} > (3/2)\sigma(q_0)$ and the gap between the results of Theorems [1](#page-5-1) and [3](#page-6-4) is similar to Example [B.1.](#page-16-4)

Assuming that $\liminf_{n\to\infty} m(n)/k(n) = (c/2) > 0$, Theorem [4](#page-8-1) can used to guarantee the performance of Algorithm [1](#page-7-0) in identifying the hidden subgraph. Under this condition, all vertices of H_n are in the c-significant set of H_n for large enough n. Note that A_{H_n} is a circulant matrix, its principal eigenvalue is $\lambda_1(A(H_n)) = 2m(n)$, corresponding eigenvector is $\mathbf{1}_{k(n)}$ and other eigenvalues are $\sum_{i=1}^{m(n)} 2 \cos(2\pi i j/k(n))$ for $j = 1, 2, ..., k(n) - 1$. Therefore, $\lambda_2(A(H_n)) \leq$ $2m(n) - \sum_{i=1}^{m(n)} 4\pi^2 i^2 / (k(n))^2 = 2m(n)(1 - 2\pi^2 m(n)^2 / (3k(n)^2)) + o(m(n))$. Hence, for large enough $n, \lambda_2(A(H_n)) \leq 2m(n)(1-\varepsilon) = \lambda_1(A(H_n))(1-\varepsilon)$ and H_n is $(\varepsilon, 1)$ -balanced in spectrum, where $\varepsilon = \pi^2 c^2/6$. For any $i \in V(H_n), \lambda_1(\mathbf{A}(H_n \backslash i)) \geq 2e(H_n \backslash i)/(k(n)-1) = 2m(n)(1-1/(k(n)-1))$. Using Cauchy's interlacing theorem, $\lambda_2(\mathbf{A}(H_n\backslash i)) \leq \lambda_2(\mathbf{A}(H_n)) \leq 2m(n)(1-\varepsilon) \leq \lambda_1(\mathbf{A}(H_n\backslash i))(1-\varepsilon_0)$ where $(1 - \varepsilon)/(1 - 1/(k(n-1)) = 1 - (\varepsilon - \varepsilon_n'), \varepsilon_n' \to 0$ as $n \to \infty$. In addition, for large enough n, for v, the leading eigenvector of $\mathbf{A}(H_n \setminus i)$, we have $v_i \ge (1 - \delta'_n)/\sqrt{k(n-1)}$ for $i = 1, 2, ..., n-1$ where $\delta'_n \to 0$ as $n \to \infty$. Therefore, for any $0 < \varepsilon' < \varepsilon$, $0 < \mu < 1$, H_n is $(\varepsilon - \varepsilon', \mu)$ -strictly balanced in spectrum, for large enough n. In addition, $\lambda_-(H_n) \geq 2m(n)(1-1/(k(n)-1))$. Thus, using Theorem [4,](#page-8-1) if $\liminf_{n\to\infty} m(n)/n^{1/2} > 9\sigma(q_0)/(2\varepsilon)$, Algorithm [1,](#page-7-0) can find the planted subgraph with high probability as $n, k(n) \to \infty$.

C Proofs: Statistical limits

We start with the following preliminary lemma.

Lemma C.1. Let, for each $n, Z : \mathcal{G}_n \to \mathbb{R}_+$ be such that,

$$
Z(G_n) = \mathbb{E}_{0,n}\{Z(G_n)\}\frac{\mathrm{d}\mathbb{P}_{1,n}}{\mathrm{d}\mathbb{P}_{0,n}}(G_n).
$$

Further let $Z_n = Z(G_n)$. Then, $\mathbb{P}_{0,n}$ and $\mathbb{P}_{1,n}$ are strongly distinguishable if and only if, under $\mathbb{P}_{0,n}$,

$$
\frac{Z_n}{\mathbb{E}_{0,n}Z_n} \xrightarrow{p} 0.
$$

They are not weakly distinguishable if and only if, along some subsequence $\{n_k\}$,

$$
\frac{Z_n}{\mathbb{E}_{0,n}Z_n} \xrightarrow{p} 1.
$$

Proof. First assume that along some subsequence $\{n_k\}$ under $\mathbb{P}_{0,n}$

$$
\frac{Z_n}{\mathbb{E}_0 Z_n} \to 1,\tag{C.1}
$$

in probability. For a test $T: \mathcal{G}_n \to \{0,1\}$, define the risk $\gamma(T)$ as

$$
\gamma_n(T) = \mathbb{P}_{0,n}(T(G_n) = 1) + \mathbb{P}_{1,n}(T(G_n) = 0).
$$

Now, for any test T we have

$$
\gamma_n(T) = \int (1 - T) d\mathbb{P}_{1,n} + \int T d\mathbb{P}_{0,n}
$$

=
$$
\int \left((1 - T) \frac{Z_n}{\mathbb{E}_0 Z_n} + T \right) d\mathbb{P}_{0,n}
$$

$$
\geq \int \left(\left(1 - 1 \left\{ \frac{Z_n}{\mathbb{E}_0 Z_n} > 1 \right\} \right) \frac{Z_n}{\mathbb{E}_0 Z_n} + 1 \left\{ \frac{Z_n}{\mathbb{E}_0 Z_n} > 1 \right\} \right) d\mathbb{P}_{0,n}.
$$

Using [\(C.1\)](#page-18-0), the last term goes to 1 as $n \to \infty$. Therefore along $\{n_k\}$

$$
\liminf_{n \to \infty} \{ \inf_T \{ \gamma_n(T) \} \} \ge 1.
$$

Which implies that for all tests T ,

$$
\limsup_{n \to \infty} \left[\mathbb{P}_{0,n}(T(G_n) = 1) + \mathbb{P}_{1,n}(T(G_n) = 0) \right] = 1.
$$

Thus, $\mathbb{P}_{0,n}$, $\mathbb{P}_{1,n}$ are not weakly distinguishable.

Now, assume that

$$
\frac{Z_n}{\mathbb{E}_0 Z_n} \to 0. \tag{C.2}
$$

As above, It is easy to see that the test $T = \mathbf{1} \{Z_n / \mathbb{E}_0 Z_n > 1\}$ satisfies

$$
\limsup_{n \to \infty} \mathbb{P}_{0,n}(T(G_n) = 1) = \limsup_{n \to \infty} \mathbb{P}_{1,n}(T(G_n) = 0) = 0.
$$

Therefore in this case $\mathbb{P}_{0,n}$, $\mathbb{P}_{1,n}$ are strongly distinguishable.

 \Box

In order to state the proof our results, given a graph $G_n \in \mathcal{G}_n$, we define $U_{G_n} : \mathcal{L}(H_n; n) \to \mathbb{N}$ by

$$
U_{G_n}(\varphi) \equiv |\varphi(E(H_n)) \cap E(G_n)|.
$$

For $n \ge v(H_n)$, we let

$$
N(H_n; G_n) \equiv \left| \left\{ \varphi \in \mathcal{L}(H_n; n) : U_{G_n}(\varphi) = e(H_n) \right\} \right|.
$$

Let $\mathbb{P}_{0,n}, \mathbb{P}_{1,n}$ be defined as in Section [1,](#page-0-0) note that

$$
\mathbb{E}_{0,n} N(H_n; G_n) = (n)_{v(H_n)} q_0^{e(H_n)}.
$$

Further, we can write

$$
\frac{\mathrm{d}\mathbb{P}_{1,n}}{\mathrm{d}\mathbb{P}_{0,n}}(G_n) = \frac{1}{(n)_{v(H_n)}} \sum_{\varphi \in \mathcal{L}(H_n;n)} \prod_{(i,j) \in E(H_n)} \left[\left(\frac{1}{q_0} \right) \mathbb{I}\{(\varphi(i), \varphi(j)) \in E(G_n)\} \right].
$$

Thus,

$$
\frac{\mathrm{d}\mathbb{P}_{1,n}}{\mathrm{d}\mathbb{P}_{0,n}}(G_n) = \frac{1}{(n)_{v(H_n)}} \left(\frac{1}{q_0}\right)^{e(H_n)} \sum_{\varphi \in \mathcal{L}(H_n;n)} \mathbb{I}\left\{|\varphi(E(H_n)) \cap E(G_n)| = |E(H_n)|\right\}.
$$

Therefore,

$$
\frac{\mathrm{d}\mathbb{P}_{1,n}}{\mathrm{d}\mathbb{P}_{0,n}}(G_n) = \frac{1}{(n)_{v(H_n)}} \left(\frac{1}{q_0}\right)^{e(H_n)} N(H_n; G_n) = \frac{1}{\mathbb{E}_{n,0}\{N(H_n; G_n)\}} N(H_n; G_n). \tag{C.3}
$$

Now we can prove Theorems [1,](#page-5-1) [2.](#page-5-2)

C.1 Proof of Theorem [1](#page-5-1)

Proof. Let \tilde{H}_n be a subgraph of H_n that satisfies $d(H_n) = e(\tilde{H}_n)/v(\tilde{H}_n)$. Using [\(C.3\)](#page-19-0), we can write

$$
\mathbb{P}_{0,n}\left(\frac{\mathrm{d}\mathbb{P}_{1,n}}{\mathrm{d}\mathbb{P}_{0,n}}(G_n) > 0\right) = \mathbb{P}_{0,n}\left(N(H_n; G_n) > 0\right)
$$
\n
$$
\leq \mathbb{P}_{0,n}(N(\tilde{H}_n; G_n) > 0)
$$
\n
$$
\leq \mathbb{E}_{0,n}N(\tilde{H}_n; G_n)
$$
\n
$$
\leq n^{v(\tilde{H}_n)}q_0^{e(\tilde{H}_n)}
$$
\n
$$
= \exp\left\{v(\tilde{H}_n)\log n\left(1 - \frac{d(H_n)\log(1/q_0)}{\log n}\right)\right\}
$$

which goes to zero as $n \to \infty$ when [\(2.2\)](#page-5-3) holds. Therefore, under the assumptions of Theorem [1,](#page-5-1) under $\mathbb{P}_{0,n}$,

$$
\frac{\mathrm{d} \mathbb{P}_{1,n}}{\mathrm{d} \mathbb{P}_{0,n}}(G_n) \xrightarrow{\ p\ } 0
$$

and using Lemma [C.1](#page-18-1) the proof is complete.

 \Box

C.2 Proof of Theorem [2](#page-5-2)

We start by stating the following lemma.

Lemma C.2. Let $\{H_n\}_{n\geq 1}, q_0, \mathbb{P}_{0,n}, \mathbb{P}_{1,n}$ be as in Theorem [1.](#page-5-1) Under the assumptions of Theorem [2;](#page-5-2) for all $\varepsilon > 0$

$$
\lim_{n \to \infty} \mathbb{P}_{0,n} \left\{ N(H_n; G_n) \le (1 - \varepsilon) \mathbb{E}_0 N(H_n; G_n) \right\} = 0.
$$
 (C.4)

Proof. Let $v(H_n) = k_n, e(H_n) = e_n$. Let

$$
X_{\varphi}(G_n) = \begin{cases} 1 & \text{if } |\varphi(E(H_n)) \cap E(G_n)| = e_n, \\ 0 & \text{otherwise.} \end{cases}
$$

Note that $N(H_n; G_n) = \sum$ $\varphi{\in}\mathcal{L}(H_n;n)$ $X_{\varphi}(G_n)$. We have

$$
\mathbb{E}_0 X_{\varphi} = p_1(e_n) \equiv q_0^{e_n}.
$$

We write $\varphi_1 \sim \varphi_2$, if $|\varphi_1(V(H_n)) \cap \varphi_2(V(H_n))| \geq 2$. We define $e_G(m) = \max_{H \subseteq G, v(H) = [m]} e(H)$. Therefore, if $|\varphi_1(V(H_n)) \cap \varphi_2(V(H_n))| = u$, we have

$$
\mathbb{E}_0 X_{\varphi_1} X_{\varphi_2} \le p_2(u, e_n) \equiv q_0^{2e_n - e_{H_n}(u)}.
$$

Define

$$
\bar{\Delta}(n, H_n) = \sum_{\varphi_1 \sim \varphi_2} \mathbb{E}_0 X_{\varphi_1} X_{\varphi_2}.
$$
 (C.5)

Therefore, we have

$$
\bar{\Delta}(n, H_n) \le \sum_{u=2}^{k_n} \frac{(n)_{2k_n-u}(k_n!)^2}{u! \left((k_n-u)!\right)^2} p_2(u, e_n).
$$

Now using the fact that for all n

$$
\sqrt{2\pi}n^{n+1/2}e^{-n} \le n! \le n^{n+1/2}e^{-n+1},
$$

we get

$$
\frac{\bar{\Delta}(n, H_n)}{(\mathbb{E}_0 N(H_n; G_n))^2} \le \sum_{u=2}^{k_n} g(u).
$$

Where,

$$
g(u) = \begin{cases} \frac{(2\pi)^{-3/2}e^{-2k_n+2}k_n^{2k_n+1}}{(n-u)^u(k_n-u)^{2(k_n-u)+1}e^{-2(k_n-u)-u}u^{u+1/2}}q_0^{-e_{H_n}(u)} & \text{if } u \le k_n-1, \\ \frac{k_n^{kn+1/2}e^{-k_n+1}(n-k_n)^{n-k_n+1/2}e^{-n+k_n+1}}{(2\pi)^{1/2}n^{n+1/2}e^{-n}}q_0^{-e_n} & \text{if } u = k_n. \end{cases} \tag{C.6}
$$

Now using Chebyshev's inequality, for all $\varepsilon>0$

$$
\mathbb{P}_{0,n} \{ N(H_n; G_n) \le (1 - \varepsilon) \mathbb{E}_0 N(H_n; G_n) \} \le \frac{\mathbb{E}_0 \left(N(H_n; G_n)^2 \right) - (\mathbb{E}_0 N(H_n; G_n))^2}{\varepsilon^2 (\mathbb{E}_0 N(H_n; G_n))^2}
$$
\n
$$
= \frac{\sum_{\varphi_1, \varphi_2} (\mathbb{E}_0 X_{\varphi_1} X_{\varphi_2} - \mathbb{E}_0 X_{\varphi_1} \mathbb{E}_0 X_{\varphi_2})}{\varepsilon^2 (\mathbb{E}_0 N(H_n; G_n))^2}
$$
\n
$$
= \frac{\sum_{\varphi_1, \varphi_2} (\mathbb{E}_0 X_{\varphi_1} X_{\varphi_2} - \mathbb{E}_0 X_{\varphi_1} \mathbb{E}_0 X_{\varphi_2})}{\varepsilon^2 (\mathbb{E}_0 N(H_n; G_n))^2}
$$
\n
$$
\le \frac{\sum_{\varphi_1, \varphi_2} \mathbb{E}_0 X_{\varphi_1} X_{\varphi_2}}{\varepsilon^2 (\mathbb{E}_0 N(H_n; G_n))^2} = \frac{\bar{\Delta}(n, H_n)}{\varepsilon^2 (\mathbb{E}_0 N(H_n; G_n))^2}.
$$
\n(C.7)

Hence, in order to complete the proof it suffices to show that

$$
\frac{\bar{\Delta}(n, H_n)}{(\mathbb{E}_0 N(H_n; G_n))^2} \le \sum_{u=2}^{k_n} g(u) \to 0
$$
\n(C.8)

as $n\to\infty.$ Note that for $g(u)$ defined as in [\(C.6\)](#page-20-0), if $u < k_n$

$$
-\frac{1}{u}\log(g(u)) \ge \log n + \log(1 - (u/n)) - \frac{e_{H_n}(u)}{u}\log(1/q_0)
$$

+ 2(k_n/u) - 2((k_n/u) - 1) - 1 - (2(k_n/u) + (1/u))\log k_n
+ (2(k_n/u) - 2 + (1/u))\log(k_n - u) + (1 + \frac{1}{2u})\log u

$$
\ge \log n - \frac{e_{H_n}(u)}{u}\log(1/q_0) - 2\log k_n + \log u + \frac{1}{2u}\log u
$$

+ $\left(-2 + \frac{1}{u} + \frac{2k_n}{u}\right)\log\left(1 - \frac{u}{k_n}\right).$

In addition,

$$
-\frac{1}{k_n}\log(g(k_n)) \ge (n/k_n + 1/(2k_n))\log n - \frac{e_n}{k_n}\log(1/q_0)
$$

$$
- (1 + 1/(2k_n))\log k_n - (n/k_n - 1 + 1/(2k_n))\log(n - k_n) + \frac{C}{k_n}
$$

$$
= \log n - \frac{e_n}{k_n}\log(1/q_0) - \log k_n - \frac{1}{2k_n}\log k_n
$$

$$
- \left(\frac{n}{k_n} - 1 + \frac{1}{2k_n}\right)\log(1 - \frac{k_n}{n}) + \frac{C}{k_n}.
$$

Letting

$$
f(u) = \log n - \frac{e_{H_n}(u)}{u} \log(1/q_0) - 2\log k_n + \log u + \frac{1}{2u} \log u + \left(-2 + \frac{2k_n + 1}{u}\right) \log\left(1 - \frac{u}{k_n}\right),\tag{C.9}
$$

for $2 \le u \le k_n - 1$, and

$$
f(k_n) = \log n - \frac{e_{H_n}(u)}{u} \log(1/q_0) - \log k_n - \frac{1}{2k_n} \log k_n - \left(\frac{n}{k_n} - 1 + \frac{1}{2k_n}\right) \log(1 - \frac{k_n}{n}), \quad (C.10)
$$

Therefore, it suffices to show that

$$
\frac{\bar{\Delta}(n, H_n)}{(\mathbb{E}_0 N(H_n; G_n))^2} \le \sum_{u=2}^{k_n} g(u)
$$
\n
$$
\le C \sum_{u=2}^{k_n} \exp\{-uf(u)\}
$$
\n
$$
\le C k_n \exp\{-\tilde{u}f(\tilde{u})\}
$$
\n
$$
= C \exp\left\{-\tilde{u}\left(f(\tilde{u}) - \frac{\log k_n}{\tilde{u}}\right)\right\} \to 0
$$
\n(C.11)

as $n \to \infty$, where $\tilde{u} = \arg \min_{2 \le u \le k_n} \{uf(u)\}.$ First note that $((2/x) - 2) \log(1 - x) \ge -2$, for $0 \leq x < 1$. Hence,

$$
\left(-2+\frac{2k_n}{u}\right)\log\left(1-\frac{u}{k_n}\right)\geq -2,
$$

for $2 \le u \le k_n - 1$. Further, since $x \log(1 - 1/x)$ is increasing for $x > 1$, for $2 \le u \le k_n - 1$,

$$
\frac{1}{u}\log(1-u/k_n) \ge \frac{1}{k_n-1}\log(1-(k_n-1)/k_n) \ge -1,
$$

for large enough k_n . In addition, $\log u + (1/(2u)) \log u \ge 0$, for $u \ge 1$. Hence, the sum of last three terms in [\(C.9\)](#page-21-0) is bounded below. Finally, note that

$$
\label{eq:2.1} \begin{split} &\frac{n}{k_n}-1+\frac{1}{2k_n}\geq 0,\\ &-\frac{\log k_n}{2k_n}\geq -1, \end{split}
$$

for large enough n . Thus, the last two terms in $(C.10)$ are also bounded below. Therefore, for $2 \leq u \leq k_n$

$$
f(u) - \frac{\log k_n}{u} \ge \log n - \frac{e_{H_n}(u)}{u} \log(1/q_0) - (5/2) \log k_n + C,\tag{C.12}
$$

for some constant C . Hence,

$$
\frac{\bar{\Delta}(n, H_n)}{(\mathbb{E}_0 N(H_n; G_n))^2} \to 0
$$

as $n \to \infty$ if

$$
\limsup_{n \to \infty} \frac{d(H_n)\log(1/q_0) + (5/2)\log v(H_n)}{\log n} < 1.
$$

This shows that the lemma holds under the assumption of Theorem [2.](#page-5-2) Now, let $u^* = \arg\min_{2 \le u \le k_n} f(u)$. Note that as we had above,

$$
f(u) \ge \log n - \frac{e_{H_n}(u)}{u} \log(1/q_0) - 2 \log k_n + C.
$$

Hence, under the assumptions of Theorem [2,](#page-5-2) $f(u^*) \to \infty$. Define

$$
f^*(u) = \begin{cases} f(u) & \text{if, } u \le u^*, \\ f(u^*) & \text{otherwise.} \end{cases}
$$

We have

$$
\sum_{u=2}^{k_n} \exp \{-uf(u)\} \le \sum_{u=2}^{k_n} \exp \{-uf^*(u)\}
$$

=
$$
\sum_{u=2}^{u^*} \exp \{-uf(u)\} + C \exp \{-u^*f(u^*)\}
$$

$$
\le u^* \exp \{-\tilde{u}f(\tilde{u})\} + C \exp \{-u^*f(u^*)\}.
$$

Where $C = (1 - e^{-f(u^*)})^{-1}$ is a constant. Therefore, it suffices that

$$
u^* \exp\{-\tilde{u} f(\tilde{u})\} + C \exp\{-u^* f(u^*)\} \rightarrow 0
$$

as $n \to \infty$. This holds when

$$
u^* f(u^*) \to +\infty,
$$

$$
u^* \exp{-\tilde{u}f(\tilde{u})} \to 0.
$$
 (C.13)

as $n \to \infty$. Note that the first condition above holds since under the assumptions of Theorem [2,](#page-5-2) $f(u^*) \to \infty$ as $n \to \infty$. Further,

$$
\log(u^* \exp\{-\tilde{u}f(\tilde{u})\}) = \log u^* - \tilde{u}f(\tilde{u}).
$$

Note that if $d(H_n) = o(\log v(H_n))$, then $\limsup_{n \to \infty} (\log u^*)/(\tilde{u} \log n) = 0$. Thus, [\(C.13\)](#page-23-0) holds when $n \to \infty$. Therefore, if $d(H_n) = o(\log v(H_n))$ the lemma holds if

$$
\limsup_{n \to \infty} \frac{v(H_n)}{n^{1/2}} = 0
$$
\n(C.14)

and this completes the proof.

Now, we can prove Theorem [2.](#page-5-2)

Proof of Theorem [2.](#page-5-2) using Lemma [C.2,](#page-20-1) under the assumptions of Theorem [2,](#page-5-2) for all $\varepsilon > 0$

$$
\lim_{n \to \infty} \mathbb{P}_{0,n} \left\{ \frac{1}{\mathbb{E}_0 N(H_n; G_n)} N(H_n; G_n) \le (1 - \varepsilon) \right\} = 0.
$$

Therefore, by taking $Z_n = N(H_n; G_n)/\mathbb{E}_0 N(H_n; G_n)$, so that $\mathbb{E}_0 Z_n = 1$ we have

$$
\mathbb{E}_0[|1 - Z_n|] = 2\mathbb{E}_0[(1 - Z_n)_+] \to 0
$$

as $n \to \infty$. Therefore, under $\mathbb{P}_{0,n}$

$$
\frac{1}{\mathbb{E}_0 N(H_n; G_n)} N(H_n; G_n) \xrightarrow{p} 1
$$

and using [\(C.3\)](#page-19-0) and Lemma [C.1,](#page-18-1) Theorem [2](#page-5-2) is proved.

 \Box

D Proofs: spectral algorithm

We start by stating the following useful theorems from random matrix theory.

Theorem 8 ([\[Tao12\]](#page-15-12), Corollary 2.3.6). Let $X \in \mathbb{R}^{n \times n}$ be a random symmetric matrix whose entries X_{ij} are independent, zero-mean, uniformly bounded random variables for $j \geq i$ and $X_{ij} = X_{ji}$ for $j < i$. There exists constants $c_1, c_2 > 0$ such that for all $t \geq c_1$

$$
\mathbb{P}\left\{\|\boldsymbol{X}\|_2 > t\sqrt{n}\right\} \leq c_1 \exp\left(-c_2 t n\right).
$$

Theorem 9 ([\[Tao12\]](#page-15-12), Theorem 2.3.24). Let $X \in \mathbb{R}^{n \times n}$ be a random symmetric matrix whose entries X_{ij} are i.i.d copies of a zero-mean random variable with variance 1 and finite fourth moment for $j \geq i$ and $X_{ij} = X_{ji}$ for $j < i$. Then, $\lim_{n \to \infty} ||\boldsymbol{X}||_2/\sqrt{n} = 2$, almost surely.

D.1 Proof of Theorem [3](#page-6-4)

First assume that G_n is generated according to the null model $\mathbb{P}_{0,n}$. Then, $\mathcal{A}_G^{q_0}$ $\frac{q_0}{G_n}$ is a random symmetric matrix with independent entries where each entry is a zero-mean Bernoulli random variable which is equal to 1 with probability q_0 and $-q_0/(1-q_0)$ with probability $1-q_0$. Using Theorem [9,](#page-24-0) $\lambda_1(\boldsymbol{A}_{G}^{q_0})$ $\mathcal{G}_{G_n}^{(0)} \leq 2.1\sigma(q_0)\sqrt{n}$ with high probability as $n \to \infty$. Therefore, $\limsup_{n \to \infty} \mathbb{P}_{0,n}(\mathcal{T}_{\text{spec}}(G_n))$ 1) = 0. Now assume that G_n is generated according to the planted model, $\mathbb{P}_{1,n}$, with parameters q_0 and H_n . Hence, $\mathbf{A}_G^{q_0}$ $_{G_n}^{q_0}$ is distributed as $\prod_n^{\mathsf{T}} A_{H_n} \prod_n^{\mathsf{T}} A_H$ where $\prod_n \in \{0,1\}^{v(H_n)\times n}$, and $(\prod_n)_{ij} = 1$ if and only if $\varphi_{0,n}(\tilde{i}) = j$. Further, \mathbf{E}_n is a random symmetric matrix with independent entries where $(E_n)_{i,j} = 0$ if $(\Pi_n^{\mathsf{T}} A_{H_n} \Pi_n)_{i,j} = 1$ and $(E_n)_{i,j}$ is a zero mean Bernoulli random variable which is equal to 1 with probability q_0 and $-q_0/(1 - q_0)$ with probability $1 - q_0$, otherwise. Let $\boldsymbol{v}, \|\boldsymbol{v}\|_2 = 1$ be the principal eigenvector of \boldsymbol{A}_{H_n} . We have

$$
\begin{aligned} \lambda_1(\boldsymbol{A}_{G_n}^{q_0}) &\geq \Big\langle \boldsymbol{\Pi}_n^{\mathsf{T}} \boldsymbol{v}, \boldsymbol{A}_{G_n}^{q_0} \boldsymbol{\Pi}_n^{\mathsf{T}} \boldsymbol{v} \Big\rangle = \Big\langle \boldsymbol{\Pi}_n^{\mathsf{T}} \boldsymbol{v}, \boldsymbol{\Pi}_n^{\mathsf{T}} \boldsymbol{A}_{H_n} \boldsymbol{\Pi}_n \boldsymbol{\Pi}_n^{\mathsf{T}} \boldsymbol{v} \Big\rangle + \Big\langle \boldsymbol{\Pi}_n^{\mathsf{T}} \boldsymbol{v}, \boldsymbol{E}_n \boldsymbol{\Pi}_n^{\mathsf{T}} \boldsymbol{v} \Big\rangle \\ & = \langle \boldsymbol{v}, \boldsymbol{A}_{H_n} \boldsymbol{v} \rangle + \Big\langle \boldsymbol{v}, \boldsymbol{\Pi}_n \boldsymbol{E}_n \boldsymbol{\Pi}_n^{\mathsf{T}} \boldsymbol{v} \Big\rangle \end{aligned}
$$

Therefore,

$$
\begin{aligned} \lim\inf_{n\to\infty}\frac{\lambda_1(\boldsymbol{A}_{G_n}^{q_0})}{\sqrt{n}} &\geq \lim\inf_{n\to\infty}\frac{\lambda_1(\boldsymbol{A}_{H_n})}{\sqrt{n}} - \limsup_{n\to\infty}\frac{\left\langle \boldsymbol{v},\boldsymbol{\Pi}_n\boldsymbol{E}_n\boldsymbol{\Pi}_n^\mathsf{T}\boldsymbol{v}\right\rangle}{\sqrt{n}} \\ &\geq 3\sigma(q_0) - \lim\sup_{n\to\infty}\frac{\lambda_1(\boldsymbol{\Pi}_n\boldsymbol{E}_n\boldsymbol{\Pi}_n^\mathsf{T})}{\sqrt{n}}. \end{aligned}
$$

Now, using Theorem [8,](#page-24-1) $\lambda_1(\Pi_n E_n \Pi_n^{\mathsf{T}}) \leq c\sqrt{v(H_n)}$, for some c, and large enough n, almost surely. Therefore, $\limsup_{n\to\infty} \frac{\lambda_1(\mathbf{\Pi}_n \mathbf{E}_n \mathbf{\Pi}_n^{\mathsf{T}}) \leq C_V v(T_n)}{\lambda_1(\mathbf{\Pi}_n \mathbf{E}_n \mathbf{\Pi}_n^{\mathsf{T}})/\sqrt{n}} = 0$ and under the alternative,

$$
\lim \inf_{n \to \infty} \frac{\lambda_1(A_{G_n}^{q_0})}{\sqrt{n}} \ge 2.1\sigma(q_0),
$$

almost surely. Hence, $\limsup_{n\to\infty} \mathbb{P}_{1,n}(T_{\text{spec}}(G_n) = 0) = 0$ and two models are strongly distinguishable using the spectral test.

D.2 Proof of Theorem [4](#page-8-1)

We start by proving some useful lemmas.

Lemma D.1. Let $A = \Pi^{\mathsf{T}} A_{H_n} \Pi + E + \tilde{E}$ where A is a symmetric n by n matrix, $\Pi \in \{0,1\}^{k \times n}$, $\Pi\Pi^{\mathsf{T}} = \mathbf{I}_k$, $\Pi\mathbf{1} = \mathbf{1}$, \boldsymbol{A}_{H_n} is k by k symmetric matrix, $k = o(n)$. Further, let \boldsymbol{E} be a random symmetric matrix with independent entries where each entry is a zero-mean Bernoulli random variable which is equal to 1 with probability p and $-p/(1-p)$ with probability $1-p$. Finally, $\tilde{E}_{i,j} = -E_{i,j}$ if $(\boldsymbol{\Pi}^\mathsf{T} \boldsymbol{A}_{H_n} \boldsymbol{\Pi})_{i,j} = 1$ and $\tilde{E}_{i,j} = 0$, otherwise. Let $\boldsymbol{v} \in \mathbb{R}^n, \boldsymbol{x} \in \mathbb{R}^k$, $\|\boldsymbol{v}\|_2 = \|\boldsymbol{x}\|_2 = 1$, be the leading eigenvectors of **A** and A_{H_n} , respectively. Assume that for some $\delta \in (0,1)$,

$$
\lambda_1({\boldsymbol{A}}_{H_n}) \geq \frac{3}{\varepsilon\delta}\sqrt{\frac{np}{1-p}}
$$

then $v = \alpha \Pi^{\mathsf{T}} x + z$ for some α such that $\alpha^2 \geq 1-\delta$ and $||z||_2^2 \leq \delta$, with high probability as $n \to \infty$.

Proof. Let $S \subseteq [n]$ be the set of i's for which the i'th column of Π is not entirely zero. We denote the complement of this set by \bar{S} . We can write $v = \Pi^{\mathsf{T}}(\alpha x + \beta y) + v_{\bar{S}} = \alpha \Pi^{\mathsf{T}} x + z$, where $y \in \mathbb{R}^k$ is such that $y \perp x$ and $\alpha^2 + \beta^2 = ||v_S||_2^2$. In addition, note that $z \perp \Pi^{\top} x$. Hence, $||z||_2^2 + \alpha^2 ||\mathbf{\Pi}^\mathsf{T} x||_2^2 = ||\mathbf{v}||_2^2 = 1.$ Thus, $||z||_2^2 = 1 - \alpha^2 ||\mathbf{\Pi}^\mathsf{T} x||_2^2 = 1 - \alpha^2$. Now, if $\alpha^2 < 1 - \delta$, then

$$
\langle {\boldsymbol v},{\boldsymbol A}{\boldsymbol v}\rangle = \alpha^2\, \langle {\boldsymbol x},{\boldsymbol A}_{H_n}{\boldsymbol x}\rangle + 2\alpha\beta\, \langle {\boldsymbol x},{\boldsymbol A}_{H_n}{\boldsymbol y}\rangle + \beta^2\, \langle {\boldsymbol y},{\boldsymbol A}_{H_n}{\boldsymbol y}\rangle + \langle {\boldsymbol v},{\boldsymbol E}{\boldsymbol v}\rangle + \langle {\boldsymbol v},{\tilde{\boldsymbol E}}{\boldsymbol v}\rangle.
$$

Since x is an eigenvector of A_{H_n} and $x \perp y$, we have $\langle x, A_{H_n} y \rangle = 0$. Now, using Theorems [8,](#page-24-1) [9,](#page-24-0) with high probability as $n \to \infty$,

$$
\langle v, \mathbf{A}v \rangle \leq \alpha^2 \lambda_1(\mathbf{A}_{H_n}) + (\|\mathbf{v}_S\|_2^2 - \alpha^2)(1 - \varepsilon)\lambda_1(\mathbf{A}_{H_n}) + (2 + o(1))\sqrt{\frac{np}{1 - p}} + c\sqrt{k}
$$

$$
\leq (1 - \delta)\lambda_1(\mathbf{A}_{H_n}) + \delta(1 - \varepsilon)\lambda_1(\mathbf{A}_{H_n}) + (2 + o(1))\sqrt{\frac{np}{1 - p}} + c\sqrt{k},
$$

and

$$
\langle \Pi^{\mathsf{T}} x, A \Pi^{\mathsf{T}} x \rangle = \langle x, \Pi A \Pi^{\mathsf{T}} x \rangle = \langle x, A_{H_n} x \rangle + \langle x, \Pi (E + \tilde{E}) \Pi^{\mathsf{T}} x \rangle \geq \lambda_1 (A_{H_n}) - c' \sqrt{k}.
$$
 (D.1)

Therefore, if $\lambda_1(A_{H_n}) \geq \frac{3}{\varepsilon \delta} \sqrt{\frac{np}{1-p}}$, then $\langle v, Av \rangle < \langle \Pi^{\mathsf{T}} x, A \Pi^{\mathsf{T}} x \rangle$ with high probability as $n \to \infty$. Hence, if $\alpha^2 < 1 - \delta$, v cannot be the leading eigenvector of **A** and the lemma is proved. \Box

The following lemma is an immediate consequence of the above lemma.

Lemma D.2. Let $\{H_n\}_{n>1}$ be a sequence of graphs that are (ε, μ) -balanced in spectrum for some $\mu > 0, \varepsilon \in (0, 1)$. Further, let $\varphi_{0,n} \in \mathcal{L}(H_n, n)$ be a labeling of H_n vertices in $[n], v(H_n) = o(n)$. Suppose that G_n is generated according to $\mathbb{P}_{1,n}(\cdot | \varphi = \varphi_0)$ as in Eq. [\(1.3\)](#page-1-0). Take v to be the leading eigenvector of $A_G^{q_0}$ $\mathcal{C}_{G_n}^{q_0}$. Let $|v_{j(1)}| \ge |v_{j(2)}| \ge \cdots \ge |v_{j(n)}|$ be the entries of \boldsymbol{v} and $S' =$ $\{j(1), j(2), \ldots, j(v(H_n))\}$. If

$$
\lambda_1(\boldsymbol{A}_{H_n}) \geq \frac{3}{\varepsilon \delta} \sqrt{\frac{nq_0}{1-q_0}},
$$

then

$$
|S' \cap \varphi_0(V(H_n))| \ge \left(1 - \frac{2\delta}{\mu^2(1-\delta)}\right)v(H_n),
$$

with high probability as $n \to \infty$.

Proof. Note that $A_G^{q_0}$ $_{G_n}^{q_0}$ is distributed as $\mathbf{\Pi}_n^{\mathsf{T}} \mathbf{A}_{H_n} \mathbf{\Pi}_n + \mathbf{E}_n + \tilde{\mathbf{E}}_n$ where $\mathbf{\Pi}_n \in \{0,1\}^{v(H_n)\times n}$, and $(\mathbf{\Pi}_n)_{ij} = 1$ if and only if $\varphi_{0,n}(i) = j$. Further, \mathbf{E}_n is a random symmetric matrix with independent entries where each entry is a zero-mean Bernoulli random variable which is equal to 1 with probability q_0 and $-q_0/(1-q_0)$ with probability $1-q_0$. Finally, $(\tilde{\mathbf{E}}_n)_{i,j} = -(\mathbf{E}_n)_{i,j}$ if $(\mathbf{\Pi}^\mathsf{T} \mathbf{A}_{H_n} \mathbf{\Pi})_{i,j} = 1$ and $(\tilde{\mathbf{E}}_n)_{i,j} = 0$, otherwise. Hence, defining x to be the leading eigenvector of \mathbf{A}_{H_n} , using Lemma [D.1,](#page-25-0) $\mathbf{v} = \tilde{\mathbf{x}} + \mathbf{z}, \ \tilde{\mathbf{x}} = \mathbf{\Pi}_{n}^{\mathsf{T}} \mathbf{x}, \ \|\mathbf{x}\|^{2} \geq 1 - \delta$ and $\mathbf{z} \perp \tilde{\mathbf{x}}$, with high probability. Let $S = \varphi_{0}(V(H_{n}))$, using the assumption that H_n is (ε, μ) -balanced in spectrum, for $i \in S$, $|\tilde{x}_i| \geq \mu \sqrt{1-\delta}/\sqrt{v(H_n)}$. Note that for $i \notin S$, $\tilde{x}_i = 0$. Therefore, for any $i \in (\bar{S} \cap S')$, there exists an index $i' \in (\bar{S'} \cap S)$ such that $z_i^2 + z_{i'}^2 \geq 2(\mu)$ $\sqrt{1-\delta}/(2\sqrt{v(H_n)})^2$. Hence, letting N be the number of indices in S' which are not in S, we have

$$
2N\left(\frac{\mu\sqrt{1-\delta}}{2\sqrt{v(H_n)}}\right)^2\leq \|{\boldsymbol{z}}\|_2^2=1-\|{\boldsymbol{x}}\|_2^2\leq \delta.
$$

Therefore, $N \leq 2\delta v(H_n)/(\mu^2(1-\delta))$ and $|S' \cap \varphi_0(V(H_n))| \geq \left(1 - \frac{2\delta}{\mu^2(1-\delta)}\right)v(H_n)$ with high probability as $n \to \infty$. \Box

Now we prove Theorem [4.](#page-8-1)

Proof of Theorem [4.](#page-8-1) First assume that $i \notin \varphi_0(V(H_n))$. Recall that $d^{(i)}$ is the number of edges between vertex *i* and vertices in S_i . Further S_i only depends on the edges induced by $V \setminus \{i\}$. Hence, we have

$$
d^{(i)} = \sum_{j=1}^{v(H_n)} X_j
$$

where $\{X_j\}$ is a sequence of i.i.d Bern (q_0) random variables. Therefore using Bernstein's inequality

$$
\sum_{i \notin \varphi_0(V(H_n))} \mathbb{P}\left\{d^{(i)} > t\right\} \le n \mathbb{P}\left\{d^{(i)} > t\right\}
$$
\n
$$
\le n \exp\left\{-\frac{(1/2)(t - v(H_n)q_0)^2}{(1/3)(t - v(H_n)q_0) + v(H_n)q_0(1 - q_0)}\right\}
$$
\n
$$
\le n \exp\left\{-\frac{(9/2)v(H_n)q_0 \log v(H_n)}{\sqrt{v(H_n)q_0 \log v(H_n) + v(H_n)q_0(1 - q_0)}}\right\}
$$
\n
$$
\le \exp\left\{\log n - 4\log v(H_n)\right\},
$$

for large enough n. Using [\(3.11\)](#page-9-2), this goes to zero as $n \to \infty$. Therefore, using union bound, the output set S of Algorithm [1](#page-7-0) will be a subset of $\varphi_0(V(H_n))$.

Next, assume that $i = \varphi_0(\tilde{i})$ and $\tilde{i} \in S_c(H_n)$ where c is as in Theorem [4.](#page-8-1) Using Lemma [D.2,](#page-25-1) $|S_i \cap \varphi_0(V(H_n))| \geq (1-\alpha)v(H_n)$. Also, note that S_i only depends on the edges induced by $V \setminus \{i\}$. Therefore, $d^{(i)}$ dominates d' , where

$$
d' = (c - \alpha)v(H_n) + \sum_{j=1}^{(1-c+\alpha)v(H_n)} X_j,
$$

in which $\{X_j\}$ is a sequence of i.i.d Bern (q_0) random variables. Note that here $\mathbb{E}d' = v(H_n)q_0 +$ $(1 - q_0)(c - \alpha)v(H_n)$. Hence, using Bernstein's inequality we get

$$
\sum_{i \in \varphi_0(S_c(H_n))} \mathbb{P}\left\{d^{(i)} \le t\right\} \le v(H_n)\mathbb{P}\left\{d' \le t\right\}
$$
\n
$$
\le v(H_n) \exp\left\{-\frac{(1/2)(v(H_n)(q_0 + (1-q_0)(c-\alpha)) - t)^2}{(1/3)t + q_0(1-q_0)(1-c+\alpha)v(H_n)}\right\}
$$
\n
$$
\le v(H_n) \exp\left\{-\frac{(1/4)(1-q_0)^2(c-\alpha)^2v(H_n)^2}{(1/3)(1-c+\alpha)v(H_n) + q_0(1-q_0)(1-c+\alpha)v(H_n)}\right\}
$$
\n
$$
\le v(H_n) \exp\left\{-C'v(H_n)\right\} \to 0,
$$

as $n, v(H_n) \to \infty$. Thus, by union bound, the output of Algorithm [1,](#page-7-0) contains all the nodes in the c-significant set of planted subgraph H_n in G_n and has no nodes which are not in the planted subgraph H_n , with high probability, as $n, v(H_n) \to \infty$. \Box

E Proofs: SDP relaxation

For simplicity we denote $v(H_n)$ by k_n . We start by proving Lemma [3.4.](#page-10-3)

E.1 Proof of Lemma [3.4](#page-10-3)

First note that every feasible Π in [\(3.13\)](#page-9-1), corresponds uniquely to an injective mapping φ from $[k]$ to [n] where $\varphi(i) = j$ if and only if $\Pi_{ji} = 1$. Based on this, we have

$$
\mathsf{Tr}\left(\boldsymbol{A}_H\boldsymbol{\Pi}^{\mathsf{T}}\boldsymbol{A}_G\boldsymbol{\Pi}\right)=\sum_{i,j=1}^k\left(\boldsymbol{A}_H\right)_{ij}\left(\boldsymbol{A}_G\right)_{\varphi(i)\varphi(j)}=\mathsf{Tr}\left((\boldsymbol{A}_G\otimes \boldsymbol{A}_H)\boldsymbol{Y}\right).
$$

where $Y = yy^{\mathsf{T}}$ and $y = \text{vec}(\mathbf{\Pi})$. Moreover Y is a rank one positive definite matrix in $\{0,1\}^{nk \times nk}$. Also,

$$
\text{Tr}(\mathbf{Y}\mathbf{J}_{nk}) = \sum_{i,j=1}^{nk} Y_{ij} = \left(\sum_{i=1}^{nk} y_i\right)^2 = k^2.
$$
 (E.1)

In addition, for $i = 1, 2, \ldots, k$

$$
\operatorname{Tr}(\mathbf{Y}(\mathbf{I}_n \otimes (\boldsymbol{e}_i \boldsymbol{e}_i^{\mathsf{T}}))) = \sum_{l=0}^{n-1} y_{kl+i} = \sum_{j=1}^n \Pi_{ji} = 1.
$$
 (E.2)

Further, for $j = 1, 2, \ldots, n$

$$
\text{Tr}(\mathbf{Y}((e_j e_j^{\mathsf{T}}) \otimes \mathbf{I}_k)) = \sum_{l=0}^{k-1} y_{(j-1)k+l} = \sum_{i=1}^k \Pi_{ji} \le 1.
$$
 (E.3)

Therefore, Y is feasible for problem (3.17) . Conversely, if Y is feasible for problem (3.17) , then $\boldsymbol{Y} = \boldsymbol{y} \boldsymbol{y}^{\mathsf{T}}$ where $\boldsymbol{y} \in \{0,1\}^{nk}$. Also, using [\(E.1\)](#page-27-0), \boldsymbol{y} has exactly k entries equal to one and $n - k$ entries equal to zero. Further, using the first equality in $(E.2)$, we deduce that for $i = 1, 2, \ldots, k$,

$$
\sum_{l=0}^{n-1} y_{kl+i} = 1.
$$

Also, using the first equality in $(E.3)$ for $j = 1, 2, \ldots, n$,

$$
\sum_{l=0}^{k-1} y_{(j-1)k+l} \leq 1.
$$

This means that the matrix $\Pi \in \{0,1\}^{n \times k}$ whose j'th row is $[y_{(j-1)k+1}, y_{(j-1)k+2}, \ldots, y_{jk}]$ has exactly one entry equal to one in each column. Therefore, Π is feasible for problem equation [\(3.13\)](#page-9-1). This completes the proof of Lemma [3.4.](#page-10-3)

E.2 Proof of Theorem [5](#page-10-5)

The following lemma about the spectrum of a random Erdős-Rényi graph is a consequence of Theorem [9.](#page-24-0)

Lemma E.1. Let $A \in \{0,1\}^{n \times n}$ be a random matrix with independent entries such that

$$
A_{ij} = \begin{cases} 1 & with \ probability \ p_{ij} \\ 0 & with \ probability \ 1 - p_{ij}, \end{cases}
$$

where $p_{ij} = p$ if $i \neq j$, $p_{ii} = 0$ and $p \in (0, 1)$ is a constant. Then,

$$
\lim_{n \to \infty} \frac{\lambda_1(\mathbf{A})}{np} = 1,
$$

$$
\limsup_{n \to \infty} \frac{-\lambda_n(\mathbf{A})}{2\sqrt{np(1-p)}} = 1,
$$

almost surely.

Lemma E.2. Let $A \in \{0,1\}^{n \times n}$ be a random matrix with independent entries such that

$$
A_{ij} = \begin{cases} 1 & with \ probability \ p_{ij} \\ 0 & with \ probability \ 1 - p_{ij}, \end{cases}
$$

where $p_{ij} = p$ if $i \neq j$, $p_{ii} = 0$ and $p \in (0, 1)$ is a constant. Let **D** be a n by n diagonal matrix such that $D_{ii} = \sum_{i=1}^{n}$ $j=1$ A_{ij} , $\boldsymbol{L} = \boldsymbol{D} - \boldsymbol{A}$. Then,

- (i) $\mathbf{L} \succeq 0$.
- (*ii*) $L1_n = 0$.
- (iii) if $\lambda_2(L)$ is the second smallest eigenvalue of L then,

$$
\lambda_2(\mathbf{L}) = np - 2(1 + o(1))\sqrt{np(1 - p)}
$$

almost surely, as $n \to \infty$.

Proof. The proof of points (i), (ii) in Lemma [E.2](#page-28-1) is standard and can be found in $[Chu97]$. Further, we can write $\mathbf{L} = \mathbb{E}\mathbf{L} + (\mathbf{D} - \mathbb{E}\mathbf{D}) - (\mathbf{A} - \mathbb{E}\mathbf{A})$ where $\mathbb{E}\mathbf{L} = np\mathbf{I}_n - p\mathbf{J}_n$ and $\lambda_2(\mathbb{E}\mathbf{L}) = \lambda_3(\mathbb{E}\mathbf{L}) =$ $\cdots = \lambda_n (EL) = np$. Therefore, point (iii) follows from this together with the bound on the spectra of $A - \mathbb{E}A$, $D - \mathbb{E}D$. \Box

Now we can state the proof of Theorem [5.](#page-10-5)

Proof of Theorem [5.](#page-10-5) Using the fact that for any graph G with adjacency matrix $A_G \in \{0,1\}^{n \times n}$, $\lambda_1(A_G) \geq -\lambda_n(A_G)$, it suffices to prove the theorem assuming that

$$
\limsup_{n\to\infty}\frac{2(\lambda_1(\boldsymbol{A}_{H_n})-\lambda_{k_n}(\boldsymbol{A}_{H_n}))\sqrt{1-q_0}}{\sqrt{nq_0}}=1-C<1.
$$

Assume that G_n is generated randomly according to $\mathbb{P}_{0,n}$. Let $\text{SDP}(G_n; H_n)$ be the sequence of the optimal values of the (random) convex programs [\(3.17\)](#page-10-4). Let D_{G_n} be a $n \times n$ diagonal matrix such that $(D_{G_n})_{ii} = \deg(i)$. In order to prove Theorem [5,](#page-10-5) we have to show that $\mathsf{SDP}(G_n; H_n) \geq 2e(H_n)$. In order to do this, we construct a sequence of matrices Y_n which are feasible for problem [\(3.17\)](#page-10-4) and $\text{Tr}((A_{G_n} \otimes A_{H_n})\mathbf{Y}_n) \geq 2e(H_n)$, with high probability as $n \to \infty$. We take

$$
\mathbf{Y}_n = \begin{cases} a_n(\mathbf{D}_{G_n} \otimes \mathbf{I}_{k_n}) + b_n(\mathbf{A}_{G_n} \otimes (\mathbf{A}_{H_n} + \mathbf{I}_{k_n})) + c_n \mathbf{J}_{nk_n} & \text{If } \lambda_{k_n}(\mathbf{A}_{H_n}) \ge (2e(H_n) - k_n^2)/k_n, \\ b_n(\mathbf{D}_{G_n} - \mathbf{A}_{G_n}) \otimes \mathbf{I}_{k_n} + b_n \mathbf{A}_{G_n} \otimes \mathbf{J}_{k_n} & \text{otherwise,} \end{cases}
$$
(E.4)

where

$$
u_n = \left[-\lambda_{k_n}(\mathbf{A}_{H_n}) - 1 + \frac{k_n \lambda_{k_n}(\mathbf{A}_{H_n}) + k_n^2 - 2e(H_n)}{nk_n^2} \right]_+,
$$

\n
$$
a_n = \frac{2e(H_n) + k_n u_n + nk_n^2 - k_n^2}{2k_n e(G_n)(nk_n - 1)},
$$

\n
$$
b_n = \frac{1}{2e(G_n)},
$$

\n
$$
c_n = \frac{k_n(k_n - 1) - 2e(H_n) - k_n u_n}{n^2 k_n^2 - nk_n}.
$$

Now, we show that Y_n is feasible for problem [\(3.17\)](#page-10-4). First, consider the case where $\lambda_{k_n}(A_{H_n}) \geq$ $\left(\frac{2e(H_n) - k_n^2}{k_n} \right)$. In this case,

$$
u_n \le k_n - 1 - 2e(H_n)/k_n \le k_n. \tag{E.5}
$$

Hence, $c_n \ge 0$. Also, $a_n \ge 0$ and $b_n \ge 0$. Thus, $\boldsymbol{Y}_n \ge 0$, entrywise. In addition,

$$
\max_{i \in V(G_n)} \deg(i) < 2nq_0,\tag{E.6}
$$

$$
e(G_n) > q_0 n^2/4,\tag{E.7}
$$

with high probability as $n \to \infty$. Thus, using the fact that, $2e(H_n) \leq k_n^2$, $n + 1 \leq 2n$ and for large enough $n, nk_n - 1 \geq nk_n/2, n^2k_n^2 - nk_n \geq n^2k_n^2/2,$

$$
a_n \left(\max_{i \in V(G_n)} \deg(i) \right) + c_n \le \frac{2nq_0(nk_n^2 + k_n u_n)}{(n^2/2)k_n q_0(nk_n/2)} + \frac{(k_n^2 - k_n) - k_n u_n}{n^2 k_n^2/2}
$$

$$
\le \frac{2n^2 k_n^2 q_0}{n^3 k_n^2 q_0/4} + \frac{2nq_0 k_n u_n}{n^3 k_n^2 q_0/4} + \frac{k_n^2}{n^2 k_n^2/2}
$$

$$
= \frac{16}{n} + \frac{8u_n}{n^2 k_n} + \frac{2}{n^2},
$$
 (E.8)

which is less than 1 for large enough n. Also, similarly, using $(E.7)$,

$$
b_n + c_n \le \frac{1}{(q_0 n^2)/2} + \frac{k_n^2}{n^2 k_n^2/2}
$$

= $\frac{2}{q_0 n^2} + \frac{2}{n^2} \le 1,$ (E.9)

for large enough n . Finally, using $(E.5)$, $(E.7)$,

$$
u_n b_n + c_n \le \frac{k_n}{(q_0 n^2)/2} + \frac{k_n^2}{n^2 k_n^2/2}
$$

=
$$
\frac{2k_n}{q_0 n^2} + \frac{2}{n^2} \le 1
$$
 (E.10)

for large enough n. Therefore, according to the construction of Y_n as in [\(E.4\)](#page-29-1), using equations $(E.8), (E.9), (E.10)$ for large enough n, $\mathbf{Y}_n \leq 1$, entrywise. Also,

$$
\text{Tr}(\mathbf{Y}_n \mathbf{J}_{nk_n}) = 2a_n e(G_n) k_n + 4b_n e(H_n) e(G_n) + 2b_n k_n u_n e(G_n) + c_n n^2 k_n^2 = k_n^2.
$$

Moreover, for $i = 1, 2, \ldots, k_n$,

$$
\mathsf{Tr}(\boldsymbol{Y}_n(\mathbf{I}_n \otimes (\boldsymbol{e}_i \boldsymbol{e}_i^{\mathsf{T}}))) = 2e(G_n)a_n + nc_n = 1.
$$

Finally, for $j = 1, 2, \ldots, n$,

$$
\operatorname{Tr}(\mathbf{Y}_n((e_j e_j^{\mathsf{T}}) \otimes \mathbf{I}_{k_n})) \le k_n a_n \left(\max_{i \in V(G_n)} \deg(i) \right) + k_n c_n
$$

$$
\le \frac{16k_n}{n} + \frac{8u_n}{n^2} + \frac{2k_n}{n^2} \le 1,
$$
 (E.11)

for large enough n . Second inequality in $(E.11)$ is by $(E.8)$. Next, we consider the case in which $\lambda_{k_n}(\mathbf{A}_{H_n}) \geq (2e(H_n) - k_n^2)/k_n$. In this case, since

$$
\frac{\max_{i \in V(G_n)} \deg(i)}{2e(G_n)} \le \frac{2nq_0}{(n^2q_0/2)} \le 1
$$

with high probability, as $n \to \infty$, $0 \le Y_n \le 1$ entrywise. Further, for $i = 1, 2, ..., k_n$ and $j = 1, 2, \ldots, n,$

$$
\mathrm{Tr}(\mathbf{Y}_n \mathbf{J}_{nk_n}) = 2b_n k_n^2 e(G_n) = k_n^2,
$$
\n
$$
\mathrm{Tr}(\mathbf{Y}(\mathbf{I}_n \otimes (\mathbf{e}_i \mathbf{e}_i^\mathsf{T}))) = 2e(G_n)b_n = 1,
$$
\n
$$
\mathrm{Tr}(\mathbf{Y}_n((\mathbf{e}_j \mathbf{e}_j^\mathsf{T}) \otimes \mathbf{I}_{k_n})) \le k_n b_n \left(\max_{i \in V(G_n)} \deg(i)\right) \le \frac{2k_n n q_0}{(n^2 q_0/2)} = \frac{4k_n}{n} \le 1,
$$

for large enough n, with high probability. Finally, we have to show that the proposed Y_n is positive semidefinite with high probability. In order to show this it is sufficient to show that

$$
\tilde{\boldsymbol{Y}}_n = 2e(G_n)\tilde{a}_n(\boldsymbol{D}_{G_n} - \boldsymbol{A}_{G_n}) \otimes \mathbf{I}_{k_n} + \boldsymbol{A}_{G_n} \otimes \tilde{\boldsymbol{A}}_{H_n} \succeq 0,
$$

where

$$
\tilde{\mathbf{A}}_{H_n} = \begin{cases} \mathbf{A}_{H_n} + (u_n + 2e(G_n)a_n)\mathbf{I}_{k_n}, & \text{if } \lambda_{k_n}(\mathbf{A}_{H_n}) \ge (2e(H_n) - k_n^2)/k_n, \\ \mathbf{J}_{k_n} & \text{otherwise,} \end{cases}
$$

and

$$
\tilde{a}_n = \begin{cases} a_n, & \text{if } \lambda_{k_n}(\mathbf{A}_{H_n}) \ge (2e(H_n) - k_n^2)/k_n, \\ b_n & \text{otherwise.} \end{cases}
$$

If
$$
\lambda_{k_n}(\mathbf{A}_{H_n}) < (2e(H_n) - k_n^2)/k_n
$$
, then $\tilde{\mathbf{A}}_{H_n} = \mathbf{J}_{k_n} \succeq 0$. Otherwise,
\n
$$
\lambda_{k_n}(\tilde{\mathbf{A}}_{H_n}) = \lambda_{k_n}(\mathbf{A}_{H_n}) + u_n + 2a_n e(G_n)
$$
\n
$$
\geq \lambda_{k_n}(\tilde{\mathbf{A}}_{H_n}) - \lambda_{k_n}(\tilde{\mathbf{A}}_{H_n}) - 1 + \frac{k_n \lambda_{k_n}(\mathbf{A}_{H_n}) + k_n^2 - 2e(H_n)}{nk_n^2}
$$
\n
$$
+ \frac{2e(H_n) + k_n u_n + nk_n^2 - k_n^2}{k_n(nk_n - 1)}
$$
\n
$$
= \frac{1}{nk_n - 1} \left(u + \lambda_{k_n}(\tilde{\mathbf{A}}_{H_n}) + 1 - \frac{k \lambda_{k_n}(\tilde{\mathbf{A}}_{H_n}) + k_n^2 - 2e(H_n)}{nk_n^2} \right) \geq 0.
$$

Therefore, $\tilde{\boldsymbol{A}}_{H_n}$ is positive semidefinite in both cases. Let z be an arbitrary vector in \mathbb{R}^{nk_n} . We can write

$$
z=z_\parallel+z_\perp
$$

where

$$
\begin{aligned} \boldsymbol{z}_{\parallel} &= \mathbf{1}_n \otimes \boldsymbol{x}, \qquad \boldsymbol{x} \in \mathbb{R}^k, \\ \boldsymbol{z}_{\perp} &= \begin{bmatrix} \boldsymbol{y}_1 \\ \boldsymbol{y}_2 \\ \vdots \\ \boldsymbol{y}_{n-1} \\ -\boldsymbol{y}_1 - \boldsymbol{y}_2 - \cdots - \boldsymbol{y}_{n-1} \end{bmatrix}, \qquad \boldsymbol{y}_i \in \mathbb{R}^k. \end{aligned}
$$

Using Lemma [E.2,](#page-28-1) z_{\parallel} is in the nullspace of $(D_{G_n} - A_{G_n}) \otimes I_{k_n}$. Therefore,

$$
\left\langle z, \tilde{\boldsymbol{Y}}_n z \right\rangle = \left\langle z_\parallel, \left(A_{G_n} \otimes \tilde{A}_{H_n} \right) z_\parallel \right\rangle \\ + 2 \left\langle z_\perp, \left(A_{G_n} \otimes \tilde{A}_{H_n} \right) z_\parallel \right\rangle \\ + \left\langle z_\perp, \left(2e(G_n) \tilde{a}_n (\boldsymbol{D}_{G_n} - A_{G_n}) \otimes \mathbf{I}_{k_n} + A_{G_n} \otimes \tilde{A}_{H_n} \right) z_\perp \right\rangle.
$$

Note that

$$
\left\langle \boldsymbol{z}_{\parallel},\left(\boldsymbol{A}_{G_n}\otimes \tilde{\boldsymbol{A}}_{H_n}\right)\boldsymbol{z}_{\parallel}\right\rangle =2e(G)\left\langle \boldsymbol{x},\tilde{\boldsymbol{A}}_{H_n}\boldsymbol{x}\right\rangle,\\ \left\langle \boldsymbol{z}_{\perp},\left(\boldsymbol{A}_{G_n}\otimes \tilde{\boldsymbol{A}}_{H_n}\right)\boldsymbol{z}_{\parallel}\right\rangle =\sum_{i=1}^{n-1}(\deg(i)-\deg(n))\left\langle \boldsymbol{y}_i,\tilde{\boldsymbol{A}}_{H_n}\boldsymbol{x}\right\rangle.
$$

Also using Lemmas [E.1,](#page-28-2) [E.2](#page-28-1) we have

$$
\langle z_\perp, \tilde{\mathbf{Y}}_n z_\perp \rangle \geq \Big(2n q_0 \tilde{a}_n e(G_n) - 2(1+o(1)) \sqrt{n q_0 (1-q_0)} \left(2\tilde{a}_n e(G_n) + \lambda_1(\tilde{\mathbf{A}}_{H_n}) \right) \Big) \|z_\perp\|^2.
$$

Note that if $\lambda_{k_n}(A_{H_n}) \geq (2e(H_n) - k_n^2)/k_n$,

$$
-\lambda_{k_n}(\mathbf{A}_{H_n})-1+\frac{k_n\lambda_{k_n}(\mathbf{A}_{H_n})+k_n^2-2e(H_n)}{nk_n^2}\geq -1.
$$

Therefore,

$$
\lambda_1(\tilde{\mathbf{A}}_{H_n}) = \lambda_1(\mathbf{A}_{H_n}) + u_n + 2a_n e(G_n)
$$

\n
$$
\leq \lambda_1(\mathbf{A}_{H_n}) - \lambda_{k_n}(\mathbf{A}_{H_n}) + \frac{k_n \lambda_{k_n}(\mathbf{A}_{H_n}) + k_n^2 - 2e(H_n)}{nk_n^2}
$$

\n
$$
+ \frac{2e(H_n) + k_n u_n + nk_n^2 - k_n^2}{k_n(nk_n - 1)}
$$

\n
$$
\leq \lambda_1(\mathbf{A}_{H_n}) - \lambda_{k_n}(\mathbf{A}_{H_n}) + 1.
$$

Otherwise, note that

$$
\lambda_1({\boldsymbol{A}}_{H_n}) \geq \frac{2e(H_n)}{k_n}.
$$

Therefore, if $\lambda_{k_n}(\mathbf{A}_{H_n}) < (2e(H_n) - k_n^2)/k_n$ then

$$
\lambda_1(\mathbf{A}_{H_n}) - \lambda_{k_n}(\mathbf{A}_{H_n}) \geq k_n \geq \lambda_1(\tilde{\mathbf{A}}_{H_n}).
$$

Thus,

$$
\lambda_1(\tilde{\boldsymbol{A}}_{H_n}) \leq \lambda_1(\boldsymbol{A}_{H_n}) - \lambda_{k_n}(\boldsymbol{A}_{H_n}) + 1.
$$

In both cases. Using the fact that, $\limsup_{n\to\infty} (2(\lambda_1(\mathbf{A}_{H_n}) - \lambda_{k_n}(\mathbf{A}_{H_n}))\sqrt{1-q_0})/\sqrt{nq_0} = 1 - C$,

$$
\langle \mathbf{z}_{\perp}, \tilde{\mathbf{Y}}_n \mathbf{z}_{\perp} \rangle \geq 2C \tilde{a}_n e(G_n) n q_0 ||\mathbf{z}_{\perp}||^2.
$$

Thus, in order to show positive semidefiniteness of $\tilde{\mathbf{Y}}_n$, it suffices to show that

$$
\boldsymbol{Y}'_n = \begin{bmatrix} 2e(G_n)\tilde{\boldsymbol{A}}_{H_n} & (\deg(1) - \deg(n))\tilde{\boldsymbol{A}}_{H_n} & (\deg(2) - \deg(n))\tilde{\boldsymbol{A}}_{H_n} & \cdots & (\deg(n-1) - \deg(n))\tilde{\boldsymbol{A}}_{H_n} \\ (\deg(1) - \deg(n))\tilde{\boldsymbol{A}}_{H_n} & 2C\tilde{a}_n e(G_n)nq_0\mathbf{I}_{k_n} & 0 & \cdots & 0 \\ (\deg(2) - \deg(n))\tilde{\boldsymbol{A}}_{H_n} & 0 & 2C\tilde{a}_n e(G_n)nq_0\mathbf{I}_{k_n} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ (\deg(n-1) - \deg(n))\tilde{\boldsymbol{A}}_{H_n} & 0 & 0 & \cdots & 2C\tilde{a}_n e(G_n)nq_0\mathbf{I}_{k_n} \end{bmatrix} \succeq 0.
$$

Note that using Bernstein's inequality,

$$
\mathbb{P}\left\{\left|\max_{i\in V(G_n)}\deg(i) - nq_0\right| \ge 2\sqrt{nq_0\log n}\right\} \le 2n \exp\left\{\frac{2nq_0\log n}{nq_0(1-q_0) + (2/3)\sqrt{nq_0\log n}}\right\}
$$

$$
\le 2\exp\left\{\log n - 2\log n\right\} = \frac{2}{n},
$$

for large enough n . Thus,

$$
\mathbb{P}\left\{\left|\max_{i\in V(G_n)}\deg(i) - nq_0\right| \ge 2\sqrt{nq_0\log n}\right\} \to 0,
$$

as $n \to \infty$. Hence, $deg(i) - deg(n) \leq 4$ $nq_0 \log n$, for all n, with high probability, as $n \to \infty$. Therefore, using Schur's theorem, since $C > 0$, we need to show that

$$
2e(G_n)\tilde{A}_{H_n} - 16\left(2C\tilde{a}_n e(G_n)nq_0\right)^{-1} n^2 q_0 \log n \tilde{A}_{H_n}^2
$$

= $C'\tilde{A}_{H_n} \left(\frac{C\tilde{a}_n (e(G_n))^2}{4n \log n} \mathbf{I}_{k_n} - \tilde{A}_{H_n}\right) \succeq 0.$ (E.12)

Where $C' > 0$. This holds, since $\tilde{A}_{H_n} \succeq 0$. Further, $C\tilde{a}_n(e(G_n))^2/(4n \log n)$ is $\Theta(n/\log n)$ and

$$
\lim_{n\to\infty}\frac{(\lambda_1(\boldsymbol{A}_{H_n})-\lambda_{k_n}(\boldsymbol{A}_{H_n}))\log n}{n}=0.
$$

Hence, Y_n is feasible for problem [\(3.17\)](#page-10-4), with high probability as $n \to \infty$. Now, note that

$$
\mathsf{Tr}\left((\boldsymbol{A}_{G_n}\otimes \boldsymbol{A}_{H_n})\boldsymbol{Y}_n\right)\geq 4e(H_n)e(G_n)b_n=2e(H_n).
$$

Thus, with high probability as $n \to \infty$ under null, the optimal value of problem [\(3.17\)](#page-10-4), SDP($G_n; H_n$), is bigger than or equal $2e(H_n)$. Note that the optimal value of (3.17) under the alternative when there is no noise is $2e(H_n)$. Therefore, under the conditions of the Theorem [5,](#page-10-5) for the test based on $\textsf{SDP}(G_n;H_n),$

$$
\mathbb{P}_{0,n}\{T(G_n)=1\} \to 1,
$$

as $n \to \infty$ and the proof is complete.

F Proofs: Multiple planted subgraphs

F.1 Proof of Theorem [6](#page-11-1)

In order to state the proofs in this section we will use the following notation. Let $\varphi_1, \varphi_2, \ldots, \varphi_m$ be labelings of $V(H_n)$ in [n]. We set

$$
M(\varphi_1, \varphi_2, \ldots, \varphi_m) \equiv \left| \bigcup_{l=1}^m \varphi_l(E(H_n)) \right|.
$$

 $\overline{1}$

 \Box

Also recall

$$
N(H_n; G_n) \equiv \left| \left\{ \varphi \in \mathcal{L}(H_n; n) : \varphi(E(H_n)) \subseteq E(G_n) \right\} \right|.
$$

We also denote $v(H_n)$ by k_n . First we prove the following Lemma which is useful in the proof of Theorem [6.](#page-11-1)

Lemma F.1. Let $\{H_n\}_{n\geq 1}$, $q_0, \mathbb{P}_{0,n}, \mathbb{P}_{1,n}$ be as in Theorem [6.](#page-11-1) Under the assumptions of Theorem [6,](#page-11-1) for all $\varepsilon > 0$

$$
\lim_{n \to \infty} \mathbb{P}_{0,n} \left\{ N(H_n; G_n)^{m_n} \le (1 - \varepsilon) (\mathbb{E}_{0,n} N(H_n; G_n))^{m_n} \right\} = 0.
$$
 (F.1)

Proof. As in [\(C.7\)](#page-21-2) in the proof of Lemma [C.2,](#page-20-1) using Chebyshev's inequality

$$
\mathbb{P}\left\{\frac{N(H_n;G_n)^{m_n}}{(\mathbb{E}_{0,n}N(H_n;G_n))^{m_n}} \le 1-\varepsilon\right\} = \mathbb{P}\left\{\frac{N(H_n;G_n)}{\mathbb{E}_{0,n}N(H_n;G_n)} \le (1-\varepsilon)^{1/m_n}\right\}
$$
\n
$$
\le \mathbb{P}\left\{\frac{N(H_n;G_n)}{\mathbb{E}_{0,n}N(H_n;G_n)} \le \left(1-\frac{\varepsilon}{m_n}\right)\right\} \le \frac{m_n^2\bar{\Delta}(n,H_n)}{\varepsilon^2(\mathbb{E}_0N(H_n;G_n))^2}.\tag{F.2}
$$

Where $\Delta(n, H_n)$ is defined in [\(C.5\)](#page-20-2). Note that, using [\(C.11\)](#page-22-0) we have

$$
\frac{m_n^2 \bar{\Delta}(n, H_n)}{(\mathbb{E}_0 N(H_n; G_n))^2} \le C \exp\left\{-\tilde{u}\left(f(\tilde{u}) - \frac{\log k_n}{\tilde{u}} - \frac{2\log m_n}{\tilde{u}}\right)\right\}
$$

where $f(u)$ is defined in [\(C.9\)](#page-21-0) and [\(C.10\)](#page-21-1) and $\tilde{u} = \arg \min_{2 \le u \le k_n} \{uf(u)\}$. Using [\(C.12\)](#page-22-1),

$$
f(\tilde{u}) - \frac{\log k_n}{\tilde{u}} - \frac{2\log m_n}{\tilde{u}} \ge \log n - \frac{e_{H_n}(\tilde{u})}{\tilde{u}}\log(1/q_0) - (5/2)\log k_n - \frac{2\log m_n}{\tilde{u}} + C
$$

for some constant C. Note that $0 \le e_{H_n}(\tilde{u})/\tilde{u} \le \min(d(H_n), \tilde{u})$. Therefore,

$$
f(\tilde{u}) - \frac{\log k_n}{\tilde{u}} - \frac{2\log m_n}{\tilde{u}} \ge \log n - \min\left(d(H_n), \tilde{u}\right) \log(1/q_0) - \frac{2\log m_n}{\tilde{u}} - (5/2)\log k_n + C. \tag{F.3}
$$

The right hand side of [\(F.3\)](#page-34-0), as a function of \tilde{u} , is minimized at $\tilde{u} = 2$ or $\tilde{u} = d(H_n)$ (we can assume, without loss of generality, $d(H_n) \geq 2$. Therefore, it is sufficient to show that the right hand side of [\(F.3\)](#page-34-0) goes to infinity for $\tilde{u} \in \{2, d(H_n)\}\$. For $\tilde{u} = 2$, the right hand side of (F.3) is equal to

$$
X = \log n \left(1 - \frac{2}{\log n} \log(1/q_0) - \frac{\log m_n}{\log n} - (5/2) \frac{\log k_n}{\log n} + o(1) \right) \to \infty
$$
 (F.4)

since by the assumption of Theorem [6](#page-11-1)

$$
\lim\sup_{n\to\infty}\frac{(5/2)\log k_n + \log m_n}{\log n} < 1.
$$

For $\tilde{u} = d(H_n)$, the right hand side of [\(F.3\)](#page-34-0) is equal to

$$
X = \log n \left(1 - \frac{d(H_n)}{\log n} \log(1/q_0) - \frac{2 \log m_n}{d(H_n) \log n} - (5/2) \frac{\log k_n}{\log n} + o(1) \right).
$$

Fix a constant $M > 0$ large enough (to be adjusted below). For $d(H_n) \leq M$ we have

$$
X \ge \log n \left(1 - \frac{M \log(1/q_0)}{\log n} - \frac{2 \log m_n}{d(H_n) \log n} - (5/2) \frac{\log k_n}{\log n} + o(1) \right)
$$

$$
\ge \log n \left(1 - \frac{2}{M} - \frac{\log m_n}{\log n} - (5/2) \frac{\log k_n}{\log n} + o(1) \right)
$$
(F.5)

where the last inequality holds for all n large enough. For $d(H_n) \geq M$, using $m_n \leq n$, we have

$$
X \ge \log n \left(1 - \frac{d(H_n) \log(1/q_0)}{\log n} - \frac{2 \log m_n}{M \log n} - (5/2) \frac{\log k_n}{\log n} + o(1) \right)
$$

$$
\ge \log n \left(1 - \frac{d(H_n) \log(1/q_0)}{\log n} - \frac{2}{M} - (5/2) \frac{\log k_n}{\log n} + o(1) \right).
$$
 (F.6)

Combining $(F.6)$, $(F.5)$, we have

$$
X \ge \log n \left(\min \left(1 - \frac{\log m_n}{\log n} - (5/2) \frac{\log k_n}{\log n}, 1 - \frac{d(H_n) \log (1/q_0)}{\log n} - (5/2) \frac{\log k_n}{\log n} \right) - \frac{2}{M} - o(1) \right)
$$

$$
\ge \left(\delta - \frac{2}{M} \right) \log n.
$$
 (F.7)

Where

$$
\delta = \min\left(\liminf_{n \to \infty} \left\{ 1 - \frac{\log m_n}{\log n} - (5/2) \frac{\log k_n}{\log n} \right\}, \liminf_{n \to \infty} \left\{ 1 - \frac{d(H_n) \log(1/q_0)}{\log n} - (5/2) \frac{\log k_n}{\log n} \right\} \right)
$$

and by assumptions of Theorem [6,](#page-11-1) $\delta > 0$. Hence, by taking $M = 4/\delta$, using [\(F.7\)](#page-35-2), we deduce that the right hand side of [\(F.3\)](#page-34-0) goes to infinity for $\tilde{u} = d(H_n)$. Combining with [\(F.4\)](#page-34-1), we deduce that under the assumptions of Theorem [6](#page-11-1)

$$
f(\tilde{u}) - \frac{\log k_n}{\tilde{u}} - \frac{2\log m_n}{\tilde{u}} \to \infty
$$

and

$$
\frac{m_n^2 \bar{\Delta}(n, H_n)}{(\mathbb{E}_0 N(H_n; G_n))^2} \to 0.
$$
 (F.8)

as $n \to \infty$. Hence, using [\(F.2\)](#page-34-2), [\(F.8\)](#page-35-3) we get that under the assumptions of Theorem [6,](#page-11-1)

$$
\lim_{n \to \infty} \mathbb{P}_{0,n} \left\{ (N(H_n; G_n))^{m_n} \le (1 - \varepsilon) (\mathbb{E}_0 N(H_n; G_n))^{m_n} \right\} = 0
$$

This completes the proof of lemma.

Now we can state the proof of Theorem [6.](#page-11-1)

 \Box

Proof of Theorem [6.](#page-11-1) For the laws $\mathbb{P}_{0,n}$, $\mathbb{P}_{1,n}$ defined as in Theorem [6](#page-11-1) we have

$$
\frac{d\mathbb{P}_{1,n}}{d\mathbb{P}_{0,n}}(G_n) = \frac{1}{(n)_{k_n}^{m_n}} \sum_{\varphi_1,\dots,\varphi_{m_n} \in \mathcal{L}(H_n,n)} \left(\frac{1}{q_0}\right)^{M(\varphi_1,\dots,\varphi_{m_n})} \mathbb{I}\left\{ \left(\bigcup_{l} \varphi_l(E(H_n))\right) \subseteq E(G_n) \right\}
$$
\n
$$
= \frac{1}{(n)_{k_n}^{m_n} q_0^{m_n e(H_n)}} \sum_{\varphi_1,\dots,\varphi_{m_n} \in \mathcal{L}(H_n,n)} \mathbb{I}\left\{ \left(\bigcup_{l} \varphi_l(E(H_n))\right) \subseteq E(G_n) \right\}
$$
\n
$$
- \frac{1}{(n)_{k_n}^{m_n} q_0^{m_n e(H_n)}} \sum_{\varphi_1,\dots,\varphi_{m_n} \in \mathcal{L}(H_n,n)} \left(1 - q_0^{m_n e(H_n) - M(\varphi_1, \varphi_2, \dots, \varphi_{m_n})}\right) \mathbb{I}\left\{ \left(\bigcup_{l} \varphi_l(E(H_n))\right) \subseteq E(G_n) \right\}
$$
\n
$$
= \frac{N(H_n; G_n)^{m_n}}{(\mathbb{E}_{0,n} N(H_n; G_n))^{m_n}} - X_n.
$$
\n(F.9)

where

$$
X_n = \frac{1}{(n)_{k_n}^{m_n} q_0^{m_n e(H_n)}} \sum_{\varphi_1, \dots, \varphi_{m_n} \in \mathcal{L}(H_n, n)} \left(1 - q_0^{m_n e(H_n) - M(\varphi_1, \varphi_2, \dots, \varphi_{m_n})}\right) \mathbb{I}\left\{ \left(\bigcup_l \varphi_l(E(H_n))\right) \subseteq E(G_n)\right\}.
$$

Note that under the assumptions of Theorem [6,](#page-11-1) using Lemma [F.1](#page-34-3)

$$
\lim_{n \to \infty} \mathbb{P}_{0,n} \left\{ (N(H_n; G_n))^{m_n} \le (1 - \varepsilon) (\mathbb{E}_0 N(H_n; G_n))^{m_n} \right\} = 0.
$$

Hence, using the same argument used in proof of Theorem [2,](#page-5-2)

$$
\frac{N(H_n; G_n)^{m_n}}{\mathbb{E}_{0,n}(N(H_n; G_n)^{m_n})} \xrightarrow{p} 1.
$$
\n(F.10)

Now, we prove that under the assumptions of Theorem [6,](#page-11-1) $X_n \stackrel{p}{\longrightarrow} 0$. Note that we have

$$
X_n \leq \frac{1}{(\mathbb{E}_{0,n}N(H_n;G_n))^{m_n}} \sum_{\varphi_1,\ldots,\varphi_{m_n}\in\mathcal{L}(H_n,n)} \mathbb{I}\left\{M(\varphi_1,\ldots,\varphi_{m_n}) < m_n e(H_n)\right\} \tag{F.11}
$$
\n
$$
= \frac{(N(H_n;G_n))^{m_n}}{(\mathbb{E}_{0,n}N(H_n;G_n))^{m_n}} \frac{1}{(N(H_n;G_n))^{m_n}} \sum_{\varphi_1,\ldots,\varphi_{m_n}\in\mathcal{L}(H_n,n)} \mathbb{I}\left\{M(\varphi_1,\ldots,\varphi_{m_n}) < m_n e(H_n)\right\} \tag{F.12}
$$

$$
\equiv \frac{(N(H_n; G_n))^{m_n}}{(\mathbb{E}_{0,n} N(H_n; G_n))^{m_n}} Q_{m_n}(H_n; G_n).
$$
\n(F.13)

Note that $Q_{m_n}(G_n; H_n)$ can be interpreted as the probability that, drawing the embeddings φ_l , $l \in \{1, \ldots, m_m\}$ independently and uniformly at random in $\mathcal{L}(H_n, h)$, at least two of them share an edge. By union bound, we have

$$
Q_{m_n}(H_n; G_n) \le m_n^2 Q_2(H_n; G_n), \tag{F.14}
$$

$$
Q_2(H_n; G_n) = \frac{N_2(H_n; G_n)}{N(H_n; G_n)^2},
$$
\n(F.15)

and

$$
N_2(H_n; G_n) = |\{(\varphi_1, \varphi_2); \varphi_1(E(H_n)) \cap \varphi_2(E(H_n)) \neq \emptyset, \varphi_i(E(H_n)) \subseteq E(G_n) \text{ for } i = 1, 2\}|.
$$

Note that

$$
\mathbb{E}_{0,n} N_2(H_n; G_n) \le \sum_{l=2}^{k_n} n^{2k_n - l} q_0^{2e(H_n) - e_{H_n}(l)} \le k_n n^{2k_n - \tilde{l}} q_0^{2e(H_n) - e_{H_n}(\tilde{l})}
$$

where $\tilde{l} = \arg \max_{2 \leq l \leq k_n} n^{2k_n - l} q_0^{2e(H_n) - e_{H_n}(l)}$ $e^{2e(H_n)-e_{H_n}(t)}$. Hence,

$$
\log \frac{m_n^2 \mathbb{E}_{0,n} N_2(H_n; G_n)}{(\mathbb{E}_{0,n} N(H_n; G))^2} \leq -\tilde{l} \left(\log n - \frac{\log k_n}{\tilde{l}} - \frac{2 \log m_n}{\tilde{l}} - \frac{e_{H_n}(\tilde{l})}{\tilde{l}} \log(1/q_0) \right)
$$

$$
\leq -\tilde{l} \left(\log n - \min \left(\tilde{l}, d(H_n) \right) \log(1/q_0) - \frac{\log k_n}{\tilde{l}} - \frac{2 \log m_n}{\tilde{l}} \right).
$$

Note that in the proof of Lemma [F.1,](#page-34-3) we proved that under the assumptions of Theorem [6,](#page-11-1) for all $2\leq l\leq k_n$

$$
\min_{2 \leq l \leq k_n} \left[\log n - \min\left(d(H_n), l\right) \log(1/q_0) - \frac{2 \log m_n}{l} - (5/2) \log k_n \right] \to \infty
$$

as $n \to \infty$. Hence, under the assumptions of Theorem [6,](#page-11-1)

$$
-\tilde{l}\left(\log n - \min\left(\tilde{l}, d(H_n)\right)\log(1/q_0) - \frac{\log k_n}{\tilde{l}} - \frac{2\log m_n}{\tilde{l}}\right) \to -\infty
$$

and

$$
\frac{m_n^2 \mathbb{E}_{0,n} N_2(H_n; G_n)}{(\mathbb{E}_{0,n} N(H_n; G_n))^2} \to 0.
$$
 (F.16)

Note that

$$
m_n^2 Q_2(H_n; G_n) = \frac{m_n^2 N_2(H_n; G_n)}{(\mathbb{E}_{0,n} N(H_n; G_n))^2} \frac{(\mathbb{E}_{0,n} N(H_n; G_n))^2}{N(H_n; G_n)^2}.
$$

Using [\(F.16\)](#page-37-0) and Markov's inequality

$$
\frac{m_n^2 N_2(H_n; G_n)}{(\mathbb{E}_{0,n} N(H_n; G_n))^2} \xrightarrow{p} 0.
$$

Hence, by [\(F.13\)](#page-36-0), [\(F.15\)](#page-36-1), and [\(F.11\)](#page-36-2)

$$
X_n \xrightarrow{p} 0.
$$

Thus, using [\(F.9\)](#page-36-3) and [\(F.10\)](#page-36-4), under $\mathbb{P}_{0,n}$

$$
\frac{\mathrm{d} \mathbb{P}_{1,n}}{\mathrm{d} \mathbb{P}_{0,n}}(G_n) \xrightarrow{p} 1
$$

and using Lemma [C.1](#page-18-1) the proof is complete.

 \Box

F.2 Proof of Theorem [7](#page-11-2)

Proof of Theorem [7.](#page-11-2) Let $N_{0,n}, N_{1,n}$ be the random variables denoting the number of edges of G_n when it is generated according to null and alternative models, respectively. We would like to show that there exists t_n^* such that $N_{0,n} < t_n^*$ and $N_{1,n} > t_n^*$ with high probability as $n \to \infty$. Let

$$
t_n^* = \frac{n(n-1)q_0}{2} + \delta_n m_n e(H_n)
$$

for some $\delta_n > 0$. Note that $N_{0,n}$ is a binomial random variable where $\mathbb{E}N_{0,n} = n(n-1)q_0/2$ and $Var(N_{0,n}) = n(n-1)q_0(1-q_0)/2$. Hence, using Chebyshev's inequality

$$
N_{0,n} \le \frac{n(n-1)q_0}{2} + C_n \sqrt{\frac{n(n-1)q_0(1-q_0)}{2}}
$$

with probability $1-1/C_n^2$. Since $\liminf_{n\to\infty} m_n e(H_n)/n = \infty$, taking $C_n \to \infty$ such that $C_n/\delta_n <$ $m_n e(H_n)/n$, we have $N_{0,n} < t_n^*$ with high probability as $n \to \infty$. Note that $N_{1,n}$ is monotonically increasing in m_n . Hence, in order to show that $N_{1,n} > t_n^*$ as $n \to \infty$ we can assume, without loss of generality, $\limsup_{n\to\infty} m_n e(H_n)/n^2 = 0$. We have

$$
N_{1,n} = X_n + m_n e(H_n) - Z_{0,n} - \sum_{i=1}^{m_n - 1} i Z_{i,n}.
$$
 (F.17)

Here X_n is denotes the number of edges in the graph before adding the copies of H_n . Hence, $\mathbb{E}X_n = n(n-1)q_0/2$ and $\text{Var}(X_n) = n(n-1)q_0(1-q_0)/2$. Further, $m_ne(H_n)$ is the total number of edges of the planted subgraphs; $Z_{0,n}$ is the number of edges in the planted subgraphs that are present before adding the subgraphs and $Z_{i,n}$ is the number of edges that are present in exactly $i+1$ different embeddings. Using Chebyshev's inequality

$$
X_n \ge \frac{n(n-1)q_0}{2} - \tilde{C}_n \sqrt{\frac{n(n-1)q_0(1-q_0)}{2}} \quad \text{with probability at least } 1 - \frac{1}{\tilde{C}_n^2},\tag{F.18}
$$

$$
Z_{0,n} \le m_n e(H_n)q_0 + \tilde{C}_n \sqrt{\frac{m_n e(H_n)q_0(1-q_0)}{2}} \quad \text{with probability at least } 1 - \frac{1}{\tilde{C}_n^2}.\tag{F.19}
$$

(The last inequality follows because $Z_{0,n}$ is dominated by a binomial random variable with parameters $m_n e(H_n)$, q_0 .)

In addition,

$$
\sum_{i=1}^{m_n-1} i \mathbb{E} Z_{i,n} \leq \sum_{i=1}^{m_n-1} \frac{i \binom{m_n}{i+1} (e(H_n))^{i+1}}{\binom{n}{2}^i} \leq \sum_{i=1}^{m_n-1} \frac{i (m_n e(H_n))^{i+1}}{n^{2i}}
$$

$$
\leq \frac{(m_n e(H_n))^2}{(n^2 - m_n e(H_n))(1 - m_n e(H_n)/n^2)} \leq \frac{2 (m_n e(H_n))^2}{n^2 - m_n e(H_n)}
$$

for large enough n. Therefore, using Markov's inequality,

$$
\sum_{i=1}^{m_n-1} iZ_{i,n} \le \varepsilon m_n e(H_n) \quad \text{with probability } 1 - \frac{2m_n e(H_n)}{\varepsilon (n^2 - m_n e(H_n))}.
$$
 (F.20)

Now using $(F.17)-(F.20)$ $(F.17)-(F.20)$ $(F.17)-(F.20)$, we can write

$$
N_{1,n} \ge \frac{n(n-1)q_0}{2} + (1-q_0-\varepsilon)m_ne(H_n) - \tilde{C}_n\sqrt{\frac{n(n-1)q_0(1-q_0)}{2}} - \tilde{C}_n\sqrt{\frac{m_ne(H_n)q_0(1-q_0)}{2}}
$$

with probability $1 - 2/\tilde{C}_n^2 - 3m_n e(H_n)/(\varepsilon n^2)$. Since $\liminf_{n \to \infty} m_n e(H_n)/n = \infty$, taking ε < $1-q_0-\delta_n$ and $\tilde{C}_n \to \infty$ such that $\tilde{C}_n/(1-q_0-\varepsilon-\delta_n) < m_n e(H_n)/(n+\sqrt{m_n e(H_n)})$, we have

$$
N_{1,n} > t_n^*
$$

with high probability as $n \to \infty$. Therefore, the two models are strongly distinguishable under the assumptions of Theorem [7](#page-11-2) and this completes the proof. \Box