

Two Higgs doublet models with an S_3 symmetry

D. Cogollo^{1,2,*} and João P. Silva^{1,†}

¹*CFTP, Departamento de Física, Instituto Superior Técnico,
Universidade de Lisboa, Avenida Rovisco Pais 1, 1049 Lisboa, Portugal*

²*Departamento de Física, Universidade Federal de Campina Grande,
Caixa Postal 10071, 58109-970, Campina Grande, PB, Brazil*

(Dated: September 27, 2021)

We study all implementations of the S_3 symmetry in the two Higgs doublet model with quarks, consistent with non-zero quark masses and a Cabibbo-Kobayashi-Maskawa (CKM) matrix which is not block diagonal. We study the impact of the various soft-breaking terms and vacuum expectation values, and find an interesting relation between the mixing angles α and β . We also show that, in this minimal setting, only two types of assignments are possible: either all field sectors are in singlets or all field sectors have a doublet.

PACS numbers: 12.60.Fr, 14.80.Ec, 14.80.-j

I. INTRODUCTION

Since the discovery of a spin 0 particle at LHC [1, 2] a crucial task in particle physics has been the determination of the number of such particles. In the Standard Model (SM), the existence of a single scalar doublet is a feature without a strong theoretical justification. As a result, multi scalar models have been studied for a long time, in particular, two Higgs doublet models (2HDM) [3, 4].

Because such models involve many new parameters, one may curtail their number by imposing extra symmetries. The identification of all possible symmetries that may be imposed on the scalar potential of the 2HDM has been performed in Refs. [5, 6]. In the notation of Ref. [6] they are: the Higgs family symmetries Z_2 , $U(1)$, and $U(2)$; and the (generalized) CP symmetries CP1, CP2, and CP3. Some steps have been taken to extend this analysis into the Yukawa sector. Ref. [7] identifies all implementations of abelian symmetries, showing that they allow for 34 distinct matrix forms, while Ref. [8] shows that there is a single implementation of a generalized CP symmetry in both scalar potential and Yukawa couplings which could be consistent with experiment. There is no classification of all possible implementations on non-abelian symmetries in both scalar and quark Yukawa sectors. This is what we target here, focusing on the simplest non-abelian group: S_3 .

There has been recent interest in specific implementations of S_3 in the 2HDM [9, 10], and many articles extending the 2HDM with extra scalar and/or fermion fields – see, for example, Refs. [11, 12]. In this article, we provide a complete classification of all possible implementation of S_3 in the 2HDM which are consistent with non-vanishing quark masses and a CKM matrix which is not block-diagonal, as required by experiment.

In Section II we introduce the S_3 representation we use, discussing in detail the features of the scalar potential implied by each possible representation of the scalar fields, the possible soft-breaking terms, and the corresponding vacuum expectation values (vev). In Section III we discuss all possible extensions of S_3 into the Yukawa sector, and in Section IV we draw our conclusions.

*E-mail: diego.cogollo@cftp.ist.utl.pt

†E-mail: jpsilva@cftp.ist.utl.pt

II. THE S_3 SYMMETRY AND THE HIGGS POTENTIAL

S_3 is the group of permutations of three objects and it has 6 elements and three irreducible representations; namely, two singlets $\mathbf{1}$, $\mathbf{1}'$, and one doublet $\mathbf{2}$. The multiplication rules are

$$\begin{aligned} \mathbf{1} \otimes \text{any} &= \text{any}, \\ \mathbf{1}' \otimes \mathbf{1}' &= \mathbf{1}, \\ \mathbf{1}' \otimes \mathbf{2} &= \mathbf{2}, \\ \mathbf{2} \otimes \mathbf{2} &= \mathbf{1} \oplus \mathbf{1}' \oplus \mathbf{2}. \end{aligned} \tag{1}$$

The representation $\mathbf{1}$ leaves the fields invariant, while fields in $\mathbf{1}'$ change their sign for odd permutations. Consider a two Higgs doublet model (2HDM) with scalar fields Φ_1 and Φ_2 .

We will denote $\Phi \sim (\mathbf{1}, \mathbf{1})$ when both scalars are in the singlet representation of S_3 . In this case we obtain the generic scalar potential of the 2HDM, which may be written as [3, 4]

$$\begin{aligned} V_H &= m_{11}^2 |\Phi_1|^2 + m_{22}^2 |\Phi_2|^2 - m_{12}^2 \Phi_1^\dagger \Phi_2 - (m_{12}^2)^* \Phi_2^\dagger \Phi_1 \\ &+ \frac{\lambda_1}{2} |\Phi_1|^4 + \frac{\lambda_2}{2} |\Phi_2|^4 + \lambda_3 |\Phi_1|^2 |\Phi_2|^2 + \lambda_4 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) \\ &+ \left[\frac{\lambda_5}{2} (\Phi_1^\dagger \Phi_2)^2 + (\lambda_6 |\Phi_1|^2 + \lambda_7 |\Phi_2|^2) (\Phi_1^\dagger \Phi_2) + \text{h.c.} \right], \end{aligned} \tag{2}$$

where ‘‘h.c.’’ stands for the hermitian conjugate. Hermiticity implies that all couplings are real, except possibly m_{12}^2 , λ_5 , λ_6 , and λ_7 . We will denote the vacuum expectation values (vevs) by $v_1/\sqrt{2}$ and $v_2/\sqrt{2}$, but we will omit the $\sqrt{2}$ in the text (though including it in actual calculations). Depending on the parameters of the potential, we may get any values for (v_1, v_2) , including the inert cases $(v, 0)$ and $(0, v)$.

If we choose the S_3 representations as $\Phi \sim (\mathbf{1}, \mathbf{1}')$, then all terms odd in Φ_2 must vanish, and we obtain the Z_2 symmetric potential: $m_{12}^2 = \lambda_6 = \lambda_7 = 0$. Usually, one includes also the terms in m_{12}^2 , which break the symmetry, but only softly. In this case, if $\arg(\lambda_5) = 2 \arg(m_{12}^2)$, then the phases can be removed. This is known as the real 2HDM and the scalar sector preserves CP. One can have nonvanishing vevs (v_1, v_2) , but the inert vevs $(v, 0)$ and $(0, v)$ are only possible if $m_{12}^2 = 0$. Alternatively, if $\arg(\lambda_5) \neq 2 \arg(m_{12}^2)$, then the phases cannot be removed. This is known as the complex 2HDM (C2HDM) and the scalar sector violates CP. In both models with softly broken Z_2 symmetry, the conditions for a bounded from below potential are [13]

$$\lambda_1 > 0, \quad \lambda_2 > 0, \quad \sqrt{\lambda_1 \lambda_2} > -\lambda_3, \quad \sqrt{\lambda_1 \lambda_2} > |\lambda_5| - \lambda_3 - \lambda_4. \tag{3}$$

We denote collectively by $\Phi \sim \mathbf{s}$ the two possibilities just discussed: $\Phi \sim (\mathbf{1}, \mathbf{1})$ and $\Phi \sim (\mathbf{1}, \mathbf{1}')$. Since the labels 1 and 2 are arbitrary, the case $\Phi \sim (\mathbf{1}', \mathbf{1})$ is automatically included. Moreover, the scalar potential remains the most general if one considers $\Phi \sim (\mathbf{1}', \mathbf{1}')$ ¹.

In order to describe how S_3 acts on doublets we must choose a representation for the group. A suitable real basis for the doublet representation of S_3 is given by [14]

$$a = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad b = \begin{pmatrix} -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{pmatrix}. \tag{4}$$

These matrices satisfy $a^2 = b^3 = (ba)^2 = 1$, showing that they indeed generate the group S_3 . In S_3 , with the basis of Eq. (4), the product of two doublets, $x = (x_1, x_2)^\top$ and $y = (y_1, y_2)^\top$, gives

$$\begin{aligned} (x \otimes y)_{\mathbf{1}} &= x_1 y_1 + x_2 y_2, \\ (x \otimes y)_{\mathbf{1}'} &= x_1 y_2 - x_2 y_1, \\ (x \otimes y)_{\mathbf{2}} &= (x_2 y_2 - x_1 y_1, x_1 y_2 + x_2 y_1)^\top. \end{aligned} \tag{5}$$

¹ The Yukawa couplings do not; but one can always reduce this to $\Phi \sim (\mathbf{1}, \mathbf{1})$. For example, if the right-handed fermions are in singlet representations of S_3 , one can change $\Phi \sim (\mathbf{1}', \mathbf{1}')$ to $\Phi \sim (\mathbf{1}, \mathbf{1})$, by simultaneously changing the representation for the right-handed fermions from $\mathbf{1}$ to $\mathbf{1}'$ and vice-versa.

Similarly, the product of the doublet x with the singlet y' of $\mathbf{1}'$ gives

$$(x \otimes y') = (-x_2 y', x_1 y')^\top. \quad (6)$$

Let us consider two scalars $\Phi = (\varphi_1, \varphi_2)^\top$ which transform as a doublet under the real basis of Eq. (4). According to Eq. (5), the relevant combinations of $\varphi_i^\dagger \varphi_j$ are

$$\begin{aligned} & |\varphi_2|^2 + |\varphi_1|^2, \\ & \varphi_1^\dagger \varphi_2 - \varphi_2^\dagger \varphi_1, \\ & (|\varphi_2|^2 - |\varphi_1|^2, \varphi_1^\dagger \varphi_2 + \varphi_2^\dagger \varphi_1)^\top, \end{aligned} \quad (7)$$

transforming, respectively, as $\mathbf{1}$, $\mathbf{1}'$, and $\mathbf{2}$. Thus, there is only one quadratic term in the potential, proportional to the first line of Eq. (7). The quartic terms come from squaring the first two lines and from the singlet combination of two doublets of the third line. Thus, the most general potential of a doublet of S_3 , consistent with the real representation of Eq. (4) is:

$$\begin{aligned} V_R = & \mu (|\varphi_2|^2 + |\varphi_1|^2) + d_1 (|\varphi_2|^2 + |\varphi_1|^2)^2 + d_2 \left(\varphi_1^\dagger \varphi_2 - \varphi_2^\dagger \varphi_1 \right)^2 \\ & + d_3 \left[(|\varphi_2|^2 - |\varphi_1|^2)^2 + \left(\varphi_1^\dagger \varphi_2 + \varphi_2^\dagger \varphi_1 \right)^2 \right]. \end{aligned} \quad (8)$$

This coincides with the generic potential in Eq. (2), subject to the conditions

$$m_{11}^2 = m_{22}^2, \quad m_{12}^2 = 0, \quad \lambda_1 = \lambda_2, \quad \lambda_5 = \lambda_1 - \lambda_3 - \lambda_4, \quad (9)$$

identified in Table I of Ref. [6] as the CP3 model.

We have studied how the scalars transform as doublets under S_3 using the real representation in Eq. (4). But other representations are possible. Imagine that the scalars Φ transform under a Higgs family symmetry S as

$$\Phi \rightarrow \Phi^S = S \Phi. \quad (10)$$

If one chooses a different basis, with new scalars Φ' given by

$$\Phi' = U \Phi, \quad (11)$$

then the specific form of the Higgs family symmetry in the new basis

$$\Phi' \rightarrow \Phi'^S = S' \Phi' \quad (12)$$

gets altered into

$$S' = U S U^\dagger. \quad (13)$$

As stressed by Ma [11], it is easier to work with a complex representation for S_3 , obtained with

$$U = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & i \\ 1 & -i \end{pmatrix} \implies U^\dagger = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -i & i \end{pmatrix}. \quad (14)$$

Applying this to the matrices in Eq. (4), we obtain

$$a_C = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad b_C = \begin{pmatrix} \omega & 0 \\ 0 & \omega^2 \end{pmatrix}, \quad (15)$$

where a_C interchanges the two fields and $\omega = e^{2i\pi/3}$ ($\omega^3 = 1$). We denote the fields in the complex basis by $\Phi' = \phi = (\phi_1, \phi_2)^\top$. In S_3 , with the basis of Eq. (15), the product of two doublets, $x = (x_1, x_2)^\top$ and $y = (y_1, y_2)^\top$, gives [11]

$$\begin{aligned} (x \otimes y)_{\mathbf{1}} &= x_1 y_2 + x_2 y_1, \\ (x \otimes y)_{\mathbf{1}'} &= x_1 y_2 - x_2 y_1, \\ (x \otimes y)_{\mathbf{2}} &= (x_2 y_2, x_1 y_1)^\top. \end{aligned} \quad (16)$$

Comparing Eqs. (5) and (16) we see the great advantage of the complex representation; the product of two doublets becomes diagonal, as does the product of three doublets, which becomes simply $111 + 222$. In some cases, such as that discussed in subsection III A below, the Yukawa couplings of the quarks are off-diagonal when using the real representation. However, one then has to diagonalize the mass matrices, realizing that this diagonalization is performed by the same unitary transformations in the up and down sectors and, thus, that the CKM matrix coincides with the unit matrix. The physics is easier to grasp in the complex representation, where some Yukawa couplings become diagonal directly. In the complex representation, the product of the doublet x with the singlet y' of $\mathbf{1}'$ gives

$$(x \otimes y') = (x_1 y', -x_2 y')^\top. \quad (17)$$

When considering the hermitian conjugated fields there is a complication arising in the complex representation. If $(\phi_1, \phi_2) \sim \mathbf{2}$, then in the real representation $(\phi_1^\dagger, \phi_2^\dagger) \sim \mathbf{2}$, but in the complex representation one has instead $(\phi_2^\dagger, \phi_1^\dagger) \sim \mathbf{2}$ [11]. Therefore, considering two scalars $\phi = (\phi_1, \phi_2)^\top$ which transform as a doublet under the complex basis of Eq. (15), the relevant combinations of $\phi_i^\dagger \phi_j$ according to Eq. (16) are

$$\begin{aligned} & |\phi_2|^2 + |\phi_1|^2, \\ & |\phi_2|^2 - |\phi_1|^2, \\ & (\phi_1^\dagger \phi_2, \phi_2^\dagger \phi_1)^\top, \end{aligned} \quad (18)$$

transforming, respectively, as $\mathbf{1}$, $\mathbf{1}'$, and $\mathbf{2}$. Thus, there is only one quadratic term in the potential, proportional to the first line of Eq. (18). The quartic terms come from squaring the first two lines and from the singlet combination of two doublets of the third line. Thus, the most general potential of a doublet of S_3 , consistent with the complex representation of Eq. (15) is [12]:

$$\begin{aligned} V_C = & \mu_1^2 (|\phi_2|^2 + |\phi_1|^2) + \frac{1}{2} \ell_1 (|\phi_2|^2 + |\phi_1|^2)^2 + \frac{1}{2} \ell_2 (|\phi_2|^2 - |\phi_1|^2)^2 \\ & + \ell_3 (\phi_1^\dagger \phi_2) (\phi_2^\dagger \phi_1). \end{aligned} \quad (19)$$

This is the same as Eq. (8), through the transformation in Eqs. (11) and (14), with $\mu_1^2 = \mu$, $\ell_1 = 2d_1$, $\ell_2 = -2d_2$, and $\ell_3 = 4d_3$. The potential obtained coincides with the generic potential in Eq. (2), subject to the conditions

$$m_{11}^2 = m_{22}^2, \quad m_{12}^2 = 0, \quad \lambda_1 = \lambda_2, \quad \lambda_5 = 0, \quad (20)$$

which may seem not to coincide with those in Eq. (9). However, they are the same conditions, but seen in different basis. This had already been pointed out in Eqs. (104)-(106) of Ref. [6], where it is shown that the CP3 model can also be obtained by imposing both $U(1)$ and the so-called Π_2 symmetry $\phi_1 \leftrightarrow \phi_2$ in the same basis. Ma and Melic [12] also include in the potential a term which breaks S_3 softly, while preserving the $\phi_1 \leftrightarrow \phi_2$ symmetry:

$$V_{\text{soft}} = -\mu_2^2 (\phi_1^\dagger \phi_2 + \phi_2^\dagger \phi_1). \quad (21)$$

This term is needed since otherwise there would be a massless pseudoscalar. But, including it precludes the inert vevs $(v, 0)$ and $(0, v)$.

We will now consider the potential $V = V_C + V_{\text{soft}}$. In terms of the new parameters, the bounded from below conditions in Eq. (3) read

$$\ell_1 + \ell_2 > 0, \quad \ell_1 > 0, \quad 2\ell_1 + \ell_3 > 0. \quad (22)$$

A convenient way to write the stationarity conditions is

$$\begin{aligned} 0 = v_2 \frac{\partial V}{\partial v_1} - v_1 \frac{\partial V}{\partial v_2} &= (v_1^2 - v_2^2) \left[\mu_2^2 + \frac{1}{2} (2\ell_2 - \ell_3) v_1 v_2 \right], \\ 0 = v_1 \frac{\partial V}{\partial v_1} - v_2 \frac{\partial V}{\partial v_2} &= (v_1^2 - v_2^2) \left[\mu_1^2 + \frac{1}{2} (\ell_1 + \ell_2) v^2 \right], \end{aligned} \quad (23)$$

where $v = \sqrt{v_1^2 + v_2^2} = 246$ GeV. Ma and Melic [12] consider the vev (v, v) , for which $\partial V / \partial v_1$ yields

$$\mu_1^2 = \mu_2^2 - \frac{1}{4} (2\ell_1 + \ell_3) v^2 \quad (24)$$

and the scalar masses

$$\begin{aligned}
m_{H^\pm}^2 &= 2\mu_2^2 - \frac{1}{2}\ell_3 v^2, \\
m_A^2 &= 2\mu_2^2, \\
m_h^2 &= \frac{1}{2}(2\ell_1 + \ell_3)v^2, \\
m_H^2 &= 2\mu_2^2 + \frac{1}{2}(2\ell_2 - \ell_3)v^2,
\end{aligned} \tag{25}$$

for the charged scalars (H^\pm), the pseudoscalar (A), the light (h) and the heavy (H) CP even scalars, respectively.

We consider also the vev² (v_1, v_2) with $v_1 \neq v_2$. Then, μ_1^2 and μ_2^2 are found from Eqs. (23). We parametrize the fields as

$$\begin{aligned}
\phi_1 &= \begin{pmatrix} c_\beta G^+ - s_\beta H^+ \\ \frac{1}{\sqrt{2}} [v_1 + \rho_1 + i(c_\beta G^0 - s_\beta A)] \end{pmatrix}, \\
\phi_2 &= \begin{pmatrix} s_\beta G^+ + c_\beta H^+ \\ \frac{1}{\sqrt{2}} [v_2 + \rho_2 + i(s_\beta G^0 + c_\beta A)] \end{pmatrix},
\end{aligned} \tag{26}$$

where G^0 and G^+ are the would-be Goldstone bosons,

$$v_1 = v c_\beta, \quad v_2 = v s_\beta, \tag{27}$$

and, thenceforth, c_θ (s_θ , t_θ) represent the cosine (sine, tangent) of whatever angle θ is the subindex. The charged scalar and pseudoscalar masses become

$$m_{H^\pm}^2 = -\ell_2 v^2, \tag{28}$$

$$m_A^2 = -\frac{1}{2}(2\ell_2 - \ell_3)v^2, \tag{29}$$

while the CP even scalar mass matrix is

$$M_n = \begin{pmatrix} \ell_1 v_1^2 + \frac{1}{2}\ell_3 v_2^2 + \ell_2(v_1^2 - v_2^2) & \frac{1}{2}(2\ell_1 + \ell_3)v_1 v_2 \\ \frac{1}{2}(2\ell_1 + \ell_3)v_1 v_2 & \ell_1 v_2^2 + \frac{1}{2}\ell_3 v_1^2 - \ell_2(v_1^2 - v_2^2) \end{pmatrix}. \tag{30}$$

Its trace and determinant are

$$m_h^2 + m_H^2 = \text{Tr}(M_n) = \frac{1}{2}(2\ell_1 + \ell_3)v^2 \tag{31}$$

$$m_h^2 m_H^2 = \text{Det}(M_n) = -\frac{1}{2}(\ell_1 + \ell_2)(2\ell_2 - \ell_3)(v_1^2 - v_2^2)^2. \tag{32}$$

Notice that, for negative ℓ_2 such that $-\ell_1 < \ell_2 < \ell_3/2$, the requirement that squared masses are positive can be made consistent with the bounded from below conditions in Eq. (22). Moreover, Eqs. (25) cannot be obtained as the limit $v_1, v_2 \rightarrow v/\sqrt{2}$ of Eqs. (28)-(32) – Eq. (32) would lead to a massless scalar; this is a singular case which must be treated separately. Eqs. (28), (29), and (31) can be inverted, to yield ℓ_1 , ℓ_2 , and ℓ_3 in terms of the physical masses:

$$\begin{aligned}
\ell_1 v^2 &= m_h^2 + m_H^2 + m_{H^\pm}^2 - m_A^2, \\
\ell_2 v^2 &= -m_{H^\pm}^2, \\
\ell_3 v^2 &= 2(m_A^2 - m_{H^\pm}^2).
\end{aligned} \tag{33}$$

Substituting into Eq. (32), we obtain

$$\cos^2(2\beta) = \frac{m_h^2 m_H^2}{m_A^2(m_h^2 + m_H^2 - m_A^2)}. \tag{34}$$

The diagonalization of M_n is performed through the transformation

$$\begin{pmatrix} \text{Re } \phi_1^0 \\ \text{Re } \phi_2^0 \end{pmatrix} = \begin{pmatrix} \rho_1 \\ \rho_2 \end{pmatrix} = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} H \\ h \end{pmatrix}. \tag{35}$$

² Recall that when the text refers to (v_1, v_2) what is really meant is $\langle \phi_k^0 \rangle = v_k/\sqrt{2}$ ($k = 1, 2$), and that this is what we use in the equations.

We find

$$\tan(2\alpha) = \frac{2\ell_1 + \ell_3}{2\ell_1 + 4\ell_2 - \ell_3} \frac{2v_1 v_2}{v_1^2 - v_2^2} = \frac{m_h^2 + m_H^2}{m_h^2 + m_H^2 - 2m_A^2} \tan(2\beta). \quad (36)$$

Using $\tan^2(2\beta) = 1/\cos^2(2\beta) - 1$ and Eq. (34), one can write $\tan(2\alpha)$ in terms of masses alone. In the real 2HDM, the coupling of the lighter scalar to two vector bosons is given by $g_{hVV} = g_{hVV}^{\text{SM}} \sin(\beta - \alpha)$, where g_{hVV}^{SM} is the SM coupling. Measurements of these couplings at LHC constrain $\beta - \alpha$ to lie very close to $\pi/2$ – the so-called alignment limit. Through Eqs. (34) and (36), this places tight constraints on the masses of A and H . In the exact alignment limit, we would be forced into $m_A = 0$. Thus, in this case the pseudoscalar must be light, running into constraints from both LEP and LHC. A full phenomenological analysis of this case is beyond the scope of this work.

We now turn to the S_3 potential with the most general real soft violations of S_3 :

$$V = \mu_1^2 (|\phi_2|^2 + |\phi_1|^2) - \mu_2^2 (\phi_1^\dagger \phi_2 + \phi_2^\dagger \phi_1) - \mu_3^2 (|\phi_2|^2 - |\phi_1|^2) \\ + \frac{1}{2} \ell_1 (|\phi_2|^2 + |\phi_1|^2)^2 + \frac{1}{2} \ell_2 (|\phi_2|^2 - |\phi_1|^2)^2 + \ell_3 (\phi_1^\dagger \phi_2) (\phi_2^\dagger \phi_1). \quad (37)$$

The stationarity conditions become

$$0 = \frac{\partial V}{\partial v_1} = (\mu_1^2 + \mu_3^2) v_1 - \mu_2^2 v_2 + \frac{v_1}{2} (\ell_1 v^2 + \ell_2 (v_1^2 - v_2^2) + \ell_3 v_2^2), \\ 0 = \frac{\partial V}{\partial v_2} = (\mu_1^2 - \mu_3^2) v_2 - \mu_2^2 v_1 + \frac{v_2}{2} (\ell_1 v^2 - \ell_2 (v_1^2 - v_2^2) + \ell_3 v_1^2), \quad (38)$$

and $v_1 = v_2$ is no longer a minimum. Although it is technically natural to have μ_3^2 small (since in the $\mu_3^2 = 0$ limit the potential gets an extra Z_2 symmetry), in which case $v_1 \sim v_2$, we will allow any value for μ_3^2 . Repeating the previous steps, we find

$$m_{H^\pm}^2 = -\ell_2 v^2 - 2\mu_3^2 \sec(2\beta), \quad (39)$$

$$m_A^2 = -\frac{1}{2} [(2\ell_2 - \ell_3) v^2 + 4\mu_3^2 \sec(2\beta)], \quad (40)$$

$$T \equiv m_h^2 + m_H^2 = \frac{1}{2} [(2\ell_1 + \ell_3) v^2 - 4\mu_3^2 \sec(2\beta)], \quad (41)$$

$$D \equiv m_h^2 m_H^2 = -\frac{v^2}{2} [(2\ell_2 - \ell_3) \cos(2\beta) ((\ell_1 + \ell_2) v^2 \cos(2\beta) + 2\mu_3^2) + 2(2\ell_1 + \ell_3) \mu_3^2 \sec(2\beta)]. \quad (42)$$

Hence,

$$\ell_1 v^2 = m_h^2 + m_H^2 + m_{H^\pm}^2 - m_A^2 + 2\mu_3^2 \sec(2\beta), \\ \ell_2 v^2 = -m_{H^\pm}^2 - 2\mu_3^2 \sec(2\beta), \\ \ell_3 v^2 = 2(m_A^2 - m_{H^\pm}^2). \quad (43)$$

These results reproduce the corresponding ones in the $\mu_3^2 \rightarrow 0$ limit. Substituting into Eq. (42) yields

$$m_h^2 m_H^2 = m_A^2 (m_h^2 + m_H^2 - m_A^2) \cos^2(2\beta) \\ - 2\mu_3^2 (m_h^2 + m_H^2) \cos(2\beta) \tan^2(2\beta) - 4(\mu_3^2)^2 \tan^2(2\beta), \quad (44)$$

providing a complicated relation between μ_3 and β . Also

$$\tan(2\alpha) = \frac{m_h^2 + m_H^2 + 4\mu_3^2 \sec(2\beta)}{m_h^2 + m_H^2 - 2m_A^2} \tan(2\beta). \quad (45)$$

Thus, for the potential of Eq. (37), the independent parameters may be taken as v^2 , β , m_h^2 , m_H^2 , m_A^2 , and $m_{H^\pm}^2$; μ_3^2 is found from Eq. (44) and α from Eq. (45). Combining Eqs. (44) and (45), we find

$$\frac{D}{c_{2\beta}^2} = m_A^2 (T - m_A^2) + \frac{T^2}{4} t_{2\beta}^2 - \left(\frac{T}{2} - m_A^2 \right)^2 t_{2\alpha}^2. \quad (46)$$

Even in the exact alignment limit ($\beta = \alpha + \pi/2$), this reduces to $m_A^2 (T - m_A^2) = D$ which has two solutions: $m_A^2 = m_h^2$; and the definitely allowed $m_A^2 = m_H^2$, consistent with the decoupling limit.

So far, we have considered only real parameters and vevs in the scalar sector. We now turn briefly to the possibility that they are complex. Since the S_3 symmetry forces $\lambda_5 = \lambda_6 = \lambda_7 = 0$, one can only introduce phases via soft breaking, by giving a phase χ to μ_2^2 and changing in Eq. (37) the term

$$-\mu_2^2(\phi_1^\dagger\phi_2 + \phi_2^\dagger\phi_1) \rightarrow -\mu_2^2(e^{i\chi}\phi_1^\dagger\phi_2 + e^{-i\chi}\phi_2^\dagger\phi_1). \quad (47)$$

The most general vev may be written as $v_1, v_2 e^{i\delta}$. Minimization with respect to δ ,

$$0 = \frac{\partial V}{\partial \delta} = \mu_2^2 v_1 v_2 \sin(\delta + \chi), \quad (48)$$

forces³ $\delta = -\chi$, while the other minimization conditions, in Eqs. (38) remain the same. Thus, we can only have complex vevs with a complex soft-breaking term.

III. YUKAWA COUPLINGS

We now turn to the Yukawa couplings, following closely the notation of Ref. [15]:

$$-\mathcal{L}_Y = \bar{q}_L(\Gamma_1\Phi_1 + \Gamma_2\Phi_2)n_R + \bar{q}_L(\Delta_1\tilde{\Phi}_1 + \Delta_2\tilde{\Phi}_2)p_R + \text{h.c.}, \quad (49)$$

where $q_L = (p_L, n_L)^\top$ is a vector in the 3-dimensional family space of left-handed doublets, and n_R (p_R) is a vector in the 3-dimensional right-handed space of charge $-1/3$ ($+2/3$) quarks. The complex 3×3 matrices $\Gamma_1, \Gamma_2, \Delta_1$, and Δ_2 contain the Yukawa couplings. In general, these matrices are not diagonal. We denote by U_α ($\alpha = d_L, d_R, u_L, u_R$) the matrices taking the quarks into the mass basis:

$$\begin{aligned} n_R &= U_{d_R} d_R, & p_R &= U_{u_R} u_R, \\ \bar{q}_L &= (\bar{p}_L, \bar{n}_L) = (\bar{u}_L U_{u_L}^\dagger, \bar{d}_L U_{d_L}^\dagger) = (\bar{u}_L V, \bar{d}_L) U_{d_L}^\dagger, \end{aligned} \quad (50)$$

where $V = U_{u_L}^\dagger U_{d_L}$ is the CKM matrix. The mass matrices become

$$\begin{aligned} \text{diag}(m_d, m_s, m_b) &= D_d = \frac{1}{\sqrt{2}} U_{d_L}^\dagger [v_1 \Gamma_1 + v_2 \Gamma_2] U_{d_R}, \\ \text{diag}(m_u, m_c, m_t) &= D_u = \frac{1}{\sqrt{2}} U_{u_L}^\dagger [v_1 \Delta_1 + v_2 \Delta_2] U_{u_R}, \end{aligned} \quad (51)$$

while eventual flavour changing neutral scalar interactions are controlled by

$$\begin{aligned} N_d &= \frac{1}{\sqrt{2}} U_{d_L}^\dagger [-v_2 \Gamma_1 + v_1 \Gamma_2] U_{d_R}, \\ N_u &= \frac{1}{\sqrt{2}} U_{u_L}^\dagger [-v_2 \Delta_1 + v_1 \Delta_2] U_{u_R}. \end{aligned} \quad (52)$$

In this section, we will absorb the $1/\sqrt{2}$ in the definitions of the parameters in the Yukawa matrices, and work with

$$Y_d = v_1 \Gamma_1 + v_2 \Gamma_2, \quad Y_u = v_1 \Delta_1 + v_2 \Delta_2, \quad (53)$$

and the hermitian matrices

$$\begin{aligned} H_d &= Y_d Y_d^\dagger = U_{d_L} \text{diag}(m_d^2, m_s^2, m_b^2) U_{d_L}^\dagger, \\ H_u &= Y_u Y_u^\dagger = U_{u_L} \text{diag}(m_u^2, m_c^2, m_t^2) U_{u_L}^\dagger. \end{aligned} \quad (54)$$

These matrices are diagonalized respectively by the matrices U_{d_L} and U_{u_L} , which must be found when fitting for the CKM matrix. This step is not needed for CP violation, which may be found directly from [16]

$$J = \text{Det}(H_d H_u - H_u H_d), \quad (55)$$

³ The other possibilities – $\mu_2^2 = 0$, or inert vacua $v_1 = 0$ or $v_2 = 0$ – will not be pursued here.

in a basis independent fashion.

Henceforth, we use the complex representation. We recall that, for the corresponding entry of the Yukawa coupling matrix to be non-vanishing, the Yukawa Lagrangian must be in the invariant singlet representation $\mathbf{1}$. We start by assuming that the two scalar fields are in singlet representations, which we denoted before by $\Phi \sim \mathbf{s}$. Thus, the product of left-handed (q_L) and right-handed (charge $+2/3$, p_R , or charge $-1/3$, n_R) fermions must also be in a singlet representation. This can be achieved by two doublets or by two singlets. Since permutations of the three fields in each sector do not lead to new structures for the Yukawa matrices, we will denote by $f \sim \mathbf{s}$ (where $f = q_L, p_R, n_R$) the following independent possibilities for the fields in each of the three generations:

$$\begin{aligned} &(\mathbf{1}, \mathbf{1}, \mathbf{1}), & (\mathbf{1}, \mathbf{1}, \mathbf{1}'), \\ &(\mathbf{1}, \mathbf{1}', \mathbf{1}'), & (\mathbf{1}', \mathbf{1}', \mathbf{1}'). \end{aligned} \tag{56}$$

Similarly, we will denote by $f \sim \mathbf{d}$ the following independent possibilities for the fields in each of the three generations:

$$(\mathbf{2}, \mathbf{1}), \quad (\mathbf{2}, \mathbf{1}'). \tag{57}$$

Let us now consider the possibility that the two scalar fields are in the doublet representation of S_3 , which we denoted before by $\Phi \sim \mathbf{2}$. Thus, the product of left-handed (q_L) and right-handed (charge $+2/3$, p_R , or charge $-1/3$, n_R) fermions must also be in a doublet representation. But this can be achieved by having both quark fields in doublets, or by having one in a doublet and another in a singlet. The possibilities for the various representations are listed in Table I. The first case (with all fields in singlets), reduces to the analysis of Z_2 and has been discussed in

Φ	\bar{q}_L	n_R	p_R
\mathbf{s}	\mathbf{s}	\mathbf{s}	\mathbf{s}
\mathbf{s}	\mathbf{d}	\mathbf{d}	\mathbf{d}
$\mathbf{2}$	\mathbf{d}	\mathbf{d}	\mathbf{d}
$\mathbf{2}$	\mathbf{d}	\mathbf{s}	\mathbf{d}
$\mathbf{2}$	\mathbf{d}	\mathbf{d}	\mathbf{s}
$\mathbf{2}$	\mathbf{d}	\mathbf{s}	\mathbf{s}
$\mathbf{2}$	\mathbf{s}	\mathbf{d}	\mathbf{d}

TABLE I: Possible representations for the scalar fields (Φ), the left-handed quark $SU(2)_L$ doublets (q_L), and the right-handed quark $SU(2)_L$ singlets (p_R and n_R).

detail in Ref. [7], where are possible implementations with abelian groups are presented.

A. Example 1: Φ in singlet; fermions in doublets

As a specific example, let us consider the possibility

$$\Phi \sim (\mathbf{1}, \mathbf{1}'), \quad \bar{q}_L \sim (\mathbf{2}, \mathbf{1}), \quad n_R \sim (\mathbf{2}, \mathbf{1}), \quad p_R \sim (\mathbf{2}, \mathbf{1}). \tag{58}$$

Because Φ is in a singlet, the product of fermions (left and right) must also be in a singlet. Using \bar{q}_L in the doublet, the product of two doublets $\bar{q}_L n_R$ in $\mathbf{2} \otimes \mathbf{2}$ is

$$\left. \begin{pmatrix} \bar{q}_{L1} \\ \bar{q}_{L2} \end{pmatrix} \otimes \begin{pmatrix} n_{R1} \\ n_{R2} \end{pmatrix} \right|_{\mathbf{1}, \mathbf{1}'} = \bar{q}_{L1} n_{R2} \pm \bar{q}_{L2} n_{R1}. \tag{59}$$

Thus, the products with the scalars into a singlet are⁴

$$\Phi_1 (\bar{q}_{L1} n_{R2} + \bar{q}_{L2} n_{R1}), \tag{60}$$

$$\Phi_2 (\bar{q}_{L1} n_{R2} - \bar{q}_{L2} n_{R1}). \tag{61}$$

⁴ We will often show the product of fields in the order that highlights the S_3 properties, even if it violates the correct $SU(2)_L$ order of $\bar{q}_L \Phi f_R$.

The remaining non vanishing term comes from

$$\Phi_1 \bar{q}_{L3} n_{R3}. \quad (62)$$

Multiplying Eqs. (60), (61), and (62) by complex coefficients a , b , and c , respectively, we find

$$Y_d = \begin{bmatrix} 0 & av_1 + bv_2 & 0 \\ av_1 - bv_2 & 0 & 0 \\ 0 & 0 & cv_1 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} av_1 - bv_2 & 0 & 0 \\ 0 & av_1 + bv_2 & 0 \\ 0 & 0 & cv_1 \end{bmatrix}. \quad (63)$$

This corresponds to the charged lepton sector of the model discussed in Refs. [9, 10], where neutrino mixing was not considered. We notice that the matrix in Eq. (63) is block diagonal. For the same reason, so would be the corresponding matrix for the up quarks, leading to a block diagonal CKM matrix, in contradiction with experiment. Thus, these S_3 assignments cannot be used for the quarks⁵. Because Φ is in a singlet and only one generation of quarks in each sector is in a singlet, this block-diagonal problem remains, regardless of the assignments of such singlets to $\mathbf{1}$ or $\mathbf{1}'$. Thus, we conclude that the case on the second line of Table I, cannot be implemented in the quark sector.

B. Example 2: doublets in all sectors

We now turn to

$$\Phi \sim \mathbf{2}, \quad \bar{q}_L \sim (\mathbf{2}, \mathbf{1}), \quad n_R \sim (\mathbf{2}, \mathbf{1}), \quad p_R \sim (\mathbf{2}, \mathbf{1}). \quad (64)$$

Because Φ is in a doublet, the product of fermions (left and right) must also be in a doublet. Using \bar{q}_L in the doublet, the product of two doublets $\bar{q}_L n_R$ in $\mathbf{2} \otimes \mathbf{2}$ is

$$\left(\begin{array}{c} \bar{q}_{L1} \\ \bar{q}_{L2} \end{array} \right) \otimes \left(\begin{array}{c} n_{R1} \\ n_{R2} \end{array} \right) \Big|_{\mathbf{2}} = \left(\begin{array}{c} \bar{q}_{L2} n_{R2} \\ \bar{q}_{L1} n_{R1} \end{array} \right). \quad (65)$$

The product with the scalar doublet into a singlet is

$$\left(\begin{array}{c} \Phi_1 \\ \Phi_2 \end{array} \right) \otimes \left(\begin{array}{c} \bar{q}_{L2} n_{R2} \\ \bar{q}_{L1} n_{R1} \end{array} \right) \Big|_{\mathbf{1}} = \Phi_1 \bar{q}_{L1} n_{R1} + \Phi_2 \bar{q}_{L2} n_{R2}, \quad (66)$$

as mentioned after Eq. (16). For a n_{R3} in a singlet, we find

$$\left(\begin{array}{c} \Phi_1 \\ \Phi_2 \end{array} \right) \otimes \left(\begin{array}{c} \bar{q}_{L1} \\ \bar{q}_{L2} \end{array} \right) \Big|_{\mathbf{1}} \otimes n_{R3} = \Phi_1 \bar{q}_{L2} n_{R3} + \Phi_2 \bar{q}_{L1} n_{R3}. \quad (67)$$

Finally, for a \bar{q}_{L3} in a singlet, we find

$$\bar{q}_{L3} \otimes \left(\begin{array}{c} \Phi_1 \\ \Phi_2 \end{array} \right) \otimes \left(\begin{array}{c} n_{R1} \\ n_{R2} \end{array} \right) \Big|_{\mathbf{1}} = \Phi_1 \bar{q}_{L3} n_{R2} + \Phi_2 \bar{q}_{L3} n_{R1}. \quad (68)$$

Since Φ is in the doublet, d_L and n_R cannot be simultaneously in the singlet. Thus, multiplying Eqs. (66), (67), and (68) by complex coefficients a , b , and c , respectively, we find

$$Y_d = \begin{bmatrix} av_1 & 0 & bv_2 \\ 0 & av_2 & bv_1 \\ cv_2 & cv_1 & 0 \end{bmatrix}. \quad (69)$$

⁵ In Refs. [9, 10] this problem is solved by keeping all quark fields in singlets; moreover, with the same assignment ($\mathbf{1}$ or $\mathbf{1}'$) for all generations within each sector. This reduces the quark sector to the usual Z_2 cases.

The matrix for the up quarks is easily found by noting that the doublet $(\Phi_1, \Phi_2)^\top$ gets substituted by $(\tilde{\Phi}_2, \tilde{\Phi}_1)^\top$, corresponding to a $v_1 \leftrightarrow v_2^*$ change. We find

$$Y_u = \begin{bmatrix} xv_2^* & 0 & yv_1^* \\ 0 & xv_1^* & yv_2^* \\ zv_1^* & zv_2^* & 0 \end{bmatrix}, \quad (70)$$

with complex coefficients x , y , and z . Besides v and $\tan\beta$, the matrices involve 6 complex coefficients, which might conceivably fit the ten known data: the 6 quark masses and the 4 parameters in the CKM matrix. We have used Eqs. (54) to check that we can generate all masses different and nonzero, and Eq. (55) to show that we can generate a nonzero CP violating phase. We note that $J \neq 0$ even if one takes the vevs to be real, implying that this model does not coincide with the CP3 model with quarks presented in Ref. [8], where a complex vev was needed in order to get a non-vanishing J . This is a further illustration of a sometimes unappreciated point: two symmetries which lead to the same scalar potential, may lead to very different models when extended into the Yukawa sector. A complete analysis of each model is beyond the scope of this work. We are interested here in mapping all possibilities consistent with S_3 which do not lead necessarily into vanishing and/or degenerate masses, or to the absence of CP violation.

We must now turn to the possibility that some among \bar{q}_{L3} , n_{L3} , and/or p_{L3} are in a $\mathbf{1}'$. Using $\eta_\alpha = +1$ when the corresponding field is in $\mathbf{1}$, and $\eta_\alpha = -1$ when the corresponding field is in $\mathbf{1}'$, we find

$$Y_d = \begin{bmatrix} av_1 & 0 & \eta_d bv_2 \\ 0 & av_2 & bv_1 \\ \eta_q cv_2 & cv_1 & 0 \end{bmatrix},$$

$$Y_u = \begin{bmatrix} xv_2^* & 0 & \eta_u yv_1^* \\ 0 & xv_1^* & yv_2^* \\ \eta_q zv_1^* & zv_2^* & 0 \end{bmatrix}. \quad (71)$$

These are all forms consistent with the assignments on the third line of Table I.

C. Example 3: singlet only on right-handed sectors

Let us consider

$$\Phi \sim \mathbf{2}, \quad \bar{q}_L \sim (\mathbf{2}, \mathbf{1}), \quad n_R \sim \mathbf{s}, \quad p_R \sim (\mathbf{2}, \mathbf{1}). \quad (72)$$

Because Φ is in a doublet, the product of fermions (left and right) must also be in a doublet. Since \bar{q}_{L3} and all n_R are in a singlet, the last line of the matrix Y_d vanishes. But this implies that the matrix H_d only has non-vanishing entries in the (1,2) sector, leading to a massless down quark. Similarly,

$$\Phi \sim \mathbf{2}, \quad \bar{q}_L \sim (\mathbf{2}, \mathbf{1}), \quad n_R \sim (\mathbf{2}, \mathbf{1}), \quad p_R \sim \mathbf{s}, \quad (73)$$

leads to a massless up quark. A combination of both problems occurs in

$$\Phi \sim \mathbf{2}, \quad \bar{q}_L \sim (\mathbf{2}, \mathbf{1}), \quad n_R \sim \mathbf{s}, \quad p_R \sim \mathbf{s}, \quad (74)$$

Thus, under the conditions considered in this paper of S_3 in a 2HDM, these three cases are ruled out.

D. Example 4: singlet only on left-handed sector

The last case to be considered is

$$\Phi \sim \mathbf{2}, \quad \bar{q}_L \sim \mathbf{s}, \quad n_R \sim (\mathbf{2}, \mathbf{1}), \quad p_R \sim (\mathbf{2}, \mathbf{1}). \quad (75)$$

Because Φ is in a doublet, the product of fermions (left and right) must also be in a doublet. Since all \bar{q}_L and n_{R3} are in a singlet, the last column of the matrix Y_d vanishes. But this implies that the matrix $Y_d^\dagger Y_d$ (which has the same eigenvalues as H_d) only has non-vanishing entries in the (1,2) sector, leading to a massless down quark. Thus, this case is also ruled out.

E. Flavour-changing Higgs couplings

Here, we comment briefly on the appearance of Flavour Changing Neutral Scalar Interactions (FCNSI). As is well known, large FCNSI with small scalar masses are precluded by measurements in the neutral meson systems. There are several ways to solve this problem. One route is to endow the heavy neutral states with large masses. This is achieved in the decoupling limit, with the added bonus that it implies the alignment limit $\beta \sim \alpha + \pi/2$, which is necessary to conform with the LHC measurements of the 125 GeV scalar. Since this is always a possibility, FCNSI cannot be used to exclude models from the realm of possibilities.

A second route, introduced by Glashow and Weinberg [17] and independently by Paschos [18], is to impose a Z_2 symmetry under which all fields of equal charge transform equally, thus implying vanishing FCNSI. Neglecting right-handed neutrinos, there are only four such possibilities – for a review, see, for example, Refs. [3, 4]. When all fields are in S_3 singlets, the situation reduces to Z_2 . When all fields of equal charge transform equally, there are no FCNSI. The cases where some fields of equal charge transform differently were discussed in Ref. [7] and do lead to FCNSI. Similarly, when there are S_3 doublets in all sectors, we do have FCNSI.

A third route is that the FCNSI are accidentally small. This may be theoretically displeasing, but cannot be excluded. A fourth route, discussed by Branco, Lavoura, and Grimus (BGL), is that the FCNSI are small because they are related with the CKM matrix elements [19]. We have checked that the models with S_3 doublets in all sectors are not of the BGL type [20].

IV. CONCLUSIONS

We study all representation assignments of S_3 in the 2HDM with quarks, consistent with the basis requirements of non-vanishing, non-degenerate masses, non-block diagonal CKM matrix and the presence of a CP violating phase. We found that there are only two implementations consistent with this simple requirements: all fields are in singlets or, else, all fields sectors have a doublet representation.

When the scalars are in an S_3 doublet, one must introduce soft-breaking terms. Even with the most general real soft-breaking term, there is a relation between α and β , shown in Eq. (46). As far as we know, this is a new result.

Finally, we point out that the viable models with doublet representations are not of the BGL type [19], where the flavour changing scalar interactions are naturally small because they are related to the CKM matrix elements [20]. It was shown in Ref. [7] that the only models based on abelian symmetries which have this property are of the type introduced originally in Ref. [19]. We now know that the non-abelian S_3 group does not provide a further example of a BGL model.

Acknowledgments

We are grateful to I. Ivanov for discussions and to D. Emmanuel-Costa for carefully reading the manuscript and for useful suggestions. This work is supported in part by the Portuguese *Fundação para a Ciência e Tecnologia* (FCT) under contract UID/FIS/00777/2013. D. C. is partly supported by the Brazilian National Council for Scientific and Technological Development (CNPq), under grants 484157/2013-2 and 201066/2015-7

-
- [1] G. Aad *et al.* [ATLAS Collaboration], “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,” *Phys. Lett. B* **716**, 1 (2012) [arXiv:1207.7214 [hep-ex]].
 - [2] S. Chatrchyan *et al.* [CMS Collaboration], “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC,” *Phys. Lett. B* **716**, 30 (2012) [arXiv:1207.7235 [hep-ex]].
 - [3] J.F. Gunion, H.E. Haber, G.L. Kane and S. Dawson, *The Higgs Hunter’s Guide* (Westview Press, Boulder, CO, 2000).
 - [4] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher, and J. P. Silva, *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rept.* **516**, 1 (2012) [arXiv:1106.0034 [hep-ph]].
 - [5] I. P. Ivanov, “Minkowski space structure of the Higgs potential in 2HDM. II. Minima, symmetries, and topology,” *Phys. Rev. D* **77**, 015017 (2008) [arXiv:0710.3490 [hep-ph]].
 - [6] P. M. Ferreira, H. E. Haber and J. P. Silva, “Generalized CP symmetries and special regions of parameter space in the two-Higgs-doublet model,” *Phys. Rev. D* **79**, 116004 (2009) [arXiv:0902.1537 [hep-ph]].
 - [7] P. M. Ferreira and J. P. Silva, “Abelian symmetries in the two-Higgs-doublet model with fermions,” *Phys. Rev. D* **83**, 065026 (2011) [arXiv:1012.2874 [hep-ph]].

- [8] P. M. Ferreira and J. P. Silva, “A Two-Higgs Doublet Model With Remarkable CP Properties,” *Eur. Phys. J. C* **69**, 45 (2010) [arXiv:1001.0574 [hep-ph]].
- [9] Y. Kajiyama, H. Okada and K. Yagyu, “Electron/Muon Specific Two Higgs Doublet Model,” *Nucl. Phys. B* **887**, 358 (2014) [arXiv:1309.6234 [hep-ph]].
- [10] A. R. Johansen and M. Sher, “Electron/muon specific two Higgs doublet model at e^+e^- colliders,” *Phys. Rev. D* **91**, 054021 (2015) [arXiv:1502.00516 [hep-ph]].
- [11] E. Ma, “Non-Abelian discrete family symmetries of leptons and quarks,” hep-ph/0409075.
- [12] E. Ma and B. Melic, “Updated S_3 model of quarks,” *Phys. Lett. B* **725**, 402 (2013) [arXiv:1303.6928 [hep-ph]].
- [13] N.G. Deshpande and E. Ma, “Pattern Of Symmetry Breaking With Two Higgs Doublets,” *Phys. Rev. D* **18**, 2574 (1978).
- [14] D. Emmanuel-Costa, private communication.
- [15] M. Nebot and J. P. Silva, “Self-cancellation of a scalar in neutral meson mixing and implications for the LHC,” *Phys. Rev. D* **92**, no. 8, 085010 (2015) [arXiv:1507.07941 [hep-ph]].
- [16] C. Jarlskog, “Commutator of the Quark Mass Matrices in the Standard Electroweak Model and a Measure of Maximal CP Violation,” *Phys. Rev. Lett.* **55**, 1039 (1985); “A Priori Definition of Maximal CP Violation,” I. Dunietz, O. W. Greenberg, and D.-D. Wu, *Phys. Rev. Lett.* **55**, 2935 (1985); F. J. Botella and L.-L. Chau, “Anticipating the Higher Generations of Quarks from Rephasing Invariance of the Mixing Matrix,” *Phys. Lett. B* **168**, 97 (1986); J. Bernabéu, G. C. Branco, and M. Gronau, “Cp Restrictions On Quark Mass Matrices,” *Phys. Lett. B* **169**, 243 (1986).
- [17] S. L. Glashow and S. Weinberg, “Natural Conservation Laws for Neutral Currents,” *Phys. Rev. D* **15**, 1958 (1977).
- [18] E. A. Paschos, “Diagonal Neutral Currents,” *Phys. Rev. D* **15**, 1966 (1977).
- [19] G. C. Branco, W. Grimus and L. Lavoura, *Phys. Lett. B* **380**, 119 (1996) [hep-ph/9601383].
- [20] We are grateful to F. Botella for discussions on this point.