Hot and dense water in the inner 25 AU of SVS13-A

C. Codella¹*, C. Ceccarelli^{2,3,1}, E. Bianchi^{1,4}, L. Podio¹, R. Bachiller⁵, B. Lefloch^{2,3}, F. Fontani¹, V. Taquet⁶, L. Testi^{1,7}

- ¹ INAF-Osservatorio Astrofisico di Arcetri, L.go E. Fermi 5, Firenze, 50125, Italy
- ² Univ. Grenoble Alpes, IPAG, F-38000 Grenoble, France
- ³ CNRS, IPAG, F-38000 Grenoble, France
- ⁴ Dipartimento di Fisica e Astronomia, Università degli Studi di Firenze, Italy
- ⁵ IGN, Observatorio Astronómico Nacional, Calle Alfonso XII, 28004 Madrid, Spain
- ⁶ Leiden Observatory, Leiden University, 9513, 2300-RA Leiden, The Netherlands
- ⁷ ESO, Karl Schwarzchild Srt. 2, 85478 Garching bei München, Germany

Accepted date. Received date; in original form date

ABSTRACT

In the context of the ASAI (Astrochemical Surveys At IRAM) project, we carried out an unbiased spectral survey in the millimeter window towards the well known low-mass Class I source SVS13-A. The high sensitivity reached (3–12 mK) allowed us to detect at least 6 HDO broad (FWHM $\sim 4–5~{\rm km~s^{-1}})$ emission lines with upper level energies up to $E_{\rm u}=837~{\rm K}.$ A non-LTE LVG analysis implies the presence of very hot (150–260 K) and dense ($\geqslant 3\times 10^7~{\rm cm^{-3}})$ gas inside a small radius ($\sim 25~{\rm AU}$) around the star, supporting, for the first time, the occurrence of a hot corino around a Class I protostar.

The temperature is higher than expected for water molecules are sublimated from the icy dust mantles (~ 100 K). Although we cannot exclude we are observing the effects of shocks and/or winds at such small scales, this could imply that the observed HDO emission is tracing the water abundance jump expected at temperatures $\sim 220-250$ K, when the activation barrier of the gas phase reactions leading to the formation of water can be overcome. We derive $X(HDO) \sim 3 \times 10^{-6}$, and a H₂O deuteration $\geq 1.5 \times 10^{-2}$, suggesting that water deuteration does not decrease as the protostar evolves from the Class 0 to the Class I stage.

Key words: Molecular data – Stars: formation – radio lines: ISM – submillimetre: ISM – ISM: molecules

1 INTRODUCTION

The origin of terrestrial water is still a source of intense debate (e.g. Ceccarelli et al. 2014; van Dischoeck et al. 2014; Altwegg et al. 2015). A key element to shed light on it, is how water evolves with time in proto-Sun analogues. Specifically, two aspects are particulary important: (1) the amount of water and its distribution in the planet formation region (a few tens of AU) of the proto-Suns, and (2) its deuterium fractionation (e.g. Ceccarelli et al. 2014; Willacy et al. 2015).

With respect to the first point, water has been detected at all stages of the Sun-like star formation process, from prestellar cores and Class 0 sources to the Solar System (e.g. Caselli et al. 2012; Ceccarelli et al. 1999; van Dishoeck et al. 2011, 2014). However, so far, we mostly have poor angular resolution observations that allowed us to detect the water

emission, but not to resolve it on small (\leq 1000 AU) scales. Only a handful of observations exists with enough spatial resolution. They show that the water emission in the envelopes of Class 0 sources is concentrated in small regions, called hot corinos (Codella et al. 2010; Persson et al. 2013, 2014; Taquet et al. 2013; Coutens et al. 2014). On the contrary, no spatially resolved observations exist for the more evolved protostars, the Class I sources. With respect to the water deuterium fractionation, again only a few measures are available in Class 0 sources (Coutens et al. 2012, 2013, 2014; Persson et al. 2013, 2014; Taquet et al. 2013), but none in Class I.

In the context of the ASAI (Astrochemical Surveys At IRAM: http://www.oan.es/asai/) project, we have carried out a systematic study of the molecular emission towards SVS13-A. This is a well studied young stellar object located in the NGC1333 star-forming region, at a distance of ~ 235 pc (Bachiller et al. 1998; Hirota et al. 2008; Lee et al. 2016).

SVS13-A is part of the system SVS13, where three millimeter sources have been identified by interferometric observations (Bachiller et al. 1998; Looney et al. 2000), called A, B, and C. The distance between A and B is 15", while C is 20" away from A. The systemic velocity of the sources A and B is between +8 km s⁻¹ and +9 km s⁻¹ (Chen et al. 2009; Lopèz-Sepulcre et al. 2015). The luminosity of SVS13-A has been estimated to be 34 L $_{\odot}$ (Chen et al. 2009; Tobin et al. 2016), where we corrected for the new estimate of the distance d=235 pc (Hirota et al. 2008).

Although SVS13-A is still deeply embedded in a large-scale (\sim 6000 AU; e.g. Lefloch et al. 1998) envelope, its extended (>0.07 outflow pc) outflow, associated with the HH7-11 chain (e.g. Lefloch et al. 1998, and references therein), and its low L_{submm}/L_{bol} ratio (\sim 0.8 %) lead to the classification as a Class I source (e.g. Chen et al. 2009 and references therein).

In this Letter, we report the detection of several lines of HDO towards SVS13-A, providing the first detection of deuterated water in a Class I source.

2 OBSERVATIONS

The present observations have been performed during several runs between 2012 and 2014 with the IRAM 30-m telescope near Pico Veleta (Spain) in the context of the Astrochemical Surveys At IRAM¹ (ASAI) Large Program. In particular, the unbiased spectral surveys at 3 mm (80116 GHz), 2 mm (129173 GHz), and 1.3 mm (200276 GHz) have been acquired using the EMIR receivers with a spectral resolution of 0.2 MHz. The observations were performed in wobbler switching mode with a throw of 180″ towards the coordinates of the SVS13-A object, namely $\alpha_{\rm J2000}=03^{\rm h}~29^{\rm m}~03^{\rm s}.29$, $\delta_{\rm J2000}=+31^{\circ}~16'~03''.8$). The pointing was checked by observing nearby planets or continuum sources and was found to be accurate to within 2″-3″. The HPBWs are in the 9″-31″ range.

The data were reduced with the GILDAS–CLASS² package. Calibration uncertainties are $\simeq 10\%$ at 3mm and $\sim 20\%$ at shorter wavelengths. All the spectra have been converted from antenna temperature to main beam temperature ($T_{\rm MB}$), using the main beam efficiencies reported on the IRAM 30-m website³.

3 RESULTS AND DISCUSSION

3.1 HDO detected lines

The ASAI unbiased survey allows us to detect 7 HDO lines (1 in the 3 mm band, 1 in the 2 mm band, and 5 in the 1.3 mm one) covering a wide range of excitation, with upper level energies $E_{\rm u}$ from 47 K to 837 K. The 81 GHz line is only tentatively detected, given the low S/N ratio. However, the following analysis will show how the $1_{1,0}-1_{1,1}$ intensity is well in agreement with those of the other lines observed at 2mm and 1.3mm. The $3_{3,1}-4_{2,2}$ transition (at $\sim 242.0~{\rm GHz})$

is the HDO line with the highest upper level energy (E₀=837 K) ever observed towards a low-mass protostar. The profiles of all detected lines are shown in Fig. 1, while Table 1 reports the results of the line Gaussian fits. The HDO emission peaks at the cloud systemic velocity, between +8.0 and +9.0 km $\rm s^{-1}$ (Chen et al. 2009; Lopèz-Sepulcre et al. 2015). The lines are quite broad, with a $FWHM \sim 4.2-4.9 \text{ km s}^{-1}$ for all the lines but the two observed at 3mm and 2mm, which are also those observed with the lowest S/N ratio and the worst spectral resolution (from 0.7 km s^{-1} to 0.2 km s^{-1} moving from 80.6 GHz and 266.2 GHz). The emission in the 1 and 2 mm bands only originates from SVS13-A, as SVS13-B, 15'' south-west, is outside the HPBW. The 3 mm band might, in principle, contain some emission from SVS13-B. However, the analysis of the measured fluxes tends to exclude a substantial contamination from SVS13-B also in this band (see below).

Finally, we searched for $\mathrm{H_{2}^{18}O}$ lines in our spectral survey and found none. The most sensitive upper limit to the $\mathrm{H_{2}O}$ column density is set by the non detection of the para— $\mathrm{H_{2}^{18}O}$ 3_{1,3}–2_{2,0} line at 203.40752 GHz⁴. We obtained a 3 σ upper limit on the peak temperature (in T_{MB} scale) of 20 mK.

3.2 Analysis of the HDO emission

We analysed the observed HDO line emission with the non-LTE LVG model by Ceccarelli et al. (2003), using the collisional coefficients for the system HDO-H₂ computed by Faure et al. (2012) and extracted from the the BASECOL database (Dubernet et al. 2013). We assumed a plane-parallel geometry and a Boltzmann distribution for the ortho-to-para H₂ ratio of 3. Note that the collisional coefficients with ortho-H₂ can be a factor 5 larger than the corresponding coefficients with para-H₂ (Faure et al. 2012), but only at low temperatures (\ll 45 K) and not at those here discussed (see below). Note also that the HDO 7_{3,4}–6_{4,3} line (with $E_{\rm u}=837$ K) has been excluded from the LVG analysis because the corresponding collisional rates have not been calculated (see later for a comparison with an LTE approach).

We run a large grid of models varying the temperature $T_{\rm kin}$ from 100 to 300 K, the H₂ density $n_{\rm H2}$ from 8×10^8 to $1 \times 10^{10} \text{ cm}^{-3}$, the HDO column density N(HDO) from 4×10^{16} to 7×10^{17} cm⁻³, and the emitting sizes θ_s from 0.05 to 10 ". The lowest χ_r^2 is obtained with $N(HDO) = 4\times10^{17}$ cm⁻², and $\theta_s = 0.2$, corresponding to ~ 50 AU. Figure 2 (upper panel) shows the χ^2_r contour plot as a function of the temperature and H₂ density with these values. The best fit solution is found for a very high temperature, $T_{\rm kin} = 150 - 260$ K, and a quite high density $n_{\rm H_2} \geqslant 3 \times 10^7 \ {\rm cm}^{-3}$. Figure 2 (lower panel) shows the goodness of the fit, namely the ratio between the measured velocity-integrated intensities and the LVG model predictions, as a function of the line upper level energy, for the best fit solution: $N(HDO) = 4 \times 10^{17} \text{ cm}^{-2}$, $\theta_s = 0.2, T_{\rm kin} = 200 \text{ K}, \text{ and } n_{\rm H_2} = 2 \times 10^8 \text{ cm}^{-3}.$ The lines are predicted to be optically thin to moderately thick. The largest opacities are ~ 2 for the four lowest lying lines (at

www.oan.es/asai

² http://www.iram.fr/IRAMFR/GILDAS

³ http://www.iram.es/IRAMES/mainWiki/Iram30mEfficiencies

⁴ from Jet Propulsion Laboratory database, http://spec.jpl.nasa.gov/home.html; Pickett et al. (1998)

Transition	$ u^{\rm a} $ (GHz)	HPBW (")	$g_{ m u}{}^a$	$E_{\rm u}{}^a$ (K)	$S\mu^{2a}$ (D ²)	$\log(A/s^{-1})^a$	rms (mK)	T_{peak}^{b} (mK)	V_{peak}^{b} (km s^{-1})	$FWHM^b$ (km s ⁻¹)	${I_{\rm int}}^b \\ ({\rm mK~km~s^{-1}})$
$1_{1,0}$ – $1_{1,1}$	80.57829	31	3	47	0.66	-5.88	5	14(2)	+9.2(0.3)	1.7(0.9)	26(9)
42,2-42,3	143.72721	17	9	319	0.73	-5.55	9	21(8)	+8.0(0.3)	3.1(0.8)	68(20)
$3_{1,2}$ – $2_{2,1}$	225.89672	11	7	168	0.69	-4.88	7	45(7)	+8.0(0.1)	4.9(0.3)	234(16)
$2_{1,1}$ – $2_{1,2}$	241.56155	10	5	95	0.36	-4.92	6	41(6)	+8.0(0.1)	4.7(0.3)	206(9)
73,4-64,3	241.97357	10	15	837	1.39	-4.82	7	17(5)	+7.7(0.3)	4.6(0.7)	83(10)
$5_{2,3}$ - $4_{3,2}$	255.05026	10	11	437	1.02	-4.75	6	40(6)	+7.8(0.1)	4.5(0.2)	199(9)
$2_{2,0}$ - $3_{1,3}$	266.16107	9	5	157	0.40	-4.76	12	31(9)	+7.4(0.3)	4.2(0.8)	141(19)

Table 1. List of HDO transitions and line properties (in $T_{\rm MB}$ scale) detected towards SVS13-A

80.58, 241.56, 266.16 and 225.90 GHz), while the other lines have opacities lower than unity.

Finally, the populations of the detected transitions are predicted to be close to LTE. Indeed, since non-LTE predictions were not possible for the higher lying line at 241.97 GHz, we also computed the LTE solution (see Fig. 3), finding a rotational temperature of 334 ± 42 K, which is larger than the kinetical temperature derived from the LVG analysis, i.e. 160-240 K considering the 1σ $\chi^2_{\rm r}$ solution. In practice, assuming the non-LTE LVG solution, the agreement between the predicted and observed intensity of the $E_{\rm u}=857$ K is within a factor 2; this is acceptable if we consider the opacity of the low- $E_{\rm u}$ lines and that the LTE condition may not apply to such a high lying line so that we cannot exclude a temperature gradient with a component with $T_{\rm kin}$ larger than 200 K. High spatial resolution observations are required to clarify this point.

3.3 Origin of the HDO emission

The non-LTE analysis reveals the presence of a hot (150-260 K), dense ($\geq 3 \times 10^7 \text{ cm}^{-3}$) and compact ($\sim 50 \text{ AU}$) component in SVS13-A. Note that the density estimate well agrees with that derived from dust continuum by Chen et al. (2009). High temperatures (218 K) have been similarly obtained by Coutens et al. (2014) applying an LTE analysis to four HDO lines detected, using the PdB array, towards the Class 0 NGC1333 IRAS2A. Also, high excitation conditions have been found through LVG analysis of H₂O lines as observed by Herschel towards both Class 0 and Class I sources (e.g. Herczeg et al. 2012; Podio et al. 2012; Karska et al. 2013; Busquet et al. 2014; Kristensen et al. 2016); in these cases, the high excitation H₂O emission has been associated with shocked gas induced by jets. Thus, the present results may indicate the presence of jet-induced shocks on the ~ 20 AU scales. However, the jet hypothesis is ruled out by: (i) the FWHM line profiles ($\simeq 4~{\rm km~s}^{-1}$) (the is usually traced by velocities larger than 10 km s⁻¹), and (ii) the compact size inferred by the LVG analysis. On the other hand, HDO emission could be emitted by shocks induced by the viscosity of a disk or it could probe the base of a disk wind (as suggested by Codella et al. 2016 for the Class 0 HH212). Again, there is no signature in the line profile suggesting an association of water with a small portion of the large disks expected to be around Class I objects (around 250 AU; e.g. Eisner et al. 2012).

In conclusion, what is HDO tracing in the SVS13-A

system? The observed high temperature and linewidth are consistent with the presence of a hot corino inside SVS13-A, where the gas is thermally heated by the central source. Of course, the definition of hot corino involves the detection of complex organic molecules (e.g. Ceccarelli et al. 2007). We anticipate that this is indeed the case for SVS13-A (Bianchi et al., in preparation). Assuming that the dust is heated by the 34 L_{\odot} central source and that the dust emission is optically thin, the dust temperature would be about 200 K at a distance of ~ 25 AU (see e.g. Ceccarelli et al. 2000, Eq. 1). Of course, if the dust opacity is thick in the innermost regions, then this value is a lower limit. Therefore, this temperature is in good agreement with the LVG analysis. Indeed, high temperatures from HDO were observed by Coutens et al. (2014) towards the Class 0 object NGC1333-IRAS2A, in agreement with the present hypothesis that HDO lines, being optically thin, probe inner regions around the protostars.

However, the hot corino interpretation has a problem. If the high temperature is caused by the thermal heating, one would expect that water is sublimated from the icy grain mantles at ~ 100 K, whereas the HDO line emission indicates a larger temperature. Why? A first possibility could be that the HDO line emission is dominated by warmer gas because of the line opacities. Indeed, even if the HDO abundance has a jump at ~ 100 K, if the lines are optically thin then the warmer regions, with higher opacities may dominate the integrated intensity. One has also to consider that the water abundance has a further jump at around 220-250 K, caused by the reactions that convert all the gaseous atomic oxygen into water and that possess activation barriers making them efficient at ≥ 220 K (Ceccarelli et al. 1996). The temperature is close to that derived from the LVG modeling, so that it is a plausible hypothesis that the gas probed by the observed HDO lines lies in a region warmer than the water desorption region because of the line opacities.

3.4 Water deuteration

Using the intensity 3σ upper limit of the para- $\mathrm{H}_2^{18}\mathrm{O}$ line at 203.40752 GHz (see §3.1) and assuming a source size of 0'.2 and a temperature of 200 K, we derive un upper limit to the $\mathrm{H}_2^{18}\mathrm{O}$ column density of $N(H_2^{18}O) \leq 8 \times 10^{17}$ cm⁻². Assuming the standard value of $^{16}\mathrm{O}/^{18}\mathrm{O} = 560$, the upper limit to the water column density is $N(H_2O) \simeq 4 \times 10^{20}$ cm⁻². Using the HDO column density previously derived, $N(HDO) = 4 \times 10^{17}$ cm⁻², leads to a lower limit to the water deuteration, $\geq 1 \times 10^{-3}$. Finally, using the derived

^a From the Jet Propulsion Laboratory database (Pickett et al. 1998). ^b The errors are the gaussian fit uncertainties.

4 C. Codella et al.




Figure 1. HDO line profiles (in main-beam temperature, $T_{\rm MB}$, scale) observed with the IRAM 30-m antenna towards SVS13-A. In each panel, both the transition and the upper level energy, $E_{\rm u}$, are reported.

 n_{H_2} density (2 × 10⁸ cm⁻³) and emitting sizes (0"2 = 50 AU in diameter) provides an estimate of the H₂ column density of ~ 1.5 × 10²³ cm⁻² and, consequently of the HDO abundance of ~ 3 × 10⁻⁶. Similarly, the upper limit to the H₂O column density can be converted into an upper limit to the water abundance, namely $\leq 3 \times 10^{-3}$. We can, therefore, increase the real lower limit to the water deuteration considering that, reasonably, the water abundance cannot be larger than about ~ 2 × 10⁻⁴. This leads to a lower limit HDO/H₂O \geq 0.015.

This upper limit is consistent with those derived so far towards Class 0 protostars, $\sim 10^{-2}$ by Coutens et al. (2012)




Figure 2. Upper panel: The $\chi^2_{\rm r}$ contour plot obtained considering the non-LTE model predicted and observed intensities of all detected HDO lines but the 241.974 GHz line with $E_{\rm u}=837~{\rm K}$ (for which no collisional rates are available). The best fit is obtained with 0".2, $N_{\rm HDO}=4\times10^{17}~{\rm cm}^{-2},~T_{\rm kin}=200~{\rm K},$ and $n_{\rm H_2}=2\times10^8~{\rm cm}^{-3})$. The 1σ and 2σ of the $\chi^2_{\rm r}$ contours are reported. Lower panel: Ratio between the observed line intensities with those predicted by the best fit model as a function of line upper level energy $E_{\rm u}$.

and Taquet et al. (2013), $0.3-8 \times 10^{-2}$, and larger than those quoted by Person et al. (2013, 2014) and Coutens (2013; 2014), $0.1-4 \times 10^{-3}$. Therefore, the deuteration of water does not seem to diminish from Class 0 to Class I sources. Yet, we conclude with a word of prudence, as this value of deuteration has been obtained taking the values of the LVG modeling. In particular, since the lines seem close to the LTE, the H₂ density could be larger and, consequently, the HDO abundance could be lower by the same factor.

4 CONCLUSIONS

The high-sensitivity of the IRAM 30-m ASAI unbiased spectral survey in the mm-window allows us to detect towards the Class I object SVS13-A a large number of HDO emission lines with upper level energies up to $E_{\rm u}=837$ K. The non-LTE LVG analysis points to hot (150–260 K), dense ($\geqslant 3 \times 10^7$ cm⁻³) gas associated with a quite small emitting region (50 AU), supporting the occurrence of a hot corino inside SVS13-A. The HDO abundance is found to be $\sim 3 \times 10^{-6}$. Although the occurrence of shocks at such small scales cannot be excluded, it is tempting to suggest we are




Figure 3. Rotation diagram for the HDO transitions reported in Table 1, including the $E_{\rm up}=837$ K, not considered in the LVG analysis (see Fig. 2). The parameters $N_{\rm up}$, $g_{\rm up}$, and $E_{\rm up}$ are respectively the column density, the degeneracy, and the energy of the upper level. The error bars on $\ln(N_{\rm up}/g_{\rm u})$ are given by the vertical bar of the symbols. The plot allows us to derive a rotational temperature of 334 ± 42 K and a total column density of $3.7\pm0.5\times10^{17}$ cm⁻².

observing for the first time the jump in the water abundance occurring at temperatures higher than 200 K, when the activation barriers of the gas phase reactions coverting oxygen into water can be overcome.

Obviously, the final answer is in the hands of future interferometric observations (e.g. using ALMA) imaging water emission around SVS13-A on scales \leq 20 AU.

ACKNOWLEDGMENTS

The authors are grateful to the IRAM staff for its help in the calibration of the PdBI data. We also thank F. Dulieu for instructive discussions. The research leading to these results has received funding from the European Commission Seventh Framework Programme (FP/2007-2013) under grant agreement N 283393 (RadioNet3). This work was partly supported by the PRIN INAF 2012 – JEDI and by the Italian Ministero dell'Istruzione, Università e Ricerca through the grant Progetti Premiali 2012 – iALMA. CCe & BL acknowledge the financial support from the French Space Agency CNES, and RB from Spanish MINECO (through project FIS2012-32096).

REFERENCES

Altwegg K., Balsiger H., Bar-Nun A., et al. 2015, Science 347, 387

Busquet G., Lefloch B., Benedettini M., et al. 2014, A&A 561, 120

Caselli P., Keto E., Bergin E.A., et al. 2012, ApJ 759, L37 Ceccarelli C., Hollenbach D.J., & Tielens A.G.G.M. 1996, ApJ 471, 400

Ceccarelli C., Caux E., Loinard L., et al. 1999, A&A 324, L21

Ceccarelli C., Castets A., Caux E., et al. 2000, A&A 355, 1129

Ceccarelli C., Maret S., Tielens A.G.G.M., Castets A., & Caux E. 2003, A&A 410, 587

Ceccarelli C., Caselli P., Bockelèe-Morvan D., et al. 2014, Protostars and Planets VI, 859

Chen X., Launhardt R., & Henning Th. 2009, ApJ 691, 1729

Codella C., Ceccarelli C., Nisini B., et al. 2010, A&A 522, L1

Coutens A., Vastel C., Caux E., et al. 2012, A&A 539, 132Coutens A., Vastel C., Cabrit S., et al. 2013, A&A 560, A39

Coutens A., Jørgensen J.K., Persson M.V., van Dishoeck E.F., Vastel C., & Taquet V. 2014, ApJ 792, L5

Dubernet M.-L., Alexander M.H., Ba Y.A., et al. 2013, A&A 553, 50

Eisner, J.A. 2012, ApJ 755, 23

Faure A., Wiesenfeld L., Scribano Y., & Ceccarelli C. 2012, MNRAS 420, 699

Herczeg G.J., Karska A., Bruderer S., et al. 2012, A&A 540, 84

Hirota T., Bushimata T., Choi Y.K., et al. 2008, PASJ 60, 37

Karska A., Bruderer S., van Dishoeck E.F., et al. 2013, A&A 552, 141

Lefloch B., Castets A., Cernicharo J., Langer W.D., & Zylka R. 1998, A&A 334, 269

Lefloch B., Cabrit S., Busquet G., et al. 2012, A&A 757, L25

López-Sepulcre A., Jaber A.A., Mendoza E., et al. 2015, MNRAS 449, 2438

Persson M.V., Jørgensen J.K., & van Dishoeck E.F. 2013, A&A 549, L3

Persson M.V., Jørgensen J.K., van Dishoeck E.F. & Harsono D. 2014, A&A 563, 74

Pickett H.M., Poynter R.L., Cohen E.A., et al. 1998, J. Quant. Spectrosc. & Rad. Transfer 60, 883

Taquet V., López-Sepulcre A., Ceccarelli C., et al. 2013, ApJ 768, L29

Taquet V., Charnley S., & Sipilä O. 2014, ApJ 791, 1

Tobin J.J., Looney L.W., Li Z.-Y., et al. 2016, ApJ 818, 73 van Dishoeck E.F., Kristensen L.E., Benz, A.O., et al. 2011, PASP 123, 138

van Dishoeck E.F., Bergin E.A., Lis D.C., & Lunine J.I. 2014, Protostars and Planets VI, 835

Vastel C., Ceccarelli C., Caux E., et al. 2010, A&A 521, L31

Willacy K., Alexander C., Ali-Dib N., et al. 2015, Space Science Reviews 197, 151

This paper has been typeset from a $\text{TEX}/\text{ E}^{\text{TEX}}$ file prepared by the author.