# Self-dual codes and the non-existence of a quasi-symmetric 2-(37, 9, 8) design with intersection numbers 1 and 3

Masaaki Harada, Akihiro Munemasa, and Vladimir D. Tonchev<sup>‡</sup>

September 10, 2018

#### Abstract

We prove that a certain binary linear code associated with the incidence matrix of a quasi-symmetric 2-(37, 9, 8) design with intersection numbers 1 and 3 must be contained in an extremal doubly even self-dual code of length 40. Using the classification of extremal doubly even self-dual codes of length 40, we show that a quasi-symmetric 2-(37, 9, 8) design with intersection numbers 1 and 3 does not exist.

### **1** Introduction

A t- $(v, k, \lambda)$  design  $\mathcal{D}$  is a pair of a set  $\mathcal{P}$  of v points and a collection  $\mathcal{B}$  of kelement subsets of  $\mathcal{P}$  (called blocks) such that every t-element subset of  $\mathcal{P}$  is contained in exactly  $\lambda$  blocks. The number of blocks that contain a given point is traditionally denoted by r, and the total number of blocks is b. We assume that t < k < v, throughout this paper. Often a t- $(v, k, \lambda)$  design is simply called a t-design. A t-design can be represented by its (block by point) incidence matrix

<sup>\*</sup>Research Center for Pure and Applied Mathematics, Graduate School of Information Sciences, Tohoku University, Sendai 980–8579, Japan

<sup>&</sup>lt;sup>†</sup>Research Center for Pure and Applied Mathematics, Graduate School of Information Sciences, Tohoku University, Sendai 980–8579, Japan.

<sup>&</sup>lt;sup>‡</sup>Department of Mathematical Sciences, Michigan Technological University Houghton, MI 49931, USA.

 $A = (a_{ij})$ , where  $a_{ij} = 1$  if the *j*th point is contained in the *i*th block and  $a_{ij} = 0$  otherwise.

A 2-design is called *quasi-symmetric* if the number of points in the intersection of any two distinct blocks takes only two values x and y (x < y). The values xand y are called the *intersection numbers*. Quasi-symmetric 2-designs have been widely studied because of their connections with strongly regular graphs and binary self-orthogonal codes (see [22], [25], [26]). Four classes of quasi-symmetric 2-designs were given in [17], namely, multiples of symmetric 2-designs, Steiner systems with  $v > k^2$ , strongly resolvable 2-designs, and residual designs of biplanes. A quasi-symmetric 2-design  $\mathcal{D}$  is called *exceptional* if neither  $\mathcal{D}$  nor its complementary design belongs to one of these four classes. A fundamental problem is to determine whether there is an exceptional quasi-symmetric 2-design for a given set of parameters. The parameters of exceptional quasi-symmetric 2-( $v, k, \lambda$ ) designs with  $2k \le v \le 70$  are listed in [21, Table 48.25].

According to [21], the existence of quasi-symmetric 2-(37, 9, 8) design with intersection numbers 1 and 3 is an open question. Bouyuklieva and Varbanov [3] proved that there is no quasi-symmetric 2-(37, 9, 8) design with intersection numbers 1 and 3 having an automorphism of order 5.

The existence of a quasi-symmetric 2-(37, 9, 8) design with intersection numbers 1 and 3 has been of interest for several reasons. The block graph of such a design, where two blocks are adjacent if they share three points, is a strongly regular graph with parameters (148, 84, 50, 44), and the existence of a graph with these parameters is still unknown (see [4], [18, Table 3]).

In [10] Jungnickel and Tonchev formulated as a conjecture a sufficient condition under which a quasi-symmetric 2- $(v, k, s\lambda)$  design with

$$k(k-1) = \lambda(v-1)$$

is necessarily the union of s identical copies of a symmetric  $2 \cdot (v, k, \lambda)$  design. This conjecture, which was later proved by Ionin and Shrikhande [9], and Sane [20], left the parameters  $2 \cdot (37, 9, 8)$  (s = 4), as the smallest open case for which a quasi-symmetric design which is not a multiple of a symmetric design, might exist. We note that the union of four identical copies of a symmetric  $2 \cdot (37, 9, 2)$ design (up to isomorphism, there exist exactly four such designs [15, page 37]), is a quasi-symmetric  $2 \cdot (37, 9, 8)$  design with intersection numbers 2 and 9.

The parameters 2-(37, 9, 8) coincide with those of a derived design with respect to a block of a (yet unknown [15, page 54]) symmetric 2-(149, 37, 9) design. If a quasi-symmetric 2-(37, 9, 8) design with intersection numbers 1 and 3

can be embedded as a derived design in a symmetric 2-(149, 37, 9) design, then the corresponding residual design will be a quasi-symmetric 2-(112, 28, 9) design with intersection numbers 6 and 8. We note that according to [15, page 54] a 2-(112, 28, 9) design is also unknown.

Self-dual codes, and more generally, self-orthogonal codes, have been used successfully for the construction and classification of quasi-symmetric designs, as well as for deriving necessary conditions for their existence [5], [7], [8], [11], [12], [16], [19], [23], [24].

The main result of this paper is the following theorem, which was established by using the recent classification of binary doubly even self-dual codes of length 40 [1].

**Theorem 1.** A quasi-symmetric 2-(37, 9, 8) design with intersection numbers 1 and 3 does not exist.

As a corollary of Theorem 1, we have the following.

**Corollary 2.** A quasi-symmetric 2-(112, 28, 9) design with intersection numbers 6 and 8 is not embeddable as a residual design in a symmetric 2-(149, 37, 9) design.

### 2 Quasi-symmetric designs and doubly even self-dual codes

A (binary) [n, k] code C is a k-dimensional vector subspace of  $\mathbb{F}_2^n$ , where  $\mathbb{F}_2$ denotes the finite field of order 2. All codes in this paper are binary. The parameter n is called the *length* of C. The *weight* wt(x) of a vector  $x \in \mathbb{F}_2^n$  is the number of nonzero components of x. The support supp(x) of a vector  $x = (x_1, \ldots, x_n) \in \mathbb{F}_2^n$  is  $\{i \mid x_i = 1\}$ . A vector of C is called a *codeword*. The minimum nonzero weight of all codewords in C is called the *minimum weight* of C, and an [n, k]code with minimum weight d is called an [n, k, d] code. A code is called *doubly even* if all codewords have weight divisible by 4.

The dual code  $C^{\perp}$  of a code C of length n is defined as  $C^{\perp} = \{x \in \mathbb{F}_2^n \mid x \cdot y = 0 \text{ for all } y \in C\}$ , where  $x \cdot y$  is the standard inner product. A code C is called *self-orthogonal* if  $C \subset C^{\perp}$ , and C is called *self-dual* if  $C = C^{\perp}$ . A doubly even code is self-orthogonal. A doubly even self-dual code of length n exists if and only if  $n \equiv 0 \pmod{8}$ . The minimum weight d(C) of a doubly even self-dual code C of length n is bounded by  $d(C) \leq 4\lfloor n/24 \rfloor + 4 \lfloor 14 \rfloor$ . A doubly even self-dual code meeting the upper bound is called *extremal*. Two codes are

*equivalent* if one can be obtained from the other by permuting the coordinates. A classification of doubly even self-dual codes was done for lengths up to 40 (see [1] and the references given therein, and the data can be found in [6]). For example, there are 16470 inequivalent extremal doubly even self-dual codes of length 40.

Let  $\mathcal{D}$  be a 2- $(v, k, \lambda)$  design. Let A be the incidence matrix of  $\mathcal{D}$ . Let  $C_1, C_2$  and  $C_3$  be the codes generated by the rows of the following matrices:

$$A, \begin{bmatrix} 1\\ A & \vdots\\ 1 \end{bmatrix}$$
$$\begin{bmatrix} 1 & 1 & 1\\ A & \vdots & \vdots\\ 1 & 1 & 1 \end{bmatrix},$$
(1)

and

respectively.

**Lemma 3** ([24]). The codes  $C_1^{\perp}$  and  $C_2^{\perp}$  have minimum weights at least  $(r+\lambda)/\lambda$  and at least (b+r)/r, respectively.

*Remark* 4. In [24, Lemma 2.3], it was claimed that  $C_2^{\perp}$  has minimum weight at least  $\min\{(r+\lambda)/\lambda, (b+r)/r\}$ . We note that  $(r+\lambda)/\lambda = (v-1)/(k-1)+1$  and (b+r)/r = v/k+1.

**Lemma 5.** Let x be a nonzero vector of  $C_3^{\perp}$ . Suppose that

$$x \notin \{(0, \dots, 0, 1, 1, 0), (0, \dots, 0, 1, 0, 1), (0, \dots, 0, 0, 1, 1)\}.$$
 (2)

Then  $\operatorname{wt}(x) \ge (b+r)/r$ .

*Proof.* Let  $x = (x_1, x_2)$  be a nonzero vector of  $C_3^{\perp}$ , where  $x_1 \in \mathbb{F}_2^v$  and  $x_2 \in \mathbb{F}_2^3$ . Suppose that  $wt(x_2)$  is even. Then  $x_1 \in C_1^{\perp}$ . From (2),  $x_1$  is a nonzero vector.

Hence,  $\operatorname{wt}(x) \ge \operatorname{wt}(x_1) \ge (r+\lambda)/\lambda = (v-1)/(k-1) + 1$  by Lemma 3.

Suppose that  $\operatorname{wt}(x_2)$  is odd. Then  $(x_1, 1) \in C_2^{\perp}$ . Hence,  $\operatorname{wt}(x) \ge \operatorname{wt}((x_1, 1)) \ge (b+r)/r$  by Lemma 3. The result follows.

Let C be a doubly even code of length  $n \equiv 0 \pmod{8}$ . By Theorem 2.2 in [13], there is a doubly even self-dual code of length n containing C.

**Lemma 6.** Let  $\mathcal{D}$  be a quasi-symmetric 2- $(v, k, \lambda)$  design with intersection numbers x and y. Suppose that  $v \equiv 5 \pmod{8}$ ,  $k \equiv 1 \pmod{4}$  and x, y are odd. Let A be the incidence matrix of  $\mathcal{D}$ . Let C be a doubly even self-dual code of length v + 3 containing the row span of the matrix (1). Then C has minimum weight at least (b + r)/r.

*Proof.* From the assumption on the parameters of  $\mathcal{D}$ , the code generated by the rows of the matrix (1) is doubly even. By Lemma 5, the doubly even self-dual code C has minimum weight at least (b+r)/r.

## **3** Quasi-symmetric 2-(37, 9, 8) designs and doubly even self-dual codes of length 40

**Theorem 7.** Let A be the incidence matrix of a quasi-symmetric 2-(37, 9, 8) design  $\mathcal{D}$  with intersection numbers 1 and 3. Let C be a doubly even self-dual [40, 20] code containing the row span of the matrix (1). Then C has minimum weight 8, and there is no codeword of weight 8 whose support contains the last three coordinates.

*Proof.* By Lemma 6, C has minimum weight 8, since the minimum weight of a doubly even self-dual code of length 40 is at most 8.

Let  $\mathcal{B}$  be the block set of  $\mathcal{D}$ . Suppose that x is a codeword of weight 8 whose support contains the last three coordinates. Let  $S = \sup(x) \setminus \{38, 39, 40\}$  and

$$n_i = |\{B \in \mathcal{B} \mid |S \cap B| = i\}| \quad (0 \le i \le 5).$$

Note that  $n_i = 0$  (i = 0, 2, 4). Moreover, if  $n_5 > 0$ , then there exists a block containing S. This means that C contains a codeword of weight 4, which contradicts the already proven fact that C has minimum weight 8. Thus  $n_5 = 0$ . Counting the number of pairs (T, B) with  $T \subset S \cap B$ , |T| = 2 and  $B \in \mathcal{B}$ , we find  $3n_3 = 80$ . This is impossible since  $n_3$  is an integer.

### 4 The search method

All computer calculations in this section were done with the help of MAGMA [2]. Starting from an extremal doubly even self-dual [40, 20, 8] code C, we take  $T \subset$ 

 $\{1, 2, ..., 40\}$  with |T| = 3 such that no codeword of weight 8 contains T in its support, and consider the following set:

$$X = \{ \operatorname{supp}(x) \setminus T \mid x \in C, \ \operatorname{wt}(x) = 12, \ \operatorname{supp}(x) \supset T \}.$$

If there is a quasi-symmetric 2-(37, 9, 8) design with intersection numbers 1 and 3, an isomorphic copy can be found in X for some choice of C and T. In order to determine whether such X contains a design, we first tested whether the graph  $\Gamma_{ij} = (V, E)$  with

$$V = \{B \in X \mid B \supset \{i, j\}\},\$$
  
$$E = \{\{B, B'\} \mid B, B' \in V, |B \cap B'| = 3\},\$$

has a clique of size  $\lambda = 8$  for distinct  $i, j \in \{1, 2, \dots, 40\} \setminus T$ .

There are 16470 inequivalent extremal doubly even self-dual codes of length 40 [1]. We verified that 15940 of them contain no triple T such that  $\Gamma_{ij}$  has a clique of size 8 for all  $i, j \in \{1, 2, ..., 40\} \setminus T$  with  $i \neq j$ .

For the remaining 530 codes, we used the following method. Fix  $i_0, j_0 \in \{1, 2, ..., 40\} \setminus T$  and enumerated all cliques of size 8 in the graph  $\Gamma_{i_0, j_0}$ . For such a clique K and distinct  $i, j \in \{1, ..., 40\} \setminus T$ , consider the subgraph  $\Gamma_{ij}^K$  of  $\Gamma_{ij}$  induced by the set of vertices

$$\{B \in X \mid B \supset \{i, j\}, |B \cap B'| \in \{1, 3\} \text{ for all } B' \in K\}.$$

If there exists a quasi-symmetric 2-(37, 9, 8) design with intersection numbers 1 and 3 whose block set is contained in X, then the block set contains a clique K of size 8 in  $\Gamma_{i_0,j_0}$ , and

$$\Gamma_{ii}^{K}$$
 has a clique of size 8 for all distinct  $i, j \in \{1, \dots, 40\} \setminus T$ . (3)

We chose the initial  $i_0$ ,  $j_0$  in such a way that  $\Gamma_{i_0,j_0}$  has minimal number of cliques of size 8, for computational efficiency. It turns out that, for each of the remaining 530 codes and each choice of a triple T, our choice of  $i_0$ ,  $j_0$  shows that no clique K of size 8 in  $\Gamma_{i_0,j_0}$  satisfies the condition (3). Therefore, we have established Theorem 1. We note that the CPU time for the entire search was approximately 300 hours.

Acknowledgment. This work was supported by JSPS KAKENHI Grant Number 15H03633. Vladimir D. Tonchev acknowledges support by NSA Grant H98230-16-1-0011.

#### References

- K. Betsumiya, M. Harada and A. Munemasa, A complete classification of doubly even self-dual codes of length 40, *Electron. J. Combin.* 19 (2012), Paper 18, 12 pp.
- [2] W. Bosma, J. Cannon and C. Playoust, The Magma algebra system I: The user language, *J. Symbolic Comput.* **24** (1997), 235–265.
- [3] S. Bouyuklieva and Z. Varbanov, Quasi-symmetric 2-(37, 9, 8) designs and self-orthogonal codes with automorphisms of order 5, *Math. Balkanica* (N.S.) 19 (2005), 33–38.
- [4] A.E. Brouwer, Parameters of Strongly Regular Graphs, Available online at http://www.win.tue.nl/~aeb/graphs/srg/srgtab.html.
- [5] A.R. Calderbank, The application of invariant theory to the existence of quasi-symmetric designs, *J. Combin. Theory Ser. A* **44** (1987), 94–109.
- [6] M. Harada and A. Munemasa, Database of Self-Dual Codes, Available online at http://www.math.is.tohoku.ac.jp/~munemasa/selfdualcodes.htm.
- [7] M. Harada, A. Munemasa and V.D. Tonchev, A characterization of designs related to an extremal doubly-even self-dual code of length 48, *Ann. Comb.* 9 (2005), 189–198.
- [8] W.C. Huffman and V.D. Tonchev, The existence of extremal [50, 25, 10] codes and quasi-symmetric 2-(49, 9, 6) designs, Des. Codes Cryptogr. 6 (1995), 97–106.
- [9] Y.J. Ionin and M.S. Shrikhande, On a conjecture of Jungnickel and Tonchev for quasi-symmetric designs, *J. Combin. Des.* **2** (1994), 49–59.
- [10] D. Jungnickel and V.D. Tonchev, "Intersection numbers of quasi-multiples of symmetric designs," in Advances in Finite Geometries and Designs, J.W.P. Hirschfeld, D.R. Hughes and J.A. Thas (Eds.), Oxford University Press, New York, 1991, pp. 227–236.
- [11] D. Jungnickel and V.D. Tonchev, On symmetric and quasi-symmetric designs with the symmetric difference property and their codes, J. Combin. Theory Ser. A 59 (1992), 40–50.

- [12] D. Jungnickel and V.D. Tonchev, Maximal arcs and quasi-symmetric designs, Des. Codes Cryptogr. 77 (2015), 365–374.
- [13] F.J. MacWilliams, N.J.A. Sloane and J.G. Thompson, Good self dual codes exist, *Discrete Math.* 3 (1972), 153–162.
- [14] C.L. Mallows and N.J.A. Sloane, An upper bound for self-dual codes, Inform. Control 22 (1973), 188–200.
- [15] R. Mathon and A. Rosa, "2- $(v, k, \lambda)$  Designs of small order," in Handbook of Combinatorial Designs (second edition), C.J. Colbourn and J.H. Dinitz (Eds.), Chapman & Hall/CRC, FL, 2007, pp. 578–582.
- [16] A. Munemasa and V.D. Tonchev, A new quasi-symmetric 2-(56, 16, 6) design obtained from codes, *Discrete Math.* 284 (2004), 231–234.
- [17] A. Neumaier, "Regular sets and quasi-symmetric 2-designs," in *Combina-torial Theory*, D. Jungnickel and K. Vedder (Eds.), Lecture Notes in Math. Vol. 969, 1982, pp. 258–275.
- [18] R.M. Pawale, M.S. Shrikhande and S.M. Nyayate, Conditions for the parameters of the block graph of quasi-symmetric designs, *Electron. J. Combin.* 22 (2015), Paper 1.36, 30 pp.
- [19] B.G. Rodrigues and V.D. Tonchev, "On quasi-symmetric 2-(64, 24, 46) designs derived from codes," in *Coding Theory and Applications*, R. Pinto et al. (Eds.), CIM Series in Mathematical Sciences 3, Springer International Publishing, Switzerland, 2015, pp. 327–333.
- [20] S. Sane, A proof of the Jungnickel-Tonchev conjecture on quasi-multiple quasi-symmetric designs, Des. Codes Cryptogr. 23 (2001), 291–296.
- [21] M.S. Shrikhande, "Quasi-symmetric designs," in Handbook of Combinatorial Designs (second edition), C.J. Colbourn and J.H. Dinitz (Eds.), Chapman & Hall/CRC, FL, 2007, pp. 578–582.
- [22] M.S. Shrikhande and S.S. Sane, *Quasi-Symmetric Designs*, London Math. Soc. Lecture Note Ser. 164, Cambridge University Press, Cambridge, 1991.
- [23] V.D. Tonchev, Quasi-symmetric 2-(31,7,7) designs and a revision of Hamada's conjecture, J. Combin. Theory Ser. A 42 (1986), 104–110.

- [24] V.D. Tonchev, Quasi-symmetric designs and self-dual codes, *Europ. J. Combin.* **7** (1986), 67–73.
- [25] V.D. Tonchev, "Codes and designs," in *Handbook of Coding Theory*, V.S. Pless and W.C. Huffman (Eds.), North-Holland, Amsterdam, 1998, pp. 1229–1267.
- [26] V.D. Tonchev, "Codes," in Handbook of Combinatorial Designs (second edition), C.J. Colbourn, J.H. Dinitz (Eds.), Chapman & Hall/CRC, FL, 2007, pp. 677–702.