



# Measurement of $CP$ violation in $B^0 \rightarrow D^+ D^-$ decays

The LHCb collaboration<sup>†</sup>

## Abstract

The  $CP$  violation observables  $S$  and  $C$  in the decay channel  $B^0 \rightarrow D^+ D^-$  are determined from a sample of proton-proton collisions at center-of-mass energies of 7 and 8 TeV, collected by the LHCb experiment and corresponding to an integrated luminosity of  $3 \text{ fb}^{-1}$ . The observable  $S$  describes  $CP$  violation in the interference between mixing and the decay amplitude, and  $C$  parametrizes direct  $CP$  violation in the decay. The following values are obtained from a flavor-tagged, decay-time-dependent analysis:

$$S = -0.54_{-0.16}^{+0.17} (\text{stat}) \pm 0.05 (\text{syst}),$$

$$C = 0.26_{-0.17}^{+0.18} (\text{stat}) \pm 0.02 (\text{syst}).$$

These values provide evidence for  $CP$  violation at a significance level of 4.0 standard deviations. The phase shift due to higher-order Standard Model corrections is constrained to a small value of

$$\Delta\phi = -0.16_{-0.21}^{+0.19} \text{ rad}.$$

Published in Phys. Rev. Lett. 117, 261801 (2016)

© CERN on behalf of the LHCb collaboration, licence CC-BY-4.0.

---

<sup>†</sup>Authors are listed at the end of this Letter.



Studies of beauty hadron decays into pairs of charm hadrons give access to a multitude of observables that probe the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix [1,2] of the Standard Model (SM). Comparisons of these observables with each other and with similar observables from beauty hadron decays to charmonia allow higher-order SM contributions, like loop diagrams, to be separated from effects caused by physics beyond the SM [3–7]. For example, under the assumption that flavor symmetries hold to a good approximation, higher-order corrections in the measurement of  $\phi_s$  in  $B_s^0 \rightarrow D_s^+ D_s^-$  [8] can be constrained.<sup>1</sup>

In the  $B^0$  meson system,  $CP$  violation in the mixing is negligible, as is the decay width difference  $\Delta\Gamma$  of the mass eigenstates [9]. In contrast, sizable  $CP$  violation from the interference between the direct (unmixed) decay into the  $CP$ -even final state  $D^+ D^-$  and the decay to the same final state after  $B^0$ - $\bar{B}^0$  mixing, or from the interference of different decay processes, leads to a decay-time-dependent decay rate of

$$\frac{d\Gamma(t, d)}{dt} \propto e^{-t/\tau} \left( 1 - dS \sin(\Delta mt) + dC \cos(\Delta mt) \right), \quad (1)$$

where  $t$  is the proper decay time,  $d$  represents the  $B^0$  flavor at production and takes a value of  $+1$  for mesons whose initial flavor is  $B^0$  and  $-1$  for  $\bar{B}^0$ ,  $\tau$  is the mean lifetime and  $\Delta m$  is the mass difference between the physical  $B^0$  meson eigenstates. The  $CP$  observables  $S$  and  $C$  are related to the  $B^0$  mixing phase  $\phi_d$  and a phase shift  $\Delta\phi$  from the decay amplitudes via  $S/\sqrt{1-C^2} = -\sin(\phi_d + \Delta\phi)$  [10]. In the SM,  $\phi_d = 2\beta$ , where  $\beta \equiv \arg[-(V_{cd}V_{cb}^*)/(V_{td}V_{tb}^*)]$  is an angle of one of the CKM unitary triangles and  $V_{qq'}$  are elements of the CKM matrix. If the  $B^0 \rightarrow D^+ D^-$  decay amplitude can be described by a dominant tree-level  $b \rightarrow c\bar{c}d$  transition, the phase shift  $\Delta\phi$  vanishes and the  $CP$  observables are given by  $C = 0$  and  $S = -\sin\phi_d$ . The value of the latter has been measured to be  $\sin\phi_d = +0.679 \pm 0.020$  [9] in  $b \rightarrow c\bar{c}s$  decays such as  $B^0 \rightarrow J/\psi K_s^0$ , in which the contribution from loop processes in the decay can be constrained to high precision [11]. In contrast, previous measurements of the  $CP$  observables in the decay  $B^0 \rightarrow D^+ D^-$  by the BaBar and Belle collaborations [12, 13] give world average values of  $S = -0.98 \pm 0.17$  and  $C = -0.31 \pm 0.14$  [9]. The values are at the edge of the physically allowed region of  $S^2 + C^2 \leq 1$ , which leaves room for a large value of  $\Delta\phi$ .

This Letter reports a measurement of  $CP$  violation in  $B^0 \rightarrow D^+ D^-$  decays with the LHCb experiment. The measurement is based on samples of  $pp$  collision data corresponding to integrated luminosities of 1 and  $2 \text{ fb}^{-1}$  at center-of-mass energies of 7 and 8 TeV, respectively, recorded by the LHCb experiment. The LHCb detector is a single-arm forward spectrometer covering the pseudorapidity range  $2 < \eta < 5$ , designed for the study of particles containing  $b$  or  $c$  quarks, and is described in detail in Refs. [14, 15]. The online event selection is performed by a trigger, which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction. Simulated events are produced with the software described in Refs. [16–21].

Candidate  $B^0 \rightarrow D^+ D^-$  decays are reconstructed through the subsequent decays  $D^+ \rightarrow K^- \pi^+ \pi^+$  and  $D^+ \rightarrow K^- K^+ \pi^+$ , with combinations of two  $D \rightarrow KK\pi$  candidates omitted due to the low branching fraction. The kaon and pion candidates, which must fulfill loose particle identification (PID) criteria, are required to have transverse momentum

---

<sup>1</sup>The inclusion of charge-conjugate processes is implied throughout the Letter, unless otherwise noted.

$p_T > 100 \text{ MeV}/c$ , to have a good track quality and to be inconsistent with originating from a primary vertex (PV). The three hadron tracks must form a good common vertex and their combined invariant mass has to be in the range  $\pm 25 \text{ MeV}/c^2$  around the known  $D^+$  mass [22]. The scalar sum of the  $p_T$  of the three hadrons has to exceed  $1800 \text{ MeV}/c$  and the  $D^+$  vertex has to be significantly displaced from all PVs. Defining  $\theta_X$  as the angle between the momentum vector of a particle  $X$  and the displacement vector from the best-matched PV to the  $X$  decay vertex,  $\cos \theta_{D^+}$  is required to be positive.

To suppress contributions from misreconstructed  $D_s^+ \rightarrow K^- K^+ \pi^+$  decays, which proceed predominantly through  $D_s^+ \rightarrow \phi \pi^+$ ,  $D^+ \rightarrow K^- \pi^+ \pi^+$  candidates are rejected if, after assigning the kaon mass hypothesis to the  $\pi^+$  with the higher  $p_T$ , the invariant mass  $m(K^- K^+)$  is within  $10 \text{ MeV}/c^2$  of the known  $\phi$  meson mass. Furthermore, if the invariant mass  $m(K^- K^+ \pi^+)$  is within  $25 \text{ MeV}/c^2$  of the known  $D_s^+$  meson mass, the requirement on the PID information of the higher- $p_T$  pion to be consistent with the pion hypothesis is tightened. Similarly, protons can be misidentified as pions, resulting in background contributions from  $\Lambda_c^+ \rightarrow K^- p \pi^+$ . To suppress these processes, the pion candidate with the higher  $p_T$  of  $D^+ \rightarrow K^- \pi^+ \pi^+$  is required to be well identified as a pion if  $|m(K^- p \pi^+) - m_{\Lambda_c^+}| < 25 \text{ MeV}/c^2$ .

Candidate  $B^0$  mesons are reconstructed from pairs of oppositely charged  $D^\pm$  candidates that form a common vertex. The scalar sum of the  $p_T$  of the  $D^\pm$  mesons must exceed  $5 \text{ GeV}/c$ . The decay time significance of each  $D^\pm$  meson, defined as its decay time divided by its estimated uncertainty, is required to be greater than 0, or greater than 3 if one of the  $D^\pm$  mesons is reconstructed in the  $K^- K^+ \pi^+$  final state. This reduces the contamination of  $B^0 \rightarrow D^- K^- K^+ \pi^+$  decays. The  $B^0$  candidate is required to have momentum  $p > 10 \text{ GeV}/c$ ,  $\cos \theta_{B^0} > 0.999$ , and to not originate from the associated PV. A fit to the full decay chain, in which the  $B^0$  production vertex is constrained to the position of the associated PV, is performed to determine the reconstructed decay time  $t'$  of the  $B^0$  candidate, which differs from the true time  $t$ . Only candidates with decay times in the range 0.25–10.25 ps are kept. The invariant mass  $m_{D^+ D^-}$  of the  $B^0$  candidate is calculated from a similar fit to the full decay chain, while additionally constraining the invariant masses of  $K^- \pi^+ \pi^+$  and  $K^- K^+ \pi^+$  to the known  $D^+$  mass, and is required to be in the range 5150–5500  $\text{MeV}/c^2$ .

Two boosted decision trees (BDTs) [23, 24], for  $B^0$  final states with two and three kaons, are used to suppress the combinatorial background. Both are trained on simulated signal samples and on background samples formed from  $B^0$  candidates at high invariant masses ( $> 5500 \text{ MeV}/c^2$ ), and exploit observables related to the kinematics of the decay, PID information, and track and vertex quality. The requirements on the BDT classifier outputs are chosen to optimize the precision of both  $CP$  observables,  $S$  and  $C$ .

To separate the remaining background from the signal a fit to the  $D^+ D^-$  invariant mass distribution is performed to calculate signal candidate weights via the *sPlot* technique [25]. The mass fit is performed simultaneously in four categories, split by the data-taking period (7 TeV, 8 TeV) and the number of kaons in the final state. The probability density function (PDF) used to parametrize the mass distribution consists of four contributions: signal,  $B_s^0 \rightarrow D^+ D^-$ , combinatorial background, and a component that includes both  $B^0 \rightarrow D_s^+ D^-$  and  $B_s^0 \rightarrow D_s^- D^+$  decays. The signal is modeled by the sum of three Crystal Ball functions [26] with a common mean. The parameters of the tails (two towards lower and one towards higher mass) and the three widths are determined from simulated samples. To account for differences in the mass resolution in simulation and data, the




Figure 1: Distribution of the reconstructed mass of all  $B^0 \rightarrow D^+ D^-$  candidates (a) and background-subtracted decay time distribution for tagged candidates (b). In plot (a) besides the data points and the projection of the full PDF (solid black) the projections of the  $B^0$  signal (dashed blue), the  $B_s^0 \rightarrow D^+ D^-$  background (short-dash-dotted turquoise), the  $B^0 \rightarrow D_s^+ D^-$  background (dotted green), the  $B_s^0 \rightarrow D_s^- D^+$  background (long-dash-three-dotted red) and the combinatorial background (long-dash-dotted purple) are shown.

width parameters are multiplied by a common scale factor, which is free to vary in the fit to data. The  $B_s^0 \rightarrow D^+ D^-$  component shares all shape parameters with the signal PDF except for the peak position, which is constrained by the known value of the difference between the  $B^0$  and the  $B_s^0$  masses [22]. Each peak in the  $B^0 \rightarrow D_s^+ D^-$  and  $B_s^0 \rightarrow D_s^- D^+$  component is described by the sum of two Crystal Ball functions (one with a tail towards lower and one with a tail towards higher masses) whose parameters are taken from simulation. The widths and the  $B^0$  peak position are free to vary in the fit while the  $B_s^0$  peak offset is constrained in the same way as that of the  $B_s^0 \rightarrow D^+ D^-$  component. The combinatorial background is parametrized with an exponential function, with separate exponents used for the final states with two or three kaons. Partially reconstructed  $B^0 \rightarrow D^{*+} D^-$  decays with  $D^{*+} \rightarrow D^+ \pi^0$ , where the neutral pion is missed, lie outside the mass range used for the fit. The equivalent  $B_s^0 \rightarrow D^{*+} D^-$  decays and decay modes with only one or no charm meson, such as  $B^0 \rightarrow D^- K^- K^+ \pi^+$ , are also neglected in the mass fit. The influence of their omission on the  $CP$  measurement is treated as a systematic uncertainty. The mass distribution is shown in Fig. 1 (a). The combined  $B^0 \rightarrow D^+ D^-$  signal yield is  $1610 \pm 50$ , of which  $1347 \pm 45$  are in the Cabibbo-favored final state with two  $D^+ \rightarrow K^- \pi^+ \pi^+$  decays.

The measurement of decay-time-dependent  $CP$  violation requires knowledge of the initial flavor of each reconstructed  $B^0$  meson. Flavor-tagging algorithms deliver a measured tag decision  $d'$  for the flavor of the  $B^0$  meson, which takes the value  $+1$  for a  $B^0$ ,  $-1$  for a  $\bar{B}^0$  initial state, and  $0$  if no decision is possible, and an estimate  $\eta$  of the probability for the tag decision to be incorrect. The latter is referred to as the mistag probability. Two classes of flavor-tagging algorithms are used: opposite-side (OS) and same-side (SS) taggers [27–29]. In  $b\bar{b}$  pair production, the dominant source of  $b$  hadrons at LHCb, the signal  $B^0$  meson is accompanied by a second  $b$  hadron. The OS taggers determine the flavor of the signal by examining the decay products of this second  $b$  hadron. The information from the decay products consists of the charge of muons or electrons produced in semileptonic decays, the charge of kaons from  $b \rightarrow c \rightarrow s$  transitions, the charge of charm hadrons from  $b \rightarrow c$  transitions, and the net charge of all decay products. The SS

taggers analyze pions and protons related to the hadronization process of the  $B^0$  meson. This is the first analysis to use the LHCb SS proton and OS charm taggers, and the first to use the new SS pion tagger.

The outputs of all OS algorithms are combined into an overall OS tagging decision and mistag estimate, and the same is done for the SS algorithms. The mistag estimates  $\eta \in \{\eta_{\text{OS}}, \eta_{\text{SS}}\}$  are calibrated using linear functions  $\omega(\eta|d)$ , so that  $\eta$  on average matches the true mistag probability  $\omega$ , which depends on the true production flavor  $d$  of the  $B^0$  meson. The calibration studies are performed with a sample of  $B^0 \rightarrow D_s^+ D^-$  decays, for which the final state determines the flavor of the  $B^0$  at decay. Since the calibration and signal channels are kinematically very similar, the calibration can be applied to the signal channel without further corrections. To ensure that the same calibration is valid for both, the same selection is used as for the signal decay with one  $D^+ \rightarrow K^- K^+ \pi^+$ , apart from requiring that the  $K^- K^+ \pi^+$  invariant mass lie within  $25 \text{ MeV}/c^2$  of the known  $D_s^+$  mass [22] and dropping the vetoes against misidentified backgrounds. Background is subtracted from the calibration sample via the *sPlot* technique [25]. The tagging calibration parameters are determined from a fit to the decay time and tag distributions of  $B^0 \rightarrow D_s^+ D^-$  candidates, in which the detection asymmetry, the production asymmetry of the  $B^0$  mesons, and the flavor-specific semileptonic asymmetry  $a_{\text{sl}}^d$  are taken into account. Here, the detection asymmetry describes the difference in reconstruction efficiency between the  $D_s^+ D^-$  and  $D_s^- D^+$  final states, and  $A_{\text{P}} \equiv [\sigma(\bar{B}^0) - \sigma(B^0)]/[\sigma(\bar{B}^0) + \sigma(B^0)]$ , where  $\sigma$  denotes the production cross-section inside the LHCb acceptance. The values of all these parameters are fixed according to the latest LHCb measurements [30, 31], and their uncertainties are treated as sources of systematic uncertainty on the calibration parameters. Further systematic uncertainties are assigned due to the calibration method, the dependence of the efficiency on decay time, the decay time resolution, and the background subtraction. More details on the calibration studies are given in the supplemental material.

In the  $B^0 \rightarrow D^+ D^-$  signal data sample, the correlation between the OS and the SS mistag estimates is found to be negligible. A small correlation of the mistag probability with decay time is seen; this is neglected in the main fit but considered as a source of systematic uncertainty.

The effective tagging efficiency is the product of the probability for reaching a tagging decision,  $\varepsilon_{\text{tag}} = (87.6 \pm 0.8) \%$ , and the square of the effective dilution  $D = 1 - 2\omega = (30.3 \pm 1.1) \%$ . Its value is  $\varepsilon_{\text{tag}} D^2 = (8.1 \pm 0.6) \%$ , the highest effective tagging efficiency to date in tagged  $CP$  violation measurements at LHCb thanks to the improved flavor-tagging algorithms and the kinematic properties of the selected  $B^0 \rightarrow D^+ D^-$  decays.

The  $CP$  violation observables  $S$  and  $C$  are determined from a multidimensional fit to the background-subtracted tag and decay time distributions of the tagged  $B^0 \rightarrow D^+ D^-$  candidates; a projection of the decay time distribution summed over the non-zero tag decisions is shown in Fig. 1 (b). The conditional PDF describing the reconstructed decay time  $t'$  and tag decisions  $\vec{d}' = (d'_{\text{OS}}, d'_{\text{SS}})$ , given a per-event decay time resolution  $\sigma_{t'}$  and per-event mistag probability estimates  $\vec{\eta} = (\eta_{\text{OS}}, \eta_{\text{SS}})$ , is

$$P\left(t', \vec{d}' \mid \sigma_{t'}, \vec{\eta}\right) \propto \epsilon(t') \left( \mathcal{P}(t, \vec{d}' \mid \vec{\eta}) \otimes \mathcal{R}(t' - t \mid \sigma_{t'}) \right), \quad (2)$$

where

$$\mathcal{P}(t, \vec{d}' \mid \vec{\eta}) \propto \sum_d \mathcal{P}(\vec{d}' \mid d, \vec{\eta}) [1 - d A_{\text{P}}] e^{-t/\tau} \{1 - d S \sin(\Delta m t) + d C \cos(\Delta m t)\}, \quad (3)$$




Figure 2: Decay-time-dependent signal yield asymmetry. The solid curve is the projection of the signal PDF given in Eq. (2).

and where  $t$  is the true decay time,  $d$  is the true production flavor,  $A_P$  is the production asymmetry, and  $\mathcal{P}(\vec{d}' | d, \vec{\eta})$  is a two-dimensional binomial PDF describing the distribution of tagging decisions given  $\vec{\eta}$  and  $d$ . Normalization factors are omitted for brevity. In the fit, the mass difference  $\Delta m$  and the lifetime  $\tau$  are constrained to their known values within uncertainties [22]. The production asymmetry  $A_P$  is constrained separately for the 7 and 8 TeV samples to the values obtained from weighting the results from the measurements in Ref. [31] according to the kinematic distribution of the  $B^0$  signal candidates. The decay time resolution model  $\mathcal{R}$  is the sum of three Gaussian functions, two of which have event-dependent widths proportional to  $\sigma_{t'}$ , and one which has a global width that describes the effect of candidates matched to a wrong PV; all three share a common mean. All parameters of the resolution model are determined from simulation. The average decay time resolution in data is 49 fs. The function  $\epsilon(t')$  describes the efficiency for all reconstruction and selection steps as a function of the reconstructed decay time. It is represented by cubic splines [32], with the spline coefficients left unconstrained in the fit.

The statistical uncertainties are estimated using the bootstrap method [33]. Individual bootstrap samples are drawn from the candidates in data that pass the full selection; the analysis procedure described above, consisting of the mass fit, background subtraction, and decay time fit, is then applied to obtain the values of the  $CP$  observables for each such sample. Two-sided 68% confidence intervals, with equal tail probabilities on either side, are obtained from the distributions of fitted parameters in the bootstrapped samples. To account for the uncertainties of the flavor-tagging calibration parameters, which are fixed in the likelihood fit, further pseudoexperiments are generated in which these flavor-tagging calibration parameters are varied within their combined statistical and systematic uncertainties. The results are then used to correct the uncertainties from the bootstrapping procedure. The  $CP$  observables are measured to be  $S = -0.54^{+0.17}_{-0.16}$  and  $C = 0.26^{+0.18}_{-0.17}$  with a correlation coefficient of  $\rho = 0.48$ . The decay-time-dependent signal yield asymmetry  $(N_{\bar{B}^0} - N_{B^0}) / (N_{\bar{B}^0} + N_{B^0})$ , where  $N_{B^0}$  is the number of  $B^0 \rightarrow D^+ D^-$  decays with a  $B^0$  flavor tag, and  $N_{\bar{B}^0}$  the number with a  $\bar{B}^0$  tag, is shown in Fig. 2.

Several sources of systematic uncertainties on the  $CP$  observables are studied with pseudoexperiments. The largest systematic uncertainty arises from neglecting backgrounds in which the final state contains only one charm meson, such as  $B^0 \rightarrow D^- K^- K^+ \pi^+$ . The yield of these backgrounds is estimated to be about 2% of the signal yield and their

impact is assessed by assuming that they maximally violate  $CP$  symmetry and have the eigenvalue opposite to the signal mode. This leads to a systematic uncertainty of  $\pm 0.05$  on  $S$  and  $\pm 0.013$  on  $C$ . Further systematic uncertainties on  $S$  are related to the assumption  $\Delta\Gamma = 0$  ( $\pm 0.014$ ), and to the modeling of the dependence of the efficiency on decay time ( $\pm 0.007$ ). For  $C$  the second largest systematic uncertainty of  $\pm 0.007$  is due to neglecting the correlation between the invariant mass and the decay time. Additional systematic uncertainties arise from the decay time resolution, the uncertainty on the knowledge of the length scale, the parametrization of the mass model, and from uncertainties on the  $B^0$  production asymmetry and mass difference  $\Delta m$ . The total systematic uncertainty, calculated as the sum in quadrature of all contributions, is  $\pm 0.05$  for  $S$  and  $\pm 0.02$  for  $C$ , with a correlation coefficient of  $\rho = -0.69$ .

In conclusion, a measurement of the  $CP$  observables  $S$  and  $C$  in the decay channel  $B^0 \rightarrow D^+ D^-$  is performed. Using the full data sample collected by the LHCb experiment during Run 1, which corresponds to a total integrated luminosity of  $3 \text{ fb}^{-1}$ , they are determined to be

$$\begin{aligned} S &= -0.54^{+0.17}_{-0.16} (\text{stat}) \pm 0.05 (\text{syst}), \\ C &= 0.26^{+0.18}_{-0.17} (\text{stat}) \pm 0.02 (\text{syst}), \end{aligned}$$

with a statistical correlation coefficient of  $\rho = 0.48$ . This result excludes the conservation of  $CP$  symmetry by 4.0 standard deviations. It is compatible with the previous measurement by the BaBar experiment of  $S = -0.63 \pm 0.36 \pm 0.05$  and  $C = -0.07 \pm 0.23 \pm 0.03$  [12] while being significantly more precise. A proper evaluation of the compatibility with the result from the Belle experiment [13] could not be performed due to its non-Gaussian uncertainties. The result presented here corresponds to  $\sin(\phi_d + \Delta\phi) = 0.56^{+0.16}_{-0.17}$ , which constrains the phase shift to the world's most precise value of  $\Delta\phi = -0.16^{+0.19}_{-0.21}$  rad, and thus implies only a small contribution from higher-order Standard Model corrections.

## Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); FOM and NWO (The Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FASO (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany), EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), Conseil Général de Haute-Savoie, Labex ENIGMASS and OCEVU, Région Auvergne (France), RFBR and Yandex LLC (Russia), GVA, XuntaGal and GENCAT (Spain), Herchel Smith Fund, The



Royal Society, Royal Commission for the Exhibition of 1851 and the Leverhulme Trust (United Kingdom).

## References

- [1] N. Cabibbo, *Unitary symmetry and leptonic decays*, Phys. Rev. Lett. **10** (1963) 531.
- [2] M. Kobayashi and T. Maskawa, *CP violation in the renormalizable theory of weak interaction*, Prog. Theor. Phys. **49** (1973) 652.
- [3] R. Fleischer, *Extracting  $\gamma$  from  $B_{s(d)} \rightarrow J/\psi K_S$  and  $B_{d(s)} \rightarrow D_{d(s)}^+ D_{d(s)}^-$* , Eur. Phys. J. **C10** (1999) 299, [arXiv:hep-ph/9903455](#).
- [4] M. Gronau, J. L. Rosner, and D. Pirjol, *Small amplitude effects in  $B^0 \rightarrow D^+ D^-$  and related decays*, Phys. Rev. **D78** (2008) 033011, [arXiv:0805.4601](#).
- [5] R. Fleischer, *Exploring CP violation and penguin effects through  $B_d^0 \rightarrow D^+ D^-$  and  $B_s^0 \rightarrow D_s^+ D_s^-$* , Eur. Phys. J. **C51** (2007) 849, [arXiv:0705.4421](#).
- [6] M. Jung and S. Schacht, *Standard model predictions and new physics sensitivity in  $B \rightarrow DD$  decays*, Phys. Rev. **D91** (2015) 034027, [arXiv:1410.8396](#).
- [7] L. Bel *et al.*, *Anatomy of  $B \rightarrow D\bar{D}$  decays*, JHEP **07** (2015) 108, [arXiv:1505.01361](#).
- [8] LHCb collaboration, R. Aaij *et al.*, *Measurement of the CP-violating phase  $\phi_s$  in  $\bar{B}_s^0 \rightarrow D_s^+ D_s^-$  decays*, Phys. Rev. Lett. **113** (2014) 211801, [arXiv:1409.4619](#).
- [9] Heavy Flavor Averaging Group, Y. Amhis *et al.*, *Averages of b-hadron, c-hadron, and  $\tau$ -lepton properties as of summer 2014*, [arXiv:1412.7515](#), updated results and plots available at <http://www.slac.stanford.edu/xorg/hfag/>.
- [10] K. De Bruyn and R. Fleischer, *A Roadmap to Control Penguin Effects in  $B_d^0 \rightarrow J/\psi K_S^0$  and  $B_s^0 \rightarrow J/\psi \phi$* , JHEP **03** (2015) 145, [arXiv:1412.6834](#).
- [11] P. Frings, U. Nierste, and M. Wiebusch, *Penguin contributions to CP phases in  $B_{d,s}$  decays to charmonium*, Phys. Rev. Lett. **115** (2015) 061802, [arXiv:1503.00859](#).
- [12] BaBar collaboration, B. Aubert *et al.*, *Measurements of time-dependent CP asymmetries in  $B^0 \rightarrow D^{(*)+} D^{(*)-}$  decays*, Phys. Rev. **D79** (2009) 032002, [arXiv:0808.1866](#).
- [13] Belle collaboration, M. Rohrken *et al.*, *Measurements of branching fractions and time-dependent CP violating asymmetries in  $B^0 \rightarrow D^{(*)\pm} D^\mp$  decays*, Phys. Rev. **D85** (2012) 091106, [arXiv:1203.6647](#).
- [14] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [15] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, Int. J. Mod. Phys. **A30** (2015) 1530022, [arXiv:1412.6352](#).

- [16] T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, Comput. Phys. Commun. **178** (2008) 852, [arXiv:0710.3820](#); T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, [arXiv:hep-ph/0603175](#).
- [17] I. Belyaev *et al.*, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, J. Phys. Conf. Ser. **331** (2011) 032047.
- [18] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [19] P. Golonka and Z. Was, *PHOTOS Monte Carlo: A precision tool for QED corrections in Z and W decays*, Eur. Phys. J. **C45** (2006) 97, [arXiv:hep-ph/0506026](#).
- [20] Geant4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; Geant4 collaboration, S. Agostinelli *et al.*, *Geant4: A simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [21] M. Clemencic *et al.*, *The LHCb simulation application, Gauss: Design, evolution and experience*, J. Phys. Conf. Ser. **331** (2011) 032023.
- [22] Particle Data Group, K. A. Olive *et al.*, *Review of particle physics*, Chin. Phys. **C38** (2014) 090001, and 2015 update.
- [23] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and regression trees*, Wadsworth international group, Belmont, California, USA, 1984.
- [24] R. E. Schapire and Y. Freund, *A decision-theoretic generalization of on-line learning and an application to boosting*, J. Comput. Syst. Sci. **55** (1997) 119.
- [25] M. Pivk and F. R. Le Diberder, *sPlot: A statistical tool to unfold data distributions*, Nucl. Instrum. Meth. **A555** (2005) 356, [arXiv:physics/0402083](#).
- [26] T. Skwarnicki, *A study of the radiative cascade transitions between the Upsilon-prime and Upsilon resonances*, PhD thesis, Institute of Nuclear Physics, Krakow, 1986, DESY-F31-86-02.
- [27] LHCb collaboration, R. Aaij *et al.*, *Opposite-side flavour tagging of B mesons at the LHCb experiment*, Eur. Phys. J. **C72** (2012) 2022, [arXiv:1202.4979](#).
- [28] LHCb collaboration, R. Aaij *et al.*, *B flavour tagging using charm decays at the LHCb experiment*, JINST **10** (2015) P10005, [arXiv:1507.07892](#).
- [29] LHCb collaboration, R. Aaij *et al.*, *New algorithms for identifying the flavour of B<sup>0</sup> mesons using pions and protons*, [arXiv:1610.06019](#), submitted to Eur. Phys. J.
- [30] LHCb collaboration, R. Aaij *et al.*, *Measurement of the semileptonic CP asymmetry in B<sup>0</sup>- $\bar{B}^0$  mixing*, Phys. Rev. Lett. **114** (2015) 041601, [arXiv:1409.8586](#).
- [31] LHCb collaboration, R. Aaij *et al.*, *Measurement of the  $\bar{B}^0$ -B<sup>0</sup> and  $\bar{B}_s^0$ -B<sub>s</sub><sup>0</sup> production asymmetries in pp collisions at  $\sqrt{s} = 7$  TeV*, Phys. Lett. **B739** (2014) 218, [arXiv:1408.0275](#).

- [32] C. de Boor, *A practical guide to splines (revised edition)*, Springer, New York, NY, USA, 2001.
- [33] B. Efron, *Bootstrap methods: Another look at the jackknife*, Ann. Statist. **7** (1979) 1.

# Supplemental material

## Tagging calibration

For the flavor tagging calibration, a fit to the background-subtracted decay time distribution of  $B^0 \rightarrow D_s^+ D^-$  candidates is used. In the first step, the mass fit, the  $B^0 \rightarrow D_s^+ D^-$  signal is modeled as two Crystal Ball functions that share a common mean, but have different widths and tail parameters that are obtained from simulations. The  $B_s^0 \rightarrow D_s^- D^+$  background component is modeled similarly and shares all shape parameters with  $B^0 \rightarrow D_s^+ D^-$  except for the peak position, which is constrained to be  $87.35 \text{ MeV}/c^2$  larger. The combinatorial background component is modeled as an exponential function. The  $B^0 \rightarrow D_s^+ D^-$  yield is found to be  $16736 \pm 134$ . The invariant mass distribution of  $B^0 \rightarrow D_s^+ D^-$  candidates is shown in Fig. 3 with the PDF projection overlaid.




Figure 3: Masses of  $B^0 \rightarrow D_s^+ D^-$  candidates and projected PDFs, shown with a linear scale on the vertical axis (a) and a logarithmic scale (b). The solid line is the PDF projection, the blue dashed line represents the  $B^0 \rightarrow D_s^+ D^-$  component, while the dash-dotted cyan (dotted green) line represents the combinatorial ( $B_s^0 \rightarrow D_s^- D^+$ ) background.

The decay time fit to the  $B^0 \rightarrow D_s^+ D^-$  candidates uses a modified version of the PDF from the  $B^0 \rightarrow D^+ D^-$  fit where  $S$  and  $C$  are fixed to zero and unity, respectively, and the production flavor variables  $d'$  and  $d$  are replaced by the mixing state (+1 if the production flavor and the reconstructed decay flavor are the same, and  $-1$  otherwise). Additional modifications are implemented to treat the production asymmetry correctly after replacing  $d'$  by the mixing state, to allow for a flavor-specific asymmetry  $a_{\text{sl}}^d$  (fixed to the latest LHCb measurement [30]) and to include an asymmetry in the detection efficiency for  $D_s^+ D^-$  and  $D_s^- D^+$  (only for the evaluation of systematic uncertainties). The  $B^0$  oscillation frequency,  $\Delta m$ , and the mean lifetime,  $\tau$ , are fixed. The associated systematic uncertainties are taken to be the changes in the calibration parameters when the quantities that were fixed are varied within their uncertainties.

The calibration function for initial  $B^0$  and  $\bar{B}^0$  mesons is given by

$$\omega(\eta|d) = p_0 + d \frac{\Delta p_0}{2} + \left( p_1 + d \frac{\Delta p_1}{2} \right) (\eta - \langle \eta \rangle). \quad (4)$$

Here,  $d$  is +1 for mesons whose initial flavor is  $B^0$  and  $-1$  for  $\bar{B}^0$ . The predicted, per-candidate mistag rate is  $\eta$  and  $\omega$  the calibrated, per-candidate mistag rate. The value of  $\eta$

when averaged over all tagged candidates is  $\langle \eta \rangle$ . The calibration parameters are  $p_0$ ,  $\Delta p_0$ ,  $p_1$ , and  $\Delta p_1$ , and are to be understood as follows. Averaging the calibrated mistag rate over all candidates,  $p_0$  is the value obtained when further averaging over  $B^0$  and  $\bar{B}^0$ , and  $\Delta p_0$  is the difference between the average mistag rates for  $B^0$  and  $\bar{B}^0$ . The coefficients  $p_1$  and  $\Delta p_1$  describe the linear relationship between  $\eta$  and  $\omega$ ; again,  $p_1$  is averaged over  $B^0$  and  $\bar{B}^0$  and  $\Delta p_1$  is the difference between  $B^0$  and  $\bar{B}^0$ . For perfectly calibrated taggers the following two relations hold

$$\begin{aligned} p_0 &= \langle \eta \rangle \\ p_1 &= 1, \end{aligned} \tag{5}$$

and the tagging asymmetries vanish:  $\Delta p_1 = \Delta p_0 = 0$ . The OS calibration parameters are determined to be

$$\begin{aligned} p_{1,\text{OS}} &= 1.07 \pm 0.07 \text{ (stat)} \pm 0.01 \text{ (syst)}, \\ p_{0,\text{OS}} &= 0.369 \pm 0.008 \text{ (stat)} \pm 0.010 \text{ (syst)}, \\ \langle \eta_{\text{OS}} \rangle &= 0.3627, \\ \Delta p_{1,\text{OS}} &= 0.03 \pm 0.11 \text{ (stat)} \pm 0.03 \text{ (syst)}, \\ \Delta p_{0,\text{OS}} &= -0.009 \pm 0.012 \text{ (stat)} \pm 0.001 \text{ (syst)}. \end{aligned} \tag{6}$$

The SS calibration parameters are determined to be

$$\begin{aligned} p_{1,\text{SS}} &= 0.84 \pm 0.09 \text{ (stat)} \pm 0.01 \text{ (syst)}, \\ p_{0,\text{SS}} &= 0.430 \pm 0.006 \text{ (stat)} \pm 0.009 \text{ (syst)}, \\ \langle \eta_{\text{SS}} \rangle &= 0.4282, \\ \Delta p_{1,\text{SS}} &= 0.07 \pm 0.13 \text{ (stat)} \pm 0.05 \text{ (syst)}, \\ \Delta p_{0,\text{SS}} &= -0.007 \pm 0.009 \text{ (stat)} \pm 0.001 \text{ (syst)}. \end{aligned} \tag{7}$$

The time-dependent raw mixing asymmetry  $(N_{\text{unmixed}} - N_{\text{mixed}})/(N_{\text{unmixed}} + N_{\text{mixed}})$ , where  $N_{\text{unmixed}}$  is the number of  $B^0 \rightarrow D_s^+ D^-$  decays with a final state that does correspond to the flavor tag, and  $N_{\text{mixed}}$  the number with a final state that does not, as measured using OS or SS taggers, is shown in Fig. 4. In the  $B^0 \rightarrow D^+ D^-$  dataset, the tagging power for events which are tagged only by OS taggers is  $(1.02 \pm 0.09) \%$ , and for events tagged only by SS taggers  $(1.36 \pm 0.19) \%$ ; for events tagged by both OS and SS taggers, the combined tagging power is  $(5.7 \pm 0.5) \%$ . These sum to a tagging power of  $(8.1 \pm 0.5) \%$  for events tagged by OS, SS, or OS and SS taggers.




Figure 4: Raw mixing asymmetry as a function of the  $B^0$  decay time for events tagged by (a) the OS tagger and (b) the SS tagger. The solid line represents the PDF projection.

## LHCb collaboration

R. Aaij<sup>40</sup>, B. Adeva<sup>39</sup>, M. Adinolfi<sup>48</sup>, Z. Ajaltouni<sup>5</sup>, S. Akar<sup>6</sup>, J. Albrecht<sup>10</sup>, F. Alessio<sup>40</sup>, M. Alexander<sup>53</sup>, S. Ali<sup>43</sup>, G. Alkhazov<sup>31</sup>, P. Alvarez Cartelle<sup>55</sup>, A.A. Alves Jr<sup>59</sup>, S. Amato<sup>2</sup>, S. Amerio<sup>23</sup>, Y. Amhis<sup>7</sup>, L. An<sup>41</sup>, L. Anderlini<sup>18</sup>, G. Andreassi<sup>41</sup>, M. Andreotti<sup>17,g</sup>, J.E. Andrews<sup>60</sup>, R.B. Appleby<sup>56</sup>, O. Aquines Gutierrez<sup>11</sup>, F. Archilli<sup>43</sup>, P. d'Argent<sup>12</sup>, J. Arnau Romeu<sup>6</sup>, A. Artamonov<sup>37</sup>, M. Artuso<sup>61</sup>, E. Aslanides<sup>6</sup>, G. Auriemma<sup>26</sup>, M. Baalouch<sup>5</sup>, I. Babuschkin<sup>56</sup>, S. Bachmann<sup>12</sup>, J.J. Back<sup>50</sup>, A. Badalov<sup>38</sup>, C. Baesso<sup>62</sup>, S. Baker<sup>55</sup>, W. Baldini<sup>17</sup>, R.J. Barlow<sup>56</sup>, C. Barschel<sup>40</sup>, S. Barsuk<sup>7</sup>, W. Barter<sup>40</sup>, V. Batozskaya<sup>29</sup>, B. Batsukh<sup>61</sup>, V. Battista<sup>41</sup>, A. Bay<sup>41</sup>, L. Beaucourt<sup>4</sup>, J. Beddow<sup>53</sup>, F. Bedeschi<sup>24</sup>, I. Bediaga<sup>1</sup>, L.J. Bel<sup>43</sup>, V. Bellee<sup>41</sup>, N. Belloli<sup>21,i</sup>, K. Belous<sup>37</sup>, I. Belyaev<sup>32</sup>, E. Ben-Haim<sup>8</sup>, G. Bencivenni<sup>19</sup>, S. Benson<sup>40</sup>, J. Benton<sup>48</sup>, A. Berezhnoy<sup>33</sup>, R. Bernet<sup>42</sup>, A. Bertolin<sup>23</sup>, F. Betti<sup>15</sup>, M.-O. Bettler<sup>40</sup>, M. van Beuzekom<sup>43</sup>, I. Bezshyiko<sup>42</sup>, S. Bifani<sup>47</sup>, P. Billoir<sup>8</sup>, T. Bird<sup>56</sup>, A. Birnkraut<sup>10</sup>, A. Bitadze<sup>56</sup>, A. Bizzeti<sup>18,u</sup>, T. Blake<sup>50</sup>, F. Blanc<sup>41</sup>, J. Blouw<sup>11</sup>, S. Blusk<sup>61</sup>, V. Bocci<sup>26</sup>, T. Boettcher<sup>58</sup>, A. Bondar<sup>36</sup>, N. Bondar<sup>31,40</sup>, W. Bonivento<sup>16</sup>, A. Borgheresi<sup>21,i</sup>, S. Borghi<sup>56</sup>, M. Borisyak<sup>35</sup>, M. Borsato<sup>39</sup>, F. Bossu<sup>7</sup>, M. Boubdir<sup>9</sup>, T.J.V. Bowcock<sup>54</sup>, E. Bowen<sup>42</sup>, C. Bozzi<sup>17,40</sup>, S. Braun<sup>12</sup>, M. Britsch<sup>12</sup>, T. Britton<sup>61</sup>, J. Brodzicka<sup>56</sup>, E. Buchanan<sup>48</sup>, C. Burr<sup>56</sup>, A. Bursche<sup>2</sup>, J. Buytaert<sup>40</sup>, S. Cadeddu<sup>16</sup>, R. Calabrese<sup>17,g</sup>, M. Calvi<sup>21,i</sup>, M. Calvo Gomez<sup>38,m</sup>, A. Camboni<sup>38</sup>, P. Campana<sup>19</sup>, D. Campora Perez<sup>40</sup>, D.H. Campora Perez<sup>40</sup>, L. Capriotti<sup>56</sup>, A. Carbone<sup>15,e</sup>, G. Carboni<sup>25,j</sup>, R. Cardinale<sup>20,h</sup>, A. Cardini<sup>16</sup>, P. Carniti<sup>21,i</sup>, L. Carson<sup>52</sup>, K. Carvalho Akiba<sup>2</sup>, G. Casse<sup>54</sup>, L. Cassina<sup>21,i</sup>, L. Castillo Garcia<sup>41</sup>, M. Cattaneo<sup>40</sup>, Ch. Cauet<sup>10</sup>, G. Cavallero<sup>20</sup>, R. Cenci<sup>24,t</sup>, M. Charles<sup>8</sup>, Ph. Charpentier<sup>40</sup>, G. Chatzikonstantinidis<sup>47</sup>, M. Chefdeville<sup>4</sup>, S. Chen<sup>56</sup>, S.-F. Cheung<sup>57</sup>, V. Chobanova<sup>39</sup>, M. Chrzaszcz<sup>42,27</sup>, X. Cid Vidal<sup>39</sup>, G. Ciezarek<sup>43</sup>, P.E.L. Clarke<sup>52</sup>, M. Clemencic<sup>40</sup>, H.V. Cliff<sup>49</sup>, J. Closier<sup>40</sup>, V. Coco<sup>59</sup>, J. Cogan<sup>6</sup>, E. Cogneras<sup>5</sup>, V. Cogoni<sup>16,40,f</sup>, L. Cojocariu<sup>30</sup>, G. Collazuol<sup>23,o</sup>, P. Collins<sup>40</sup>, A. Comerma-Montells<sup>12</sup>, A. Contu<sup>40</sup>, A. Cook<sup>48</sup>, G. Coombs<sup>40</sup>, S. Coquereau<sup>8</sup>, G. Corti<sup>40</sup>, M. Corvo<sup>17,g</sup>, C.M. Costa Sobral<sup>50</sup>, B. Couturier<sup>40</sup>, G.A. Cowan<sup>52</sup>, D.C. Craik<sup>52</sup>, A. Crocombe<sup>50</sup>, M. Cruz Torres<sup>62</sup>, S. Cunliffe<sup>55</sup>, R. Currie<sup>55</sup>, C. D'Ambrosio<sup>40</sup>, F. Da Cunha Marinho<sup>2</sup>, E. Dall'Occo<sup>43</sup>, J. Dalseno<sup>48</sup>, P.N.Y. David<sup>43</sup>, A. Davis<sup>59</sup>, O. De Aguiar Francisco<sup>2</sup>, K. De Bruyn<sup>6</sup>, S. De Capua<sup>56</sup>, M. De Cian<sup>12</sup>, J.M. De Miranda<sup>1</sup>, L. De Paula<sup>2</sup>, M. De Serio<sup>14,d</sup>, P. De Simone<sup>19</sup>, C.-T. Dean<sup>53</sup>, D. Decamp<sup>4</sup>, M. Deckenhoff<sup>10</sup>, L. Del Buono<sup>8</sup>, M. Demmer<sup>10</sup>, D. Derkach<sup>35</sup>, O. Deschamps<sup>5</sup>, F. Dettori<sup>40</sup>, B. Dey<sup>22</sup>, A. Di Canto<sup>40</sup>, H. Dijkstra<sup>40</sup>, F. Dordei<sup>40</sup>, M. Dorigo<sup>41</sup>, A. Dosil Suárez<sup>39</sup>, A. Dovbnya<sup>45</sup>, K. Dreimanis<sup>54</sup>, L. Dufour<sup>43</sup>, G. Dujany<sup>56</sup>, K. Dungs<sup>40</sup>, P. Durante<sup>40</sup>, R. Dzhelyadin<sup>37</sup>, A. Dziurda<sup>40</sup>, A. Dzyuba<sup>31</sup>, N. Déleage<sup>4</sup>, S. Easo<sup>51</sup>, M. Ebert<sup>52</sup>, U. Egede<sup>55</sup>, V. Egorychev<sup>32</sup>, S. Eidelman<sup>36</sup>, S. Eisenhardt<sup>52</sup>, U. Eitschberger<sup>10</sup>, R. Ekelhof<sup>10</sup>, L. Eklund<sup>53</sup>, Ch. Elsasser<sup>42</sup>, S. Ely<sup>61</sup>, S. Esen<sup>12</sup>, H.M. Evans<sup>49</sup>, T. Evans<sup>57</sup>, A. Falabella<sup>15</sup>, N. Farley<sup>47</sup>, S. Farry<sup>54</sup>, R. Fay<sup>54</sup>, D. Fazzini<sup>21,i</sup>, D. Ferguson<sup>52</sup>, V. Fernandez Albor<sup>39</sup>, A. Fernandez Prieto<sup>39</sup>, F. Ferrari<sup>15,40</sup>, F. Ferreira Rodrigues<sup>1</sup>, M. Ferro-Luzzi<sup>40</sup>, S. Filippov<sup>34</sup>, R.A. Fini<sup>14</sup>, M. Fiore<sup>17,g</sup>, M. Fiorini<sup>17,g</sup>, M. Firlej<sup>28</sup>, C. Fitzpatrick<sup>41</sup>, T. Fiutowski<sup>28</sup>, F. Fleuret<sup>7,b</sup>, K. Fohl<sup>40</sup>, M. Fontana<sup>16,40</sup>, F. Fontanelli<sup>20,h</sup>, D.C. Forshaw<sup>61</sup>, R. Forty<sup>40</sup>, V. Franco Lima<sup>54</sup>, M. Frank<sup>40</sup>, C. Frei<sup>40</sup>, J. Fu<sup>22,q</sup>, E. Furfaro<sup>25,j</sup>, C. Färber<sup>40</sup>, A. Gallas Torreira<sup>39</sup>, D. Galli<sup>15,e</sup>, S. Gallorini<sup>23</sup>, S. Gambetta<sup>52</sup>, M. Gandelman<sup>2</sup>, P. Gandini<sup>57</sup>, Y. Gao<sup>3</sup>, L.M. Garcia Martin<sup>68</sup>, J. García Pardiñas<sup>39</sup>, J. Garra Tico<sup>49</sup>, L. Garrido<sup>38</sup>, P.J. Garsed<sup>49</sup>, D. Gascon<sup>38</sup>, C. Gaspar<sup>40</sup>, L. Gavardi<sup>10</sup>, G. Gazzoni<sup>5</sup>, D. Gerick<sup>12</sup>, E. Gersabeck<sup>12</sup>, M. Gersabeck<sup>56</sup>, T. Gershon<sup>50</sup>, Ph. Ghez<sup>4</sup>, S. Gianì<sup>41</sup>, V. Gibson<sup>49</sup>, O.G. Girard<sup>41</sup>, L. Giubega<sup>30</sup>, K. Gizdov<sup>52</sup>, V.V. Gligorov<sup>8</sup>, D. Golubkov<sup>32</sup>, A. Golutvin<sup>55,40</sup>, A. Gomes<sup>1,a</sup>, I.V. Gorelov<sup>33</sup>, C. Gotti<sup>21,i</sup>, M. Grabalosa Gándara<sup>5</sup>, R. Graciani Diaz<sup>38</sup>, L.A. Granado Cardoso<sup>40</sup>, E. Graugés<sup>38</sup>, E. Graverini<sup>42</sup>, G. Graziani<sup>18</sup>, A. Grecu<sup>30</sup>, P. Griffith<sup>47</sup>, L. Grillo<sup>21,40,i</sup>, B.R. Gruberg Cazon<sup>57</sup>, O. Grünberg<sup>66</sup>, E. Gushchin<sup>34</sup>, Yu. Guz<sup>37</sup>, T. Gys<sup>40</sup>, C. Göbel<sup>62</sup>, T. Hadavizadeh<sup>57</sup>,

C. Hadjivasiliou<sup>5</sup>, G. Haefeli<sup>41</sup>, C. Haen<sup>40</sup>, S.C. Haines<sup>49</sup>, S. Hall<sup>55</sup>, B. Hamilton<sup>60</sup>, X. Han<sup>12</sup>,  
 S. Hansmann-Menzemer<sup>12</sup>, N. Harnew<sup>57</sup>, S.T. Harnew<sup>48</sup>, J. Harrison<sup>56</sup>, M. Hatch<sup>40</sup>, J. He<sup>63</sup>,  
 T. Head<sup>41</sup>, A. Heister<sup>9</sup>, K. Hennessy<sup>54</sup>, P. Henrard<sup>5</sup>, L. Henry<sup>8</sup>, J.A. Hernando Morata<sup>39</sup>,  
 E. van Herwijnen<sup>40</sup>, M. Heß<sup>66</sup>, A. Hicheur<sup>2</sup>, D. Hill<sup>57</sup>, C. Hombach<sup>56</sup>, H. Hopchev<sup>41</sup>,  
 W. Hulsbergen<sup>43</sup>, T. Humair<sup>55</sup>, M. Hushchyn<sup>35</sup>, N. Hussain<sup>57</sup>, D. Hutchcroft<sup>54</sup>, V. Iakovenko<sup>46</sup>,  
 M. Idzik<sup>28</sup>, P. Ilten<sup>58</sup>, R. Jacobsson<sup>40</sup>, A. Jaeger<sup>12</sup>, J. Jalocha<sup>57</sup>, E. Jans<sup>43</sup>, A. Jawahery<sup>60</sup>,  
 F. Jiang<sup>3</sup>, M. John<sup>57</sup>, D. Johnson<sup>40</sup>, C.R. Jones<sup>49</sup>, C. Joram<sup>40</sup>, B. Jost<sup>40</sup>, N. Jurik<sup>61</sup>,  
 S. Kandybei<sup>45</sup>, W. Kanso<sup>6</sup>, M. Karacson<sup>40</sup>, J.M. Kariuki<sup>48</sup>, S. Karodia<sup>53</sup>, M. Kecke<sup>12</sup>,  
 M. Kelsey<sup>61</sup>, I.R. Kenyon<sup>47</sup>, M. Kenzie<sup>49</sup>, T. Ketel<sup>44</sup>, E. Khairullin<sup>35</sup>, B. Khanji<sup>21,40,i</sup>,  
 C. Khurewathanakul<sup>41</sup>, T. Kirn<sup>9</sup>, S. Klaver<sup>56</sup>, K. Klimaszewski<sup>29</sup>, S. Koliiev<sup>46</sup>, M. Kolpin<sup>12</sup>,  
 I. Komarov<sup>41</sup>, R.F. Koopman<sup>44</sup>, P. Koppenburg<sup>43</sup>, A. Kosmyntseva<sup>32</sup>, A. Kozachuk<sup>33</sup>,  
 M. Kozeiha<sup>5</sup>, L. Kravchuk<sup>34</sup>, K. Kreplin<sup>12</sup>, M. Kreps<sup>50</sup>, P. Krokovny<sup>36</sup>, F. Kruse<sup>10</sup>,  
 W. Krzemien<sup>29</sup>, W. Kucewicz<sup>27,l</sup>, M. Kucharczyk<sup>27</sup>, V. Kudryavtsev<sup>36</sup>, A.K. Kuonen<sup>41</sup>,  
 K. Kurek<sup>29</sup>, T. Kvaratskheliya<sup>32,40</sup>, D. Lacarrere<sup>40</sup>, G. Lafferty<sup>56</sup>, A. Lai<sup>16</sup>, D. Lambert<sup>52</sup>,  
 G. Lanfranchi<sup>19</sup>, C. Langenbruch<sup>9</sup>, T. Latham<sup>50</sup>, C. Lazzeroni<sup>47</sup>, R. Le Gac<sup>6</sup>, J. van Leerdam<sup>43</sup>,  
 J.-P. Lees<sup>4</sup>, A. Leflat<sup>33,40</sup>, J. Lefrançois<sup>7</sup>, R. Lefèvre<sup>5</sup>, F. Lemaitre<sup>40</sup>, E. Lemos Cid<sup>39</sup>, O. Leroy<sup>6</sup>,  
 T. Lesiak<sup>27</sup>, B. Leverington<sup>12</sup>, Y. Li<sup>7</sup>, T. Likhomanenko<sup>35,67</sup>, R. Lindner<sup>40</sup>, C. Linn<sup>40</sup>,  
 F. Lionetto<sup>42</sup>, B. Liu<sup>16</sup>, X. Liu<sup>3</sup>, D. Loh<sup>50</sup>, I. Longstaff<sup>53</sup>, J.H. Lopes<sup>2</sup>, D. Lucchesi<sup>23,o</sup>,  
 M. Lucio Martinez<sup>39</sup>, H. Luo<sup>52</sup>, A. Lupato<sup>23</sup>, E. Luppi<sup>17,g</sup>, O. Lupton<sup>57</sup>, A. Lusiani<sup>24</sup>, X. Lyu<sup>63</sup>,  
 F. Machefert<sup>7</sup>, F. Maciuc<sup>30</sup>, O. Maev<sup>31</sup>, K. Maguire<sup>56</sup>, S. Malde<sup>57</sup>, A. Malinin<sup>67</sup>, T. Maltsev<sup>36</sup>,  
 G. Manca<sup>7</sup>, G. Mancinelli<sup>6</sup>, P. Manning<sup>61</sup>, J. Maratas<sup>5,v</sup>, J.F. Marchand<sup>4</sup>, U. Marconi<sup>15</sup>,  
 C. Marin Benito<sup>38</sup>, P. Marino<sup>24,t</sup>, J. Marks<sup>12</sup>, G. Martellotti<sup>26</sup>, M. Martin<sup>6</sup>, M. Martinelli<sup>41</sup>,  
 D. Martinez Santos<sup>39</sup>, F. Martinez Vidal<sup>68</sup>, D. Martins Tostes<sup>2</sup>, L.M. Massacrier<sup>7</sup>,  
 A. Massafferri<sup>1</sup>, R. Matev<sup>40</sup>, A. Mathad<sup>50</sup>, Z. Mathe<sup>40</sup>, C. Matteuzzi<sup>21</sup>, A. Mauri<sup>42</sup>, B. Maurin<sup>41</sup>,  
 A. Mazurov<sup>47</sup>, M. McCann<sup>55</sup>, J. McCarthy<sup>47</sup>, A. McNab<sup>56</sup>, R. McNulty<sup>13</sup>, B. Meadows<sup>59</sup>,  
 F. Meier<sup>10</sup>, M. Meissner<sup>12</sup>, D. Melnychuk<sup>29</sup>, M. Merk<sup>43</sup>, A. Merli<sup>22,q</sup>, E. Michielin<sup>23</sup>,  
 D.A. Milanes<sup>65</sup>, M.-N. Minard<sup>4</sup>, D.S. Mitzel<sup>12</sup>, A. Mogini<sup>8</sup>, J. Molina Rodriguez<sup>62</sup>,  
 I.A. Monroy<sup>65</sup>, S. Monteil<sup>5</sup>, M. Morandin<sup>23</sup>, P. Morawski<sup>28</sup>, A. Mordà<sup>6</sup>, M.J. Morello<sup>24,t</sup>,  
 J. Moron<sup>28</sup>, A.B. Morris<sup>52</sup>, R. Mountain<sup>61</sup>, F. Muheim<sup>52</sup>, M. Mulder<sup>43</sup>, M. Mussini<sup>15</sup>,  
 D. Müller<sup>56</sup>, J. Müller<sup>10</sup>, K. Müller<sup>42</sup>, V. Müller<sup>10</sup>, P. Naik<sup>48</sup>, T. Nakada<sup>41</sup>, R. Nandakumar<sup>51</sup>,  
 A. Nandi<sup>57</sup>, I. Nasteva<sup>2</sup>, M. Needham<sup>52</sup>, N. Neri<sup>22</sup>, S. Neubert<sup>12</sup>, N. Neufeld<sup>40</sup>, M. Neuner<sup>12</sup>,  
 A.D. Nguyen<sup>41</sup>, T.D. Nguyen<sup>41</sup>, C. Nguyen-Mau<sup>41,n</sup>, S. Nieswand<sup>9</sup>, R. Niet<sup>10</sup>, N. Nikitin<sup>33</sup>,  
 T. Nikodem<sup>12</sup>, A. Novoselov<sup>37</sup>, D.P. O'Hanlon<sup>50</sup>, A. Oblakowska-Mucha<sup>28</sup>, V. Obraztsov<sup>37</sup>,  
 S. Ogilvy<sup>19</sup>, R. Oldeman<sup>49</sup>, C.J.G. Onderwater<sup>69</sup>, J.M. Otalora Goicochea<sup>2</sup>, A. Otto<sup>40</sup>,  
 P. Owen<sup>42</sup>, A. Oyanguren<sup>68</sup>, P.R. Pais<sup>41</sup>, A. Palano<sup>14,d</sup>, F. Palombo<sup>22,q</sup>, M. Palutan<sup>19</sup>,  
 J. Panman<sup>40</sup>, A. Papanestis<sup>51</sup>, M. Pappagallo<sup>14,d</sup>, L.L. Pappalardo<sup>17,g</sup>, W. Parker<sup>60</sup>,  
 C. Parkes<sup>56</sup>, G. Passaleva<sup>18</sup>, A. Pastore<sup>14,d</sup>, G.D. Patel<sup>54</sup>, M. Patel<sup>55</sup>, C. Patrignani<sup>15,e</sup>,  
 A. Pearce<sup>56,51</sup>, A. Pellegrino<sup>43</sup>, G. Penso<sup>26</sup>, M. Pepe Altarelli<sup>40</sup>, S. Perazzini<sup>40</sup>, P. Perret<sup>5</sup>,  
 L. Pescatore<sup>47</sup>, K. Petridis<sup>48</sup>, A. Petrolini<sup>20,h</sup>, A. Petrov<sup>67</sup>, M. Petruzzo<sup>22,q</sup>,  
 E. Picatoste Olloqui<sup>38</sup>, B. Pietrzyk<sup>4</sup>, M. Piekies<sup>27</sup>, D. Pinci<sup>26</sup>, A. Pistone<sup>20</sup>, A. Piucci<sup>12</sup>,  
 S. Playfer<sup>52</sup>, M. Plo Casasus<sup>39</sup>, T. Poikela<sup>40</sup>, F. Polci<sup>8</sup>, A. Poluektov<sup>50,36</sup>, I. Polyakov<sup>61</sup>,  
 E. Polycarpo<sup>2</sup>, G.J. Pomery<sup>48</sup>, A. Popov<sup>37</sup>, D. Popov<sup>11,40</sup>, B. Popovici<sup>30</sup>, S. Poslavskii<sup>37</sup>,  
 C. Potterat<sup>2</sup>, E. Price<sup>48</sup>, J.D. Price<sup>54</sup>, J. Prisciandaro<sup>39</sup>, A. Pritchard<sup>54</sup>, C. Prouve<sup>48</sup>,  
 V. Pugatch<sup>46</sup>, A. Puig Navarro<sup>41</sup>, G. Punzi<sup>24,p</sup>, W. Qian<sup>57</sup>, R. Quagliani<sup>7,48</sup>, B. Rachwal<sup>27</sup>,  
 J.H. Rademacker<sup>48</sup>, M. Rama<sup>24</sup>, M. Ramos Pernas<sup>39</sup>, M.S. Rangel<sup>2</sup>, I. Raniuk<sup>45</sup>, G. Raven<sup>44</sup>,  
 F. Redi<sup>55</sup>, S. Reichert<sup>10</sup>, A.C. dos Reis<sup>1</sup>, C. Remon Alepuz<sup>68</sup>, V. Renaudin<sup>7</sup>, S. Ricciardi<sup>51</sup>,  
 S. Richards<sup>48</sup>, M. Rihl<sup>40</sup>, K. Rinnert<sup>54</sup>, V. Rives Molina<sup>38</sup>, P. Robbe<sup>7,40</sup>, A.B. Rodrigues<sup>1</sup>,  
 E. Rodrigues<sup>59</sup>, J.A. Rodriguez Lopez<sup>65</sup>, P. Rodriguez Perez<sup>56</sup>, A. Rogozhnikov<sup>35</sup>, S. Roiser<sup>40</sup>,  
 A. Rollings<sup>57</sup>, V. Romanovskiy<sup>37</sup>, A. Romero Vidal<sup>39</sup>, J.W. Ronayne<sup>13</sup>, M. Rotondo<sup>19</sup>,  
 M.S. Rudolph<sup>61</sup>, T. Ruf<sup>40</sup>, P. Ruiz Valls<sup>68</sup>, J.J. Saborido Silva<sup>39</sup>, E. Sadykhov<sup>32</sup>, N. Sagidova<sup>31</sup>,



B. Saitta<sup>16,f</sup>, V. Salustino Guimaraes<sup>2</sup>, C. Sanchez Mayordomo<sup>68</sup>, B. Sanmartin Sedes<sup>39</sup>, R. Santacesaria<sup>26</sup>, C. Santamarina Rios<sup>39</sup>, M. Santimaria<sup>19</sup>, E. Santovetti<sup>25,j</sup>, A. Sarti<sup>19,k</sup>, C. Satriano<sup>26,s</sup>, A. Satta<sup>25</sup>, D.M. Saunders<sup>48</sup>, D. Savrina<sup>32,33</sup>, S. Schael<sup>9</sup>, M. Schellenberg<sup>10</sup>, M. Schiller<sup>40</sup>, H. Schindler<sup>40</sup>, M. Schlupp<sup>10</sup>, M. Schmelling<sup>11</sup>, T. Schmelzer<sup>10</sup>, B. Schmidt<sup>40</sup>, O. Schneider<sup>41</sup>, A. Schopper<sup>40</sup>, K. Schubert<sup>10</sup>, M. Schubiger<sup>41</sup>, M.-H. Schune<sup>7</sup>, R. Schwemmer<sup>40</sup>, B. Sciascia<sup>19</sup>, A. Sciubba<sup>26,k</sup>, A. Semennikov<sup>32</sup>, A. Sergi<sup>47</sup>, N. Serra<sup>42</sup>, J. Serrano<sup>6</sup>, L. Sestini<sup>23</sup>, P. Seyfert<sup>21</sup>, M. Shapkin<sup>37</sup>, I. Shapoval<sup>45</sup>, Y. Shcheglov<sup>31</sup>, T. Shears<sup>54</sup>, L. Shekhtman<sup>36</sup>, V. Shevchenko<sup>67</sup>, A. Shires<sup>10</sup>, B.G. Siddi<sup>17,40</sup>, R. Silva Coutinho<sup>42</sup>, L. Silva de Oliveira<sup>2</sup>, G. Simi<sup>23,o</sup>, S. Simone<sup>14,d</sup>, M. Sirendi<sup>49</sup>, N. Skidmore<sup>48</sup>, T. Skwarnicki<sup>61</sup>, E. Smith<sup>55</sup>, I.T. Smith<sup>52</sup>, J. Smith<sup>49</sup>, M. Smith<sup>55</sup>, H. Snoek<sup>43</sup>, M.D. Sokoloff<sup>59</sup>, F.J.P. Soler<sup>53</sup>, B. Souza De Paula<sup>2</sup>, B. Spaan<sup>10</sup>, P. Spradlin<sup>53</sup>, S. Sridharan<sup>40</sup>, F. Stagni<sup>40</sup>, M. Stahl<sup>12</sup>, S. Stahl<sup>40</sup>, P. Stefko<sup>41</sup>, S. Stefkova<sup>55</sup>, O. Steinkamp<sup>42</sup>, S. Stemmler<sup>12</sup>, O. Stenyakin<sup>37</sup>, S. Stevenson<sup>57</sup>, S. Stoica<sup>30</sup>, S. Stone<sup>61</sup>, B. Storaci<sup>42</sup>, S. Stracka<sup>24,p</sup>, M. Straticiu<sup>30</sup>, U. Straumann<sup>42</sup>, L. Sun<sup>59</sup>, W. Sutcliffe<sup>55</sup>, K. Swientek<sup>28</sup>, V. Syropoulos<sup>44</sup>, M. Szczekowski<sup>29</sup>, T. Szumlak<sup>28</sup>, S. T'Jampens<sup>4</sup>, A. Tayduganov<sup>6</sup>, T. Tekampe<sup>10</sup>, G. Tellarini<sup>17,g</sup>, F. Teubert<sup>40</sup>, E. Thomas<sup>40</sup>, J. van Tilburg<sup>43</sup>, M.J. Tilley<sup>55</sup>, V. Tisserand<sup>4</sup>, M. Tobin<sup>41</sup>, S. Tolc<sup>49</sup>, L. Tomassetti<sup>17,g</sup>, D. Tonelli<sup>40</sup>, S. Topp-Joergensen<sup>57</sup>, F. Toriello<sup>61</sup>, E. Tournefier<sup>4</sup>, S. Tourneur<sup>41</sup>, K. Trabelsi<sup>41</sup>, M. Traill<sup>53</sup>, M.T. Tran<sup>41</sup>, M. Tresch<sup>42</sup>, A. Trisovic<sup>40</sup>, A. Tsaregorodtsev<sup>6</sup>, P. Tsopelas<sup>43</sup>, A. Tully<sup>49</sup>, N. Tuning<sup>43</sup>, A. Ukleja<sup>29</sup>, A. Ustyuzhanin<sup>35</sup>, U. Uwer<sup>12</sup>, C. Vacca<sup>16,f</sup>, V. Vagnoni<sup>15,40</sup>, A. Valassi<sup>40</sup>, S. Valat<sup>40</sup>, G. Valenti<sup>15</sup>, A. Vallier<sup>7</sup>, R. Vazquez Gomez<sup>19</sup>, P. Vazquez Regueiro<sup>39</sup>, S. Vecchi<sup>17</sup>, M. van Veghel<sup>43</sup>, J.J. Velthuis<sup>48</sup>, M. Veltri<sup>18,r</sup>, G. Veneziano<sup>41</sup>, A. Venkateswaran<sup>61</sup>, M. Vernet<sup>5</sup>, M. Vesterinen<sup>12</sup>, B. Viaud<sup>7</sup>, D. Vieira<sup>1</sup>, M. Vieites Diaz<sup>39</sup>, X. Vilasis-Cardona<sup>38,m</sup>, V. Volkov<sup>33</sup>, A. Vollhardt<sup>42</sup>, B. Voneki<sup>40</sup>, A. Vorobyev<sup>31</sup>, V. Vorobyev<sup>36</sup>, C. Vob<sup>66</sup>, J.A. de Vries<sup>43</sup>, C. Vázquez Sierra<sup>39</sup>, R. Waldi<sup>66</sup>, C. Wallace<sup>50</sup>, R. Wallace<sup>13</sup>, J. Walsh<sup>24</sup>, J. Wang<sup>61</sup>, D.R. Ward<sup>49</sup>, H.M. Wark<sup>54</sup>, N.K. Watson<sup>47</sup>, D. Websdale<sup>55</sup>, A. Weiden<sup>42</sup>, M. Whitehead<sup>40</sup>, J. Wicht<sup>50</sup>, G. Wilkinson<sup>57,40</sup>, M. Wilkinson<sup>61</sup>, M. Williams<sup>40</sup>, M.P. Williams<sup>47</sup>, M. Williams<sup>58</sup>, T. Williams<sup>47</sup>, F.F. Wilson<sup>51</sup>, J. Wimberley<sup>60</sup>, J. Wishahi<sup>10</sup>, W. Wislicki<sup>29</sup>, M. Witek<sup>27</sup>, G. Wormser<sup>7</sup>, S.A. Wotton<sup>49</sup>, K. Wraight<sup>53</sup>, S. Wright<sup>49</sup>, K. Wyllie<sup>40</sup>, Y. Xie<sup>64</sup>, Z. Xing<sup>61</sup>, Z. Xu<sup>41</sup>, Z. Yang<sup>3</sup>, H. Yin<sup>64</sup>, J. Yu<sup>64</sup>, X. Yuan<sup>36</sup>, O. Yushchenko<sup>37</sup>, K.A. Zarebski<sup>47</sup>, M. Zavertyaev<sup>11,c</sup>, L. Zhang<sup>3</sup>, Y. Zhang<sup>7</sup>, Y. Zhang<sup>63</sup>, A. Zhelezov<sup>12</sup>, Y. Zheng<sup>63</sup>, A. Zhokhov<sup>32</sup>, X. Zhu<sup>3</sup>, V. Zhukov<sup>9</sup>, S. Zucchelli<sup>15</sup>.

<sup>1</sup>Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

<sup>2</sup>Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

<sup>3</sup>Center for High Energy Physics, Tsinghua University, Beijing, China

<sup>4</sup>LAPP, Université Savoie Mont-Blanc, CNRS/IN2P3, Annecy-Le-Vieux, France

<sup>5</sup>Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

<sup>6</sup>CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

<sup>7</sup>LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

<sup>8</sup>LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

<sup>9</sup>I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

<sup>10</sup>Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

<sup>11</sup>Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

<sup>12</sup>Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

<sup>13</sup>School of Physics, University College Dublin, Dublin, Ireland

<sup>14</sup>Sezione INFN di Bari, Bari, Italy

<sup>15</sup>Sezione INFN di Bologna, Bologna, Italy

<sup>16</sup>Sezione INFN di Cagliari, Cagliari, Italy

<sup>17</sup>Sezione INFN di Ferrara, Ferrara, Italy

<sup>18</sup>Sezione INFN di Firenze, Firenze, Italy

<sup>19</sup>Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy

<sup>20</sup>Sezione INFN di Genova, Genova, Italy

<sup>21</sup>Sezione INFN di Milano Bicocca, Milano, Italy

- <sup>22</sup> *Sezione INFN di Milano, Milano, Italy*
- <sup>23</sup> *Sezione INFN di Padova, Padova, Italy*
- <sup>24</sup> *Sezione INFN di Pisa, Pisa, Italy*
- <sup>25</sup> *Sezione INFN di Roma Tor Vergata, Roma, Italy*
- <sup>26</sup> *Sezione INFN di Roma La Sapienza, Roma, Italy*
- <sup>27</sup> *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- <sup>28</sup> *AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- <sup>29</sup> *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- <sup>30</sup> *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- <sup>31</sup> *Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia*
- <sup>32</sup> *Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*
- <sup>33</sup> *Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- <sup>34</sup> *Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia*
- <sup>35</sup> *Yandex School of Data Analysis, Moscow, Russia*
- <sup>36</sup> *Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia*
- <sup>37</sup> *Institute for High Energy Physics (IHEP), Protvino, Russia*
- <sup>38</sup> *ICCUB, Universitat de Barcelona, Barcelona, Spain*
- <sup>39</sup> *Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- <sup>40</sup> *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- <sup>41</sup> *Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- <sup>42</sup> *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- <sup>43</sup> *Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- <sup>44</sup> *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- <sup>45</sup> *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- <sup>46</sup> *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- <sup>47</sup> *University of Birmingham, Birmingham, United Kingdom*
- <sup>48</sup> *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- <sup>49</sup> *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- <sup>50</sup> *Department of Physics, University of Warwick, Coventry, United Kingdom*
- <sup>51</sup> *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- <sup>52</sup> *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- <sup>53</sup> *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- <sup>54</sup> *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- <sup>55</sup> *Imperial College London, London, United Kingdom*
- <sup>56</sup> *School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- <sup>57</sup> *Department of Physics, University of Oxford, Oxford, United Kingdom*
- <sup>58</sup> *Massachusetts Institute of Technology, Cambridge, MA, United States*
- <sup>59</sup> *University of Cincinnati, Cincinnati, OH, United States*
- <sup>60</sup> *University of Maryland, College Park, MD, United States*
- <sup>61</sup> *Syracuse University, Syracuse, NY, United States*
- <sup>62</sup> *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to <sup>2</sup>*
- <sup>63</sup> *University of Chinese Academy of Sciences, Beijing, China, associated to <sup>3</sup>*
- <sup>64</sup> *Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China, associated to <sup>3</sup>*
- <sup>65</sup> *Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to <sup>8</sup>*
- <sup>66</sup> *Institut für Physik, Universität Rostock, Rostock, Germany, associated to <sup>12</sup>*
- <sup>67</sup> *National Research Centre Kurchatov Institute, Moscow, Russia, associated to <sup>32</sup>*
- <sup>68</sup> *Instituto de Física Corpuscular (IFIC), Universitat de Valencia-CSIC, Valencia, Spain, associated to <sup>38</sup>*
- <sup>69</sup> *Van Swinderen Institute, University of Groningen, Groningen, The Netherlands, associated to <sup>43</sup>*
- <sup>a</sup> *Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil*
- <sup>b</sup> *Laboratoire Leprince-Ringuet, Palaiseau, France*
- <sup>c</sup> *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*
- <sup>d</sup> *Università di Bari, Bari, Italy*
- <sup>e</sup> *Università di Bologna, Bologna, Italy*
- <sup>f</sup> *Università di Cagliari, Cagliari, Italy*

- <sup>g</sup> *Università di Ferrara, Ferrara, Italy*  
<sup>h</sup> *Università di Genova, Genova, Italy*  
<sup>i</sup> *Università di Milano Bicocca, Milano, Italy*  
<sup>j</sup> *Università di Roma Tor Vergata, Roma, Italy*  
<sup>k</sup> *Università di Roma La Sapienza, Roma, Italy*  
<sup>l</sup> *AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*  
<sup>m</sup> *LIFAEELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*  
<sup>n</sup> *Hanoi University of Science, Hanoi, Viet Nam*  
<sup>o</sup> *Università di Padova, Padova, Italy*  
<sup>p</sup> *Università di Pisa, Pisa, Italy*  
<sup>q</sup> *Università degli Studi di Milano, Milano, Italy*  
<sup>r</sup> *Università di Urbino, Urbino, Italy*  
<sup>s</sup> *Università della Basilicata, Potenza, Italy*  
<sup>t</sup> *Scuola Normale Superiore, Pisa, Italy*  
<sup>u</sup> *Università di Modena e Reggio Emilia, Modena, Italy*  
<sup>v</sup> *Iligan Institute of Technology (IIT), Iligan, Philippines*