The Discovery of a Low-Luminosity SPIRAL DRAGN.

D. D. Mulcahy¹, M. Y. Mao¹, I. Mitsuishi², A. M. M. Scaife¹, A. O. Clarke¹, Y. Babazaki², H. Kobayashi², R. Suganuma², H. Matsumoto³, and Y. Tawara²

- Jodrell Bank Centre for Astrophysics, Alan Turing Building, School of Physics and Astronomy, The University of Manchester, Oxford Road, Manchester, M13 9PL, U.K. e-mail: david.mulcahy@manchester.ac.uk
- Division of Particle and Astrophysical Science, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi, 464-8602 JAPAN
- Kobayashi-Maskawa Institute for the Origin of Particles and the Universe, Nagoya University, Furo-cho, Chikusa-ku, Nagoya,

ABSTRACT

Standard galaxy formation models predict that large-scale double-lobed radio sources, known as DRAGNs, will always be hosted by elliptical galaxies. In spite of this, in recent years a small number of spiral galaxies have also been found to host such sources. These so-called spiral DRAGNs are still extremely rare, with only ~ 5 cases being widely accepted. Here we report on the serendipitous discovery of a new spiral DRAGN in data from the Giant Metrewave Radio Telescope (GMRT) at 322 MHz. The host galaxy, MCG+07-47-10, is a face-on late-type Sbc galaxy with distinctive spiral arms and prominent bulge suggesting a high black hole mass. Using WISE infra-red and GALEX UV data we show that this galaxy has a star formation rate of 0.16-0.75 M_{\odot} yr⁻¹, and that the radio luminosity is dominated by star-formation. We demonstrate that this spiral DRAGN has similar environmental properties to others of this class, but has a comparatively low radio luminosity of $L_{1.4\,\mathrm{GHz}} = 1.12 \times 10^{22}\,\mathrm{W\ Hz^{-1}}$, two orders of magnitude smaller than other known spiral DRAGNs. We suggest that this may indicate the existence of a previously unknown low-luminosity population

Key words. Galaxies: spiral – Galaxies: jets – Radio continuum: galaxies

The University of Manchester, Oxford Road, Manchester, M13 e-mail: david.mulcahy@manchester.ac.uk

2 Division of Particle and Astrophysical Science, Nagoya Univers

3 Kobayashi-Maskawa Institute for the Origin of Particles and tachi, 464-8602 JAPAN

Received 16 August 2016; accepted 13 September 2016

ABST

Standard galaxy formation models predict that large-scale double-elliptical galaxies. In spite of this, in recent years a small number of so-called spiral DRAGNs are still extremely rare, with only ~ 5 or discovery of a new spiral DRAGN in data from the Giant Metropy MCG+07-47-10, is a face-on late-type Sbc galaxy with distinctive mass. Using WISE infra-red and GALEX UV data we show that the radio luminosity is dominated by star-formation. We demonstrothers of this class, but has a comparatively low radio luminosity than other known spiral DRAGNs. We suggest that this may indicated of spiral DRAGNs.

Key words. Galaxies: spiral – Galaxies: jets – Radio continuum: Spiral DRAGNs (Double-lobed Radio sources Associated with Galactic Nuclei, Leahy 1993) are spiral galaxies that host large-scale double-lobed radio sources. The existence of such sources is in contradiction to our existing models of galaxy formation (e.g. Hopkins et al. 2008), which predict that DRAGNs should be hosted exclusively by elliptical galaxies. Indeed, until recently, observations of DRAGNs in the local Universe confirmed this expectation (e.g. Matthews et al. 1964; Urry & Padovani 1995; Best et al. 2005).

Elliptical galaxies are formed as a result of mergers, the phenomenology of which also triggers the formation of DRAGNs (Chiaberge & Marconi 2011; Chiaberge et al. 2015). However, a spiral galaxy's structure cannot withstand a major merger. Moreover, morphological transition from spiral to elliptical is thought to be a one-way process, at least in the local Universe. Consequently, the standard galaxy formation model does not predict the existence of spiral DRAGNs.

quently, the standard galaxy formation model does not predict the existence of spiral DRAGNs.

Nonetheless, a number of spiral DRAGNs have been discovered in recent years (e.g. Ledlow et al. 2001; Hota et al. 2011; Bagchi et al. 2014; Mao et al. 2015; Singh et al. 2015). While the first three spiral DRAGN discoveries were serendipitous, Mao et al. (2015) performed the first systematic search for these sources. Using Galaxy Zoo (Lintott et al. 2008) morphological classifications were cross-matched with the Faint Images of the Radio Sky at Twenty-Centimeters (FIRST, Becker et al. 1995) and the NRAO VLA Sky-Survey (NVSS, Condon et al. 1998). In this study, only one spiral DRAGN was found above L_{1.4 GHz} $= 10^{23} \,\mathrm{W\,Hz^{-1}}$. Singh et al. (2015) performed a similar analy-

sis using the spiral galaxy catalogue of Meert et al. (2015) and reported the identification of four spiral DRAGNs in those data, including one that was previously known and three that were unknown. However, the precise identification of spiral DRAGN hosts remains contentious in the literature and to date the existence of only 5 spiral DRAGNs are widely accepted.

DRAGNs with spiral hosts may represent a rare phenomenon of elliptical galaxies transitioning back into spirals through accretion of gas and stars, perhaps from a companion. A key question is whether spiral DRAGNs are a result of non-standard physical properties, a result of their environment, or perhaps a combination of their nature and nurture. Studying spiral DRAGNs, as well as establishing their numbers more exactly, is vital in order to reconcile their role in standard galaxy formation theories.

In this Letter, we present the discovery of a new spiral DRAGN at 325 MHz with the Giant Meterwave Radio Telescope (GMRT; Swarup 1990). In Sect. 2 we outline the data processing and imaging steps. In Sect. 3, we present the discovery of this new spiral DRAGN with a description of both its radio morphology and that of the host galaxy, as understood from available multi-wavelength data. In Sect. 4, using Infra-red and UV data we demonstrate that the host galaxy is star-forming and we compare its star formation rate to other spiral DRAGNs. Finally, in Sect. 5 we discuss the nature of this object and state our conclusions. In this work we assume a ACDM cosmology with $H_0 = 69.6 \,\mathrm{km s^{-1} Mpc^{-1}}$, $\Omega_{\rm m} = 0.286$ and $\Omega_{\Lambda} = 0.714$ (Bennett et al. 2014), which we use for the calculation of distance, luminosity and star formation rate. At a redshift of z = 0.017, these values result in a conversion of 0.348 kpc/". All uncertainties are quoted at 1 σ .

2. Observations and Data Reduction

Observations of the galaxy groups NGC7618 & UGC 12491 were performed in full synthesis mode with the Giant Meterwave Radio Telescope (GMRT) at 325 MHz. The GMRT is a full aperture synthesis telescope located near Pune, India (Swarup 1990). It consists of 30 steerable dishes that are 45 m in diameter, with a longest interferometric baseline of 25.5 km, and 14 additional antennas located in a central 1 sq. km, to provide dense uv coverage at short spacings. The telescope operates at six frequencies: 150, 230, 325, 610, and 1420 MHz. In this work the 325 MHz receiver was utilised. The Full Width Half Maximum (FWHM) of the GMRT primary beam at 325 MHz is approximately $81' \pm 4'$.

The aforementioned observation took place on 16th August 2014 under project code 25.059 (PI Mitsuishi). The observation was a single pointing with a phase center of (J2000) 23^h 19^m 19^s +42° 54′ 50″. The bright radio source 3C48 was used as a flux calibrator, tied to the Scaife & Heald (2012) flux density scale.

The dataset was reduced using the Source Peeling and Atmospheric Modelling (SPAM) software (Intema et al. 2009). The SPAM software performed an initial phase calibration and astrometry correction using a sky model derived from the NVSS (Condon et al. 1998), followed by three rounds of direction-independent phase-only self-calibration. Following this, SPAM performed facet-based direction-dependent calibration on strong sources within the primary beam FWHM, and utilised the solutions from this calibration to fit a global ionospheric model. During this calibration process, various automated flagging routines were used between cycles of imaging and self-calibration in order to reduce residual RFI and clip statistical outliers.

The final primary beam corrected image towards MCG+07-47-10 is shown in Fig. 1 and has a resolution of 9.4×8.3 ". Images were made using an AIPS ROBUST value of -1.0. This parameter value was selected to achieve an optimal compromise between sensitivity and resolution. More details of these observations and data reduction can be found in Mitsuishi et al., in prep.

In order to isolate the integrated flux density of the separate source components (radio lobes and core) the source finding software, the Python Blob Detection and Source Measurement software, (PYBDSM¹) was first used to identify confusing compact sources. To do this, a local rms noise value (σ) was calculated in order to identify all significant peaks above a threshold of 7σ . Following standard practice, PYBDSM then forms islands of contiguous emission down to a threshold of 5σ and an elliptical Gaussian is fitted to each island, of which 625 were found in the entire field. Each Gaussian was then subtracted from the image and the residual data used to determine the integrated flux of the radio lobes. The measured integrated fluxes are presented in Table. 1. The figures shown in this Letter are not the source subtracted images.

3. Discovery of spiral DRAGN - MCG+07-47-10

At a distance of approximately 22' from the field phase centre, a Fanaroff-Riley type II radio galaxy is identified. This radio galaxy is shown in Fig. 1, where the lobes of this object are denoted as "A" and "B" and the central host galaxy, MCG+07-47-10, as "C". Closeups of the individual radio lobes are shown in

Fig. 2. Co-ordinates for each of the lobes and the host galaxy are listed in Table 1. In particular, lobe "B" displays weak extended emission tracing back to the host galaxy (see Fig. 2, Right). Such weak emission is only now detectable due to the high sensitivity of this observation and would have not been detected in surveys such as the NVSS or FIRST.




Figure 1: GMRT data at 322 MHz. The local rms noise around the source is approximately $\sigma_{\rm rms} = 68\,\mu{\rm Jy/beam}$. Contours are shown at 3, 5, 8, 10, 12, 24 & 48 $\sigma_{\rm rms}$. The GMRT synthesized beam has dimensions of 9.4×8.3" and is shown in the bottom left corner as a filled-in ellipse.

On inspection of the Digitized Sky Survey (DSS), the host galaxy (denoted "C" in Fig. 1), MCG+07-47-10, is a late type galaxy with a classification of Sbc (Vorontsov-Vel'Yaminov & Arkhipova 1968), a galaxy with spiral arms and a bulge. The limited resolution of the DSS means that the presence of a bar cannot be ruled out. The optical DSS image of this galaxy can be seen in Fig. 3 with the 5σ radio contour overlaid. The spiral arms of this face-on galaxy are clearly visible and extended radio emission is seen coincident with the optical disk. If this emission was solely due to the core of the AGN, one would expect highly compact emission, unresolved by the GMRT in these data. It can be seen from Fig. 1 that the radio emission in this case is clearly resolved, with a diameter measured from the 3 σ contour of approximately 45", equivalent to 5 synthesized beam widths. Such extended radio emission indicates that active star formation is occurring throughout the disk and is due to the injection of Cosmic Ray electrons (CREs) into the interstellar medium (ISM) via the action of Supernovae. These CREs produce non-thermal synchrotron emission, which is observable here.

This area is not covered by the Sloan Digital Sky Survey (SDSS²; Alam et al. 2015) and currently no spectroscopic redshift information is available for this galaxy. However, MCG+07-47-10 is located close to the galaxy groups NGC 7618 (z=0.0173) and UGC 12491 (z=0.0173), the central galaxies of two nearby, roughly equal mass galaxy groups. Kraft et al. (2006) found through X-ray observations that these galaxy groups are in the process of merging. Assuming that MCG+07-47-10 is associated with one of these galaxy groups, we adopt an approximate redshift of $z\approx0.017$. At this redshift, MCG+07-47-10 would be located at a projected distance of 584 kpc (28') from NGC 7618 and 375 kpc (18') from UGC 12491. Given

¹ https://dl.dropboxusercontent.com/u/1948170/html/index.html

² http://www.sdss.org/

Src.	ID	Right Ascension	Declination	1.4 GHz Flux	322 MHz Flux	Spectral
		(J2000)	(J2000)	(mJy)	(mJy)	Index
A	Northern Lobe	23 ^h 18 ^m 06 ^s	+43° 18′ 18″	7.8±1.3	20.0±2.0	-0.6±0.3
В	Southern Lobe	23 ^h 18 ^m 43 ^s	+43° 12′ 20′′	5.5 ± 0.5	16.1±1.6	-0.73 ± 0.21
C	MCG+07-47-10	23 ^h 18 ^m 33 ^s	+43° 14′ 49′′	3.9 ± 0.4	12.4 ± 1.5	-0.79 ± 0.24

Table 1: Locations of the radio lobes and host galaxy of the spiral DRAGN with integrated fluxes and spectral indices.




Figure 2: Left: Zoomed in image of the northern lobe and host galaxy from Fig. 1. Right: Zoomed in image of the southern lobe and host galaxy from Fig. 1. The measured rms noise (σ) around the source is approximately 68 μ Jy/beam. The contours for both images are at levels of 3, 5, 8, 10, 12, 24, 48 × σ . The resolution is 9.4×8.3" and is shown in the bottom right/left corner as a filled-in ellipse.




Figure 3: Optical DSS image of the host galaxy (MCG+07-47-10) for the spiral DRAGN. The contour shows the 5σ level of the 322 MHz radio data.

a virial radius for these galaxy groups of approximately 600-800 kpc (Kraft et al. 2006), MCG+07-47-10 would therefore be found within the virial radius of both groups.

From the GMRT image, we find that this new spiral DRAGN has an angular linear extent of $\sim 9'$, which equates to a physical size of 207 kpc at our assumed redshift. Additionally, the optical diameter of MCG+07-47-10 is found to be 12.02 kpc.

At an assumed redshift of z=0.017 and measured radio flux density values in Table. 1 indicate a radio luminosity at 322 MHz for this proposed spiral DRAGN (lobes and host galaxy) of approximately $L_{322\,\mathrm{MHz}}=3.15\times10^{22}\,\mathrm{Watts}\,\mathrm{Hz}^{-1}$ and $L_{1.4\,\mathrm{GHz}}=1.12\times10^{22}\,\mathrm{Watts}\,\mathrm{Hz}^{-1}$. This is significantly weaker to other spiral DRAGNS such as J1649+2635 with $L_{1.4\,\mathrm{GHz}}=1.03\times10^{24}\,\mathrm{Watts}\,\mathrm{Hz}^{-1}$ (Mao et al. 2015).

3.1. Additional radio data and integrated fluxes

MCG+07-47-10 and both lobes are detected by the NVSS (Condon et al. 1998), but not by the VLA Low-frequency Sky Survey redux (VLSSr, Lane et al. 2014; Cohen et al. 2007), TIFR GMRT Sky Survey (TGSS ADR, Intema et al. 2016) nor the WEsterbork Northern Sky Survey (WENSS, Rengelink et al. 1997). In addition, the FIRST survey does not have coverage of this area of the sky. In the NVSS, the three main components are detected and the catalogue integrated fluxes for each component are shown in Table. 1.

From the combination of the NVSS data and the GMRT data presented in this work, the host galaxy is found to have a spectral index of $\alpha = -0.79 \pm 0.24$, which is a typical value for a star-forming spiral galaxies ($\alpha = -0.74 \pm 0.03$, Gioia et al. 1982).

4. Star formation rate

MCG+07-47-10 is clearly detected in all four WISE bands (Wright et al. 2010). Using colour-colour relationships (Lacy et al. 2004; Yan et al. 2013), the measured magnitude difference [W1]-[W2] = 0.184, indicates that MCG+07-47-10 is classified as a star-forming galaxy, with values greater than 0.8 signifying an AGN. Normal spiral galaxies are seen to have a range of magnitude differences of $\approx 0.0\text{-}0.6$ (Wright et al. 2010).

Using the derived relations between star formation rate (SFR) and observed luminosity in the WISE W3 and W4 bands (Jarrett et al. 2013), we find the SFR for MCG+07-47-10 at $12\mu m$ to be $\Sigma_{12}=0.75\,M_{\odot}\,yr^{-1}$ and at $22\,\mu m$ to be $\Sigma_{22}=0.51\,M_{\odot}\,yr^{-1}$. In addition, MCG+07-47-10 is also detected in the near UV by GALEX (Martin et al. 2005). Based on Kennicutt (1998), we find the UV SFR for MCG+07-47-10 to be $\Sigma_{UV}=0.16\,M_{\odot}\,yr^{-1}$. Assuming no contribution from an AGN, the SFR for MCG+07-47-10 derived from the radio data presented here is $\Sigma_{1.4\,GHz}=0.63\,M_{\odot}\,yr^{-1}$ (Condon 1992, Eq.21), intermediate to the IR and UV derived SFRs. There are large uncertainties

associated with SFR indicators (Hopkins et al. 2003), thus we estimate the SFR to be in the range of 0.16-0.75 M_{\odot} yr⁻¹.

The SFR rate in galaxies can show an enormous range from virtually zero in gas poor ellipticals and dwarf-galaxies to $0.63\,M_\odot\,yr^{-1}$ in gas rich spirals (Kennicutt 1998). For comparison MCG+07-47-10 has a similar SFR to the nearby Sb galaxy, NGC2683, with has a SFR $\approx 0.36\,M_\odot\,yr^{-1}$ determined by Irwin et al. (2012). Compared to other spiral DRAGNS, Mao et al. (2015) found J1649+2635's SFR to be $\approx 0.26\text{-}2.6\,M_\odot\,yr^{-1}$. The two spiral DRAGNs found by Singh et al. (2015) (J0836+0532,J1159+5820) were found to have 9.99 and 2.89 $M_\odot\,yr^{-1}$ respectively. MCG+07-47-10 with a SFR range of 0.16-0.75 $M_\odot\,yr^{-1}$ would be seen to be in the lower range of the SFR of J1649+2635 and other spiral DRAGNs.

5. Discussion & Conclusions

This paper presents the discovery of the spiral DRAGN MCG+07-47-10. The host galaxy has clearly defined spiral arms, a prominent bulge, and hosts a 188 kpc DRAGN.

The radio source, first identified in NVSS, was not previously classified as a DRAGN, but rather three separate radio sources. The deep GMRT observation presented in this paper detects the low surface brightness emission connecting the radio components, thus identifying this radio source as a DRAGN for the first time.

The central component of the radio emission, from the host galaxy, appears extended in both the NVSS and GMRT data. These new GMRT data show resolved radio emission emanating from the entirety of the host galaxy, as opposed to only the core. This suggests that the radio emission is not solely due to the presence of an AGN. Moreover, the total integrated radio flux density for the host galaxy gives a SFR that is in good agreement with SFRs calculated from the IR emission. Resolved radio emission observed throughout the disk suggests that most, if not all the radio emission across the disk is due to star-formation.

We find that the luminosity of this spiral DRAGN ($L_{1.4\,\text{GHz}}$ = 1.12×10^{22} Watts Hz⁻¹) to be significantly lower than other spiral DRAGNS. However, without an reliable redshift this could be misleading. If we take the median redshift from the main galaxy sample from SDSS of z = 0.1 (Strauss et al. 2002), we would calculate a luminosity of $L_{1.4\,\mathrm{GHz}} = 4.25 \times 10^{23}\,\mathrm{Watts\,Hz^{-1}}$. This is still lower than that found in other spiral DRAGNS such as J1649+2635 with $L_{1.4\,\mathrm{GHz}}=1.03\times10^{24}\,\mathrm{Watts}\,\mathrm{Hz}^{-1}$ (Mao et al. 2015). A redshift of 0.1 would imply an optical diameter of 57 kpc for MCG+07-47-10 and the angular linear extent of the entire spiral DRAGN of ~ 9' would equate to a physical size of ≈ 1 Mpc. In order for MCG+07-47-10 to have a comparable luminosity to J1649+2635, MCG+07-47-10 would need to have a redshift of $z \approx 0.17$. This redshift would make the physical size of the DRAGN associated with MCG+07-47-10 significantly larger than 1 Mpc. We note that this would not necessarily be unusual for the class as two previously identified spiral DRAGNS (Hota et al. 2011; Bagchi et al. 2014) show evidence of mega-parsec structure. However, the host galaxy, MCG+07-47-10, would then have an optical size of approximately 100 kpc, which would be exceptionally large compared to similar analogues.

The low-luminosity and low surface brightness of the DRAGN may suggest that the radio emission is old. One possible scenario leading to the formation of this spiral DRAGN is that the host of a low-luminosity DRAGN has had gas injected onto it, perhaps through a merger, and this gas has triggered star-

formation and built up spiral arms. Further modelling and observations at several frequencies are required to test this theory.

Assuming that the host galaxy, MCG+07-47-10, is associated with the two galaxy groups in its immediate vicinity (located within the virial radius of NGC 7618 and UGC 12491 (Kraft et al. 2006)), it would appear to reside in a similarly intermediate density environment to other known spiral DRAGNs (Mao et al. 2015). This supports the idea that spiral DRAGNs require specific environments to form and that a moderately overdense environment is conducive for mergers to trigger a process that transitions ellipticals back to spirals.

In conclusion, we have presented the discovery of a new spiral DRAGN from observations with the GMRT at 322 MHz. We have shown that the host galaxy is a star-forming spiral galaxy. Given currently available information, we have demonstrated that the radio luminosity of this new source is significantly lower than that of other spiral DRAGNs, about two orders of magnitude, with a value of $L_{1.4\,\mathrm{GHz}}=1.12\times10^{22}\,\mathrm{Watts\,Hz^{-1}}$. This may indicate the existence of a previously unknown population of low-luminosity spiral DRAGNs and we suggest that future radio surveys may be used to expand this sample further.

Acknowledgements. DM, AMS & AC gratefully acknowledge support from ERCStG 307215 (LODESTONE). MYM acknowledges support from EC H2020-MSCA-IF-2014 660432 (spiral DRAGNs). We thank the anonymous referee for their helpful comments which improved this paper. We acknowledge the usage of the HyperLeda database (http://leda.univ-lyon1.fr) and NASA/IPAC Extragalactic Database. The Digitized Sky Surveys were produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166. The images of these surveys are based on photographic data obtained using the Oschin Schmidt Telescope on Palomar Mountain and the UK Schmidt Telescope. The plates were processed into the present compressed digital form with the permission of these institutions.

References

Alam, S., Albareti, F. D., Allende Prieto, C., et al. 2015, ApJS, 219, 12
Bagchi, J., Vivek, M., Vikram, V., et al. 2014, ApJ, 788, 174
Becker, R. H., White, R. L., & Helfand, D. J. 1995, ApJ, 450, 559
Bennett, C. L., Larson, D., Weiland, J. L., & Hinshaw, G. 2014, ApJ, 794, 135
Best, P. N., Kauffmann, G., Heckman, T. M., et al. 2005, MNRAS, 362, 25
Chiaberge, M., Gilli, R., Lotz, J. M., & Norman, C. 2015, ApJ, 806, 147
Chiaberge, M. & Marconi, A. 2011, MNRAS, 416, 917
Cohen, A. S., Lane, W. M., Cotton, W. D., et al. 2007, AJ, 134, 1245
Condon, J. J. 1992, ARA&A, 30, 575
Condon, J. J., Cotton, W. D., Greisen, E. W., et al. 1998, AJ, 115, 1693
Gioia, I. M., Gregorini, L., & Klein, U. 1982, A&A, 116, 164
Hopkins, A. M., Miller, C. J., Nichol, R. C., et al. 2003, ApJ, 599, 971
Hopkins, P. F., Hernquist, L., Cox, T. J., & Kereš, D. 2008, ApJS, 175, 356
Hota, A., Sirothia, S. K., Ohyama, Y., et al. 2011, MNRAS, 417, L36
Intema, H. T., Jagannathan, P., Mooley, K. P., & Frail, D. A. 2016, ArXiv e-prints
[arXiv:1603.04368]
Intema, H. T., van der Tol, S., Cotton, W. D., et al. 2009, A&A, 501, 1185
Irwin, J., Beck, R., Benjamin, R. A., et al. 2012, AJ, 144, 43
Jarrett, T. H., Masci, F., Tsai, C. W., et al. 2013, AJ, 145, 6
Kennicutt, Jr., R. C. 1998, ARA&A, 36, 189
Kraft, R. P., Jones, C., Nulsen, P. E. J., & Hardcastle, M. J. 2006, ApJ, 640, 762
Lacy, M., Storrie-Lombardi, L. J., Sajina, A., et al. 2004, ApJS, 154, 166
Lane, W. M., Cotton, W. D., van Velzen, S., et al. 2014, MNRAS, 440, 327
Leahy, J. P. 1993, in Lecture Notes in Physics, Berlin Springer Verlag, Vol. 421,
Jets in Extragalactic Radio Sources, ed. H.-J. Röser & K. Meisenheimer, 1
Ledlow, M. J., Owen, F. N., Yun, M. S., & Hill, J. M. 2001, ApJ, 552, 120
Lintott, C. J., Schawinski, K., Slosar, A., et al. 2008, MNRAS, 389, 1179
Mao, M. Y., Owen, F., Duffin, R., et al. 2015, MNRAS, 446, 4176
Martin, D. C., Fanson, J., Schiminovich, D., et al. 2005, ApJ, 619, L1
Matthews, T. A., Morgan, W. W., & Schmidt, M. 1964, ApJ, 140, 35
Meert, A., Vikram, V., & Bernardi,

Yan, L., Donoso, E., Tsai, C.-W., et al. 2013, AJ, 145, 55