

The IC-indices of Some Complete Multipartite Graphs

Chin-Lin Shiue ^{*} Hui-Chuan Lu [†]

September 18, 2018

Abstract

A coloring of a connected graph G is a function f mapping the vertex set of G into the set of all integers. For any subgraph H of G , we denote the sum of the values of f on the vertices of H as $f(H)$. If for any integer $k \in \{1, 2, \dots, f(G)\}$, there exists an induced connected subgraph H of G such that $f(H) = k$, then the coloring f is called an IC-coloring of G . The IC-index of G , denoted as $M(G)$, is the maximum value of $f(G)$ over all possible IC-colorings f of G . In this paper, we present a useful method from which a lower bound on the IC-index of any complete multipartite graph can be derived. Subsequently, we show that, for $m \geq 2$ and $n \geq 2$, our lower bound on $M(K_{1(n),m})$ is the exact value of it.

Keywords: IC-coloring; IC-index; complete multipartite graph

1 Introduction

Given a connected simple graph G , a *coloring* of G is a function f mapping $V(G)$ into \mathbb{N} . For any subgraph H of G , we denote the sum $\sum_{v \in V(H)} f(v)$ as $f(H)$. If for any integer $k \in \{1, 2, 3, \dots, f(G)\}$, there exists an induced connected subgraph H of G such that $f(H) = k$, then the coloring f is called an *IC-coloring* of G . Every connected graph G admits a trivial IC-coloring which assigns the value 1 to every vertex of G . The highest possible value of $f(G)$ is referred to as the *IC-index* of a graph G , denoted as $M(G)$, that is,

$$M(G) = \max\{f(G) \mid f \text{ is an IC-coloring of } G\}.$$

An IC-coloring f satisfying $f(G) = M(G)$ is called a *maximal IC-coloring* of G . In this paper, we only consider simple graphs. For the terminologies and notations in graph theory, please refer to [13].

The problem of IC-coloring of finite graphs originated from the postage stamp problem in number theory, which has been studied in some literature [1, 2, 3, 4, 5, 6, 8, 9, 10, 12]. In 1992, Glenn Chappel formulated the IC-coloring problem as a "subgraph sums problem" and showed that $M(C_n) \leq n^2 - n + 1$. Later, in 1995, Penrice [7] introduced the IC-coloring as the stamp covering and showed that $M(K_n) = 2^n - 1$ for $n \geq 1$ and $M(K_{1,n}) = 2^n + 2$ for $n \geq 2$. In 2005, Salehi et. al.[8] proved that $M(K_{2,n}) = 3 \cdot 2^n + 1$ for $n \geq 2$. Along with the result by Shiue and Fu [11] who showed that $M(K_{m,n}) = 3 \cdot 2^{m+n-2} - 2^{m-2} + 2$,

^{*}Department of Applied Mathematics, Chung Yuan Christian University, Chung Li, Taiwan 32023. Research supported in part by NSC-99-2115-M033-001.

[†]**Corresponding author;** Center for Basic Required Courses, National United University, Maioli, Taiwan 36003. E-mail:hjlu@nuu.edu.tw. Research supported in part by NSC-102-2115-M-239-002.

for $2 \leq m \leq n$, in 2008, the problem regarding complete bipartite graphs was completely settled. In this present paper, we deal with *complete multipartite graphs*. A complete multipartite graph K_{m_1, m_2, \dots, m_k} is a graph whose vertex set can be partitioned into k partite sets V_1, V_2, \dots, V_k , where $|V_i| = m_i$ for all $i \in \{1, 2, \dots, k\}$, such that there are no edges within each V_i and any two vertices from different partite sets are adjacent. A complete multipartite graph with k partite sets is called a *complete k -partite graph* as well. We also denote as $K_{1(n), m_{n+1}, m_{n+2}, \dots, m_k}$, $n \leq k$, the complete k -partite graphs in which there are n partite sets which are of size one and the rest $(k - n)$ partite sets have sizes m_{n+1}, m_{n+2}, \dots , and m_k . Therefore $K_{1(n)}$ represents the complete graph K_n .

In this paper, we introduce some useful lemmas in Section 2. In Section 3, our main results are presented. We start with a useful proposition which gives a lower bound on the IC-index of the join of an independent set and a given connected graph. Consequently, a lower bound on the IC-index of any complete mutlipartite graph can be deduced. We shall show that our lower bound on $M(K_{1(n), m})$ is indeed the exact value of it for $m \geq 2$ and $n \geq 2$. A concluding remark will be given in Section 4.

2 Preliminaries

Some basic know results from [11] are introduced in this section. They are very useful in the discussion of our main results. For brevity, we let $[1, \ell]$, $\ell \in \mathbb{N}$, denote the set $\{1, 2, \dots, \ell\}$. A sequence consisting of 0's and 1's is called a *binary sequence*.

Lemma 2.1. [11] *If a_1, a_2, \dots, a_n are n positive integers which satisfy that $a_1 = 1$ and $a_i \leq a_{i+1} \leq \sum_{j=1}^i a_j + 1$ for all $i \in [1, n - 1]$. Then, for each $\ell \in [1, \sum_{j=1}^n a_j]$, there exists a binary sequence c_1, c_2, \dots, c_n such that $\ell = \sum_{j=1}^n c_j a_j$.*

Lemma 2.2. [11] *If s_0, s_1, \dots, s_n is a sequence of integers, then for each $i \in [1, n]$ there exists an integer $r_i \in \mathbb{Z}$ such that $s_i = \sum_{j=0}^{i-1} s_j + r_i$ and $\sum_{j=0}^n s_j = 2^n s_0 + \sum_{j=1}^n 2^{n-j} r_j$.*

Lemma 2.3. [11] *Let $V(G) = \{u_1, u_2, \dots, u_n\}$. If f is an IC-coloring of G such that $f(u_i) \leq f(u_{i+1})$ for all $i \in [1, n - 1]$, then $f(u_1) = 1$ and $f(u_{i+1}) \leq \sum_{j=1}^i f(u_j) + 1$ for all $i \in [1, n - 1]$.*

Lemma 2.4. [11] *Let f be an IC-coloring of a graph G such that $f(u_i) < f(u_{i+1})$ for $i \in [1, n - 1]$, where $V(G) = \{u_1, u_2, \dots, u_n\}$. For each pair (i_1, i_2) where $1 \leq i_1 < i_2 \leq n$, if $f(u_{i_1}) = \sum_{j=1}^{i_1-1} f(u_j) + 1$ and $u_{i_1} u_{i_2} \notin E(G)$, then either $f(u_{i_2}) \leq \sum_{j=1}^{i_2-1} f(u_j) - f(u_{i_1})$ or $f(u_{i_2+1}) \leq f(u_{i_1}) + f(u_{i_2})$.*

Lemma 2.5. [11] *Let r_1, r_2, \dots, r_n be n numbers. If there are two integers i and k such that $1 \leq i < k \leq n$ and $r_i < r_k$, then*

$$\sum_{j=1}^n 2^{n-j} r_j < \sum_{j=1}^n 2^{n-j} r_j - (2^{n-i} r_i + 2^{n-k} r_k) + (2^{n-i} r_k + 2^{n-k} r_i).$$

Lemma 2.6. [11] *Let f be an IC-coloring of a graph G . If G has ℓ induced connected subgraphs and there are $2k$ distinct induced connected subgraphs $H_1, G_1, H_2, G_2, \dots, H_k, G_k$ of G such that $f(H_i) = f(G_i)$ for all $i \in [1, k]$, then $f(G) \leq \ell - k$.*

3 Main Result

We start this section with a useful method for deriving a meaningful lower bound on the IC-index of the join of an independent set and a given graph. Subsequently, we show that the lower bound on $M(K_{1(n),m})$ derived from our method, for $m \geq 2$ and $n \geq 2$, also serves as an upper bound on it. This determines the exact value of $M(K_{1(n),m})$.

3.1 Lower Bounds on the IC-indices of Complete Multipartite Graphs

For the derivation of lower bounds, we view complete multipartite graphs as being generated by a graph operation starting with graphs with some vertices and no edges. The *join* of two disjoint graphs H_0 and H_1 , written $H_0 \vee H_1$, is the graph with vertex set $V = V(H_0) \cup V(H_1)$ and edge set

$$E = E(H_0) \cup E(H_1) \cup \{(u, v) \mid u \in V(H_0), v \in V(H_1)\}.$$

Let O_m be the graph with m vertices and no edges, then the join of O_1 and K_n , or $O_1 \vee K_n$, is the complete graph K_{n+1} and the graph $O_m \vee K_n$ is exactly the complete multipartite graph $K_{1(n),m}$. Observe that the join of O_m and O_n is the complete bipartite graph $K_{m,n}$. The join of O_m and K_{n_1, n_2} forms the complete tripartite graph $K_{n_1, n_2, m}$. Since joining O_m with a complete $(k-1)$ -partite graph generates a complete k -partite graph, we are concerned about how the value of the IC-index of a graph changes as we joining O_m to that graph.

Proposition 3.1. *If g is an IC-coloring of a connected graph G , then there exists an IC-coloring f of $O_m \vee G$ such that $f(O_m \vee G) = 2^m g(G) + 1$ for $m \geq 1$.*

Proof. Let $V_0 = V(O_m) = \{w_1, w_2, \dots, w_m\}$ and $V_1 = V(G) = \{v_1, v_2, \dots, v_n\}$. We define f on $V(O_m \vee G)$ as $f(v_i) = g(v_i)$ for $i \in [1, n]$, $f(w_j) = 2^{j-1}g(G)$ for $j \in [1, m-1]$ and $f(w_m) = 2^{m-1}g(G) + 1$. The value of $f(O_m \vee G)$ can be calculated as follows.

$$\begin{aligned} f(G) &= g(G) + \sum_{j=1}^m 2^{j-1}g(G) + 1 \\ &= g(G) + (2^m - 1)g(G) + 1 \\ &= 2^m g(G) + 1. \end{aligned}$$

Given any integer $k \in [1, 2^m g(G) + 1]$, we need to identify an induced connected subgraph H of $O_m \vee G$ such that $f(H) = k$. Since g is an IC-coloring of G and $f(v) = g(v)$ for each vertex in G , the desired induced connected subgraph exists for each $k \in [1, g(G)]$. For $k \in [g(G) + 1, 2^{m-1}g(G)]$, we rewrite k into the form $k = qg(G) + r$ where $1 \leq r \leq g(G)$. Hence, $1 \leq q \leq 2^{m-1} - 1$ and then there exists a binary sequence c_1, c_2, \dots, c_{m-1} , which are not all zero, such that $q = \sum_{i=1}^{m-1} c_i 2^{i-1}$. It follows that $qg(G) = \sum_{i=1}^{m-1} c_i 2^{i-1}g(G) = \sum_{i=1}^{m-1} c_i f(w_i)$. Now, let H' be a connected subgraph of G such that $f(H') = r$ and let $W = V(H') \cup \{w_i \mid c_i = 1, i \in [1, m-1]\}$. Since H' is connected, the subgraph H induced by W is also connected and satisfies $f(H) = k$. Next, if $k = 2^{m-1}g(G) + 1$, then the subgraph H induced by the single vertex w_m fits our need. Finally, for $k \in [2^{m-1}g(G) + 2, 2^m g(G) + 1]$, we write k as $k = (2^{m-1}g(G) + 1) + k'$, where $1 \leq k' \leq 2^{m-1}g(G)$. By the above argument, there is a connected subgraph H' such that $f(H') = k'$ and $V(H') \cap V_1 \neq \emptyset$. Let $W = V(H') \cup \{w_m\}$, then the subgraph H induced by W is connected and we have $f(H) = k$. The result follows. \square

Using the known result $M(K_n) = 2^n - 1$ for $n \geq 1$ [7], lower bounds on the IC-indices of $K_{1(n),m}$ and $K_{1(n),m_1,m_2,\dots,m_\ell}$ can be easily obtained by Proposition 3.1.

Corollary 3.2. $M(K_{1(n),m}) = M(O_m \vee K_n) \geq 2^{m+n} - 2^m + 1$ for $m \geq 1$ and $n \geq 1$.

For $m = 1$, this lower bound matches the known value of the IC-index of K_{n+1} . We will show that this inequality is in fact an equality when $m \geq 2$ and $n \geq 2$.

Corollary 3.3. For $m_1 \geq m_2 \geq \dots \geq m_\ell$,

$$\begin{aligned} M(K_{1(n),m_\ell,m_{\ell-1},\dots,m_2,m_1}) &= M(O_{m_1} \vee O_{m_2} \vee \dots \vee O_{m_\ell} \vee K_n) \\ &\geq (2^{m_1}(2^{m_2}(\dots(2^{m_\ell}(2^n - 1) + 1)\dots) + 1) + 1). \end{aligned}$$

Furthermore, since the IC-index of a bipartite graph is known, a lower bound on $M(K_{m_1,m_2,m_3})$ can be derived as well. By successively applying Proposition 3.1, a lower bound on the IC-index of any complete multipartite graph can be easily found.

3.2 The Exact Value of the IC-index of $K_{1(n),m}$

Now, we consider the graph $K_{1(n),m}$, or $O_m \vee K_n$. We shall show that the IC-index of this graph is $2^{m+n} - 2^m + 1$. In the remainder of this paper, we let $G = K_{1(n),m} = O_m \vee K_n$, $V_0 = V(O_m)$ and $V_1 = V(K_n)$. We introduce some properties of the graph $K_{1(n),m}$ and any maximal IC-coloring of it first.

Lemma 3.4. $K_{1(n),m}$ has $(2^{m+n} - 2^m + m)$ induced connected subgraphs.

Proof. Any induced connected subgraph H of G must satisfy exactly one of the following three conditions: (i) $V(H) \subseteq V_1$ and $V(H) \neq \emptyset$; (ii) $V(H) \subseteq V_0$ and $|V(H)| = 1$; (iii) $V(H) \cap V_1 \neq \emptyset$ and $V(H) \cap V_0 \neq \emptyset$. Therefore, the number of induced connected graphs of G is $(2^n - 1) + m + (2^m - 1)(2^n - 1) = 2^{m+n} - 2^m + m$. \square

Proposition 3.5. If f is a maximal IC-coloring of $K_{1(n),m}$, then $f(u) \neq f(v)$ for each pair of distinct vertices u and v in $V(K_{1(n),m})$.

Proof. Suppose that there are two distinct vertices u and v in $V(G)$ such that $f(u) = f(v)$. For subsets of vertices $V'_0 \subseteq V_0 \setminus \{u, v\}$ and $V'_1 \subseteq V_1 \setminus \{u, v\}$, we denote as H_u and H_v the subgraphs of G induced by $V'_0 \cup V'_1 \cup \{u\}$ and $V'_0 \cup V'_1 \cup \{v\}$ respectively. Then, $H_u \neq H_v$ and $f(H_u) = f(H_v)$. Let S be the set of all possible pairs (V'_0, V'_1) such that H_u and H_v are both connected, and let $p = |S|$. One can see from Lemma 2.6 and Lemma 3.4 that $f(G) \leq (2^{m+n} - 2^m + m) - p$.

Observe that any connected subgraph H of G must satisfy exactly one of the two conditions: (i) $V(H) \cap V_1 \neq \emptyset$, or (ii) $V(H) \cap V_1 = \emptyset$ and $|V(H) \cap V_0| = 1$. Base on this fact, we now evaluate the number p in the following three possible cases.

Case (a): if $u, v \in V_1$, then $p = 2^{|V_1 \setminus \{u,v\}|} \cdot 2^{|V_0|} = 2^{m+n-2}$.

Case (b): if $u, v \in V_0$, then either $V'_1 \neq \emptyset$ or $V'_0 = V'_1 = \emptyset$. Thus we have $p = 2^{|V_0 \setminus \{u,v\}|} \cdot (2^{|V_1|} - 1) + 1 = 2^{m-2}(2^n - 1) + 1 = 2^{m+n-2} - 2^{m-2} + 1$.

Case (c): if exactly one of u and v is in V_0 , then either $V'_1 \neq \emptyset$ or $V'_0 = V'_1 = \emptyset$. It follows that $p = 2^{m-1}(2^{n-1} - 1) + 1 = 2^{m+n-2} - 2^{m-1} + 1$.

The value of p in Case (c) is the minimum among these three cases. This leads to an upper bound on $f(G)$ as follows.

$$\begin{aligned} f(G) &\leq (2^{m+n} - 2^m + m) - p \\ &\leq (2^{m+n} - 2^m + m) - (2^{m+n-2} - 2^{m-1} + 1) \\ &= 2^{m+n} - 2^{m+n-2} - 2^{m-1} + m - 1. \end{aligned}$$

Since $m, n \geq 2$, we have $2^{m+n-2} \geq 2^m$ and $-2^{m-1} + m - 1 < 0$. This implies that $f(G) < 2^{m+n} - 2^m + 1$ which is a contradiction to Corollary 3.2 and we have the result. \square

Now we are in a position to prove that the lower bound given in Corollary 3.2 also serves as an upper bound on $M(K_{1(n),m})$. However, the proof is too involved that we need some more notations to facilitate the whole discussion process. In what follows, for the given maximal IC-coloring f of G , we always assume that $\{u_1, u_2, \dots, u_{m+n}\}$ is the vertex set of G such that $f(u_i) < f(u_{i+1})$ for all $i \in [1, m+n-1]$. Thus, $f(u_1) = 1$ and $f(u_2) = 2$ are always true. Besides, we also define $f_0 = 0$ and denote the sum $\sum_{j=1}^i f(u_j)$ as f_i for $i \in [1, m+n]$. Let us introduce some useful facts first.

Lemma 3.6. *If f is a maximal IC-coloring of $K_{1(n),m}$, then*

- (1) $f_j \leq 2^{j-i}(f_i + 1) - 1$ for every pair (i, j) such that $1 \leq i < j \leq m+n$.
- (2) $f_i \geq 3 \cdot 2^{i-2}$ for each $i \in [2, m+n]$.

Proof. (1) Given $i \in [1, m+n-1]$, we let $s_0 = f_i$ and $s_k = f(u_{i+k})$ for each $k \in [1, m+n-i]$. By Lemma 2.3, we have $s_k \leq f_{i+k-1} + 1 = \sum_{\ell=0}^{k-1} s_\ell + 1$. For each $j \in [i+1, m+n]$, one can see from Lemma 2.2 that

$$\begin{aligned} f_j &= \sum_{\ell=0}^{j-i} s_\ell \\ &\leq 2^{j-i} f_i + \sum_{\ell=1}^{j-i} 2^{j-i-\ell} \cdot 1 \\ &= 2^{j-i}(f_i + 1) - 1. \end{aligned}$$

(2) Suppose that there exists some $i \in [2, m+n]$ such that $f_i \leq 3 \cdot 2^{i-2} - 1$. According to part(1), we have

$$\begin{aligned} f(G) &= f_{m+n} \\ &\leq 2^{m+n-i}(f_i + 1) - 1 \\ &\leq 2^{m+n-i}(3 \cdot 2^{i-2}) - 1 \\ &= 2^{m+n} - 2^{m+n-2} - 1 \\ &< 2^{m+n} - 2^m + 1. \end{aligned}$$

This contradicts to Corollary 3.2 and we have the result. \square

Lemma 3.7. *Suppose that f is a maximal IC-coloring of $K_{1(n),m}$. Let $s_0 = f_0 = 0$ and $s_i = f(u_i)$ for $i \in [1, m+n]$. If each r_i is the integer such that $s_i = \sum_{j=0}^{i-1} s_j + r_i$, $i \in [1, m+n]$, then $r_i \leq 1$. Furthermore, if $r_i \leq 0$ for all $i \in \{j \mid u_j \in V_0 \setminus \{u_{k_0}\}\}$, where $k_0 = \max \{j \mid u_j \in V_0\}$, then $f(K_{1(n),m}) \leq 2^{m+n} - 2^m + 1$.*

Proof. The first result is trivial from Lemma 2.3. To prove that second result, we describe the IC-coloring \bar{f} defined in the proof of Proposition 3.1 for $O_m \vee K_n$ explicitly. Let $V_0 = V(O_m) = \{w_1, w_2, \dots, w_m\}$ and $V_1 = V(K_n) = \{v_1, v_2, \dots, v_n\}$. Since we are given a maximal IC-coloring g of K_n defined as $g(v_i) = 2^{i-1}$ for each $v_i \in V_1$ [7], we have an IC-coloring \bar{f} of $O_m \vee K_n$ defined as $\bar{f}(v_i) = g(v_i) = 2^{i-1}$ for $i \in [1, n]$, $\bar{f}(w_j) = 2^{j-1}(2^n - 1)$ for $j \in [1, m-1]$ and $\bar{f}(w_m) = 2^{m-1}(2^n - 1) + 1$. Then $\bar{f}(O_m \vee K_n) = 2^{m+n} - 2^m + 1$. Let us

rearrange the vertices of G into a new order $\{\bar{u}_1, \bar{u}_2, \dots, \bar{u}_{m+n}\}$ such that $\bar{f}(\bar{u}_i) < \bar{f}(\bar{u}_{i+1})$ for all $i \in [1, m+n-1]$. Then $\bar{f}(\bar{u}_i) = \sum_{j=1}^{i-1} \bar{f}(\bar{u}_j) + \bar{r}_i$, where $\bar{r}_i = 1$ for $i \in [1, n] \cup \{n+m\}$, and $\bar{r}_i = 0$ for $i \in [n+1, n+m-1]$. According to the assumption in this Lemma, $r_i \leq 0$ for all $i \in \{j \mid u_j \in V_0 \setminus \{u_{k_0}\}\}$. If $r_{m+n} = 1$, then among r_1, r_2, \dots, r_{m+n} there are at most $(n+1)$ of them taking the value one and at least $(m-1)$ of them having the values no more than zero. By comparing the distributions of the 1's in \bar{r}_i 's and r_i 's, we conclude from Lemma 2.5 that $f(G)$ does not exceed the value $\bar{f}(G) = 2^{m+n} - 2^m + 1$. If $r_{m+n} = 0$, then there are at most n r_i 's being one and at least m r_i 's not exceeding zero. Lemma 2.5 again guarantees the truth of the inequality $f(G) \leq \bar{f}(G)$ in this case. \square

Next, we consider the case where the assumption in Lemma 3.7 is violated, namely, there is a r_i having the value one for some $i \in \{j \mid u_j \in V_0 \setminus \{u_{k_0}\}\}$.

Lemma 3.8. *Let f be a maximal IC-coloring of $K_{1(n),m}$. Let $s_0 = f_0 = 0$, $s_i = f(u_i)$ and $r_i = s_i - \sum_{j=0}^{i-1} s_j$ for $i \in [1, m+n]$. Suppose that $S_1 = \{i < k_0 \mid r_i = 1 \text{ and } u_i \in V_0\} \neq \emptyset$, where $k_0 = \max \{j \mid u_j \in V_0\}$. Let $i_1 = \min S_1$, $t = |\{u_i \in V_0 \mid i < i_1\}|$ and $S_2 = \{i \geq i_1 + 1 \mid u_i \in V_0 \text{ and } f(u_i) > f_{i-1} - f(u_{i_1})\}$.*

- (i) *If $S_2 = \emptyset$, then $f(K_{1(n),m}) \leq 2^{m+n} - 2^m + 1$.*
- (ii) *If $S_2 \neq \emptyset$, then $f(K_{1(n),m}) \leq 2^{m+n} - 2^{m+n-2}(1 - 2^{-(i_2-i_1)}) - (2^t - 1)(3 \cdot 2^{m+n-i_1-1} + 2^{m+n-i_2-1})$, where $i_2 = \min S_2$.*
- (iii) *If $S_2 \neq \emptyset$ and $n \geq 3$, then $f(K_{1(n),m}) \leq 2^{m+n} - 2^m + 1$.*

Proof. Observe that, among $r_1, r_2, \dots, r_{i_1-1}$, there are at most $(i_1 - t - 1)$ of them having the value one. By Lemma 2.2 and Lemma 2.5 we have that

$$\begin{aligned} f_{i_1-1} &= 2^{i_1-1} s_0 + \sum_{j=1}^{i_1-1} 2^{i_1-1-j} r_j \\ &\leq 2^{i_1-1} \cdot 0 + \sum_{j=1}^{i_1-t-1} 2^{i_1-1-j} \cdot 1 \\ &= 2^{i_1-1} - 2^t \dots\dots\dots (1) \end{aligned}$$

and

$$\begin{aligned} f_{i_1} &= f_{i_1-1} + f(u_{i_1}) \\ &= 2f_{i_1-1} + 1 \\ &\leq 2^{i_1} - 2^{t+1} + 1 \dots\dots\dots (2) \end{aligned}$$

(i) Consider the case $s'_0 = f_{i_1}$ and $s'_j = f(u_{i_1+j})$ for $j \in [1, m+n-i_1]$. Let $r'_j = s'_j - \sum_{\ell=0}^{j-1} s'_\ell$, then $r'_j \leq 1$. Furthermore, if $S_2 = \emptyset$, that is, $f(u_{i_1+j}) \leq f_{i_1+j-1} - f(u_{i_1})$ for all $u_{i_1+j} \in V_0$, then $f(u_{i_1+j}) - \sum_{i=1}^{i_1+j-1} f(u_i) \leq -f(u_{i_1}) \leq -1$. Since $r'_j \leq -1$ for all $j \in \{j \leq m+n-i_1 \mid u_{i_1+j} \in V_0\}$, there are at most $(n - (i_1 - t - 1))$ 1's in the values of $r'_1, r'_2, \dots, r'_{m+n-i_1}$. By Lemma 2.2, 2.5 and Inequality (2), we have

$$\begin{aligned} f(G) &= \sum_{j=0}^{m+n-i_1} s'_j \\ &= 2^{m+n-i_1} f_{i_1} + \sum_{j=1}^{m+n-i_1} 2^{m+n-i_1-j} r'_j \\ &\leq 2^{m+n-i_1} (2^{i_1} - 2^{t+1} + 1) + \sum_{j=1}^{n-(i_1-t-1)} 2^{m+n-i_1-j} \cdot 1 + \\ &\quad \sum_{j=n-(i_1-t-1)+1}^{m+n-i_1} 2^{m+n-i_1-j} (-1) \\ &\leq 2^{m+n-i_1} (2^{i_1} - 2^{t+1} + 1) + 2^{m+n-i_1} - 2^{m-t-1} - (2^{m-t-1} - 1) \\ &\leq 2^{m+n} + 2^{m+n-i_1+1} - 2^{m+n-i_1+t+1} - 2^{m-t} + 1. \end{aligned}$$

Note that $n \geq i_1 - t - 1 \geq 0$ and $t \geq 0$. If $n = i_1 - t - 1$ or $t = 0$, then $f(G) \leq 2^{m+n} - 2^m + 1$.

Otherwise, $n \geq i_1 - t$ and $t \geq 1$, then $2^{m+n-i_1+1} - 2^{m+n-i_1+t+1} \leq 2^{m+n-i_1+t} - 2^{m+n-i_1+t+1} = -2^{m+n-i_1+t} \leq -2^m$. This implies that $f(G) \leq 2^{m+n} - 2^m - 2^{m-t} + 1 < 2^{m+n} - 2^m + 1$ and the result in (i) is asserted.

(ii) If $S_2 \neq \emptyset$, then $f(u_{i_2}) > f_{i_2-1} - f(u_{i_1})$. By Lemma 2.4 we have that $f(u_{i_2+1}) \leq f(u_{i_1}) + f(u_{i_2})$ and then

$$\begin{aligned} f_{i_2+1} &= f_{i_2-1} + f(u_{i_2}) + f(u_{i_2+1}) \\ &\leq f_{i_2-1} + 2f(u_{i_2}) + f(u_{i_1}) \\ &\leq f_{i_2-1} + 2(f_{i_2-1} + 1) + (f_{i_1-1} + 1) \\ &= 3f_{i_2-1} + f_{i_1-1} + 3 \dots \dots \dots (3) \end{aligned}$$

Now, the desired upper bound on $f(G)$ can be derived as follows.

$$\begin{aligned} f(G) &= f_{m+n} \leq 2^{m+n-(i_2+1)}(f_{i_2+1} + 1) && \text{(by Lemma 3.6(1))} \\ &\leq 2^{m+n-(i_2+1)}[(3f_{i_2-1} + f_{i_1-1} + 3) + 1] && \text{(by Inequality (3))} \\ &\leq 2^{m+n-(i_2+1)}[3(2^{i_2-i_1}(f_{i_1-1} + 1) - 1) + f_{i_1-1} + 4] && \text{(by Lemma 3.6(1))} \\ &= 2^{m+n-(i_2+1)}[(3 \cdot 2^{i_2-i_1} + 1)(f_{i_1-1} + 1)] \\ &= 2^{m+n-(i_2+1)}(3 \cdot 2^{i_2-i_1} + 1)(2^{i_1-1} - (2^t - 1)) && \text{(by Inequality (1))} \\ &= 3 \cdot 2^{m+n-2} + 2^{m+n-(i_2-i_1)-2} - (2^t - 1)(3 \cdot 2^{m+n-i_1-1} + 2^{m+n-i_2-1}) \\ &= 2^{m+n} - 2^{m+n-2}(1 - 2^{-(i_2-i_1)}) - (2^t - 1)(3 \cdot 2^{m+n-i_1-1} + 2^{m+n-i_2-1}) \dots \dots \dots (4) \end{aligned}$$

(iii) Since $i_2 - i_1 \geq 1$, $2^{m+n-2}(1 - 2^{-(i_2-i_1)}) \geq 2^{m+n-3}$. It follows that $f(G) \leq 2^{m+n} - 2^{m+n-3} < 2^{m+n} - 2^m + 1$ when $n \geq 3$. The proof is completed. \square

We have shown that $f(G) \leq 2^{m+n} - 2^m + 1$ is valid in many cases. With some further discussion, the final conclusion can be achieved.

Theorem 3.9. $M(K_{1(n),m}) = 2^{m+n} - 2^m + 1$.

Proof. Let f be a maximal IC-coloring of G . We adopt the notation used in Lemma 3.8. By Lemma 3.7 and Lemma 3.8, it suffices to show that $f(G) \leq 2^{m+n} - 2^m + 1$ holds when $S_2 \neq \emptyset$ and $n = 2$. First note that when $n = 2$, we have $0 \leq i_1 - t - 1 \leq 2$ and the upper bound on $f(G)$ in (4) can be rewritten as

$$\begin{aligned} f(G) &\leq 2^{m+2} - 2^m(1 - 2^{-(i_2-i_1)}) - (2^t - 1)(3 \cdot 2^{m-i_1+1} + 2^{m-i_2+1}) \\ &= (2^{m+2} - 2^m) + 2^{m-(i_2-i_1)} - 3 \cdot 2^{m-i_1+t+1}(1 - 2^{-t}) + 2^{m-i_2+1}(2^t - 1) \\ &= 2^{m+2} - 2^m - 3 \cdot 2^{m-(i_1-t-1)}(1 - 2^{-t}) + 2^{m-(i_2-i_1)}(1 - 2^{-i_1+1})(2^t - 1) \dots \dots \dots (5) \end{aligned}$$

(1) If $t \geq 2$, then

$$\begin{aligned} f(G) &\leq (2^{m+2} - 2^m) - 3 \cdot 2^{m-2}(1 - \frac{1}{4}) + 2^{m-1} \\ &< 2^{m+2} - 2^m + 1. \end{aligned}$$

(2) If $t = 1$ and $i_1 \leq 3$, then

$$\begin{aligned} f(G) &\leq (2^{m+2} - 2^m) - 3 \cdot 2^{m-1}(1 - 2^{-1}) + 2^{m-1}(1 - 2^{-2}(2 - 1)) \\ &= (2^{m+2} - 2^m) + (3/4 - 3/2) \cdot 2^{m-1} \\ &< 2^{m+2} - 2^m + 1. \end{aligned}$$

(3) We have so far consider all but the following four cases: $(i_1, t) = (1, 0), (2, 0), (3, 0), (4, 1)$.

Let us have a closer investigation of the value $f(G)$ again before literally starting the discussion of these cases. Now, consider the the situation where $s_0 = f_0$, $s_i = f(u_i)$ and $s_i = \sum_{j=0}^{i-1} s_j + r_i$ for $i \in [1, i_2 - 1]$. From the definition of i_1 and i_2 , we see that $r_i \leq 0$ for all $i \in \{i \leq i_2 - 1 \mid u_i \in V_0\} \setminus \{i_1\}$. Since $n = 2$, among $r_1, r_2, \dots, r_{i_2-1}$, there are at most three of them taking the value one. We therefore have

$$\begin{aligned} f_{i_2-1} &\leq 2^{i_2-1} s_0 + \sum_{j=1}^{i_2-1} 2^{i_2-1-j} \cdot r_j \\ &\leq 2^{i_2-2} + 2^{i_2-3} + 2^{i_2-4} \\ &= 7 \cdot 2^{i_2-4}. \end{aligned}$$

Making use of this fact and Inequality (1), an upper bound on f_{i_2+1} can be derived from Inequality (3) as follows.

$$\begin{aligned} f_{i_2+1} &\leq 3f_{i_2-1} + f_{i_1-1} + 3 \\ &\leq 21 \cdot 2^{i_2-4} + 2^{i_1-1} - 2^t + 3. \end{aligned}$$

This leads to an upper bound on $f(G)$.

$$\begin{aligned} f(G) &< 2^{m+2-(i_2+1)}(f_{i_2+1} + 1) && \text{(by Lemma 3.6(1))} \\ &= 2^{m+2-(i_2+1)}(21 \cdot 2^{i_2-4} + 2^{i_1-1} - 2^t + 4) \\ &= (21 \cdot 2^{m-3}) + [2^{m-(i_2-i_1)} - 2^{m+1-i_2+t} + 2^{m+3-i_2}] \\ &= (2^{m+2} - 2^m - 3 \cdot 2^{m-3}) + [2^{m-(i_2-i_1)}(1 - 2^{-i_1+t+1} + 2^{-i_1+3})] \dots \dots \dots (6) \end{aligned}$$

Now, let us have the discussion for the remaining four cases.

Case 1. $(i_1, t) = (1, 0)$.

In this case, $f(u_{i_1}) = f(u_1) = 1$. If $i_2 \geq 5$, then Inequality (6) gives

$$\begin{aligned} f(G) &< (2^{m+2} - 2^m) - 3 \cdot 2^{m-3} + 2^{m-4}(1 - 2^0 + 2^2) \\ &< 2^{m+2} - 2^m + 1. \end{aligned}$$

When $i_2 = 4$, since $f(u_2) = 2 > f_1 - f(u_1)$ and $f(u_3) > f(u_2) = f_2 - f(u_1)$, we have $\{u_2, u_3\} \subseteq V_1$ by the definition of i_2 . So, $V_1 = \{u_2, u_3\}$ and $V_0 = \{u_i \mid i \in \{1\} \cup [4, m+n]\}$. Now, $f(u_3) \leq f_2 + 1 = 4$ and $f_3 \leq 7$. Since $i_2 = 4$, we know that $f(u_4) > f_3 - f(u_1)$ holds. Lemma 2.4 guarantees that $m+n \geq 5$ and $f(u_5) \leq f(u_1) + f(u_4) \leq 9$, giving $f_5 = f_3 + f(u_4) + f(u_5) \leq f_3 + (f_3 + 1) + f(u_5) \leq 7 + 8 + 9 \leq 24$. Now, suppose that $m+n \geq 6$, then the fact $f_6 \geq 3 \cdot 2^{6-2} = 48$ from Lemma 3.6(2) implies that $f(u_6) \geq 48 - f_5 \geq 24$, which means $f(u_6) > f_5 - f(u_1)$. One can then obtain from Lemma 2.4 that $m+n \geq 7$ and $f(u_7) \leq f(u_1) + f(u_6) \leq 1 + (f_5 + 1) = 26$. However, this leads to $f_7 = f_5 + f(u_6) + f(u_7) \leq 24 + 25 + 26 = 75 < 3 \cdot 2^{7-2}$, contradicting to Lemma 3.6(2). We therefore conclude that the only possible situation is " $m+n = 5$ " and $f(G) = f_5 \leq 24 < 2^{m+2} - 2^m + 1$.

When $i_2 = 3$, from the definition of i_2 , we have $u_2 \in V_1$ and also $f(u_4) = f(u_{i_2+1}) \leq f(u_{i_1}) + f(u_{i_2}) = f(u_1) + f(u_3)$ by Lemma 2.4. Now, $2 = f(u_2) < f(u_3) \leq f_2 + 1 = 4$. From Lemma 3.6(2), we see that $f_4 \geq 3 \cdot 2^{4-2} = 12$ and thus $12 \leq f_4 = f_2 + f(u_3) + f(u_4) \leq 3 + f(u_3) + (f(u_1) + f(u_3)) = 2f(u_3) + 4$ which implies that $f(u_3) = 4$. It follows that $f_3 = 7$ and $f(u_4) \leq f(u_1) + f(u_3) = 5$ by Lemma 2.4 because $f(u_3) > f_2 - f(u_1)$ and $u_1 u_3 \notin E(G)$. Since $f(u_4) > f(u_3) = 4$, we see that $f(u_4) = 5$ and $f_4 = 12$. Suppose that $m+n \geq 5$, then Lemma 3.6(2) gives $f_5 \geq 3 \cdot 2^{5-2} = 24$. One can see that $f(u_5) = f_5 - f_4 \geq 24 - 12 = 12 > 9 = f(u_4) + f(u_3)$ and $f(u_4) > f_3 - f(u_3)$. Lemma 2.4 then guarantees that u_4 must be in V_1 . We therefore conclude that $V_1 = \{u_2, u_4\}$ and

$V_0 = \{u_i \mid i \in \{1, 3\} \cup [5, m+n]\}$. Next, since $f(u_5) \geq 12 > f_4 - f(u_1)$, we have from Lemma 2.4 that $m+n \geq 6$ and $f(u_6) \leq f(u_1) + f(u_5) \leq 1 + (f_4 + 1) \leq 14$. However, this leads to $f_6 \leq f_4 + f(u_5) + f(u_6) \leq 12 + (12 + 1) + 14 = 39 < 3 \cdot 2^{6-2}$ which is a contradiction to Lemma 3.6(2). Therefore the only possible situation is " $m+n = 4$ " and then $f(G) = f_4 = 12 \leq 2^{m+2} - 2^m + 1$.

If $i_2 = 2$, then $f(u_{i_1}) = f(u_1) = 1$ and $f(u_{i_2}) = f(u_2) = 2 = f_1 + 1$. Since $f(u_3) = f(u_{i_2+1}) \leq f(u_{i_1}) + f(u_{i_2}) = 3$ and $f(u_3) > f(u_2) = 2$, we have $f(u_3) = 3$ and then $f(u_3) > f_2 - f(u_1)$. In addition, the fact $f_4 \geq 3 \cdot 2^{4-2} = 12$ implies that $f(u_4) = f_4 - f_3 \geq 12 - 6 = 6 > 4 = f(u_3) + f(u_1)$. From Lemma 2.4, we see that u_3 must be in V_1 . When $m = 2$, $f(G) = f_4 \leq 1 + 2 + 3 + (f_3 + 1) = 13 = 2^{m+2} - 2^m + 1$. Next, we consider the situation when $m \geq 3$. Note that the inequality $f_5 \geq 3 \cdot 2^{5-2} = 24$ gives $f(u_5) = f_5 - f_4 \geq 24 - 13 = 11 > 8 \geq f(u_1) + f(u_4)$. Along with the fact that $f(u_4) \geq 6 > 5 = f_3 - f(u_1)$, one can see that u_4 must also be in V_1 from Lemma 2.4. Therefore, we have $V_1 = \{u_3, u_4\}$ and $V_0 = \{u_i \mid i \in \{1, 2\} \cup [5, m+n]\}$. Suppose that $f(u_5) \geq 12$, then $f(u_5) > 11 \geq f_4 - f(u_2)$. Lemma 2.4 guarantees that $m+n \geq 6$ and $f(u_6) \leq f(u_2) + f(u_5) \leq 2 + (f_4 + 1) \leq 16$. However, these facts lead to $f_6 = f_4 + f(u_5) + f(u_6) \leq 13 + 14 + 16 = 43 < 3 \cdot 2^{6-2}$, contradicting to Lemma 3.6(2). Hence, $f(u_5) = 11$ is asserted. Now, suppose again that $m+n \geq 6$. The inequality $f_6 \geq 3 \cdot 2^{6-2} = 48$ from Lemma 3.6(2) implies $f(u_6) = f_6 - f_5 \geq 48 - 24 = 24 > f_5 - f(u_1)$. So we have from Lemma 2.4 that $m+n \geq 7$ and $f(u_7) \leq f(u_1) + f(u_6) \leq 1 + (f_5 + 1) \leq 26$. However, this gives $f_7 \leq f_5 + f(u_6) + f(u_7) \leq f_5 + (f_5 + 1) + 26 \leq 75 < 3 \cdot 2^{7-2}$, contradicting to Lemma 3.6(2) again. Therefore, " $m+n = 5$ " is the only possible situation and then $f(G) = f_5 = 24 < 2^{m+2} - 2^m + 1$.

Case 2. $(i_1, t) = (2, 0)$.

In this case, $f(u_1) = 1$, $f(u_2) = 2$ and $u_1 \in V_1$. If $i_2 \geq 5$, then the upper bound on $f(G)$ in (6) can be rewritten as

$$f(G) \leq (2^{m+2} - 2^m) - 3 \cdot 2^{m-3} + 2^{m-3}(1 - 2^{-1} + 2^1) < 2^{m+2} - 2^m + 1.$$

If $i_2 = 4$, then since $f(u_3) > 1 = f_2 - f(u_2)$, one can see that u_3 must be in V_1 from the definition of i_2 . Therefore we have $V_1 = \{u_1, u_3\}$ and $V_0 = \{u_i \mid i \in \{2\} \cup [4, m+n]\}$. Note that $f(u_4) \leq f_3 + 1 = 8$. From the definition of i_2 and Lemma 2.4, we have $f(u_5) \leq f(u_2) + f(u_4) = 2 + f(u_4)$. Now, $3 \cdot 2^{5-2} \leq f_5 = f_3 + f(u_4) + f(u_5) \leq (1 + 2 + 4) + f(u_4) + (2 + f(u_4))$. This implies that $f(u_4) = 8 = f_3 + 1$ and $f(u_5) \leq 10$. Hence, $f(u_5) > f(u_4) > 7 \geq f_3 = f_4 - f(u_4)$. It follows from Lemma 2.4 that $f(u_6) \leq f(u_4) + f(u_5) \leq 8 + 10 = 18$ and then $f_6 = f_4 + f(u_5) + f(u_6) = 15 + 10 + 18 = 41 < 3 \cdot 2^{6-2}$ which contradicts to Lemma 3.6(2). We therefore conclude that $i_2 = 4$ can never occur in this case.

If $i_2 = 3$, then $u_1 \in V_1$ and $f(u_3) \leq f_2 + 1 = 4$. From the definition of i_2 and by virtue of Lemma 2.4, we have $f(u_4) \leq f(u_2) + f(u_3) \leq 6$. Suppose that $m+n \geq 5$. Since Lemma 3.6(2) gives that $f_4 \geq 3 \cdot 2^{4-2} = 12$ and $f_4 \leq 1 + 2 + f(u_3) + (f(u_2) + f(u_3)) = 5 + 2f(u_3)$, we obtain that $f(u_3) = 4$. Now, the inequality $f_5 \geq 3 \cdot 2^{5-2} = 24$ from Lemma 3.6(2) leads to $f(u_5) = f_5 - f_4 \geq 24 - (1 + 2 + 4 + 6) = 11 > f(u_3) + f(u_4)$. One can be sure that u_4 must be in V_1 by Lemma 2.4 because $f(u_4) > f(u_3) = 4 > f_3 - f(u_3)$ and $f(u_3) = f_2 + 1$. Hence, we have $V_1 = \{u_1, u_4\}$ and $V_0 = \{u_i \mid i \in \{2, 3\} \cup [5, m+n]\}$. In addition, since $f(u_5) \geq 11 > f_4 - f(u_3)$, we know from Lemma 2.4 that $m+n \geq 6$ and $f(u_6) \leq f(u_3) + f(u_5) \leq 4 + (f_4 + 1) \leq 18$. These facts together lead to $f_6 = f_4 + f(u_5) + f(u_6) \leq 13 + 14 + 18 = 45 < 3 \cdot 2^{6-2}$ which contradicts to Lemma 3.6(2). We therefore conclude that the inequality $m+n \geq 5$ is impossible to be true in Case 3. So,

$m + n = 4$ and then $f(G) = f_4 \leq 13 = 2^{m+2} - 2^m + 1$.

Case 3. $(i_1, t) = (3, 0)$.

In this case, $V_1 = \{u_1, u_2\}$ can be easily seen from the definition of i_1 . So, $V_0 = \{u_i | i \in [3, m+n]\}$. Now, we have $f(u_1) = 1$, $f(u_2) = 2$ and $f(u_3) = f_2 + 1 = 4$. Let $k = f(G) - (f(u_1) + f(u_2))$ and let H be an induced subgraph of G such that $f(H) = k$. If $V_0 \setminus V(H) \neq \emptyset$, we denote as i_0 the minimum element in $\{i | u_i \in V_0 \setminus V(H)\}$, then $i_0 \geq 3$ and $V(H) \subseteq V_1 \cup \{u_i | i \geq i_0 + 1\}$. Therefore, $f(H) \leq f(u_1) + f(u_2) + \sum_{i=i_0+1}^{m+n} f(u_i) \leq f(u_1) + f(u_2) + \sum_{i=4}^{m+n} f(u_i) < \sum_{i=3}^{m+n} f(u_i) = f(H)$, giving a contradiction. We then have that $V_0 \subseteq V(H)$. Besides, since $\sum_{i=3}^{m+n} f(u_i) = k$, we conclude that $V(H) = V_0$ and H is disconnected. Therefore, Case 3 is impossible to occur.

Case 4. $(i_1, t) = (4, 1)$.

In this case, $V_1 = \{u_1, u_2\}$ and $V_0 = \{u_i | i \in [3, m+n]\}$ are also true. Now, $f(u_1) = 1$, $f(u_2) = 2$ and $f(u_3) = f_2 = 3$ and $f(u_4) = f_3 + 1 = 7$. Let $k = f(G) - (f(u_1) + f(u_2) + f(u_3))$. In exactly the same way as we used in Case 3, one can show that Case 4 is not possible either.

Since $f(G) \leq 2^{m+2} - 2^m + 1$ is valid in all possible situations. The proof is completed. \square

4 Conclusion

In this work, we have given lower bounds on the IC-index of all complete multipartite graphs and have shown that our lower bound on $M(K_{1(n),m})$ is the exact value of it. Our coloring constructed in Proposition 3.1 is indeed a qualified maximal IC-coloring.

For further study of this problem, one can try to show that the lower bound given in Proposition 3.3 will also be an upper bound on $M(K_{1(n),m_\ell, m_{\ell-1}, \dots, m_2, m_1})$ for all $\ell \geq 2$. We conjecture that the inequality in Proposition 3.3 is in fact an equality in the case where $\ell \geq 2$, $m_\ell \geq 2$ and $n \geq 2$.

References

- [1] R. Alter, J. A. Brnett, A postage stamp problem, *Amer. Math. Monthly* 87(1980), 206-210 .
- [2] R. Guy, The Postage Stamp Problem, *Unsolved Problems in Number Theory*, second ed. 123-127, Springer, New York, 1994.
- [3] R. L. Heimer, H. Langenbach, The stamp problem, *J. Recreational Math.* 7(1974), 235-250.
- [4] C. T. Long, N. Woo, On bases for the set of integers, *Duke Math. J.* 38(1971), 583-590.
- [5] W. F. Lunnon, A postage stamp problem, *Comput. J.* 12(1969), 377-380.

- [6] S. Mossige, The postage problem: an algorithm to determine the h -range of the h -range formula on the extremal basis problem for $k = 4$, *Math. Comput.* 69(2000), 325-337.
- [7] S. G. Penrice, Some new graph labeling problems: a preliminary report, *DIMACS Tech. Rep.* 95-26(1995), 1-9.
- [8] E. Salehi, Sin-Min Lee, M. Khatirinejad, IC-colorings and IC-indices of graphs, *Discrete Math.* 299(2005), 297-310.
- [9] E. S. Selmer, On the postage stamp problem with the three stamp denominations, *Math. Scand.* 47(1980), 29-71.
- [10] R. G. Stanton, J. G. Kalbfleisch, R. C. Mullin, Some tables for the postage stamp problem, *Proceedings of 4th Manitoba Conference on Numerical Math.*, Utilitas Math. Pub., Winnipeg, (1974), 351-356.
- [11] C. L. Shiue and H. L. Fu, The IC-indices of complete bipartite graphs, *Electronic J. of Combinatorial* 15(2008), R43.
- [12] A. Stöhr, Gelöste und ungelöste Fragen über Basen der natürlichen Zahlenreihe I, II, *J. Reine Angew. Math.* 194(1955), 111-140.
- [13] D. B. West, *Introduction to graph theory*, Prentice-Hall, Upper Saddle River, NJ, 2001.