

# Relative $m$ -ovoids of elliptic quadrics

Antonio Cossidente

Dipartimento di Matematica Informatica ed Economia

Università della Basilicata

Contrada Macchia Romana

I-85100 Potenza

Italy

antonio.cossidente@unibas.it

Francesco Pavese

Dipartimento di Meccanica Matematica e Management

Politecnico di Bari

Via Orabona, 4

I-70125 Bari

Italy

francesco.pavese@poliba.it

## Abstract

Let  $\mathcal{Q}^-(2n+1, q)$  be an elliptic quadric of  $\text{PG}(2n+1, q)$ . A *relative  $m$ -ovoid* of  $\mathcal{Q}^-(2n+1, q)$  (with respect to a parabolic section  $\mathcal{Q} := \mathcal{Q}(2n, q) \subset \mathcal{Q}^-(2n+1, q)$ ) is a subset  $\mathcal{R}$  of points of  $\mathcal{Q}^-(2n+1, q) \setminus \mathcal{Q}$  such that every generator of  $\mathcal{Q}^-(2n+1, q)$  not contained in  $\mathcal{Q}$  meets  $\mathcal{R}$  in precisely  $m$  points. A relative  $m$ -ovoid having the same size as its complement (in  $\mathcal{Q}^-(2n+1, q) \setminus \mathcal{Q}$ ) is called a *relative hemisystem*. We show that a nontrivial relative  $m$ -ovoid of  $\mathcal{Q}^-(2n+1, q)$  is necessarily a relative

hemisystem, forcing  $q$  to be even. Also, we construct an infinite family of relative hemisystems of  $\mathcal{Q}^-(4n+1, q)$ ,  $n \geq 2$ , admitting  $\mathrm{PSp}(2n, q^2)$  as an automorphism group. Finally, some applications are given.

**Keywords:** elliptic quadric, relative hemisystem, strongly regular graph

**MSC 2010:** 05B25, 51A50, 51E20, 05E30

## 1 Introduction

In 2011 Penttinen and Williford [12] introduced the notion of *relative hemisystem* of a Hermitian surface  $\mathcal{H}(3, q^2)$ ,  $q$  even. A relative hemisystem of  $\mathcal{H}(3, q^2)$  with respect to a symplectic subgeometry  $\mathcal{W}(3, q) \subset \mathcal{H}(3, q^2)$  is a set  $\mathcal{S}$  of generators of  $\mathcal{H}(3, q^2)$  disjoint from  $\mathcal{W}(3, q)$  such that each point of  $\mathcal{H}(3, q^2) \setminus \mathcal{W}(3, q)$  lies on  $q/2$  generators of  $\mathcal{S}$ . Relative hemisystems are interesting in view of their connection with primitive  $Q$ -polynomial association schemes not arising from distance regular graphs [12, Theorem 3]. Up to date, three infinite families of relative hemisystems of  $\mathcal{H}(3, q^2)$  and a sporadic example are known [12], [4], [5], [6].

In a recent paper Bamberg and Lee [1] generalized the concept of a relative hemisystem to a *relative  $m$ -cover* of a generalized quadrangle. Let  $\Gamma$  be a generalized quadrangle with a subquadrangle  $\Gamma'$ . A relative  $m$ -cover of  $\Gamma$  is a set  $\mathcal{S}$  of external lines with respect to  $\Gamma'$  such that each point of  $\Gamma \setminus \Gamma'$  lies on  $m$  members of  $\mathcal{S}$ . Bamberg and Lee showed that a relative  $m$ -cover of a generalized quadrangle of order  $(q^2, q)$  with respect to a subquadrangle of order  $(q, q)$  must be a hemisystem. In particular, they were able to prove an analogue of Segre's result [13], [8, Section 19.3]. See also [7]. Indeed they showed that a nontrivial relative  $m$ -cover of  $\mathcal{H}(3, q^2)$ , with respect to a symplectic subgeometry  $\mathcal{W}(3, q)$  is a relative hemisystem, forcing  $q$  to be even [1, Theorem 1.1, Corollary 1.2].

An  $m$ -ovoid of a finite polar space  $\mathcal{P}$  is a set of points which meets every generator of  $\mathcal{P}$  in precisely  $m$  points [2].

Dual of the generalized quadrangles  $\mathcal{H}(3, q^2)$  and  $\mathcal{W}(3, q)$  are the elliptic quadric  $\mathcal{Q}^-(5, q)$  and the parabolic quadric  $\mathcal{Q}(4, q)$ , respectively. A relative  $m$ -

cover of  $\mathcal{H}(3, q^2)$  corresponds then to a set of points  $\mathcal{R}$  of  $\mathcal{Q}^-(5, q) \setminus \mathcal{Q}(4, q)$  such that each generator of  $\mathcal{Q}^-(5, q)$  not on  $\mathcal{Q}(4, q)$  meets  $\mathcal{R}$  in  $m$  points. In this way, the dual of a relative  $m$ -cover of  $\mathcal{H}(3, q^2)$  can be considered an “affine”  $m$ -ovoid of  $\mathcal{Q}^-(5, q)$ .

In this paper we introduce the concept of relative  $m$ -ovoid of  $\mathcal{Q}^-(2n + 1, q)$  with respect to a parabolic section  $\mathcal{Q} := \mathcal{Q}(2n, q) \subset \mathcal{Q}^-(2n + 1, q)$  as a generalization of the dual of a relative  $m$ -cover of  $\mathcal{H}(3, q^2)$ . A relative  $m$ -ovoid of  $\mathcal{Q}^-(2n + 1, q)$  (with respect to  $\mathcal{Q}$ ) is a subset  $\mathcal{R}$  of points of  $\mathcal{Q}^-(2n + 1, q) \setminus \mathcal{Q}$  such that every generator of  $\mathcal{Q}^-(2n + 1, q)$  not contained in  $\mathcal{Q}$  meets  $\mathcal{R}$  in precisely  $m$  points. A relative  $m$ -ovoid having the same size as its complement (in  $\mathcal{Q}^-(2n + 1, q) \setminus \mathcal{Q}$ ) is called a *relative hemisystem*.

The paper is organized as follows. In Section 2 we show that a nontrivial relative  $m$ -ovoid of  $\mathcal{Q}^-(2n + 1, q)$  is necessarily a relative hemisystem, forcing  $q$  to be even. This generalizes the main result of [1]. In Section 3 we construct an infinite family of relative hemisystems of  $\mathcal{Q}^-(4n + 1, q)$ ,  $n \geq 2$ , admitting  $\text{PSp}(2n, q^2)$  as an automorphism group. In Section 4 we show that a relative hemisystem of  $\mathcal{Q}^-(2n + 1, q)$  gives rise to non-classical  $(q^n - 1)/(q - 1)$ -ovoids of the symplectic polar space  $\mathcal{W}(2n + 1, q)$ , generalizing [6, Proposition 3.1, Corollary 3.2] and, assuming  $q = 2$  and  $n \geq 3$ , to a strongly regular graph with parameters  $v = 2^{n-1}(2^n - 1)$ ,  $k = (2^{n-2} - 1)(2^n + 1)$ ,  $\lambda = 2(2^{n-1} + 1)(2^{n-3} - 1)$ ,  $\mu = 2^{n-1}(2^{n-2} - 1)$ .

Throughout the paper we will use the term  $n$ -space to refer to an  $n$ -dimensional projective space.

## 2 Relative $m$ -ovoids of $\mathcal{Q}^-(2n + 1, q)$

Let  $\text{PG}(2n + 1, q)$  be the projective space over the finite field  $\text{GF}(q)$  and let  $\mathcal{Q}^-(2n + 1, q)$  be an elliptic quadric of  $\text{PG}(2n + 1, q)$ . Let  $\mathcal{Q} = \mathcal{Q}(2n, q)$  be a parabolic quadric embedded in  $\mathcal{Q}^-(2n + 1, q)$ . Let  $\perp$  denote the polarity of  $\text{PG}(2n + 1, q)$  induced by  $\mathcal{Q}^-(2n + 1, q)$ . A *generator* of  $\mathcal{Q}^-(2n + 1, q)$  is a totally singular  $(n - 1)$ -space. A generator of  $\mathcal{Q}^-(2n + 1, q)$  either is contained in  $\mathcal{Q}$  or

meets  $\mathcal{Q}$  in an  $(n-2)$ -space. Let  $\mathcal{B}$  denote the set of generators of  $\mathcal{Q}^-(2n+1, q)$  not contained in  $\mathcal{Q}$ . Then  $|\mathcal{B}| = q(q^n - 1) \prod_{i=2}^n (q^i + 1)$ .

**Definition 2.1.** A relative  $m$ -ovoid of  $\mathcal{Q}^-(2n+1, q)$  (with respect to  $\mathcal{Q}$ ) is a subset  $\mathcal{R}$  of points of  $\mathcal{Q}^-(2n+1, q) \setminus \mathcal{Q}$  such that every generator of  $\mathcal{Q}^-(2n+1, q)$  not contained in  $\mathcal{Q}$  meets  $\mathcal{R}$  in  $m$  points.

A relative  $m$ -ovoid is said to be *nontrivial* or *proper* if  $m \neq 0$  or  $m \neq q^{n-1}$ . A relative  $m$ -ovoid having the same size as its complement (in  $\mathcal{Q}^-(2n+1, q) \setminus \mathcal{Q}$ ) is called a *relative hemisystem*.

**Lemma 2.2.** A relative  $m$ -ovoid  $\mathcal{R}$  of  $\mathcal{Q}^-(2n+1, q)$  has size  $mq(q^n - 1)$ .

*Proof.* We count in two ways the pairs  $(P, b)$ , where  $P \in \mathcal{R}$ ,  $b \in \mathcal{B}$  and  $P \in b$ :

$$|\mathcal{R}| \prod_{i=2}^n (q^i + 1) = mq(q^n - 1) \prod_{i=2}^n (q^i + 1)$$

and hence  $|\mathcal{R}| = mq(q^n - 1)$ . □

**Lemma 2.3.** Let  $\mathcal{Q}'$  be a parabolic quadric  $\mathcal{Q}(2n, q)$  embedded in  $\mathcal{Q}^-(2n+1, q)$  and let  $P$  be a point of  $\mathcal{Q}^-(2n+1, q)$ .

- a) If  $\mathcal{Q} \cap \mathcal{Q}'$  is an elliptic quadric  $\mathcal{Q}^-(2n-1, q)$ , then  $|\mathcal{R} \cap \mathcal{Q}'| = m(q^n + 1)$ .
- b) If  $\mathcal{Q} \cap \mathcal{Q}'$  is a hyperbolic quadric  $\mathcal{Q}^+(2n-1, q)$ , then  $|\mathcal{R} \cap \mathcal{Q}'| = m(q^n - 1)$ .
- c) If  $\mathcal{Q} \cap \mathcal{Q}'$  is a cone, then  $|\mathcal{R} \cap \mathcal{Q}'| = mq^n$ .
- d) If  $P \in \mathcal{R}$ , then  $|\mathcal{R} \cap P^\perp| = m(q^n + 1) - q^n$ .
- e) If  $P \notin \mathcal{Q} \cup \mathcal{R}$ , then  $|\mathcal{R} \cap P^\perp| = m(q^n + 1)$ .
- f) If  $P \in \mathcal{Q}$ , then  $|\mathcal{R} \cap P^\perp| = m(q^n - q)$ .

*Proof.* In the first three cases, we count in two ways the pairs  $(P, b)$ , where  $P \in \mathcal{R} \cap \mathcal{Q}'$ ,  $b \in \mathcal{B}$  is a generator of  $\mathcal{Q}'$  and  $P \in b$ :

a)

$$|\mathcal{R} \cap \mathcal{Q}'| \prod_{i=1}^{n-1} (q^i + 1) = m \prod_{i=1}^n (q^i + 1)$$

and hence  $|\mathcal{R} \cap \mathcal{Q}'| = m(q^n + 1)$ .

b)

$$|\mathcal{R} \cap \mathcal{Q}'| \prod_{i=1}^{n-1} (q^i + 1) = m \left( \prod_{i=1}^n (q^i + 1) - 2 \prod_{i=1}^{n-1} (q^i + 1) \right)$$

and hence  $|\mathcal{R} \cap \mathcal{Q}'| = m(q^n - 1)$ .

c)

$$|\mathcal{R} \cap \mathcal{Q}'| \prod_{i=1}^{n-1} (q^i + 1) = m \left( \prod_{i=1}^n (q^i + 1) - \prod_{i=1}^{n-1} (q^i + 1) \right)$$

and hence  $|\mathcal{R} \cap \mathcal{Q}'| = mq^n$ .

d) We count in two ways the pairs  $(T, b)$ , where  $T \in \mathcal{R} \cap (P^\perp \setminus \{P\})$ ,  $b \in \mathcal{B}$ ,  $P \in b$  and  $T \in b$ :

$$|\mathcal{R} \cap (P^\perp \setminus \{P\})| \prod_{i=2}^{n-1} (q^i + 1) = (m - 1) \prod_{i=2}^n (q^i + 1)$$

and hence  $|\mathcal{R} \cap P^\perp| = m(q^n + 1) - q^n$ .

In the remaining two cases, we count in two ways the pairs  $(T, b)$ , where  $T \in \mathcal{R} \cap P^\perp$ ,  $b \in \mathcal{B}$ ,  $P \in b$  and  $T \in b$ :

e)

$$|\mathcal{R} \cap P^\perp| \prod_{i=2}^{n-1} (q^i + 1) = m \prod_{i=2}^n (q^i + 1)$$

and hence  $|\mathcal{R} \cap P^\perp| = m(q^n + 1)$ .

f)

$$|\mathcal{R} \cap P^\perp| \prod_{i=2}^{n-1} (q^i + 1) = m \left( \prod_{i=2}^n (q^i + 1) - \prod_{i=1}^{n-1} (q^i + 1) \right)$$

and hence  $|\mathcal{R} \cap P^\perp| = m(q^n - q)$ . □

**Theorem 2.4.** *Let  $\mathcal{R}$  be a proper relative  $m$ -ovoid of  $\mathcal{Q}^-(2n+1, q)$ . Then  $q$  is even and  $\mathcal{R}$  is a relative hemisystem.*

*Proof.* Let  $\mathcal{Q}^-(2n-1, q)$  be an elliptic quadric contained in  $\mathcal{Q}$ . Let  $\mathcal{F}$  be the pencil of quadrics embedded in  $\mathcal{Q}^-(2n+1, q)$  having as base locus  $\mathcal{Q}^-(2n-1, q)$ . Then  $\mathcal{F}$  contains, apart from  $\mathcal{Q}$ ,  $q-2$  parabolic quadrics and two cones. Let  $P_1, P_2$  be the vertices of the two cones. We distinguish three cases according as

$P_1, P_2 \in \mathcal{R}, P_1, P_2 \notin \mathcal{R}, P_1 \in \mathcal{R}, P_2 \notin \mathcal{R}$ , respectively. If the first case occurs then, from Lemma 2.3 a), d), we have

$$(q-2)m(q^n+1) + 2m(q^n+1) - 2q^n = mq(q^n-1)$$

and hence  $m = q^{n-1}$ . If the second case occurs then, from Lemma 2.3 a), e), we have

$$(q-2)m(q^n+1) + 2m(q^n+1) = mq(q^n-1)$$

and hence  $m = 0$ . If the third case occurs then, from Lemma 2.3 a), d), e), we have

$$(q-2)m(q^n+1) + 2m(q^n+1) - q^n = mq(q^n-1)$$

and hence  $2m = q^{n-1}$ . It follows that  $q$  is even and  $m = q^{n-1}/2$ .  $\square$

**Remark 2.5.** Theorem 2.4 does not hold true for hyperbolic quadrics as shown in the following example. Let  $\mathcal{Q}^-(3, q) \subset \mathcal{Q} := \mathcal{Q}(4, q) \subset \mathcal{Q}^+(5, q)$ ,  $q$  even. There exists a pencil  $\mathcal{P}$  of parabolic quadrics having  $\mathcal{Q}^-(3, q)$  as base locus. A plane of  $\mathcal{Q}^+(5, q)$  meets a parabolic quadric of  $\mathcal{P}$  in a line having necessarily a point on  $\mathcal{Q}^-(3, q)$  and hence meets each of the  $q$  sets  $\mathcal{Q}(4, q) \setminus \mathcal{Q}^-(3, q)$  in  $q$  points, i.e., each of them is a “relative  $q$ -ovoid” of  $\mathcal{Q}^+(5, q)$  with respect to  $\mathcal{Q}$ . Of course, gluing together  $i$  of such sets, a “relative  $iq$ -ovoid” is obtained,  $1 \leq i \leq q$ . Hence there exist “relative  $m$ -ovoids” of  $\mathcal{Q}^+(5, q)$  with respect to  $\mathcal{Q}$  that are not relative hemisystems.

In view of Theorem 2.4, we will assume for the remaining part of the paper that  $q$  is even.

Let  $G$  denote the stabilizer of  $\mathcal{Q}^-(2n+1, q)$  in  $\text{PGL}(2n+2, q)$ . Let  $\Sigma$  be the hyperplane of  $\text{PG}(2n+1, q)$  containing  $\mathcal{Q}$  and let  $N = \Sigma^\perp$ . Note that  $N$  is the nucleus of  $\mathcal{Q}$ . Let  $\tau$  be the involution of  $G$  fixing  $\Sigma$  pointwise and  $N$  linewise.

**Corollary 2.6.** *If  $\mathcal{R}$  is a relative hemisystem of  $\mathcal{Q}^-(2n+1, q)$ , then  $|\mathcal{R} \cap \mathcal{R}^\tau| = 0$ .*

*Proof.* With the same notation introduced in the proof of Theorem 2.4 we only observe that the line  $P_1P_2$  contains  $N$ .  $\square$

**Remark 2.7.** Note that Theorem 2.4 and Corollary 2.6 generalize [1, Theorem 1.1 (a), (b)] when the generalized quadrangle is classical.

### 3 Relative hemisystems of $\mathcal{Q}^-(4n + 1, q)$

In this Section we construct a relative hemisystem of  $\mathcal{Q}^-(4n + 1, q)$  admitting the group  $\text{PSp}(2n, q^2)$  as an automorphism group.

Let  $V$  be a  $2n$ -dimensional vector space over  $\text{GF}(q^2)$  and let  $k$  be an alternating form with associated polar space  $\mathcal{W}(2n - 1, q^2)$ . The vector space  $V$  can be considered as a  $4n$ -dimensional vector space  $V'$  over  $\text{GF}(q)$  via the inclusion  $\text{GF}(q) \subset \text{GF}(q^2)$ . Composition of  $k$  with the trace map  $T : z \in \text{GF}(q^2) \mapsto z + z^q \in \text{GF}(q)$  provides a new alternating form  $k'$  on  $V'$  and so we obtain a new polar space  $\mathcal{W}(4n - 1, q)$ . Now each point in  $\text{PG}(2n - 1, q^2)$  corresponds to a 1-dimensional vector space in  $V$ , which in turn corresponds to a 2-dimensional vector space in  $V'$ , that is a 1-dimensional projective space of  $\text{PG}(4n - 1, q)$ . Extending this map from points of  $\text{PG}(2n - 1, q^2)$  to subspaces of  $\text{PG}(2n - 1, q^2)$ , we obtain an injective map from subspaces of  $\text{PG}(2n - 1, q^2)$  to certain subspaces of  $\text{PG}(4n - 1, q)$ :

$$\phi : \text{PG}(2n - 1, q^2) \rightarrow \text{PG}(4n - 1, q).$$

The map  $\phi$  is called the  $\text{GF}(q)$ -linear representation of  $\text{PG}(2n - 1, q^2)$ .

A *line spread* of a polar space  $\mathcal{P}$  is a collection  $\mathcal{S}$  of mutually disjoint lines of  $\mathcal{P}$  such that each point of  $\mathcal{P}$  is contained in an element of  $\mathcal{S}$ . The set  $\mathcal{D} = \{\phi(P) \mid P \in \mathcal{W}(2n - 1, q^2)\}$  is an example of line spread of  $\mathcal{W}(4n - 1, q)$ . The incidence structure whose points are the elements of  $\mathcal{D}$  and whose lines are the generators of  $\mathcal{W}(4n - 1, q)$  spanned by two distinct elements of  $\mathcal{D}$ , is isomorphic to  $\mathcal{W}(2n - 1, q^2)$ . Denote by  $\rho$  the symplectic polarity of  $\text{PG}(2n - 1, q^2)$  associated with  $\mathcal{W}(2n - 1, q^2)$  and by  $\rho'$  the symplectic polarity of  $\text{PG}(4n - 1, q)$  associated with  $\mathcal{W}(4n - 1, q)$

Since  $q$  is even, the symplectic polar space  $\mathcal{W}(4n - 1, q)$  is isomorphic to the parabolic quadric  $\mathcal{Q} = \mathcal{Q}(4n, q)$ . This isomorphism can be realized by projection

of  $\mathcal{Q}$  from its nucleus  $N$  onto a hyperplane not containing  $N$  [14]. Denote by  $\mu$  the bijection between  $\mathcal{W}(4n-1, q)$  and  $\mathcal{Q}$ . Thus,  $\mu(\mathcal{D})$  is a line spread of  $\mathcal{Q}$  admitting  $\text{PSp}(2n, q^2)$  as an automorphism group.

Let  $\Sigma$  be the ambient projective space of  $\mathcal{Q}$  and consider  $\Sigma$  as a hyperplane of a  $\text{PG}(4n+1, q)$ . Let  $\mathcal{Q}^-(4n+1, q)$  be an elliptic quadric of  $\text{PG}(4n+1, q)$  such that  $\mathcal{Q}^-(4n+1, q) \cap \Sigma = \mathcal{Q}$ . In this way, if  $G$  denotes the stabilizer of  $\mathcal{Q}^-(4n+1, q)$  in  $\text{PGL}(4n+2, q)$ , by Witt's theorem, we can say that there exists a subgroup  $K$  of  $G$ , isomorphic to  $\text{PSp}(2n, q^2)$ , that stabilizes the line spread  $\mu(\mathcal{D})$  of  $\mathcal{Q}$ .

Let  $\perp$  denote the orthogonal polarity of  $\text{PG}(4n+1, q)$  associated with  $\mathcal{Q}^-(4n+1, q)$ . Then  $N = \Sigma^\perp$ . Let  $\tau$  be the involution of  $G$  fixing  $\Sigma$  pointwise and  $N$  linewise. We are interested in the action of  $K$  on points of  $\mathcal{Q}^-(4n+1, q) \setminus \mathcal{Q}$ .

**Proposition 3.1.** *The group  $K$  has two orbits  $\mathcal{O}_1, \mathcal{O}_2$  of the same size on points of  $\mathcal{Q}^-(4n+1, q) \setminus \mathcal{Q}$ . Moreover,  $\mathcal{O}_1^\perp = \mathcal{O}_2$ .*

*Proof.* Let  $P_1 \in \mathcal{Q}^-(4n+1, q) \setminus \mathcal{Q}$ , then  $P_1^\perp$  meets  $\mathcal{Q}$  in an elliptic quadric  $\mathcal{Q}^-(4n-1, q)$ . An easy counting argument shows that  $\mathcal{Q}^-(4n-1, q)$  contains exactly  $|\mathcal{Q}^-(2n-1, q^2)|$  lines of  $\mu(\mathcal{D})$ . If  $\mathcal{E}$  is an elliptic quadric of  $\text{PG}(2n-1, q^2)$  inducing the polarity  $\rho$ , then  $\phi(\mathcal{E})$  defines a set consisting of  $|\mathcal{E}|$  lines of  $\mathcal{D}$  that are contained in an elliptic quadric  $\mathcal{E}'$  of  $\text{PG}(4n-1, q)$  that in turn induces the polarity  $\rho'$ . Note that  $\mu(\mathcal{E}')$  is an elliptic quadric  $\mathcal{Q}^-(4n-1, q)$  contained in  $\mathcal{Q}$ . Moreover, the lines of  $\mu(\mathcal{D})$  contained in  $\mu(\mathcal{E}')$  span a  $\text{PG}(4n-1, q)$ . On the other hand the number of elliptic quadrics of  $\text{PG}(2n-1, q^2)$  inducing the polarity  $\rho$  equals the number of elliptic quadrics of  $\text{PG}(4n-1, q)$  contained in  $\mathcal{Q}$ , indeed

$$|\text{PSp}(2n, q^2) : \text{PGO}^-(2n, q^2)| = |\text{PSp}(4n, q) : \text{PGO}^-(4n, q)| = \frac{q^{2n}(q^{2n}-1)}{2}.$$

It follows that there exists a one-to-one correspondence between the elliptic quadrics of  $\text{PG}(2n-1, q^2)$  inducing the polarity  $\rho$  and the elliptic quadrics of  $\text{PG}(4n-1, q)$  contained in  $\mathcal{Q}$ . Since  $\text{PGO}^-(2n, q^2)$  is maximal in  $K$  [9], we have that the stabilizer of  $P_1$  in  $K$  is isomorphic to  $\text{PGO}^-(2n, q^2)$  and hence from the Orbit-Stabilizer theorem  $|P_1^K| = q^{2n}(q^{2n}-1)/2$ .

Let  $\ell$  be the line joining  $P_1$  and  $N$  and let  $\ell \cap \mathcal{Q}^-(4n+1, q) = \{P_1, P_2\}$ . Since  $K$  fixes  $N$ , if there was an element  $\alpha \in K$  such that  $P_1^\alpha = P_2$ , then  $\alpha$  would fix  $\ell$  and hence  $\alpha$  would lie in the stabilizer of  $\ell$  in  $K$ . On the other hand, since  $P_1^\perp \cap \Sigma = \ell^\perp$ , the stabilizer of  $\ell$  in  $K$  coincides with the stabilizer of  $P_1$  in  $K$ , which acts identically on  $\ell$ . It follows that  $P_2 \notin P_1^K$ . In particular  $K$  has two orbits  $\mathcal{O}_1$  and  $\mathcal{O}_2$  on points of  $\mathcal{Q}^-(4n+1, q) \setminus \mathcal{Q}$ , both of size  $q^{2n}(q^{2n}-1)/2$ , and if  $\ell$  is a secant line to  $\mathcal{Q}^-(4n-1, q)$  containing  $N$ , then  $|\ell \cap \mathcal{O}_1| = |\ell \cap \mathcal{O}_2| = 1$ .  $\square$

From now on we assume that  $n \geq 2$ . We are now interested in determining the orbits of the group  $K$  on the set  $\mathcal{L}$  consisting of lines of  $\mathcal{Q}^-(4n+1, q)$  not contained in  $\mathcal{Q}$ .

**Proposition 3.2.** *The group  $K$  has three orbits on lines of  $\mathcal{L}$ .*

*Proof.* With the same notation introduced in Proposition 3.1, let  $P_1 \in \mathcal{O}_1$  and let  $P_2 = P_1^\tau$ . Through  $P_1$  there pass  $|\mathcal{Q}^-(4n-1, q)|$  lines of  $\mathcal{L}$  and, as already observed,  $\Gamma = P_1^\perp \cap \Sigma$  meets  $\mathcal{Q}^-(4n+1, q)$  in a  $\mathcal{Q}^-(4n-1, q)$  that corresponds to an elliptic quadric  $\mathcal{Q}^-(2n-1, q^2)$  inducing the polarity  $\rho$ . Note that  $N \notin \Gamma$ . Hence  $\mathcal{Q}^-(4n-1, q)$  contains  $|\mathcal{Q}^-(2n-1, q^2)|$  lines of  $\mu(\mathcal{D})$ . Let  $X$  denote the set of points covered by the lines of  $\mu(\mathcal{D})$  that are contained in  $\mathcal{Q}^-(4n-1, q)$ .

Let  $\ell$  be a line of  $\mathcal{Q}^-(4n+1, q)$  passing through  $P_1$  and meeting  $\mathcal{Q}$  at the point  $P$ . Let  $\ell_P$  be the unique line of  $\mu(\mathcal{D})$  through  $P$  and let  $\ell' = P_2P$ . The plane  $\pi = \langle \ell, \ell_P \rangle$  is either contained in  $\mathcal{Q}^-(4n+1, q)$  or shares with  $\mathcal{Q}^-(4n+1, q)$  the  $2q+1$  points of  $\ell \cup \ell_P$ . Let  $\pi' = \langle \ell_P, P_2 \rangle$

If  $P \in X$ , then  $\pi$  is contained in  $\mathcal{Q}^-(4n+1, q)$  and the solid  $\langle \pi, P_2 \rangle$  meets  $\mathcal{Q}^-(4n+1, q)$  in the union of the two planes,  $\pi$  and  $\pi'$ . Let  $L = \phi^{-1}(\mu^{-1}(\ell_P))$ . Then  $L \in \mathcal{Q}^-(2n-1, q^2)$ . If  $r$  is a secant line to  $\mathcal{Q}^-(2n-1, q^2)$  through  $L$ , then  $\mu(\phi(r))$  is a set of  $q^2+1$  lines of  $\mu(\mathcal{D})$  partitioning the point set of a  $\mathcal{Q}(4, q)$  contained in  $\mathcal{Q}$ . In particular  $\ell_P \in \mu(\phi(r))$  and  $\mathcal{Q}(4, q)$  meets  $\mathcal{Q}^-(4n-1, q)$  in a hyperbolic quadric  $\mathcal{Q}^+(3, q)$ . Indeed, since a line of  $\mu(\mathcal{D})$  is either contained in  $\mathcal{Q}^-(4n-1, q)$  or it is tangent to it and  $\mathcal{Q}(4, q)$  contains  $q^2+1$  lines of  $\mu(\mathcal{D})$ , it follows that  $|\mathcal{Q}(4, q) \cap \mathcal{Q}^-(4n-1, q)| = q^2 - 1 + 2(q+1) = (q+1)^2$ . Let  $\mathcal{E} = \mathcal{Q}^-(4n-5, q)$  be the elliptic quadric  $\langle \mathcal{Q}^+(3, q) \rangle^{\perp_{|\mathcal{R}}}$  and

let  $\mathcal{Q}^+(5, q) = \langle \mathcal{E} \rangle^\perp \cap \mathcal{Q}^-(4n+1, q)$ . By construction  $\pi$  and  $\pi'$  are planes of  $\mathcal{Q}^+(5, q)$  and  $\mathcal{Q}(4, q) \subset \mathcal{Q}^+(5, q)$ . The stabilizer of  $\mathcal{Q}(4, q)$  in  $K$  certainly contains a group  $H \simeq \text{PSp}(2, q^2) \times 1 \leq \text{PSp}(2, q^2) \times \text{PSp}(2n-2, q^2)$  acting in its natural representation on the  $q^2+1$  lines of  $\mu(\mathcal{D})$  contained in  $\mathcal{Q}(4, q)$ . Since  $N \in \langle \mathcal{Q}(4, q) \rangle$ , we have that  $\langle \mathcal{Q}(4, q) \rangle^\perp$  is contained in  $\Sigma$ . Also, since  $H$  fixes  $\langle \mathcal{Q}(4, q) \rangle^\perp$  pointwise, it follows that  $H$  stabilizes  $\mathcal{Q}^+(5, q)$ . Under the Klein map, the action of  $H$  on points of  $\mathcal{Q}^+(5, q)$  corresponds to the action of the stabilizer of an elliptic quadric  $\mathcal{Q}^-(3, q)$  in  $\text{PGL}(4, q)$  on lines of  $\text{PG}(3, q)$ . Then  $H$  has two orbits of size  $q^2(q^2+1)/2$  on points of  $\mathcal{Q}^+(5, q) \setminus \mathcal{Q}(4, q)$  corresponding to lines of  $\text{PG}(3, q)$  that are secant or external to  $\mathcal{Q}^-(3, q)$ , respectively. Since  $\pi$  and  $\pi'$  share a line, they belong to different systems of generators of  $\mathcal{Q}^+(5, q)$ . We may assume, that  $\pi$  is a Latin plane and  $\pi'$  is a Greek plane. Hence the  $q^2$  points of  $\pi \setminus \mathcal{Q}(4, q)$  (resp.  $\pi' \setminus \mathcal{Q}(4, q)$ ) correspond to  $q^2$  secant lines (resp. external lines) to  $\mathcal{Q}^-(3, q)$  and therefore they lie in the same  $H$ -orbit. Since  $H \leq K$ , the line  $\ell$  contains  $q$  points of  $\mathcal{O}_1$ . Analogously, the line  $\ell'$  contains  $q$  points of  $\mathcal{O}_2$ .

If  $P \notin X$ , then  $\pi$  shares with  $\mathcal{Q}^-(4n+1, q)$  the  $2q+1$  points of  $\ell \cup \ell_P$ . Let  $L = \phi^{-1}(\mu^{-1}(\ell_P))$ . Then  $L \notin \mathcal{Q}^-(2n-1, q^2)$ . If  $r$  is an external line to  $\mathcal{Q}^-(2n-1, q^2)$  through  $L$ , then  $\mu(\phi(r))$  is a set of  $q^2+1$  lines of  $\mu(\mathcal{D})$  partitioning the point set of a  $\mathcal{Q}(4, q)$  contained in  $\mathcal{Q}$ . In particular  $\ell_P \in \mu(\phi(r))$  and  $\mathcal{Q}(4, q)$  meets  $\mathcal{Q}^-(4n-1, q)$  in an elliptic quadric  $\mathcal{Q}^-(3, q)$ . Indeed, since a line of  $\mu(\mathcal{D})$  is either contained in  $\mathcal{Q}^-(4n-1, q)$  or it is tangent to it and  $\mathcal{Q}(4, q)$  contains  $q^2+1$  lines of  $\mu(\mathcal{D})$ , it follows that  $|\mathcal{Q}(4, q) \cap \mathcal{Q}^-(4n-1, q)| = q^2+1$ . Let  $\mathcal{H} = \mathcal{Q}^+(4n-5, q)$  be the quadric  $\langle \mathcal{Q}^-(3, q) \rangle^{\perp \text{tr}}$  and let  $\mathcal{Q}^-(5, q) = \langle \mathcal{H} \rangle^\perp \cap \mathcal{Q}^-(4n+1, q)$ . By construction  $\ell$  and  $\ell'$  are lines of  $\mathcal{Q}^-(5, q)$  and  $\mathcal{Q}(4, q) \subset \mathcal{Q}^-(5, q)$ . The stabilizer of  $\mathcal{Q}(4, q)$  in  $K$  certainly contains a group  $H \simeq \text{PSp}(2, q^2) \times 1 \leq \text{PSp}(2, q^2) \times \text{PSp}(2n-2, q^2)$  acting in its natural representation on the  $q^2+1$  lines of  $\mu(\mathcal{D})$  contained in  $\mathcal{Q}(4, q)$ . Since  $N \in \langle \mathcal{Q}(4, q) \rangle$ , we have that  $\langle \mathcal{Q}(4, q) \rangle^\perp$  is contained in  $\Sigma$ . Also, since  $H$  fixes  $\langle \mathcal{Q}(4, q) \rangle^\perp$  pointwise, it follows that  $H$  stabilizes  $\mathcal{Q}^-(5, q)$ . Under the Klein map, the action of  $H$  on points of  $\mathcal{Q}^-(5, q)$  corresponds to the action of the stabilizer of an elliptic quadric  $\mathcal{Q}^-(3, q)$

contained in a symplectic polar space  $\mathcal{W}(3, q)$  embedded in a Hermitian surface  $\mathcal{H}(3, q^2)$  in  $\text{PGU}(4, q^2)$  on lines of  $\mathcal{H}(3, q^2)$ . From [12],  $H$  has two orbits,  $\mathcal{R}_1$  and  $\mathcal{R}_2$ , of size  $q^2(q^2 - 1)/2$  on points of  $\mathcal{Q}^-(5, q) \setminus \mathcal{Q}(4, q)$  and each of them is a relative hemisystem of  $\mathcal{Q}^-(5, q)$  with respect to  $\mathcal{Q}(4, q)$ . In particular, we may assume that  $P_i \in \mathcal{R}_i$ ,  $i = 1, 2$  and, since  $H \leq K$ , we have that  $\mathcal{R}_i \subset \mathcal{O}_i$ ,  $i = 1, 2$ . Also, from [12], the group  $H$  acts transitively on lines of  $\mathcal{Q}^-(5, q)$  that are not contained in  $\mathcal{Q}(4, q)$ . Therefore  $\ell$  and  $\ell'$  lie in the same  $H$ -orbit and both contain  $q/2$  points of  $\mathcal{O}_1$  and  $q/2$  points of  $\mathcal{O}_2$ .

Let  $\mathcal{L}_i$  be the set of lines of  $\mathcal{L}$  containing  $q$  points of  $\mathcal{O}_i$ ,  $i = 1, 2$  and  $\mathcal{L}_3$  be the set of lines of  $\mathcal{L}$  containing  $q/2$  points of  $\mathcal{O}_1$  and  $\mathcal{O}_2$ . Through the point  $P_1$  (resp.  $P_2$ ), there pass  $(q^{2n-2} - 1)(q^{2n} + 1)/(q - 1)$  lines of  $\mathcal{L}_1$  (resp.  $\mathcal{L}_2$ ) and  $q^{2n-2}(q^{2n} + 1)$  lines of  $\mathcal{L}_3$ . From Proposition 3.1, the stabilizer of  $P_1$  (resp.  $P_2$ ) in  $K$  is a group  $J$  isomorphic to  $\text{PGO}^-(2n, q^2)$ . The group  $J$  has two orbits on points of  $\mathcal{Q}^-(4n - 1, q)$ , namely  $X$  and its complement. Therefore  $J$  (and hence  $K$ ) permutes in a single orbit the lines of  $\mathcal{L}_i$  passing through  $P_1$  (resp.  $P_2$ ),  $i = 1, 2, 3$ . Hence, if  $\ell_i \in \mathcal{L}_i$ , we have that

$$|\ell_i^K| = \frac{q^{2n}(q^{2n} - 1)(q^{2n-2} - 1)(q^{2n} + 1)}{2q(q - 1)} = \frac{q^{2n-1}(q^{2n-2} - 1)(q^{4n} - 1)}{2(q - 1)}, i = 1, 2,$$

whereas, if  $\ell_3 \in \mathcal{L}_3$ , we have that

$$|\ell_3^K| = \frac{2q^{2n}(q^{2n} - 1)q^{2n-2}(q^{2n} + 1)}{2q} = q^{4n-3}(q^{4n} - 1).$$

On the other hand,

$$\sum_{i=1}^3 |\ell_i^K| = |\mathcal{L}| = \frac{q^{2n-1}(q^{2n-1} - 1)(q^{4n} - 1)}{q - 1}$$

and hence  $\mathcal{L}_i = \ell_i^K$  is an orbit under the action of  $K$ ,  $i = 1, 2, 3$ .  $\square$

We are ready to prove the main result of this Section.

**Theorem 3.3.**  $\mathcal{O}_i$  is a relative hemisystem of  $\mathcal{Q}^-(4n + 1, q)$ ,  $i = 1, 2$ .

*Proof.* Let  $g$  be a generator of  $\mathcal{Q}^-(4n + 1, q)$  not contained in  $\mathcal{Q}$  and let  $g'$  be the  $(2n-2)$ -space obtained by intersecting  $g$  with  $\mathcal{Q}$ . Let  $P$  be a point of  $g \setminus g'$  and let

$\mathcal{Q}^-(4n-1, q) = P^\perp \cap \mathcal{Q}$ . Then  $g'$  is a generator of  $\mathcal{Q}^-(4n-1, q)$  and, as already observed,  $\mathcal{Q}^-(4n-1, q)$  contains  $|\mathcal{Q}^-(2n-1, q^2)|$  lines of  $\mu(\mathcal{D})$ . Let  $X$  denote the set of points covered by the lines of  $\mu(\mathcal{D})$  that are contained in  $\mathcal{Q}^-(4n-1, q)$ . From [11, Theorem 1.1, Table 1], the set  $X$  is a  $(q^{2n-2} - 1)/(q - 1)$ -ovoid of  $\mathcal{Q}^-(4n-1, q)$ , i.e., every generator of  $\mathcal{Q}^-(4n-1, q)$  meets  $X$  in  $(q^{2n-2} - 1)/(q - 1)$  points. If  $P \in \mathcal{O}_1$ , then  $g$  contains  $(q^{2n-2} - 1)/(q - 1)$  lines of  $\mathcal{L}_1$  passing through  $P$  and  $q^{2n-2}$  lines of  $\mathcal{L}_3$  passing through  $P$ . It follows that  $g$  contains

$$\frac{(q-1)(q^{2n-2}-1)}{q-1} + \left(\frac{q}{2}-1\right)q^{2n-2} + 1 = \frac{q^{2n-1}}{2}$$

points of  $\mathcal{O}_1$  and

$$\frac{q}{2}q^{2n-2} = \frac{q^{2n-1}}{2}$$

points of  $\mathcal{O}_2$ . Analogously, if  $P \in \mathcal{O}_2$ . □

**Remark 3.4.** The relative hemisystems of  $\mathcal{Q}^-(4n+1, q)$  constructed in Theorem 3.3 admit  $\text{PSp}(2n, q^2)$  as an automorphism group.

**Remark 3.5.** With the aid of Magma [3] we checked that  $\mathcal{Q}^-(7, 2)$  has no relative hemisystems.

## 4 Applications

In this section we show that associated to a relative hemisystem of  $\mathcal{Q}^-(2n+1, q)$ ,  $n \geq 2$ , is a  $\left(\frac{q^n-1}{q-1}\right)$ -ovoid of  $\mathcal{W}(2n+1, q)$ . Also, we provide a construction of a strongly regular graph with parameters  $v = 2^{n-1}(2^n-1)$ ,  $k = (2^{n-2}-1)(2^n+1)$ ,  $\lambda = 2(2^{n-1}+1)(2^{n-3}-1)$ ,  $\mu = 2^{n-1}(2^{n-2}-1)$  from a relative hemisystem of  $\mathcal{Q}^-(2n+1, 2)$ ,  $n \geq 3$ . Note that, when  $n$  is even, this graph has the same parameters as the strongly regular graph constructed in [10], by considering as vertices the elliptic sections of a parabolic quadric of  $\text{PG}(n, 4)$ .

### 4.1 $\left(\frac{q^n-1}{q-1}\right)$ -ovoids of $\mathcal{W}(2n+1, q)$ , $q$ even

In  $\mathcal{W}(2n+1, q)$ ,  $q$  even, the classical example of  $\left(\frac{q^n-1}{q-1}\right)$ -ovoid is an elliptic quadric  $\mathcal{Q}^-(2n+1, q)$  polarizing to the symplectic polarity of  $\mathcal{W}(2n+1, q)$ .

Our examples of  $\left(\frac{q^n-1}{q-1}\right)$ -ovoids are examples of *quasi-quadrics* in the sense that they have the same size and the same intersection numbers with respect to hyperplanes of a non-degenerate quadric of  $\text{PG}(2n+1, q)$ .

Let  $\mathcal{Q}^-(2n+1, q)$ ,  $n \geq 2$ , be an elliptic quadric of  $\text{PG}(2n+1, q)$ , let  $\mathcal{Q} = \mathcal{Q}(2n, q)$  be a parabolic quadric embedded in  $\mathcal{Q}^-(2n+1, q)$  and let  $\mathcal{R}$  be a relative hemisystem of  $\mathcal{Q}^-(2n+1, q)$  with respect to  $\mathcal{Q}$ . Since  $q$  is even, the polarity induced by  $\mathcal{Q}^-(2n+1, q)$  gives rise to a symplectic polar space  $\mathcal{W}(2n+1, q)$ . If  $\delta$  is a non-zero element of  $\text{GF}(q)$  such that  $X^2 + X + \delta = 0$  is irreducible over  $\text{GF}(q)$ , then we may assume that  $\mathcal{Q}^-(2n+1, q)$  is given by

$$X_1X_{2n+2} + X_2X_{2n+1} + \dots + X_nX_{n+3} + X_{n+1}^2 + X_{n+1}X_{n+2} + \delta X_{n+2}^2 = 0,$$

whereas  $\mathcal{Q}$  is given by

$$X_1X_{2n+2} + X_2X_{2n+1} + \dots + X_nX_{n+3} + X_{n+1}^2 = 0.$$

Let  $\mathcal{Q}^+(2n+1, q)$  be the hyperbolic quadric given by

$$X_1X_{2n+2} + X_2X_{2n+1} + \dots + X_nX_{n+3} + X_{n+1}^2 + X_{n+1}X_{n+2} = 0.$$

Then the quadrics  $\mathcal{Q}^-(2n+1, q)$  and  $\mathcal{Q}^+(2n+1, q)$  share  $\mathcal{Q}(2n, q)$  and generate a pencil  $\mathcal{P}$  comprising  $q/2$  elliptic quadrics,  $q/2$  hyperbolic quadrics and a cone  $\mathcal{C}$  having as basis the parabolic quadric  $\mathcal{Q}(2n, q)$  and as vertex its nucleus  $N$ . Since  $q$  is even, an orthogonal polarity gives rise to a symplectic polarity. In particular, the orthogonal polarities induced by the non-degenerate quadrics of  $\mathcal{P}$  in  $\text{PG}(2n+1, q)$  all polarize to the same symplectic polarity  $\perp$  of  $\text{PG}(2n+1, q)$  which defines the symplectic polar space  $\mathcal{W}(2n+1, q)$ .

Each elliptic quadric  $\mathcal{Q}_i^-$  of  $\mathcal{P}$  is partitioned as follows:  $\mathcal{Q}_i^- = \mathcal{R}_i \cup \mathcal{R}_i^r \cup \mathcal{Q}$ ,  $i = 1 \dots q/2$ , where with  $\mathcal{R}_i$  and  $\mathcal{R}_i^r$  we have denoted the points on  $\mathcal{Q}_i^-$  corresponding to  $\mathcal{R}$  and its complement in  $\mathcal{Q}_i^- \setminus \mathcal{Q}$ , respectively.

Let  $\mathcal{Q}_i^-$  and  $\mathcal{Q}_j^-$  be two elliptic quadrics of  $\mathcal{P}$ . Let  $\mathcal{A}$  be one of the two subsets  $\mathcal{R}_i$  and  $\mathcal{R}_i^c$  on  $\mathcal{Q}_i^-$  and let  $\mathcal{B}$  be one of the two subsets  $\mathcal{R}_j$  and  $\mathcal{R}_j^c$  on  $\mathcal{Q}_j^-$ ,  $i \neq j$ . We show that  $\mathcal{X} = \mathcal{A} \cup \mathcal{B} \cup \mathcal{Q}$  is a  $(q^n - 1)/(q - 1)$ -ovoid of  $\mathcal{W}(2n+1, q)$ . Let  $\pi$  be a totally isotropic  $n$ -space of  $\mathcal{W}(2n+1, q)$ . Three possibilities occur:

1.  $\pi$  meets  $\mathcal{Q}$  in an  $(n-1)$ -space and  $\pi$  is contained in  $N^\perp$ . There are  $\prod_{i=1}^n (q^i + 1)$  such  $n$ -spaces and none of these meets  $\mathcal{X}$  furtherly.
2.  $\pi$  meets  $\mathcal{Q}$  in an  $(n-1)$ -space and  $\pi$  is not contained in  $N^\perp$ . Then  $\pi$  will be contained in a hyperbolic quadric of  $\mathcal{P}$  and  $\pi$  does not meet  $\mathcal{X}$  furtherly.
3.  $\pi$  meets  $\mathcal{Q}$  in an  $(n-2)$ -space  $\sigma$ . Then  $\pi \cap \mathcal{Q}_i^-$  is an  $(n-1)$ -space  $\sigma_i$  and  $\pi \cap \mathcal{Q}_j^-$  is an  $(n-1)$ -space  $\sigma_j$ ,  $i \neq j$ . In this case  $\sigma_i \cap \sigma_j = \sigma$ . Since  $\mathcal{A}$  and  $\mathcal{B}$  are relative hemisystems of  $\mathcal{Q}_i$  and  $\mathcal{Q}_j$ , respectively, we have that  $\sigma_i$  meets  $\mathcal{A}$  in  $q^{n-1}/2$  points and  $\sigma_j$  meets  $\mathcal{B}$  in  $q^{n-1}/2$  points. Then,  $\pi$  meets  $\mathcal{X}$  in  $(q^n - 1)/(q - 1)$  points.

We have proved the following proposition.

**Proposition 4.1.** *A relative hemisystem of  $\mathcal{Q}^-(2n+1, q)$  gives rise to  $2^{q/2}$  non-classical  $(q^n - 1)/(q - 1)$ -ovoids of  $\mathcal{W}(2n+1, q)$ .*

**Corollary 4.2.** *The set  $\mathcal{X}$  is an elliptic quasi-quadric.*

*Proof.* The set  $\mathcal{X}$  has the same size of an elliptic quadric  $\mathcal{Q}^-(2n+1, q)$  of  $\text{PG}(2n+1, q)$  and from [2, Theorem 11] is a two-character set with intersection numbers  $h_1 = (q^n - 1)(q^n + 1)/(q - 1)$  and  $h_2 = (q^n - 1)(q^n + 1)/(q - 1) - q^n$ .  $\square$

The above construction of non-classical  $(q^n - 1)/(q - 1)$ -ovoids of  $\mathcal{W}(2n+1, q)$  could be realized by taking on the two elliptic quadrics  $\mathcal{Q}_i^-$  and  $\mathcal{Q}_j^-$  two subsets  $\mathcal{A}$  and  $\mathcal{B}$  corresponding to two non-isomorphic relative hemisystems.

## 4.2 A strongly regular graph

Let  $\mathcal{R}$  be a relative hemisystem of  $\mathcal{Q}^-(2n+1, 2)$ ,  $n \geq 3$ . Here  $\perp$  denotes the orthogonal polarity of  $\text{PG}(2n+1, 2)$  induced by  $\mathcal{Q}^-(2n+1, 2)$ . Let  $\mathcal{G}$  be the graph having as vertices the points of  $\mathcal{R}$  with two vertices  $x, y$  being adjacent if  $xy$  is a line of  $\mathcal{Q}^-(2n+1, 2)$ .

**Proposition 4.3.** *The graph  $\mathcal{G}$  is strongly regular with parameters  $v = 2^{n-1}(2^n - 1)$ ,  $k = (2^{n-2} - 1)(2^n + 1)$ ,  $\lambda = 2(2^{n-1} + 1)(2^{n-3} - 1)$ ,  $\mu = 2^{n-1}(2^{n-2} - 1)$ .*

*Proof.* The number of vertices is  $v = 2^{n-1}(2^n - 1)$  and from Lemma 2.3 d) we have that  $k = (2^{n-2} - 1)(2^n + 1)$ . Let  $x, y \in \mathcal{R}$  such that  $x$  is not adjacent to  $y$ . Then  $y \in (x^\tau)^\perp \setminus \{x^\tau\}$ . We have that  $\mu = |x^\perp \cap y^\perp \cap \mathcal{R}|$  and  $x^\perp \cap y^\perp$  is an elliptic quadric  $\mathcal{Q}^-(2n-1, 2)$  meeting  $\mathcal{Q}$  in the cone  $P\mathcal{Q}^-(2n-3, 2)$ , where  $P = x^\tau y \cap \mathcal{Q}$ . There are three hyperplanes through  $x^\perp \cap y^\perp$ , namely,  $x^\perp$ ,  $y^\perp$  and  $T^\perp$ , where  $T = \Sigma \cap xy$  and  $\Sigma = \langle \mathcal{Q} \rangle$ . Then  $T^\perp$  meets  $\mathcal{Q}$  in a cone  $P\mathcal{Q}(2n-2, 2)$  containing  $P\mathcal{Q}^-(2n-3, 2)$ . Hence, from Lemma 2.3 c), d),  $|T^\perp \cap \mathcal{R}| = 2^{2n-2}$  and  $|x^\perp \cap \mathcal{R}| = |y^\perp \cap \mathcal{R}| = 2^{n-2}(2^n + 1) - 2^n$ . It follows that

$$\mu + 2(2^{n-2}(2^n + 1) - 2^n - \mu) + 2^{2n-2} - \mu = 2^{n-1}(2^n - 1)$$

and hence  $\mu = 2^{n-1}(2^{n-2} - 1)$ . Let  $x, y \in \mathcal{R}$  such that  $x$  is adjacent to  $y$ . Then  $y \in x^\perp \cap \mathcal{R} \setminus \{x\}$ . We have that  $\lambda = |x^\perp \cap y^\perp \cap \mathcal{R} \setminus \{x, y\}|$  and  $x^\perp \cap y^\perp$  is a cone  $\ell\mathcal{Q}^-(2n-3, 2)$ , where  $\ell = xy$ , meeting  $\mathcal{Q}$  in the cone  $T\mathcal{Q}^-(2n-3, 2)$ , with  $T = \ell \cap \mathcal{Q}$ . There are three hyperplanes through  $x^\perp \cap y^\perp$ , namely,  $x^\perp$ ,  $y^\perp$  and  $T^\perp$ . Hence, from Lemma 2.3 d), f),  $|T^\perp \cap \mathcal{R}| = 2^{n-1}(2^{n-1} - 1)$  and  $|x^\perp \cap \mathcal{R}| = |y^\perp \cap \mathcal{R}| = 2^{n-2}(2^n + 1) - 2^n$ . It follows that

$$\lambda + 2 + 2(2^{n-2}(2^n + 1) - 2^n - \lambda - 2) + (2^{n-1}(2^{n-1} - 1) - \lambda - 2) = 2^{n-1}(2^n - 1)$$

and hence  $\lambda = 2^{2n-3} - 2^n + 2^{n-2} - 2 = 2(2^{n-1} + 1)(2^{n-3} - 1)$ .  $\square$

**Remark 4.4.** The graph  $\mathcal{G}$  arising from the relative hemisystem of  $\mathcal{Q}^-(9, 2)$  constructed in Theorem 3.3 has the same parameters of the graph constructed in [10] by considering the elliptic sections of a parabolic quadric of  $\text{PG}(4, 4)$ . With the aid of Magma [3] we checked that the two graphs have the same automorphism group isomorphic to  $2 \cdot \text{PSp}(4, 4)$ . Hence it is plausible that the two graphs are isomorphic.

## References

- [1] J. Bamberg, M. Lee, A relative  $m$ -cover of a Hermitian surface is a relative hemisystem, *arXiv 1608.03055v1*.

- [2] J. Bamberg, S. Kelly, M. Law, T. Penttila, Tight sets and  $m$ -ovoids of finite polar spaces, *J. Combin. Theory Ser. A* 114 (7) (2007), 1293-1314.
- [3] J. Cannon, C. Playoust, *An introduction to MAGMA*, University of Sidney, Sidney, Australia, 1993.
- [4] A. Cossidente, Relative hemisystems on the Hermitian surface, *J. Algebraic Combin.* 38 (2013) 2, 275-284.
- [5] A. Cossidente, A new family of relative hemisystems on the Hermitian surface, *Des. Codes Cryptogr.* 75 (2015) 2, 213-221.
- [6] A. Cossidente, F. Pavese, Intriguing sets of  $\mathcal{W}(5, q)$ ,  $q$  even, *J. Combin. Theory Ser. A* 127 (2014), 303-313.
- [7] A. Cossidente, T. Penttila, Hemisystems on the Hermitian surface, *J. Lond. Math. Soc.* 72 (2005), 731-741.
- [8] J. W. P. Hirschfeld, *Finite Projective Spaces of Three Dimensions*, Oxford University Press, Oxford, 1991.
- [9] R.H. Dye, On the maximality of the orthogonal groups in the symplectic groups in characteristic two, *Math. Z.* 172 (1980) 3, 203-212.
- [10] X. Hubaut, R. Metz, A class of strongly regular graphs related to orthogonal groups, *Combinatorics 1981 (Rome, 1981)*, 469-472, *Ann. Discrete Math.*, 18, North-Holland, Amsterdam-New York, 1983.
- [11] S. Kelly, Constructions of intriguing sets of polar spaces from field reduction and derivation, *Des. Codes Cryptogr.* 43 (2007) 1, 1-8.
- [12] T. Penttila, J. Williford, New families of  $Q$ -polynomial association schemes, *J. Combin. Theory Ser. A* 118 (2011), 502-509.
- [13] B. Segre, Forme e geometrie Hermitiane con particolare riguardo al caso finito, *Ann. Mat. Pura Appl.* 70 (1965), 1-201.
- [14] J.A. Todd, As it might have been, *Bull. London Math. Soc.* 2 (1970) 4, 1-4.