

SCExAO and GPI *YJH* Band Photometry and Integral Field Spectroscopy of the Young Brown Dwarf Companion to HD 1160

E. Victor Garcia^{1,2,36}, Thayne Currie³, Olivier Guyon^{3,4,5}, Keivan G. Stassun^{2,6}, Nemanja Jovanovic³, Julien Lozi³, Tomoyuki Kudo³, Danielle Doughty⁷, Josh Schlieder⁸, J. Kwon⁹, T. Uyama¹⁰, M. Kuzuhara^{4,11,12}, J. C. Carson¹³, T. Nakagawa⁹, J. Hashimoto^{4,11}, N. Kusakabe^{4,11}, L. Abe¹⁴, W. Brandner¹⁵, T. D. Brandt^{16,17}, M. Feldt¹⁵, M. Goto¹⁸, C. A. Grady^{19,20,21}, Y. Hayano³, M. Hayashi¹¹, S. S. Hayashi^{3,22}, T. Henning¹⁴, K. W. Hodapp²³, M. Ishii¹¹, M. Iye¹¹, M. Janson²⁴, R. Kandori¹¹, G. R. Knapp²⁵, T. Matsuo²⁶, M. W. McElwain¹⁸, S. Miyama²⁷, J.-I. Morino¹¹, A. Moro-Martin^{28,29}, T. Nishimura³, T.-S. Pyo³, E. Serabyn³⁰, T. Suenaga^{10,22}, H. Suto^{4,11}, R. Suzuki¹¹, Y. H. Takahashi^{11,30}, H. Takami³¹, M. Takami³¹, N. Takato³, H. Terada¹¹, C. Thalmann³², E. L. Turner^{25,33}, M. Watanabe³⁴, J. Wisniewski³⁵, T. Yamada⁹, T. Usuda¹¹, and M. Tamura^{4,10,11}

-
- ¹Lowell Observatory, Flagstaff, AZ, 86001, USA; eugenio.v.garcia@gmail.com
- ²Department of Physics and Astronomy, Vanderbilt University, Nashville, TN
- ³⁶Lawrence Livermore National Laboratory, L-210, 7000 East Avenue, Livermore, CA 94550, USA
- ³Subaru Telescope, National Astronomical Observatory of Japan, 650 N. A'ohoku Place, Hilo, HI 96720, USA
- ⁴Astrobiology Center of NINS, 2-21-1 Osawa, Mitaka, Tokyo, 181-8588, Japan
- ⁵Steward Observatory, University of Arizona, 933 N Cherry Ave., Tucson, AZ 85721, USA
- ⁶Department of Physics, Fisk University, Nashville, TN 37208, USA
- ⁷College of Optical Sciences, University of Arizona, Tucson, AZ 85721
- ⁸NASA Postdoctoral Program Fellow, NASA Ames Research Center, Moffett Field, CA
- ⁹Department of Space Astronomy and Astrophysics, Institute of Space & Astronautical Science (ISAS), Japan Aerospace Exploration Agency (JAXA), 3-1-1 Yoshinodai, Chuo-ku, Sagamihara, Kanagawa 252-5210, Japan
- ¹⁰Department of Astronomy, University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan
- ¹¹National Astronomical Observatory of Japan, 2-21-1, Osawa, Mitaka, Tokyo, 181-8588, Japan
- ¹²Department of Earth and Planetary Sciences, Tokyo Institute of Technology, Ookayama, Meguro-ku, Tokyo, 152-8551, Japan
- ¹³Department of Physics and Astronomy, College of Charleston, 58 Coming St., Charleston, SC 29424, USA
- ¹⁴Laboratoire Lagrange (UMR 7293), Université de Nice-Sophia Antipolis, CNRS, Observatoire de la Côte d'Azur, 28 Avenue Valrose, F-06108 Nice Cedex 2, France
- ¹⁵Max Planck Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany
- ¹⁶Astrophysics Department, Institute for Advanced Study, Princeton, NJ, USA
- ¹⁷NASA Sagan Fellow
- ¹⁸Universitäts-Sternwarte München, Ludwig-Maximilians-Universität, Scheinerstr. 1, 81679 München, Germany
- ¹⁹Exoplanets and Stellar Astrophysics Laboratory, Code 667, Goddard Space Flight Center, Greenbelt, MD 20771, USA
- ²⁰Eureka Scientific, 2452 Delmer, Suite 100, Oakland, CA 96002, USA
- ²¹Goddard Center for Astrobiology
- ²²Department of Astronomical Science, SOKENDAI (The Graduate University for Advanced Studies), 2-21-1 Osawa, Mitaka, Tokyo, 181-8588, Japan
- ²³Institute for Astronomy, University of Hawaii, 640 N. A'ohoku Place, Hilo, HI 96720, USA

ABSTRACT

We present high signal-to-noise ratio, precise YJH photometry and Y band ($0.957 - 1.120 \mu\text{m}$) spectroscopy of HD 1160 B, a young substellar companion discovered from the Gemini NICI Planet Finding Campaign, using the Subaru Coronagraphic Extreme Adaptive Optics instrument and the Gemini Planet Imager. HD 1160 B has typical mid-M dwarf-like infrared colors and a spectral type of $M5.5_{-0.5}^{+1.0}$, where the blue edge of our Y band spectrum rules out earlier spectral types. Atmospheric modeling suggests HD 1160 B having an effective temperature of $3000\text{--}3100 \text{ K}$, a surface gravity of $\log g = 4\text{--}4.5$, a radius of $1.55 \pm 0.10 R_{\text{J}}$, and a luminosity of $\log L/L_{\odot} = -2.76 \pm 0.05$. Neither the primary’s Hertzsprung-Russell diagram position nor atmospheric modeling of HD 1160 B show evidence for a sub-solar metallicity. The interpretation of the HD 1160 B depends on which stellar system components are used to estimate an age. Considering HD 1160 A, B and C jointly, we derive an age of $80\text{--}125 \text{ Myr}$, implying that HD 1160 B straddles the hydrogen-burning limit ($70\text{--}90 M_{\text{J}}$). If we consider HD 1160 A alone, younger ages ($20\text{--}125 \text{ Myr}$) and a brown dwarf-like mass ($35\text{--}90 M_{\text{J}}$) are

²⁴Department of Astronomy, Stockholm University, AlbaNova University Center, SE-10691, Stockholm, Sweden

²⁵Department of Astrophysical Science, Princeton University, Peyton Hall, Ivy Lane, Princeton, NJ 08544, USA

²⁶Department of Earth and Space Science, Graduate School of Science, Osaka University, 1-1 Machikaneyama, Toyonaka, Osaka, 560-0043, Japan

²⁷Hiroshima University, 1-3-2, Kagamiyama, Higashihiroshima, Hiroshima, 739-8511, Japan

²⁸Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

²⁹Center for Astrophysical Sciences, Johns Hopkins University, Baltimore, MD 21218, USA

³⁰Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, 171-113, USA

³¹Institute of Astronomy and Astrophysics, Academia Sinica, P.O. Box 23-141, Taipei 10617, Taiwan

³²Swiss Federal Institute of Technology (ETH Zurich), Institute for Astronomy, Wolfgang-Pauli-Strasse 27, CH-8093 Zurich, Switzerland

³³Kavli Institute for Physics and Mathematics of the Universe, The University of Tokyo, 5-1-5, Kashiwanoha, Kashiwa, Chiba, 277-8568, Japan

³⁴Department of CosmoSciences, Hokkaido University, Kita-ku, Sapporo, Hokkaido, 060-0810, Japan

³⁵H. L. Dodge Department of Physics & Astronomy, University of Oklahoma, 440 W Brooks St Norman, OK 73019, USA

possible. Interferometric measurements of the primary, a precise GAIA parallax, and moderate resolution spectroscopy can better constrain the system’s age and how HD 1160 B fits within the context of (sub)stellar evolution.

1. Introduction

In addition to about a dozen superjovian-mass extrasolar planets (e.g. Marois et al. 2008; Lagrange et al. 2010; Kraus and Ireland 2012; Rameau et al. 2013; Quanz et al. 2013; Kuzuhara et al. 2013; Carson et al. 2013; Currie et al. 2014a, 2015; Macintosh et al. 2015), the past decade of ground-based *direct imaging* observations of young stars have also revealed numerous higher mass (ratio) and (often) wider separation brown dwarf companions (Chauvin et al. 2005; Biller et al. 2010; Ireland et al. 2011). Their typically more modest companion-to-primary contrasts make their detection and precise photometric/spectroscopic calibration possible without having to correct for throughput/signal losses inherent in the algorithms needed to detect more extreme contrast-ratio exoplanets (e.g. Marois et al. 2010). They, and free-floating brown dwarfs of comparable ages, can reveal general trends in infrared colors and spectra that clarify the properties of young substellar atmospheres, can serve as important tests of the same atmosphere models used to characterize bona fide imaged exoplanets, and may provide insights into the formation of substellar objects (e.g. Faherty et al. 2016; Kratter et al. 2010; Reggiani and Meyer 2011).

One particularly interesting system with a brown dwarf companion is HD 1160, located at a distance of 103.1 ± 5.3 pc (van Leeuwen 2007) with an A0V primary that is comparable in mass to at least some stars around which planets have been imaged (e.g. β Pic). Targeted by the NICI Planet-Finding Campaign (Liu et al. 2010), HD 1160 is orbited by a pair of substellar companions: HD 1160 B at ~ 80 au and HD 1160 C at 530 au (Nielsen et al. 2012, hereafter N12).

N12’s photometry for HD 1160 B and spectrum for HD 1160 C suggested that the former is a $L0 \pm 2$ brown dwarf and the latter a M3.5 star. N12 used HD 1160 B’s and C’s red $J - K_s$ colors, HD 1160 A’s luminosity, and the system’s large space motion to argue that all components are young: ~ 10 – 100 Myr. By comparing HD 1160 B and C’s derived luminosity to predictions from substellar cooling models given the system’s age, N12 derived a mass of $24 - 45 M_J$ for HD 1160 B and 0.18 – $0.25 M_\odot$ for HD 1160 C.

HD 1160 B was revisited as part of the first results from the Spectro-Polarimetric High-contrast Exoplanet REsearch (SPHERE, Beuzit et al. 2008) at the Very Large Telescope (VLT) by Maire et al. (2016, hereafter M16). M16 observed HD 1160 B with VLT/SPHERE

to obtain the first spectrum for the object, with resolution of $R \approx 30$ covering the wavelength range of $1.0 - 1.6 \mu\text{m}$. Based on HD 1160 B’s spectral shape, M16 re-classify its spectral type as $M6_{-0.5}^{+1}$ and derive an effective temperature of $3000 \pm 100 \text{ K}$ but do not constrain the object’s surface gravity their SPHERE YJH spectrum. Based largely on a wider range and systematically older age estimate ($30 - 300 \text{ Myr}$), they estimate a mass of $39 - 144 M_J$ based on bolometric luminosity and $69 - 172 M_J$ based on effective temperature.

Two key questions surround the nature of HD 1160’s companions. The first is metallicity. By comparing HD 1160 A’s Hertzsprung-Russel (HR) diagram position to predictions from the Siess et al. (2000) isochrones, N12 found that the primary is only matched to isochrones with a substantially subsolar metallicity ($[\text{Fe}/\text{H}] = -0.3$, in conflict with the very red near-infrared colors of HD 1160 B and C. While M16 derived bluer infrared colors for HD 1160 B and C, they found atmosphere models assuming a subsolar metallicity fit HD 1160 B’s spectrophotometry the best. As most stars in the solar neighborhood have a near-solar metallicity, new analysis is needed to clarify whether or not HD 1160 A has a non-solar metallicity. *The atmospheric properties of HD 1160 B, including metallicity, could be clarified with new spectrophotometry with greater precision covering a wider wavelength baseline.*

The second question is the system’s age. Different age estimates for HD 1160 result in quite different inferred masses for the two low-mass companions. HD 1160 B’s interpretation changes from a low-mass, high mass ratio brown dwarf (for a young age) to an object straddling the brown dwarf/stellar boundary (for an older age). *The age of the HD 1160 system could be better clarified with a more in-depth and systematic analysis of possible age indicators for the primary and the two companions.*

In this work, we present high-precision photometry of the HD 1160 system obtained with the Subaru Coronagraphic Extreme Adaptive Optics (SCEXAO) project in the YJH passbands and integral field spectroscopy with the Gemini Planet Imager (GPI) in the Y band (§2). We compare HD 1160 B’s spectrum and new photometry to that of young brown dwarfs and field objects and match its photometry/spectra using the BT-Settl atmosphere models (§3) to address whether or not HD 1160 B clearly shows evidence for a subsolar metallicity and derive an updated luminosity for both it and HD 1160 C. We use an array of age estimates to better clarify the age of the system (§4). After modeling HD 1160 B’s atmosphere and better constraining HD 1160 C’s luminosity, we use our updated age estimate to more precisely infer HD 1160 BC’s mass (§5). Section 6 summarizes our findings and identifies future work that can better clarify the properties of this system.

2. Data

2.1. Subaru Coronagraphic Extreme Adaptive Optics *YJH* Imaging

The Subaru Coronagraphic Extreme Adaptive Optics (SCEExAO) is a high contrast imaging platform designed for the discovery and characterization of faint companions (Jovanovic et al. 2015). A partial correction of the low order modes of the wavefront is performed by the Subaru facility adaptive optics instrument (AO188), and a final correction, including the high-order modes is performed by a visible-light pyramid wavefront sensor and a 2000-element deformable mirror inside SCEExAO. Ultimately, the project is slated to yield 90+% Strehl ratios at 1.6 μm on bright stars ($I(\text{mag}) < 9\text{--}10$).

On 31 October 2015, we observed HD 1160 with SCEExAO coupled with the near infrared HiCIAO camera (Tamura 2009) in the *Y*, *J*, and *H* broadband filters ($\sim 8.0 \text{ mas pixel}^{-1}$)¹. All observations were obtained in angular differential imaging (ADI) mode (Marois et al. 2006) but with a very small amount of field rotation ($\Delta\text{P.A.} \sim 1.5\text{--}2^\circ$).

While our instantaneous Strehl ratios were in the 80–90% range, strong low frequency vibrations induced by the telescope drive system degraded AO correction quality to $\sim 60\text{--}70\%$ Strehl over the course of $> 0.1\text{s}$ observations. These vibrations are now partially mitigated by a linear-quadratic-gaussian (LQG) controller, yielding slightly improved performance (Lozi et al. 2016). Upcoming improvements – using the higher-stroke AO188 deformable mirror driven by multiple accelerometers to offload tip-tilt variations – should eliminate these vibrations. Nevertheless, the partial correction offered by SCEExAO substantially improves upon the image quality achieved with discovery images of HD 1160 B obtained in the near-infrared (NIR) from Gemini/NICI.

Our observing log is detailed in Table 1. We cycled through the three near-infrared passbands using a combination of short, lightly saturated exposures and unsaturated images obtained using HiCIAO’s neutral density filters. Basic image processing followed standard methods employed in Currie et al. (2011, 2014a) for flat fielding, dark subtraction, destriping, radial profile subtraction, and combining images. As HD 1160 B/C were plainly visible in the raw data, we did not perform any point-spread function (PSF) subtraction but instead simply removed the azimuthally-averaged seeing halo.

Despite the short integration times and no PSF subtraction techniques applied, both HD 1160 B and C were easily recovered with a signal-to-noise ratio (SNR) of ~ 70 and > 200 ,

¹The plate scale when using AO188 and HiCIAO is $9.5 \text{ mas pixel}^{-1}$, but for SCEExAO and HiCIAO, the plate scale is $8.3 \text{ mas pixel}^{-1}$.

respectively, in YJH passbands (Figure 1). While processing the data with advanced PSF subtraction techniques like A-LOCI or KLIP (Currie et al. 2015; Soummer et al. 2012) yielded stronger detections and lower residuals at smaller separations, they also partially attenuate the companion signal. For simplicity, we analyze only the halo-subtracted images which lack companion signal loss.

Our SCEXAO photometry of HD 1160 B are shown in the left columns of Table 2: $m(Y, J, H) = 15.30 \pm 0.04$, 14.69 ± 0.05 , and 14.21 ± 0.02 . For HD 1160 C, we derive $m(Y, J, H) = 13.74 \pm 0.02$, 13.37 ± 0.02 , and 12.81 ± 0.02 . Our errors consider (in quadrature) the SNR of the detection (computed as in Currie et al. 2015), uncertainties in the primary magnitude, and a minor contribution of uncertainties in the attenuation of the neutral density filter.

We use aperture photometry to measure the brightness of HD 1160 A (in unsaturated images obtained with the neutral density filter) and HD 1160 BC. For the fainter HD 1160 B companion, we derived the contrast between it and the primary using a range of aperture diameters comparable to the FWHM of a point source in each passband. For the contrast of the brighter, wider-separation HD 1160 C component, we used larger apertures enclosing much more ($> 90\%$) of the primary and companion signals to guard against flux loss due to PSF degradation (e.g. vignetting of the field).

Our SCEXAO HD 1160 B J and H photometry shows good agreement with SPHERE measurements (Table 2, third column). SPHERE’s brightness estimates are systematically brighter by 1.8σ (0.22 ± 0.12 mag) at J -band and $\approx 2.8\sigma$ (0.23 ± 0.08 mag) at H -band. A good portion of the discrepancy at H -band is due to SPHERE’s spectrum only covering half the H -band filter. Small differences at J band may be due to SPHERE’s flux calibration. SPHERE’s spectro-photometry had to be corrected for signal loss due to aggressive processing; while the HiCIAO neutral density filter is flat across the bandpass, SPHERE’s exhibits a slight wavelength dependent attenuation at J band.

In contrast, our photometry strongly disagrees with the measurements of N12 by 1.14 mag at J and 0.44 at H (Table 2, columns three and six)². Similar discrepancies between NICI photometry and other instruments (Boccaletti et al. 2013) have been observed previously. The difference may be due to inaccurate flux normalization, possibly from mismeasured attenuation through the coronagraphic mask, or an inaccurate subtraction of the sky

²For another estimate of HD 1160 B’s J -band photometry, we downloaded and reduced archival J -band Keck/NIRC2 data for HD 1160 from 2004-07-14 (P.I. I. Song) using image processing methods identical to those described in this work. Our Keck detection has a substantially lower SNR than SCEXAO and SPHERE. However, it agrees with the SCEXAO and SPHERE photometry against the NICI photometry: $m_J = 14.54 \pm 0.24$.

background annulus surrounding the companion (as was the case for early photometry of ROXs 42B “c”, see Currie et al. 2014b).

2.2. Gemini Planet Imager Y band Low-Resolution Spectroscopy

2.2.1. Observations and Basic Processing

HD 1160 B was observed with the *Gemini Planet Imager* (Macintosh et al. 2006, 2014) (GPI) in Y band ($0.957 - 1.120 \mu\text{m}$, $R \approx 37$) on 17 November 2013 in $\sim 1''$ seeing by the GPI Verification and Commissioning team (Table 1). The observations consist of nine 88.7 s exposures in IFS mode (Larkin et al. 2014) using GPI’s apodized pupil Lyot coronagraph with a pixel scale of $14.14 \pm 0.01 \text{ mas pixel}^{-1}$ (Konopacky et al. 2014). As shown in Figure 2, the GPI Y band filter extends $0.1 \mu\text{m}$ past typical Y band filters (See the appendix of Liu et al. 2012). It also extends blueward of the SPHERE Y band data from M16, thus expanding the wavelength range at which HD 1160 B can be detected and analyzed beyond that offered by either SPHERE or SCExAO observations. The observations were obtained in ADI mode. However, like the SCExAO observations, the GPI sequence covered little parallactic motion ($\Delta\text{P.A.} \sim 4.3^\circ$).

We processed the images using the GPI data reduction pipeline version 1.3.0 (Maire et al. 2010, 2012; Perrin et al. 2014). The pipeline requires the location and spectral solution for every lenslet on the HAWAII-2RG detector. These lenslet locations were determined by using a cross correlation between the deep argon calibration source images available on the GPI public webpage³. The elevation of the telescope differed between the images of HD 1160 B and the daytime argon calibration lamp sequence. We calculated a shift to determine the overall change of the wavelength solution between the daytime calibrations and that appropriate for the observations of HD 1160 B.

We further used the GPI data reduction pipeline to apply dark corrections, remove bad pixels, track satellite spot locations, convert each microspectra into a 37 channel spectral cube ($0.957 - 1.120 \mu\text{m}$), correct for distortion using the publicly available distortion solution¹, and correct for atmospheric differential refraction. Each of the 9 individual raw observations were processed in an identical way. HD 1160 B is visible in individual spectral channels of each data cube (Figure 3).

³<http://www.gemini.edu/sciops/instruments/gpi/public-data>

2.2.2. Extraction of HD 1160 B Spectrum and Calibration

The final, distortion-corrected, de-rotated, time-averaged data cube of HD 1160 has a smooth, slowly-varying halo at HD 1160 B’s angular separation (left panel, Figure 3). We remove this slow-varying background using a median high pass filter (right panel, Figure 3) with a box size of 11 pixels ($\approx 5\lambda/D$). We manually confirmed that this filtering did not result in a measurable loss in signal. Our final data cube has an SNR ~ 90 detection for HD 1160 B in the wavelength-collapsed cube and SNR > 10 for each channel.

To flux-calibrate our final data cube and extract HD 1160 B’s flux-calibrated spectrum, we again followed steps outlined in the GPI data reduction pipeline. First, we extracted the flux-calibrated spectra of the four satellite spots using the laboratory measured satellite-to-star flux ratio of 2.345×10^{-4} from Wang et al. (2014) and assumed an A0V template spectrum from the Pickles stellar library (Pickles 1998) to characterize the host star spectrum of HD 1160 A (N12)⁴. We adopted a 2-pixel aperture radius, a 5–20 pixel annulus to define the sky background, and the standard deviation of the four satellite spot flux density measurements as the uncertainty in absolute calibration in each spectral channel⁵.

To extract the spectrum of HD 1160 B, we used the same aperture radius and background annulus used to extract the satellite spot spectra. The errors in the extracted and flux-calibrated HD 1160 B spectrum draw from the uncertainty in the spot calibration and the uncertainty in the background annulus signal surrounding HD 1160 B⁶. We trimmed the HD 1160 B spectrum of the first and last few channels ($< 0.953 \mu\text{m}$ and $> 1.12 \mu\text{m}$), which had absolute calibration uncertainties $\gtrsim 10\%$. The resulting flux calibrated spectrum spanning $0.957 - 1.120 \mu\text{m}$ is shown in Figure 4. From the wavelength-collapsed data cube, we estimate a *Y* band magnitude of $m(Y) = 15.13 \pm 0.21$, consistent with our more precise SCEXAO measurements (Table 2, third column).

⁴HD 1160 A is a well-known IR photometric standard star (Elias et al. 1982).

⁵The upper-left spot spectrum contributed the most to the uncertainty in each spectral channel and in the wavelength-collapsed image, which is $\sim 20\%$ fainter than the average of the other three spots.

⁶For a separate spectral extraction independent of the GPI data reduction pipeline, we performed aperture photometry using `aper.pro`⁷ at the location of the companion, for each wavelength slice, assuming the background to be zero. The two methods show strong agreement, to within the calculated errors/channel.

2.3. Rereduced Keck/NIRC2 M' Photometry

M16 found significant discrepancies between their SPHERE/IRDIS photometry and that from N12 in the J and H passbands. Additionally, even after adopting the revised HD 1160 B NaCo L' photometry from M16 (~ 0.2 mag fainter), the implied $L' - M'$ color is ~ -0.7 , characteristic of L/T transition objects but far too blue for a mid M dwarf like HD 1160 B (see Galicher et al. 2011).

Therefore, we rereduced the same archival Keck/NIRC2 M' data first analyzed by N12 (HD 1160 C was outside of the field of view). Basic processing followed standard steps we have previously used for thermal IR data (e.g. Currie et al. 2011) but without PSF subtraction. Briefly, after applying a linearity correction to the data, we constructed a sky frame for a given image from the 5 nearest (in time) images where the star is in a different dither position. After sky subtraction, we corrected for distortion, registered each image to a common center, subtracted off the radial intensity profile for each image, averaged the set of profile-subtracted images using $3\text{-}\sigma$ outlier rejection, and applied a $5 \lambda/D$ moving-box median filter to the combined image. As the combined image shows some residuals of the PSF halo, we computed the SNR of the detection in a conservative fashion (not assuming that the image at HD 1160 B's location is photon noise dominated) as we have typically done with high-contrast imaging data sets (e.g. Currie et al. 2011).

Our re-reduction yields a SNR ~ 4.6 detection of HD 1160 B with a measured apparent magnitude of $m(M') = 13.81 \pm 0.24$ (Table 2, far-right column), where the intrinsic SNR of the detection contributes almost all of the photometric uncertainty. HD 1160 B is about 63% brighter at M' than reported in N12. The implied $L'-M'$ color of $\sim -0.21 \pm 0.23$ is in agreement with expected colors for mid M to early L objects (Galicher et al. 2011).

3. Characterization of HD 1160 B and C

3.1. Analysis of the HD 1160 B and C Photometry

To compare HD 1160 B and C to field MLT dwarfs, we adopt our SCEXAO YJH photometry. We assume a distance of 103.1 ± 5.3 pc (van Leeuwen 2007), yielding absolute magnitudes of $Y=10.23 \pm 0.11$, $J=9.63 \pm 0.10$ and $H=9.14 \pm 0.10$ mag. Figure 5 compares our HD 1160 B photometry to the sequence of MLT dwarfs from Dupuy and Liu (2012), and M-dwarf standards of Kirkpatrick et al. (2010) and references therein. We plot YJH colors of $J - H=0.48 \pm 0.05$, $Y - J=0.61 \pm 0.06$ colors and the absolute magnitudes in Y , J , passbands for HD 1160 B as the blue star in Figure 5. In all cases, HD 1160 B's photometric

points are consistent with the field dwarfs sequence. We find HD 1160 B’s colors in agreement with the M5–M6 spectral standards of Kirkpatrick et al. (2010) (transition of green and red diamonds in Figure 5). Similarly, as shown by the pink star in Figure 5, we find the near-IR colors and the YJH absolute magnitudes of HD 1160 C in agreement with M3–M4 spectral standards of Kirkpatrick et al. (2010).

As a further check, we compare HD 1160 B’s nominal colors to those if we adopt photometry extracted from the SPHERE spectrum in JH (Table 2) and photometry from partial coverage in Y band ($m(Y) = 15.19 \pm 0.02$; $M(Y)=10.12 \pm 0.02$). The SPHERE-derived photometry is slightly redder in $Y - J$ but otherwise suggests that HD 1160 B is consistent with the field sequence (pink and blue triangles, Figure 5).

3.2. Analysis of the HD 1160 B Spectrum

To compare HD 1160 B’s full near-infrared spectrum to that from other objects, we construct a merged YJH spectrum from GPI and SPHERE data. We applied an offset to the GPI spectrum such that its photometry integrated over Y band perfectly matched the SCExAO Y band photometry and a separate offset to the SPHERE spectrum such that its band-integrated photometry matched that of SCExAO’s as well. We adopt the GPI (SPHERE) measurements shortwards (longwards) of $1.12 \mu m$.

As implied by N12 and M16 and shown in §4 of this paper, HD 1160’s age is likely intermediate between that of low mass members of young star forming regions such as Upper Sco (11 ± 2 Myr, Pecaut et al. 2012) and older field objects (> 1 Gyr). For spectral typing HD 1160 B, we therefore compare the GPI+SPHERE spectrum to field M dwarf spectral standards from the SPEX library (Rayner et al. 2003) and Upper Sco members (Dawson et al. 2014).

We perform the spectral typing by binning the template NIR spectra to the resolution of the GPI and SPHERE spectra and computing the χ^2 as:

$$\chi^2 = \sum_{i=1}^{i=57} \frac{(F_{i,\text{obs}} - F_{i,\text{template}})^2}{\sigma_{i,\text{obs}}^2 + \sigma_{i,\text{template}}^2} \quad (1)$$

where flux F is in watts $m^{-2} \mu m^{-1}$ and index i corresponds to one of the 57 spectral channels of the flux calibrated GPI and SPHERE spectra. We weighted the GPI and SPHERE spectra equally.

As shown in Figure 6, the GPI and SPHERE spectrum shape is poorly reproduced for Upper Sco M8 member 2M16101–28563 for spectral channels $< 1.0 \mu m$ and $1.3\text{--}1.4 \mu m$.

The disagreement is greater for L0.8 Upper Sco member 2M16195–28322 (top of Figure 6). On the other hand, M5.2–M6.8 Upper Sco members have $\Delta\chi^2$ within the 95% confidence interval for 57 degrees of freedom (see Numerical Recipes, Press et al. 1992). We therefore find the spectral type of HD 1160 B as compared to Upper Sco Members to be $M6_{-0.8}^{+0.8}$.

Similarly, Figure 7 compares the GPI and SPHERE spectra of HD 1160 B to M2–M8 field SPEX spectra standards from Kirkpatrick et al. (2010). We find that the overall spectral shape is best reproduced by M5 standard 2M01532+36314 or M6 standard 2M13272+09464. The overall shape of the spectrum is poorly reproduced by the linear slope of M4 or earlier standards across 0.95–1.6 μm . Standard spectra later than M6 poorly reproduce the smooth portion of the HD 1160 B spectrum at 1.3–1.4 μm as well as the slope at $< 1.0 \mu\text{m}$. Therefore, we find the spectral type of HD 1160 B as compared to field M dwarfs to be $M5.5_{-0.5}^{+0.5}$, in agreement with the spectral types derived for HD 1160 B compared to Upper Sco members above. We adopt a spectral type for HD 1160 B of $M5.5_{-0.5}^{+1.0}$ as an average between the dwarf and Upper Sco spectral types, finding agreement with the M16 spectral type of $M6_{-0.5}^{+1.0}$.

3.3. Atmospheric Modeling of HD 1160 B

Atmosphere Models Considered. To explore HD 1160 B’s atmospheric properties, we compare the object’s broadband photometry and *YJH* GPI+SPHERE spectrum to predictions from planet atmosphere models, adopting a range of effective temperatures and surface gravities. We consider the BT-Settl 2013 models of Allard et al. (2012) with temperatures of 2500–3500 K, surface gravities of $\log g = 3.5–5.5$ dex and solar metallicity. Using the same atmosphere models, M16 concluded that spectral energy distribution (SED) modeling favors a subsolar metallicity for HD 1160 B. Using revised/new photometry and spectroscopy, we assess whether or not HD 1160 B may yet have a solar metallicity.

Fitting Method. Following previous approaches (e.g. Currie et al. 2014b), we fit our *YJH* (SCEXAO, this work) K_s (SPHERE, M16) L' (NaCo, M16) M' (Keck, this work) photometry and *YJH* GPI+SPHERE spectroscopy. To model photometry, we compare the measured flux densities with predicted ones from the model spectra convolved with the MKO filter functions. The BT-Settl grid has significantly higher spectral resolution than the data ($R \approx 37$, $\text{SNR} \gg 10$ per channel). Thus, to model the spectrum of HD 1160 B, we rebin the BT-Settl spectra to the resolution of our extracted spectrum. We treat the planet radius as a free parameter, varying it from $R \sim 0.75 R_J$ to $R \sim 2.25 R_J$, which spans the wide range in radii predicted from models (Baraffe et al. 2003).

We identify the set of BT-Settl models consistent with the spectra or photometry at the

95% confidence limits given the number of degrees of freedom. We adopt the measured photometric error, listed in Table 2, for model fitting. We incorporate our distance uncertainty of ± 5.3 pc (van Leeuwen 2007) in the uncertainty estimate for the planet radius. We do not fit for T_{eff} and $\log g$ parameters via χ^2 minimization; rather, we compare the χ^2 to a grid of atmosphere models with pre-determined T_{eff} and $\log g$, treating the radius as a scaling parameter. Given six photometric points and four degrees of freedom, the 95% cutoff is $\chi_{lim}^2 = 9.49$. Given 57 spectral channels and 55 degrees of freedom, models with $\chi^2 \leq 73.3$ are consistent with the data.

Photometry and Spectroscopy Fitting Results. Table 6 summarizes our model-fitting results to the $YJHK_sL'M'$ photometry and $0.95 - 1.6 \mu\text{m}$ GPI+SPHERE spectroscopy⁸. Figures 8 and 9 display the χ^2 distribution for our model fits to the photometry and spectroscopy, respectively. Figure 10 displays an acceptably-fitting model compared to the $YJHK_sL'M'$ photometry and $0.95 - 1.6 \mu\text{m}$ spectroscopy.

For fitting the photometry alone, our best fit model has a reduced chi-square $\chi_{\text{red}}^2 = 0.63$ (d.o.f = $6 - 2 = 4$). Our 95% cutoff for photometry fits corresponds to surface gravities of $\log g = 3.5 - 5.5$ dex, and effective temperature of $T_{\text{eff}} = 2900 - 3500$ K. Our model fits to the photometry are rather degenerate given our small number (6) of photometry points and the similarity of broadband colors for objects at $T_{\text{eff}} = 2900 - 3500$ K.

However, combining fits to the photometry and spectroscopy significantly constrains the temperature and surface gravity of HD 1160 B. Our best-fit model for the spectrum has a reduced chi-square $\chi_{\text{red}}^2 = 0.92$ (d.o.f = $57 - 2 = 55$) with a surface gravity of $\log g = 4.5$ dex and effective temperature of $T_{\text{eff}} = 3000$ K. The solutions generally agree with M16 and cluster around $T_{\text{eff}} = 3000 - 3100$ K and $\log g = 4 - 4.5$, although we cannot rule out a 2900 K, $\log g = 4.5$ model nor a 3000 K, $\log g = 5.0$ model. We identify a suite of acceptably fitting models with a *solar* metallicity, not a *subsolar* metallicity. Thus, subsolar metallicities are not required to reproduce HD 1160 B’s spectrophotometry.

3.4. Inferred Properties of HD 1160 B from Best-fitting models

The BT-Settl models matching HD 1160 B’s photometry and spectroscopy allow us to infer a radius for the object. Compared to the default radius we used ($1.2 R_J$), our fits require

⁸The SPHERE and NICI photometry at K_s agree to within errors. As a separate check on these results, we downloaded and reduced K_s data from the Keck/NIRC2 archive and derived photometry that agreed with both measurements, within errors

radii larger by factors of $1.22 - 1.37\times$ and $0.98 - 1.42\times$ for the spectra and photometry respectively. The set of parameter space fitting photometry and spectra simultaneously covers $R = 1.55 \pm 0.10 R_J$ using our model fits to the HD 1160 B GPI+SPHERE spectrum, where errors in scaling the radius of our atmosphere models include the distance uncertainty error of $\approx 5\%$. HD 1160 B’s luminosity, from our best-fit effective temperature and radius from the spectroscopy, is $\log L/L_\odot = -2.76 \pm 0.05$ dex, in good agreement with $\log L/L_\odot = -2.81 \pm 0.10$ dex for HD 1160 B from M16.

3.5. Physical Characteristics of HD 1160 C

We adopt an effective temperature of $T_{\text{eff}} = 3260 \pm 100$ for HD 1160 C using the $M3.5 \pm 0.5$ SPEX/IRTF spectral type from N12. We obtain this T_{eff} by averaging the T_{eff} of an M3 and M4 5 – 30 Myr star from the relations of Pecaut and Mamajek (2013) (c.f. Table 5). Given that HD 1160 C is likely not 5 – 30 Myr, the spectral type- T_{eff} relations in Table 5 of Pecaut and Mamajek (2013) may not apply. The spectral type- T_{eff} relations for 50 – 300 Myr M-dwarfs (like HD 1160 C) is expected to lie in between the relations for young stars and field stars. As an additional check, using the spectral type- T_{eff} relations for field stars from Pecaut and Mamajek (2013) gives an effective temperature of $T_{\text{eff}} = 3300 \pm 100$ for HD 1160 C, which is within error of our adopted T_{eff} .

We use our J band absolute magnitude of 8.29 ± 0.10 mag and a bolometric correction of $BC_J = 1.875 \pm 0.035$ from the relations of Pecaut and Mamajek (2013) for young stars to estimate a bolometric magnitude of 10.16 ± 0.11 mag. We calculate the bolometric correction $BC_J = 1.875 \pm 0.035$ by averaging the bolometric corrections of a M3 and M4 star from Table 5 of Pecaut and Mamajek (2013). We compute a corresponding luminosity of $\log L/L_\odot = -2.16 \pm 0.04$ dex, using a bolometric magnitude of 4.755 for the Sun⁹, in fair agreement with $\log L/L_\odot = -2.05 \pm 0.06$ dex from M16.

4. Analysis of the HD 1160 System Age and Metallicity

4.1. Motivation

The age of the HD 1160 system is critical for interpreting the nature of the primary’s two companions. Furthermore, HD 1160 provides a tight constraint on evolutionary models:

⁹<https://sites.google.com/site/mamajeksstarnotes/basic-astronomical-data-for-the-sun>

the models should predict a common age given the observed effective temperatures and luminosities for three stars simultaneously. Given that HD 1160 A is on the main sequence, its age cannot be greater than the 300 Myr main-sequence lifetime of an A0 star (e.g. Siess et al. 2000). N12 derive a system age of 10 – 100 Myr, while M16 derive a much wider age estimate of 30 – 300 Myr.

Here, we re-investigate the age of the system based on the derived T_{eff} and $\log L/L_{\odot}$ for HD 1160 A, B and C. For stars close to the hydrogen burning limit and brown dwarfs just below it, recent results suggest difficulties in deriving ages and masses from HR diagram positions (Dupuy et al. 2009, 2010; Kraus et al. 2015; Dupuy et al. 2015, 2016), while ages derived from the upper main sequence are arguably preferable (Soderblom et al. 2014). Thus, we report two possible age ranges: one considering HD 1160 A alone and one considering all three components jointly.

Our derived system parameters are detailed in Table 4. We estimate the temperature and luminosity of HD 1160 A as follows. We fit the SED of HD 1160 A using the NextGen atmosphere models of Hauschildt et al. (1999). The photometry that we fit from the literature spans a wavelength range of 0.16–22 μm (Figure 11 and Table 5). Our best-fit model atmosphere implies an effective temperature of $T_{\text{eff}} = 9011 \pm 85$ K. The bolometric flux obtained by integrating the model SED provides an estimated $\log L/L_{\odot} = 1.12 \pm 0.07$.

4.2. Methodology

To derive the range of acceptable ages, we compare the system components’ luminosities and temperatures to evolutionary models. For HD 1160 A and C, we consider the Dartmouth evolutionary models (Dotter et al. 2008; Feiden et al. 2011), the Yonsei-Yale (Y^2) models (Yi et al. 2003; Spada et al. 2013), and the MESA Isochrones and Stellar Tracks (MIST), which use the Modules for Experimental Astrophysics (Paxton et al. 2011, 2013; Choi et al. 2016; Dotter 2016). We also consider the Baraffe15 models (Baraffe et al. 2015) for HD 1160 B and C. To arrive at a final age range for HD 1160 A and C, we take the median of the ranges derived from each individual evolutionary model comparison.

All evolutionary models assume solar composition and metallicity. Our atmospheric modeling (§3.3) suggests that HD 1160 B’s spectrophotometry is consistent with a solar metallicity atmosphere. Provided that other components of HD 1160 can match solar metallicity isochrones, there is no need to posit a subsolar metallicity for the system¹⁰.

¹⁰A substantially subsolar metallicity is unlikely for other reasons. Nearby young star clusters are nearly

Given that HD 1160 A has an unknown rotational velocity $v \sin i$, we also consider the effect of rotation on the derived ages. Rapidly rotating stars such as Altair, Vega and substellar companion host Kappa And A exhibit temperature changes of $\sim 1000 - 2000 \text{ K}$ from the stellar pole to equator. This effect is due to gravity darkening, which is directly observed via optical/IR interferometry (Monnier et al. 2007, 2012; Jones et al. 2016) and modelled extensively by Espinosa Lara and Rieutord (2011). Given this large temperature change across the star’s surface, only a rapidly rotating star’s disk-integrated apparent luminosity is observable, which varies with inclination. Jones et al. (2016) observed a 16% difference in the true and apparent luminosity of Kappa And A, a difference that impacts the star’s best-estimated age. HD 1160 A may be a rapidly rotating star, and thus only its apparent luminosity and apparent effective temperature are directly observable without interferometry. Therefore, we derive a Bayesian age estimation for HD 1160 A from the Brandt and Huang (2015) server ¹¹ which incorporates these uncertainties. Brandt and Huang (2015)’s method utilizes the Geneva evolutionary models (Georgy et al. 2013), the atmosphere models of Castelli and Kurucz (2004), and orientation-dependent effective temperatures of Espinosa Lara and Rieutord (2011).

4.3. Results

Our comparisons of HD 1160 to evolutionary models are shown in Figures 12 and 13. Evolutionary models suggest a young age for HD 1160 A: 13 – 150 Myr, 20 – 80 Myr, and 20–125 Myr for the Dartmouth, Yale-Yonsei, and MIST models, respectively, where we quote age ranges consistent with $1-\sigma$ uncertainties in temperature and luminosity. The Bayesian techniques of Brandt and Huang (2015) also point towards a young age. Bayesian methods are consistent with this estimate, with a ‘best-estimated’ age (mean of the age \times probability) of $\sim 58 \text{ Myr}$, and reject ages $\gtrsim 200 \text{ Myr}$ at 90% confidence.

Model comparisons to HD 1160’s lower-mass components generally favor older ages. The Dartmouth, Yale-Yonsei, MIST and Baraffe models favor ages of $> 50 \text{ Myr}$, $> 75 \text{ Myr}$, $> 70 \text{ Myr}$, and $> 45 \text{ Myr}$ for HD 1160 C. The Baraffe models imply an age of $> 80 \text{ Myr}$ for HD 1160 B. In most cases, the lower limit on the components’ luminosity implies stellar

all within $[\text{Fe}/\text{H}] \approx 0.1$ dex of solar metallicity (see continually updated catalog from Dias et al. 2002). The youngest nearby star-forming regions may be slightly sub-solar, ($[\text{Fe}/\text{H}] \approx -0.1$) but with very little dispersion (Santos et al. 2008); slightly older moving groups like AB Dor appear slightly metal rich $[\text{Fe}/\text{H}] \approx 0.1$ (Biazzo et al. 2012). Thus, the small dispersion in metallicity amongst young groups makes $[\text{Fe}/\text{H}] = -0.3$ unlikely.

¹¹www.bayesianstellarparameters.info

objects on/near the main sequence, precluding us from assigning age upper limits.

Therefore, considering HD 1160 A alone, we derive an age range of $\approx 20\text{--}125$ Myr for the system. Considering HD 1160 ABC jointly implies a system age of $80\text{--}125$ Myr. That much of the model age ranges for HD 1160 A vs. HD 1160 BC do not overlap at the $1\text{-}\sigma$ level (Fig. 13) may suggest discrepancies between models and the true physical properties of young M-dwarfs (Feiden et al. 2011; Kraus et al. 2015).

5. Masses of HD 1160 B and C

Given an age of $80\text{--}125$ Myr from analyzing HD 1160 A, B and C jointly, and luminosities of $\log L/L_\odot = -2.76 \pm 0.05$ and $\log L/L_\odot = -2.16 \pm 0.04$, we derive mass ranges of $70\text{--}90 M_J$ and $175\text{--}200 M_J$ for HD 1160 B and C respectively, using the Baraffe et al. (2015) models. Considering instead the age range derived from HD 1160 A alone, we estimate masses of $35\text{--}90 M_J$ for B and $110\text{--}200 M_J$ for HD 1160 C.

Mass estimates derived from our atmospheric modeling are broadly consistent with those derived from luminosity evolution. The best-fit radius and surface gravity for HD 1160 B in turn implies a mass of $\sim 29_{-20.8}^{+63.8} M_J$. Surface gravity is the key factor, as models with $\log g = 4.0$ imply a planet-mass companion ($8\text{--}9 M_J$), those with $\log g = 4.5$ imply a $\sim 30 M_J$ brown dwarf and those with $\log g = 5.0$ imply an object slightly above the hydrogen burning limit of $70\text{--}80 M_J$ (Dieterich et al. 2014)¹².

Thus, the interpretation of HD 1160 B changes depending on which system component(s) we use to estimate an age. Using only HD 1160 A to calibrate the system’s age, HD 1160 B’s allowable mass range mostly covers the substellar regime. But ages derived from both HD 1160 A, B and C preclude the youngest ages ($20\text{--}80$ Myr), and suggest that HD 1160 B straddles the hydrogen burning limit.

Our mass ranges for HD 1160 B agree with those of M16, although our mass upper limit is smaller due to our somewhat lower due to upper age limit of ~ 125 Myr for the HD 1160 system. However, our mass ranges for HD 1160 B are wider than the mass range of $24\text{--}45 M_J$ from N12 using *J*-band photometry, and generally skew towards larger values.

¹²As point of comparison, the surface gravity predicted for HD 1160 B by the Baraffe models for a $30 Myr$ -old, $35 M_J$ object is $\log g \sim 4.4$ and for a $125 Myr$ -old, $90 M_J$ object is $\log g \sim 4.9$. While a subset of these surface gravities are consistent with results from our atmospheric modeling, the predicted temperatures are significantly lower ($2600\text{--}2850 K$), discrepancies that may point to additional challenges in predicting the evolution of young substellar objects and the lowest-mass stars (see also Kraus et al. 2015)

6. Conclusions

We present new and precise SCEXAO near-infrared photometry, reprocessed thermal infrared photometry, and Y band GPI integral field spectroscopy of the HD 1160 system, which hosts two low-mass companions discovered by N12. Combining our data with recently-published near-IR spectroscopy from SPHERE, we constrain the infrared colors of HD 1160 BC and constrain the atmospheric properties of HD 1160 B. After revisiting the age of the HD 1160 system using multiple diagnostics, we determine the likely intrinsic properties of HD 1160 B and C.

Our study yields the following results:

- Our SCEXAO photometry refines estimates for HD 1160 B’s brightness in major infrared passbands. Our photometry in general agrees with the SPHERE 1.0 – 1.6 μm spectrum of M16, although it is systematically fainter by 15 – 25% for HD 1160 B. Like M16, our SCEXAO photometry for HD 1160 B is in strong disagreement with the J and H photometry of Nielsen et al. (2012) by 1.14 and 0.44 mag respectively; our re-derived M' photometry likewise shows differences.
- These differences, in turn, lead to a reinterpretation of HD 1160 B and C’s infrared colors. Specifically, HD 1160 BC are not discrepant or redder than the field sequence but completely consistent with the colors expected for field dwarfs with early-to-mid M spectral types.
- We revise the spectral type of HD 1160 B to $M5.5_{-0.5}^{+1.0}$, earlier than proposed by N12 based purely on near-infrared colors but generally consistent with the results from M16 based on longer-wavelength spectra.
- Our atmospheric modeling finds a best-fit temperature for HD 1160 B of 3000-3100 K and a surface gravity of $\log g = 4-4.5$, although we cannot rule out slightly cooler temperatures or slightly higher surface gravities over some portion of parameter space.
- In contrast with the results from M16, there is no evidence from atmospheric modeling for HD 1160 B having a subsolar metallicity, as the best-fitting (solar metallicity) models are consistent with the data, having a reduced $\chi^2 \sim 1$.
- We find different answers for the system’s age depending on which components are analyzed. Considering HD 1160 A, B and C jointly, we estimate an age of 80–125 Myr, implying masses of 70–90 M_J for HD 1160 B, making it an object straddling the hydrogen burning limit. Considering HD 1160 A alone, we estimate an age of 20–125

Myr, a wider range than that listed in N12 and younger than that estimated in M16. Given this age, we derive a mass of 35–90 M_J for HD 1160 B. Estimated masses for HD 1160 B from atmospheric modeling are consistent with these ranges but allow for slightly lower masses. Our analysis is then consistent with HD 1160 B being a relatively intermediate-to-high mass brown dwarf and HD 1160 C being a low-mass star.

HD 1160 is an important triple system, with one high-mass star, one very low-mass star, and one brown dwarf/extremely low mass star. As such, it presents an excellent laboratory for studying (sub)stellar evolution and testing our understanding of luminosity evolution. Our results may hint at challenges ahead, as age estimates derived from HD 1160 A, although formally consistent with those from HD 1160 BC, tend to be younger than those derived from the system’s lower-mass companions. This trend is an inversion of that seen for ages derived for the youngest clusters/moving groups: e.g. ages derived for Upper Scorpius intermediate-mass stars are older than those derived for its lower-mass stars (e.g. Pecaut et al. 2012; Preibisch et al. 2002). Interferometric measurements of HD 1160 A’s angular diameter, combined with a GAIA parallax more precise than Hipparcos, would clarify the primary’s true luminosity and temperature, allowing us to place more firm constraints on its age based on its HR diagram position.

HD 1160 B is one of many young substellar objects that may provide key insights into young substellar object formation and atmospheres. Its mass ratio (~ 0.015 – 0.040 or 0.027 – 0.040) perhaps indicates formation by protostellar disk fragmentation/disk instability (e.g. Boss 2011), or is consistent with the lowest-mass objects formed like binary stars (Reggiani and Meyer 2011). Indeed, HD 1160 B’s mass range and the semi-major axis is compatible with a disk-instability scenario of formation, provided the disk is $\sim 50\%$ the star’s mass (Figure C.1 in Bonnefoy et al. 2014).

As shown by works like Currie et al. (2013, 2014a) and most recently (and exhaustively) by Faherty et al. (2016), low-gravity L- and T-type brown dwarfs and directly imaged planets across a wide range of spectral types and temperatures appear to depart from the field dwarf sequence. HD 1160 B, an earlier spectral type object, does not yet show differences. Identifying where/why young and low-gravity objects begin to depart from the field provides some insight into and empirical constraint on the evolution of substellar atmospheres.

While the masses inferred for HD 1160 B from atmospheric modeling appear consistent with those implied by luminosity evolution models, our results hint that follow-up spectroscopy could better constrain the object’s surface gravity and (by inference) mass. The implied mass of HD 1160 B varies by a factor of ten from the lowest to highest surface gravity from our suite of acceptably fitting models. As described in Allers and Liu (2013), higher resolution spectroscopy focused on line transitions such as Na I and K I could better clarify

how HD 1160 B’s surface gravity compares to that of other brown dwarfs with a range of ages. Facilities such as Keck/OSIRIS and VLT/SINFONI could potentially provide these data.

Table 1
Observation log of HD 1160

Telescope	Instrument	UT date	Band	Wavelength (μm)	t_{int} (s)	N_{exp}	Mode
Subaru	SCE _x AO/HiCIAO	10-31-2015	<i>Y</i>	0.97 – 1.07	1.5	11	Photometry
Subaru	SCE _x AO/HiCIAO	”	<i>Y</i> +ND	0.97 – 1.07	1.5	6	Photometry
Subaru	SCE _x AO/HiCIAO	”	<i>J</i>	1.17 – 1.33	1.5	5	Photometry
Subaru	SCE _x AO/HiCIAO	”	<i>J</i> +ND	1.17 – 1.33	1.5	5	Photometry
Subaru	SCE _x AO/HiCIAO	”	<i>H</i>	1.49 – 1.78	1.5	5	Photometry
Subaru	SCE _x AO/HiCIAO	”	<i>H</i> +ND	1.49 – 1.78	1.5	5	Photometry
Gemini-South	GPI/IFS	11-17-2013	<i>Y</i>	0.957 – 1.120	88.7	9	IFS

Observation log of HD 1160. ND stands for “neutral density filter”: these are flux calibration observations obtained for HD 1160 with the primary unsaturated and in the linear count regime. The neutral density filters are $0.388 \pm 0.008\%$, $0.590 \pm 0.004\%$ and $0.854 \pm 0.002\%$ for *Y*, *J* and *H*, respectively.

Table 2
Near-to-Mid IR Photometry of HD 1160 B and C

Object	Band (MKO)	SCE _x AO/HiCIAO (mag)	GEMINI/GPI (mag)	VLT/SPHERE (mag)	Gemini/NICI (mag)	Keck/NIRC2 (mag)
HD 1160 B	<i>Y</i>	15.30 ± 0.04^1	$15.13 \pm 0.21^{1,4}$			
	<i>J</i>	14.69 ± 0.05^1		14.47 ± 0.11^2	15.83 ± 0.10^3	
	<i>H</i>	14.21 ± 0.02^1		13.97 ± 0.08^2	14.65 ± 0.08^3	
	<i>K_s</i>			13.96 ± 0.12^2	14.12 ± 0.05^3	
	<i>M'</i>					13.81 ± 0.24^1
HD 1160 C	<i>Y</i>	13.74 ± 0.02^1				
	<i>J</i>	13.37 ± 0.02^1			13.31 ± 0.04^3	
	<i>H</i>	12.81 ± 0.02^1			12.54 ± 0.03^3	
	<i>K_s</i>				12.18 ± 0.06^3	

Photometry.

¹This work.

² The SPHERE *J* and *H* photometry is estimated from the IFS spectrum over the standard MKO *J* and *H* filter bandpasses. The *H* band portion of the SPHERE spectrum is incomplete, covering only the blue half of the standard MKO *H* band wavelength range. The *K_s* band photometry is adopted from Maire et al. (2016). Note that we also adopt their *L'* measurement of $m(L') = 13.60 \pm 0.10$ for HD 1160 B.

³Nielsen et al. (2012)

⁴Note that the GPI *Y* band filter extends to slightly longer wavelengths than HiCIAO.

Table 3
Model Fitting Results

	Photometry	Spectroscopy
Wavelength Range	$YJHK_sL'M'$	0.95-1.6 μm
T_{eff} K (95% Confidence)	2900 – 3500	3000 – 3100K
$\log g$ dex (95% Confidence)	3.5 – 5.5	2900 K (for $\log g = 4.5$) 4.0 – 4.5 5.0 (for 3000 K)
Models fitting both	$T_{\text{eff}} = 3000 - 3100$, $\log g = 4.0 - 4.5$	
Photometry and Spectra:	$T_{\text{eff}} = 2900$ K, $\log g = 4.5$; $T_{\text{eff}} = 3000$ K, $\log g = 5.0$	

Table 4
Properties of the HD 1160 System

Property	HD 1160 A	HD 1160 B	HD 1160 C	Unit
R.A. (ep J2000)		00 : 15 : 57.3025 ⁴		
Dec. (ep J2000)		+04 : 15 : 04.018 ⁴		
Distance		103.1 ± 5.3 ³		pc
μ_α		21.15 ± 0.62 ⁴		mas yr ⁻¹
μ_δ		-14.20 ± 0.24 ⁴		mas yr ⁻¹
Age, HD 1160 A		20 – 125		Myr
Age, HD 1160 A, B and C		80 – 125		Myr
M_Y (MKO)	1.89 ± 0.10 ⁶	10.23 ± 0.11	8.67 ± 0.10	mag
M_J	1.91 ± 0.10	9.63 ± 0.10	8.29 ± 0.10	mag
M_H	1.94 ± 0.10	9.14 ± 0.10	7.75 ± 0.10	mag
M_{K_s}	1.97 ± 0.10	8.88 ± 0.12 ⁵	7.15 ± 0.12 ¹	mag
ΔY (MKO)		8.34 ± 0.04	6.78 ± 0.01	mag
ΔJ		7.72 ± 0.02	6.38 ± 0.01	mag
ΔH		7.20 ± 0.02	5.80 ± 0.01	mag
ΔK		6.91 ± 0.10 ⁵	5.14 ± 0.06 ¹	mag
SpT	A0 ¹	M5.5 ^{+1.0} _{-0.5}	M3.5 ± 0.5 ¹	
T_{eff}	9011 ± 85	3000 – 3100	3260 ± 100 ²	K
Log L/L_\odot	1.12 ± 0.07	-2.76 ± 0.05	-2.16 ± 0.04	dex
Log(g)		4.0 – 4.5		dex
Mass	2.2 ¹			M_\odot
		35–90 ⁷	110–200 ⁷	M_J
		70–90 ⁸	175–200 ⁸	M_J

¹Nielsen et al. (2012).

²Maire et al. (2016).

³ van Leeuwen (2007).

⁴van Leeuwen (2007).

⁵ Calculated by Maire et al. (2016) at the K_s passband using the SPHERE spectrum and IRDIS K_1 and K_2 photometry.

⁶ Calculated from our SED fit to HD 1160 A photometry (see §4).

⁷ mass ranges from Baraffe et al. (2015) models using an age derived for HD 1160 A alone (see §4).

⁸ mass ranges from Baraffe et al. (2015) using an age derived for HD 1160 A, B and C (see §4).

Table 5
Observed Fluxes of the Spectral Energy Distribution of HD 1160 A

Wavelength [μm]	Width [μm]	Flux [ergs/s/cm ⁻² $\times 10^{-10}$]
0.274	0.030	111.000 \pm 2.740
0.236	0.030	117.000 \pm 6.150
0.197	0.030	130.000 \pm 8.250
0.157	0.030	67.000 \pm 4.230
0.422	0.145	403.361 \pm 3.715
0.535	0.167	317.359 \pm 2.923
0.358	0.062	224.376 \pm 2.067
0.450	0.091	423.056 \pm 3.896
0.556	0.086	289.911 \pm 2.670
0.357	0.056	198.323 \pm 1.827
0.487	0.129	367.277 \pm 3.383
0.626	0.134	227.942 \pm 2.099
0.765	0.137	155.036 \pm 1.428
0.907	0.140	112.999 \pm 1.041
1.273	0.152	58.606 \pm 0.540
1.671	0.237	26.964 \pm 0.248
2.171	0.255	13.514 \pm 0.124
3.353	0.663	4.753 \pm 0.232
4.603	1.042	1.745 \pm 0.031
11.561	5.507	0.112 \pm 0.010
22.800	4.101	0.014 \pm 0.002

We thank Anna-Lise Marie for detailed, helpful discussions about and access to SPHERE HD 1160 B spectra. We thank Eric Mamajek for discussions about the metallicity of nearby, young stars. We thank Davy Kirkpatrick for providing NIR spectra of M-dwarf spectral standards. We thank Federico Spada for discussions and tests of the Yonsei-Yale models. EVG would like to acknowledge the gracious support of his Lowell Pre-doctoral Fellowship by the BF foundation and the excellent support of Fisk-Vanderbilt Bridge Program. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. This research has made use of the Keck Observatory Archive (KOA), which is operated by the W. M. Keck Observatory and the NASA Exoplanet Science Institute (NExSci), under contract with the National Aeronautics and Space Administration. The authors acknowledge support

Table 6
Age estimations of the HD 1160 System

Evidence	Age Constraint
HD 1160 A is a main-sequence early A star	$\lesssim 300$ Myr
Darhmouth Models	13 – 150 Myr for HD 1160 A; > 50 Myr for HD 1160 C
Yonsei-Yale Models	20 – 80 Myr for A; > 75 Myr for C
Baraffe Models	> 80 Myr for B; > 45 Myr for C
MIST Models	20 – 125 Myr for A; and > 70 Myr for C
Bayesian analysis of Geneva Models	~ 58 Myr (best-estimated age)
	$\lesssim 200$ Myr (90% confidence) for HD 1160 A
Adopted Age HD 1160 A only	20–125 Myr
Adopted Age HD 1160 A, B and C	80–125 Myr

from the JSPS (Grant-in-Aid for Research #23103002, #23340051, and #26220704). This work was partially supported by the Grant-in-Aid for JSPS fellows (Grant Number 25-8826). M. Janson acknowledges support of the U.S. National Science Foundation, under Award No. 1009203. This work performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344 with document release number LLNL-JRNL-701012-DRAFT. We wish to acknowledge the pivotal cultural role and reverence that the summit of Maunakea has always had within the indigenous Hawaiian community. We are most fortunate to have the privilege to conduct scientific observations from this mountain.

REFERENCES

- F. Allard, D. Homeier, B. Freytag, and C. M. Sharp. Atmospheres From Very Low-Mass Stars to Extrasolar Planets. In C. Reylé, C. Charbonnel, and M. Schultheis, editors, *EAS Publications Series*, volume 57 of *EAS Publications Series*, pages 3–43, Nov. 2012. doi: 10.1051/eas/1257001.
- K. N. Allers and M. C. Liu. A Near-infrared Spectroscopic Study of Young Field Ultracool Dwarfs. *ApJ*, 772:79, Aug. 2013. doi: 10.1088/0004-637X/772/2/79.
- S. M. Ammons, S. E. Robinson, J. Strader, G. Laughlin, D. Fischer, and A. Wolf. The N2K Consortium. IV. New Temperatures and Metallicities for More than 100,000 FGK Dwarfs. *ApJ*, 638:1004–1017, Feb. 2006. doi: 10.1086/498490.
- I. Baraffe, G. Chabrier, T. S. Barman, F. Allard, and P. H. Hauschildt. Evolutionary models

- for cool brown dwarfs and extrasolar giant planets. The case of HD 209458. *A&A*, 402:701–712, May 2003. doi: 10.1051/0004-6361:20030252.
- I. Baraffe, D. Homeier, F. Allard, and G. Chabrier. New evolutionary models for pre-main sequence and main sequence low-mass stars down to the hydrogen-burning limit. *A&A*, 577:A42, May 2015. doi: 10.1051/0004-6361/201425481.
- J.-L. Beuzit, M. Feldt, K. Dohlen, D. Mouillet, P. Puget, F. Wildi, L. Abe, J. Antichi, A. Baruffolo, P. Baudoz, A. Boccaletti, M. Carbillet, J. Charton, R. Claudi, M. Downing, C. Fabron, P. Feautrier, E. Fedrigo, T. Fusco, J.-L. Gach, R. Gratton, T. Henning, N. Hubin, F. Joos, M. Kasper, M. Langlois, R. Lenzen, C. Moutou, A. Pavlov, C. Petit, J. Pragt, P. Rabou, F. Rigal, R. Roelfsema, G. Rousset, M. Saisse, H.-M. Schmid, E. Stadler, C. Thalmann, M. Turatto, S. Udry, F. Vakili, and R. Waters. SPHERE: a ‘Planet Finder’ instrument for the VLT. In *Ground-based and Airborne Instrumentation for Astronomy II*, volume 7014 of *Proc. SPIE*, page 701418, July 2008. doi: 10.1117/12.790120.
- K. Biazzo, V. D’Orazi, S. Desidera, E. Covino, J. M. Alcalá, and M. Zusi. Elemental abundances of low-mass stars in nearby young associations: AB Doradus, Carina Near and Ursa Major. *MNRAS*, 427:2905–2916, Dec. 2012. doi: 10.1111/j.1365-2966.2012.22132.x.
- B. A. Biller, M. C. Liu, Z. Wahhaj, E. L. Nielsen, L. M. Close, T. J. Dupuy, T. L. Hayward, A. Burrows, M. Chun, C. Ftaclas, F. Clarke, M. Hartung, J. Males, I. N. Reid, E. L. Shkolnik, A. Skemer, M. Tecza, N. Thatte, S. H. P. Alencar, P. Artymowicz, A. Boss, E. de Gouveia Dal Pino, J. Gregorio-Hetem, S. Ida, M. J. Kuchner, D. Lin, and D. Toomey. The Gemini NICI Planet-finding Campaign: Discovery of a Close Substellar Companion to the Young Debris Disk Star PZ Tel. *ApJ*, 720:L82–L87, Sept. 2010. doi: 10.1088/2041-8205/720/1/L82.
- A. Boccaletti, A.-M. Lagrange, M. Bonnefoy, R. Galicher, and G. Chauvin. Independent confirmation of β Pictoris b imaging with NICI. *A&A*, 551:L14, Mar. 2013. doi: 10.1051/0004-6361/201321255.
- M. Bonnefoy, T. Currie, G.-D. Marleau, J. E. Schlieder, J. Wisniewski, J. Carson, K. R. Covey, T. Henning, B. Biller, P. Hinz, H. Klahr, A. N. Marsh Boyer, N. Zimmerman, M. Janson, M. McElwain, C. Mordasini, A. Skemer, V. Bailey, D. Defrère, C. Thalmann, M. Skrutskie, F. Allard, D. Homeier, M. Tamura, M. Feldt, A. Cumming, C. Grady, W. Brandner, C. Helling, S. Witte, P. Hauschildt, R. Kandori, M. Kuzuhara, M. Fukagawa, J. Kwon, T. Kudo, J. Hashimoto, N. Kusakabe, L. Abe,

- T. Brandt, S. Egner, O. Guyon, Y. Hayano, M. Hayashi, S. Hayashi, K. Hodapp, M. Ishii, M. Iye, G. Knapp, T. Matsuo, K. Mede, M. Miyama, J.-I. Morino, A. Moro-Martin, T. Nishimura, T. Pyo, E. Serabyn, T. Suenaga, H. Suto, R. Suzuki, Takahashi, M. Takami, N. Takato, H. Terada, D. Tomono, E. Turner, M. Watanabe, T. Yamada, H. Takami, and T. Usuda. Characterization of the gaseous companion κ Andromedae b. New Keck and LBTI high-contrast observations. *A&A*, 562:A111, Feb. 2014. doi: 10.1051/0004-6361/201322119.
- A. P. Boss. Formation of Giant Planets by Disk Instability on Wide Orbits Around Protostars with Varied Masses. *ApJ*, 731:74, Apr. 2011. doi: 10.1088/0004-637X/731/1/74.
- T. D. Brandt and C. X. Huang. Bayesian Ages for Early-type Stars from Isochrones Including Rotation, and a Possible Old Age for the Hyades. *ApJ*, 807:58, July 2015. doi: 10.1088/0004-637X/807/1/58.
- J. Carson, C. Thalmann, M. Janson, T. Kozakis, M. Bonnefoy, B. Biller, J. Schlieder, T. Currie, M. McElwain, M. Goto, T. Henning, W. Brandner, M. Feldt, R. Kandori, M. Kuzuhara, L. Stevens, P. Wong, K. Gainey, M. Fukagawa, Y. Kuwada, T. Brandt, J. Kwon, L. Abe, S. Egner, C. Grady, O. Guyon, J. Hashimoto, Y. Hayano, M. Hayashi, S. Hayashi, K. Hodapp, M. Ishii, M. Iye, G. Knapp, T. Kudo, N. Kusakabe, T. Matsuo, S. Miyama, J. Morino, A. Moro-Martin, T. Nishimura, T. Pyo, E. Serabyn, H. Suto, R. Suzuki, M. Takami, N. Takato, H. Terada, D. Tomono, E. Turner, M. Watanabe, J. Wisniewski, T. Yamada, H. Takami, T. Usuda, and M. Tamura. Direct Imaging Discovery of a "Super-Jupiter" around the Late B-type Star κ And. *ApJ*, 763:L32, Feb. 2013. doi: 10.1088/2041-8205/763/2/L32.
- F. Castelli and R. L. Kurucz. New Grids of ATLAS9 Model Atmospheres. *ArXiv Astrophysics e-prints*, May 2004.
- G. Chauvin, A.-M. Lagrange, B. Zuckerman, C. Dumas, D. Mouillet, I. Song, J.-L. Beuzit, P. Lowrance, and M. S. Bessell. A companion to AB Pic at the planet/brown dwarf boundary. *A&A*, 438:L29–L32, Aug. 2005. doi: 10.1051/0004-6361:200500111.
- J. Choi, A. Dotter, C. Conroy, M. Cantiello, B. Paxton, and B. D. Johnson. Mesa Isochrones and Stellar Tracks (MIST). I. Solar-scaled Models. *ApJ*, 823:102, June 2016. doi: 10.3847/0004-637X/823/2/102.
- T. Currie, A. Burrows, Y. Itoh, S. Matsumura, M. Fukagawa, D. Apai, N. Madhusudhan, P. M. Hinz, T. J. Rodigas, M. Kasper, T.-S. Pyo, and S. Ogino. A Combined Subaru/VLT/MMT 1-5 μ m Study of Planets Orbiting HR 8799: Implications for

- Atmospheric Properties, Masses, and Formation. *ApJ*, 729:128, Mar. 2011. doi: 10.1088/0004-637X/729/2/128.
- T. Currie, A. Burrows, N. Madhusudhan, M. Fukagawa, J. H. Girard, R. Dawson, R. Murray-Clay, S. Kenyon, M. Kuchner, S. Matsumura, R. Jayawardhana, J. Chambers, and B. Bromley. A Combined Very Large Telescope and Gemini Study of the Atmosphere of the Directly Imaged Planet, β Pictoris b. *ApJ*, 776:15, Oct. 2013. doi: 10.1088/0004-637X/776/1/15.
- T. Currie, S. Daemgen, J. Debes, D. Lafreniere, Y. Itoh, R. Jayawardhana, T. Ratzka, and S. Correia. Direct Imaging and Spectroscopy of a Candidate Companion Below/Near the Deuterium-burning Limit in the Young Binary Star System, ROXs 42B. *ApJ*, 780:L30, Jan. 2014a. doi: 10.1088/2041-8205/780/2/L30.
- T. Currie, T. Muto, T. Kudo, M. Honda, T. D. Brandt, C. Grady, M. Fukagawa, A. Burrows, M. Janson, M. Kuzuhara, M. W. McElwain, K. Follette, J. Hashimoto, T. Henning, R. Kandori, N. Kusakabe, J. Kwon, K. Mede, J.-i. Morino, J. Nishikawa, T.-S. Pyo, G. Serabyn, T. Suenaga, Y. Takahashi, J. Wisniewski, and M. Tamura. Recovery of the Candidate Protoplanet HD 100546 b with Gemini/NICI and Detection of Additional (Planet-induced?) Disk Structure at Small Separations. *ApJ*, 796:L30, Dec. 2014b. doi: 10.1088/2041-8205/796/2/L30.
- T. Currie, R. Cloutier, S. Brittain, C. Grady, A. Burrows, T. Muto, S. J. Kenyon, and M. J. Kuchner. Resolving the HD 100546 Protoplanetary System with the Gemini Planet Imager: Evidence for Multiple Forming, Accreting Planets. *ApJ*, 814:L27, Dec. 2015. doi: 10.1088/2041-8205/814/2/L27.
- R. M. Cutri and et al. VizieR Online Data Catalog: WISE All-Sky Data Release (Cutri+ 2012). *VizieR Online Data Catalog*, 2311, 2012.
- R. M. Cutri, M. F. Skrutskie, S. van Dyk, C. A. Beichman, J. M. Carpenter, T. Chester, L. Cambresy, T. Evans, J. Fowler, J. Gizis, E. Howard, J. Huchra, T. Jarrett, E. L. Kopan, J. D. Kirkpatrick, R. M. Light, K. A. Marsh, H. McCallon, S. Schneider, R. Stiening, M. Sykes, M. Weinberg, W. A. Wheaton, S. Wheelock, and N. Zacarias. VizieR Online Data Catalog: 2MASS All-Sky Catalog of Point Sources (Cutri+ 2003). *VizieR Online Data Catalog*, 2246, June 2003.
- P. Dawson, A. Scholz, T. P. Ray, D. E. Peterson, D. Rodgers-Lee, and V. Geers. Near-infrared spectroscopy of young brown dwarfs in upper Scorpius. *MNRAS*, 442:1586–1596, Aug. 2014. doi: 10.1093/mnras/stu973.

- W. S. Dias, B. S. Alessi, A. Moitinho, and J. R. D. Lépine. New catalogue of optically visible open clusters and candidates. *A&A*, 389:871–873, July 2002. doi: 10.1051/0004-6361:20020668.
- S. B. Dieterich, T. J. Henry, W.-C. Jao, J. G. Winters, A. D. Hosey, A. R. Riedel, and J. P. Subasavage. The Solar Neighborhood. XXXII. The Hydrogen Burning Limit. *AJ*, 147:94, May 2014. doi: 10.1088/0004-6256/147/5/94.
- A. Dotter. MESA Isochrones and Stellar Tracks (MIST) 0: Methods for the Construction of Stellar Isochrones. *ApJS*, 222:8, Jan. 2016. doi: 10.3847/0067-0049/222/1/8.
- A. Dotter, B. Chaboyer, D. Jevremović, V. Kostov, E. Baron, and J. W. Ferguson. The Dartmouth Stellar Evolution Database. *ApJS*, 178:89-101, Sept. 2008. doi: 10.1086/589654.
- T. J. Dupuy and M. C. Liu. The Hawaii Infrared Parallax Program. I. Ultracool Binaries and the L/T Transition. *ApJS*, 201:19, Aug. 2012. doi: 10.1088/0067-0049/201/2/19.
- T. J. Dupuy, M. C. Liu, and M. J. Ireland. Dynamical Mass of the Substellar Benchmark Binary HD 130948BC. *ApJ*, 692:729–752, Feb. 2009. doi: 10.1088/0004-637X/692/1/729.
- T. J. Dupuy, M. C. Liu, B. P. Bowler, M. C. Cushing, C. Helling, S. Witte, and P. Hauschildt. Studying the Physical Diversity of Late-M Dwarfs with Dynamical Masses. *ApJ*, 721:1725–1747, Oct. 2010. doi: 10.1088/0004-637X/721/2/1725.
- T. J. Dupuy, M. C. Liu, S. K. Leggett, M. J. Ireland, K. Chiu, and D. A. Golimowski. The Mass-Luminosity Relation in the L/T Transition: Individual Dynamical Masses for the New J-band Flux Reversal Binary SDSSJ105213.51+442255.7AB. *ApJ*, 805:56, May 2015. doi: 10.1088/0004-637X/805/1/56.
- T. J. Dupuy, J. Forbrich, A. Rizzuto, A. W. Mann, K. Aller, M. C. Liu, A. L. Kraus, and E. Berger. High-Precision Radio and Infrared Astrometry of LSPM J1314+1320AB - II: Testing Pre-Main-Sequence Models at the Lithium Depletion Boundary with Dynamical Masses. *ArXiv e-prints*, May 2016.
- J. H. Elias, J. A. Frogel, K. Matthews, and G. Neugebauer. Infrared standard stars. *AJ*, 87:1029–1034, July 1982. doi: 10.1086/113185.
- F. Espinosa Lara and M. Rieutord. Gravity darkening in rotating stars. *A&A*, 533:A43, Sept. 2011. doi: 10.1051/0004-6361/201117252.

- J. K. Faherty, A. R. Riedel, K. L. Cruz, J. Gagne, J. C. Filippazzo, E. Lambrides, H. Fica, A. Weinberger, J. R. Thorstensen, C. G. Tinney, V. Baldassare, E. Lemonier, and E. L. Rice. Population Properties of Brown Dwarf Analogs to Exoplanets. *ArXiv e-prints*, May 2016.
- G. A. Feiden, B. Chaboyer, and A. Dotter. Accurate Low-mass Stellar Models of KOI-126. *ApJ*, 740:L25, Oct. 2011. doi: 10.1088/2041-8205/740/1/L25.
- R. Galicher, C. Marois, B. Macintosh, T. Barman, and Q. Konopacky. M-band Imaging of the HR 8799 Planetary System Using an Innovative LOCI-based Background Subtraction Technique. *ApJ*, 739:L41, Oct. 2011. doi: 10.1088/2041-8205/739/2/L41.
- C. Georgy, S. Ekström, A. Granada, G. Meynet, N. Mowlavi, P. Eggenberger, and A. Maeder. Populations of rotating stars. I. Models from 1.7 to 15 M_{\odot} at $Z = 0.014$, 0.006, and 0.002 with $\Omega/\Omega_{\text{crit}}$ between 0 and 1. *A&A*, 553:A24, May 2013. doi: 10.1051/0004-6361/201220558.
- P. H. Hauschildt, F. Allard, and E. Baron. The NextGen Model Atmosphere Grid for 3000–10,000 K. *ApJ*, 512:377–385, Feb. 1999. doi: 10.1086/306745.
- P. C. Hewett, S. J. Warren, S. K. Leggett, and S. T. Hodgkin. The UKIRT Infrared Deep Sky Survey ZY JHK photometric system: passbands and synthetic colours. *MNRAS*, 367:454–468, Apr. 2006. doi: 10.1111/j.1365-2966.2005.09969.x.
- M. J. Ireland, A. Kraus, F. Martinache, N. Law, and L. A. Hillenbrand. Two Wide Planetary-mass Companions to Solar-type Stars in Upper Scorpius. *ApJ*, 726:113, Jan. 2011. doi: 10.1088/0004-637X/726/2/113.
- J. Jones, R. J. White, S. Quinn, M. Ireland, T. Boyajian, G. Schaefer, and E. K. Baines. The Age of the Directly Imaged Planet Host Star κ Andromedae Determined from Interferometric Observations. *ApJ*, 822:L3, May 2016. doi: 10.3847/2041-8205/822/1/L3.
- N. Jovanovic, F. Martinache, O. Guyon, C. Clergeon, G. Singh, T. Kudo, V. Garrel, K. Newman, D. Doughty, J. Lozi, J. Males, Y. Minowa, Y. Hayano, N. Takato, J. Morino, J. Kuhn, E. Serabyn, B. Norris, P. Tuthill, G. Schworer, P. Stewart, L. Close, E. Huby, G. Perrin, S. Lacour, L. Gauchet, S. Vievard, N. Murakami, F. Oshiyama, N. Baba, T. Matsuo, J. Nishikawa, M. Tamura, O. Lai, F. Marchis, G. Duchene, T. Kotani, and J. Woillez. The Subaru Coronagraphic Extreme Adaptive Optics System: Enabling High-Contrast Imaging on Solar-System Scales. *PASP*, 127:890–910, Oct. 2015. doi: 10.1086/682989.

- J. D. Kirkpatrick, D. L. Looper, A. J. Burgasser, S. D. Schurr, R. M. Cutri, M. C. Cushing, K. L. Cruz, A. C. Sweet, G. R. Knapp, T. S. Barman, J. J. Bochanski, T. L. Roellig, I. S. McLean, M. R. McGovern, and E. L. Rice. Discoveries from a Near-infrared Proper Motion Survey Using Multi-epoch Two Micron All-Sky Survey Data. *ApJS*, 190:100–146, Sept. 2010. doi: 10.1088/0067-0049/190/1/100.
- Q. M. Konopacky, S. J. Thomas, B. A. Macintosh, D. Dillon, N. Sadakuni, J. Maire, M. Fitzgerald, S. Hinkley, P. Kalas, T. Esposito, C. Marois, P. J. Ingraham, F. Marchis, M. D. Perrin, J. R. Graham, J. J. Wang, R. J. De Rosa, K. Morzinski, L. Pueyo, J. K. Chilcote, J. E. Larkin, D. Fabrycky, S. J. Goodsell, B. R. Oppenheimer, J. Patience, L. Saddlemyer, and A. Sivaramakrishnan. Gemini planet imager observational calibrations V: astrometry and distortion. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, volume 9147 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, page 84, July 2014. doi: 10.1117/12.2056646.
- K. M. Kratter, R. A. Murray-Clay, and A. N. Youdin. The Runts of the Litter: Why Planets Formed Through Gravitational Instability Can Only Be Failed Binary Stars. *ApJ*, 710:1375–1386, Feb. 2010. doi: 10.1088/0004-637X/710/2/1375.
- A. L. Kraus and M. J. Ireland. LkCa 15: A Young Exoplanet Caught at Formation? *ApJ*, 745:5, Jan. 2012. doi: 10.1088/0004-637X/745/1/5.
- A. L. Kraus, A. M. Cody, K. R. Covey, A. C. Rizzuto, A. W. Mann, and M. J. Ireland. The Mass-Radius Relation of Young Stars. I. USco 5, an M4.5 Eclipsing Binary in Upper Scorpius Observed by K2. *ApJ*, 807:3, July 2015. doi: 10.1088/0004-637X/807/1/3.
- M. Kuzuhara, M. Tamura, T. Kudo, M. Janson, R. Kandori, T. D. Brandt, C. Thalmann, D. Spiegel, B. Biller, J. Carson, Y. Hori, R. Suzuki, A. Burrows, T. Henning, E. L. Turner, M. W. McElwain, A. Moro-Martín, T. Suenaga, Y. H. Takahashi, J. Kwon, P. Lucas, L. Abe, W. Brandner, S. Egner, M. Feldt, H. Fujiwara, M. Goto, C. A. Grady, O. Guyon, J. Hashimoto, Y. Hayano, M. Hayashi, S. S. Hayashi, K. W. Hodapp, M. Ishii, M. Iye, G. R. Knapp, T. Matsuo, S. Mayama, S. Miyama, J.-I. Morino, J. Nishikawa, T. Nishimura, T. Kotani, N. Kusakabe, T.-S. Pyo, E. Serabyn, H. Suto, M. Takami, N. Takato, H. Terada, D. Tomono, M. Watanabe, J. P. Wisniewski, T. Yamada, H. Takami, and T. Usuda. Direct Imaging of a Cold Jovian Exoplanet in Orbit around the Sun-like Star GJ 504. *ApJ*, 774:11, Sept. 2013. doi: 10.1088/0004-637X/774/1/11.
- A.-M. Lagrange, M. Bonnefoy, G. Chauvin, D. Apai, D. Ehrenreich, A. Boccaletti, D. Grata-dour, D. Rouan, D. Mouillet, S. Lacour, and M. Kasper. A Giant Planet Im-

- aged in the Disk of the Young Star β Pictoris. *Science*, 329:57, July 2010. doi: 10.1126/science.1187187.
- J. E. Larkin, J. K. Chilcote, T. Aliado, B. J. Bauman, G. Brims, J. M. Canfield, A. Cardwell, D. Dillon, R. Doyon, J. Dunn, M. P. Fitzgerald, J. R. Graham, S. Goodsell, M. Hartung, P. Hibon, P. Ingraham, C. A. Johnson, E. Kress, Q. M. Konopacky, B. A. Macintosh, K. G. Magnone, J. Maire, I. S. McLean, D. Palmer, M. D. Perrin, C. Quiroz, F. Rantakyrö, N. Sadakuni, L. Saddlemyer, A. Serio, S. Thibault, S. J. Thomas, P. Vallee, and J. L. Weiss. The integral field spectrograph for the Gemini planet imager. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, volume 9147 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, page 1, July 2014. doi: 10.1117/12.2056504.
- M. C. Liu, Z. Wahhaj, B. A. Biller, E. L. Nielsen, M. Chun, L. M. Close, C. Ftaclas, M. Hartung, T. L. Hayward, F. Clarke, I. N. Reid, E. L. Shkolnik, M. Tecza, N. Thatte, S. Alencar, P. Artymowicz, A. Boss, A. Burrows, E. de Gouveia Dal Pino, J. Gregorio-Hetem, S. Ida, M. J. Kuchner, D. Lin, and D. Toomey. The Gemini NICI Planet-Finding Campaign. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, volume 7736 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, page 1, July 2010. doi: 10.1117/12.858358.
- M. C. Liu, T. J. Dupuy, B. P. Bowler, S. K. Leggett, and W. M. J. Best. Two Extraordinary Substellar Binaries at the T/Y Transition and the Y-band Fluxes of the Coolest Brown Dwarfs. *ApJ*, 758:57, Oct. 2012. doi: 10.1088/0004-637X/758/1/57.
- J. Lozi, O. Guyon, N. Jovanovic, G. Singh, G. Goebel, B. Norris, and H. Okita. SPHERE: a ‘Planet Finder’ instrument for the VLT. In *Characterizing and mitigating vibrations for SCExAO*, volume 9909 of *Proc. Soc. Photo-Opt. Instrum. Eng.*, page 9909, June 2016.
- B. Macintosh, J. Graham, D. Palmer, R. Doyon, D. Gavel, J. Larkin, B. Oppenheimer, L. Saddlemyer, J. K. Wallace, B. Bauman, J. Evans, D. Erikson, K. Morzinski, D. Phillion, L. Poyneer, A. Sivaramakrishnan, R. Soummer, S. Thibault, and J.-P. Veran. The Gemini Planet Imager. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, volume 6272 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, page 0, June 2006. doi: 10.1117/12.672430.
- B. Macintosh, J. R. Graham, T. Barman, R. J. De Rosa, Q. Konopacky, M. S. Marley, C. Marois, E. L. Nielsen, L. Pueyo, A. Rajan, J. Rameau, D. Saumon, J. J. Wang,

- J. Patience, M. Ammons, P. Arriaga, E. Artigau, S. Beckwith, J. Brewster, S. Bruzzone, J. Bulger, B. Burningham, A. S. Burrows, C. Chen, E. Chiang, J. K. Chilcote, R. I. Dawson, R. Dong, R. Doyon, Z. H. Draper, G. Duchêne, T. M. Esposito, D. Fabrycky, M. P. Fitzgerald, K. B. Follette, J. J. Fortney, B. Gerard, S. Goodsell, A. Z. Greenbaum, P. Hibon, S. Hinkley, T. H. Cotten, L.-W. Hung, P. Ingraham, M. Johnson-Groh, P. Kalas, D. Lafreniere, J. E. Larkin, J. Lee, M. Line, D. Long, J. Maire, F. Marchis, B. C. Matthews, C. E. Max, S. Metchev, M. A. Millar-Blanchaer, T. Mittal, C. V. Morley, K. M. Morzinski, R. Murray-Clay, R. Oppenheimer, D. W. Palmer, R. Patel, M. D. Perrin, L. A. Poyneer, R. R. Rafikov, F. T. Rantakyro, E. L. Rice, P. Rojo, A. R. Rudy, J.-B. Ruffio, M. T. Ruiz, N. Sadakuni, L. Saddlemyer, M. Salama, D. Savransky, A. C. Schneider, A. Sivaramakrishnan, I. Song, R. Soummer, S. Thomas, G. Vasisht, J. K. Wallace, K. Ward-Duong, S. J. Wiktorowicz, S. G. Wolff, and B. Zuckerman. Discovery and spectroscopy of the young jovian planet 51 Eri b with the Gemini Planet Imager. *Science*, 350:64–67, Oct. 2015. doi: 10.1126/science.aac5891.
- B. A. Macintosh, A. Anthony, J. Atwood, B. Bauman, A. Cardwell, K. Caputa, J. Chilcote, R. J. De Rosa, D. Dillon, R. Doyon, J. Dunn, D. Erickson, M. P. Fitzgerald, D. T. Gavel, R. Galvez, S. Goodsell, J. Graham, A. Z. Greenbaum, M. Hartung, P. Hibon, P. Ingraham, D. Kerley, Q. Konopacky, K. Labrie, J. Larkin, J. Maire, F. Marchis, C. Marois, M. Millar-Blanchaer, K. Morzinski, A. Nunez, R. Oppenheimer, D. Palmer, J. Pazder, M. Perrin, L. A. Poyneer, L. Pueyo, C. Quiroz, F. Rantakyro, V. Reshetov, L. Saddlemyer, N. Sadakuni, D. Savransky, A. Serio, A. Sivaramakrishnan, M. Smith, R. Soummer, S. Thomas, J. K. Wallace, J. Wang, J. Weiss, S. Wiktorowicz, and S. G. Wolff. The Gemini planet imager: first light and commissioning. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, volume 9148 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, page 0, Aug. 2014. doi: 10.1117/12.2056709.
- A.-L. Maire, M. Bonnefoy, C. Ginski, A. Vigan, S. Messina, D. Mesa, R. Galicher, R. Gratton, S. Desidera, T. G. Kopytova, M. Millward, C. Thalmann, R. U. Claudi, D. Ehrenreich, A. Zurlo, G. Chauvin, J. Antichi, A. Baruffolo, A. Bazzon, J.-L. Beuzit, P. Blanchard, A. Boccaletti, J. de Boer, M. Carle, E. Cascone, A. Costille, V. De Caprio, A. Delboulbé, K. Dohlen, C. Dominik, M. Feldt, T. Fusco, J. H. Girard, E. Giro, D. Gisler, L. Gluck, C. Gry, T. Henning, N. Hubin, E. Hugot, M. Jaquet, M. Kasper, A.-M. Lagrange, M. Langlois, D. Le Mignant, M. Llored, F. Madec, P. Martinez, D. Mawet, J. Milli, O. Möller-Nilsson, D. Mouillet, T. Moulin, C. Moutou, A. Origné, A. Pavlov, C. Petit, J. Pragt, P. Puget, J. Ramos, S. Rochat, R. Roelfsema, B. Salasnich, J.-F. Sauvage, H. M. Schmid, M. Turatto, S. Udry, F. Vakili, Z. Wahhaj, L. Weber, and

- F. Wildi. First light of the VLT planet finder SPHERE. II. The physical properties and the architecture of the young systems PZ Telescopii and HD 1160 revisited. *A&A*, 587:A56, Mar. 2016. doi: 10.1051/0004-6361/201526594.
- J. Maire, M. D. Perrin, R. Doyon, E. Artigau, J. Dunn, D. T. Gavel, J. R. Graham, D. Lafrenière, J. E. Larkin, J.-F. Lavigne, B. A. Macintosh, C. Marois, B. Oppenheimer, D. W. Palmer, L. A. Poyneer, S. Thibault, and J.-P. Véran. Data reduction pipeline for the Gemini Planet Imager. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, volume 7735 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, page 31, July 2010. doi: 10.1117/12.858028.
- J. Maire, M. D. Perrin, R. Doyon, J. Chilcote, J. E. Larkin, J. L. Weiss, C. Marois, Q. M. Konopacky, M. Millar-Blanchaer, J. R. Graham, J. Dunn, R. Galicher, F. Marchis, S. J. Wiktorowicz, K. Labrie, S. J. Thomas, S. J. Goodsell, F. T. Rantakyro, D. W. Palmer, and B. A. Macintosh. Test results for the Gemini Planet Imager data reduction pipeline. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, volume 8451 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, page 3, Sept. 2012. doi: 10.1117/12.926202.
- C. Marois, D. Lafrenière, R. Doyon, B. Macintosh, and D. Nadeau. Angular Differential Imaging: A Powerful High-Contrast Imaging Technique. *ApJ*, 641:556–564, Apr. 2006. doi: 10.1086/500401.
- C. Marois, B. Macintosh, T. Barman, B. Zuckerman, I. Song, J. Patience, D. Lafrenière, and R. Doyon. Direct Imaging of Multiple Planets Orbiting the Star HR 8799. *Science*, 322:1348, Nov. 2008. doi: 10.1126/science.1166585.
- C. Marois, B. Macintosh, and J.-P. Véran. Exoplanet imaging with LOCI processing: photometry and astrometry with the new SOSIE pipeline. In *Adaptive Optics Systems II*, volume 7736 of *Proc. SPIE*, page 77361J, July 2010. doi: 10.1117/12.857225.
- J. D. Monnier, M. Zhao, E. Pedretti, N. Thureau, M. Ireland, P. Muirhead, J.-P. Berger, R. Millan-Gabet, G. Van Belle, T. ten Brummelaar, H. McAlister, S. Ridgway, N. Turner, L. Sturmann, J. Sturmann, and D. Berger. Imaging the Surface of Altair. *Science*, 317:342, July 2007. doi: 10.1126/science.1143205.
- J. D. Monnier, X. Che, M. Zhao, S. Ekström, V. Maestro, J. Aufdenberg, F. Baron, C. Georgy, S. Kraus, H. McAlister, E. Pedretti, S. Ridgway, J. Sturmann, L. Sturmann, T. ten Brummelaar, N. Thureau, N. Turner, and P. G. Tuthill. Resolving Vega

- and the Inclination Controversy with CHARA/MIRC. *ApJ*, 761:L3, Dec. 2012. doi: 10.1088/2041-8205/761/1/L3.
- E. L. Nielsen, M. C. Liu, Z. Wahhaj, B. A. Biller, T. L. Hayward, A. Boss, B. Bowler, A. Kraus, E. L. Shkolnik, M. Tecza, M. Chun, F. Clarke, L. M. Close, C. Ftaclas, M. Hartung, J. R. Males, I. N. Reid, A. J. Skemer, S. H. P. Alencar, A. Burrows, E. de Gouveia Dal Pino, J. Gregorio-Hetem, M. Kuchner, N. Thatte, and D. W. Toomey. The Gemini NICI Planet-Finding Campaign: Discovery of a Multiple System Orbiting the Young A Star HD 1160. *ApJ*, 750:53, May 2012. doi: 10.1088/0004-637X/750/1/53.
- B. Paxton, L. Bildsten, A. Dotter, F. Herwig, P. Lesaffre, and F. Timmes. Modules for Experiments in Stellar Astrophysics (MESA). *ApJS*, 192:3, Jan. 2011. doi: 10.1088/0067-0049/192/1/3.
- B. Paxton, M. Cantiello, P. Arras, L. Bildsten, E. F. Brown, A. Dotter, C. Mankovich, M. H. Montgomery, D. Stello, F. X. Timmes, and R. Townsend. Modules for Experiments in Stellar Astrophysics (MESA): Planets, Oscillations, Rotation, and Massive Stars. *ApJS*, 208:4, Sept. 2013. doi: 10.1088/0067-0049/208/1/4.
- M. J. Pecaut and E. E. Mamajek. Intrinsic Colors, Temperatures, and Bolometric Corrections of Pre-main-sequence Stars. *ApJS*, 208:9, Sept. 2013. doi: 10.1088/0067-0049/208/1/9.
- M. J. Pecaut, E. E. Mamajek, and E. J. Bubar. A Revised Age for Upper Scorpius and the Star Formation History among the F-type Members of the Scorpius-Centaurus OB Association. *ApJ*, 746:154, Feb. 2012. doi: 10.1088/0004-637X/746/2/154.
- M. D. Perrin, J. Maire, P. Ingraham, D. Savransky, M. Millar-Blanchaer, S. G. Wolff, J.-B. Ruffio, J. J. Wang, Z. H. Draper, N. Sadakuni, C. Marois, A. Rajan, M. P. Fitzgerald, B. Macintosh, J. R. Graham, R. Doyon, J. E. Larkin, J. K. Chilcote, S. J. Goodsell, D. W. Palmer, K. Labrie, M. Beaulieu, R. J. De Rosa, A. Z. Greenbaum, M. Hartung, P. Hibon, Q. Konopacky, D. Lafreniere, J.-F. Lavigne, F. Marchis, J. Patience, L. Pueyo, F. T. Rantakyro, R. Soummer, A. Sivaramakrishnan, S. Thomas, K. Ward-Duong, and S. Wiktorowicz. Gemini Planet Imager observational calibrations I: Overview of the GPI data reduction pipeline. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, volume 9147 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, page 3, July 2014. doi: 10.1117/12.2055246.

- A. Pickles and É. Depagne. All-Sky Spectrally Matched UBVRI-ZY and ugriz Magnitudes for Stars in the Tycho2 Catalog. *PASP*, 122:1437–1464, Dec. 2010. doi: 10.1086/657947.
- A. J. Pickles. A Stellar Spectral Flux Library: 1150-25000 Å. *PASP*, 110:863–878, July 1998. doi: 10.1086/316197.
- T. Preibisch, A. G. A. Brown, T. Bridges, E. Guenther, and H. Zinnecker. Exploring the Full Stellar Population of the Upper Scorpius OB Association. *AJ*, 124:404–416, July 2002. doi: 10.1086/341174.
- W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery. *Numerical recipes in FORTRAN. The art of scientific computing*. Cambridge: University Press —c1992, 1992.
- S. P. Quanz, A. Amara, M. R. Meyer, M. A. Kenworthy, M. Kasper, and J. H. Girard. A Young Protoplanet Candidate Embedded in the Circumstellar Disk of HD 100546. *ApJ*, 766:L1, Mar. 2013. doi: 10.1088/2041-8205/766/1/L1.
- J. Rameau, G. Chauvin, A.-M. Lagrange, T. Meshkat, A. Boccaletti, S. P. Quanz, T. Currie, D. Mawet, J. H. Girard, M. Bonnefoy, and M. Kenworthy. Confirmation of the Planet around HD 95086 by Direct Imaging. *ApJ*, 779:L26, Dec. 2013. doi: 10.1088/2041-8205/779/2/L26.
- J. T. Rayner, D. W. Toomey, P. M. Onaka, A. J. Denault, W. E. Stahlberger, W. D. Vacca, M. C. Cushing, and S. Wang. SpeX: A Medium-Resolution 0.8-5.5 Micron Spectrograph and Imager for the NASA Infrared Telescope Facility. *PASP*, 115:362–382, Mar. 2003. doi: 10.1086/367745.
- M. M. Reggiani and M. R. Meyer. Binary Formation Mechanisms: Constraints from the Companion Mass Ratio Distribution. *ApJ*, 738:60, Sept. 2011. doi: 10.1088/0004-637X/738/1/60.
- N. C. Santos, C. Melo, D. J. James, J. F. Gameiro, J. Bouvier, and J. I. Gomes. Chemical abundances in six nearby star-forming regions. Implications for galactic evolution and planet searches around very young stars. *A&A*, 480:889–897, Mar. 2008. doi: 10.1051/0004-6361:20079083.
- L. Siess, E. Dufour, and M. Forestini. An internet server for pre-main sequence tracks of low- and intermediate-mass stars. *A&A*, 358:593–599, June 2000.
- D. R. Soderblom, L. A. Hillenbrand, R. D. Jeffries, E. E. Mamajek, and T. Naylor. Ages of Young Stars. *Protostars and Planets VI*, pages 219–241, 2014. doi: 10.2458/azu_uapress_9780816531240-ch010.

- R. Soummer, L. Pueyo, and J. Larkin. Detection and Characterization of Exoplanets and Disks Using Projections on Karhunen-Loève Eigenimages. *ApJ*, 755:L28, Aug. 2012. doi: 10.1088/2041-8205/755/2/L28.
- F. Spada, P. Demarque, Y.-C. Kim, and A. Sills. The Radius Discrepancy in Low-mass Stars: Single versus Binaries. *ApJ*, 776:87, Oct. 2013. doi: 10.1088/0004-637X/776/2/87.
- M. Tamura. Subaru Strategic Exploration of Exoplanets and Disks with HiCIAO/AO188 (SEEDS). In T. Usuda, M. Tamura, and M. Ishii, editors, *American Institute of Physics Conference Series*, volume 1158 of *American Institute of Physics Conference Series*, pages 11–16, Aug. 2009. doi: 10.1063/1.3215811.
- G. I. Thompson, K. Nandy, C. Jamar, A. Monfils, L. Houziaux, D. J. Carnochan, and R. Wilson. *Catalogue of stellar ultraviolet fluxes. A compilation of absolute stellar fluxes measured by the Sky Survey Telescope (S2/68) aboard the ESRO satellite TD-1*. Unknown, 1978.
- F. van Leeuwen. Validation of the new Hipparcos reduction. *A&A*, 474:653–664, Nov. 2007. doi: 10.1051/0004-6361:20078357.
- J. J. Wang, A. Rajan, J. R. Graham, D. Savransky, P. J. Ingraham, K. Ward-Duong, J. Patience, R. J. De Rosa, J. Bulger, A. Sivaramakrishnan, M. D. Perrin, S. J. Thomas, N. Sadakuni, A. Z. Greenbaum, L. Pueyo, C. Marois, B. R. Oppenheimer, P. Kalas, A. Cardwell, S. Goodsell, P. Hibon, and F. T. Rantakyrö. Gemini planet imager observational calibrations VIII: characterization and role of satellite spots. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, volume 9147 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, page 55, July 2014. doi: 10.1117/12.2055753.
- S. K. Yi, Y.-C. Kim, and P. Demarque. The Y^2 Stellar Evolutionary Tracks. *ApJS*, 144: 259–261, Feb. 2003. doi: 10.1086/345101.




Figure 1: SCEXAO observations of HD 1160 in *Y* (left), *J* (middle), and *H* (right). Diffracted light from the secondary spider is masked in each panel. The four bright, elongated speckles $2''$ from the star are scattered light from SCEXAO, the result of quilting from the deformable mirror, which can be used to aid image registration. The companions HD 1160 B and C are detected at a high SNR in each image.




Figure 2: A comparison of the GPI Y band filter and the HiCIAO Y band filter. The GPI filter is slightly wider on both the blue and red end in comparison to the HiCIAO filter. This leads to slightly brighter photometry for GPI Y band as compared to our SCEXAO/HiCIAO photometry for HD 1160 B.




Figure 3: The wavelength-collapsed, de-rotated, stack of temporal frames for Gemini Planet Imager Y band observations of HD 1160 B. To remove the slowly varying background due to the smooth speckle halo, the original image (left) is high-pass filtered (right) using a median filter with a box size of 11 pixels ($\sim 5\lambda/D$).




Figure 4: Comparison between the GPI Y band spectrum of HD 1160 B ($0.957 - 1.120 \mu\text{m}$, black squares) and the SPHERE spectrum (red triangles) covering $1.0 - 1.6 \mu\text{m}$. There is good agreement between the SPHERE spectra and our higher resolution GPI spectrum (see §2.2.2) within errors, although the SPHERE spectra are slightly brighter over the Y bandpass. The GPI spectra errors are larger due to the higher resolution (the wavelength bins are $\sim 5.4 \text{ nm}$ for GPI and $\sim 18.5 \text{ nm}$ for SPHERE). We use a combined GPI and SPHERE spectrum for our atmosphere modeling (see 3.3).




Figure 5: Color-Magnitude diagrams comparing SCEXAO/HiCIAO (this work), Gemini/NICI photometry (Nielsen et al. 2012), and the integrated-spectra photometry of SPHERE observations (Maire et al. 2016). The M3–M6 main sequence is from dwarf standards of Kirkpatrick et al. (2010). The M6–T9 main-sequence dwarfs that form the color-magnitude sequence is adopted from Dupuy and Liu (2012). At Y , Dupuy and Liu (2012) integrated spectra for the M6–T9 dwarfs to compute the photometry using the UKIRT/WFCAM Y band (Hewett et al. 2006).




Figure 6: The combined SPHERE and GPI spectra of HD 1160 B are best reproduced by the SPEX spectra of M5.2-M6.8 members of Upper Sco from Dawson et al. (2014). M8 and later members of Upper Sco do not reproduce the semi-smooth slope of the SPHERE spectrum from 1.2-1.6 μm . The SPEX spectra were binned to the resolution of the GPI and SPHERE respectively, and scaled to the corresponding flux levels to minimize the χ^2 . Given that the GPI spectrum is higher resolution and the SPHERE spectrum has greater wavelength coverage, we weighted both equally in our spectral typing procedure (see §3.2). F_λ is normalized by a factor of 10^{-15} . The constant is an offset in unit steps of 1.7×10^{-15} $\text{ergs}/\text{cm}^2/\text{s}$.




Figure 7: Similar to figure 6, the combined SPHERE and GPI spectra of HD 1160 B are best reproduced by the SPEX spectra of M5–M6 field standards from Kirkpatrick et al. (2010) (c.f. Figure 9). Standards M4 or earlier spectra do not reproduce the overall shape of the HD 1160 spectrum, while standards M7 and later also do not accurately produce the shape of the HD 1160 spectrum from 1.3–1.6 μm . The method of comparing standard spectral templates to the spectrum of HD 1160 is identical to figure 6 and detailed in §3.2. F_λ is ed by a factor of 10^{-15} . The constant is an an offset in unit steps of 1.9×10^{-15} ergs/cm²/s.




Figure 8: We find the photometry unable to constrain HD 1160 B’s effective temperature to better than $T_{\text{eff}}=2900 - 3500 \text{ K}$ and surface gravity $\log(g)$ to 3.5 – 5.5 dex. χ^2 distributions for fitting HD 1160 B photometry as a function effective temperature and surface gravity. The horizontal dashed red line corresponds to the 95% (2σ) confidence limit, identifying acceptably fitting models. Data points (circles) below this line represent BT-settl models that are in good agreement with the photometry of HD 1160 B.




Figure 9: Same as Figure 8 except for the spectroscopy. With the $0.95\text{--}1.6\ \mu\text{m}$ spectroscopy we find the best fits for an effective temperature $T_{\text{eff}} = 3000\text{--}3100\ \text{K}$ and surface gravity $\log(g) = 4.0\text{--}4.5$ dex, while $T_{\text{eff}} = 2900\ \text{K}$, $\log g = 4.5$ and $T_{\text{eff}} = 3000\ \text{K}$, $\log g = 5.0$ are also marginally acceptable.




Figure 10: Example model fitting both the photometry (left) and spectroscopy (right) of HD 1160 B: $T_{\text{eff}} = 3000 \text{ K}$, $\log g = 4.0$. The photometric points are denoted by gray dots. In the left panel, the model is depicted as a thick black line and horizontal magenta lines shows the predicted photometry from the model in photometric bandpasses. In the right panel, the spectrum is denoted by a grey line, and the model is depicted as a magenta dashed line.




Figure 11: SED fit to catalog photometry spanning a wavelength range of 0.16–22 μm with a NextGen atmosphere model. Red symbols are the observed fluxes with errors (horizontal ‘error bars’ are the photometric passband widths) and blue symbols are the model fluxes. We obtain a best-fit effective temperature of $T_{\text{eff}} = 9011 \pm 85 \text{ K}$ and luminosity $\log L/L_\odot = 1.12 \pm 0.07$ dex.




Figure 12: A comparison of the effective temperature and luminosities of HD 1160 A (red point, derived from SED fitting in Figure 11), and HD 1160 C (light blue point, see §3.5) to stellar models. The Dartmouth, Yonsei-Yale and MIST evolutionary models do not extend into the sub-stellar ($< 0.1 M_{\odot}$) regime, precluding a reliable age estimate for HD 1160 B, while the Baraffe15 models do not extend to the mass of HD 1160 A.




Figure 13: Ages derived from the Baraffe15, DSEP, MIST, and Yonsei-Yale isochrones (see Figure 12) for HD 1160 A, B and C. The model ages are derived from the observed T_{eff} and $\log L/L_{\odot}$ for each star. The bright green dotted line denotes the 300 Myr cut off age for the system, given that HD 1160 A is a main-sequence star (Siess et al. 2000).