Graphs of kei and their diameters

Matthew Ashford[∗]

November 1, 2021

Abstract

A kei on [n] can be thought of as a set of maps $(f_x)_{x \in [n]}$, where each f_x is an involution on [n] such that $(x) f_x = x$ for all x and $f_{(x) f_y} = f_y f_x f_y$ for all x and y. We can think of kei as loopless, edge-coloured multigraphs on $[n]$ where we have an edge of colour y between x and z if and only if $(x) f_y = z$; in this paper we show that any component of diameter d in such a graph must have at least 2^d vertices and contain at least 2^{d-1} edges of the same colour. We also show that these bounds are tight for each value of d .

1 Introduction

A kei (or involutive quandle) is a pair (X, \triangleright) , where X is a non-empty set and $\triangleright: X \times X \to X$ is a binary operation such that:

- 1. For any $y, z \in X$, there exists $x \in X$ such that $z = x \triangleright y$;
- 2. Whenever we have $x, y, z \in X$ such that $x \triangleright y = z \triangleright y$, then $x = z$;
- 3. For any $x, y, z \in X$, $(x \triangleright y) \triangleright z = (x \triangleright z) \triangleright (y \triangleright z);$
- 4. For any $x \in X$, $x \triangleright x = x$;
- 5. For any $x, y \in X$, $(x \triangleright y) \triangleright y = x$.

Note that conditions 1 and 2 above are equivalent to the statement that for each y, the map $x \mapsto x \triangleright y$ is a bijection on X.

A quandle is a pair (X, \triangleright) satisfying conditions 1–4 above, while a rack is a pair (X, \triangleright) satisfying conditions 1–3. As mentioned in [\[2\]](#page-8-0), racks originally developed from correspondence between J.H. Conway and G.C. Wraith in 1959, while quandles were introduced independently by Joyce [\[4\]](#page-8-1) and Matveev [\[5\]](#page-8-2) in 1982 as invariants of knots, and kei were first studied by Takasaki [\[8\]](#page-8-3) in 1943. Fenn and Rourke [\[3\]](#page-8-4) provide a history of racks and quandles, while Nelson [\[6\]](#page-8-5) gives an

[∗]Mathematical Institute, University of Oxford, Radcliffe Observatory Quarter, Woodstock Road, Oxford OX2 6GG, UK. E-mail: matthew.ashford1@btinternet.com.

overview of how these structures relate to other areas of mathematics; a recent paper by Stanovsk`y [\[7\]](#page-8-6) gives a thorough survey of the history of research on kei.

As a first example of a kei, note that for any set X, if we define $x \triangleright y = x$ for all $x, y \in X$, we obtain a kei known as the *trivial kei* T_X . Let G be a group and let X be the set of all involutions of G. If we define a binary operation ⊳ : $X \times X \to X$ by $x \triangleright y := y^{-1}xy$, then (X, \triangleright) is a kei; it is an example of a *conjugation quandle*. For a further example, define a binary operation on [n] by setting $i \triangleright j := 2j - i$ (mod *n*); ([*n*], ρ) is known as a *dihedral kei*.

For any kei (X, \triangleright) , we can define a set of involutions $(f_y)_{y \in X}$ by setting $(x) f_y =$ $x \triangleright y$ for all x and y. The following well-known result (see for example, [\[3\]](#page-8-4), [\[2\]](#page-8-0)) gives the correct conditions for a collection of maps $(f_y)_{y \in X}$ to define a kei.

Proposition 1.1. Let X be a set and $(f_x)_{x\in X}$ be a collection of functions each with domain and co-domain X. Define a binary operation $\triangleright : X \times X \to X$ by $x \triangleright y := (x) f_y$. Then (X, \triangleright) is a kei if and only if f_y is an involution for each $y \in X$ and the following conditions hold: for all $y, z \in X$ we have

$$
f_{(y)f_z} = f_z f_y f_z,\tag{1.1}
$$

and for all $x \in X$ we have

$$
(x)f_x = x.\t\t(1.2)
$$

Proof. As noted earlier, each f_y is a bijection; it remains to show that items 3 and 4 in the definition of a kei are equivalent to [\(1.1\)](#page-1-0) and [\(1.2\)](#page-1-1) respectively, while item 5 is equivalent to the statement that each f_y is an involution. This is essentially a reworking of the definition and we omit the simple details. □

This means that we can just as well define a kei on a set X by the set of maps $(f_y)_{y\in X}$, providing they are all involutions satisfying [\(1.1\)](#page-1-0) and [\(1.2\)](#page-1-1). We will move freely between the two definitions, with $x \triangleright y = (x) f_y$ for all $x, y \in X$ unless otherwise stated.

Now observe that any kei on X can be represented by a multigraph on X ; we give each vertex a colour and then put an edge of colour i from vertex j to vertex k if and only if $(j) f_i = k$. This is well-defined as each f_i is an involution, so $(j) f_i = k$ if and only if $(k) f_i = j$. We then remove all loops from the graph; i.e. if $(j) f_i = j$ we don't have an edge of colour i incident to j.

It will be helpful to recast the representation of kei by multigraphs in a slightly different setting. Let V be a finite set and let $\sigma \in \text{Sym}(V)$ be an involution; then we can define a simple graph G_{σ} on V by letting $uv \in E(G_{\sigma})$ if and only if $u \neq v$ and $(u)\sigma = v$. As σ is a disjoint product of transpositions, we see that G_{σ} consists of a partial matching and some isolated vertices. We can now extend this definition to the case of multiple involutions in a natural way.

Definition 1.2. Suppose $\Sigma = {\sigma_1, \ldots, \sigma_k} \subseteq \text{Sym}(V)$ is a set of involutions on a set V. Define a loopless multigraph $G_{\Sigma} = (V, E)$ with a k-edge-colouring by putting an edge of colour i from u to v if and only if $u \neq v$ and $(u)\sigma_i = v$.

Figure 1: Graphical representations of two kei. Both of these are subquandles of conjugation quandles; S_3 on the left and S_4 on the right.

We also define the *reduced* graph G_{Σ}^{0} to be the simple graph on V obtained by setting $e = uv \in E(G_{\Sigma}^0)$ if and only if there is at least one edge from u to v in G_{Σ} .

Observe that if $\Sigma' \subseteq \Sigma$, then $G_{\Sigma'}$ is a subgraph of G_{Σ} . Now let us return specifically to kei.

Definition 1.3. Let $K = (X, \triangleright)$ be a kei, and let $(f_y)_{y \in X}$ be the associated maps. For any $S \subseteq X$, define $\Sigma_S = \{f_y | y \in S\}$. Then by G_S we mean the multigraph G_{Σ_S} in the sense of Definition [1.2;](#page-1-2) G_S thus has an associated $|S|$ -edge-colouring, although if $|S| = 1$ we may not necessarily consider G_S as being coloured. We will also write $G_K = G_X$, indicating the graph for the whole kei.

Figure [1](#page-2-0) gives two examples of graphs representing kei. Before stating our main result, we will need some more definitions. Let (X, \triangleright) be a kei; then a *subkei* of (X, \triangleright) is a kei $(Y, \triangleright|_{Y \times Y})^1$ $(Y, \triangleright|_{Y \times Y})^1$ where $Y \subseteq X$. Thus a subset $Y \subseteq X$ forms a subkei if and only if for all $y, z \in Y$, $(z) f_y \in Y$. For any $T \subseteq X$ and $u, v \in X$, denote by $d_T(u, v)$ the graph distance between u and v in the graph G_T . As G_T^0 is the simple graph on X formed by ignoring all colours and multiple edges, $d_T(u, v)$ is clearly identical to the graph distance between u and v in the reduced graph G_T^0 . We will prove the following result.

Theorem 1.4. Let (X, \triangleright) be a kei, and let (S, \triangleright) be a subkei. Let $C \subseteq X$ span a component of G_S (or equivalently of G_S^0) and suppose that $G_S[C]$ has diameter d. Then $|C| \geq 2^d$, and further there exists some $k \in S$ such that there are at least 2^{d-1} edges of colour k in $G_S[C]$.

The remainder of the paper is organised as follows. In Section [2](#page-3-0) we show how to construct a large number of shortest paths between two vertices u and v that are connected in G_S . In Section [3](#page-4-0) we will show that any sequence of colours occurring in order (from u to v) on a shortest path corresponds to a different vertex connected to u, which will prove the result. In Section [4](#page-7-0) we give examples to show that Theorem [1.4](#page-2-2) is tight for all values of d.

¹The notation ⊳ $|Y \times Y|$ in the above context will always be abbreviated to ⊳, with the restriction to the subset Y left implicit.

Figure 2: An example with $i = 4$. The colour c_i is represented by red, with the colours on the lower path being the colours c' , c'' etc. found as we progress from right to left along the upper path.

2 Shortest paths

We begin by showing that there are many shortest paths between pairs of vertices.

Lemma 2.1. Let $S \subseteq X$ be a subkei, and let $u, v \in X$ such that $d_S(u, v) = d \in$ $[2,\infty);$ let $u = v_0v_1 \cdots v_d = v$ be a path of length d in G_S between u and v. For each i, let c_i be the colour of the $v_{i-1}v_i$ edge on the path, so $(v_i)f_{c_i} = v_{i-1}$. Now fix some $i \geqslant 2$ and define $w_k = (v_k) f_{c_i}$ for $k = 0, \ldots, i - 2$. Then the vertices $w_0, \ldots, w_{i-2}, v_0, \ldots, v_d$ are all distinct and $v_0w_0 \cdots w_{i-2}v_i \cdots v_d$ is a (shortest) uv path in G_S .

Proof. (See Figure [2](#page-3-1) throughout). For any $j \leq i-2$, denote by $P(j)$ the statement that the vertices $w_j, \ldots, w_{i-2}, v_0, \ldots, v_d$ are all distinct, and that the path $v_0 \cdots v_j w_j \cdots w_{i-2} v_i \cdots v_d$ is a (shortest) uv-path in G_S . Then the lemma is the statement $P(0)$; we will prove that $P(j)$ holds for all j by reverse induction.

Firstly, as $w_{i-2} = (v_{i-2})f_{c_i}$ and $v_i = (v_{i-1})f_{c_i}$ we see that $w_{i-2} \notin \{v_{i-1}, v_i\}$ (or there would be a vertex with two incident edges of the same colour). If we had $w_{i-2} = v_l$ for some $l > i$ then $v_0 \cdots v_{i-2} v_l \cdots v_d$ would be a uv -path in G_S of length $d - (l - i + 1) < d$, a contradiction. Now observe that

$$
w_{i-2} := (v_{i-2})f_{c_i} = (v_{i-1})f_{c_{i-1}}f_{c_i} = (v_i)f_{c_i}f_{c_{i-1}}f_{c_i} = (v_i)f_{c'},
$$

where $c' = (c_{i-1})f_{c_i}$; note that $c' \in S$ as S is a subkei and so there is an edge in G_S between v_i and w_{i-2} . So now suppose $w_{i-2} = v_l$ for $l < i-1$; then $v_0 \cdots v_l =$ $w_{i-2}v_i\cdots v_d$ is a uv-path in G_S of length $d-(i-l-1)$, again a contradiction. So $w_{i-2} \neq v_i$ for any l, and we can use the $v_{i-2}w_{i-2}$ and $w_{i-2}v_i$ edges, of colours c_i and c' respectively, to construct a uv-path $v_0 \cdots v_{i-2} w_{i-2} v_i \cdots v_d$ of length d in G_S . This establishes $P(i-2)$.

So now take $j < i - 2$ and suppose $P(j + 1)$ holds. As $v_{j+1}w_{j+1}, \ldots, v_{i-2}w_{i-2}$ and $v_{i-1}v_i$ are all edges of colour $c_i, w_j \neq v_l$ for $j < l \leq i$ and it is also not equal to any other w_k . As before, if $w_j = v_l$ for $l > i$ we can construct a uv-path in G_S of length $d - (l - i + 1)$, a contradiction. Now observe that

$$
w_j := (v_j) f_{c_i} = (v_{j+1}) f_{c_{j+1}} f_{c_i} = (w_{j+1}) f_{c_i} f_{c_{j+1}} f_{c_i} = (w_{j+1}) f_{c''},
$$

where $c'' = (c_{j+1})f_{c_i} \in S$, so there is an edge in G_S between w_j and w_{j+1} . Now suppose $w_j = v_l$ for some $l \leq j$; then $v_0 \cdots v_l = w_j w_{j+1} \cdots w_{i-2} v_i \cdots v_d$ is a uv path in G_S of length $d - (j - l + 1)$, a contradiction (note the existence of the $w_{j+1}v_i$ -path follows from the assumed $P(j+1)$). Hence $w_j \neq v_l$ for any l, and we can use the v_jw_j and w_jw_{j+1} edges, of colours c_i and c'' respectively, to construct a uv-path $v_0 \cdots v_j w_j \cdots w_{i-2} v_i \cdots v_d$ of length d in G_S . Thus $P(j)$ holds, and so by reverse induction $P(0)$ holds, proving the lemma. П

Corollary 2.2. Let P be any shortest uv-path in G_S . Then no two edges of P have the same colour.

Proof. Suppose there is a path $u = v_0 \cdots v_d = v$ and a colour $c \in S$ such that $(v_{j-1})f_c = v_j$ and $(v_{i-1})f_c = v_i$ for some $j < i-1$ (we can't have $j = i-1$ as then v_j is incident to two edges of colour c). From the lemma, with $c = c_i$, $w_j = (v_j) f_{c_i} \neq v_l$ for any l, but here $w_j = (v_j) f_c = v_{j-1}$, a contradiction. \Box

3 Sequences of elements

Suppose we are considering a shortest path $P : (u = v_0, \ldots, v_d = v)$ in G_S . In the light of Corollary [2.2,](#page-4-1) and to ease notation, we will assume without loss of generality that $X = [n]$ and $[d] \subseteq S$, and that the edge between v_{i-1} and v_i in P is of colour *i*. Now for each strictly increasing sequence $\mathbf{s} = (a_1, \ldots, a_r)$ of elements from $[d]$, define

$$
u_{\mathbf{s}}=(u)f_{a_1}\cdots f_{a_r},
$$

where we note $u_s \in S$ as S is a subkei. Now define $U_0 = \{u\}$ and for $r = 1, \ldots, d$, define

$$
U_r = \{u_s \mid \mathbf{s} \text{ is strictly increasing, } |\mathbf{s}| = r\},
$$

so in particular $U_d = \{v\}$. Now let $\mathbf{e}_i = (1, \ldots, i)$ for all i; then $v_i = (u)f_1 \cdots f_i =$ $u_{\mathbf{e}_i}$ and so $v_i \in U_i$ for all i. We also have the property that as **s** is increasing, $a_i \geq i$ for any i, with equality if and only if the subsequence consisting of the first i terms of **s** is the canonical sequence e_i .

We will show that any strictly increasing sequence s can appear at the start of a shortest uv-path, and that any such path passes sequentially through each of U_0, U_1, \ldots, U_d .

Lemma 3.1. Let $S \subseteq X$ be a subkei, and let $u, v \in X$ be such that $d_S(u, v) =$ $d \in [2,\infty)$; let $u = v_0v_1 \cdots v_d = v$ be a path of length d in G_S between u and v, where the $v_{i-1}v_i$ edge is of colour i for each i. Let $\mathbf{s} = (a_1, \ldots, a_r)$ be a strictly increasing sequence of elements from [d]. Then there is a shortest path P_s : (u = $x_0x_1 \cdots x_{a_r-1}x_{a_r} = v_{a_r} \cdots v_d$ such that $x_k \in U_k$ for $1 \leq k \leq a_r$ and the $x_{k-1}x_k$ edge in P_s is of colour a_k for $1 \leq k \leq r$ (which means in particular that $x_r = u_s$).

Proof. We shall prove the following stronger statement by induction on r : there exists a shortest path P_s : $(u = x_0x_1 \cdots x_{a_r-1}x_{a_r} = v_{a_r} \cdots v_d)$ such that for $1 \leqslant k < a_r$

$$
x_k = u_t \tag{3.1}
$$

for a (strictly increasing) sequence t of length k whose largest element is at most a_r , and that for $1 \leq k \leq r$ the $x_{k-1}x_k$ edge in P_s is of colour a_k . These statements clearly imply the result.

First consider the base case $r = 1$; if $s = e_1 = (1)$ then the original path P suffices as the first edge is of colour 1, so $v_1 = (u)f_1 = u_{(1)}$. Hence we may assume $a_1 > 1$. Then applying Lemma [2.1](#page-3-2) to P with $i = a_1$, there exists a shortest path P' : $(u = v_0w_0\cdots w_{a_1-2}v_{a_1}\cdots v_d)$ in G_S , with an edge of colour a_1 between v_k and w_k for $0 \leq k \leq a_1 - 2$; in particular, the first edge of P' is of colour a_1 . Now for $k \leq a_1 - 2$, $w_k = (v_k) f_{a_1} = (u_{\mathbf{e}_k}) f_{a_1}$, so $w_k = u_{(\mathbf{e}_k, a_1)} \in U_{k+1}$. So we obtain [\(3.1\)](#page-5-0) by setting $x_k = w_{k-1}$ for $1 \leq k < a_1$ and putting $P_s = P'$.

So now take $r > 1$ and assume the result for smaller r. If $a_r = r$ we have $\mathbf{s} = \mathbf{e}_r$, and in this situation the original path P suffices as the first r edges are of colours 1, ..., r and $v_k \in U_k$ for $1 \leq k \leq r$. So we can assume that $a_r > r$.

By applying the inductive hypothesis to the sequence $\mathbf{s}' = (a_1, \ldots, a_{r-1})$ we see that there exists a path

$$
P_{\mathbf{s}'} : (u = y_0 y_1 \cdots y_{a_{r-1}-1} y_{a_{r-1}} = v_{a_{r-1}} \cdots v_d)
$$

such that, for $1 \leq k \leq a_{r-1}$, $y_k = u_t$ for a strictly increasing sequence t whose largest element is at most a_{r-1} . We also have that the $y_{k-1}y_k$ edge is of colour a_k for $1 \leq k < r$, but we don't know the colours of the edges $y_{k-1}y_k$ for $r \leq k \leq a_{r-1}$. But as $a_r > a_{r-1}$ the $v_{a_r-1}v_{a_r}$ edge of colour a_r is still present in $P_{s'}$; hence we can apply Lemma [2.1](#page-3-2) to $P_{s'}$, with $i = a_r$, to obtain a shortest path P'' : $(u =$ $v_0w_0 \cdots w_{a_r-2}v_{a_r}\cdots v_d$ in G_S , with an edge of colour a_r between y_k and w_k for $0 \leq k \leq \min\{a_{r-1}, a_r - 2\}$ (note that $a_r - 2 < a_{r-1}$ if and only if $a_r = a_{r-1} + 1$). We aim to show that for all $0 \leq k \leq a_r - 2$, $w_k = u_{\mathbf{t'}}$ for some sequence $\mathbf{t'}$ of length $k+1$ whose largest element is at most a_r . The proof will vary slightly depending on whether $a_r = a_{r-1} + 1$ (see Figure [3\)](#page-6-0) or $a_r > a_{r-1} + 1$ (see Figure [4\)](#page-6-1).

Fix a k such that $0 \le k \le a_{r-1} - 1 \le a_r - 2$; by the inductive hypothesis (specifically [\(3.1\)](#page-5-0)) $y_k = u_t$ for some sequence t whose kth and largest element is at most a_{r-1} . But $w_k = (y_k) f_{a_r} = (u_t) f_{a_r}$, so $w_k = u_{(t,a_r)}$; as $a_r > a_{r-1}$ we have $w_k \in U_{k+1}$, where $w_k = u_{t'}$ for some sequence t' whose largest element is at most a_r . If $a_r = a_{r-1} + 1$ then we have considered all the vertices w_1, \ldots, w_{a_r-2} .

Now suppose that $a_r > a_{r-1} + 1$; in this case we also have an edge of colour a_r between v_k and w_k for $a_{r-1} \leq k \leq a_r - 2$. But as before, $w_k = (v_k) f_{a_r}$ $(u_{\mathbf{e}_k})f_{a_r} = u_{(\mathbf{e}_k,a_r)}$ for any such k, so $w_k \in U_{k+1}$ and $w_k = u_{\mathbf{t}'}$ for some sequence t' whose largest element is at most a_r . Thus we have the desired result on w_k for the entire range $0 \leq k \leq a_r - 2$.

Now as $a_r > r$ and $a_{r-1} \geq r-1$ we have $\min\{a_{r-1}, a_r - 2\} \geq r-1$, so there is always an edge of colour a_r between y_{r-1} and w_{r-1} . Consider the path P'''

Figure 3: Suppose $\mathbf{s} = (2,3)$ and that edges of colour 2 are red while those of colour 3 are blue. The top path $P_{s'}$ corresponds to the sequence (2), and the bottom path P'' is a result of applying Lemma [2.1.](#page-3-2) The new path P''' is shown in bold.

Figure 4: Suppose we now have the sequence $s = (2, 3, 5)$, with edges of colour 5 being violet. The top path $P_{s'}$ is the (relabelled) bold path from Figure [3,](#page-6-0) while the bottom path is again a result of applying Lemma [2.1;](#page-3-2) note the additional edges. The new path P''' is shown in bold.

obtained by using the first $r-1$ edges of $P_{s'}$, the edge $y_{r-1}w_{r-1}$ and then the remaining edges of P'' ; we obtain

$$
P''' : (uy_1 \cdots y_{r-1}w_{r-1} \cdots w_{a_r-2}v_{a_r} \cdots v_d),
$$

where the first r edges are of colours a_1, \ldots, a_r . We get the result by putting $x_k = y_k \in U_k$ for $1 \leq k < r$ and $x_k = w_{k-1} \in U_k$ for $r \leq k < a_r$, and setting $P_{\rm s}=P^{\prime\prime\prime}$. □

We can now use this result to show that different sequences correspond to distinct vertices.

Corollary 3.2. Let s and t be distinct, strictly increasing sequences from $[d]$. Then $u_s \neq u_t$.

Proof. Let $|s| = q$ and $|t| = r$, where we may assume $q \leq r$, and suppose that $u_{s} = u_{t}$. Write $s = (a_{1}, \ldots, a_{q})$ and $t = (b_{1}, \ldots, b_{r})$ and consider the paths $P_{\bf s}$: $(ux_1 \cdots x_{a_q} = v_{a_q} \cdots v_d)$ and $P_{\bf t}$: $(uy_1 \cdots y_{b_r} = v_{b_r} \cdots v_d)$ as described in Lemma [3.1;](#page-4-2) note that $x_q = u_s = u_t = y_r$. Thus we may replace the uy_r -segment of P_t with the ux_q-segment of P_s to obtain a uv-walk in G_S of length $d - (r - q)$, a contradiction if $q < r$. This shows that $U_q \cap U_r = \emptyset$ for $q \neq r$.

So suppose $q = r$ (so $x_r = y_r$) and for now that $a_r \neq b_r$; we may assume that $a_r < b_r$. Now replace the ux_r -segment of P_s with the uy_r -segment of P_t to obtain a new uv-path

$$
P': (uy_1 \cdots y_r = x_r \cdots x_{a_r-1} x_{a_r} = v_{a_r} \cdots v_{b_r-1} v_{b_r} \cdots v_d),
$$

where we note this is a path as both P_s and P_t pass sequentially through the pairwise disjoint sets U_0, U_1, \ldots, U_d . But both the $y_{r-1}y_r$ and $v_{b_r-1}v_{b_r}$ edges on P' are of colour b_r , contradicting Corollary [2.2.](#page-4-1)

Now note that in any case $x_{r-1} = (x_r) f_{a_r}$ and $y_{r-1} = (y_r) f_{b_r}$, so if $a_r = b_r$ we have $x_{r-1} = y_{r-1}$. An easy inductive argument (and the fact that $s \neq t$) shows that there exists some $p \leq r$ such that $x_i = y_i$ for $p \leq i \leq r$ but $a_p \neq b_p$, so we may apply the above argument to the sequences $\mathbf{s}' = (a_1, \ldots, a_p)$ and $\mathbf{t}' = (b_1, \ldots, b_p)$ to get a contradiction, thus proving the result.

This allows us to prove the main result.

Proof of Theorem [1.4.](#page-2-2) If $d = 0$ or 1 then the result is trivial, so suppose $d \ge 2$. $G_S[C]$ has diameter d, so there exist $u, v \in C$ such that $d_S(u, v) = d$; as before, assume that $X = [n]$ and $[d] \subseteq S$, and that there is a shortest wv-path using edges of colours $1, \ldots, d$ sequentially. There are 2^d strictly increasing sequences of elements from [d] (including the empty sequence) and from Corollary [3.2](#page-6-2) these correspond to 2^d distinct vertices (including u). Hence $|C| \geq 2^d$.

Note that there are 2^{d-1} strictly increasing sequences from [d−1]; adding d to the end of such a sequence s' gives a strictly increasing sequence s from [d]. But then $(u_{s})f_d = u_s$ so the edge $u_{s'}u_s$ is of colour d for each such sequence s'; as all vertices are distinct this gives us 2^{d-1} edges of colour d. \Box

4 Extremal examples

In this section we construct a family of extremal examples, one for each d. Let $X_d := \{u_1, \ldots, u_d\} \cup \{0, 1\}^d$ be a set of $2^d + d$ elements, and for each $v \in \{0, 1\}^d$ and $S \subseteq [d]$, define v_S to be the element of $\{0,1\}^d$ obtained by changing only the coordinates in S, i.e.

$$
\{i \in [d] \mid (v_S)_i \neq v_i\} = S.
$$

We will now define a set of bijections on X_d ; for each $1 \leq i \leq d$, set $(u_j) f_{u_i} = u_j$ for all j and $(v)f_{u_i} = v_{\{i\}}$ for all $v \in \{0,1\}^d$. We will also set $f_v = \iota$ for all $v \in \{0,1\}^d$.

Note that $(v_{\{i\}})_{\{i\}} = v$ and thus each map is an involution; as we also have $(x) f_x = x$ for each $x \in X_d$, we need only show that [\(1.1\)](#page-1-0) holds to prove that we have defined a kei K. Note that for any distinct $i, j \in [d]$ and $v \in \{0, 1\}^d$,

$$
(v)f_{u_i}f_{u_j} = (v_{\{i\}})f_{u_j} = v_{\{i,j\}} = (v_{\{j\}})f_{u_i} = (v)f_{u_j}f_{u_i},
$$

and it follows that all the maps $(f_x)_{x \in X_d}$ commute. Hence (1.1) reduces to the statement that $f(y)f_z = f_y$ for all y, z ; this is easily seen to be true.

Figure 5: An example of a kei on X_3 .

Now consider the graph G_K ; by construction, G_K consists of d isolated vertices and a coloured copy of the cube Q_d (see Figure [5\)](#page-8-7). The cube has diameter d, size 2^d and contains 2^{d-1} edges of each colour u_1, \ldots, u_d .

Acknowledgements

This paper forms part of the author's Ph.D. thesis [\[1\]](#page-8-8), supervised by Oliver Riordan; his assistance in preparing this paper is gratefully acknowledged. The thesis contains some applications of this result.

References

- [1] Matthew Ashford. Graphs of Algebraic Objects. PhD thesis, University of Oxford, 2016.
- [2] Simon R Blackburn. Enumerating finite racks, quandles and kei. The Electronic Journal of Combinatorics, 20(3):P43, 2013.
- [3] Roger Fenn and Colin Rourke. Racks and links in codimension two. Journal of Knot theory and its Ramifications, 1(04):343–406, 1992.
- [4] David Joyce. A classifying invariant of knots, the knot quandle. Journal of Pure and Applied Algebra, 23(1):37–65, 1982.
- [5] Sergei Vladimirovich Matveev. Distributive groupoids in knot theory. Sbornik: Mathematics, 47(1):73–83, 1984.
- [6] Sam Nelson. The combinatorial revolution in knot theory. Notices of the AMS, 58(11):1553–1561, 2011.
- [7] David Stanovsky. The origins of involutory quandles. arXiv preprint [arXiv:1506.02389](http://arxiv.org/abs/1506.02389), 2015.
- [8] Mituhisa Takasaki. Abstractions of symmetric functions. Tohuko Mathematical Journal, 49:143–207, 1943.