

# TRANSFORMING A MATRIX INTO A STANDARD FORM

AKIHIRO MUNEMASA AND PRITTA ETRIANA PUTRI

ABSTRACT. We show that every matrix all of whose entries are in a fixed subgroup of the group of units of a commutative ring with identity is equivalent to a standard form. As a consequence, we improve the proof of Theorem 5 in D. Best, H. Kharaghani, H. Ramp [Disc. Math. 313 (2013), 855–864].

## 1. INTRODUCTION

Throughout this note, we let  $R$  be a commutative ring with identity. We fix a subgroup  $T$  of the group of units of  $R$ , and set  $T_0 = T \cup \{0\}$ . The set of  $m \times n$  matrices with entries in  $T_0$  is denoted by  $T_0^{m \times n}$ . If  $T = \{z \in \mathbb{C} : |z| = 1\}$ , then  $W \in T_0^{n \times n}$  is called a *unit weighing matrix of order  $n$  with weight  $w$*  provided that  $WW^* = wI$  where  $W^*$  is the transpose conjugate of  $W$ . Unit weighing matrices are introduced by D. Best, H. Kharaghani, and H. Ramp in [1, 2]. Moreover, a unit weighing matrix is known as a unit Hadamard matrix if  $w = n$  (see [3]). A unit weighing matrix in which every entry is in  $\{0, \pm 1\}$  is called a *weighing matrix*. We refer the reader to [4] for an extensive discussion of weighing matrices, and to [5] for more information on applications of weighing matrices.

The study on the number of inequivalent unit weighing matrices was initiated in [1]. Also, observing the number of weighing matrices in standard form leads to an upper bound on the number of inequivalent unit weighing matrices [1]. In this work, we will introduce a standard form of an arbitrary matrix in  $T_0^{m \times n}$  and show that every matrix in  $T_0^{m \times n}$  is equivalent to a matrix in standard form.

We equip  $T_0$  with a total ordering  $\prec$  satisfying  $\min(T_0) = 1$  and  $\max(T_0) = 0$ . Moreover, let  $\mathbf{a} = (a_1, \dots, a_n)$  and  $\mathbf{b} = (b_1, \dots, b_n)$  be arbitrary row vectors with entries in  $T_0$ . If  $k$  is the smallest index such that  $a_k \neq b_k$ , then we write  $\mathbf{a} < \mathbf{b}$  provided  $a_k \prec b_k$ . We write  $\mathbf{a} \leq \mathbf{b}$  if  $\mathbf{a} < \mathbf{b}$  or  $\mathbf{a} = \mathbf{b}$ . If  $\mathbf{a}_1, \dots, \mathbf{a}_m$  are row vectors of a matrix  $A \in T_0^{m \times n}$  and  $\mathbf{a}_1 < \dots < \mathbf{a}_m$ , then we say that the rows of  $A$  are in *lexicographical order*.

**Definition 1.1.** We say that a matrix in  $T_0^{m \times n}$  is in *standard form* if the following conditions are satisfied:

- (S1) The first non-zero entry in each row is 1.
- (S2) The first non-zero entry in each column is 1.
- (S3) The first row is ones followed by zeros.

---

*Date:* November 11, 2016.

*2010 Mathematics Subject Classification.* 15B34, 05B20.

*Key words and phrases.* monomial matrix, weighing matrix, Hadamard matrix, permutation matrix.

(S4) The rows are in lexicographical order according to  $\prec$ .

The subset of  $T_0^{m \times m}$  consisting of permutation matrices, nonsingular diagonal matrices and monomial matrices, are denoted respectively, by  $\mathbb{P}_m, \mathbb{D}_m$  and  $\mathbb{M}_m$ . Then  $\mathbb{M}_m = \mathbb{P}_m \mathbb{D}_m$ .

**Definition 1.2.** For  $A, B \in T_0^{m \times n}$ , we say that  $A$  is *equivalent* to  $B$  if there exist monomial  $T_0$ -matrices  $M_1$  and  $M_2$  such that  $M_1 A M_2 = B$ .

We will restate the proof of [1, Theorem 5] as the following algorithm.

**Algorithm 1.3.** Let  $W$  be an arbitrary unit weighing matrix.

- (1) We multiply each  $i$ th row of  $W$  by  $r_i^{-1}$  where  $r_i$  is the first non-zero entry in  $i$ th row. Denote the obtained matrix by  $W^{(1)}$ .
- (2) Let  $c_j$  be the first non-zero entry in  $j$ th column of  $W^{(1)}$ . Let  $W^{(2)}$  obtained from  $W^{(1)}$  by multiplying each  $j$ th column by  $c_j^{-1}$ .
- (3) Permute the columns of  $W^{(2)}$  so that the first row has  $w$  ones. Denote the resulting matrix by  $W^{(3)}$ .
- (4) Let  $W^{(4)}$  be a matrix obtained from  $W^{(3)}$  by sorting the rows of  $W^{(3)}$  lexicographically with the ordering  $\prec$ .

Then  $W^{(4)}$  is in standard form.

The steps (1)–(4) in Algorithm 1.3 was used in order to prove Theorem 5 in [1]. However, we provide a counterexample to show that this algorithm does not produce a standard form.

**Counterexample 1.4.** The matrix

$$W = \begin{bmatrix} 1 & -i & i & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & i & i \\ 1 & 0 & 0 & -1 & -i & i \\ 1 & 0 & 0 & -1 & i & -i \\ 0 & 1 & 1 & 0 & -i & -i \\ 1 & i & -i & 1 & 0 & 0 \end{bmatrix}$$

is a unit weighing matrix, where  $i$  is a 4th root of unity in  $\mathbb{C}$ . Also, we equip the set  $\{0, \pm i, \pm 1\}$  with a total ordering  $\prec$  defined by  $1 \prec -1 \prec i \prec -i \prec 0$ . Since the first nonzero entry in each row of  $W$  is one,  $W^{(1)} = W$ . Applying step (2), we obtain

$$W^{(2)} = \begin{bmatrix} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & i & -i & 0 & 1 & 1 \\ 1 & 0 & 0 & -1 & -1 & 1 \\ 1 & 0 & 0 & -1 & 1 & -1 \\ 0 & i & -i & 0 & -1 & -1 \\ 1 & -1 & -1 & 1 & 0 & 0 \end{bmatrix}.$$

Notice that the first row of  $W^{(2)}$  is all ones followed by zeros. So,  $W^{(3)} = W^{(2)}$ . Finally, by applying the last step of the algorithm, we have

$$W^{(4)} = \begin{bmatrix} 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & -1 & -1 & 1 & 0 & 0 \\ 1 & 0 & 0 & -1 & 1 & -1 \\ 1 & 0 & 0 & -1 & -1 & 1 \\ 0 & i & -i & 0 & 1 & 1 \\ 0 & i & -i & 0 & -1 & -1 \end{bmatrix}.$$

We see that  $W^{(4)}$  is not in standard form. So, we conclude that the algorithm does not produce a matrix in standard form as claimed.

This counterexample shows that the additional steps are needed to complete the proof of Theorem 5 in [1]. In the next section, we will prove a more general theorem than [1, Theorem 5] by showing that every matrix in  $T_0^{m \times n}$  is equivalent to a matrix that is in standard form.

## 2. MAIN THEOREM

In addition to the conditions (S1)–(S4) in Definition 1.1, we will consider the following condition:

(S3)' The first nonzero row is ones followed by zeros.

Note that (S3)' is weaker than (S3). The condition (S3)' is crucial in the proof of Lemma 2.1, where we encounter a matrix whose first row consists entirely of zeros.

**Lemma 2.1.** *Let*

$$A = [A_1 \ A_2] \in T_0^{m \times (n_1 + n_2)},$$

where  $A_i \in T_0^{m \times n_i}$ ,  $i = 1, 2$ . Then there exist  $P \in \mathbb{P}_m$  and  $M \in \mathbb{M}_{n_2}$  such that  $PA_2M$  satisfies (S2) and (S3)', and  $[PA_1 \ PA_2M]$  satisfies (S4).

*Proof.* Without loss of generality, we may assume  $A_1$  satisfies (S4). Then there exist row vectors  $\mathbf{a}_1, \dots, \mathbf{a}_k$  of  $A_1$  such that  $\mathbf{a}_1 < \dots < \mathbf{a}_k$ , and positive integers  $m_1, \dots, m_k$  such that

$$A_1 = \begin{bmatrix} \mathbf{1}_{m_1}^\top & & \\ & \ddots & \\ & & \mathbf{1}_{m_k}^\top \end{bmatrix} \begin{bmatrix} \mathbf{a}_1 \\ \vdots \\ \mathbf{a}_k \end{bmatrix},$$

where  $\sum_{i=1}^k m_i = m$ . Write

$$A_2 = \begin{bmatrix} B_1 \\ \vdots \\ B_k \end{bmatrix},$$

where  $B_i \in T_0^{m_i \times n_2}$  for  $i = 1, 2, \dots, k$ . We may assume  $B_1 \neq 0$ , since otherwise the proof reduces to establishing the assertion for the matrix  $A$  with the first  $m_1$  rows deleted. Let  $\mathbf{b}$  be a row vector of  $B_1$  with maximum number of nonzero components. Then there exists  $M \in \mathbb{M}_{n_2}$  such that the vector  $\mathbf{b}M$  constitutes ones followed by zeros. Moreover, for each  $i \in \{1, \dots, k\}$ , there exists  $P_i \in \mathbb{P}_{m_i}$  such that the rows of  $P_i B_i M$  are in lexicographic order. It follows that  $\mathbf{b}M$  is the first row of  $P_1 B_1 M$ , that

is also the first row of  $PA_2M$ . Set  $P = \text{diag}(P_1, \dots, P_k)$ . Then  $PA_2M$  satisfies (S3). Since  $PA_1 = A_1$ , we see that  $[PA_1 \ PA_2M]$  satisfies (S4).

With the above notation, we prove the assertion by induction on  $n_2$ . First we treat the case where  $\mathbf{b}M = \mathbf{1}$ . This in particular includes the case where  $n_2 = 1$ , the starting point of the induction. In this case, the first row of  $PA_2M$  is  $\mathbf{1}$ , hence  $PA_2M$  satisfies (S2). The other assertions have been proved already.

Next we consider the case where  $\mathbf{b}M = [\mathbf{1}_{n_2-n'_2} \ \mathbf{0}_{n'_2}]$ , with  $0 < n'_2 < n_2$ . Define  $A'_1 \in T_0^{m \times (n_1+n_2-n'_2)}$  and  $A'_2 \in T_0^{m \times n'_2}$  by setting  $[A'_1 \ A'_2]$  to be the matrix obtained from  $[A_1 \ PA_2M]$  by deleting the first row. By inductive hypothesis, there exist  $P' \in \mathbb{P}_{m-1}$  and  $M' \in \mathbb{M}_{n'_2}$  such that  $P'A'_2M'$  satisfies (S2) and (S3)', and  $[P'A'_1 \ P'A'_2M']$  satisfies (S4). By our choice of  $\mathbf{b}$ , the row vector  $\mathbf{b}M$  is lexicographically the smallest member among the rows of  $P_1B_1M$ , and the same is true among the rows of the matrix  $P_1B_1M''$ , where

$$M'' = M \begin{bmatrix} I_{n_2-n'_2} & 0 \\ 0 & M' \end{bmatrix}.$$

It follows that the matrix

$$\begin{bmatrix} 1 & 0 \\ 0 & P' \end{bmatrix} [A_1 \ PA_2M''] = \begin{bmatrix} * & 0 \\ P'A'_1 & P'A'_2M' \end{bmatrix}$$

satisfies (S4). Set

$$P'' = \begin{bmatrix} 1 & 0 \\ 0 & P' \end{bmatrix} P.$$

Since  $P'A'_2M'$  satisfies (S2), while the first row of  $P''A_2M''$  is the same as that of  $PA_2M$  which is  $[\mathbf{1}_{n_2-n'_2} \ \mathbf{0}_{n'_2}]$ , the matrix  $P''A_2M''$  satisfies both (S2) and (S3)'. We have already shown that the matrix  $[P''A_1 \ P''A_2M'']$  satisfies (S4).  $\square$

**Lemma 2.2.** *Under the same assumption as in Lemma 2.1, there exist  $M_1 \in \mathbb{M}_m$  and  $M_2 \in \mathbb{M}_{n_2}$  such that  $[M_1A_1 \ M_1A_2M_2]$  satisfies (S1) and (S4), and  $M_1A_2M_2$  satisfies (S2) and (S3)'.*

*Proof.* We will prove the assertion by induction on  $m$ . Suppose  $m = 1$ . It is clear that every single row vector always satisfies (S4). Also, every single row vector satisfying (S3)' necessarily satisfies (S2). Now, if  $A_1 = 0$  or  $n_1 = 0$ , then there exists  $M_2 \in \mathbb{M}_{n_2}$  such that  $A_2M_2$  satisfies (S3)' and hence (S1) is satisfied. If  $A_1 \neq 0$ , then there exist  $a \in T$  and  $M_2 \in \mathbb{M}_{n_2}$  such that  $aA_1$  satisfies (S1) and  $aA_2M_2$  satisfies (S3)'.

Assume the assertion is true up to  $m - 1$ . First, we consider the case where  $A_1 = 0$  or  $n_1 = 0$ . Without loss of generality, we may assume  $A_2 \neq 0$ . Furthermore, we may assume that the first row and the first column of  $A_2$  are ones followed by zeros. Then there exists  $P' \in \mathbb{P}_{n_2}$  such that

$$A_2P' = \left[ \begin{array}{ccc|c} 1 & \mathbf{1} & 0 & 0 \\ \mathbf{1}^T & B_1 & B_2 & 0 \\ 0 & C_1 & & C_2 \end{array} \right]$$

where  $B_2 \in T_0^{m_1 \times t}$  has no zero column. By Lemma 2.1, there exist  $P \in \mathbb{P}_{m_1}$  and  $M \in \mathbb{M}_t$  such that  $PB_2M$  satisfies (S2) and (S3)' and  $[PB_1 \ PB_2M]$  satisfies (S4).

Let

$$C'_1 = C_1 \begin{bmatrix} I_{n_2-n'_2-t-1} & 0 \\ 0 & M \end{bmatrix}.$$

By inductive hypothesis, there exist  $M'_1 \in \mathbb{M}_{m-m_1-1}$ , and  $M'_2 \in \mathbb{M}_{n'_2}$  such that  $[M'_1 C'_1 \quad M'_1 C_2 M'_2]$  satisfies (S1) and (S4), and  $M'_1 C_2 M'_2$  satisfies (S2) and (S3)'. By setting

$$M_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & P & 0 \\ 0 & 0 & M'_1 \end{bmatrix}, \quad M_2 = P' \begin{bmatrix} I_{n_2-n'_2-t} & 0 & 0 \\ 0 & M & 0 \\ 0 & 0 & M'_2 \end{bmatrix},$$

the matrix  $M_1 A_2 M_2$  satisfies (S1)–(S4).

Next we consider the case  $A_1 \neq 0$ . Without loss of generality, we may assume that the first nonzero column in  $A_1$  is ones followed by zeros. Write

$$A_1 = \begin{bmatrix} & \mathbf{1}^T & B_1 \\ 0_{m \times t} & 0 & D_1 \end{bmatrix}$$

for some  $t < n_1$ , with  $B_1 \in T_0^{m_1 \times (n_1-t-1)}$  and  $D_1 \in T_0^{m_2 \times (n_1-t-1)}$  for some  $m_1, m_2$  with  $m_1 + m_2 = m$  and  $m_2 < m$ . Then there exists  $P' \in \mathbb{P}_{n_2}$  such that

$$A_2 P' = \begin{bmatrix} B_2 & 0_{m_1 \times n'_2} \\ D_2 & C_2 \end{bmatrix}$$

for some  $n'_2 \geq 0$ , where  $B_2 \in T_0^{m_1 \times (n_2-n'_2)}$  has no zero column. By Lemma 2.1, there exist  $P \in \mathbb{P}_{m_1}$  and  $M \in \mathbb{M}_{n_2-n'_2}$  such that  $P B_2 M$  satisfies (S2) and (S3)' and  $[P B_1 \quad P B_2 M]$  satisfies (S4). Let  $C_1 = [D_1 \quad D_2 M]$ . Then by inductive hypothesis, there exist  $M'_1 \in \mathbb{M}_{m_2}$  and  $M'_2 \in \mathbb{M}_{n'_2}$  such that  $[M'_1 C_1 \quad M'_1 C_2 M'_2]$  satisfies (S1) and (S4), and  $M'_1 C_2 M'_2$  satisfies (S2) and (S3)'. By setting

$$M_1 = \begin{bmatrix} P & 0 \\ 0 & M'_1 \end{bmatrix}, \quad M_2 = P' \begin{bmatrix} M & 0 \\ 0 & M'_2 \end{bmatrix},$$

the proof is complete.  $\square$

**Theorem 2.3.** *Every matrix in  $T_0^{m \times n}$  is equivalent to a matrix that is in standard form.*

*Proof.* Let  $W \in T_0^{m \times n}$ . Setting  $A_1 = \emptyset$  and  $A_2 = W$  in Lemma 2.2, we see that  $W$  is equivalent to a matrix that is in standard form.  $\square$

**Corollary 2.4.** *Every unit weighing matrix is equivalent to a unit weighing matrix that is in standard form.*

*Proof.* Setting  $T = \{z \in \mathbb{C} : |z| = 1\}$ , the proof is immediate from Theorem 2.3.  $\square$

#### ACKNOWLEDGEMENT

The second author would like to acknowledge the Hitachi Global Foundation for providing a scholarship grant.

## REFERENCES

- [1] D. Best, H. Kharaghani, H. Ramp, On unit matrices with small weight, *Disc. Math.* 313 (2013), 855–864.
- [2] D. Best, H. Kharaghani, H. Ramp, Mutually unbiased weighing matrices, *Des. Codes Cryptogr.* 76 (2015), 237–256.
- [3] D. Best, H. Kharaghani, *Cryptogr. Commun.* 2 (2010), 199–209.
- [4] R. Craigen, H. Kharaghani, Orthogonal designs in: *Handbook of Comb. Des.* (C.J. Colbourn and J.H. Dinitz., eds.), 2nd Ed., pp. 280–295, Chapman & Hall/CRC Press, Boca Raton, FL, 2007.
- [5] C. Koukouvinos, J. Seberry, Weighing matrices and their applications, *JSPI* 62(1997), 91–101.

RESEARCH CENTER FOR PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCES, TOHOKU UNIVERSITY

*E-mail address:* `munemasa@math.is.tohoku.ac.jp`

RESEARCH CENTER FOR PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION SCIENCES, TOHOKU UNIVERSITY

*E-mail address:* `pritta@ims.is.tohoku.ac.jp`