

SDSS-IV MaNGA: The Impact of Diffuse Ionized Gas on Emission-line Ratios, Interpretation of Diagnostic Diagrams, and Gas Metallicity Measurements

Kai Zhang,^{1*} Renbin Yan¹, Kevin Bundy², Matthew Bershadsky³, L. Matthew Haffner³, René Walterbos⁴, Roberto Maiolino^{5,6}, Christy Tremonti³, Daniel Thomas⁷, Niv Drory⁸, Amy Jones⁹, Francesco Belfiore^{5,6}, Sebastian F. Sánchez¹⁰, Aleksandar M. Diamond-Stanic³, Dmitry Bizyaev^{11,12}, Christian Nitschelm¹³, Brett Andrews¹⁴, Jon Brinkmann¹¹, Joel R. Brownstein¹⁵, Edmond Cheung², Cheng Li^{16,17}, David R. Law¹⁸, Alexandre Roman Lopes¹⁹, Daniel Oravetz¹¹, Kaike Pan¹¹, Thaisa Storchi-Bergmann^{20,21}, Audrey Simmons¹¹

(Affiliations can be found after the references)

30 January 2017

ABSTRACT

Diffuse Ionized Gas (DIG) is prevalent in star-forming galaxies. Using a sample of 365 nearly face-on star-forming galaxies observed by MaNGA, we demonstrate how DIG in star-forming galaxies impacts the measurements of emission line ratios, hence the interpretation of diagnostic diagrams and gas-phase metallicity measurements. At fixed metallicity, DIG-dominated low $\Sigma_{\text{H}\alpha}$ regions display enhanced $[\text{S II}]/\text{H}\alpha$, $[\text{N II}]/\text{H}\alpha$, $[\text{O II}]/\text{H}\beta$, and $[\text{O I}]/\text{H}\alpha$. The gradients in these line ratios are determined by metallicity gradients and $\Sigma_{\text{H}\alpha}$. In line ratio diagnostic diagrams, contamination by DIG moves H II regions towards composite or LI(N)ER-like regions. A harder ionizing spectrum is needed to explain DIG line ratios. Leaky H II region models can only shift line ratios slightly relative to H II region models, and thus fail to explain the composite/LI(N)ER line ratios displayed by DIG. Our result favors ionization by evolved stars as a major ionization source for DIG with LI(N)ER-like emission.

DIG can significantly bias the measurement of gas metallicity and metallicity gradients derived using strong-line methods. Metallicities derived using N2O2 are optimal because they exhibit the smallest bias and error. Using O3N2, R_{23} , $\text{N2}=[\text{N II}]/\text{H}\alpha$, and N2S2H α (Dopita et al. 2016) to derive metallicities introduces bias in the derived metallicity gradients as large as the gradient itself.

The strong-line method of Blanc et al. (2015; IZI hereafter) cannot be applied to DIG to get an accurate metallicity because it currently contains only H II region models which fail to describe the DIG.

Key words: galaxies: surveys – galaxies: evolution – galaxies: fundamental parameters – galaxies: ISM – galaxies: abundances – galaxies: active

1 INTRODUCTION

Diffuse ionized gas (DIG hereafter) is an important gas component in star-forming galaxies. It was first identified in our Milky Way (MW) off the disk, and known as the Reynolds layer (Reynolds 1984). DIG is a major part of ionized gas in

MW. In terms of mass, it is about 30% of the MW neutral hydrogen (Reynolds 1990, 1991). DIG is also found in external galaxies both in extra-planar halos (e.g., Dettmar 1990; Rand et al. 1990; Rand 1996; Hoopes et al. 1999; Rossa & Dettmar 2003a,b) and in the disk (e.g., Monnet 1971; Zurita et al. 2000; Oey et al. 2007). The contribution of DIG to the total emission line flux for face-on galaxies is substantial (e.g., Walterbos & Braun 1994; Ferguson et al. 1996;

* Contact e-mail: zkdtckk@gmail.com

Hoopes, Walterbos, & Greenawalt 1996; Greenawalt et al. 1998). For 109 star-forming galaxies in the SINGG sample, the DIG fraction in $H\alpha$ flux is 0.59 ± 0.19 , and this ratio depends on the $H\alpha$ surface brightness of the whole galaxy (Oey et al. 2007). DIG is important in understanding the ionized gas in star-forming galaxies.

The differences between DIG and H II regions are not only in emission intensity, but also in emission line ratios, indicating different physical conditions. The $[S II] \lambda\lambda 6717, 6731/H\alpha$, $[N II] \lambda 6548/H\alpha$, ($[S II]/H\alpha$ and $[N II]/H\alpha$ hereafter, e.g. Reynolds 1985a; Hoopes & Walterbos 2003; Madsen et al. 2006), $[O I] \lambda 6300/H\alpha$ and $[O II] \lambda 3727/H\beta$ ($[O I]/H\alpha$ and $[O II]/H\beta$ hereafter, e.g. Voges & Walterbos 2006; Haffner et al. 1999) are found to be enhanced in DIG relative to H II regions. More quantitatively, $[S II]/H\alpha$, $[N II]/H\alpha$, $[O II]/H\alpha$, $[O I]/H\alpha$ ratios correlate negatively with $H\alpha$ flux in the MW and other galaxies (Reynolds et al. 1998; Haffner et al. 1999; Tüllmann et al. 2000; Hausen et al. 2002; Voges & Walterbos 2006; Madsen et al. 2006; Blanc et al. 2009). In high spatial resolution observations (pc to tens of pc, such as those by The Wisconsin H-Alpha Mapper (WHAM)) the line ratio vs $\Sigma_{H\alpha}$ relation is a gradual transition from H II regions to DIG. In low spatial resolution observations (hundreds of pc or kpc, such as integral-field studies of nearby galaxies) this relation is due to the mixing of H II regions and DIG within resolution elements. Different fractions of DIG and H II region contribution, combined with the different line ratios for these two types of regions would naturally produce the trend we see. Observed relations between line ratios and $\Sigma_{H\alpha}$ provide an empirical method to separate DIG dominated and H II region dominated regions (Blanc et al. 2009; Kreckel et al. 2016; Kaplan et al. 2016).

DIG line ratios cannot be explained by models of H II regions. On the classical BPT diagram (Baldwin et al. 1981; Veilleux & Osterbrock 1987; Kewley et al. 2001, 2006; Kauffmann et al. 2003), the location of an H II region is mainly determined by its metallicity and ionization parameter (e.g., Kewley et al. 2002; Dopita et al. 2013). DIG shows a lower ionization parameter than H II regions, which explains partly if not mostly the enhancement of $[N II]/H\alpha$, $[S II]/H\alpha$, and $[O II]/H\alpha$ in DIG. However, just varying these two parameters cannot fully produce the line ratios seen in DIG (e.g., Galarza et al. 1999; Hoopes & Walterbos 2003; Kaplan et al. 2016). A third variable is needed. Also, the line ratios of DIG *within* a galaxy often vary much more than those of H II regions. For example, in our Milky Way, DIG $[S II]/H\alpha$ ratios display a dispersion of 0.13 (in linear space), compared with a dispersion of 0.03 for H II regions (Madsen et al. 2006). Temperature has been proposed as the strongest factor explaining the variations of DIG line ratios because it can explain the coherent variation of $[O II] \lambda 3727/H\alpha$, $[N II] \lambda 6548/H\alpha$, and $[S II]/H\alpha$ (Haffner et al. 1999; Mierkiewicz et al. 2006). The variation of temperature is a result of the balance between heating and cooling, which also requires a physical explanation. Besides photoionization, an additional source of heating might be important especially when the density is low (e.g., Reynolds & Cox 1992). We will show in this paper that a harder ionizing spectrum could easily explain the DIG line ratios we observe in star-forming galaxies. The harder spectrum is capable of

producing partially-ionized regions that emit strong $[S II]$, $[N II]$, $[O II]$ and $[O I]$, and an increase in temperature.

The DIG can impact our interpretation of the line-ratio diagnostic diagrams of galaxies for either integrated or spatially-resolved spectroscopy. The BPT diagrams, for example, are widely used to diagnose the physical properties of ionized gas and separate different types of galaxies (Baldwin et al. 1981; Veilleux & Osterbrock 1987; Kewley et al. 2001, 2006; Kauffmann et al. 2003). When metallicity, ionization parameter, density, or ionizing spectrum change, the location of ionized gas on the BPT diagram changes (e.g., Dopita et al. 2000, 2013; Kewley et al. 2002, 2013a,b). When the N/O ratio is high, it is possible for a star-forming galaxy to be classified as composite galaxy (Pérez-Montero & Contini 2009; Pérez-Montero et al. 2013, 2016). The enhanced $[N II]/H\alpha$ and $[S II]/H\alpha$ of DIG would move the position of a star-forming galaxy toward the composite or LI(N)ER region (e.g., Sarzi et al. 2006; Stasińska et al. 2008; Yan & Blanton 2012; Kehrig et al. 2012; Singh et al. 2013; Gomes et al. 2016; Belfiore et al. 2016a,b) on the BPT diagrams. Consequently, understanding DIG helps us understand the nature of galaxies classified as composites or LI(N)ERs by the BPT diagrams.

The study of DIG is also critical for gas-phase metallicity measurements in star-forming galaxies. Metallicity calibrations are generally based on H II region models which have certain assumptions. Some of these assumptions are not valid for DIG, and hence lead to biased metallicity measurements when DIG is present. The *metallicity* and *ionization parameter* ($q = \frac{\text{ionizing photon flux}}{N_e} = U \times c$) determine the line ratios of a H II region, and these two parameters are correlated (Dopita et al. 2006). Metallicity also determines the shape of the ionizing spectrum. In DIG, however, the correlation between q and metallicity no longer holds, since DIG has a much lower ionization parameter than H II regions, and the ionizing spectrum shape can change. Given the very different line ratios such as $[N II]/H\alpha$, $[O III] \lambda 5007/H\beta$ ($[O III]/H\beta$ hereafter) and $[S II]/H\alpha$ for DIG and H II regions, biases in metallicities derived from strong line methods are inevitable. These biases potentially contribute to the large dispersion in metallicity measurements found in the literature. This will influence metallicity gradient measurements (Sánchez et al. 2014; Ho et al. 2015), metallicities at the outskirts of galaxies (Moran et al. 2012), the mass-metallicity relation in the local universe (e.g. McClure & van den Bergh, 1968; Lequeux et al. 1979; Garnett 2002; Tremonti et al. 2004; Lee et al. 2006) and at high redshift (Erb et al. 2006; Maiolino et al. 2008; Mannucci et al. 2009), and the mass-metallicity-SFR fundamental plane (Mannucci et al. 2010; Yates et al. 2012; Andrews & Martini 2013). For single-fiber surveys such as SDSS, the observed emission lines come from a combination of DIG and H II regions, and the bias introduced by DIG is uncertain and hard to quantify. With the help of integral field spectroscopy (IFS), we can study how the presence of DIG impacts metallicity measurements in detail.

In this paper, we demonstrate the prevalence of DIG in star-forming galaxies from the MaNGA survey. The large sample and full optical wavelength coverage enable us to explore the optical line ratios for DIG for an unprecedented number of star-forming galaxies. Section 2 describes our sample and the emission line measurements; while Sec-

tion 3.1 shows the $[\text{S II}]/\text{H}\alpha$ vs $\Sigma_{\text{H}\alpha}$ relation that illustrates the dominance of DIG in low $\Sigma_{\text{H}\alpha}$ regions. Section 3 demonstrates how DIG impact line ratios like $[\text{S II}]/\text{H}\alpha$, $[\text{N II}]/\text{H}\alpha$, $[\text{O II}]/\text{H}\beta$, $[\text{O I}]/\text{H}\alpha$, $[\text{O III}]/[\text{O II}]$, $[\text{O III}]/\text{H}\beta$. Section 4 tests the leaky H II region model as well as one in which hot evolved stars serve as the ionization source of the DIG, using different line ratios and diagnostic diagrams. Section 5 presents our study of how the DIG impacts metallicity derived using strong line methods: N2, N2O2, R_{23} , O3N2 and N2S2H α , and IZI. Section 6 contains a discussion of these results, summarized then in Section 7. We use a cosmology with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3$, and $\Omega_\Lambda = 0.7$ throughout this paper.

2 SAMPLE AND MEASUREMENTS

2.1 MaNGA Overview

MaNGA (Mapping Nearby Galaxies at APO) (Bundy et al. 2015) is one of the three core programs in the Sloan Digital Sky Survey-IV (SDSS-IV). It aims at obtaining Integrated Field Spectroscopy (IFS) of 10,000 nearby galaxies. The survey employs the BOSS spectrographs (Smee et al. 2013) on the 2.5m Sloan Foundation Telescope (Gunn et al. 2006). The spectrographs provide a spectral coverage from 3600Å to 10300Å at a resolution around $R \sim 2000$. The fiber feed system has been re-designed (Drory et al. 2015) to accommodate 1423 fibers that are bundled into five 127-fiber bundles, two 91-fiber bundles, four 61-fiber bundles, four 37-fiber bundles, two 19-fiber bundles, twelve 7-fiber mini-bundles, and 92 sky fibers. All bundles with 19 or more fibers are used for galaxy targets, while the mini-bundles are used to observe standard stars providing a flux calibration to an accuracy better than 5% (Yan et al. 2016a). Each individual fiber is 2" in diameter. The filling factor of fibers in the bundles is 56%. The observations are done with a dithering scheme to achieve a complete spatial coverage and near critical sampling of the PSF (Law et al. 2015).

The MaNGA sample is designed to have a roughly flat i-band absolute magnitude distribution (Wake et al. in prep). The reconstructed datacubes have 2.5" PSF and 0.5" spaxel (Law et al. 2015, 2016; Yan et al. 2016a,b). The sample has a redshift range of $0.01 < z < 0.15$, meaning the targets cover a factor of 15 in distance, and consequently the physical resolution also span a range of 15. We select more luminous galaxies, which are larger, at higher redshift. The typical physical resolution is 1-2 kpc. The physical resolution is highly correlated with luminosity which means that we need to be aware that any results we find which might depend on physical resolution will be specific to a specific range of luminosity and vice versa. The primary sample, which covers to 1.5 effective radius (R_e) in major axis, comprise 2/3 of the sample while the secondary sample, which covers to 2.5 R_e comprise the remaining 1/3. Massive galaxies, which are bigger, are selected at higher redshift. All plates are exposed until we reach a $(S/N)^2$ of 20 per pixel per fiber in the g-band continuum for a galactic-extinction-corrected g-band fiber magnitude of 22, and a $(S/N)^2$ of 36 per pixel per fiber in the i-band continuum for a galactic-extinction-corrected i-band fiber magnitude of 21 (Yan et al. 2016b). MaNGA provides a benchmark of resolved ionized gas properties, stellar

population, and dynamical evolution in the local universe (e.g., Belfiore et al. 2015; Li et al. 2015; Wilkinson et al. 2015).

2.2 Sample Selection

In this paper, we use a sample of 81 regular survey plates observed before summer of 2015. There are 1391 unique galaxies observed. This corresponds to the sample that was released in DR13. The photometry data is from NASA-Sloan Atlas catalog¹ (NSA Catalog). By applying a color cut of $M_u - M_r < 2.1$ (M_u and M_r are rest-frame absolute magnitudes without extinction correction) we select only blue galaxies. This resulted in a sample of 592 galaxies. To remove edge-on galaxies we apply the cut $b/a > 0.5$, where b and a are the minor and major axis of the Sersic model. Edge-on galaxies suffer from strong projection effects and are also prone to severe extinction. These criteria leave us with 365 galaxies. AGNs are not eliminated because we care about DIG that is far away from the center. From this sample we choose three galaxies that have large internal variations of H α surface brightness and best spatial resolution (127 fiber IFUs) to demonstrate the impact of DIG on line ratios, interpretation of diagnostic diagrams, and metallicity measurements. We discuss in Section 3.4 and Section 5.7 that the impact of DIG is prevalent in all star-forming galaxies in our sample.

2.3 Spectrum Fitting

We start with the datacubes produced by the MaNGA data reduction pipeline (Law et al. 2016). We first construct Voronoi bins of the spectra by requiring the S/N in r band in each bin to be greater than 30 per Å. The covariance between spaxels is not accounted for when binning. With BC03² (Bruzual & Charlot 2003), we produce 14 continuum templates for SSPs with $Z_\odot = 0.02$ and 0.008 and ages = 13, 7, 2, 1, 0.5, 0.25, and 0.125 Gyr. For the stacked spectrum in each bin, we fit combinations of these simple stellar populations (SSPs), fitting velocity and relative amplitudes to spectra combined from spaxels in a Voronoi bin. We then derive the stellar velocity dispersion using the *vdispfit.pro* in IDLSPEC2D³ package. 20Å windows around $[\text{O II}] \lambda 3727$, H β , $[\text{O III}] \lambda 4959$, $[\text{O III}] \lambda 5007$, $[\text{O I}] \lambda 6300$, $[\text{N II}] \lambda 6548$, H α , $[\text{N II}] \lambda 6583$, and $[\text{S II}] \lambda \lambda 6717, 6731$ are masked during the continuum fitting. We smooth the observed spectrum with a 50 pixel window and then subtract the smoothed spectrum from the original spectrum. This smoothing basically removes remaining spectrophotometric calibration mismatch between data and templates as well as effects of internal extinction. The smoothing scale is $\sim 3000 \text{ km s}^{-1}$. By doing this, we are left with only small scale spectral variations such as absorption lines and emission lines, which we refer to as the 'line feature spectrum'. The same technique is applied to the SSP templates to get the 'line feature templates' that only contains small scale information. A linear regression is performed to fit the 'line feature spectrum' with the sum of

¹ <http://www.nsatlas.org>

² <http://www2.iap.fr/users/charlot/bc2003/>

³ http://spectro.princeton.edu/idlspec2d_doc.html

'line feature templates' at a number of velocity offsets relative to the systematic redshift of the galaxy. The velocity grid ranges from -450 km s^{-1} to 450 km s^{-1} with an interval of 30 km s^{-1} . For each Voronoi bin at each velocity, the least-square fitting yields a χ^2 for that fit. A quadratic curve is fitted to the χ^2 vs velocity curve to find the velocity offset yielding the minimum χ^2 . For each spaxel in one Voronoi bin, we use the fitted stellar continuum, the combination of 14 BC03 templates, as one template, and adjust the amplitude to fit the continuum in each spaxel so that we can measure the emission-line ratios on a spaxel by spaxel basis. The residual emission-line only spectrum is stored for refined emission line fitting. We use single gaussians to fit the [O I], H β , [O III], [N II] $\lambda 6548$, H α , [N II] $\lambda 6583$, and [S II] respectively, and we use two gaussians to fit [O II]. The line ratio of [N II] $\lambda 6583$ /[N II] $\lambda 6548$ is fixed to 3. Velocity dispersions and central velocity shifts of all emission lines are tied to be the same. The H α surface brightness maps for the 3 representative galaxies are shown in Figure 1. The errors of the line strength are obtained by the MPFIT package which only includes the fitting errors (Markwardt 2009). A S/N cut of 5 is applied to emission lines used.

3 VARIATION OF LINE RATIOS WITH H α SURFACE BRIGHTNESS

3.1 Separation of regions dominated by DIG and H II regions

The first step to study DIG is separating DIG from H II regions. The best way to isolate DIG is to identify individual H II regions and subtract them out (e.g., Walterbos & Braun 1994; Zurita et al. 2000, 2002; Thilker, Walterbos, & Braun 2002; Sánchez et al. 2012). Due to the limited spatial resolution of MaNGA, we can not resolve individual H II regions in one galaxy. Our reconstructed PSF is about $2.5''$ FWHM ($2''$ covers 1 kpc at $z=0.025$) while the size of a typical H II region is a few to hundreds of pc (Kennicutt 1984; Garay & Lizano 1999; Kim & Koo 2001; Hunt & Hirashita 2009). So the light in one spaxel is always a mixture of H II region emission and the surrounding DIG. Instead we separate DIG dominated regions and H II regions dominated regions. In Figure 2, we see that the low surface brightness regions have $[S II]/H\alpha \sim 0.5 - 1.0$ while the high surface brightness regions show $[S II]/H\alpha \sim 0.2 - 0.4$, depending on the metallicity. Since $[S II]/H\alpha$ is sensitive to metallicity, and there are metallicity gradients in galaxies, we separate all the spaxels in each galaxy into different radial bins: $[0,0.6]$, $[0.6,1.2]$, $[1.2,1.8]$, and $[1.8,2.4]R_e$. The $[S II]/H\alpha$ vs $\Sigma_{H\alpha}$ relations are similar in all radial bins, meaning the variation of $[S II]/H\alpha$ is not caused by metallicity variation, but reflects the transition from DIG dominated low $\Sigma_{H\alpha}$ regions to H II region dominated high $\Sigma_{H\alpha}$ regions. This figure illustrates that $\Sigma_{H\alpha}$ can be used to separate the two different kinds of regions: low $\Sigma_{H\alpha}$ DIG dominated regions and high $\Sigma_{H\alpha}$ H II region dominated regions.

3.2 $[S II]/H\alpha$, $[N II]/H\alpha$, $[O II]/H\beta$, and $[O I]/H\alpha$

Galaxies show metallicity variations, in particular in the form of radial gradients (e.g. Sánchez et al. 2014; Ho et al.

2015). It is crucial to control metallicity before exploring how the line ratios vary with other parameters like H α surface brightness because metallicity is a major source of variation in line ratios. We plot line ratios as a function of radius, and color code the dots with H α surface brightness. We assume that the metallicity is constant within annuli for each galaxy but may be changing with radius.

We explore the impact of DIG on $[S II]/H\alpha$ first. We color-code the dots by H α surface brightness and show how $[S II]/H\alpha$ changes with radius in Figure 3. At a fixed radius, we see a rainbow pattern such that low $\Sigma_{H\alpha}$ regions have higher $[S II]/H\alpha$. This is a direct demonstration that $[S II]/H\alpha$ is enhanced in DIG after controlling for metallicity.

We then explore how DIG impacts $[N II]/H\alpha$. $[N II]/H\alpha$ is used as a metallicity indicator since Nitrogen is a secondary element and proportional to Z^2 while H α is not sensitive to metallicity (e.g. Storch-Bergmann et al., 1994, van Zee et al., 1998, Denicoló et al., 2002). We color-code the dots by H α surface brightness and show how $[N II]/H\alpha$ changes with radius in Figure 4. If we control for H α surface brightness by looking at the dots with the same color, $[N II]/H\alpha$ decreases towards large radius, reflecting a metallicity gradient. Due to the presence of an H α surface brightness gradient, the different surface brightness bins usually trace different parts of the galaxy. However, at the radius where they overlap, we see a rainbow pattern. At a fixed radius, DIG dominated low surface brightness region show a higher $[N II]/H\alpha$. If we use N2 to derive the metallicity, the enhancement means that the metallicity would be overestimated in those spaxels with a high DIG fraction and vice-versa. The impact of DIG on the metallicity derived using $[N II]/H\alpha$ is given in Section 5.1.

We show how $[O II]/H\beta$ changes with radius and H α surface brightness in Figure 5. The extinction correction is not applied because it is not reliable when the emission line, especially H β , is weak. Besides, a foreground dust screen may not be the appropriate model for DIG. Finally, We show in Figure 6 that $H\beta/H\alpha$ does not depend on H α surface brightness, meaning extinction will not produce any line ratios change between DIG and H II regions.

$[O II]/H\beta$ generally increases with radius at fixed H α surface brightness due to a metallicity gradient. At fixed radius, the low H α surface brightness regions have higher $[O II]/H\beta$. It is interesting that $[O II]/H\beta$ and $[N II]/H\alpha$ both increase with decreasing surface brightness at fixed radius while they change reversely with radius at fixed H α surface brightness. The opposite variation trends of $[O II]/H\beta$ and $[N II]/H\alpha$ with radius at fixed surface brightness are consistent with a metallicity variation. With $[O/H]$ above solar value, $[N II]/H\alpha$ increases with metallicity due to the addition of secondary Nitrogen while $[O II]/H\beta$ drops with increasing metallicity because of decreasing temperature. The positive correlation between $[O II]/H\beta$ and $[N II]/H\alpha$ with surface brightness at fixed radius is seemingly consistent with temperature variation. Mierkiewicz et al. (2006) show that $[O II]/H\alpha$ and $[N II]/H\alpha$ correlate positively with temperature variation and they concluded that the variation of line ratios is driven by temperature variations (Haffner et al. 1999; Haffner et al. 2009). However, the variation of temperature is a result of the balance between heating and cooling which, itself, needs a physical explanation. Besides, some




Figure 1. $H\alpha$ surface brightness plot and SDSS optical image for each galaxy. The cyan ellipses in the upper represent are 1.5 and 2.5 times the effective radius. The hexagons in the lower row demonstrate the location of the IFU bundles on the object.




Figure 2. $[S\text{II}]/H\alpha$ vs $H\alpha$ surface brightness plot for each galaxy. Different color represent different annuli. The solid line is the $[S\text{II}]/H\alpha$ vs $H\alpha$ surface brightness relation for Milky Way (Madsen et al. 2006). We see that high and low $H\alpha$ surface brightness regions have different $[S\text{II}]/H\alpha$. The red and blue bars show the typical error at $\log \Sigma_{H\alpha}=38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log \Sigma_{H\alpha}=38 \text{ erg s}^{-1} \text{ kpc}^{-2}$ for individual spaxels. The length of the error bars is 2σ .

significant assumptions need to be made to use $[O\text{II}]/H\beta$ and $[N\text{II}]/H\alpha$ as temperature tracers. The biggest assumption is that $[O\text{II}]$ and $H\beta$ are co-spatial. However, we only resolve galaxies to kpc scale for MaNGA survey, thus we see the integrated emission from all layers of ISM around an ionizing source. The co-spatial assumption may not hold for DIG. $[N\text{II}]/H\alpha$ line ratios are often used as a metallicity indicator for extra-galactic studies, while it may be used as a temperature indicator for galactic uses. Both metallicity and temperature are relevant to $[N\text{II}]/H\alpha$ emission. Finally, a decrease in ionization parameter and/or a harder ionizing spectrum towards lower $\Sigma_{H\alpha}$ will also translate into enhancement of $[O\text{II}]/H\beta$ and $[N\text{II}]/H\alpha$. This is further explored in Section 4.

$[O\text{I}]$ is an important line to diagnose the physical properties of ionized gas (Dopita et al. 2000; Kewley et al. 2006). $[O\text{I}]$ is detected in DIG of M33 and $[O\text{I}]/H\alpha$ is found to correlate negatively with emission measure (Voges & Walterbos 2006). $[O\text{I}]/H\alpha$ is a strong function of temperature and the hardness of the ionizing spectrum. $[O\text{I}]$ has a high critical density ($n_{[O\text{I}]} = 10^{6.3} \text{ cm}^{-3}$), thus it is emitted in high density neutral and partially ionized regions. $[O\text{I}]/H\alpha$ is also an excellent tracer of shocks (Dopita et al. 2000; Kewley et al. 2002; Allen et al. 2008). In Figure 7, we show how the $[O\text{I}]/H\alpha$ changes with $H\alpha$ surface brightness and radius for our three galaxies. For all galaxies, $[O\text{I}]/H\alpha$ increases with radius. At a fixed radius, $[O\text{I}]/H\alpha$ is enhanced for low surface brightness regions, indicating $[O\text{I}]/H\alpha$ is higher in DIG.

We note the contrast between DIG and H II region is large, $[\text{O I}]/\text{H}\alpha$ drops ~ 1 dex with an increase of 1 dex in $\Sigma_{\text{H}\alpha}$.

3.3 $[\text{O III}]/[\text{O II}]$ and $[\text{O III}]/\text{H}\beta$

$[\text{O III}]/[\text{O II}]$ is a good ionization parameter proxy when metallicity is controlled (Kewley et al. 2002, see also Figure 18). We show how the $[\text{O III}]/[\text{O II}]$ values of low surface brightness spaxels differ from the high surface brightness spaxels for our sample galaxies in Figure 8. This relation has large dispersion but there is a clear trend. At a fixed $\text{H}\alpha$ surface brightness, $[\text{O III}]/[\text{O II}]$ increases as we go to larger radii due to a metallicity gradient (see also Figure 14). At a fixed radius, $[\text{O III}]/[\text{O II}]$ mostly decreases with decreasing surface brightness. The result suggests that DIG has a lower ionization parameter than H II regions.

$[\text{O III}]/\text{H}\beta$ is another frequently used ionization parameter proxy. Its dependence on metallicity and ionizing spectrum hardness is much stronger than $[\text{O III}]/[\text{O II}]$. How $[\text{O III}]/\text{H}\beta$ of DIG differs from that of H II regions depends on the specific physical properties of the ISM. DIG can show either higher (Wang, Heckman, & Lehnert 1997; Rand 1998, 2000; Collins & Rand 2001; Otte 2001, 2002; Otte, Gallagher, & Reynolds 2002) or lower (MW: Reynolds 1985b; M31: Greenawalt et al. 1997; Galarza, Walterbos, & Braun 1999) $[\text{O III}]/\text{H}\beta$ than H II regions. The study of our whole sample of 395 galaxies in Section 3.4 confirms the diverse behavior of $[\text{O III}]/\text{H}\beta$ in DIG. In the previous paragraph we have shown that DIG has lower ionization parameter than H II regions. At a fixed metallicity, a decrease in ionization parameter will result in a decrease of $[\text{O III}]/\text{H}\beta$ for $q < 7.5$ if no other parameter in the photoionization model is changed (see Figure 15). For $q > 7.5$, $[\text{O III}]/\text{H}\beta$ is roughly constant and independent of ionization parameter. This diverse behavior of $[\text{O III}]/\text{H}\beta$ means a third parameter other than metallicity and ionization parameter is needed, as will be discussed in Section 4. The 3rd galaxy: 8243-12704 shows a negative $[\text{O III}]/[\text{O II}]$ and $[\text{O III}]/\text{H}\beta$ gradient. This is not consistent with an inverse metallicity gradient at the center because the $[\text{N II}]/[\text{O II}]$ gradient is negative in this range. This could be due to AGN activity, but the strength of the AGN is not strong enough to produce AGN-like line ratios on the BPT diagram. These kind of sources will be explored in future work.

3.4 Line ratios vs $\Sigma_{\text{H}\alpha}$ relation

3.4.1 Line ratios vs $\Sigma_{\text{H}\alpha}$ relation: The Whole Sample

To quantify the variation of the line ratios as a function of $\text{H}\alpha$ surface brightness for the whole sample, we select all spaxels that have $[0.4R_e, 0.6R_e]$ in each galaxy. We then normalize the log line ratios vs log $\Sigma_{\text{H}\alpha}$ relation of each galaxy by subtracting from all the spaxels the median log line ratio and the median log $\Sigma_{\text{H}\alpha}$ of the set of spaxels. All the spaxels in a galaxy are weighted by $\frac{1}{N}$, where N is the number of valid spaxels in this galaxy. Figure 10 shows $\Delta \log$ line ratios vs $\Delta \log \Sigma_{\text{H}\alpha}$ relation for the whole sample by combining all galaxies. $[\text{S II}]/\text{H}\alpha$, $[\text{N II}]/\text{H}\alpha$, $[\text{O II}]/\text{H}\beta$, $[\text{O I}]/\text{H}\alpha$ are higher in DIG dominated low $\Sigma_{\text{H}\alpha}$ regions. $[\text{O III}]/[\text{O II}]$ is slightly lower in DIG while $[\text{O III}]/\text{H}\beta$ is mildly higher in DIG.

Furthermore, we perform linear regression to the line

ratio vs $\text{H}\alpha$ surface brightness relation in narrow annuli for individual galaxies. For each galaxy, we get the slope of the linear regression at three radial bins: $[0.4R_e, 0.6R_e]$, $[0.8R_e, 1.0R_e]$, and $[1.3R_e, 1.5R_e]$. The distribution of slopes for different line ratios at three radii are shown in black, blue, and red in Figure 11. A negative slope means the line ratio is enhanced in DIG while a positive slope means the line ratio is lower in DIG. We see that the slope peaks around ~ -0.3 for $\log [\text{S II}]/\text{H}\alpha$, $\log [\text{N II}]/\text{H}\alpha$ and $\log [\text{O II}]/\text{H}\beta$. The slopes for these three ratios are rarely positive. The slope peaks at -1.0 for $\log [\text{O I}]/\text{H}\alpha$, which is the most significant among the line ratios explored here. The slopes for $[\text{O III}]/[\text{O II}]$ is mostly positive, and its dispersion is larger than the dispersion for $[\text{S II}]/\text{H}\alpha$, $[\text{N II}]/\text{H}\alpha$, and $[\text{O II}]/\text{H}\beta$. The larger dispersion tells us that the ionization parameter varies significantly. The slope distribution for $\Delta \log [\text{O III}]/\text{H}\beta$ vs $\Delta \log \Sigma_{\text{H}\alpha}$ peaks at ~ 0 , but the distribution is skewed to negative values. There are more galaxies with higher $[\text{O III}]/\text{H}\beta$ in DIG than galaxies with lower $[\text{O III}]/\text{H}\beta$ in DIG. The dispersion is even larger than that for $[\text{O III}]/[\text{O II}]$. This illustrates that $[\text{O III}]/\text{H}\beta$ is very diverse in DIG. As shown in Section 3.3, DIG shows higher $[\text{O III}]/\text{H}\beta$ than H II regions in some galaxies, while it shows lower or similar $[\text{O III}]/\text{H}\beta$ for other galaxies. Figure 11 demonstrates that DIG with higher $[\text{S II}]/\text{H}\alpha$, $[\text{N II}]/\text{H}\alpha$, $[\text{O II}]/\text{H}\beta$, $[\text{O I}]/\text{H}\alpha$ and lower $[\text{O III}]/[\text{O II}]$ than H II regions is prevalent among all galaxies in our sample. $[\text{O III}]/\text{H}\beta$ could be significantly higher or lower than H II regions. The contamination of DIG influences every star-forming galaxy.

3.4.2 Line ratios vs $\Sigma_{\text{H}\alpha}$ relation: Split by Stellar Mass

In Figure 12 and Figure 13, we show the $\Delta \log$ line ratio vs $\Delta \log \Sigma_{\text{H}\alpha}$ relations for galaxies with stellar mass less than $10^{9.43}$ and higher than $10^{10.08}$ respectively. These are the one third least massive and the one third most massive galaxies in our sample. The slopes of $\Delta \log [\text{S II}]/\text{H}\alpha$, $[\text{N II}]/\text{H}\alpha$, $[\text{O II}]/\text{H}\beta$, $[\text{O I}]/\text{H}\alpha$, and $[\text{O III}]/[\text{O II}]$ vs $\Delta \log \Sigma_{\text{H}\alpha}$ relations do not depend much on stellar mass. However, massive galaxies show a significantly negative $\Delta \log [\text{O III}]/\text{H}\beta$ vs $\Delta \log \Sigma_{\text{H}\alpha}$ relation while less massive galaxies show a positive $\Delta \log [\text{O III}]/\text{H}\beta$ vs $\Delta \log \Sigma_{\text{H}\alpha}$ relation. In other words, $[\text{O III}]/\text{H}\beta$ of DIG is always enhanced relative to H II regions in the most massive galaxies. A leaky H II region model can not produce high $[\text{O III}]/\text{H}\beta$ relative to H II regions (Section 4.2). The dependence of the $\Delta \log [\text{O III}]/\text{H}\beta$ vs $\Delta \log \Sigma_{\text{H}\alpha}$ relation on stellar mass may indicate different ionization mechanisms in galaxies of different masses. We leave the study of the physical reason behind the dependence of $\Delta \log$ line ratios vs $\Delta \log \Sigma_{\text{H}\alpha}$ relation on stellar mass for future studies.

4 DIAGNOSTIC DIAGRAMS

4.1 Comparison of line ratios with H II region grids

In Figure 14, we plot the line ratios for each spaxel on the $[\text{O III}]/[\text{O II}]$ vs $[\text{N II}]/[\text{O II}]$ diagram (Dopita et al. 2000). $[\text{N II}]/[\text{O II}]$ is sensitive to metallicity because N is a secondary element. At high metallicity, $[\text{O II}]$ is suppressed due




Figure 3. $[\text{S II}]/\text{H}\alpha$ as a function of radius. The dispersion at fixed radius is about 0.2 dex. But when the dots are color-coded by $\text{H}\alpha$ surface brightness as shown in the colorbar, we see a beautiful rainbow pattern. The dispersion is significantly reduced at fixed $\Sigma_{\text{H}\alpha}$. This is because DIG that dominates the low $\Sigma_{\text{H}\alpha}$ region has higher $[\text{S II}]/\text{H}\alpha$. The red and blue bars show the typical line ratio error at $\log \Sigma_{\text{H}\alpha} = 39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log \Sigma_{\text{H}\alpha} = 38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ for individual spaxels. The length of the error bars is 2σ . We show in Figure 11 that the impact of DIG is prevalent in all star-forming galaxies in our sample, not only in the three galaxies we show.




Figure 4. $[\text{N II}]/\text{H}\alpha$ as a function of radius. We see $[\text{N II}]/\text{H}\alpha$ generally drops toward large radius due to a metallicity gradient. The dispersion at fixed radius is about 0.2 dex. But when the dots are color-coded by $\text{H}\alpha$ surface brightness as shown in the colorbar, we see a beautiful rainbow pattern. This shows $[\text{N II}]/\text{H}\alpha$ is enhanced in DIG. The red and blue bars show the typical line ratio error at $\log \Sigma_{\text{H}\alpha} = 39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log \Sigma_{\text{H}\alpha} = 38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ for individual spaxels. The length of the error bars is 2σ .




Figure 5. $[\text{O II}]/\text{H}\beta$ as a function of radius. The dots are color-coded by $\Sigma_{\text{H}\alpha}$. We see rainbow patterns as in Figure 4, indicating $[\text{O II}]/\text{H}\beta$ is enhanced in DIG. The red and blue bars show the typical line ratio error at $\log \Sigma_{\text{H}\alpha} = 39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log \Sigma_{\text{H}\alpha} = 38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ for individual spaxels. The length of the error bars is 2σ .




Figure 6. $H\beta/H\alpha$ as a function of radius. The dots are color-coded by $\Sigma_{H\alpha}$. We don't see any dependence of $H\beta/H\alpha$ on $\Sigma_{H\alpha}$ at a given radius, implying the extinction correction is not the reason for the different line ratios of DIG and H II regions. The red and blue bars show the typical line ratio error at $\log\Sigma_{H\alpha}=39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log\Sigma_{H\alpha}=38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ for individual spaxel. The length of the error bars is 2σ .




Figure 7. $[O\text{I}]/H\alpha$ as a function of radius. The dots are color-coded by $\Sigma_{H\alpha}$. We see rainbow patterns as in Figure 4, indicating $[O\text{I}]/H\alpha$ is enhanced in DIG. The red and blue bars show the typical line ratio error at $\log\Sigma_{H\alpha}=39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log\Sigma_{H\alpha}=38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ for individual spaxels. The length of the error bars is 2σ .




Figure 8. $[O\text{III}]/[O\text{II}]$ as a function of radius. The dots are color-coded by $\Sigma_{H\alpha}$. At a given radius, DIG shows lower $[O\text{III}]/[O\text{II}]$, meaning lower ionization parameter. The red and blue bars show the typical line ratio error at $\log\Sigma_{H\alpha}=39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log\Sigma_{H\alpha}=38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ for individual spaxels. The length of the error bars is 2σ .




Figure 9. $[\text{O III}]/\text{H}\beta$ as a function of radius. The dots are color-coded by $\Sigma_{\text{H}\alpha}$. At a given radius, DIG might show higher or lower $[\text{O III}]/\text{H}\beta$ than H II regions depending on the specific situation of a galaxy. The red and blue bars show the typical line ratio error at $\log \Sigma_{\text{H}\alpha} = 39$ $\text{erg s}^{-1} \text{kpc}^{-2}$ and $\log \Sigma_{\text{H}\alpha} = 38.5$ $\text{erg s}^{-1} \text{kpc}^{-2}$ for individual spaxels. The length of the error bars is 2σ .




Figure 10. $\Delta \log$ (line ratios) vs $\Delta \log \Sigma_{\text{H}\alpha}$ relation at $[0.4R_e, 0.6R_e]$ for all galaxies. We normalize the log line ratios vs $\log \Sigma_{\text{H}\alpha}$ relation by subtracting the median log line ratio and median $\log \Sigma_{\text{H}\alpha}$. All the spaxels in a galaxy are weighted by $\frac{1}{N}$, where N is the number of valid spaxels in this galaxy. Contour levels are 5, 40, 70, 85, and 92 percentile. $[\text{S II}]/\text{H}\alpha$, $[\text{N II}]/\text{H}\alpha$, $[\text{O II}]/\text{H}\beta$, $[\text{O I}]/\text{H}\alpha$ are higher in DIG dominated low $\Sigma_{\text{H}\alpha}$ regions. $[\text{O III}]/[\text{O II}]$ is slightly lower in DIG while $[\text{O III}]/\text{H}\beta$ is mildly higher in DIG.

to the decrease in temperature. $[\text{O III}]/[\text{O II}]$ is a very good proxy of the ionization parameter. We over-plot the most up-to-date grids for H II regions from Dopita et al. (2013) with $\kappa = Inf$, which means the electrons obey the Maxwell-Boltzmann distribution. Our data fall within the grids. However, when we turn to the BPT diagrams in Figure 15 and

16, we see that while the high $\text{H}\alpha$ surface brightness spaxels are still consistent with the grid prediction, DIG dominated low $\text{H}\alpha$ surface brightness regions are located outside the model grids. Specifically, the $[\text{N II}]/\text{H}\alpha$ and $[\text{S II}]/\text{H}\alpha$ are significantly enhanced in DIG. Previous work found a similar phenomenon for DIG (e.g., Galarza et al. 1999; Hoopes &




Figure 11. The distribution of $\Delta \log$ line ratios vs $\Delta \log \Sigma_{H\alpha}$ slope at three radii. The black line is for the radius at $[0.4R_e, 0.6R_e]$, the blue line is for the radius at $[0.8R_e, 1.0R_e]$, and the red line is for the radius at $[1.3R_e, 1.5R_e]$. The green dashed line is the slope=0 line for reference. For almost all star-forming galaxies in our sample, DIG shows higher $[S II]/H\alpha$ $[N II]/H\alpha$ $[O II]/H\beta$, $[O I]/H\alpha$ and lower $[O III]/[O II]$ than H II regions. $[O III]/H\beta$ in DIG can be higher or lower than in H II regions.

Walterbos 2003; Kaplan et al. 2016). This means that H II region models with only a lower q can not explain the low ionization line ratios like $[S II]/H\alpha$ and $[N II]/H\alpha$. Apart from metallicity and ionization parameter, there must be other parameter(s) that govern the behavior of the emission line.

The location of DIG on the BPT diagrams gives us some clue to what the other parameters could be. DIG enters the composite or LI(N)ER regions on the $[O III]/H\beta$ vs $[N II]/H\alpha$ and $[O III]/H\beta$ vs $[S II]/H\alpha$ diagrams. AGN and LI(N)ER are characterized by their hard ionizing spectrum. A hard spectrum is capable of producing a large partially-ionized region which enhances the $[S II]/H\alpha$ and $[N II]/H\alpha$, just like what we observe. Thus, a harder ionizing spectrum is one possible answer to the line ratios enhancement.

4.2 Photoionization Grid

To get a quantitative idea of how hardness can change line ratios, we use CLOUDY (Ferland et al. 1998) to calculate new grids for a series of incident spectra. We test two models that are capable of producing a hard ionizing spectrum: a leaky H II region model and a hot evolved star model. The leaky H II region model can have a harder ionizing spectrum because the lower energy part is more likely to be absorbed by the IGM (e.g. Giammanco et al. 2005). Hot evolved stars, like pAGB stars, are characterized by very high tempera-

tures, yielding a hard ionizing spectrum. The input spectra to test the two models include (see Figure 17):

- (1) The spectrum of an O star with $T_{eff} = 42,300K$, $\log g = 4.22$, solar metallicity. (Tlusty OSTAR2002, Lanz & Hubeny 2003; magenta line)
- (2) Spectrum in (1) filtered through a gas cloud with $n_e = 100 \text{ cm}^{-3}$ and $N_H = 10^{18} \text{ cm}^{-2}$. The resulting spectrum includes both the transmitted spectrum and the emission from the cloud itself. (orange line)
- (3) Spectrum in (1) filtered through a gas cloud with $n_e = 100 \text{ cm}^{-3}$ and $N_H = 10^{18.5} \text{ cm}^{-2}$ (green line)
- (4) Spectrum in (1) filtered through a gas cloud with $n_e = 100 \text{ cm}^{-3}$ and $N_H = 10^{18.7} \text{ cm}^{-2}$ (cyan line)
- (5) Spectrum in (1) filtered through a gas cloud with $n_e = 100 \text{ cm}^{-3}$ and $N_H = 10^{18.9} \text{ cm}^{-2}$ (blue line)
- (6) The spectrum of a 13 Gyr SSP generated with BC03. (red line)

The metallicity of the ionized cloud is that of Orion Nebula (Baldwin et al. 1991; Rubin et al. 1991; Osterbrock et al. 1992; Rubin et al. 1993). The density of the cloud for model (1) is 10 cm^{-3} , and 1 cm^{-3} for model (2), (3), (4), (5) and (6). We run the calculation for $\log U = -4.5$ to -2.0 with an interval of 0.5 dex ($\log U = \log q/c$, c is the speed of light).




Figure 12. The $\Delta \log$ (line ratios) vs $\Delta \log \Sigma_{H\alpha}$ relation at $[0.4R_e, 0.6R_e]$ for the one third most massive galaxies. We normalize the log line ratios vs $\log \Sigma_{H\alpha}$ relation by subtracting the median log line ratio and median $\log \Sigma_{H\alpha}$. All the spaxels in a galaxy are weighted by $\frac{1}{N}$, where N is the number of valid spaxels in the galaxy. $[S II]/H\alpha$, $[N II]/H\alpha$, $[O II]/H\beta$, $[O I]/H\alpha$ are higher in DIG dominated low $\Sigma_{H\alpha}$ regions. $[O III]/[O II]$ is slightly lower in DIG while $[O III]/H\beta$ is significantly higher in DIG. $[O III]/H\beta$ in DIG of massive galaxies is different from that in less massive galaxies.

Model (1) represents a typical H II region. Models (2), (3), (4), and (5) are for testing the leaky H II region model. The hardening of the ionizing spectrum is obvious when compared with (1). Model (6) simulates LI(N)ER-like ionization by an old stellar population which has been proposed to explain the emission seen in a large fraction of passive galaxies (e.g., Sarzi et al 2006; Stasińska et al. 2008; Yan & Blanton 2012; Kehrig et al. 2012; Singh et al. 2013; Gomes et al. 2016; Belfiore et al. 2016a,b). Evolved stellar populations have a very hard spectrum. In Figure 18, we show the diagnostic diagrams for the different incident spectra listed above. The O star model lies near the Kauffmann demarcation on the $[O III]/H\beta$ vs $[N II]/H\alpha$ and $[O III]/H\beta$ vs $[S II]/H\alpha$ diagrams as expected. The 13 Gyr SSP grid shows significant higher $[S II]/H\alpha$, $[N II]/H\alpha$, and $[O II]/H\alpha$ relative to H II regions with the same ionization parameters due to a more extended partially-ionized region. Models (2) and (3) show a harder ionizing spectrum than H II region, but their line ratios are similar to those for H II regions. Model (4) exhibits enhancement in $[N II]/H\alpha$, $[S II]/H\alpha$, and $[O II]/H\alpha$ relative to H II regions. The enhancements are small, typically less than 0.2 dex. However, model (5), which has the hardest spectrum blueward of 912\AA shows the lowest $[N II]/H\alpha$, $[S II]/H\alpha$, and $[O II]/H\alpha$. This is because even though those photons lower than 13.6 eV cannot ionize Hydrogen, they

have enough energy to excite an electron in the Hydrogen atom from ground level ($n=1$) to $n=3$ or $n=4$ level. When they jump back, they can jump to $n=2$ which will produce $H\alpha$ and $H\beta$ lines. This is fluorescent production. In model (5), photons pass through a very high column density which absorbs more than 99.9% of all the ionizing photons. When the transmitted spectrum is forced to have $\log U > -4.5$, we obtain an unphysically high continuum redward of 912\AA , which leads to large fluorescent production. It is the enhancement of $H\alpha$ due to fluorescent production that gives rise to lower $[N II]/H\alpha$, $[S II]/H\alpha$, $[O II]/H\alpha$, and $[O I]/H\alpha$.

Considering the large amount of ionizing photons from OB stars and the spatial correlation of DIG and H II regions, leaky H II regions are likely a major mechanism for producing DIG. However, our results disfavor leaky H II region models to account for ionization of ALL DIG. DIG has a lower ionization parameter than H II regions, a decrease in ionization parameter leads to enhancement of $[N II]/H\alpha$, $[S II]/H\alpha$, $[O II]/H\alpha$, and $[O I]/H\alpha$, but a decrease in $[O III]/H\beta$ at the same time. Leaky H II region models can not produce LI(N)ER-like emission. The cyan line shows enhancement of these four line ratios, but it needs fine tuning of the filtering column density. DIG that shows LI(N)ER-like emission needs another ionization source. We favor evolved stars as a major ionization source for DIG because only ionization by




Figure 13. The $\Delta \log$ (line ratios) vs $\Delta \log \Sigma_{H\alpha}$ relation at $[0.4R_e, 0.6R_e]$ for the one third least massive galaxies. We normalize the log line ratios vs $\log \Sigma_{H\alpha}$ relation by subtracting the median log line ratio and median $\log \Sigma_{H\alpha}$. All the spaxels in a galaxy are weighted by $\frac{1}{N}$, where N is the number of valid spaxels in the galaxy. $[S II]/H\alpha$, $[N II]/H\alpha$, $[O II]/H\beta$, $[O I]/H\alpha$ are higher in DIG dominated low $\Sigma_{H\alpha}$ regions. $[O III]/[O II]$ is slightly lower in DIG while $[O III]/H\beta$ is slightly lower in DIG.




Figure 14. $[O III]/[O II]$ vs $[N II]/[O II]$ diagram (Dopita et al. 2000, 2013). The grids are from Dopita et al. (2013) with $\kappa = Inf$. The nearly-horizontal lines are for constant ionization parameters, and the nearly vertical lines are for constant metallicities. The labels denote the metallicities and ionization parameters ($\log q$) for the grid.

evolved stars (red line, Model 6) can produce enhancement of $[S II]/H\alpha$, $[N II]/H\alpha$, $[O II]/H\alpha$, $[O I]/H\alpha$, and $[O III]/H\beta$ even when the ionization parameter drops. Evolved stars are prevalent all over galaxies, and their contribution to ionizing photons may prevail. Flores-Fajard et al. (2011) proposed hot low-mass evolved stars (HOLMES) as an impor-

tant ionization source for the extra-planar diffuse ionized gas in edge-on galaxies. For NGC 891, HOLMES begin to contribute more than 50% of the ionizing photons at scale height $|z| > 2$ kpc. We propose here that the evolved stars are not only capable of ionizing the extraplanar gas, but also an important ionizing source for the DIG near the disk.




Figure 15. $[\text{O III}]/\text{H}\beta$ vs $[\text{N II}]/\text{H}\alpha$ diagram for each galaxy. Different colors represent different $\Sigma_{\text{H}\alpha}$. The solid and dashed lines are the demarcation line from Kauffmann et al. (2003) and Kewley et al. (2001). The grids are from Dopita et al. (2013) with $\kappa = \text{Inf}$. DIG can not be covered by the grids, suggesting HII region models with variations of metallicity and ionization parameter can not produce the DIG line ratios.




Figure 16. $[\text{O III}]/\text{H}\beta$ vs $[\text{S II}]/\text{H}\alpha$ diagram for each galaxy. Different colors represent different $\Sigma_{\text{H}\alpha}$. The solid lines are the demarcation lines to separate Seyfert galaxies from LINER in Kewley et al. (2006). The grids are from Dopita et al. (2013) with $\kappa = \text{Inf}$. The grids do not cover DIG, suggesting HII region models with variations of metallicity and ionization parameter can not produce the DIG line ratios.

4.3 Photoionization Grids for SSP at different ages

We have shown in last subsection that a 13 *Gyr* SSP can produce the LI(N)ER-like emission we see for DIG. One interesting question is: How would the line ratios change as a SSP ages? To answer this question, we generate SSPs at 1 *Myr*, 3 *Myr*, 9 *Myr*, 0.125 *Gyr*, 7 *Gyr*, and 13 *Gyr* using BC03, and use these spectra as the incident ionizing spectra for CLOUDY. The spectra from 0.125 *Gyr* to 13 *Gyr* change only slightly. The metallicity is that of the Orion Nebula, and the density is 1 cm^{-3} . We run the calculation for $\log U = -4.5$ to -2.0 with an interval of 0.5 dex. The incident spectra are shown in Figure 19. The output line ratios are plotted in the same color as their incident spectra in Figure 20. At 1 *Myr*, the line ratios are located at the HII region part on the diagnostic diagrams. At 3 *Myr*, the line ratios are similar to those of 1 *Myr*. At 9 *Myr*, however, we see a significant decrease in $[\text{N II}]/\text{H}\alpha$, $[\text{S II}]/\text{H}\alpha$, $[\text{O II}]/\text{H}\alpha$, $[\text{O III}]/\text{H}\beta$, $[\text{O III}]/[\text{O II}]$, and an increase in $[\text{N II}]/[\text{O II}]$. This is due to the significant decrease in 200-500Å spectrum hard-

ness as OB stars age (Levesque et al. 2010). At 125 *Myr*, the line ratios are already in the LI(N)ER/AGN region. This means once OB stars die, the line ratios of the ionized gas would turn into “LI(N)ER/AGN like” very fast. The line ratios do not change very much afterwards. This sheds light on the interpretation of LI(N)ER/AGN like emission line ratios in galaxies. When OB stars are alive, their light dominates the ionizing spectrum and the ionized gas located in the star-forming regions dominates on the diagnostic diagrams. When OB stars cease to form, their contribution disappears very fast, and the light of hot evolved stars begins to dominate after tens of *Myr*, and the ionized gas is located in the LI(N)ER/AGN regions on the diagnostic diagrams. We have shown in Section 4.2 that a filtered spectrum can not reproduce the DIG line ratios with an increase of $[\text{O III}]/\text{H}\beta$. A higher $[\text{O III}]/\text{H}\beta$ in DIG than in HII regions means the ionization of DIG may not be linked to OB stars directly but linked to the hot evolved stars.

We are not claiming that evolved stellar populations are the major ionizing source of DIG. Overall, the radiative




Figure 17. The incident spectra for our models described in Section 4.2. All spectra are normalized to the flux at 1000Å. The magenta line is a 42,300K, $\log g=4.22$, solar metallicity O star generated by *Thusty* (Lanz & Hubeny 2003). The orange, green, cyan, and blue lines are the spectra by filtering the O star spectrum through column density of 10^{18} cm^{-1} , $10^{18.5} \text{ cm}^{-1}$, $10^{18.7} \text{ cm}^{-1}$, $10^{18.9} \text{ cm}^{-1}$, simulating the leaky H II regions. The red line is the spectrum of a 13Gyr Simple Stellar Population generated by BC03.




Figure 18. The BPT diagrams, $\log [\text{O III}]/\text{H}\beta$ vs $\log [\text{O I}]/\text{H}\alpha$ diagram, $\log [\text{O III}]/[\text{O II}]$ vs $\log [\text{N II}]/[\text{O II}]$ diagram, and $\log [\text{O III}]/[\text{O II}]$ vs $\log [\text{O II}]/\text{H}\alpha$ diagrams for different incident spectra shown in Figure 17. The color scheme is the same as in Figure 17. The dots with the same color have different ionization parameters from $\log U=-4.5$ to $\log U=-2$ with an interval of 0.5 dex. The leaky H II region models (orange, green, cyan lines) show very similar line ratios to the O star model (magenta line) at the same ionization parameter. We note that DIG has lower ionization parameters than H II regions. Leakly H II region models can not produce enhancement of $[\text{S II}]/\text{H}\alpha$, $[\text{N II}]/\text{H}\alpha$, $[\text{O II}]/\text{H}\alpha$ and $[\text{O III}]/\text{H}\beta$ with a decrease in ionization parameter in most cases. The cyan line shows enhancement of these three line ratios, but it needs fine tuning of the filtering column density. The blue line even shows decreases in $[\text{S II}]/\text{H}\alpha$, $[\text{N II}]/\text{H}\alpha$, $[\text{O II}]/\text{H}\alpha$ and $[\text{O III}]/\text{H}\beta$, because of large H α fluorescent production, which is unphysical. Only ionization by evolved stellar populations (red line) can produce enhancement of $[\text{S II}]/\text{H}\alpha$, $[\text{N II}]/\text{H}\alpha$, $[\text{O II}]/\text{H}\alpha$ and $[\text{O III}]/\text{H}\beta$ even when the ionization parameter drops.

and mechanical energy from hot evolved stars falls short of the energy budget of diffuse ionized gas in star-forming galaxies (Reynolds 1984; Ferguson et al. 1996; Binette et al. 1994). Even if the energy emitted by hot evolved stars meets the energy requirement of DIG, O stars can provide at least one order of magnitude more ionizing photons than hot evolved stars in star-forming galaxies (Reynolds 1984; Ferguson et al. 1996). Indeed, DIG is found to be linked to H II regions both along the disk and in vertical direction (e.g. Ferguson et al. 1996; Zurita et al. 2000; 2002; Rossa & Dettmar 2003a,b). In star-forming galaxies, hot evolved stars cannot compete with O stars in total ionizing photons production. However, we see in two cases that hot evolved stars might be a major contributor of ionizing photons. 1) In low surface brightness regions that are located far away from H II regions. The density of ionizing photons from the H II regions has dropped significantly so the hot evolved stars begin to dominate. One example is in regions at large vertical height (extra-planar gas at high $|z|$). Other low surface brightness regions that show higher $[\text{O III}]/\text{H}\beta$ than H II regions probably belong to this category as well. 2) In galaxies where OB stars have died: post-starburst galaxies and quiescent/passive galaxies. Quiescent galaxies are known to show LI(N)ER emissions (Yan et al. 2006; Sarzi et al. 2006; Stasińska et al. 2008; Kehrig et al. 2012; Singh et al. 2013; Gomes et al. 2016; Belfiore et al. 2015, 2016a,b). We predict DIG to be prevalent in post-starburst galaxies, and they will show LI(N)ER/AGN like emission.

5 IMPACT OF DIG ON METALLICITY MEASUREMENT

There are several methods to measure gas-phase metallicity in star-forming galaxies using strong lines in the optical, such as $\text{N2}=[\text{N II}]/\text{H}\alpha$, $R_{23}=(\text{[O II]} \lambda 3727 + \text{[O III]} \lambda 4959 + \text{[O III]} \lambda 5007)/\text{H}\beta$, $\text{O3N2}=(\text{[O III]}/\text{H}\beta)/(\text{[N II]}/\text{H}\alpha)$ (e.g., Alloin et al. 1979; Zaritsky et al. 1994; Pilyugin, 2001; Denicoló et al. 2002; Pettini & Pagel, 2004; Pilyugin & Thuan, 2005) and $\text{N2O2}=(\text{[N II]}/\text{[O II]})$ (Dopita et al. 2000; Kewley et al. 2001), $\text{N2S2H}\alpha=8.77+\log[\text{N II}]/\text{[S II]}+0.264 \times \log[\text{N II}]/\text{H}\alpha$ (Dopita et al. 2016). If the line ratios of DIG are different from these in H II regions, the metallicity measurement will inevitably be biased because these metallicity indicators are a combination of several line ratios. The bias depends on the fraction of emission contributed by DIG, which increases towards low surface brightness regions.

5.1 $\text{N2}=[\text{N II}]/\text{H}\alpha$

If we use N2 to derive the metallicity, the enhancement seen in DIG means that the metallicity will be overestimated in those spaxels with a high DIG fraction. This is especially important for observation of galaxies at high redshift when only the emission lines near $\text{H}\alpha$ are available. We see a typical $[\text{N II}]/\text{H}\alpha$ enhancement of 0.2 dex. We use the equation in Pettini & Pagel (2004) to convert $[\text{N II}]/\text{H}\alpha$ to metallicity: $12+\log(\text{O}/\text{H})=9.37+2.03 \times \text{N2}+1.26 \times \text{N2}^2+0.32 \times \text{N2}^3$, $\text{N2}=\log([\text{N II}]/\text{H}\alpha)$. Assuming a H II region has $\text{N2}=-0.5$, $\text{Z}(\text{N2})=8.63$ in a local galaxy, we obtain $\text{Z}(\text{N2})=8.865$ for DIG with the same metallicity because $\text{N2}=-0.3$. The typical

metallicity gradient for a local star-forming galaxy is about $-0.1 \text{ dex } R_e^{-1}$ (Sánchez et al. 2014; Ho et al. 2015), meaning metallicity drops by ~ 0.2 from the center to $2R_e$. The bias due to DIG would flatten the metallicity gradient from $-0.1 \text{ dex } R_e^{-1}$ to 0 if DIG totally dominates at $2R_e$. If we assume a metal poor H II region has $\text{N2}=-1.5$, $\text{Z}(\text{N2})=8.08$ in a high redshift galaxy, DIG with the same metallicity has $\text{N2}=-1.3$ and we would obtain $\text{Z}(\text{N2})=8.157$. The metallicity gradient will be flattened as well. N2 is usually used for measuring metallicity and metallicity gradients at high redshift (e.g., Wuyts et al. 2016). Our result indicates that the presence of DIG potentially flattens N2 derived metallicity gradients at high redshift.

5.2 $\text{N2O2}=[\text{N II}]/[\text{O II}]$

N2O2 is a good metallicity indicator because it is not sensitive to ionization parameter or ionizing spectrum hardness but to metallicity (Dopita et al. 2000, 2013; Kewley et al. 2002). The caveat of N2O2 is that it relies on the Nitrogen-to-Oxygen abundance ratio (N/O). Nitrogen and α elements have different enrichment timescales, hence their ratio depends on the star formation history and several other parameters (e.g. Vincenzo et al. 2016) as well as mixing issues (e.g. Belfiore et al. 2015), especially for low mass galaxies and galaxy outskirts, where there are prominent variations of N/O vs O/H relative to metal rich systems and central regions. Moreover, these diagnostics are sensitive to metallicity only at $12+\log(\text{O}/\text{H})>8.3$, i.e. where N/O is a strong function of metallicity, but they become essentially insensitive to metallicity at $12+\log(\text{O}/\text{H})<8.3$, i.e. in the regime where N/O is nearly constant. We plot metallicities derived using $\text{N2O2}=[\text{N II}]/[\text{O II}]$ as a function of effective radius and color-code the dots with $\text{H}\alpha$ surface brightness in Figure 21. N2O2 is converted to metallicity using $\log(\text{O}/\text{H})+12=\log[1.54020+1.26602 \times R+0.167977 \times R^2]+8.93$, $R=\log[\text{N II}]/[\text{O II}]$ (Dopita et al. 2013). At fixed radius, $\text{Z}(\text{N2O2})$ in the low surface brightness bin is similar to or a little bit higher than that in the high surface brightness bin. As stated in Section 3.2, $[\text{N II}]/[\text{O II}]$ only depends on the N/O abundance ratio and temperature (Dopita et al. 2000, 2013). It is not subject to the ionization parameter and ionizing spectrum shape variation. This makes it an excellent metallicity indicator. The weak dependence of $[\text{N II}]/[\text{O II}]$ on $\text{H}\alpha$ surface brightness at fixed radius (Figure 21) demonstrates it is a good metallicity indicator even in the presence of DIG. However, we note that N2O2 is subject to variation of N/O ratio (Pérez-Montero & Contini 2009; Pérez-Montero et al. 2013, 2016), temperature variation, and uncertainty of extinction correction. $\text{Z}(\text{N2O2})$ could be enhanced or suppressed for DIG in some regions, maybe due to N/O variation and temperature variation. The measurement of extinction involves an accurate measurement of $\text{H}\beta$ which is not an easy task since the stellar continuum shows deep $\text{H}\beta$ absorption. This is even harder when the emission line is weak and S/N is low, such as in the outskirts of galaxies. Besides, $\text{H}\alpha$ and $\text{H}\beta$ arise from the fully ionized region, while $[\text{N II}]$ and $[\text{O II}]$ are from the partially-ionized region. The extinction derived using $\text{H}\alpha$ and $\text{H}\beta$ may not necessarily be the same as the extinction experienced by the partially-ionized region. From Figure 6, we don't see any signs of deviation of $\text{H}\beta/\text{H}\alpha$ for different $\text{H}\alpha$ surface brightness spaxels, so the result that




Figure 19. The incident spectra for SSP at 0.001 Gyr, 0.003 Gyr, 0.009 Gyr, 0.125 Gyr, 7 Gyr, and 13 Gyr with solar metallicity, normalized to flux at 6000Å. We can see that OB stars dominate the ionizing spectrum in the beginning, and wane after tens of Myr. After that the ionizing spectrum is dominated by emission from evolved stellar populations and it hardens.




Figure 20. BPT diagrams, $\log [\text{O III}]/\text{H}\beta$ vs $\log [\text{O I}]/\text{H}\alpha$, $\log [\text{O III}]/[\text{O II}]$ vs $\log [\text{N II}]/[\text{O II}]$, and $\log [\text{O III}]/[\text{O II}]$ vs $\log [\text{O II}]/\text{H}\alpha$, for different incident spectra of SSPs at different ages. The color scheme is the same as Figure 19. The dots with the same color have different ionization parameters from $\log U = -4.5$ to $\log U = -2$ with an interval of 0.5 dex. At Age < tens of Myr, the ionizing spectrum softens as OB stars evolve and die. $[\text{S II}]/\text{H}\alpha$, $[\text{N II}]/\text{H}\alpha$, $[\text{O II}]/\text{H}\alpha$ and $[\text{O III}]/\text{H}\beta$ all decrease. After the OB stars die, the ionizing spectrum is dominated by hot evolved stars, and the spectrum hardens. We see increases of $[\text{S II}]/\text{H}\alpha$, $[\text{N II}]/\text{H}\alpha$, $[\text{O II}]/\text{H}\alpha$ and $[\text{O I}]/\text{H}\alpha$. This figure shows how the ionized gas moves on the diagnostic diagrams as the stellar population ages.

DIG have similar N2O2 as H II region is robust. We see in Figure 6 that $H\beta/H\alpha$ shows a gradient with positive slope even though it does not depend on $\Sigma_{H\alpha}$. The gradient in $H\beta/H\alpha$ will change the $Z(N2O2)$ gradient slope. To correct this, we suggest getting an $H\beta/H\alpha$ gradient by fitting the $H\beta/H\alpha$ of individual spaxels and apply the overall extinction gradient to $[N II]/[O II]$ to get the correct metallicity gradient. In this way, we avoid the very noisy $H\beta/H\alpha$ when $\Sigma_{H\alpha}$ is low when correcting extinction.

A key for probing abundance is to use lines from different elements with the same ionization potential. The ionization potential for H^0 , O^0 , O^+ , O^{++} , S^+ and N^+ are 13.598eV, 13.618eV, 35.121eV, 54.936eV, 23.337eV, and 29.601eV respectively. Ionization potential of $[O II]$ and $[N II]$ are similar: 35.121eV and 29.601eV. This makes $[N II]/[O II]$ insensitive to ionization parameter, thus a good metallicity indicator. The complexity in using $[N II]/[O II]$ is the extinction correction and the N/O ratio variation at fixed O/H.

To quantify the change of the metallicity measurements as a function of $H\alpha$ surface brightness, we do a linear regression fit to the derived metallicities vs $H\alpha$ surface brightness relation in a narrow annulus. The distribution of slopes for different metallicity indicators at three radii are shown in black, blue, and red in Figure 29. The metallicity is independent of surface brightness. The reason we are seeing a dependence of metallicity on $H\alpha$ surface brightness is due to contamination by DIG. For $Z(N2O2)$, the slope distribution is narrow, and peaks at ~ 0.05 . The bias in metallicity using N2O2 is small and the error is small because N2O2 does not vary much among DIG. DIG has marginally lower $Z(N2O2)$, suggesting marginally higher temperature.

5.3 R_{23}

$R_{23} = ([O II] \lambda 3727 + [O III] \lambda 4959 + [O III] \lambda 5007) / H\beta$ is a commonly used metallicity indicator (McGaugh 1991; Zaritsky et al. 1994; Pilyugin 2001; Kewley & Dopita 2002; Kobulnicky & Kewley 2004). The advantage is it does not depend on the N/O ratio. The disadvantage is that the metallicity is double-valued at a given R_{23} . In order to use R_{23} to measure Z, one has to first determine the ionization parameter and which branch it is on. Therefore, it always needs to be used in conjunction with $[O III]/[O II]$ and N2O2 (or similar). We use the method of Kewley & Ellison (2008) to determine which branch a spaxel is on and derive the metallicity. Figure 22 show how the Z derived from R_{23} changes with radius and $H\alpha$ surface brightness. One can see that low SB regions have lower measured $Z(R_{23})$ values. We have shown in Section 3.2 that DIG has higher $[O II]/H\beta$, lower $[O III]/[O II]$, and on average similar $[O III]/H\beta$ as H II regions. The combination of the three line ratios leads to the result that $Z(R_{23})$ is biased to be lower in DIG.

For the whole sample, the $Z(R_{23})$ vs $\Sigma_{H\alpha}$ slope distribution peaks at ~ 0.2 and the dispersion is large. The contribution of DIG would not only bias the metallicity measurement systematically, but also introduce large metallicity measurement uncertainties, as the scatter in the metallicity vs. radius relation is large. We note that the gas-phase metallicity of star-forming galaxies shows a universal gradient of $-0.1 \text{ dex } R_e^{-1}$. Meanwhile, the slope of $Z(R_{23})$ vs $\Sigma_{H\alpha}$ is ~ 0.2 . According to Figure 22, $\Sigma_{H\alpha}$ of DIG at $2R_e$ is about $38 \text{ erg s}^{-1} \text{ kpc}^{-2}$. Compared to a $\Sigma_{H\alpha} = 39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ H II

region, the $Z(R_{23})$ bias for DIG would be 0.2. Note the drop of metallicity from the center to $2R_e$ is only 0.2 dex. The metallicity gradient would be $-0.2 \text{ dex } R_e^{-1}$ if the outskirts of the galaxy is pure DIG. In other words, the metallicity measured using $Z(R_{23})$ is biased systematically by $\sim -0.1 \text{ dex } R_e^{-1}$.

5.4 O3N2

O3N2 is sensitive to oxygen abundance, and it is not impacted by extinction. The disadvantage of O3N2 is it depends on N/O (Pérez-Montero & Contini 2009; Pérez-Montero et al. 2013, 2016) and the ionization parameter. In Figure 23, we show how DIG would impact metallicity measurements made from $O3N2 = ([O III]/H\beta) / ([N II]/H\alpha)$. We see in Figure 23 that O3N2 is higher in DIG for some galaxies and lower or similar in other galaxies. The contamination by DIG could be responsible for a substantial portion of the scatter in metallicity measurements. When confined only to the high surface brightness regions, the metallicity gradient derived using O3N2 is similar to the ones using R_{23} or N2O2. To make robust metallicity gradient measurements, one has to properly isolate H II regions and correct for DIG contamination.

Because the surface brightness generally decreases towards large radii, the metallicity gradient derived using O3N2 at large radii might be flatter or steeper than that derived using N2O2 if we include all the spaxels. For example, in MaNGA galaxy 8313-12702, we get a metallicity of $-0.15 \text{ dex } R_e^{-1}$ using $Z(N2O2)$. Using only the high surface brightness regions (red dots) and $Z(O3N2)$ gives the same result. However, if we include everything, the metallicity gradient derived using O3N2 will be $-0.1 \text{ dex } R_e^{-1}$. The bias is significant at least for this galaxy. $Z(O3N2)$ is biased in the other direction for MaNGA galaxy 8603-12704. Similarly, Mast et al. (2014) found that the presence of DIG would bias the abundance gradient when using O3N2 to derive metallicity. The strength of the bias depends a lot on the assumed calibrator. We note the degree of bias is not the same for all galaxies, and we definitely can give a robust metallicity gradient measurement using high surface brightness regions. The σ of the distribution of $Z(O3N2)$ vs $\Sigma_{H\alpha}$ slope at fixed radius is ~ 0.1 . This translates to a metallicity gradient dispersion of $\pm 0.05 \text{ dex } R_e^{-1}$ through a similar analysis as in Section 5.3, consistent with the different biases we see in different galaxies. For a large sample of galaxies (Sánchez et al. 2014, Ho et al. 2015), the bias for different galaxies may cancel out when deriving average metallicity gradients.

5.5 N2S2H α

Dopita et al. (2016) proposed a new metallicity proxy using only $[N II] \lambda 6548$, $[S II] \lambda \lambda 6717, 6731$, and $H\alpha$: $12 + \log(O/H) = Z(N2S2H\alpha) = 8.77 + \log[N II]/[S II] + 0.264 \times \log[N II]/H\alpha$. It is especially suitable for metallicity measurements of high redshift galaxies whose spectral wavelength coverage is limited because the 4 lines used are close to each other. This estimator is almost linear up to an abundance of $12 + \log(O/H) = 9.05$. The caveat is the calibration of this metallicity proxy is based on a well defined N/O and O/H relation. This calibration fails for any systems deviating from




Figure 21. Metallicity derived using N2O2 as a function of radius. $Z(\text{N2O2})$ drops with radius due to a metallicity gradient. We see that the derived metallicity does not depend on the $\Sigma_{\text{H}\alpha}$ at a fixed radius, and the dispersion at a fixed radius is similar to the measuring error. Red and blue bars are errors at $\log \Sigma_{\text{H}\alpha}=39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log \Sigma_{\text{H}\alpha}=38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ respectively. The length of the error bars is 2σ . We show in Section 5.7 that the impact of DIG is prevalent in all star-forming galaxies in our sample, not confined to the three galaxies we show.




Figure 22. Metallicity derived using R_{23} as a function of radius. $Z(R_{23})$ drops with radius due to metallicity gradient. At fixed radius, $Z(R_{23})$ decreases with $\Sigma_{\text{H}\alpha}$, indicating $Z(R_{23})$ is systematically biased low for DIG. The measuring errors at $\log \Sigma_{\text{H}\alpha}=39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log \Sigma_{\text{H}\alpha}=38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ are indicated using red and blue bars. The dispersion of $Z(R_{23})$ is about twice that of $Z(\text{N2O2})$. The metallicity gradient would be systematically biased by $-0.1 \text{ dex } R_e^{-1}$ if the bias in DIG is not accounted for.




Figure 23. Metallicity derived using O3N2 as a function of radius. $Z(\text{O3N2})$ drops with radius due to a metallicity gradient. At fixed radius, $Z(\text{O3N2})$ could either decrease or increase with $\Sigma_{\text{H}\alpha}$. The measuring errors at $\log \Sigma_{\text{H}\alpha}=39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log \Sigma_{\text{H}\alpha}=38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ are indicated using red and blue bars. The dispersion of $Z(\text{O3N2})$ is similar to that of $Z(\text{N2O2})$. The metallicity gradient would be biased by $\pm 0.05 \text{ dex } R_e^{-1}$ if the bias in DIG is not considered.

the assumed N/O–O/H relation. We explore how DIG biases the metallicities derived using N2S2H α in Figure 24. At fixed radius, we don't see a significant offset in Z(N2S2H α) between DIG and H II regions. This is because Z(N2S2H α) is insensitive to variation of ionization parameter and spectral hardness (Dopita et al. 2016), very similar to N2O2. Compared with Z(N2O2), the dispersion of Z(N2S2H α) is larger. This is understandable because:

(a) Four lines are used in Z(N2S2H α) while only 2 lines are used to derive Z(N2O2). The measurement errors enter into the derivation of metallicities.

(b) The relative abundances of N, S, and O are involved in Z(N2S2H α) while Z(N2O2) only include N/O.

According to the upper right panel of Figure 29, The distribution of Z(N2S2H α) vs $\Sigma_{H\alpha}$ slope is very similar to that of Z(O3N2) in its centroid and dispersion. The slope distribution for Z(N2S2H α) has a wider wing, meaning there are some sources that show significant biases in Z(N2S2H α) for DIG and H II regions. This is similar to the scatter introduced by DIG on O3N2.

5.6 IZI

IZI (Blanc et al. 2015) uses strong nebular emission lines to derive the Bayesian posterior probability density function for metallicity and ionization parameter based on a series of H II region models. In Figure 25, we plot the metallicities derived using IZI as a function of radius. We input [O II] λ 3727, [O III] λ 4959, [O III] λ 5007, H α , [N II] λ 6583, and [S II] λ 6717, 6731 for the calculation. All the emission line fluxes are corrected for reddening using the Balmer decrement H α /H β assuming a Milky Way extinction curve (Fitzpatrick 1999). We use the values derived with the "output joint mode" in IZI. The dots are color-coded by $\Sigma_{H\alpha}$. For high $\Sigma_{H\alpha}$ regions, IZI gives similar metallicity gradients as the other metallicity indicators. However, at a fixed radius, different surface brightness regions (different colors in the plots) have large metallicity discrepancies. For low $\Sigma_{H\alpha}$ regions, the metallicities derived from IZI can vary by 0.2 dex among themselves. This is at least partly because IZI currently only adopts H II region grids. These models make an inherent assumption that the ionizing spectrum shape is fixed by the temperature of the OB stars, which is determined by the metallicity. However, for DIG this assumption does not hold. We have shown that metallicity+low q for an H II region model can not produce the line ratios we see in DIG. Thus, IZI is very vulnerable to contamination by DIG because currently it only contains H II region models. However, if one were to include a DIG model in IZI and other Bayesian codes to derive metallicities (e.g., HII-CHI-mistry; Pérez-Montero 2014; BOND: Vale Asari et al 2016), then they may be able to give accurate estimates of metallicity and ionization parameter even with contamination of DIG.

Furthermore, we input only extinction–corrected [N II] and [O II] into IZI to see if IZI could give a less biased metallicity in DIG. [N II] and [O II] can only be combined to N2O2 to estimate metallicity. Figure 26 shows that the metallicity derived through IZI using [N II] and [O II] only is better than metallicity derived using all strong emission lines available. The dispersion at a fixed radius is smaller and the dependence on $\Sigma_{H\alpha}$ at a fixed radius is weaker. However, Z(N2O2) in Figure 21 still shows a tighter metallicity gradi-

ent than Z(IZI) [N II]+[O II]. This is because we do not apply extinction corrections to individual spaxels when calculating Z(N2O2). IZI metallicity with extinction–uncorrected [N II] and [O II] input alone gives very similar result as Z(N2O2).

5.7 Comparison of different metallicity estimators

5.7.1 Metallicities vs $\Sigma_{H\alpha}$ relation: The Whole Sample

To quantify the bias introduced by DIG on strong–line metallicity measurements, we examine the derived metallicities as a function of $\Sigma_{H\alpha}$ for the whole sample. We select all spaxels in $[0.4R_e, 0.6R_e]$, subtract from all metallicity and $\log \Sigma_{H\alpha}$ measurements their respective medians, then plot all galaxies together in the same plot. The spaxels in each galaxy are weighted by $\frac{1}{N}$, where N is the number of valid spaxels in this galaxy. Figure 28 shows the weighted ΔZ vs $\Delta \log \Sigma_{H\alpha}$ relation for the whole sample. A non–zero slope of the relation means metallicities are biased by DIG contribution, and the dispersion of this relation in y direction reflects how reliable a metallicity proxy is for an individual galaxy. The $\Delta Z(N2O2)$ vs $\Delta \Sigma_{H\alpha}$ relation has a slope of ~ 0.05 . The slope of $\Delta Z(R_{23})$ vs $\Delta \log \Sigma_{H\alpha}$, $\Delta Z(O3N2)$ vs $\Delta \log \Sigma_{H\alpha}$, $\Delta Z(N2S2H\alpha)$ vs $\Delta \log \Sigma_{H\alpha}$, $\Delta Z(N2)$ vs $\Delta \log \Sigma_{H\alpha}$, relations are ~ 0.2 , -0.05 , ~ 0 , and -0.2 . We define the dispersion as the interval between 90% percentiles contours at $\Delta \log \Sigma_{H\alpha} = 0$. For Z(N2O2), the dispersion is 0.2, and for Z(R_{23}), Z(O3N2), Z(N2S2H α) and Z(N2), the dispersions are 0.35, 0.23, 0.37 and 0.25. Based on slopes and dispersions, Z(N2O2) is optimal because the slope is mild and the dispersion is smallest. Z(R_{23}) has the most significant slope and large dispersion. Z(O3N2) has smallest slope and a dispersion only slightly larger than Z(N2O2). Z(N2S2H α) has a nearly 0 slope but the dispersion is the largest. Z(N2) is derived using [N II]/H α , thus very vulnerable to DIG contamination.

We also do linear regression to the derived metallicities vs $\log \Sigma_{H\alpha}$ relation in narrow radial annuli so the metallicity is essentially fixed. The distribution of slopes for different metallicity indicators at three radii are shown in black, blue, and red in Figure 29. The intrinsic metallicity is independent of surface brightness. The reason we are seeing a dependence of derived metallicities on H α surface brightness is due to contamination by DIG. A slope of 0 means the DIG does not impact the metallicity measurements, and a positive slope means DIG biases the metallicity measurement low. The centroid of the distribution reflects the systematic offset of that metallicity indicator for the whole sample, while the dispersion of the distribution reflects the metallicity error it brings for individual galaxies. For Z(N2O2), the slope distribution is narrow, and peaks at $\sim 0.05 \text{ dex}^{-1}$. The bias in metallicity using N2O2 is small and the error is small because N2O2 does not vary much among DIG. DIG regions have marginally lower Z(N2O2), suggesting slightly higher temperature. The slope distribution of Z(N2S2H α) is very similar to that of Z(N2O2), suggesting Z(N2S2H α) to be a robust metallicity estimator even in the presence of DIG. For Z(R_{23}), the slope distribution peaks at 0.2 dex^{-1} and dispersion is large. The contribution of DIG would not only bias the metallicity measurements systematically, but also introduce a large metallicity measurement error. This can be seen in the tightness of metallicity vs radius relation. For




Figure 24. Metallicity derived using $[\text{N II}]/[\text{S II}]$ and $[\text{O II}]/\text{H}\alpha$ proposed by Dopita et al. (2016) as a function of radius. We see that the derived metallicity does not depend on the $\Sigma_{\text{H}\alpha}$ at a fixed radius, and the dispersion at a fixed radius is similar to the measuring error (red and blue bars are errors at $\log \Sigma_{\text{H}\alpha}=39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log \Sigma_{\text{H}\alpha}=38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ respectively). $Z(\text{N2S2H}\alpha)$ performs very similar to $Z(\text{O3N2})$.




Figure 25. Metallicity derived using IZI by Blanc et al. (2015) as a function of radius. The dots are color-coded by $\Sigma_{\text{H}\alpha}$. For high $\Sigma_{\text{H}\alpha}$ regions, the metallicity gradients are similar to those derived using other metallicity indicators. However, at fixed radius, different surface brightness regions have large metallicity discrepancies. For low $\Sigma_{\text{H}\alpha}$ regions, the metallicities derived from IZI can vary by 0.2 dex themselves. This demonstrates that IZI is not robust to derive the metallicity of DIG. Note the metallicity lower limit is set to 8, so the cut-off at $Z \sim 8.3$ is not due to a high metallicity lower limit.




Figure 26. Metallicity derived using IZI by Blanc et al. (2015) as a function of radius using only extinction corrected $[\text{N II}]$ and $[\text{O II}]$. The impact of DIG is much less than when using all strong emission lines available. The red and blue bars show the typical line ratio errors at $\log \Sigma_{\text{H}\alpha}=39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log \Sigma_{\text{H}\alpha}=38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ for individual spaxels. The length of the error bars is 2σ .




Figure 27. Metallicity derived using IZI by Blanc et al. (2015) as a function of radius using only extinction corrected [N II], [O II], and [O III]. The result is indistinguishable from the result using only [N II] and [O II]. The impact of DIG is much less than when using all strong emission lines available. The red and blue bars show the typical line ratio error at $\log \Sigma_{\text{H}\alpha} = 39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ and $\log \Sigma_{\text{H}\alpha} = 38.5 \text{ erg s}^{-1} \text{ kpc}^{-2}$ for individual spaxels. The length of the error bars is 2σ .

$Z(\text{O3N2})$, the distribution of slope is broad, and peaks at $\sim -0.05 \text{ dex}^{-1}$. The magnitude of the systematic metallicity measurement bias is similar for $Z(\text{O3N2})$ and $Z(\text{N2O2})$, but $Z(\text{O3N2})$ in DIG varies much more significantly. Using $Z(\text{O3N2})$ will not bias the measurement if we average a large sample of galaxies, but it will inevitably bias the metallicity in individual galaxies as shown in Section 5.4. The large variation of $Z(\text{O3N2})$ and $Z(R_{23})$ in DIG is expected because these two indicators involve ionization parameters. For an H II region, this does not matter because the metallicity and ionization parameter are linked to each other. This assumption does not hold for DIG. The variation in the ionization parameter in DIG enters the metallicity measurement. The impact of DIG on using $Z(\text{N2})$, $Z(R_{23})$ and $Z(\text{O3N2})$ metallicity gradients is discussed in Section 5.1, Section 5.3 and Section 5.4 respectively. For $Z(\text{IZI})$, the metallicities derived for DIG are systematically lower. Besides, the dispersion of the slope distribution is broadest, with $\sigma \sim 0.3$. This means IZI not only systematically biases the metallicity of DIG lower, but also introduces large errors into the metallicity measurements.

5.7.2 Metallicities vs $\Sigma_{\text{H}\alpha}$ relation: Split by Stellar Mass

In Figure 30 and Figure 31, we show the normalized Z vs $\Sigma_{\text{H}\alpha}$ relations for galaxies with stellar mass less than $10^{9.43}$ and higher than $10^{10.08}$ respectively. They are the one third least massive and one third most massive galaxies in our sample. The $\Delta Z(\text{N2O2})$, $\Delta Z(R_{23})$, $\Delta Z(\text{N2S2H}\alpha)$ and $\Delta Z(\text{N2})$ vs $\Delta \log \Sigma_{\text{H}\alpha}$ relations are similar in the most massive and least massive galaxies. However, the $\Delta Z(\text{O3N2})$ vs $\Delta \log \Sigma_{\text{H}\alpha}$ relations in the most massive galaxies have positive slopes while those in the least massive galaxies have negative slopes. This is mostly due to the dependence of the $\Delta \log [\text{O III}]/\text{H}\beta$ vs $\Delta \log \Sigma_{\text{H}\alpha}$ relation on stellar mass. The dependence of DIG's impact on stellar mass means it is crucial to take care of DIG contamination when comparing the metallicity and metallicity gradient of galaxies of different masses using $Z(\text{O3N2})$.

In summary,

(i) Metallicities derived using N2O2 are optimal because they exhibit the smallest bias and error.

(ii) Metallicities derived using the O3N2 or N2S2H α (Dopita et al. 2016) for DIG can be significantly higher or lower than those for H II regions. Using O3N2 or N2S2H α to derive metallicity can bias the metallicity gradient by $\pm 0.05 R_e^{-1}$ for an individual galaxy if the contamination by DIG is not accounted for. For a large sample of galaxies (Sánchez et al. 2014, Ho et al. 2015), the bias for different galaxies may cancel out when deriving average metallicity gradients.

(iii) R_{23} derived metallicities for DIG are lower than those for H II regions due to a lower ionization parameter. Using R_{23} to derive metallicity will systematically bias the metallicity gradient by $\sim -0.1 R_e^{-1}$ because of DIG.

(iv) Using $\text{N2} = [\text{N II}]/\text{H}\alpha$ to derive metallicities will systematically bias the metallicity gradient by $\sim 0.05\text{--}0.1 \text{ dex } R_e^{-1}$, considering that DIG typically shows 0.2 dex higher $[\text{N II}]/\text{H}\alpha$.

(v) IZI works well for the H II region dominated regions, but fails for deriving metallicities of DIG, probably because IZI currently only contains H II region models.

5.8 Metallicity Gradients in the local universe and at high redshift

For the local universe, we usually have [O II], [O III], H β , [N II], H α , and [S II] to derive metallicities. Recent works find that the local star-forming galaxies exhibit a universal metallicity gradient of $-0.1 \text{ dex } R_e^{-1}$ within $2 R_e$ (e.g. Sánchez et al. 2014; Ho et al. 2015). These works employ O3N2 (Sánchez et al. 2014, Ho et al. 2015) and N2O2 (Ho et al. 2015) to derive the metallicity. Sánchez et al. (2014) isolated H II regions from DIG and used O3N2 to derive the metallicities using H II regions. The contamination of DIG is small. Ho et al. (2015) rejected the DIG-dominated spaxels by imposing $\text{S/N} > 3$, and $Z(\text{O3N2})$ and $Z(\text{N2O2})$ give consistent metallicities and metallicity gradients. The metallicity gradient at $R < 2R_e$ is robust against DIG contamination. Using MaNGA data and the robust metallicity estimator N2O2 to remeasure metallicities would be worthwhile to re-examine the metallicity gradient at large radius.




Figure 28. ΔZ vs $\Delta \log \Sigma_{\text{H}\alpha}$ relation at $[0.4R_e, 0.6R_e]$ for all galaxies. We normalize the ΔZ vs $\Delta \log \Sigma_{\text{H}\alpha}$ relation by subtracting the median Z and median $\log \Sigma_{\text{H}\alpha}$. All the spaxels in a galaxy are weighted by $\frac{1}{N}$, where N is the number of valid spaxels in the galaxy. A non-zero slope of the relation means metallicities are biased by DIG contribution, and the dispersion of this relation in y direction reflect how reliable a metallicity proxy is for an individual galaxy.

At high-redshift, we need a new calibration of strong line metallicity indicators because the physical properties are different in high redshift galaxies compared to local galaxies. High redshift star-forming galaxies are systematically offset towards higher $[\text{O III}]/\text{H}\beta$ on the BPT diagram relative to the local star-forming galaxy locus (Shapley et al. 2005; Erb et al. 2006; Liu et al. 2008; Brinchmann et al. 2008; Hainline et al. 2009; Wright et al. 2010; Trump et al. 2011; Kewley et al. 2013a,b; Steidel et al. 2014). The line ratio shifts of high-redshift star-forming galaxies are similar to DIG. Based on photoionization models, Steidel et al. (2014) concluded that the offset on the BPT diagram of the $z \sim 2.3$ star-forming galaxies is caused by a combination of harder stellar ionizing radiation field, higher ionization parameter, and higher N/O at a given O/H compared to most

local galaxies. A new calibration between N2 and O3N2 was obtained. A higher N/O abundance has been claimed to explain the offset in the BPT Diagram (Masters et al. 2014, Shapley et al. 2015, Sanders et al. 2016). There is debate whether the metallicity gradients at high redshift are flatter or steeper (Yuan et al. 2011; Swinbank et al. 2012; Jones et al. 2013, 2015; Leethochawalit et al. 2016; Wuyts et al. 2016). It takes time and resources to obtain all $[\text{O III}]$, $\text{H}\beta$, $[\text{N II}]$, $\text{H}\alpha$ for a galaxy at high redshift. In most cases, either $[\text{O III}]+\text{H}\beta$ or $[\text{N II}]+\text{H}\alpha$ is observed. The few lines available limit the metallicity estimator to either R_{23} or N2, two indicators that are impacted most by the presence of DIG. This makes the metallicity measurements at high- z vulnerable to contamination of DIG. However, we note current studies are unlikely to be seriously affected by DIG because in general




Figure 29. The distribution of metallicities vs $\Sigma(H\alpha)$ slope at three radii. The black line is for radius within $[0.4R_e, 0.6R_e]$, the blue line is for radius in $[0.8R_e, 1.0R_e]$, and the red line is for the radius in $[1.3R_e, 1.5R_e]$. The green dashed line has slope=0 line for reference. The metallicity is independent of surface brightness; the reason we are seeing a dependence of metallicity on $H\alpha$ surface brightness is due to contamination by DIG.

they are forming stars at an incredibly high rate and they are very well represented by pure H II region spectra. Future studies at high redshift probing low surface brightness limits (e.g., with JWST) may have to take into account the contribution of DIG. One needs to be cautious when making comparisons of high- z and low- z line ratios because DIG matters more at low z .

6 OTHER RELEVANT ISSUES

6.1 Line ratios vs $H\alpha$ surface brightness relation for an individual H II region

Relaño et al. (2010) studied the line ratios and $H\alpha$ surface brightness of NGC 595, one of the most luminous H II re-

gions in M33. The scale of NGC 595 is about 300pc. By comparing the $H\alpha$ surface brightness distribution shown in their Figure 3 and the $[N II]/H\alpha$, $[S II]/H\alpha$, and $[O III]/H\beta$ maps in Figure 6, we can see that the low surface brightness regions, which are located far away from the ionizing stars, have high $[N II]/H\alpha$, $[S II]/H\alpha$, and lower $[O III]/H\beta$. The line ratios vs. $H\alpha$ surface brightness relations are derived on a scale of hundreds of pc around an individual H II region. Madsen et al. (2006) saw the same trend in their observation of Milky Way. The resolution of the WHAM survey is 1 degree, which corresponds to 90 pc at 5 kpc. So the physical scale of WHAM is tens to hundred parsecs. Do these local relations on scales of hundreds of parsecs extend to kpc scale probed by MaNGA? MaNGA's spatial resolution of 1 kpc means each PSF may include tens or even




Figure 30. ΔZ vs $\Delta \log \Sigma_{\text{H}\alpha}$ relation at $[0.4R_e, 0.6R_e]$ for the one third most massive galaxies in our sample. We normalize the Z vs $\log \Sigma_{\text{H}\alpha}$ relation by subtracting the median Z and median $\log \Sigma_{\text{H}\alpha}$. All the spaxels in a galaxy are weighted by $\frac{1}{N}$, where N is the number of valid spaxels in the galaxy.

hundreds of H II regions. If the H α surface brightness vs line ratios relation is confined to 300pc scales, when smeared by a kpc-scale PSF, the line ratio variation will be smoothed out and disappear. In other words, if DIG is dominated by regions within 10pc of each H II region, then we would not see the line ratios vs $\Sigma_{\text{H}\alpha}$ relation after smoothing it with a kpc-sized PSF. *The smearing effect can produce a surface brightness gradient but can not produce an anti-correlation between surface brightness and line ratios.* The fact that we see a surface brightness vs line ratio relation on kpc-scales means that these relations for individual H II region can not be local, but extend to kpc scales. The DIG we see is domi-

nated by regions far away (kpc) from individual H II regions. In edge-on galaxies, we do see that DIG can extend to a few kpc above the galaxy plane (e.g. Rand 1996, 1997; Rossa & Dettmar 2000; Kehrig et al. 2012; Jones et al. 2016).

Then what does the line ratio vs H α surface brightness relation tell us? What we need to keep in mind is that we are looking at the average properties of the galaxy at our resolution. As discussed in detail in Appendix A, this relation is preserved for individual H II regions no matter what resolution we use to observe the galaxy. However, under poor resolution, many H II regions are mixed, and their relations are also mixed. We don't know how many H II regions there




Figure 31. ΔZ vs $\Delta \log \Sigma_{H\alpha}$ relation at $[0.4R_e, 0.6R_e]$ for the one third least massive galaxies in our sample. We normalize the Z vs $\log \Sigma_{H\alpha}$ relation by subtracting the median Z and median $\log \Sigma_{H\alpha}$. All the spaxels in a galaxy are weighted by $\frac{1}{N}$, where N is the number of valid spaxels in the galaxy.

are in one spaxel and what their line ratio vs $H\alpha$ surface brightness relations looks like. What we see is the average properties within 1 kpc, and how the line ratios and surface brightness change coherently on this scale. The curvature of the relation does tell us that an $H\text{II}$ region is not a simple Strömgen sphere but includes an extra component of DIG which has different line ratios. At 1 kpc scale, we can model the ISM as a combination of DIG and $H\text{II}$ regions with their respective line ratios. This relation also illustrates the feasibility of using $H\alpha$ surface brightness to roughly separate DIG and $H\text{II}$ regions.

6.2 Shedding Light on Composite Region and LI(N)ER

Any factors that change the line ratios can influence the positions on the diagnostic diagrams (e.g., Zhang et al. 2008; Pérez-Montero & Contini 2009). In Figure 15 and 16, we explored the distribution of each spaxel on the BPT diagram: $[\text{O III}]/H\beta$ vs $[\text{N II}]/H\alpha$ and $[\text{O III}]/H\beta$ vs $[\text{S II}]/H\alpha$ (Baldwin et al. 1981; Veilleux & Osterbrock 1987). In the $[\text{O III}]/H\beta$ vs $[\text{N II}]/H\alpha$ diagram, we plot the demarcations from Kewley et al. (2001) and Kauffmann et al. (2003) to identify the ionizing source properties. We color-code the

dots by $H\alpha$ surface brightness, and the dots in composite & LI(N)ER region have low $H\alpha$ surface brightness. As discussed in Section 4, a hardened O star spectrum filtered by ISM can not produce LI(N)ER-like emissions, while hot evolved stars like pAGB stars can produce the LI(N)ER-like line ratios. In Belfiore et al. (2016a), they classified galaxies that show LI(N)ER-like emission into two classes: cLIER and eLIER. cLIER shows LI(N)ER-like emission in the center, while eLIER shows extended LI(N)ER-like emission all over the galaxy. In some galaxies in our sample, the line ratios could extend to LI(N)ER regions on the BPT diagrams (See also Galarza et al. 1999; Kaplan et al. 2016). We suspect these two phenomenon may have a common origin. However, the galaxies that show eLIER emission and cLIER could have different stellar population and gas metallicity from the sample we study in this paper. Therefore a lot more work is needed to prove if they are really the same. We defer this to a future paper. The Shocked POststarburst Galaxy Survey (SPOGS, Alatalo et al. 2016a,b) explores a sample of galaxies selected from SDSS Data Release 7 that show LI(N)ER like emission line ratios and post-starburst spectral features. A similar phenomenon is found by Yan et al. (2006). After OB stars die, the spectrum of the galaxy is dominated by spectral features of A stars, characterizing the post-starburst signature. According to our grids, the decrease in OB stars and increase LI(N)ER-like emission line ratios are naturally linked through increasing photoionization by evolved stars. DIG, cLIER, and SPOGS are very similar to each other, and the study of their similarity and distinctions can greatly help us understand the ionized gas in different types of galaxies. From our figures, DIG dominated regions can mimic composite or even AGN spectra. If the selection of AGN is purely based on the BPT diagram, star-forming galaxies with significant DIG could be incorrectly classified as AGN. DIG and AGNs might be distinguished using the WHAN diagram (Cid Fernandes et al 2010, 2011) due to their different $EW(H\alpha)$ and $[N II]/H\alpha$ distributions. This will be explored further in the future. Our analysis demonstrates the importance of considering the presence of DIG while making optical line ratio classifications. Additionally, the analysis illustrates the complexity caused by DIG in interpreting the line ratio diagnostic diagrams. The so-called composite galaxies or LI(N)ERs could be coming from a variety of objects.

Shocks could also produce LI(N)ER-like line ratios. Shocked gas with velocities less than 500 km s^{-1} occupy the LI(N)ER part of the diagnostic diagrams (Farage et al. 2010; Rich et al. 2011), while shocked gas with velocities greater than 500 km s^{-1} fall in the Seyfert part of the diagram (Allen et al. 2008; Kewley et al. 2013). Shock model can produce the enhanced $[S II]/H\alpha$, $[N II]/H\alpha$, $[O II]/H\beta$, $[O I]/H\alpha$ and $[O III]/H\beta$ we see. Additionally, the temperature of shocked gas is elevated (Allen et al. 2008). To test if shocks are indeed a major ionization source for DIG, high spectral resolution spectra (~ 4000) are needed to separate different velocity components and constrain the kinematics. Our spectral resolution ($R \sim 2000$) limits our ability to test for shocks. We leave the question of the full nature of ionization sources of DIG for further studies.

6.3 How to minimize the impact of DIG

The best way to minimize the impact of DIG is to separate DIG and H II regions spatially. With high spatial resolution IFS data like CALIFA, it is possible to resolve individual H II regions to lower the contamination of DIG. For low spatial resolution data like MaNGA and SAMI, the individual spaxels are covering kpc scale regions, thus the emission is a mix of H II regions and DIG. The mixing also happens for MUSE data of high redshift galaxies. For these data, selecting high $\Sigma_{H\alpha}$ spaxels would mitigate the impact of DIG. As we have shown in this paper, the high $\Sigma_{H\alpha}$ regions have H II region line ratios. Besides, the metallicity gradient derived using only the high $\Sigma_{H\alpha}$ regions for different strong line method: $Z(N2O2)$, $Z(R_{23})$, $Z(O3N2)$, $Z(N2S2H\alpha)$ are consistent with each other. This means the high $\Sigma_{H\alpha}$ regions, even though contaminated by DIG due to beam smearing, are H II region dominated. A $\Sigma_{H\alpha}$ cut is a robust and easy way to reduce the impact of DIG. The exact value of $\Sigma_{H\alpha}$ cut depends on the spatial resolution. For our MaNGA survey, $\Sigma_{H\alpha} > 39 \text{ erg s}^{-1} \text{ kpc}^{-2}$ select reliable H II region dominated spaxels. An equivalent width (EW) cut is not recommended since EW depends on metallicity (e.g., Tresse et al. 1999). For low metallicity regions, for example in low metallicity galaxies or the outskirts of a galaxy, EW will be high due to the low metallicity. A EW cut suitable for the center of a galaxy will select DIG contaminated spaxels at the outskirts of this galaxy. So selecting high $\Sigma_{H\alpha}$ regions is a reliable and convenient way to minimize DIG contamination.

7 SUMMARY AND CONCLUSIONS

We selected a sample of 365 blue face-on galaxies from 1391 galaxies observed by MaNGA, and illustrated the impact of DIG on line ratios, interpretation of diagnostic diagrams, and metallicity measurements. We find that $H\alpha$ surface brightness is a good indicator to separate H II regions from DIG. DIG shows distinct properties as listed below:

- (i) $[S II]/H\alpha$, $[N II]/H\alpha$, $[O II]/H\beta$, and $[O I]/H\alpha$ are enhanced in DIG relative to H II regions.
- (ii) DIG has lower $[O III]/[O II]$, indicating lower ionization parameter. $[O III]/H\beta$ of DIG can be higher or lower than H II regions.
- (iii) On BPT diagrams, contamination by DIG moves H II regions towards composite or LI(N)ER-like regions. A harder ionizing spectrum is needed to explain DIG line ratios.
- (iv) Leaky H II region models only shift the line ratios slightly relative to H II region models, thus fail to explain composite/LI(N)ER line ratios displayed by DIG. Leaky H II region models cannot explain the $[O III]/H\beta$ but do pretty well for the other line ratios.
- (v) Our result favors ionization by evolved stars as a major ionization source for DIG with LI(N)ER-like emission.
- (vi) Metallicities derived using $N2O2 = [N II]/[O II]$ are optimal because they exhibit the smallest bias and scatter.
- (vii) Metallicities derived using the $O3N2 = ([O III]/H\beta)/([N II]/H\alpha)$ or $N2S2H\alpha = 8.77 + \log [N II]/[S II] + 0.264 \times \log [N II]/H\alpha$ (Dopita et al. 2016) for DIG can be significantly higher or lower than those for H II regions. Using O3N2 or N2S2H α to derive metallicities can bias the metallicity gradient by $\pm 0.05 \text{ dex } R_e^{-1}$ for

an individual galaxy if the contamination by DIG is not accounted for. R_{23} derived metallicities for DIG are lower than those for H II regions due to a lower ionization parameter. Using R_{23} to derive metallicities will systematically bias the metallicity gradient by ~ -0.1 dex R_e^{-1} because of DIG. Using $N2=[N\text{II}]/H\alpha$ to derive metallicities will systematically bias the metallicity gradient by ~ 0.05 - 0.1 dex R_e^{-1} , considering that DIG typically shows 0.2 dex higher $[N\text{II}]/H\alpha$.

(viii) The metallicities in high redshift galaxies are mostly derived using R_{23} or $N2$, rendering their metallicity and metallicity gradient measurements most vulnerable to the impact of DIG. Using $Z(N2S2H\alpha)$ for high redshift galaxies is more robust to prevent the contamination by DIG. For most of the recent high- z observations, the contamination by DIG is probably not severe because we only see high $\Sigma_{H\alpha}$ regions. When comparing the metallicities of high- z galaxies with local galaxies, one needs to use caution since DIG might impact metallicity measurements of local galaxies.

ACKNOWLEDGEMENTS

KB is supported by World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan and by JSPS KAKENHI Grant Number 15K17603. MAB acknowledges support from NSF AST-1517006. RM acknowledge support by the Science and Technology Facilities Council (STFC) and the ERC Advanced Grant 695671 "QUENCH?". C.A.T. acknowledges support from National Science Foundation of the United States grant no. 1412287 DB was supported by grant RSF 14-50-00043. AD acknowledges support from The Grainger Foundation. Funding for the Sloan Digital Sky Survey IV has been provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the Participating Institutions. SDSS- IV acknowledges support and resources from the Center for High-Performance Computing at the University of Utah. The SDSS web site is www.sdss.org. SDSS-IV is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS Collaboration including the Brazilian Participation Group, the Carnegie Institution for Science, Carnegie Mellon University, the Chilean Participation Group, the French Participation Group, Harvard-Smithsonian Center for Astrophysics, Instituto de Astrofísica de Canarias, The Johns Hopkins University, Kavli Institute for the Physics and Mathematics of the Universe (IPMU) / University of Tokyo, Lawrence Berkeley National Laboratory, Leibniz Institut für Astrophysik Potsdam (AIP), Max-Planck-Institut für Astronomie (MPIA Heidelberg), Max-Planck-Institut für Astrophysik (MPA Garching), Max-Planck-Institut für Extraterrestrische Physik (MPE), National Astronomical Observatory of China, New Mexico State University, New York University, University of Notre Dame, Observatório Nacional / MCTI, The Ohio State University, Pennsylvania State University, Shanghai Astronomical Observatory, United Kingdom Participation Group, Universidad Nacional Autónoma de México, University of Arizona, University of Colorado Boulder, University of Oxford, University of Portsmouth, University of Utah, University of Virginia, Uni-

versity of Washington, University of Wisconsin, Vanderbilt University, and Yale University.

REFERENCES

- Alatalo K., et al., 2016, arXiv, arXiv:1604.01122
 Alatalo K., et al., 2016, arXiv, arXiv:1601.05085
 Allen M. G., Groves B. A., Dopita M. A., Sutherland R. S., Kewley L. J., 2008, ApJS, 178, 20-55
 Alloin D., Collin-Souffrin S., Joly M., Vigroux L., 1979, A&A, 78, 200
 Andrews B. H., Martini P., 2013, ApJ, 765, 140
 Baldwin, J. A., Phillips, M. M., & Terlevich, R. 1981, PASP, 93, 5
 Baldwin, J. A., Ferland, G. J., Martin, P. G., et al. 1991, ApJ, 374, 580
 Belfiore, F., Maiolino, R., Bundy, K., et al. 2015, MNRAS, 449, 867
 Belfiore F., et al., 2016a, MNRAS, 461, 3111
 Belfiore F., et al., 2016b, arXiv, arXiv:1609.01737
 Binette L., Magris C. G., Stasińska G., Bruzual A. G., 1994, A&A, 292, 13
 Blanc, G. A., Heiderman, A., Gebhardt, K., Evans, N. J., II, & Adams, J. 2009, ApJ, 704, 842-862
 Blanc, G. A., Kewley, L., Vogt, F. P. A., & Dopita, M. A. 2015, ApJ, 798, 99
 Brinchmann, J., Pettini, M., & Charlot, S. 2008, MNRAS, 385, 769
 Bruzual, G., & Charlot, S. 2003, MNRAS, 344, 1000
 Bundy, K., Bershady, M. A., Law, D. R., et al. 2015, ApJ, 798, 7
 Collins, J. A., & Rand, R. J. 2001, ApJ, 551, 57
 Cid Fernandes R., Stasińska G., Schlickmann M. S., Mateus A., Vale Asari N., Schoenell W., Sodré L., 2010, MNRAS, 403, 1036
 Cid Fernandes R., Stasińska G., Mateus A., Vale Asari N., 2011, MNRAS, 413, 1687
 Denicoló, G., Terlevich, R., & Terlevich, E. 2002, MNRAS, 330, 69
 Dettmar R.-J., 1990, A&A, 232, L15
 Dopita, M. A., Kewley, L. J., Heisler, C. A., & Sutherland, R. S. 2000, ApJ, 542, 224
 Dopita, M. A., Fischera, J., Sutherland, R. S., et al. 2006, ApJs, 167, 177
 Dopita, M. A., Sutherland, R. S., Nicholls, D. C., Kewley, L. J., & Vogt, F. P. A. 2013, ApJs, 208, 10
 Dopita, M. A., Kewley, L. J., Sutherland, R. S., & Nicholls, D. C. 2016, ApJS, 361, 61
 Drory, N., MacDonald, N., Bershady, M. A., et al. 2015, AJ, 149, 77
 Erb, D. K., Shapley, A. E., Pettini, M., et al. 2006, ApJ, 644, 813
 Farage C. L., McGregor P. J., Dopita M. A., Bicknell G. V., 2010, ApJ, 724, 267
 Ferguson, A. M. N., Wyse, R. F. G., Gallagher, J. S., III, & Hunter, D. A. 1996, AJ, 111, 2265
 Ferland G. J., Korista K. T., Verner D. A., Ferguson J. W., Kingdon J. B., Verner E. M., 1998, PASP, 110, 761
 Fitzpatrick E. L., 1999, PASP, 111, 63
 Flores-Fajardo N., Morisset C., Stasińska G., Binette L., 2011, MNRAS, 415, 2182
 Galarza, V. C., Walterbos, R. A. M., & Braun, R. 1999, AJ, 118, 2775
 Garay, G., & Lizano, S. 1999, PASP, 111, 1049
 Garnett, D. R. 2002, ApJ, 581, 1019
 Giammanco C., Beckman J. E., Cedrés B., 2005, A&A, 438, 599
 Gomes J. M., et al., 2016, A&A, 588, A68
 Greenawalt, B., Walterbos, R. A. M., & Braun, R. 1997, ApJ, 483, 666

- Greenawalt, B., Walterbos, R. A. M., Thilker, D., & Hoopes, C. G. 1998, *ApJ*, 506, 135
- Gunn, J. E., Siegmund, W. A., Mannery, E. J., et al. 2006, *AJ*, 131, 2332
- Haffner, L. M., Reynolds, R. J., & Tufte, S. L. 1999, *ApJ*, 523, 223
- Haffner, L. M., Dettmar, R.-J., Beckman, J. E., et al. 2009, *Reviews of Modern Physics*, 81, 969
- Hainline, K. N., Shapley, A. E., Kornei, K. A., et al. 2009, *ApJ*, 701, 52
- Hausen, N. R., Reynolds, R. J., & Haffner, L. M. 2002, *AJ*, 124, 3336
- Ho, I.-T., Kudritzki, R.-P., Kewley, L. J., et al. 2015, *MNRAS*, 448, 2030
- Hoopes, C. G., Walterbos, R. A. M., & Greenawalt, B. E. 1996, *AJ*, 112, 1429
- Hoopes C. G., Walterbos R. A. M., Rand R. J., 1999, *ApJ*, 522, 669
- Hoopes, C. G., & Walterbos, R. A. M. 2003, *ApJ*, 586, 902
- Hunt, L. K., & Hirashita, H. 2009, *A&A*, 507, 1327
- Jones, T., Ellis, R. S., Richard, J., & Jullo, E. 2013, *ApJ*, 765, 48
- Jones, T., Wang, X., Schmidt, K. B., et al. 2015, *AJ*, 149, 107
- Jones A., et al., 2016, *arXiv*, arXiv:1612.03920
- Kaplan K. F., et al., 2016, *arXiv*, arXiv:1606.05713
- Kauffmann, G., Heckman, T. M., Tremonti, C., et al. 2003, *MNRAS*, 346, 1055
- Kennicutt, R. C., Jr. 1984, *ApJ*, 287, 116
- Kewley, L. J., Heisler, C. A., Dopita, M. A., & Lumsden, S. 2001, *ApJs*, 132, 37
- Kewley, L. J., & Dopita, M. A. 2002, *ApJs*, 142, 35
- Kewley, L. J., Groves, B., Kauffmann, G., & Heckman, T. 2006, *MNRAS*, 372, 961
- Kewley, L. J., & Ellison, S. L. 2008, *ApJ*, 681, 1183
- Kewley L. J., Dopita M. A., Leitherer C., Davé R., Yuan T., Allen M., Groves B., Sutherland R., 2013, *ApJ*, 774, 100
- Kim, K.-T., & Koo, B.-C. 2001, *ApJ*, 549, 979
- Kobulnicky, H. A., & Kewley, L. J. 2004, *ApJ*, 617, 240
- Kreckel K., Blanc G. A., Schinnerer E., Groves B., Adamo A., Hughes A., Meidt S., 2016, *arXiv*, arXiv:1603.08009
- Kehrig C., et al., 2012, *A&A*, 540, A11
- Lanz T., Hubeny I., 2003, *ApJS*, 146, 417
- Law, D. R., Yan, R., Bershady, M. A., et al. 2015, *AJ*, 150, 19
- Law D. R., et al., 2016, *AJ*, 152, 83
- Lee, H., Skillman, E. D., Cannon, J. M., et al. 2006, *ApJ*, 647, 970
- Leethochawalit N., Jones T. A., Ellis R. S., Stark D. P., Richard J., Zitrin A., Auger M., 2016, *ApJ*, 820, 84
- Lequeux, J., Peimbert, M., Rayo, J. F., Serrano, A., & Torres-Peimbert, S. 1979, *A&A*, 80, 155
- Levesque, E. M., Kewley, L. J., & Larson, K. L. 2010, *AJ*, 139, 712
- Li, C., Wang, E., Lin, L., et al. 2015, *ApJ*, 804, 125
- Liu, X., Shapley, A. E., Coil, A. L., Brinchmann, J., & Ma, C.-P. 2008, *ApJ*, 678, 758-779
- Madsen, G. J., Reynolds, R. J., & Haffner, L. M. 2006, *ApJ*, 652, 401
- Maiolino, R., Nagao, T., Grazian, A., et al. 2008, *A&A*, 488, 463
- Mannucci, F., Cresci, G., Maiolino, R., et al. 2009, *MNRAS*, 398, 1915
- Mannucci, F., Cresci, G., Maiolino, R., Marconi, A., & Gnerucci, A. 2010, *MNRAS*, 408, 2115
- Markwardt, C. B. 2009, *Astronomical Data Analysis Software and Systems XVIII*, 411, 251
- Mast D., et al., 2014, *A&A*, 561, A129
- Masters D., et al., 2014, *ApJ*, 785, 153
- McClure, R. D., & van den Bergh, S. 1968, *AJ*, 73, 313
- McGaugh, S. S. 1991, *ApJ*, 380, 140
- Mierkiewicz, E. J., Reynolds, R. J., Roesler, F. L., Harlander, J. M., & Jaehnig, K. P. 2006, *ApJl*, 650, L63
- Monnet, G. 1971, *A&A*, 12, 379
- Moran S. M., et al., 2012, *ApJ*, 745, 66
- Oey, M. S., Meurer, G. R., Yelda, S., et al. 2007, *ApJ*, 661, 801
- Osterbrock, D. E., Tran, H. D., & Veilleux, S. 1992, *ApJ*, 389, 305
- Otte, B. 2001, Ph.D. Thesis, 3233
- Otte, B., Reynolds, R. J., Gallagher, J. S., III, & Ferguson, A. M. N. 2001, *ApJ*, 560, 207
- Otte, B., Gallagher, J. S., III, & Reynolds, R. J. 2002, *ApJ*, 572, 823
- Pettini, M., & Pagel, B. E. J. 2004, *MNRAS*, 348, L59
- Pérez-Montero, E., & Contini, T. 2009, *MNRAS*, 398, 949
- Pérez-Montero E., et al., 2013, *A&A*, 549, A25
- Pérez-Montero E., 2014, *MNRAS*, 441, 2663
- Pérez-Montero E., et al., 2016, *arXiv*, arXiv:1608.04677
- Pilyugin, L. S. 2001, *A&A*, 369, 594
- Pilyugin, L. S., & Thuan, T. X. 2005, *ApJ*, 631, 231
- Rand R. J., Kulkarni S. R., Hester J. J., 1990, *ApJ*, 352, L1
- Rand, R. J. 1996, *ApJ*, 462, 712
- Rand R. J., 1997, *ApJ*, 474, 129
- Rand, R. J. 1998, *ApJ*, 501, 137
- Rand, R. J. 2000, *ApJl*, 537, L13
- Relaño, M., Monreal-Ibero, A., Vílchez, J. M., & Kennicutt, R. C. 2010, *MNRAS*, 402, 1635
- Reynolds R. J., 1984, *ApJ*, 282, 191
- Reynolds, R. J. 1985, *ApJ*, 294, 256
- Reynolds, R. J. 1985, *ApJl*, 298, L27
- Reynolds R. J., 1990, *ApJ*, 349, L17
- Reynolds R. J., 1991, *ApJ*, 372, L17
- Reynolds R. J., Cox D. P., 1992, *ApJ*, 400, L33
- Reynolds, R. J., Hausen, N. R., Tufte, S. L., & Haffner, L. M. 1998, *ApJl*, 494, L99
- Rich J. A., Kewley L. J., Dopita M. A., 2011, *ApJ*, 734, 87
- Rossa, J., & Dettmar, R.-J. 2003, *A&A*, 406, 493
- Rossa, J., & Dettmar, R.-J. 2003, *A&A*, 406, 505
- Rubin, R. H., Simpson, J. P., Haas, M. R., & Erickson, E. F. 1991, *ApJ*, 374, 564
- Rubin, R. H., Dufour, R. J., & Walter, D. K. 1993, *ApJ*, 413, 242
- Sánchez, S. F., Rosales-Ortega, F. F., Marino, R. A., et al. 2012, *A&A*, 546, A2
- Sánchez, S. F., Rosales-Ortega, F. F., Iglesias-Páramo, J., et al. 2014, *A&A*, 563, A49
- Sanders R. L., et al., 2016, *ApJ*, 816, 23
- Sarzi, M., Falcón-Barroso, J., Davies, R. L., et al. 2006, *MNRAS*, 366, 1151
- Shapley A. E., Coil, A. L., Ma, C.-P., & Bundy, K. 2005, *ApJ*, 635, 1006
- Shapley A. E., et al., 2015, *ApJ*, 801, 88
- Singh, R., van de Ven, G., Jahnke, K., et al. 2013, *A&A*, 558, A43
- Smee, S. A., Gunn, J. E., Uomoto, A., et al. 2013, *AJ*, 146, 32
- Stasińska, G., Vale Asari, N., Cid Fernandes, R., et al. 2008, *MNRAS*, 391, L29
- Steidel, C. C., Rudie, G. C., Strom, A. L., et al. 2014, *ApJ*, 795, 165
- Storchi-Bergmann, T., Calzetti, D., & Kinney, A. L. 1994, *ApJ*, 429, 572
- Swinbank, A. M., Sobral, D., Smail, I., et al. 2012, *MNRAS*, 426, 935
- Thilker, D. A., Walterbos, R. A. M., Braun, R., & Hoopes, C. G. 2002, *AJ*, 124, 3118
- Tremonti, C. A., Heckman, T. M., Kauffmann, G., et al. 2004, *ApJ*, 613, 898
- Tresse L., Maddox S., Loveday J., Singleton C., 1999, *MNRAS*, 310, 262
- Trump, J. R., Weiner, B. J., Scarlata, C., et al. 2011, *ApJ*, 743, 144

- Tüllmann R., Dettmar R.-J., 2000, *A&A*, 362, 119
- Vale Asari N., Stasińska G., Morisset C., Cid Fernandes R., 2016, *MNRAS*, 460, 1739
- van Zee, L., Salzer, J. J., & Haynes, M. P. 1998, *ApJl*, 497, L1
- Veilleux, S., & Osterbrock, D. E. 1987, *ApJs*, 63, 295
- Vincenzo F., Belfiore F., Maiolino R., Matteucci F., Ventura P., 2016, *MNRAS*, 458, 3466
- Voges, E. S., & Walterbos, R. A. M. 2006, *ApJl*, 644, L29
- Walterbos, R. A. M., & Braun, R. 1994, *ApJ*, 431, 156
- Wang J., Heckman T. M., Lehnert M. D., 1997, *ApJ*, 491, 114
- Wilkinson, D. M., Maraston, C., Thomas, D., et al. 2015, *MNRAS*, 449, 328
- Wright, S. A., Larkin, J. E., Graham, J. R., & Ma, C.-P. 2010, *ApJ*, 711, 1291
- Wuyts, E., Wisnioski, E., Fossati, M., et al. 2016, *arXiv:1603.01139*
- Yan, R., Newman, J. A., Faber, S. M., et al. 2006, *ApJ*, 648, 281
- Yan, R., & Blanton, M. R. 2012, *ApJ*, 747, 61
- Yan, R., Tremonti, C., Bershady, M. A., et al. 2016, *AJ*, 151, 8
- Yan R., et al., 2016, *arXiv*, *arXiv:1607.08613*
- Yates R. M., Kauffmann G., Guo Q., 2012, *MNRAS*, 422, 215
- Yuan, T.-T., Kewley, L. J., Swinbank, A. M., Richard, J., & Livermore, R. C. 2011, *ApJl*, 732, L14
- Zaritsky, D., Kennicutt, R. C., Jr., & Huchra, J. P. 1994, *ApJ*, 420, 87
- Zhang K., Wang T., Dong X., Lu H., 2008, *ApJ*, 685, L109
- Zurita, A., Beckman, J. E., Rozas, M., & Ryder, S. 2002, *A&A*, 386, 801
- Zurita, A., Rozas, M., & Beckman, J. E. 2000, *A&A*, 363, 9

¹Department of Physics and Astronomy, University of Kentucky, 505 Rose Street, Lexington, KY 40506, USA

²Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU, WPI), Todai Institutes for Advanced Study, the University of Tokyo, Kashiwa 277-8583, Japan

³Department of Astronomy, University of Wisconsin-Madison, 475 N. Charter Street, Madison, WI, 53706, USA

⁴Department of Astronomy, New Mexico State University, Las Cruces, NM 88003, USA

⁵Cavendish Laboratory, University of Cambridge, 19 J. J. Thomson Avenue, Cambridge CB3 0HE, United Kingdom

⁶Kavli Institute for Cosmology, University of Cambridge, Madingley Road, Cambridge, United Kingdom

⁷Institute of Cosmology & Gravitation, University of Portsmouth, Dennis Sciamia Building, Portsmouth, PO1 3FX, UK

⁸Department of Astronomy, University of Texas at Austin, Austin, TX 78712, USA

⁹Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, D-85748 Garching, Germany

¹⁰Instituto de Astronomía, Universidad Nacional Autónoma de México, A.P. 70-264, 04510, México, D.F., México

¹¹Apache Point Observatory and New Mexico State University, P.O. Box 59, Sunspot, NM, 88349-0059, USA

¹²Sternberg Astronomical Institute, Moscow State University, Moscow, Russia

¹³Unidad de Astronomía, Universidad de Antofagasta, Avenida Angamos 601, Antofagasta 1270300, Chile

¹⁴PITT PACC, Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, PA 15260, USA

¹⁵Department of Physics and Astronomy, University of Utah, 115 S. 1400 E., Salt Lake City, UT 84112, USA

¹⁶Tsinghua Center of Astrophysics & Department of

Physics, Tsinghua University, Beijing 100084, China

¹⁷Shanghai Astronomical Observatory, Nandan Road 80, Shanghai 200030, China

¹⁸Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

¹⁹Departamento de Física, Facultad de Ciencias, Universidad de La Serena, Cisternas 1200, La Serena, Chile

²⁰Universidade Federal do Rio Grande do Sul, IF, CP 15051, Porto Alegre 91501-970, RS, Brazil

²¹Laboratório Interinstitucional de e-Astronomia - LIneA, Rua Gal. José Cristino 77, Rio de Janeiro, RJ - 20921-400, Brazil

APPENDIX A: BEAM SMEARING EFFECT

Most of the H II region is smaller than 1 kpc, and the size depends on the density of the gas (e.g. Hunt & Hirashita 2009). Under MaNGA resolution, the mixing of H II region with DIG and other H II region is unavoidable. What's the effect of this mixing? We have seen in Section 3.1 that in one annulus, the line ratio vs surface brightness relation is tight. What would happen if we mix 2 regions on the line ratio vs surface brightness relation? From an analytical calculation, the mixed point is still on the same relation. However, if we mix two or more H II regions together, what would the line ratio vs H α surface brightness relation like? We show in Figure A1 for a simulation of observing our galaxies with poorer resolution. The left panel is the original $\Sigma(\text{H}\alpha)$ map, and the middle panel is the smeared $\Sigma(\text{H}\alpha)$ map. In the right panel, we show the comparison of original (black dots) and smeared (red dots) line ratio vs H α surface brightness relation. The H α flux map and [S II] flux map is convolved with a 2D gaussian with σ pixels listed in the middle panels of each row. We see several fact from this exercise:

(i) The beam smearing effect does not change the line ratio vs SB relation for one H II region, so the curve we derive using MaNGA data is likely to preserve when higher resolution data is obtained. In other word, this relation is intrinsic, reflecting the relationship between line ratio and surface brightness. Our resolution of ~ 1 kpc means one H α flux peak in our map is probably a mixture of several H II regions.

(ii) It is impossible to disentangle DIG from H II region from the curve alone.

(iii) The smearing makes the relation tighter and shorter.

(iv) When mixing two or more H II regions together, their line ratio vs SB relations also merge to an intermediate one.

The smearing process is an averaging process. We are witnessing more and more of the average properties of the H II region and DIG of the galaxy as we go to poorer resolution. The overall smeared line ratio vs H α surface brightness relation is tighter than before. So the tightness of the relation we see should be partly due to the beam smearing effect.




Figure A1. Left panel: the original $\Sigma(H\alpha)$ map, middle panel: the smeared $\Sigma(H\alpha)$ map. Right panel: the comparison of original (black dots) and smeared (red dots) line ratio vs $H\alpha$ surface brightness relation. The $H\alpha$ flux map and $[N II]$ flux map is convolved with a 2D gaussian with σ pixels listed in the middle panels of each row.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.