

# Complexity in the light curves and spectra of slow-evolving superluminous supernovae

C. Inserra<sup>1,2\*</sup>, M. Nicholl<sup>3</sup>, T.-W. Chen<sup>4</sup>, A. Jerkstrand<sup>5</sup>, S. J. Smartt<sup>1</sup>, T. Krühler<sup>4</sup>, J. P. Anderson<sup>6</sup>, C. Baltay<sup>7</sup>, M. Della Valle<sup>8</sup>, M. Fraser<sup>9</sup>, A. Gal-Yam<sup>10</sup>, L. Galbany<sup>11</sup>, E. Kankare<sup>1</sup>, K. Maguire<sup>1</sup>, D. Rabinowitz<sup>7</sup>, K. Smith<sup>1</sup>, S. Valenti<sup>12</sup> and D. R. Young<sup>1</sup>

<sup>1</sup>*Astrophysics Research Centre, School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, UK*

<sup>2</sup>*Department of Physics & Astronomy, University of Southampton, Southampton, Hampshire, SO17 1BJ, UK*

<sup>3</sup>*Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, Massachusetts 02138, USA*

<sup>4</sup>*Max-Planck-Institut für Extraterrestrische Physik, Giessenbachstraße 1, 85748, Garching, Germany*

<sup>5</sup>*Max Planck Institute for Astrophysics, Garching, Karl-Schwarzschild-Str. 1, Postfach 1317, D-85741 Garching, Germany*

<sup>6</sup>*European Southern Observatory, Alonso de Córdova 3107, Vitacura, Casilla 190001, Santiago, Chile*

<sup>7</sup>*Physics Department, Yale university, New Haven, CT 06511, USA*

<sup>8</sup>*Capodimonte Observatory, Salita Moiariello 16, I-80131, Napoli, Italy*

<sup>9</sup>*School of Physics, O'Brien Centre for Science North, University College Dublin, Belfield, Dublin 4, Ireland*

<sup>10</sup>*Benoziyo Center for Astrophysics, Weizmann Institute of Science, 76100 Rehovot, Israel*

<sup>11</sup>*Physics and Astronomy Department, University of Pittsburgh, Pittsburgh, PA 15260, USA*

<sup>12</sup>*Dept. of Physics, University of California Davis, 1 Shields Ave, Davis, CA 95616, USA*

Received.....; accepted.....

## ABSTRACT

A small group of the newly discovered superluminous supernovae show broad and slowly evolving light curves. Here we present extensive observational data for the slow-evolving superluminous supernova LSQ14an, which brings this group of transients to four in total in the low redshift Universe ( $z < 0.2$ ; SN 2007bi, PTF12dam, SN 2015bn). We particularly focus on the optical and near-infrared evolution during the period from 50 days up to 400 days from peak, showing that they are all fairly similar in their light curve and spectral evolution. LSQ14an shows broad, blue-shifted [O III]  $\lambda\lambda 4959, 5007$  lines, as well as a blue-shifted [O II]  $\lambda\lambda 7320, 7330$  and [Ca II]  $\lambda\lambda 7291, 7323$ . Furthermore, the sample of these four objects shows common features. Semi-forbidden and forbidden emission lines appear surprisingly early at 50–70 days and remain visible with almost no variation up to 400 days. The spectra remain blue out to 400 days. There are small, but discernible light curve fluctuations in all of them. The light curve of each shows a faster decline than  $^{56}\text{Co}$  after 150 days and it further steepens after 300 days. We also expand our analysis presenting X-ray limits for LSQ14an and SN 2015bn and discuss their diagnostic power. These features are quite distinct from the faster evolving superluminous supernovae and are not easily explained in terms of only a variation in ejecta mass. While a central engine is still the most likely luminosity source, it appears that the ejecta structure is complex, with multiple emitting zones and *at least* some interaction between the expanding ejecta and surrounding material.

**Key words:** supernovae: general – supernovae: individual: LSQ14an, SN2015bn, PTF12dam, SN2007bi – X-rays: general – stars: mass loss

## 1 INTRODUCTION

A new class of intrinsically bright supernovae, labelled superluminous supernovae (SLSNe; Quimby et al. 2011; Gal-Yam 2012), was unveiled during last decade. They were orig-

inally identified as those supernovae (SNe) showing  $M \sim -21$  mag (e.g. Smith et al. 2007; Pastorello et al. 2010; Quimby et al. 2011; Chomiuk et al. 2011; Gal-Yam 2012; Inserra et al. 2013; Howell et al. 2013) and divided in three different types (see the early review of Gal-Yam 2012). They now tend to be separated and classified based on their spectrophotometric behaviour rather than a simple magnitude threshold, (e.g.

\* E-mail: c.inserra@qub.ac.uk (CI)

Papadopoulos et al. 2015; Inserra et al. 2016b; Prajs et al. 2017; Lunnan et al. 2016). Those exhibiting hydrogen-free ejecta are grouped together as SLSNe I, while those showing hydrogen as SLSNe II.

SLSNe I, although intrinsically rare (Quimby et al. 2013; McCrum et al. 2015; Prajs et al. 2017) are the most commonly studied and usually found in dwarf, metal poor galaxies (e.g. Lunnan et al. 2014; Leloudas et al. 2015a; Angus et al. 2016; Perley et al. 2016; Chen et al. 2016). At about 30 days after peak, these SNe tend to show strong resemblances to normal or broad-lined type Ic SNe at peak luminosity (e.g. Pastorello et al. 2010; Liu & Modjaz 2016) and thus are also usually referred to as SLSNe Ic (e.g. Inserra et al. 2013; Nicholl et al. 2013). Their light curves show significant differences in width and temporal evolution with the rise times being correlated with the declines rates (Nicholl et al. 2015b). Whether or not there are two distinct groups, and two different physical mechanisms for the fast-evolving (e.g. SN 2010gx) and slow-evolving (e.g. SN 2007bi) explosions remains to be seen.

In addition, thanks to modern imaging surveys we have been able to differentiate between two groups of hydrogen rich SLSNe, which are rarer by volume than SLSNe I. The first is that of strongly interacting SNe such as SN 2006gy (Smith et al. 2007; Ofek et al. 2007; Agnoletto et al. 2009) of which the enormous luminosity is undeniably powered by the interaction of supernova ejecta with dense circumstellar (CSM) shells and hence usually referred to as SLSNe IIn to highlight the similarity with normal type IIn. Whereas, the second group is that of bright, broad lined H-rich SNe labelled SLSNe II, such as SN 2008es (Gezari et al. 2009; Miller et al. 2009), which have shallow but persistent and broad Balmer lines (see Inserra et al. 2016b, for a review). They show spectroscopic evolution that is similar, in some ways to type II-L supernovae, but the evolution happens on a longer timescale.

The mechanisms responsible to power the luminosities of SLSNe I & II is still debated. However, the most favoured is that of a magnetar as a central engine, depositing additional energy into the supernova ejecta (Woosley 2010; Kasen & Bildsten 2010; Dessart et al. 2012). The popularity of this scenario is due to its ability to reproduce a variety of observables for all the types (e.g. Inserra et al. 2013; Nicholl et al. 2013; Inserra et al. 2016b). However, alternative scenarios such as the accretion onto a central black hole (Dexter & Kasen 2013), the interaction with a dense circumstellar medium (e.g. Chevalier & Irwin 2011) or between two massive shells (pulsational pair instability, Woosley et al. 2007), and a pair instability explosion (e.g. Kozyreva & Blinnikov 2015) are still feasible alternatives.

Due to their intrinsic brightness, SLSNe I are emerging as high redshift probes and might be a new class of distance indicators (Inserra & Smartt 2014; Papadopoulos et al. 2015; Scovaccicchi et al. 2016). To use them with the new generation of telescopes such as the Large Synoptic Survey Telescope (LSST) or Euclid we need novel approaches to better understand their progenitor scenario, reveal the mechanism powering the light curve and investigate on the nature of their differences. These new approaches include analyses of large dataset (e.g. Nicholl et al. 2015b), spectropolarimetric studies (Leloudas et al. 2015b; Inserra et al. 2016b,c) or de-

voted spectroscopic modelling of photospheric (Mazzali et al. 2016) and nebular spectroscopy (Jerkstrand et al. 2017).

The slowly evolving group of SLSNe I, of which the first to be discovered was SN 2007bi (Gal-Yam et al. 2009; Young et al. 2010), is small in number but of particular interest because the light curves decline at a rate consistent with powering by  $^{56}\text{Co}$  during first 100-200 days after maximum. A very large amount of explosively produced  $^{56}\text{Ni}$  is required to power the peak through  $^{56}\text{Co}$  decay (to produce stable  $^{56}\text{Fe}$ ). With estimates in the region of  $4-6M_{\odot}$  this led Gal-Yam et al. (2009) to propose a pair-instability explosion origin for SN 2007bi (in combination with spectral analysis) from a very massive progenitor star above  $M_{\text{ZAMS}} \simeq 150M_{\odot}$ . Hence any new objects with similar light curve and spectra behaviour are of interest to study over long durations.


Here we present an extensive dataset for LSQ14an, a slow-evolving SLSNe I discovered by the La Silla QUEST survey and studied by the Public ESO Spectroscopic Survey for Transient Objects (PESSTO<sup>1</sup>, Smartt et al. 2015). To draw a meaningful comparison, we compare it with a sample of slow-evolving SLSNe I at low redshift ( $z < 0.2$ ). There are three objects that share very similar characteristics that make up this comparison - SN 2007bi (Gal-Yam et al. 2009; Young et al. 2010), PTF12dam (Nicholl et al. 2013; Chen et al. 2015; Vreeswijk et al. 2017), SN 2015bn (Nicholl et al. 2016a,b). These objects all have rest-frame spectra coverage up to at least 8000 Å and late time spectroscopy and photometry up to 400 days. We will also discuss the higher redshift, and perhaps more extreme PS1-14bj (Lunnan et al. 2016). These comparisons allow us to further investigate the explosion mechanism and the progenitors of these transients, as well as the role of interaction with a circumstellar medium.

## 2 LSQ14AN OBSERVATIONS AND DATA REDUCTION

LSQ14an was discovered on UT 2014 January 02.3 by the La Silla-Quest (LSQ) survey (Baltay et al. 2013). The LSQ camera used a broad composite filter covering the SDSS  $g$  and  $r$  bands, from 4000 Å to 7000 Å and the pipeline initially reported a discovery magnitude of  $m \sim 19.1$  mag in this filter. The object was promptly classified within 24hrs by PESSTO as a SLSN roughly 50 days after peak brightness at  $z \sim 0.16$  with similarities to the slow-evolving PTF12dam and SN 2007bi (Leget et al. 2014). A more careful photometric analysis and calibration of the images from the QUEST camera to SDSS  $r$ -band, resulted in an initial magnitude of  $r = 18.60 \pm 0.08$  mag, as reported in Table A2 (see Section 5 for details of the photometry).

Optical spectrophotometric follow-up was obtained with the New Technology Telescope (NTT)+EFOSC2 as part of PESSTO and supplemented with other facilities. Photometry with EFOSC2 was obtained using  $BVR+gri$  filters and was augmented with observations from the Liverpool Telescope (LT)+IO:O in  $gri$  filters. The EFOSC2 images were reduced (trimmed, bias subtracted and flat-fielded) using the PESSTO pipeline (Smartt et al. 2015),

<sup>1</sup> www.pessto.org



**Figure 1.** The left panel shows the  $r$ -band EFOSC2 image of LSQ14an taken on 2014 Jan 31. The right panel shows the colour composite of the host galaxy from the Pan-STARRS1 survey (constructed from  $g_{P1}r_{P1}z_{P1}$  filters). The two images are astrometrically aligned, with the centroid position of LSQ14an marked with the white cross hairs on the right. This illustrates that the SN is coincident with the dwarf galaxy host and that spectra of LSQ14an will be contaminated by the host galaxy flux since it is unresolved. The image quality of the EFOSC2 image is  $0.9''$  FWHM and the PS1  $r_{P1}$  image is  $1.2''$ . North is up and east to the left.

while the LT images were reduced automatically by the LT instrument-specific pipeline. One epoch of near-infrared (NIR) imaging was obtained with the NTT and SOFI of Isaac (SOFI) instrument. These data were also reduced using the PESSTO pipeline. To remove the contribution of the host galaxy to the SN photometry (see Section 3 for a data on the host) we applied a template subtraction technique (with the `HOTPANTS`<sup>2</sup> package based on the algorithm presented in Alard 2000). We used EFOSC2  $BVR$  images taken on 2015 April 17 (MJD=57129, or an estimated +460d after peak), and the PS1  $gri$  images taken before 2013 June 24 (see Section 3) as the reference templates. On the other hand, we did not use any  $JHK$  templates for our single epoch of observations since the contamination is on the order of a few per cent and has no significant effect on our analysis. Since instruments with different passbands were used for the follow-up of LSQ14an we applied a passband correction (P-correction, Inserra et al. 2016b), which is similar to the S-correction (Stritzinger et al. 2002; Pignata et al. 2004), and allows photometry to be standardised to a common system, which in our case is SDSS AB mags for  $gri$  and Bessell Vega mags for  $BVR$ . This procedure takes into account the filter transmission function and the quantum efficiency of the detector, as well as the intrinsic spectrum of LSQ14an, but does *not* include the reflectance or transmission of other optical components, as this is relatively flat across the optical range (see Inserra et al. 2016b, for further details). Optical and NIR photometry are reported in Tables A2 & A3 together with the coordinates and magnitudes of the sequence stars used (see Table A4).

Optical and near-infrared spectra were taken during the PESSTO follow-up campaign and were supplemented with two spectra from X-Shooter on the ESO Very Large Telescope (VLT). The log of spectral observations (eight separate epochs), with resolutions and wavelength ranges is given

in Table A1. Two further spectra were taken with the VLT at later epochs of (+365 and +410d) when the SN was in the nebular phase and are analysed in detail in Jerkstrand et al. (2017). The PESSTO spectra were reduced in standard fashion as described in Smartt et al. (2015). The X-Shooter spectra were reduced in the same manner as described for the nebular spectra in Jerkstrand et al. (2017). We used the custom made pipeline described in Kriehler et al. (2015) which takes the ESO pipeline produced 2D spectral products (Modigliani et al. 2010) and uses optimal extraction with a Moffat profile fit. This pipeline produced flux calibrated spectra with rebinned dispersions of  $0.4 \text{ \AA pix}^{-1}$  in the UVB+VIS arms and  $0.6 \text{ \AA pix}^{-1}$  in the NIR arm. The PESSTO spectra taken up to April 2014 are available from the public data release Spectroscopic Survey Data Release 2 (SSDR2) at ESO. Steps to retrieve the data from the ESO Science Archive Facility are available on the PESSTO website. The final EFOSC2 spectrum on 21 August 2014 will be released in the upcoming SSDR3. All spectra will also be available on WISeREP<sup>3</sup> (Yaron & Gal-Yam 2012).

The *Swift* satellite observed LSQ14an with the X-ray Telescope (XRT) on four epochs (PIs: Margutti & Inserra) listed in Table A5. For each XRT observation of LSQ14an we extracted images over the 0.3-10 keV band and all data were reduced using the HEASARC<sup>4</sup> software package. We measured the observed counts in each image in apertures of 5 pixel radius ( $11.8''$ ), while the background in a large region (approximately 100 pixel radius), at a location free of bright X-ray sources. We then corrected this flux for the PSF contained within the aperture radius. *Swift* also observed another nearby, slowly evolving SLSN, namely SN 2015bn. The monitoring of this was described in Nicholl et al. (2016a,b) and analysis of the nebular spectra was presented alongside LSQ14an in Jerkstrand et al. (2017). For comparison, we retrieved the *Swift* X-ray data (5 epochs were taken, PI: Margutti) and applied similar reductions and measurements as for LSQ14an. We additionally combined all the available observations in a single stacked image in order to have a deeper measurement. All measurements for the two SLSNe are listed in Table A5.

### 3 HOST GALAXY, LOCATION AND REDSHIFT

The host galaxy was detected in the Pan-STARRS1  $3\pi$  survey (it is outside the SDSS footprint) in all five filters  $grizy_{P1}$  (see Tonry et al. 2012, for a description of the filter set). Images in each of the filters were taken from the PS1 Processing Version 2 (PS1.PV2) stacks. All the individual images which make up these stacked images were taken before MJD 56467 (2013 June 24). This is 4 months before the estimated peak date of late October 2013 and hence there is unlikely to be any SN flux in the images. Magnitude measurements of the host were carried out using aperture photometry with an aperture of  $2''$ , and the zero points were determined with 15-18 PS1 reference stars (Schlafly et al.

<sup>3</sup> <http://wiserep.weizmann.ac.il/home>

<sup>4</sup> NASA High Energy Astrophysics Science Archive Research Center.

<sup>2</sup> <http://www.astro.washington.edu/users/becker/hotpants.html>


2012; Magnier et al. 2013) in the field resulting in the magnitudes listed in Table 1

As discussed in Jerkstrand et al. (2017), the position of LSQ14an was measured at  $\alpha = 12^{\text{h}}53^{\text{m}}47^{\text{s}}.82$ ,  $\delta = -29^{\circ}31'27''.6$  (J2000) in the astrometrically calibrated EFOSC2 images (with SExtractor). In comparison, the centroid of the host galaxy, PSO J193.4492-29.5243, was measured at  $\alpha = 12^{\text{h}}53^{\text{m}}47^{\text{s}}.81$ ,  $\delta = -29^{\circ}31'27''.4$  (J2000). The difference between the two is  $0.2''$ , which illustrates that LSQ14an is effectively coincident with the dwarf galaxy host (as illustrated in Fig 1). Ultraviolet (UV) photometry of the host was obtained with the *Swift* satellite + UltraViolet and Optical Telescope (UVOT) (programme ID 33205; P.I. Margutti). The images were co-added before aperture magnitudes were measured following the prescription of Poole et al. (2008). A  $5''$  aperture was used to maximise the signal-to-noise ratio (S/N). The UV magnitudes were determined by averaging the measurements of MJD 56740.52 (2014 July 03) and MJD 56998.41 (2014 December 07) when the SN flux was already dimmer than that of the host. We measured  $uvw2 = 21.94 \pm 0.14$ ,  $uvm2 = 21.72 \pm 0.15$ ,  $uvw1 = 21.44 \pm 0.16$  mag in the AB mag system.

The host galaxy emission lines from LSQ14an are strong and conspicuously narrower than the SN emission features and they do not evolve in our spectral series. We used  $\text{H}\alpha$  and other galaxy lines to refine the host redshift to  $z = 0.1637 \pm 0.0001$ , equivalent to a luminosity distance of  $D_L = 766$  Mpc assuming a fiducial cosmology of  $H_0 = 72$   $\text{km s}^{-1}$ ,  $\Omega_M = 0.27$  and  $\Omega_\Lambda = 0.73$ . The foreground Galactic reddening toward LSQ14an is  $E_G(B-V) = 0.07$  mag from the Schlafly & Finkbeiner (2011) dust maps. The available SN spectra do not show Na I D lines from the host galaxy hence we adopt a total reddening along the line of sight towards LSQ14an of  $E_{\text{tot}}(B-V) = 0.07$  mag.

To estimate physical diameters of the host galaxy we assumed  $(\text{galaxy observed FWHM})^2 = (\text{observed PSF FWHM})^2 + (\text{intrinsic galaxy FWHM})^2$ . The measured FWHM of the galaxy and the PSF were  $1.65''$  and  $1.21''$ , respectively (with the stellar PSF FWHM averaged over 16 reference stars in  $r$ -band images, located within a  $2'$  radius of the host in the images). The angular size distance at this redshift ( $D_A = 565.6$  Mpc) implies a physical diameter of the galaxy of  $\sim 3.09$  kpc.

From the VLT+X-Shooter spectrum taken on 2014 June 21 (see Table A1) we measured the observed emission line fluxes of PSO J193.4492-29.5243 (see Figure 2, top panel). Emission line measurements were made after fitting a low-order polynomial function to the continuum and subtracting off this contribution. This continuum flux is composed of the underlying stellar population of the host and the SN flux. No significant stellar absorption was seen in the spectrum, so we did not use stellar population model subtraction method. We fitted Gaussian line profiles of those emission lines using the QUB custom built PROCSPPEC environment within IDL. We let the full width at half-maximum (FWHM) vary for the strong lines and fixed the FWHM for weak lines adopting the FWHM of nearby stronger, single transitions. The equivalent width (EW) of lines were determined after spectrum normalisation. However, we caution that the normalised continuum has SN flux contamination and therefore the EW of each line is a minimum value (rather than a real line strength compared to the galaxy continuum).



**Figure 2.** *Top:* Spectrum of the host of LSQ14an taken by VLT+X-Shooter on 2014 June 21. The spectrum in the merged data from the UVB, VIS and NIR arms, corrected to rest-frame. Host lines are labelled in the spectrum. *Bottom:* The best-fit SED of the host of LSQ14an from MAGPHYS (da Cunha, Charlot, & Elbaz 2008) shown in black line. The colour circles show photometry of LSQ14an host in the order of the Swift  $uvw2$ ,  $uvm2$ ,  $uvw1$  and Pan-STARRS1  $grizy$  filters.

Uncertainties were calculated from the line profile, EW and rms of the continuum, following the equation from Gonzalez-Delgado et al. (1994). The observed flux measurement and related parameters are listed in Table A6. The lines are identified as those commonly seen in star-forming galaxies, such as Green Peas (e.g. Amorín et al. 2012) and other SLSN hosts (e.g. see Chen et al. 2015, for a comprehensive line list for the host of PTF12dam).

We used the “direct”  $T_e$  method, following the prescription of Nicholls et al. (2013) and assuming Maxwell-Boltzmann distribution, based on the detection of  $[\text{O III}] \lambda 4363$  line and found  $T_e([\text{O III}]) = 13700 \pm 1200$  K. We can then infer a  $T([\text{O II}]) = 13100 \pm 800$  K, resulting in an oxygen abundance of  $12 + \log(\text{O}/\text{H}) = 8.04 \pm 0.08$ , equivalent to  $0.22 Z_\odot$  considering a solar value of 8.69 (Asplund et al. 2009). We also checked the electron temperature at the lower-ionisation zone using the ratio of the nebular lines  $[\text{O II}] \lambda \lambda 7320, 7330$  and thus retrieved  $T_e([\text{O II}])$  of  $\sim 7100$  K which is much lower than the inferred  $T([\text{O II}])$ . We believe

**Table 1.** LSQ14an host galaxy observed magnitudes (AB system) and properties derived from the spectral energy distribution and late time spectra (see Appendix 3 for further information).


$uvw2$ (mag)	$21.94 \pm 0.14$
$wvm2$ (mag)	$21.72 \pm 0.15$
$uvw1$ (mag)	$21.44 \pm 0.16$
$g_{P1}$ (mag)	$20.93 \pm 0.11$
$r_{P1}$ (mag)	$20.50 \pm 0.12$
$i_{P1}$ (mag)	$20.56 \pm 0.10$
$z_{P1}$ (mag)	$20.42 \pm 0.14$
$y_{P1}$ (mag)	$20.43 \pm 0.19$
Physical diameter (kpc)	3.09
H $\alpha$ Luminosity (erg s $^{-1}$ )	$2.41 \times 10^{41}$
SFR ( $M_{\odot}$ yr $^{-1}$ )	1.19
Stellar mass (log( $M/M_{\odot}$ ))	$8.6^{+0.2}_{-0.1}$
sSFR (Gyr $^{-1}$ )	2.98
$12 + \log(O/H)$ ( $T_e$ )	$8.04 \pm 0.08$
$12 + \log(O/H)$ (PP04 N2)	$8.02 \pm 0.03$
$12 + \log(O/H)$ (PP04 O3N2)	$7.96 \pm 0.01$
log (S/O)	$-1.94 \pm 0.15$
log (N/O) ( $T_e$ )	$-1.74 \pm 0.18$

the discrepancy is due to the low S/N of the [O II] lines with a large SN flux contamination and hence we do not use the  $T[O II]$ . We used the open-source PYTHON code PYMCZ from Bianco et al. (2016) to alternatively calculate metallicity of the host of LSQ14an with different strong line calibrations, and we listed them as follows:  $12 + \log(O/H) = 8.02 \pm 0.02$  from the N2 method (Pettini & Pagel 2004) and  $12 + \log(O/H) = 7.96 \pm 0.01$  of the O3N2 method (Pettini & Pagel 2004). The R23 value of 0.9 is at an insensitive regime, so we do not use this scale here.

We employed the MAGPHYS stellar population model program of da Cunha, Charlot, & Elbaz (2008) to estimate the stellar mass from the observed photometry (after foreground extinction correction) of the host galaxy. MAGPHYS gives the best-fit model spectrum ( $\chi^2_{red} = 0.316$ ) and the total stellar mass of the hosts (see Figure 2). The median and the  $1\sigma$  range of the stellar mass of LSQ14an host are  $4.0^{+2.5}_{-1.1} \times 10^8 M_{\odot}$ , which is consistent with the measurements of Schulze et al. (2016) for the same host, and that retrieved in SLSNe hosts (e.g. Lunnan et al. 2014; Leloudas et al. 2015a; Perley et al. 2016).

We converted the H $\alpha$  luminosity of  $2.41 \times 10^{41}$  erg s $^{-1}$  to calculate the star-formation rate (SFR) using the conversion of Kennicutt (1998) and a Chabrier IMF<sup>5</sup>. We found a SFR of LSQ14an host SFR =  $1.19 M_{\odot}$  yr $^{-1}$ , which is higher than the average, but still consistent with other slow-evolving SLSNe I (e.g. Chen et al. 2015; Leloudas et al. 2015a). In this case our estimate is somewhat lower than that reported by Schulze et al. (2016). Considering the stellar mass previously evaluated, we determined the specific SFR (sSFR) of 2.98 Gyr $^{-1}$  for the host of LSQ14an. A summary of LSQ14an host galaxy properties is reported in Table 1.

<sup>5</sup> The tool employed to evaluate the SFR uses a Chabier IMF, while Kennicutt (1998) a Salpeter IMF. We took that in account dividing the SFR by a factor of 1.6.




**Figure 3.** Top: comparison of LSQ14an classification spectrum with those of other slow-evolving SLSNe I (SN 2007bi, PTF12dam, SN 2015bn; Gal-Yam et al. 2009; Young et al. 2010; Nicholl et al. 2013, 2016a) around 50 days post maximum suggesting LSQ14an was discovered at  $\sim 55$  days from maximum light. Bottom: same LSQ14an spectrum of the top panel compared with fast-evolving SLSNe I (SN 2011ke, SN 2012il and CSS121015; Inserra et al. 2013; Benetti et al. 2014) in order to highlight the differences. The grey area identify the [O III] lines observed in emission in LSQ14an spectra, while the cyan in the bottom panel identifies the region of the [Ca II]  $\lambda\lambda 7291, 7323$ .

## 4 SPECTRA

The first spectra taken of LSQ14an by PESTO (Leget et al. 2014) already suggested an epoch well after maximum brightness. The absence of the characteristic broad O II absorption lines observed in SLSNe, at and before the peak epoch, together with a temperature of  $\sim 8000$  K - derived from the blackbody fit to the continuum of our spectra - confirm the initial classification phase. In order to secure the phase with more precision and estimate the date of the peak of the light curve, we compared our classification spectrum with other slow-evolving SLSNe I. As highlighted in the top panel of Figure 3, the LSQ14an spectrum taken on the 2nd January 2014 is very similar to those of the three other objects with broad light curves: SN 2007bi, PTF12dam and SN 2015bn at 50 to 60d after peak.

At this phase, the emission lines that are developing and becoming pronounced are similar to those that have quantitatively been identified and modelled by Jerkstrand et al. (2017). The Mg I]  $\lambda 4571$  line is strong, and this usually begins to appear about  $\sim 35$  days post peak (Inserra et al. 2013; Nicholl et al. 2016a). The region between  $5000 \text{ \AA}$  and  $5800 \text{ \AA}$  shows a broad emission feature which is likely



**Figure 4.** Spectral evolution of LSQ14an. The phase of each spectra is relative to the assumed peak epoch. The  $\oplus$  symbol marks the positions of the strongest telluric absorptions. The most prominent features are labelled, together with the position (dashed magenta line) of the NIR Ca II, which is not observed.

a blend of Mg I  $\lambda 5180$ , [Fe II]  $\lambda 5250$  and [O I]  $\lambda 5577$ . There is a distinct lack of absorption that could be attributed to Si II  $\lambda 6355$  and the forbidden doublet of [O I]  $\lambda\lambda 6300, 6363$  is beginning to emerge in emission (see Figure 4 for line identification). Finally, the [Ca II]  $\lambda\lambda 7291, 7323$  line strength and the equivalent width of O I  $\lambda 7774$  both support an epoch of the spectrum of +50-55 days after maximum in the rest-frame<sup>6</sup>. As a consequence, assuming a phase of +55d for the first spectrum, we estimate the peak epoch to have been MJD  $56596.3 \pm 5.0$  (around 2013 Oct 31). This date is consistent with our analysis of the light curve evolution (see Section 5).

For comparison, we also showed the earliest PESSTO spectrum of LSQ14an together with faster evolving SLSNe

(bottom panel of Figure 3). SN 2011ke and SN 2012il decline much faster than the other three comparison objects and are similar to SN 2010gx in their overall evolution (see Inserra et al. 2013, for a discussion of these two objects). CSS121015 is a peculiar SLSN I which shows weak signs of interaction with a hydrogen CSM that fade with time, displaying broad lines similar to those of SLSNe I (Benetti et al. 2014). In this comparison, the noticeable differences are that the Mg I line has a larger equivalent width (a factor of 2 higher in LSQ14an than the others), and the fast-evolving SLSNe do not show the forbidden [O I] and [Ca II] at this phase. In truth, a clear detection of [Ca II] has only been recently reported for a fast-evolving SLSNe (at +151d from maximum in Gaia16apd, Kangas et al. 2016, a SLSNe that shows a late photometric behaviour intermediate between the fast- and slow-evolving events), but see Section 4.2 for a more detailed analysis of the elements contributing to the feature. The continuum of LSQ14an, and the other slowly


<sup>6</sup> All the phases are reported in rest-frame unless otherwise stated.

evolving objects above are also significantly bluer than the fast-evolving and stay blue for longer (more than a factor of 3 in time) as already noted for SN 2015bn by (Nicholl et al. 2016a). The prominent emission feature at 5200 Å to 5600 Å has a markedly different profile shape in LSQ14an - sharper with a peak consistent with it being a blend of Mg I  $\lambda$ 5180 and [Fe II]  $\lambda$ 5250. This profile is similar to that marking the beginning of the pseudo-continuum, dominated by Fe lines, in interacting SNe such as SNe 2005gl and 2012ca (Gal-Yam et al. 2007; Inserra et al. 2016a) but in the SLSNe case the feature is 200 Å bluer and not as sharp as in the interacting transients.

In Figure 4 we show the optical spectroscopic evolution of LSQ14an with nine spectra taken from +55d to +253d after the estimated peak epoch. At all epochs, the spectra are “pseudo-nebular”, showing permitted, semi-forbidden and forbidden lines. However it is remarkable that the lines that are unambiguously identified in Jerkstrand et al. (2017) through modelling the nebular phase spectra at +300-400d are already visible and prominent at +50d (cfr. Figure 5). We note that the epoch of the first spectrum is supported by the previous spectroscopic comparison of Figure 3 and also by LSQ14an photometric evolution (cfr. Section 5). The strongest emission line features are almost all the same in these spectra which are separated by a long time interval up to a year (see Figure 5 in this paper and fig. 7 of Jerkstrand et al. 2017). The strong blue continuum is a feature of both LSQ14an, and SN 2015bn even after subtraction of the blue host galaxy. A similar behaviour is also shown by SN 2015bn on an almost equivalent time baseline of LSQ14an (Figure 5). The strongest emission lines identified at late time are already in the +72d spectrum (and in any spectrum after  $\sim$ 50d post peak; see Nicholl et al. 2016a). The only small difference is the [O I]  $\lambda$ 6300, 6364 that appears later in SN 2015bn (Jerkstrand et al. 2017). We note that the continuum flux of the host galaxy becomes comparable to the SN at +150d and later, therefore the last two spectra at +200d and +252d will contain host galaxy flux.

The near infrared coverage of X-Shooter at +78d and +200d allows the identification of the line at 9200 Å line as O I  $\lambda$ 9263. This line was identified as O I in the modelling of the +315d spectrum of SN 2015bn and the +365d spectrum of LSQ14an by Jerkstrand et al. (2017). This is a recombination line ( $2s^2 2p^3 ({}^4S^{\circ}) 3p$ ) and we measure a full width at half maximum (FWHM) of  $v \sim 8000 \text{ km s}^{-1}$  in the +200d spectrum. The O I  $\lambda$ 7774 is another recombination line which decays from the lower state of the O I  $\lambda$ 9263 transition. As expected, the two lines have the same velocity.

We also measured the FWHM velocity of the [O I] and [Ca II], which are usually the strongest forbidden lines in nebular spectra of supernovae. They display an almost constant velocity with average  $v \sim 7500 \text{ km s}^{-1}$  for [O I] and  $v \sim 7400 \text{ km s}^{-1}$  for [Ca II]. The calcium line, clearly not Gaussian, is noticeably asymmetric (see Section 4.2) throughout the evolution and is much stronger than [O I]. The feature could be a blend of [O II]  $\lambda$ 7320, 7330 and [Ca II]  $\lambda$ 7291, 7323. A more detailed discussion of this and the implications for the explosion mechanism are presented in Jerkstrand et al. (2017). However, we already notice that lines coming from the same elements show different velocities and how semi-forbidden and forbidden lines are observed



**Figure 5.** *Top:* LSQ14an at +78d (black) and +364d (blue, after subtraction of the galaxy model, and removal of host galaxy lines). Note that the +364d spectrum (Jerkstrand et al. 2017) has been scaled by 2.3 to match the earlier one. The lines have not changed in almost a year. The same is true if the +50d and +409d (Jerkstrand et al. 2017) spectra are compared. *Bottom:* similar comparison of before but for SN 2015bn (Nicholl et al. 2016a; Jerkstrand et al. 2017).

earlier and at higher temperatures than normal stripped envelope SNe.

What is surprising is the absence of the Ca II NIR triplet  $\lambda$ 8498, 8542, 8662 up to 200 days (see Figure 4), whereas it is present at +365d in Jerkstrand et al. (2017). Since it requires high temperatures, this feature is usually seen quite early and disappears after 100 days. On the contrary, in LSQ14an the evolution is the opposite and in some extent similar to that of SN 2015bn where the NIR Ca grows in strength from 100 days to more than 300d (Nicholl et al. 2016b; Jerkstrand et al. 2017). The two X-Shooter spectra have a coverage out to  $2.0 \mu\text{m}$  in the rest-frame, but they are featureless beyond  $1.0 \mu\text{m}$  and display no strong and identifiable emission lines.

The narrowest emission lines which are lines observable throughout the spectroscopic evolution, are the Balmer, [O II] and [O III] lines due to the host galaxy. They have a constant velocity of  $v = 150 \pm 20 \text{ km s}^{-1}$  (measured from the two X-Shooter spectra) comparable with those of the host galaxies of SLSNe in previous studies (e.g. Leloudas et al. 2015a).

**Table 2.** Breakdown of the oxygen and calcium lines analysis from our X-Shooter spectra of LSQ14an (see Sections 4.1 & 4.2).

Phase (day)	+78.2	+200.0
[O I] $\lambda\lambda 6300, 6363$		
shift ( $\text{km s}^{-1}$ )	-	-
$v$ (FWHM, $\text{km s}^{-1}$ )	7500	7500
$n_e$ ( $\text{cm}^{-3}$ ) <sup>†</sup>	$\sim 10^9$	$\sim 10^9$
[O III] $\lambda 4363$		
shift ( $\text{km s}^{-1}$ )*	-1400	-1400
$v$ (FWHM, $\text{km s}^{-1}$ )	3200	3200
$n_e$ ( $\text{cm}^{-3}$ )	$\sim 10^7$	$\sim 10^7$
[O III] $\lambda\lambda 4959, 5007$		
shift ( $\text{km s}^{-1}$ )*	-1400	-1400
$v$ (FWHM, $\text{km s}^{-1}$ )	3500	3500
$n_e$ ( $\text{cm}^{-3}$ )	$\sim 10^7$	$\sim 10^7$
[Ca II] $\lambda\lambda 7291, 7323$		
shift ( $\text{km s}^{-1}$ )*	-1300	0
$v$ (FWHM, $\text{km s}^{-1}$ )	7400	7400
$n_e$ ( $\text{cm}^{-3}$ ) <sup>†</sup>	$> 10^9$	$> 10^9$
[O II] $\lambda\lambda 7320, 7330$ + [Ca II] <sup>‡</sup>		
shift ( $\text{km s}^{-1}$ )*	-1900/-1700	-1400/0
$v$ (FWHM, $\text{km s}^{-1}$ )	1700/2800	1700/2800
$n_e$ ( $\text{cm}^{-3}$ ) <sup>†</sup>	$\sim 10^9$ / $> 10^9$	$\sim 10^9$ / $> 10^9$

\* w.r.t. the centroid of the line or main doublet


† data derived from the estimates reported in Jerkstrand et al. (2017)

‡ left values are related to [O II], while the right to [Ca II]

#### 4.1 [O III] $\lambda 4363$ and $\lambda\lambda 4959, 5007$ lines

A peculiarity shown by the spectroscopic evolution of LSQ14an is the presence of a broad component of [O III]  $\lambda 4363$  and  $\lambda\lambda 4959, 5007$  (see Figure 6). In the first two PESSTO spectra with EFOSC2, these oxygen lines are prominent but the resolution of the Grism#13 prevented easy identification of emission broader than the host galaxy lines (see Figure 6). The first X-Shooter spectrum taken at +78d shows that [O III]  $\lambda 4363$  and  $\lambda\lambda 4959, 5007$  have a broad component arising in the supernova ejecta. The existence and origin of these lines as broad [O III] was first pointed out by Lunnan et al. (2016, for PS1-14bj which is a very slow-evolving SLSN) and they have been analysed in detail in Jerkstrand et al. (2017) in the later nebula spectra.

The average velocities of the components, as measured from the two X-Shooter spectra are  $v_{\lambda 4363} \sim 3200 \text{ km s}^{-1}$  and  $v_{\lambda\lambda 4959, 5007} \sim 3500 \text{ km s}^{-1}$  (see also Figure 6). Although these broad lines are severely blended with the galaxy emission lines in the lower resolution PESSTO spectra, we measure no velocity evolution in the FWHM of the features over the whole period (similarly to the other lines discussed in Section 4). The velocities of [O III] we measure from the X-Shooter spectra are a factor 2-3 less than those of the [O I] forbidden lines and the O I recombination lines. Even if we assume a low-mass, dilute oxygen region, allowing a higher energy deposition than the other slow-evolving SLSNe (Jerkstrand et al. 2017), this would not take in account their intrinsic different velocities, which could be explained with multiple emitting regions. Something similar was suggested



**Figure 6.** *Left:* LSQ14an [O III]  $\lambda 4363$  evolution from the two X-Shooter spectra at +78.2 and +200.0 days. *Right:* LSQ14an [O III]  $\lambda\lambda 4950, 5007$  evolution. The zero velocity is set at 5007 Å, since that is the strongest component of the [O III] doublet. The vertical dashed blue lines denote the centroids of the lines. H $\gamma$  (right panel) and H $\delta$  (left panel) are also visible.

in case of SN 2015bn but for different oxygen lines (Nicholl et al. 2016b).

The measured ratio for these collisionally excited lines in the optically thin limit can provide some information about the electron density of the emitting region where those are formed

$$\frac{f_{\lambda\lambda 4959, 5007}}{f_{\lambda 4363}} = \frac{7.90 \exp(3.29 \times 10^4 / T)}{1 + (4.50 \times 10^{-4} \times n_e / T^{1/2})}, \quad (1)$$

from Osterbrock & Ferland (2006), where  $T$  is the temperature and  $n_e$  is the electron density of the medium. We measured  $f_{\lambda\lambda 4959, 5007} / f_{\lambda 4363} = 1.8$  from our X-Shooter

spectra on an average of two measurements with a Gaussian fit. The ratio is consistent with that reported by Lunan et al. (2016) for PS1-14bj ( $< 3$ ) and hence less than what is typically observed in gaseous nebulae (Osterbrock & Ferland 2006) and nebular type IIIn (e.g. SN 1995N, Fransson et al. 2002). We initially assumed the ejecta temperature ( $T \sim 8000$  K) and found  $n_e = 6.2 \times 10^7 \text{ cm}^{-3}$  for the emitting region of the [O III] lines. If we increase the assumed temperature to  $T = 20000$  K the density decreases to  $n_e = 8.0 \times 10^6 \text{ cm}^{-3}$ . By comparison, the [O I]  $\lambda\lambda 6300, 6363$  arising from the inner region of the ejecta has  $n_e \sim 10^9 \text{ cm}^{-3}$  (as measured for late spectra of LSQ14an,  $\sim 200$ d later, by Jerkstrand et al. 2017, and having taken in account that density evolves as  $t^{-3}$ ). These lines are mainly collisionally excited with thermal ions at all times, as they are close to the ground state (see Jerkstrand et al. 2017, for a more in depth analysis).


Although [O III] and [O I] can come from the same physical region, the different densities ( $n_e([\text{O III}]) < n_e([\text{O I}])$ ), velocities ( $v([\text{O III}]) < v([\text{O I}])$ ) and the absence of a comparable strong [O I]  $\lambda 7300$  would suggest that these lines are formed in two different regions (see Section 8 for the interpretation).

In the magnetar scenario, the [O III] lines could come from an oxygen-rich ejecta layer which is ionised from the X-ray energy coming from the inner engine. This would be very similar to the scenario of a pulsar wind nebula expanding in a H- and He-free supernova gas, for which the second strongest lines predicted are [O III]  $\lambda\lambda 4959, 5007$  (Chevalier & Fransson 1992). On the other hand, the strongest lines observable in a H-free pulsar wind nebula should be those of [S III]  $\lambda\lambda 9069, 9532$ <sup>7</sup>. These are not observed in our NIR spectra, suggesting it may not be the X-ray heating from the magnetar that causes the presence of the [O III] lines.

As shown in Figure 6, we also note that the peak of the [O III]  $\lambda 4363$  is blueshifted by  $\sim 1400 \text{ km s}^{-1}$  compared to the rest-frame of the SN and the centroid of the [O I] lines (see Table 2 for a comparison between the velocities of the oxygen lines). Similarly the [O III]  $\lambda\lambda 4959, 5007$  lines have a centroid which appears shifted of  $\sim 1400 \text{ km s}^{-1}$ . These lines and shifts are clear in the high resolution X-Shooter spectra, but are not easily discerned in the lower resolution EFOOSC2 spectra.

#### 4.2 Forbidden lines around 7300Å

Explosively produced calcium would likely come from an O-burning zone rather than a C-burning zone, where the [O I]  $\lambda\lambda 6300, 6363$  lines would form. In slow-evolving SLSNe [Ca II]  $\lambda\lambda 7291, 7323$  has been observed in emission from surprisingly early stage: about 50-70 days from maximum (e.g. SNe 2007bi, 2015bn; Gal-Yam et al. 2009; Young et al. 2010; Nicholl et al. 2016a). We see similar behaviour from LSQ14an. [Ca II] appears almost at the same phase in normal stripped envelope SNe (after 60-70 days, e.g. SNe 1994I, 2007gr; Filippenko et al. 1995; Hunter et al. 2009) and later in broad line SNe (after 100d, e.g. SN 1998bw; Patat et al.



**Figure 7.** LSQ14an [Ca II]+[O I] evolution from the first epoch at +55.0 days to the last at +252.8d. A double gaussian fit (red dashed line) to [Ca II] is shown for the +59.7d epoch to highlight the skewed shape of the profile. The two vertical black dashed lines denote  $\lambda\lambda 7291, 7323$  at rest-frame, while the solid green mark [O II]  $\lambda\lambda 7320, 7330$ . The zero velocity is set at the centroid of the [Ca II] doublet. Phase with respect to the maximum is also shown.

<sup>7</sup> We note that if hydrogen is also present in the nebula, [O III]  $\lambda 5007$  and [S II]  $\lambda 6723$  are stronger than [S III] (Kirshner et al. 1989). However, in our case also [S II] is not observed.


2001), which however have less massive ejecta than the slow-evolving SLSNe. Also in fast-evolving SLSNe it appears at least after 100 days from peak (Kangas et al. 2016). Hence in SLSNe, where it is detected, it seems to appear earlier than expected.

The [Ca II] profile shows a component which has a peak at a velocity of  $-1300 \text{ km s}^{-1}$  with respect to rest-frame [Ca II]. Although we are somewhat limited by the EFOSC2 resolution, the blue shifted peak is visible in all spectra from +55d to +253d (see Figure 7). The X-Shooter spectra show the line profile well resolved and indicate that this is not a simple blue shifting of the peak of [Ca II]. The three most likely explanations are either asymmetry in the [Ca II] emitting region, a second [Ca II] emitting region which is at a different velocity, or another ionic component. Figure 7 shows a double gaussian fit to the broad [Ca II] in the second epoch (FWHM=  $7400 \text{ km s}^{-1}$ ) with the blue peak visible. If this is [Ca II], then the blueshift of  $-1300 \text{ km s}^{-1}$  is similar to that observed in the [O III] lines. Therefore the region that is producing the [O III] lines may also be emitting [Ca II].

There is an iron line, [Fe II]  $\lambda 7155$ , but this transition is  $120 \text{ \AA}$  too far blue and the X-Shooter profile would not support that identification. Furthermore we would also have expected to observe [Fe II]  $\lambda 1.257 \mu\text{m}$  with a similar strength, but the spectra at that wavelength are featureless. Jerkstrand et al. (2017) showed that the strength of the overall emission feature of [Ca II] is much stronger in LSQ14an than in the other two comparator objects (SN 2015bn and SN 2007bi). Their models at +365 and +410 days indicate that there is likely a significant contribution from [O II]  $\lambda\lambda 7320, 7330$ . If the blue peak is indeed dominated by [O II]  $\lambda\lambda 7320, 7330$  then it is blue shifted by  $-1920 \text{ km s}^{-1}$  and has a width of around  $1700 \text{ km s}^{-1}$  (from a blended two gaussian fit as evaluated by the 200d epoch). It is possible that this is [O II] from the same physical region that is producing the narrow [O III] lines, since the blueshift is similar. The velocity width of the line is lower than that estimated for [O III] (of order  $2900 \text{ km s}^{-1}$ ), but the blending of both these lines with the broad [Ca II] component means that both estimates are uncertain. It is certainly clear that both are narrower than the widths of main components of [Ca II] and [O I], which are of order  $7000\text{-}8000 \text{ km s}^{-1}$  and that the feature exhibits a blue-shift.

Surprisingly, LSQ14an is not the only slow-evolving SLSNe I showing this profile. As highlighted in Figure 8, SN 2015bn<sup>8</sup> shows a similar blue-shifted peak to that of LSQ14an, whereas that of SN 2007bi is centred at the rest wavelength. In PTF12dam, despite the low S/N of the spectra at that wavelength region, [Ca II] also appears to be centred at the rest wavelength (see extended data fig. 3 in Nicholl et al. 2013). Therefore we can conclude that the blue-shifted feature associated with [Ca II]  $\lambda\lambda 7291, 7323$  is not a unique feature of LSQ14an but is also not ubiquitous in all slow-evolving SLSNe I.

A blue-shifted emission from [Ca II] is not uncommon in core collapse SNe interacting with a CSM at late times. Type II SNe 1998S, 2007od (Pozzo et al. 2004; Inserra et al.



**Figure 8.** Blue-shifted [Ca II]+[O II] profile of LSQ14an compared to the same line of other two slow-evolving SLSNe at similar epoch. The two vertical black dashed lines denote  $\lambda\lambda 7291, 7323$  at rest-frame, while the solid green mark [O II]  $\lambda\lambda 7320, 7330$ .


2011) showed a similar blue-shifted [Ca II] profile, as well as type Ib/c (e.g. Taubenberger et al. 2009; Milisavljevic et al. 2010). This is usually interpreted as a consequence of dust formation in ejecta, which causes the dimming of the red wings of line profiles due to the attenuation of the emission originating in the receding layers, or residual line opacity especially for stripped envelope SNe as also shown by models (see Jerkstrand et al. 2015). However, in the case of LSQ14an, only [Ca II]+[O II] and [O III] show a blue-shifted peak and only [Ca II]+[O II] shows a clearly skewed profile. If the explanation were dust, then it would suggest that the dust distribution is not homogenous and most likely in clumps. Hence we favour the component being either narrow [O II] arising from the same region as the [O III] lines (see Table 2), another component of [Ca II] also located physically with the narrow [O III] emitting region, or an effect of the residual line opacity. See Section 8 for more discussion and interpretation.

## 5 LIGHT AND BOLOMETRIC CURVES

### 5.1 Light curves

The LSQ14an light curve shows a decline in all the available bands since the discovery epoch. This confirms that the SN was discovered past peak. However, the decline rate is not uniform and the uncertainty in the photometric measurements is small enough to detect clear changes in the decline. In the  $r$ -band we observe a  $1.8 \text{ mag } 100\text{d}^{-1}$  decline ( $2.1 \text{ mag } 100\text{d}^{-1}$  in rest-frame) until 91 days past the rest-frame assumed peak, followed by a slower decrease of

<sup>8</sup> We chose an earlier spectrum of SN 2015bn, with respect to that of LSQ14an, due to its high S/N. However, the skewed [Ca II] is also observable later than 50 days (see Nicholl et al. 2016a).




**Figure 9.** *Top:* *griBVRJHK* light curves of LSQ14an in the observer frame with respect to the assumed maximum. The epochs of spectroscopy are marked with black, dashed lines, while the discovery with the letter ‘D’. *Bottom:* Comparison of the *r*-band magnitude of well sampled slow-evolving SLSNe.

1.2 mag 100d<sup>-1</sup> (1.4 mag 100d<sup>-1</sup> in rest-frame) from 114 to 257 days and a faster decrease of 2.7 mag 100d<sup>-1</sup> (3.1 mag 100d<sup>-1</sup> in rest-frame) in the last phase from 365 to 408 days. All declines are more rapid than that of <sup>56</sup>Co to <sup>56</sup>Fe, which should be at  $\sim 1.1$  mag 100d<sup>-1</sup> in case of full trapping (Wheeler & Benetti 2000). In the bottom panel of Figure 9, we show LSQ14an *r*-band absolute light curve compared with those of four slow-evolving SLSNe with data during the same phase. All five are similar, with the resemblance between LSQ14an, SN 2015bn and PTF12dam being particularly close. This suggests that our assumed peak epoch from the spectra comparison should be correct within the uncertainties above stated. Previously SN 2007bi has been assumed, or proposed to be a unique object which requires a physical interpretation as a pair-instability SN (Gal-Yam et al. 2009), but it is clear that there is a class of these slow-evolving SLSNe (as initially proposed by Gal-Yam 2012), which LSQ14an (and the others) are part.

## 5.2 Luminosity, temperature and radius

Our LSQ14an imaging dataset lacks UV coverage and has only one epoch of NIR coverage. Hence, only an optical pseudo-bolometric can be built (see Inserra et al. 2016b, for further details in the construction of the bolometric



**Figure 10.** *Panel A:* bolometric light curve of LSQ14an obtained by integrating the flux over the optical band (open circles) and from UV to NIR (black filled circles). *Panel B:* Bolometric light curve of LSQ14an compared with those of SN 2007bi, PTF12dam, SN 2015bn. The <sup>56</sup>Co to <sup>56</sup>Fe decay slope is shown as a dashed black line. See text for further information. *Panel C:* temperature evolution of best-fit blackbody models evaluated for photometry (black circles) and spectroscopy (open squares) compared to that of the well observed SN 2015bn (cyan solid line) and PTF12dam (orange dashed line). *Panel D:* evolution of blackbody radius (black circles), compared to that of the same SLSNe of panel C.


light curve). Applying the  $K$ -correction<sup>9</sup>, fitting the available spectral energy density (SED) and integrating the flux from 1500 Å to 25000 Å should provide a good approximation to the full bolometric (see Inserra et al. 2013, 2016a). Pseudo and total bolometric light curves are shown in the top panel of Figure 10, comparing the bolometric light curve of LSQ14an with those of other slow-evolving SLSNe I, namely SN 2007bi, PTF12dam, SN 2015bn. They are all very similar, with decline rates from peak to 150 days that match the radioactive decay of  $^{56}\text{Co}$  to  $^{56}\text{Fe}$ . After 150 days, all of them deviate from the  $^{56}\text{Co}$  rate, as already noted in the case of SN 2015bn by (Nicholl et al. 2016b). LSQ14an is the less noticeable in this behaviour, possibly due to the sparse sampling in that phase combined with the light curve oscillations (see Section 6). If they were mostly powered by  $^{56}\text{Ni}$  then they can not be fully trapped. This argues strongly against them being pair-instability explosions, since the massive ejecta required for this explosion mechanism inherently implies full trapping up to 500 days after maximum light (Jerkstrand, Smartt, & Heger 2016). Hence, while at first sight the approximate match to  $^{56}\text{Co}$  is appealing to invoke nickel-powered explosions, it quantitatively does not fit with massive pair-instability explosions. We also see some evidence that after 250–300 days, there may be a steepening in the decline slope in all four. The decline at times after 300 days is roughly consistent with a power-law of index  $\alpha = 5$  ( $L \sim t^{-5}$ ), typical of an adiabatic expansion (a.k.a. Sedov-Taylor phase), which is unexpected at this phase (see Section 8).

The blackbody fit to the photometry delivers information about the temperature and radius evolution. We also used a blackbody fit on our spectra in order to increase the available measurements. We then compared our results with the well sampled PTF12dam (Chen et al. 2015; Vreeswijk et al. 2017) and SN 2015bn (Nicholl et al. 2016a,b). LSQ14an shows an overall steady temperature of 8000K, which is slightly higher than that measured in SN 2015bn and PTF12dam but still consistent within the errors.

The evolution of the blackbody radius of LSQ14an shows an overall similar behaviour to that of PTF12dam and to SN 2015bn up to 150 days. After 150d it remains roughly constant until the end of our dataset in a similar fashion to that of PTF12dam, while SN 2015bn steadily decreases. A small increase of the radius is noticeable from  $\sim 60$ d to  $\sim 75$ d, which corresponds to the first of the two possible fluctuations observed in LSQ14an light curve. Such behaviour was also observed for the undulations in SN 2015bn (Nicholl et al. 2016a).

## 6 LIGHT CURVE UNDULATIONS IN HYDROGEN-POOR SLOW-EVOLVING SLSNE I

All the individual LSQ14an light curves show some low level evidence of undulations. To explore this further, we fitted a first order polynomial to the bolometric light curve. The



**Figure 11.** *Top:* Residuals, after subtracting fits to the declining bolometric light curve, showing undulations over the selected phase in the case of slow-evolving SLSNe. *Bottom:* LSQ14an residuals compared with those of four fast-evolving SLSNe, which do not show any appreciable fluctuations, with the possible exception of SSS120810.

choice of using the bolometric allows us to avoid an analysis dominated by  $K$ -correction or line evolution within a particular pass-band. We chose to focus from 50 to almost 250 days excluding data after 250 days due to sparse sampling. The choice of a linear function to fit is due to the fact that slow-evolving SLSNe I light curves are usually well reproduced by models having a uniform, steady decline over the analysed baseline. We then subtracted the fit from the light curve over that timescale. The residuals are shown on the bottom panel of Figure 11.

The residuals appear to have at least one clear fluctuation phase from 50 to 200 days in rest-frame. At 75 days we observe a fluctuation of 0.06 dex from the linear fit. Another less distinct oscillation (0.04 dex) is visible around 170 days. Nevertheless, there is a residual pattern which oscillates across the linear decline fit with an amplitude that appears larger than the errors on the flux measurements.

We then compared the LSQ14an residual analysis with those of our selected slow-evolving SLSNe I sample. The phase analysed here is after that of the ‘knee’ observed in SN 2015bn (Nicholl et al. 2016a). We also observe a small fluctuation from 75 until 120 days in the SN 2015bn data, similar to that of LSQ14an but with lower amplitude (0.02 - 0.03 dex). Nonetheless, our estimated errors are smaller

<sup>9</sup> In this paper all the absolute magnitudes have been  $K$ -corrected with the SNAKE code (Inserra et al. 2016b).

than the observed amplitude for this SN too. PTF12dam may also have some undulations present, but in this case the amplitude is comparable or less than the errors. Fluctuations are also observed for PS1-14bj, which shows a 0.03 dex fluctuation at around 100 days and a more noticeable one (0.07 dex) around 220 days. For all objects the undulations are small, with an average of 0.05 dex, but they are generally greater than the photometric errors (0.01-0.03 dex depending on the phase and distance of the object). On the other hand, SN 2007bi shows stronger oscillations from 0.05 (at 60 days) to 0.12 dex (at 115 days), a factor ten bigger than the errors. Following on from the analysis of SN 2015bn by Nicholl et al. (2016a) which first highlighted these undulations, it may be that light curve fluctuations are common place in slow-evolving SLSNe I, but that their magnitude is highly variable from object to object. On the other hand, we also checked if such behaviour was present in the fast-evolving SLSNe I. We used data of the well sampled SNe 2010gx and 2011ke (Pastorello et al. 2010; Inserra et al. 2013) together with LSQ12dlf and SSS120810, which have data up to  $\sim 140$ d (Nicholl et al. 2014). We used a linear fit in the case of SN 2010gx, LSQ12dlf and SS120810 and a second order polynomial in the case of SN 2011ke. The difference in the latter is due to the fact that the SN light curve undergoes a noticeable transition to the tail phase just around  $\sim 50$  days (see Inserra et al. 2013) and a linear fit would have produced a bogus oscillation in the residuals. SSS120810 seems to experience a genuine oscillation around  $\sim 70$  days, as already noticed by Nicholl et al. (2014), but the rise is sharp and the width is small, especially when compared with the slow-evolving. LSQ12dlf could have experienced an oscillation similar in amplitude to those of before, or alternatively could be the beginning of the tail phase, but the errors and the lack of later data prevent any conclusion. In general, we do not see any clear trend in the residuals resembling those of slow-evolving SLSNe I, even though the time coverage is too short and future work should analyse earlier epochs (e.g.  $0 < \text{phase (d)} < 60$ ) where fast-evolving SLSNe are better sampled.

In the magnetar scenario, light curve oscillations could be the consequence of the magnetar ionisation front. In particular, when O II and O IV layers driven by the hard radiation field of the magnetar reach the ejecta surface a change of the continuum opacity will happen (Metzger et al. 2014). Lower ionisation states generally penetrate further and reach the surface earlier. The less ionised layer should reach the surface in  $\sim 10$  days from explosion (well before our detections of the light curve deviations). The other should happen at the same time of an X-ray shock breakout that, considering the ejecta masses and spin period of slow-evolving SLSNe I, should happen more than a year after peak (see Section 7). Hence, a sudden rise in opacity, would be non trivial to account for and would require a radiative transfer calculation to test the feasibility of this scenario. Multiple changes in opacity, which would be required for SNe 2007bi and 2015bn, would be more difficult to explain.

Another option suggested to explain the pre-peak undulation of SN 2015bn (Nicholl et al. 2016a) is that the fluctuation is powered by oxygen recombination, which would be below 12000 K (Hatano et al. 1999; Quimby et al. 2013; Inserra et al. 2013), in a similar fashion to what occurs in type II SNe within the hydrogen layer. Since the oscillations


after 50 days already happen at an almost constant temperature of  $T_{\text{off}} \approx 8000\text{K}$  (see Section 5.2) it would be difficult to explain all of them with this interpretation.

On the other hand, oscillations have been observed in SN light curves of interacting transients such as type Ibn, IIn or SN impostors (e.g. SNe 1998S, 2005ip, 2005la, 2009ip, 2011hw; SNhunt248 Fassia et al. 2000; Smith et al. 2009; Fox et al. 2009; Stritzinger et al. 2012; Pastorello et al. 2008; Fraser et al. 2013; Margutti et al. 2014; Martin et al. 2015; Smith et al. 2012; Pastorello et al. 2015; Kankare et al. 2015). To reproduce the short time scale oscillations observed in slow-evolving SLSNe (amplitude of 50-100 days) with ejected masses of 7-16  $M_{\odot}$  (Chen et al. 2015; Nicholl et al. 2016a; Inserra et al. 2016c) may be difficult since massive ejecta should not produce small scale changes in the light curve (e.g. Gawryszczak et al. 2010; Fraser et al. 2013). This, together with the small increase in the radius, would disfavour an interaction scenario such as that of pulsational pair instability SNe (PPISNe) where thermonuclear outbursts due to a recurring pair-instability release massive shells of material will collide with each other if the latter shells have velocity higher than the initial (Woosley et al. 2007). As a consequence, a viable scenario would need a relatively low mass CSM ( $< 1 M_{\odot}$ ) as that proposed by Moriya et al. (2015) for the late interacting slow-evolving SLSN iPTF13ehe (Yan et al. 2015). Indeed, assuming an average half-period of the fluctuation  $t=25$  days, average luminosity  $L=10^{43}$  erg  $s^{-1}$  and  $v=7000$  km  $s^{-1}$  (this work, Gal-Yam et al. 2009; Young et al. 2010; Nicholl et al. 2013, 2016a) and using the scaling relation  $L \approx M_{\text{CSM}} v^2 / 2t$  (Quimby et al. 2007; Smith & McCray 2007) we find that a CSM of no more than  $M_{\text{CSM}} \sim 0.04 M_{\odot}$  could be responsible for each fluctuation, similar to the findings of SN 2015bn (Nicholl et al. 2016a). If the light curve oscillations are periodic, they could be the consequence of a close binary system that drives the stripping of the progenitor stars (or the companion) causing a heterogeneous density structure of the CSM (e.g. Weiler et al. 1992; Moriya et al. 2015).

## 7 X-RAY LIMITS IN SLOW-EVOLVING SLSNE I

An alternative method to investigate the SLSNe progenitor scenario and powering mechanism is that of X-ray observations. Both the magnetar and the interaction scenario would predict X-ray emission but at different luminosities and at different phases.

In the case of the magnetar scenario, the pulsar wind inflates a hot cavity behind the expanding stellar ejecta, which is the pulsar wind nebula. Electron/positron pairs cool through synchrotron emission and inverse Compton scattering, producing X-rays inside the nebula. These X-rays ionise the inner regions of the ejecta, driving an ionisation front that propagates outwards with time (Metzger et al. 2014). LSQ14an shows a light curve similar to that of SN 2015bn and hence we can assume similar magnetar and ejecta parameters to those retrieved by Nicholl et al. (2016a,  $P = 1.7 - 2.1$  ms,  $B_{14} = 0.9 - 1.0$  G and  $M_{\text{ej}} = 8 - 15 M_{\odot}$ ) in their magnetar fit (for the magnetar semi analytic code see



**Figure 12.** SLSNe I X-ray luminosity limits (grey stars) and stacked limits (black open stars) as reported by Levan et al. (2013, but evaluated using the cosmology adopted here), compared with those of the available slow-evolving SLSNe I LSQ14an, SN 2015bn (this work) and PTF12dam (Levan et al. 2013), the single X-ray detection of SLSN I SCP06F6 and a sample of SNe detections (Ofek et al. 2013). X-ray limits in case of break-out from the pulsar nebula ionisation front (blue dashed dotted line) and interaction (red dashed lines for the two non massive CSM cases and red dotted for the massive CSM scenario) are also shown. See text for further details.

prescription of Inserra et al. 2013)<sup>10</sup>. The mass range is similar to that estimated by other work on slow-evolving SLSNe (Nicholl et al. 2013; Chen et al. 2015), with the only exception of PS1-14bj (Lunnan et al. 2016) showing redder spectra, longer rise-time and fainter peak luminosity than the bulk of slow-evolving SLSNe I. Thus, for our analysis, we can use an average mass of  $12 M_{\odot}$  for LSQ14an, SN 2015bn and in general for all slow-evolving SLSNe I. Following the prescription of Metzger et al. (2014, their equations 5, 6, 56 and A11) we would expect an X-ray break-out of  $L_X \sim 7.0 \times 10^{42}$  erg s<sup>-1</sup> at  $t_X > 700$ <sup>11</sup> days from explosion that corresponds to  $\gtrsim 630$  days from maximum light for an average rise-time of about 70 days.

Alternatively, in the interaction scenario the shock waves created by the interaction between the SN ejecta and the CSM heat gas to X-ray emitting temperatures and can possibly accelerate particles to relativistic energies. The presence of undulations in the light curves, would disfavour the presence of a massive circumstellar medium ( $7-19 M_{\odot}$ ; Chatzopoulos et al. 2013; Nicholl et al. 2016a), which would also power the peak luminosity (Chevalier & Irwin 2011). In contrast, a less massive circumstellar medium ( $<1 M_{\odot}$ , see Section 6) could be more suited to explain LSQ14an and other SLSNe I observables. Generally, during the interaction process between SN ejecta and CSM, the conversion of the kinetic energy into radiation is affected by complicated hydrodynamic and thermal processes including thin shell instabilities (Vishniac 1994), the Rayleigh–Taylor instability of the decelerating cool dense shell, the CSM clumpiness, mixing, and energy exchange between cold and hot compo-

nents via radiation and thermal conductivity (Chandra et al. 2015). Here we use a simplified approach in which the X-ray luminosity of both shocks is equal to the total kinetic luminosity times the radiation efficiency ( $\alpha_X > 0.1$ ; Moriya et al. 2013; Chandra et al. 2015). As a consequence, the energy mass relation would be  $E \propto M^{0.7}$ , assuming an outer power law of the density of the ejecta  $n=10$ .

This is equal to  $L_X \sim 2.3 - 57.3 \times 10^{41}$  erg s<sup>-1</sup> for a  $0.04-1.0 M_{\odot}$  CSM interacting for 200 days (the time when we observe the fluctuations) and to  $L_X \sim 1.7 \times 10^{43}$  erg s<sup>-1</sup> for  $13 M_{\odot}$  interacting for 400 days, which corresponds to the massive CSM case. In both cases we assumed  $\alpha = 0.1$  and an ejecta always more massive than CSM (e.g. Chatzopoulos et al. 2013; Nicholl et al. 2014). We note that in interacting SNe there is a higher efficiency in the optical (where  $\alpha \geq 0.5$ ; Moriya et al. 2013) than in X-ray (Ofek et al. 2014; Chandra et al. 2015), hence the highest luminosity achievable in our case would be five times those above. On the other hand, to estimate our X-ray luminosities we did not consider any photoelectric absorption. In our case such contribution is hard to evaluate and varies with time, decreasing the soft X-ray output after the start of interaction (see Chevalier & Fransson 2003). This absorption would increase in case of massive CSM with respect to normal interacting SNe ( $CSM \lesssim 1 M_{\odot}$ ), but the X-ray emission would increase too. Hence, the values reported above have to be treated as higher limits of  $L_X$  and they should be similar to the X-ray luminosity emitted around maximum optical light, or soon before that, in case of interaction with a massive CSM (see Chevalier & Fransson 2003).

We report here the measurements for LSQ14an and SN 2015bn spanning 124–876 days from assumed maximum for the former, and from -27 to 628 days for the latter. In order to place meaningful constraints, we chose observations having at least 4000s exposure or covering the late phase of the light curve ( $>300$  days since maximum). To date, no observation has been carried out later than

<sup>10</sup> The code is available at <https://star.pst.qub.ac.uk/wiki/doku.php/users/ajerkstrand/start>

<sup>11</sup> We note that these calculations are highly sensitive to the ejecta mass, that however seems similar for all slow-evolving SLSNe I.

those reported here and no X-ray emission has been detected in the data here reported, neither for the stacked images of each SN (evaluated adding all the frames together and assuming an average phase, see Table A5)<sup>12</sup>. For both SNe we assumed the H-column density to the Galactic value ( $n_{\text{H}}(\text{LSQ14an}) = 4.16 \times 10^{20} \text{ cm}^{-2}$  and  $n_{\text{H}}(\text{SN 2015bn}) = 1.22 \times 10^{20} \text{ cm}^{-2}$ ) since SLSNe are hosted in galaxies with little to no internal reddening (see Lunnan et al. 2014; Leloudas et al. 2015a; Perley et al. 2016). We used a generic spectral model with a photon index  $\Gamma = 2$  (Levan et al. 2013) to estimate the luminosity limits. We note that limits can change by a factor of  $\sim 3$  in flux for a variation of the photon index of order of ten. We also used a thermal model with  $kT = 1.55$  (equal to that used for SCP06F6; Levan et al. 2013, which is the only SLSN detected in X-ray) and hence a factor of 1.2 to 10 lower than those used in H-rich interacting SNe (e.g. Chandra et al. 2012; Ofek et al. 2014; Chandra et al. 2015).

Figure 12 shows the average limits between the two methods with the power law method usually resulting in higher luminosities than the thermal one, but still on the same order of magnitude. Only the stacked limits, LSQ14an limit at 876d, SN 2015bn limit at 259d and limits at  $>571\text{d}$ , are below the magnetar threshold reported above (dotted dashed blue line in Figure 12). SN 2015bn phase is from  $\sim 400$  to  $\sim 30$  days earlier than that of the expected ionisation front break-out, while LSQ14an is  $\sim 300$  days after and the break-out does not last so long at that luminosity. Indeed it declines with time (see Metzger et al. 2014). However, one needs to be very careful in interpreting the stacked limits in a meaningful way. They are only quantitatively meaningful if they are taken to represent a constant level of flux across the epochs which are stacked. In other words they by no means rule out time variable flux between the observed points. All limits are below the X-ray luminosity (dotted red line) that we should have observed in case of interaction with a dense and massive CSM or collision between two massive shells as suggested by the PPIsNe scenario. However, in the case of a massive CSM, the X-ray luminosity should decrease from peak epoch due to photoelectric absorption (Chevalier & Fransson 2003) and after a year should be a factor of two to fifteen less than that at peak. Hence, the late time limits could not be so constraining<sup>13</sup>. On the other hand, in case of interaction with a less massive CSM ( $0.04 M_{\odot}$ , lower dashed red line) we would have expected an X-ray emission at a similar phase to the fluctuations and hence earlier than 250 days. Such X-ray would have luminosities lower than our limits. This is fulfilled for both limits and stacked limits and all slow-evolving SLSNe. Although we have made many

simplifying assumptions, the most simple interpretations is that the X-ray limits disfavour the scenario in which the interaction of ejecta with a massive CSM is the powering mechanism for this SLSN. However both the magnetar engine and interaction with a less massive CSM could play a role in the spectrophotometric behaviour of slow-evolving SLSNe I.

## 8 DISCUSSION ON THE CHARACTERISTICS OF SLOW-EVOLVING SLSNE I

The LSQ14an dataset is limited to post-peak, but the density of the spectra and light curve monitoring and the two excellent X-Shooter spectra offer interesting further information to better understand the explosion scenarios of slow-evolving SLSNe I. In summary it is a slow-evolving SLSN of type I or Ic. There are a group of, at least, four low redshift ( $z < 0.2$ ) SLSNe I (or SLSNe Ic) that show very similar spectrophotometric evolution SN 2007bi, PTF12dam, SN 2015bn and LSQ14an. They have very broad light curves that initially ( $< 150\text{d}$  after peak) decline at a rate similar to <sup>56</sup>Co, although it is unlikely that this is the true power source of the luminosity. They have higher redshift analogues such as LSQ14bdq ( $z = 0.3450$ ; Nicholl et al. 2015a) PS1-11ap ( $z = 0.5240$ ; McCrum et al. 2014), DES13S2cmm ( $z = 0.6630$ ; Papadopoulos et al. 2015), iPTF13ajg ( $z = 0.7403$ ; Vreeswijk et al. 2014), SN 2213-1745 and SN 1000+0216 ( $z = 2.0458$  and  $z = 3.8993$ , respectively; Cooke et al. 2012). PS1-14bj ( $z = 0.5215$ ; Lunnan et al. 2014) is also similar but has the most extreme in terms of its light curve width, while iPTF13ehe ( $z = 0.3434$ ; Yan et al. 2015) shows a late time re-brightening due to interaction with a H-shell (something similar could have happened also to LSQ14bdq, Schulze private communications). In Sections 4.1 & 4.2 we reported two unusual characteristics of the spectra of LSQ14an, some of which had been observed before, as well as a somewhat common feature in the spectral evolution of this group of SLSNe. Furthermore, in Sections 5 & 6 we described distinctive features of LSQ14an light curves that seem shared by the sample of slow-evolving SLSNe I.

### 8.1 Spectroscopic characteristics

The first spectroscopic characteristic is the presence of [O III] lines which are significantly broader than the host galaxy, unambiguously formed in the SN ejecta and blueshifted compared to the rest of the ejecta (see Section 4.1 and Table 2). The second is the existence of a strong blueshifted component associated with the prominent [Ca II] line (see Section 4.2).

In LSQ14an and PS1-14bj, the [O III] lines have  $v \sim 3000 \pm 300 \text{ km s}^{-1}$ . In the case of LSQ14an, this is significantly lower than that of the bulk of the ejecta as traced by the strong [O I] and O I lines, around  $8000 \text{ km s}^{-1}$  (Mg I) is broader at  $\sim 12000 \text{ km s}^{-1}$ , although it may be a blend). Furthermore, the density of the [O III] emitting region is likely much less than that where [O I] is formed. This would suggest multiple emitting regions for the above ions. Ionised elements show line profiles much narrower than the neutral ones, strengthening the hypothesis that they come from different regions and likely from a region interior to that where

<sup>12</sup> For both SNe we only stacked the first three epochs since it would have been deceptive to also add the later one, which were carried out from five months to more than a year after the end of our optical campaign.

<sup>13</sup> The photoelectric absorption strongly depends on the column density of the cool gas behind the reverse shock. This is difficult to evaluate in a case of a massive ( $\gtrsim 10 M_{\odot}$ ) H- and He-free (or deficient) shell in a sub-solar metallicity environment since it would require assumptions on each term of the formulas. Hence, we avoid any proper calculation of this effect but we would like to warn the reader that photoelectric absorption cannot be ignored and would somehow decrease the X-ray emission at late time.

the neutral lines are formed. Since [O I] travel faster than [O III] (see Section 4.1 and Table 2), another possibility is to have a fast-dense outer region and a slow-diffuse region similar to an evacuated inner cavity, which could be another indirect evidence of a magnetar inner engine.

In the magnetar or inner engine scenario, these blue-shifted lines and multiple emitting regions could be explained if the ejecta was fragmented or axis-symmetric and the observer was in the direction of the axis of symmetry (or both). The latter would happen since only the nearer (blue) component is observed, while red photons would be scattered perpendicular to the observer. An axisymmetric ejecta has been observed for SN 2015bn (Inserra et al. 2016c), while a certain degree of fragmentation, and also asphericity of the ejecta has been observed in 2D magnetar models (Chen, Woosley, & Sukhbold 2016). On the other hand, if we assume a certain degree of interaction with a small amount of material (see Section 6), the blue-shift could be due to an additional contribution to the line coming from the cool dense shell (CDS), consequence of the shock propagating in the ejecta. This is blue-shifted because the CDS is moving at a velocity of  $\sim 1500 \text{ km s}^{-1}$  and is optically thick in the continuum, thus line photons emitted in the red are absorbed by the near side of the CDS (Dessart, Audit, & Hillier 2015). Since a high dilution factor is required to model [O III] and [O II] (Jerkstrand et al. 2017), this could be achieved in a diluted interaction region. We already mentioned as [O II] and [O III] lines are much narrower than neutral lines, which in the simplest scenario means an origin in the inner, slower ejecta. Lines from the inner ejecta are normally more prone to obscuration by a residual opacity (a residual “photosphere”), which can also explain their suppressed red wings. This would be another piece of circumstantial evidence for a hot, ionised inner region and thereby central powering.

Finally, an important point is that the spectra of slow-evolving SLSNe I at 50–70 days, surprisingly show almost identical emission features to those seen in the nebular phase at 300–400 days. The spectra at each phase have quite different continuum properties (the early spectra being much bluer), but the ionic transitions are the same. [O I]  $\lambda\lambda 6300, 6363$  significantly increases in strength with time (as expected in CC-SNe), but others such as [Ca II] and [O III] are relatively constant in luminosity. This implies that the line forming regions of these lines have values of density, temperature and opacity that remain roughly constant over this extended period of evolution. That could also be a consequence of a nearly constant powering mechanism. We see that the emission lines which are redder than  $5500 \text{ \AA}$  show a shift in the centroid of the peak between their values at 50–70 days and those at  $>300$  days (see Figure 5). For example in SN 2015bn the [O I] and O I centroids move toward bluer wavelengths, while [Ca II] moves to the red. Multiple emitting regions may thus be present not only in LSQ14an but may be a more common feature in slow-evolving SLSNe I. If slow-evolving SLSNe I are the more massive version of the fast-evolving as suggested by Nicholl et al. (2015b), sharing similar explosion energies and the same mechanism responsible for the enormous luminosities, we would expect to observe nebular lines later than the fast-evolving, as observed for normal stripped envelope. For example in SN 2009jf, the onset of emission from forbidden calcium is delayed until

85 days from peak (Valenti et al. 2011). However, nebular emission lines (e.g. [O I] or [Ca II]) in fast-evolving SLSNe I have only been observed at  $\sim 150$  days after peak (Kangas et al. 2016)<sup>14</sup> and not before  $\sim 100$  days (Quimby et al. 2011; Inserra et al. 2013).


Another important characteristic shown by slow-evolving SLSNe I is that their early time spectra are always bluer than the fast-evolving and the late time spectra are still remarkably blue even past 200 days (see fig. 7 in Jerkstrand et al. 2017, and Figure 5 here). They are significantly bluer than stripped envelope SNe, or the faster evolving SLSNe as also noted by Nicholl et al. (2015b, 2016b) for SN 2015bn. That could be explained if we assume a fairly constant  $8000 \pm 1000 \text{ K}$  blackbody type of emission with the observed forbidden and semi-forbidden lines in addition (as seen in the photometric fits in Figure 10). This type of underlying continuum emission is common scenario in SNe dominated by interaction, however in slow-evolving SLSNe I we still lack the presence of narrow lines forming in the unshocked material. Alternatively, this could be explained by multiple optically thick lines creating a blackbody-like continuum with the forbidden and semi-forbidden lines forming in more external region than those creating the pseudo blackbody continuum. As highlighted by Jerkstrand et al. (2017), a dense CSM shell of  $\sim 10M_{\odot}$  could reproduce the multiple emission regions and the low filling factor required, but published models are still not able to reproduce the bulk of velocities and the absence of narrow lines.

## 8.2 Light curve characteristics

We also highlighted that while the light curve declines at a rate similar to  $^{56}\text{Co}$  until 150 days, it increases (for all the objects) from 150 days onwards with a possible further steepening after  $\sim 300\text{d}$ . The change in decline after 150 days disfavors the pair instability scenario for slow-evolving SLSNe I since PISN explosion should exhibit full gamma-ray trapping and hence closely follow the decay of  $^{56}\text{Co}$  to  $^{56}\text{Fe}$  up to 500 days (Jerkstrand, Smartt, & Heger 2016). Furthermore, there are some likely real undulations (see Section 6) in the bolometric luminosity. These peculiarities could be present in the higher redshift analogues, but a combination of lower quality data at red wavelengths (observer frame) and late time (rest-frame) due to their high redshift prevent a proper investigation.

First of all we consider the possibility that these features are a consequence of the magnetar scenario. The only possibility would be a change in the density of the ejecta due to the magnetar ionisation front, linked to oxygen ionised states (mainly O II and O IV, Metzger et al. 2014), and moving toward the surface of the ejecta. This happens twice but the timescale is not consistent for both since the former occurs 3–4 weeks after the explosion, while the other at least 700 days after explosion. Moreover the second is also coincident with an X-ray breakout that has not been observed during the time frame in which we observed the undulations.

<sup>14</sup> We note that Kangas et al. (2016) reported that SLSN Gaia16apd shows a fast decline at early phase similar to SNe 2010gx and 2011ke but a late ( $>130\text{d}$ ) behaviour between the fast- and slow-evolving events.



**Figure 13.** The late time bolometric evolution of LSQ14an, together with those of slow-evolving SLSNe I SN 2007bi, PTF12dam, SN 2015bn. The green dashed ( $n = 3$ ) and red dotted ( $n = 4$ ) lines show the decline similar to a snowplow phase, while the solid blue line that similar to an adiabatic (Sedov–Taylor) phase. The black dot-dashed line shows the expected  $^{56}\text{Co}$  decay. The x axis is in semi-log scale to highlight the decline and the differences in the power law indexes.

Secondly, we examine the possibility that the phenomenon is due to some ejecta - CSM interaction. However even if it is, it is unlikely that this CSM interaction powers the full luminosity of the light curve since our X-ray analysis (see Section 7 for the assumed caveats) does not support the interaction with a massive CSM shell ( $\sim 13 M_{\odot}$ ).

We also see some evidence of an increase in the decline rates after 300 days (see Figure 13). This is observed in LSQ14an, SN 2015bn and PTF12dam and possibly even SN 2007bi, even though for this object the increase is less certain. This decline is steeper than  $t^{-3}$  or  $t^{-4}$ , which is known as the snow-plow phase and it has been observed in H-rich interacting SNe (e.g. Ofek et al. 2014; Fransson et al. 2014; Inserra et al. 2016a). Figure 13 shows that the decline observed in slow-evolving SLSNe I after 300 days seems to follow a power law of  $t^{-5}$ , which is the same slope of an adia-

batic expansion phase (the Sedov-Taylor phase). This would suggest conservation of energy that, at this time, could be achieved when the density is low enough and the gas cannot cool quickly. An adiabatic expansions at late time of a SN evolution happens when the mass of gas swept up becomes greater than the mass of ejecta and the kinetic energy of the original exploded envelope is transferred to the swept up gas, and that is heated up by the shock wave. This would require a large amount of interstellar medium or CSM. However, spectra and X-ray analysis of the sample do not show any of the *primary* observables usually seen in interacting SNe (e.g. narrow lines or X-ray emission).

In the case of LSQ14an, considering a hypothetical rise time from explosion to peak of approximately 70 days (Nicholl et al. 2013, 2016a) and using the simple formalism of  $R_{\text{shock}} = 2.1 \times 10^{13} (E/\rho)^{1/5} t^{2/5}$  (Reynolds 2008), where  $t = 370$  days from explosion and  $\rho \sim 10^8$  (Jerkstrand et al. 2017, and Section 4.1), we derive a distance of  $1.2 \times 10^{23}$  cm for the radius of the shock, which is at least a factor of  $10^8$  higher than the SN radius. This suggests it is too early for SLSNe to experience a “natural” adiabatic phase, that indeed usually happens after 200yrs in normal supernovae (Reynolds 2008). The approximate calculation above is also valid (in terms of rough values) for SN 2015bn and PTF12dam. A possible alternative is that these slow-evolving SLSNe experience a further decrease of the  $\gamma$ -ray trapping (see also Chen et al. 2015; Nicholl et al. 2016b). A further investigation is needed to understand if this behaviour belongs to all slow-evolving SLSNe I and what is the physical reason. We note that none of the fast-evolving SLSNe I discovered up to date has observations at  $> 300$  days, which prevent a similar investigation.

## 9 SUMMARY AND DEDUCTIONS

We have presented an extensive dataset on LSQ14an, covering its spectrophotometric behaviour from 55 to 408 days together with X-ray limits spanning over the same timescale and up to  $\sim 870$  days. We show that there are now four well observed, low redshift SLSNe I which are similar in their photometric and spectroscopic evolution: SN2007bi, PTF12dam, SN2015bn and LSQ14an. The combination of the datasets provides some new insights into these SNe.

LSQ14an was classified after maximum light and exhibits spectroscopic peculiarities such as broad ( $v \sim 3500$  km s $^{-1}$ ), blue-shifted [O III]  $\lambda\lambda 4959, 5007$  lines, a blue-shifted peak of [O II]  $\lambda\lambda 7320, 7330$  (or [O II] + [Ca II]  $\lambda\lambda 7291, 7323$ ), as well as the late appearance of NIR Ca and the remarkable fact that the lines observed at late time (300-400 days after maximum) are unambiguously shown already at 50 days. The spectra remain surprisingly blue and the blackbody temperature fits to the photometric flux shows a lack of cooling over 300 days. The density ( $n_e \sim 10^6$  cm $^{-3}$ ) and line velocity ( $v \sim 3000$  km s $^{-1}$ ) of these oxygen features are different than the bulk of ejecta ( $n_e \sim 10^9$  cm $^{-3}$ ,  $v \sim 8000$  km s $^{-1}$ ), suggesting that multiple emitting regions are responsible for the features observed in the spectra. This could be caused by an axis-symmetric or fragmented ejecta or by an additional component arising in a circumstellar material. Furthermore, an in depth analysis of the light curve shows hints of undulations as observed in other similar SLSNe I (e.g. SN 2015bn),

as well a decline faster than  $^{56}\text{Co}$  after 150 days and even faster after 300d (times measured from estimated maximum light). We investigated X-ray upper limits, presenting new limits for LSQ14an and SN 2015bn, and spectrophotometric behaviour with those of nearby slow-evolving SLSNe I showing similar coverage. The deepest limits (not stacked) disfavour the circumstellar interaction with a massive CSM ( $\sim 13 M_{\odot}$ ) that is usually needed to reproduce the slow-evolving SLSNe light curves.

We noticed that some of these observational features are not unique to LSQ14an but are common in, or even exclusive to, all slow-evolving SLSNe I observed to date

- they show semi-forbidden and forbidden lines from 50-70 days to peak with little to no variation up to 400 days. The strongest evolution is seen in the strengths of the [O I]  $\lambda\lambda 6300, 6363$  lines. The centroid of oxygen and calcium lines appear to move in time. In addition, ionised elements show line profiles different from the neutral ones and likely from a region interior to that where the neutral lines are formed, and where occultation effects can be easily produced. This suggests that multiple emitting regions are responsible of the overall spectra.

- [Ca II] appears earlier than in broad-line type Ic SNe and fast-evolving SLSNe. In two cases (SN 2015bn and LSQ14an) it exhibits a blue-shifted peak. In LSQ14an the line is likely dominated by a [O I] component. The Ca II NIR triplet is observed increasing with time, from 100 days onward, in contrast to what shown in normal stripped envelope SNe.

- the overall light curve evolution of these four objects are similar. Initially, all of them decay at a rate consistent with  $^{56}\text{Co}$  powering. But after 150 days their bolometric light curves decline faster than the fully trapped  $^{56}\text{Co}$  to  $^{56}\text{Fe}$  decay. Moreover, after  $\sim 300$  days from maximum light the bolometric light curves show a further steepening in the decline consistent with a power law of  $t^{-5}$ . The light curves overall are not consistent with massive ejecta from PISNe that fully trap the gamma rays from  $^{56}\text{Co}$  decay up to 500 days.

- light curve fluctuations seem to be common in the slow-evolving. They can be generally explained with changes in the density structure of the expanding material. Since they appear to occur more than once and at different phases in each object, they are likely due to interaction with a clumpy medium. The medium should have  $M < 0.04 M_{\odot}$  to explain the oscillations observed in four out of five inspected slow-evolving SLSNe I and to account for the X-ray limits here reported.

There remains a challenge to explain all these observables, together with those previously reported to date. For example the observed pre peak bumps, axis-symmetric ejecta, late spectra similar to broad line type Ic and reproducible by  $\sim 10 M_{\odot}$  of oxygen (Nicholl et al. 2015a; Smith et al. 2016; Nicholl & Smartt 2016; Inserra et al. 2016c; Jerkstrand et al. 2017; Nicholl et al. 2016b), with a common scenario. The two most appealing mechanisms for these slowly evolving SLSNe are still central engine (likely a magnetar) and the interaction, even though for the latter some observables are not in agreement with the classical observed behaviour in interacting transients. However, some further considerations can be made. At least a small degree of in-

teraction seems to happen in slow-evolving SLSNe I. If the slow and fast evolving SLSNe I share the same nature (i. e. mechanism and maybe progenitor system), the difference between them cannot be due exclusively to a difference in mass as previously suggested (Nicholl et al. 2015b) since that would not reflect the spectral evolution.

To understand if fast and slow-evolving SLSNe I are a similar or different kind of transients we need further, and more statistical, investigations. Further studies should also focus on a more detailed early (when we have available both the types) spectra analysis and modelling in order to increase the available information.

## ACKNOWLEDGMENTS

CI thanks Ragnhild Lunnan for sharing the data of PS1-14bj and Stuart Sim for stimulating discussions. This work is based on observations collected at the European Organisation for Astronomical Research in the Southern Hemisphere, Chile as part of PESSTO, (the Public ESO Spectroscopic Survey for Transient Objects Survey) ESO program 188.D-3003, 191.D-0935, 197.D-1075, and the X-Shooter programmes 093.D-0229, 092.D-0555. The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC Grant agreement no [291222]. SJS acknowledges funding from STFC grants ST/I001123/1 and ST/L000709/1. C.I. thanks the organisers and participants of the Munich Institute for Astro- and Particle Physics (MIAPP) workshop "Supernovae: The Outliers" for stimulating discussions. This research was supported by the Munich Institute for Astro- and Particle Physics (MIAPP) of the DFG cluster of excellence "Origin and Structure of the Universe". MF acknowledges the support of a Royal Society - Science Foundation Ireland University Research Fellowship. KM acknowledges support from the STFC through an Ernest Rutherford Fellowship. TWC and TK acknowledge the support through the Sofia Kovalevskaja Award to P. Schady from the Alexander von Humboldt Foundation of Germany. The Liverpool Telescope is operated on the island of La Palma by Liverpool John Moores University in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias with financial support from the UK Science and Technology Facilities Council. The Pan-STARRS1 Surveys (PS1) have been made possible through contributions of the Institute for Astronomy, the University of Hawaii, the Pan-STARRS Project Office, the Max-Planck Society and its participating institutes, the Max Planck Institute for Astronomy, Heidelberg and the Max Planck Institute for Extraterrestrial Physics, Garching, The Johns Hopkins University, Durham University, the University of Edinburgh, Queen's University Belfast, the Harvard-Smithsonian Center for Astrophysics, the Las Cumbres Observatory Global Telescope Network Incorporated, the National Central University of Taiwan, the Space Telescope Science Institute, the National Aeronautics and Space Administration under Grant No. NNX08AR22G issued through the Planetary Science Division of the NASA Science Mission Directorate, the National Science Foundation under Grant No. AST-1238877, the University of Mary-

land, and Eotvos Lorand University (ELTE) and the Los Alamos National Laboratory.

## REFERENCES

- Agnoletto, I., Benetti, S., Cappellaro, E., et al. 2009, *ApJ*, 691, 1348
- Alard C., 2000, *A&AS*, 144, 363
- Amorín R., Pérez-Montero E., Vílchez J. M., Papaderos P., 2012, *ApJ*, 749, 185
- Angus, C. R., Levan, A. J., Perley, D. A., et al. 2016, *MNRAS*, 458, 84
- Asplund M., Grevesse N., Sauval A. J., Scott P., 2009, *ARA&A*, 47, 481
- Baltay, C., Rabinowitz, D., Hadjiyska, E., et al. 2013, *PASP*, 125, 683
- Bianco F. B., Modjaz M., Oh S. M., Fierroz D., Liu Y. Q., Kewley L., Graur O., 2016, *A&C*, 16, 54
- Benetti S., et al., 2014, *MNRAS*, 441, 289
- Chandra P., Chevalier R. A., Chugai N., Fransson C., Irwin C. M., Soderberg A. M., Chakraborti S., Immler S., 2012, *ApJ*, 755, 110
- Chandra P., Chevalier R. A., Chugai N., Fransson C., Soderberg A. M., 2015, *ApJ*, 810, 32
- Chatzopoulos, E., Wheeler, J. C., Vinko, J., Horvath, Z. L., & Nagy, A. 2013, *ApJ*, 773, 76
- Chen K.-J., Woosley S. E., Sukhbold T., 2016, *ApJ*, 832, 73
- Chen, T.-W., Smartt, S. J., Jerkstrand, A., et al. 2015, *MNRAS*, 452, 1567
- Chen T.-W., Smartt S. J., Yates R. M., Nicholl M., Krühler T., Schady P., Dennefeld M., Inserra C., 2016, *arXiv:1605.04925*
- Chevalier, R. A., & Irwin, C. M. 2011, *ApJL*, 729, L6
- Chevalier R. A., Fransson C., 1992, *ApJ*, 395, 540
- Chevalier R. A., Fransson C., 2003, *LNP*, 598, 171
- Chomiuk, L., Chornock, R., Soderberg, A. M., et al. 2011, *ApJ*, 743, 114
- Cooke J., et al., 2012, *Natur*, 491, 228
- da Cunha E., Charlot S., Elbaz D., 2008, *MNRAS*, 388, 1595
- Dessart, L., Hillier, D. J., Waldman, R., Livne, E., & Blondin, S. 2012, *MNRAS*, 426, L76
- Dessart L., Audit E., Hillier D. J., 2015, *MNRAS*, 449, 4304
- Dexter, J., & Kasen, D. 2013, *ApJ*, 772, 30
- Fassia A., et al., 2000, *MNRAS*, 318, 1093
- Filippenko A. V., et al., 1995, *ApJ*, 450, L11
- Fox O., et al., 2009, *ApJ*, 691, 650
- Fransson C., et al., 2002, *ApJ*, 572, 350
- Fransson C., et al., 2014, *ApJ*, 797, 118
- Fraser M., et al., 2013, *MNRAS*, 433, 1312
- Gal-Yam A., et al., 2007, *ApJ*, 656, 372
- Gal-Yam A., et al., 2009, *Natur*, 462, 624
- Gal-Yam A., 2012, *Sci*, 337, 927
- Gawryszczak A., Guzman J., Plewa T., Kifonidis K., 2010, *A&A*, 521, A38
- Gezari S., et al., 2009, *ApJ*, 690, 1313
- Gonzalez-Delgado R. M., et al., 1994, *ApJ*, 437, 239
- Hatano K., Branch D., Fisher A., Millard J., Baron E., 1999, *ApJS*, 121, 233
- Howell D. A., et al., 2013, *ApJ*, 779, 98
- Hunter D. J., et al., 2009, *A&A*, 508, 371
- Inserra C., et al., 2011, *MNRAS*, 417, 261
- Inserra C., et al., 2013, *ApJ*, 770, 128
- Inserra C., Smartt S. J., 2014, *ApJ*, 796, 87
- Inserra C., et al., 2016a, *MNRAS*, 459, 2721
- Inserra C., et al., 2016b, *arXiv*, *arXiv:1604.01226*
- Inserra C., Bulla M., Sim S. A., Smartt S. J., 2016, *ApJ*, 831, 79
- Jerkstrand A., et al., 2017, *ApJ*, 835, 13
- Jerkstrand A., Smartt S. J., Heger A., 2016, *MNRAS*, 455, 3207
- Jerkstrand A., Ergon M., Smartt S. J., Fransson C., Sollerman J., Taubenberger S., Bersten M., Spyromilio J., 2015, *A&A*, 573, A12
- Kangas T., et al., 2016, *arXiv*, *arXiv:1611.10207*
- Kankare E., et al., 2015, *A&A*, 581, L4
- Kasen, D., & Bildsten, L. 2010, *ApJ*, 717, 245
- Kennicutt R. C., Jr., 1998, *ARA&A*, 36, 189
- Kirshner R. P., Morse J. A., Winkler P. F., Blair W. P., 1989, *ApJ*, 342, 260
- Kozyreva A., Blinnikov S., 2015, *MNRAS*, 454, 4357
- Krühler T., et al., 2015, *A&A*, 581, A125
- Leget P.-F., et al., 2014, *ATel*, 5718,
- Leloudas, G., Schulze, S., Krühler, T., et al. 2015a, *MNRAS*, 449, 917
- Leloudas G., et al., 2015b, *ApJ*, 815, L10
- Levan A. J., Read A. M., Metzger B. D., Wheatley P. J., Tanvir N. R., 2013, *ApJ*, 771, 136
- Liu Y.-Q., Modjaz M., 2016, *arXiv*, *arXiv:1612.07321*
- Lunnan R., et al., 2014, *ApJ*, 787, 138
- Lunnan R., et al., 2016, *ApJ*, 831, 144
- Magnier E. A., et al., 2013, *ApJS*, 205, 20
- Margutti R., et al., 2014, *ApJ*, 780, 21
- Martin J. C., Hamsch F.-J., Margutti R., Tan T. G., Curtis I., Soderberg A., 2015, *AJ*, 149, 9
- Mazzali, P. A., Sullivan, M., Pian, E., Greiner, J., & Kann, D. A. 2016, *MNRAS*,
- McCrum M., et al., 2014, *MNRAS*, 437, 656
- McCrum M., et al., 2015, *MNRAS*, 448, 1206
- Metzger B. D., Vurm I., Hascoët R., Beloborodov A. M., 2014, *MNRAS*, 437, 703
- Milisavljevic D., Fesen R. A., Gerardy C. L., Kirshner R. P., Challis P., 2010, *ApJ*, 709, 1343
- Miller A. A., et al., 2009, *ApJ*, 690, 1303
- Modigliani A., et al., 2010, *SPIE*, 7737, 773728
- Moriya T. J., Blinnikov S. I., Baklanov P. V., Sorokina E. I., Dolgov A. D., 2013, *MNRAS*, 430, 1402
- Moriya T. J., Liu Z.-W., Mackey J., Chen T.-W., Langer N., 2015, *A&A*, 584, L5
- Nicholl M., et al., 2013, *Natur*, 502, 346
- Nicholl M., et al., 2014, *MNRAS*, 444, 2096
- Nicholl M., et al., 2015a, *ApJ*, 807, L18
- Nicholl M., et al., 2015b, *MNRAS*, 452, 3869
- Nicholl M., Smartt S. J., 2016, *MNRAS*, 457, L79
- Nicholl M., et al., 2016a, *ApJ*, 826, 39
- Nicholl M., et al., 2016b, *ApJ*, 828, L18
- Nicholls D. C., Dopita M. A., Sutherland R. S., Kewley L. J., Palay E., 2013, *ApJS*, 207, 21
- Ofek, E. O., Cameron, P. B., Kasliwal, M. M., et al. 2007, *ApJL*, 659, L13
- Ofek E. O., et al., 2013, *ApJ*, 763, 42
- Ofek E. O., et al., 2014, *ApJ*, 781, 42

- Osterbrock D. E., Ferland G. J., 2006, *agna.book*,  
 Papadopoulos A., et al., 2015, *MNRAS*, 449, 1215  
 Pastorello A., et al., 2008, *MNRAS*, 389, 131  
 Pastorello A., et al., 2010, *ApJ*, 724, L16  
 Pastorello A., et al., 2015, *MNRAS*, 449, 1954  
 Patat F., et al., 2001, *ApJ*, 555, 900  
 Pettini M., Pagel B. E. J., 2004, *MNRAS*, 348, L59  
 Perley D. A., et al., 2016, *ApJ*, 830, 13  
 Pignata G., et al., 2004, *MNRAS*, 355, 178  
 Poole T. S., et al., 2008, *MNRAS*, 383, 627  
 Pozzo M., Meikle W. P. S., Fassia A., Geballe T., Lundqvist  
 P., Chugai N. N., Sollerman J., 2004, *MNRAS*, 352, 457  
 Prajs S., et al., 2017, *MNRAS*, 464, 3568  
 Quimby R. M., Aldering G., Wheeler J. C., Höflich P., Ak-  
 erlof C. W., Rykoff E. S., 2007, *ApJ*, 668, L99  
 Quimby R. M., et al., 2011, *Natur*, 474, 487  
 Quimby, R. M., Yuan, F., Akerlof, C., & Wheeler, J. C.  
 2013, *MNRAS*, 431, 912  
 Reynolds S. P., 2008, *ARA&A*, 46, 89  
 Schlafly E. F., Finkbeiner D. P., 2011, *ApJ*, 737, 103  
 Schlafly E. F., et al., 2012, *ApJ*, 756, 158  
 Schulze S., et al., 2016, *arXiv*, arXiv:1612.05978  
 Scovacricchi D., Nichol R. C., Bacon D., Sullivan M., Prajs  
 S., 2016, *MNRAS*, 456, 1700  
 Smartt S. J., et al., 2015, *A&A*, 579, A40  
 Smith, N., Li, W., Foley, R. J., et al. 2007, *ApJ*, 666, 1116  
 Smith N., McCray R., 2007, *ApJ*, 671, L17  
 Smith N., et al., 2009, *ApJ*, 695, 1334  
 Smith N., Mauerhan J. C., Silverman J. M., Gane-  
 shalingam M., Filippenko A. V., Cenko S. B., Clubb K. I.,  
 Kandrashoff M. T., 2012, *MNRAS*, 426, 1905  
 Smith M., et al., 2016, *ApJ*, 818, L8  
 Stritzinger M., et al., 2002, *AJ*, 124, 2100  
 Stritzinger M., et al., 2012, *ApJ*, 756, 173  
 Taubenberger S., et al., 2009, *MNRAS*, 397, 677  
 Tonry J. L., et al., 2012, *ApJ*, 750, 99  
 Valenti S., et al., 2011, *MNRAS*, 416, 3138  
 Vishniac E. T., 1994, *ApJ*, 428, 186  
 Vreeswijk P. M., et al., 2014, *ApJ*, 797, 24  
 Vreeswijk P. M., et al., 2017, *ApJ*, 835, 58  
 Weiler K. W., van Dyk S. D., Pringle J. E., Panagia N.,  
 1992, *ApJ*, 399, 672  
 Wheeler J. C., Benetti S., 2000, *asqu.book*, 451  
 Woosley, S. E. 2010, *ApJL*, 719, L204  
 Woosley, S. E., Blinnikov, S., & Heger, A. 2007, *Nature*,  
 450, 390  
 Yan L., et al., 2015, *ApJ*, 814, 108  
 Yaron, O., & Gal-Yam, A. 2012, *PASP*, 124, 668  
 Young D. R., et al., 2010, *A&A*, 512, A70

**APPENDIX A: TABLES**

**Table A1.** Journal of spectroscopic observations of LSQ14an.

Date yy/mm/dd	MJD	Phase* (days)	Range (observer-frame) (Å)	Range (rest-frame) (Å)	Resolution (Å)	Instrumental Configuration
14/01/02	56660.31	55.0	3665–9285	3150–7980	18	NTT+EFOSC2+gm13
14/01/08	56665.81	59.7	3360–10010	2890–8600	14/13	NTT+EFOSC2+gm11/16
14/01/29	56687.28	78.2	3095–24785	2660–21300	1.0/1.1/3.3	VLT+XSHOOTER+UV/OPT/NIR
14/01/29	56687.31	78.2	3360–10010	2890–8600	14/13	NTT+EFOSC2+gm11/16
14/02/21	56710.70	98.3	3360–10010	2890–8600	14/13	NTT+EFOSC2+gm11/16
14/03/08	56725.16	110.7	3665–9285	3150–7980	18	NTT+EFOSC2+gm13
14/04/22	56770.02	149.3	3665–9285	3150–7980	18	NTT+EFOSC2+gm13
14/06/21	56829.09	200.0	3095–24785	2660–21300	1.0/1.1/3.3	VLT+XSHOOTER+UV/OPT/NIR
14/08/21	56890.49	252.8	3665–9285	3150–7980	18	NTT+EFOSC2+gm13

\* Phase with respect to the assumed maximum.

**Table A2.** *gri* light curves of LSQ14an.

Date yy/mm/dd	MJD	Phase* (days)	<i>g</i>	<i>r</i>	<i>i</i>	Instrument <sup>†</sup>
14/01/02	56659.28	54.1		18.60 (0.08)		LSQ
14/01/04	56661.28	55.8		18.62 (0.08)		LSQ
14/01/06	56663.30	57.6		18.64 (0.08)		LSQ
14/01/14	56672.19	65.2	18.93 (0.13)	18.92 (0.19)	18.89 (0.17)	LT
14/01/28	56686.23	77.3	19.15 (0.11)	19.12 (0.11)	18.99 (0.12)	LT
14/02/01	56689.24	79.9	19.12 (0.13)	19.13 (0.14)	18.99 (0.12)	NTT
14/02/07	56696.24	85.9	19.11 (0.17)	19.14 (0.11)	19.09 (0.12)	LT
14/02/13	56702.11	90.9	19.19 (0.16)	19.13 (0.14)	19.24 (0.15)	LT
14/03/12	56729.03	114.1	19.35 (0.18)	19.36 (0.14)	19.40 (0.16)	LT
14/04/21	56769.17	148.6	19.68 (0.12)	20.02 (0.14)	19.83 (0.12)	NTT
14/05/16	56793.93	169.8	20.06 (0.14)	20.13 (0.14)	20.13 (0.14)	LT
14/05/19	56796.89	172.4	20.03 (0.12)	20.19 (0.12)	20.18 (0.13)	LT
14/05/23	56800.88	175.8	20.11 (0.12)	20.32 (0.12)	20.45 (0.14)	LT
14/05/27	56804.91	179.3	20.07 (0.11)	20.41 (0.12)	20.49 (0.14)	LT
14/06/03	56811.96	185.3	20.07 (0.12)	20.46 (0.13)	20.55 (0.15)	LT
14/08/25	56894.99	256.7	21.55 (0.11)	21.46 (0.14)	21.60 (0.23)	NTT
14/12/29	57021.30	365.2	22.50 (0.13)	22.49 (0.15)	22.82 (0.19)	NTT
15/01/26	57049.30	389.3	22.98 (0.15)	23.03 (0.19)	23.18 (0.25)	NTT
15/02/17	57071.20	408.1	23.64 (0.19)	23.83 (0.22)	24.06 (0.32)	NTT

\* Phase with respect to the assumed maximum

† LSQ = La Silla Quest; LT = Liverpool Telescope + IO:O; NTT = ESO NTT + EFOSC2

**Table A3.** *BVRJHK* light curve of LSQ14an measured after template subtraction (from 2015-04-17) using HOTPANTS

Date yy/mm/dd	MJD	Phase* (days)	<i>B</i>	<i>V</i>	<i>R</i>	Instrument <sup>†</sup>
14/01/02	56660.31	55.0		19.00 (0.10)		NTT
14/01/08	56665.81	59.7		19.10 (0.13)		NTT
14/01/29	56687.28	78.2		19.49 (0.09)		NTT
14/02/21	56710.70	98.3		19.63 (0.12)		NTT
14/03/08	56725.16	110.7		19.68 (0.18)		NTT
14/04/22	56770.02	149.3		20.13 (0.14)		NTT
14/08/25	56894.99	256.7	21.85 (0.10)	21.50 (0.03)	21.31 (0.09)	NTT
14/12/29	57021.30	365.2	22.72 (0.15)	22.47 (0.06)	22.39 (0.10)	NTT
15/01/26	57049.30	389.3	23.21 (0.13)	22.99 (0.10)	22.93 (0.15)	NTT
15/02/17	57071.20	408.1	23.95 (0.20)	23.46 (0.16)	23.84 (0.19)	NTT
Date yy/mm/dd	MJD	Phase* (days)	<i>J</i>	<i>H</i>	<i>K</i>	Instrument <sup>†</sup>
14/04/23	56771.04	150.16	19.93 (0.12)	20.04 (0.12)	20.08 (0.20)	NTT

\* Phase with respect to the assumed maximum † NTT = ESO NTT + EFOSC2

**Table A4.** Magnitudes in *griBVR* of the local sequence stars in the field of LSQ14an.

RA	Dec	<i>g</i>	<i>r</i>	<i>i</i>
12:53:52.095	-29:32:20.47	16.36 (0.05)	15.92 (0.05)	15.74 (0.05)
12:53:49.806	-29:31:58.25	17.61 (0.07)	16.37 (0.05)	15.56 (0.05)
12:53:49.136	-29:32:15.38	17.11 (0.05)	16.61 (0.04)	16.38 (0.04)
12:53:48.019	-29:31:37.48	17.78 (0.07)	16.51 (0.04)	15.67 (0.05)
12:53:44.110	-29:32:04.44	16.76 (0.06)	16.32 (0.05)	16.13 (0.05)
12:53:52.622	-29:29:23.17	16.38 (0.05)	15.93 (0.04)	15.75 (0.05)
12:53:52.300	-29:29:36.81	16.22 (0.05)	15.59 (0.04)	15.27 (0.05)
12:53:44.595	-29:29:30.65	17.76 (0.08)	17.22 (0.04)	16.98 (0.03)
RA	Dec	<i>B</i>	<i>V</i>	<i>R</i>
12:53:52.672	-29:30:12.65	20.56 (0.13)	19.60 (0.08)	18.95 (0.07)
12:53:40.273	-29:30:15.14	19.35 (0.12)	18.52 (0.10)	18.02 (0.10)
12:53:40.049	-29:30:54.67	18.93 (0.11)	18.22 (0.09)	17.76 (0.09)
12:53:44.735	-29:31:10.91	20.88 (0.12)	20.41 (0.10)	20.04 (0.08)
12:53:47.986	-29:31:37.54	18.62 (0.14)	17.16 (0.10)	16.19 (0.08)
12:53:49.102	-29:32:15.83	17.67 (0.13)	16.95 (0.10)	16.48 (0.08)
12:53:55.383	-29:32:04.58	20.59 (0.14)	20.00 (0.10)	19.62 (0.09)

**Table A5.** Log of X-ray observations of LSQ14an and SN 2015bn

LSQ14an					
Date yy/mm/dd	MJD	Phase* (days)	Exposure (s)	$F_X$ (erg cm <sup>-2</sup> s <sup>-1</sup> )	$L_X$ (erg s <sup>-1</sup> )
14/03/24	56740.52	123.9	9000	<1.0e-13	<7.0e42
14/07/03	56841.04	210.3	5400	<2.6e-13	<1.8e43
14/12/07	56998.41	345.5	7600	<1.8e-13	<1.3e43
16/08/16	57616.01	876.3	9500	<5.6e-14	<3.9e42
Stacked					
	56859.70 <sup>†</sup>	226.3 <sup>†</sup>	22000	<4.1e-14	<2.9e42
SN 2015bn					
Date yy/mm/dd	MJD	Phase* (days)	Exposure (s)	$F_X$ (erg cm <sup>-2</sup> s <sup>-1</sup> )	$L_X$ (erg s <sup>-1</sup> )
15/02/19	57072.29	-26.7	6000	<2.7e-13	<8.6e42
15/04/19	57131.26	26.3	6700	<8.7e-14	<2.8e42
16/01/03	57390.02	258.6	8700	<3.6e-14	<1.1e42
16/03/02	57449.07	311.6	2800	<3.7e-13	<1.2e43
16/07/04	57573.21	423.1	3900	<3.2e-13	<1.0e43
16/12/16	57738.87	571.8	8900	<3.6e-14	<1.1e42
16/12/31	57753.58	585.0	9940	<3.0e-14	<9.5e41
17/01/19	57772.01	601.5	9920	<3.2e-14	<1.0e42
17/02/06	57788.56	616.5	8950	<3.5e-14	<1.1e42
17/02/17	57801.71	628.3	8900	<5.0e-14	<1.6e42
Stacked					
	57323.17 <sup>‡</sup>	198.6 <sup>‡</sup>	28000	<2.4e-14	<7.6e41

\* Phase with respect to the assumed maximum

† Average date/MJD between the first three exposures †† Average date/MJD between the first four exposures

Line	$\lambda$ ( $\text{\AA}$ )	Observed flux ( $10^{-17}$ erg s $^{-1}$ cm $^{-2}$ )	Error ( $10^{-17}$ erg s $^{-1}$ cm $^{-2}$ )	RMS ( $10^{-17}$ erg s $^{-1}$ cm $^{-2}$ )	EW ( $\text{\AA}$ )	FWHM ( $\text{\AA}$ )
He I	3188	3.92	1.98	0.52	0.55	1.88
He I	3704	1.43	0.91	0.29	0.20	1.29
[O II]	3726	65.65	2.39	0.42	8.94	1.64
[O II]	3729	92.03	2.69	0.42	12.46	1.63
H12	3750	1.22	1.03	0.34	0.16	1.24
H11	3770	1.70	0.98	0.32	0.23	1.24
H10	3798	2.89	0.73	0.23	0.43	1.28
H9	3835	6.79	1.44	0.37	0.91	1.76
[Ne III]	3868	34.63	1.76	0.37	4.51	1.54
He I+H8	3889	12.27	0.96	0.25	1.65	1.50
[Ne III]	3967	7.40	0.93	0.27	0.94	1.36
H7	3969	9.65	0.95	0.27	1.23	1.36
[N II]+He I	4026	0.82	0.44	0.16	0.12	1.00
H $\delta$	4102	17.14	0.90	0.22	2.88	1.54
H $\gamma$	4340	35.26	1.29	0.24	5.16	1.86
[O III]	4363	5.43	1.03	0.30	0.82	1.86
He I	4471	4.35	0.87	0.20	0.86	2.44
H $\beta$	4861	68.47	2.06	0.24	22.68	2.03
[O III]	4959	129.20	3.10	0.37	23.43	1.96
[O III]	5007	345.70	4.65	0.34	68.15	2.03
He I	5876	7.75	0.57	0.11	4.90	2.25
[O I]	6300	4.88	0.96	0.19	2.12	2.90
[S III]	6312	1.56	0.85	0.18	0.65	2.90
[O I]	6364	1.36	0.43	0.11	0.51	2.15
[N II]	6548	2.11	1.61	0.35	1.01	2.73
H $\alpha$	6563	234.30	3.37	0.20	105.19	2.73
[N II]	6584	5.25	0.54	0.11	2.67	2.73
He I	6678	1.96	0.33	0.07	1.43	2.72
[S II]	6717	13.83	0.78	0.11	10.87	2.74
[S II]	6731	10.21	0.68	0.11	8.19	2.74
He I	7065	1.87	0.41	0.09	1.36	2.75
[Ar III]	7136	4.89	1.16	0.22	3.10	2.82
[O II]	7320	0.95	0.51	0.12	0.19	2.47
[O II]	7330	1.05	0.52	0.12	0.21	2.47
[S III]	9069	15.74	1.05	0.15	10.96	3.28
[S III]	9532	36.59	2.86	0.31	24.45	3.42

**Table A6.** Observed emission line measurements of the host galaxy of LSQ14an from the spectrum taken by VLT + Xshooter at +200.0d (no applying any reddening corrections). The EW of each line is a minimum, not a real line strength due to the SN contamination of the continuum.