arXiv:1702.08890v2 [math.AG] 10 Dec 2017 [arXiv:1702.08890v2 \[math.AG\] 10 Dec 2017](http://arxiv.org/abs/1702.08890v2)

DIMENSION OF THE SPACE OF CONICS ON FANO HYPERSURFACES

KATSUHISA FURUKAWA

ABSTRACT. R. Beheshti showed that, for a smooth Fano hypersurface X of degree ≤ 8 over the complex number field \mathbb{C} , the dimension of the space of lines lying in X is equal to the expected dimension.

We study the space of conics on X . In this case, if X contains some linear subvariety, then the dimension of the space can be larger than the expected dimension.

In this paper, we show that, for a smooth Fano hypersurface X of degree ≤ 6 over $\mathbb C$, and for an irreducible component R of the space of conics lying in X, if the 2-plane spanned by a general conic of R is not contained in X , then the dimension of R is equal to the expected dimension.

1. INTRODUCTION

Let $X \subset \mathbb{P}^n$ be a hypersurface of degree d over the complex number field \mathbb{C} . We define $R_e(X)$ to be the space of smooth rational curves of degree e in \mathbb{P}^n lying in X, which is an open subscheme of the Hilbert scheme Hilb $e^{t+1}(X)$. The number

$$
(n+1-d)e+n-4
$$

is called the *expected dimension* of $R_e(X)$, where the dimension of $R_e(X)$ at C is greater than or equal to this number if there exists $C \in R_e(X)$ such that X is smooth along C.

The space of rational curves on a *general* Fano hypersurface have been studied by many authors ([\[5\]](#page-13-0), [\[8\]](#page-13-1), [\[9\]](#page-13-2), [\[14\]](#page-14-0), [\[4\]](#page-13-3), [\[12\]](#page-14-1) in characteristic zero; [\[10,](#page-13-4) V, §4], [\[7\]](#page-13-5) in any characteristic). At least for $e = 1, 2$, it is well known that $R_e(X)$ has the expected dimension if X is a general Fano hypersurface.

On the other hand, it is difficult to know about $R_e(X)$ for *any smooth* $X \subset \mathbb{P}^n$. For n exponentially large in d , T. D. Browning and P. Vishe [\[5\]](#page-13-0) showed that the space of rational curves of any degree e on smooth X has the expected dimension.

For any n in the case of $e = 1$, as an answer of the question which was asked by O. Debarre and J. de Jong independently, R. Beheshti [\[2\]](#page-13-6) showed that $R_1(X)$ has the expected dimension if $X \subset \mathbb{P}^n$ is a smooth Fano hypersurface of degree $d \leq 6$.

²⁰¹⁰ Mathematics Subject Classification. 14C05, 14J70, 14N05.

Key words and phrases. space of rational curves, Fano hypersurface, expected dimension.

J. M. Landsberg and C. Robles [\[11\]](#page-13-7) gave another proof for the same degree $d \leq 6$. Beheshti [\[3\]](#page-13-8) later showed the same statement for $d \leq 8$.

In the case of $e = 2$, A. Collino, J. P. Murre, G. E. Welters [\[6\]](#page-13-9) studied $R_2(X)$ for a smooth quadric 3-fold $X \subset \mathbb{P}^4$; in this case, $R_2(X)$ has the expected dimension. Note that $\text{Hilb}^{2t+1}(X)$ is connected for any smooth hypersurface $X \subset \mathbb{P}^n$ if the expected dimension is positive [\[7,](#page-13-5) Proposition 5.6].

In this paper, we study the dimension of $R_2(X)$ for a smooth Fano hypersurface of degree $d \leq 6$. Our main result is the following.

Theorem 1.1. Let $X \subset \mathbb{P}^n$ be a smooth Fano hypersurface of degree $d \leq 6$ over $\mathbb{C}.$ Let $R \neq \emptyset$ be an irreducible component of $R_2(X)$ such that

$$
(1) \t\langle C \rangle \not\subset X \text{ for general } C \in R,
$$

where $\langle C \rangle = \mathbb{P}^2 \subset \mathbb{P}^n$ *is the* 2*-plane spanned by* C*.* Then $\dim(R)$ *is equal to the* $expected\ dimension\ 3n-2d-2.$

The dimension of $R_2(X)$ can be greater than the expected one when X contains certain linear varieties (see [Example 3.17\)](#page-13-10); this is the reason why we assume the condition [\(1\).](#page-1-0) The statement of [Theorem 1.1](#page-1-1) does not hold for $d \geq 10$ (see [Example 3.18\)](#page-13-11); thus it may need some conditions stronger than [\(1\)](#page-1-0) for larger d.

The paper is organized as follows. We assume that $\dim R$ is greater than the expected dimension, and take $Y := Loc(R) \subset X$ to be the locus swept out by conics of R. The codimension of Y is ≥ 2 in X due to a result of Beheshti [\[3\]](#page-13-8) (see [Remark 2.2\)](#page-2-0). Then it is sufficient to investigate the case when $(d, \dim Y) =$ $(6, n-3)$. In §[2,](#page-1-2) we consider the linear subvariety $\langle Y \rangle \subset \mathbb{P}^n$ spanned by Y, and show that the codimension of $\langle Y \rangle$ is ≤ 1 in \mathbb{P}^n by using projective techniques [\(Proposition 2.6\)](#page-4-0). Let $\mathbb{T}_x X \subset \mathbb{P}^n$ be the embedded tangent space to X at x. In §[3,](#page-6-0) considering the subset $R_x^* \subset R$ consisting of conics C such that $x \in C \subset \mathbb{T}_x X$, we show that $\langle Loc(R_x^*)\rangle$ is an $(n-3)$ -plane [\(Proposition 3.6\)](#page-7-0), and show that $Loc(R_x^*)$ is a quadric hypersurface in $\langle Loc(R_x^{\star}) \rangle = \mathbb{P}^{n-3}$ [\(Corollary 3.9\)](#page-9-0). In particular, our problem is reduced to the case $n = d = 6$. Using such quadrics, we give the proof of [Theorem 1.1](#page-1-1) step by step.

Acknowledgments. The author would like to thank Professor Hiromichi Takagi for many helpful comments and advice. The author was supported by the Grantin-Aid for JSPS fellows, No. 16J00404.

2. The locus swept out by conics

We use the following notations. For a Fano hypersurface $X \subset \mathbb{P}^n$ of degree $d \leq 6$, we take an irreducible component $R \neq \emptyset$ of $R_2(X)$ such that $\langle C \rangle \not\subset X$ for general $C \in R_2(X)$. We denote by \overline{R} the closure in Hilb^{2t+1}(X). Let

$$
\mathcal{U} := \{ (C, x) \in R \times X \mid x \in C \}
$$

be the universal family of R, and let $\pi : \mathfrak{U} \to R$ and $ev : \mathfrak{U} \to X$ be the first and second projections. For a subset $A \subset R$, we write $\mathcal{U}_A := \pi^{-1}(A)$ and $\text{Loc}(A) :=$ $\overline{\mathrm{ev}(\mathfrak{U}_A)} = \overline{\bigcup_{C \in A} C} \subset X.$

We write $R_x \subset R$ to be the set of $C \in R$ passing through $x \in X$, and write $R_{xy} = R_x \cap R_y \subset R$, the set of $C \in R$ passing through $x, y \in X$.

We set $\langle S_1 \cdot \cdot \cdot S_m \rangle \subset \mathbb{P}^n$ to be the linear variety spanned by subsets $S_1, \ldots, S_m \subset$ \mathbb{P}^n . For example, $\langle xy \rangle \subset \mathbb{P}^n$ is the line passing through $x, y \in \mathbb{P}^n$, and $\langle C \rangle \subset \mathbb{P}^n$ is the 2-plane spanned by C for a conic $C \subset \mathbb{P}^n$.

The condition [\(1\)](#page-1-0) in [Theorem 1.1](#page-1-1) gives the following basic property for R_{xy} .

Lemma 2.1. Let R'_{xy} be an irreducible component of R_{xy} such that $\langle C \rangle \not\subset X$ for $general C \in R'_{xy}$. Then

$$
\mathfrak{U}_{R_{xy}'} \to \mathrm{Loc}(R_{xy}') \subset X
$$

is generically finite; moreover a fiber at $z \in Loc(R'_{xy})$ *is of positive dimension only* $if \langle xyz \rangle \subset X$. In particular, dim $Loc(R'_{xy}) = dim(R'_{xy}) + 1$.

Proof. If dim $R'_{xy} = 0$, then the assertion follows immediately. Assume dim $R'_{xy} \geqslant$ 1. Then dim $\text{Loc}(R'_{xy}) \geq 2$. Take $C \in R'_{xy}$ and $z \in C$ such that $M := \langle xyz \rangle =$ $\langle C \rangle \not\subset X$. Then $\check{M} \cap X$ is a union of finitely many curves. Since any conic $\tilde{C} \in \pi(\mathrm{ev}^{-1}(z) \cap \mathfrak{U}_{R'_{xy}})$ satisfies $\tilde{C} \subset M$, it coincides with a component of $M \cap X$. Hence the fiber $ev^{-1}(z) \cap \mathcal{U}_{R'_{xy}}$ must be a finite set.

We set $Y := \text{Loc}(R) \subset X$, the locus swept out by conics $C \in R$, which is non-linear since $\langle C \rangle \not\subset X$ for general C. Let us consider the projection

(2)
$$
ev^{(2)}: \mathcal{U} \times_R \mathcal{U} \simeq \{ (C, x, y) \in R \times Y \times Y \mid x, y \in C \} \to Y \times Y
$$

whose fiber at $(x, y) \in Y \times Y$ is isomorphic to R_{xy} . Considering $\mathcal{U} \times_R \mathcal{U} \to R$, we have dim $\mathcal{U} \times_R \mathcal{U} = r + 2$. Note that $ev^{(2)}$ is dominant if and only if $Loc(R_x) = Y$ holds for general $x \in Y$.

Remark 2.2. Assume that dim R is greater than the expected dimension. Then the locus Y is much smaller than X. More precisely, by a result of R. Beheshti [\[3,](#page-13-8) Theorem 3.2(b)], it holds that dim $Y \leq n-3$.

We immediately have dim $Y \leq n-2$; this is because if $X = Y = \text{Loc}(R)$ (i.e., $\dim Y = n - 1$) in characteristic zero, then R has a free curve C and then R must have expected dimension. Beheshti's result gives the inequity which is sharper than this.

Lemma 2.3. If $d \leq 6$ and $r = \dim R$ is greater than the expected dimension, then $(d, \dim Y) = (6, n-3)$.

Proof. We have dim $Y \leq n-3$ due to Beheshti's result as we saw in [Remark 2.2.](#page-2-0) Let $(C, x, y) \in \mathcal{U} \times_R \mathcal{U}$ be general. Since $Loc(R_{xy}) \subset Y$, it follows from the morphism [\(2\)](#page-2-1) and [Lemma 2.1](#page-2-2) that $(r + 2 - 2 \dim(Y)) + 1 \leq \dim(Y)$. Hence $r+3-3\dim(Y) \leq 0$. By assumption, $r \geq 3n-2d-1$. Thus

$$
3n - 2d + 2 - 3\dim(Y) \leq 0.
$$

Since dim $Y \leq n - 3$, we have $11 - 2d \leq 0$; hence $d = 6$. Therefore $n - 10/3 \leq$ $dim(Y)$; hence $dim(Y) = n-3$.

By the above lemma, let us study the case $(d, \dim Y) = (6, n-3)$, and assume that $r := \dim R$ is greater than the expected dimension, that is to say,

$$
r \geqslant 3n - 13.
$$

Lemma 2.4. $ev^{(2)}$ *is dominant. Therefore* $Loc(R_x) = Y$ *for general* $x \in Y$ *.*

Proof. For general $(x, y) \in im(ev^{(2)})$, we have dim $R_{xy} = (r + 2) - dim(im(ev^{(2)}))$, which implies dim $R_{xy} + \dim(\text{im}(ev^{(2)})) = (r+2) \ge 3n - 11$.

Suppose that $ev^{(2)}$ is not dominant, that is, dim(im($ev^{(2)}$)) < 2(n – 3). Then $Loc(R_{xy}) \neq Y$, which implies dim $R_{xy} + 1 <$ dim $Y = n - 3$ because of [Lemma 2.1.](#page-2-2) Then dim R_{x} +dim(im(ev⁽²⁾)) $\leqslant (n-5)+(2(n-3)-1) = 3n-12$, a contradiction. \Box

Note that for any $x, y \in Y$

(4)
$$
\dim R_{xy} \geqslant r + 2 - 2 \dim(Y) \geqslant n - 5.
$$

Hereafter we will use several projective techniques in order to study Y .

Remark 2.5. We sometimes consider the *Gauss map* of a variety $Z \subset \mathbb{P}^n$, which is a rational map

$$
\gamma_Z: Z \dashrightarrow \mathbb{G}(\dim Z, \mathbb{P}^n),
$$

sending a smooth point $x \in Z$ to the embedded tangent space $\mathbb{T}_x Z \subset \mathbb{P}^n$ at x. A general fiber of γ_Z is a linear variety of \mathbb{P}^n in characteristic zero (in particular, irreducible). The map γ_Z is a finite morphism if Z is smooth. See [\[15,](#page-14-2) I, §2].

We write $(\mathbb{P}^n)^{\vee} = \mathbb{G}(n-1,\mathbb{P}^n)$, the space of hyperplanes of \mathbb{P}^n . For a linear subvariety $A \subset \mathbb{P}^n$, we denote by $A^* \subset (\mathbb{P}^n)^{\vee}$ the set of hyperplanes containing A. In addition, for a subset $B \subset \mathbb{P}^n$, we set $Cone_A(B) := \overline{\bigcup_{x \in B} \langle A, x \rangle} \subset \mathbb{P}^n$, the cone of B with vertex A.

Let us consider the linear variety $\langle Y \rangle \subset \mathbb{P}^n$ spanned by the locus $Y \subset X$. The following proposition states $\langle Y \rangle$ cannot be so small in \mathbb{P}^n .

Proposition 2.6. *Assume* $(d, \dim Y) = (6, n-3)$ *and the formula* [\(3\)](#page-3-0). *Then* $\langle Y \rangle$ *is of dimension* $\geq n-1$ *.*

Proof. Since Y is non-linear, we immediately have $\dim(Y) > n-3$. Now assume that $\langle Y \rangle$ is an $(n-2)$ -plane. We need to show two claims.

Claim 2.7. *It holds that* $Y = \langle Y \rangle \cap X$ *and* dim(Sing Y) ≤ 1 *. In particular,* $Y \subset \langle Y \rangle = \mathbb{P}^{n-2}$ *is a hypersurface whose degree is equal to* deg $X = d = 6$ *.*

Proof. For $x \in \langle Y \rangle \cap X$, it holds that $x \in Sing(\langle Y \rangle \cap X)$ if and only if $\langle Y \rangle \subset \mathbb{T}_x X$. It means that $\gamma_X(\text{Sing}(\langle Y \rangle \cap X)) \subset \langle Y \rangle^*$ for the Gauss map $\gamma_X : X \to (\mathbb{P}^n)^{\vee}$, where $\langle Y \rangle^* \subset (\mathbb{P}^n)^{\vee}$ is the set of hyperplanes containing $\langle Y \rangle$. Since X is smooth, γ_X is a finite morphism. Since $\dim(Y)^* = 1$, we have $\dim(\text{Sing}(\langle Y \rangle \cap X)) \leq 1$. If there exists an irreducible component $Y' \subset \langle Y \rangle \cap X \subset \langle Y \rangle = \mathbb{P}^{n-2}$ such that $Y' \neq Y$, then we have $\dim(Y' \cap Y) \geq n-4 \geq 2$, which is a contradiction since $Y' \cap Y \subset \text{Sing}(\langle Y \rangle \cap X)$. Thus $\langle Y \rangle \cap X = Y$.

Claim 2.8. *A general* $C \in R$ *satisfies* $C \cap \text{Sing } Y \neq \emptyset$.

Proof. Suppose that a general conic $C \in R$ satisfies $C \cap \text{Sing } Y = \emptyset$. Since $Y =$ Loc(R) and the characteristic is zero, C is free in Y. Then $R_2(Y)$ is smooth at C and has the expected dimension $3(n-2)-2 \deg Y - 2$. This contradicts that $R \subset R_2(Y)$ is of dimension $> 3n - 2d - 2$, where deg $Y = \deg X = d$ because of [Claim 2.7.](#page-4-1)

From [Claim 2.8,](#page-4-2) we may indeed assume

$$
(5) \tC \cap S \neq \emptyset
$$

for an irreducible component $S \subset \text{Sing } Y$. Note that $\dim S \leq 1$.

First we consider the case $n > 6$. Let $C_0 \in R$ be a general conic such that $\langle C_0 \rangle \not\subset X$, and take $x, y \in C_0 \setminus S$ be general. From [\(4\),](#page-3-1) we have dim $R_{xy} \geq$ $n-5 \geqslant 2$. Let $R'_{xy} \subset R_{xy}$ be an irreducible component containing C_0 , and take $(R'_{xy})^{\circ} \subset R'_{xy}$ to be the set of C satisfying $\langle C \rangle \not\subset X$. Then, for any $C \in (R'_{xy})^{\circ}$ and $s \in C \cap S$, we have $\langle xyz \rangle = \langle C \rangle \not\subset X$. From [Lemma 2.1,](#page-2-2) for the morphism

$$
\widetilde{\operatorname{ev}}:=\operatorname{ev}|_{\mathfrak{U}_{(R_{xy}')^\circ}}:\mathfrak{U}_{(R_{xy}')^\circ}\rightarrow \operatorname{Loc}(R_{xy}'),
$$

the preimage $\widetilde{\text{ev}}^{-1}(S)$ is of dimension ≤ 1 . Since $\dim R'_{xy} \geq 2$, we have $\pi(\widetilde{\text{ev}}^{-1}(S)) \neq$ R'_{xy} , which means that $C \cap S = \emptyset$ for general $C \in R'_{xy}$, a contradiction.

Next we consider the case $n = 6$, and complete the proof in the following four steps. Note that $\langle Y \rangle = \mathbb{P}^4 \subset \mathbb{P}^6$.

Step 1. We show that $S \not\subset \langle C_0 \rangle$ for general $C_0 \in R$, and also show that $S \not\subset M$ for a general 3-plane $M \subset \langle Y \rangle$ containing C_0 .

Suppose $S \subset \langle C_0 \rangle$ for general $C_0 \in R$, and take $x, y \in Y$ be general points. We can assume $y \notin \text{Cone}_x(S)$. Since dim $R_{xy} \geqslant n-5 \geqslant 1$, taking general $C, C' \in R_{xy}$ with $C \neq C'$, we have $S \subset \langle C \rangle \cap \langle C' \rangle = \langle xy \rangle$, a contradiction.

If $S \subset M$ for a general 3-plane M containing C_0 , then we can also take another general $\tilde{M} \neq M$, and then $\langle C_0 \rangle = M \cap \tilde{M} \supset S$, a contradiction.

Step 2. We consider
$$
\langle Y \rangle^{\vee} := \mathbb{G}(3, \langle Y \rangle)
$$
, the set of 3-planes in $\langle Y \rangle = \mathbb{P}^4$. Let

$$
W = \{ (C, M) \in R \times \langle Y \rangle^{\vee} \mid C \subset M \},
$$

which is a \mathbb{P}^1 -bundle over R; in particular, dim $W \geq 6$. Let $\text{pr}_2 : W \to \langle Y \rangle^{\vee}$ be the projection to the second factor.

For general $(C_0, M) \in W$, take R^M to be an irreducible component of $R \cap R_2(M)$ containing C_0 . We may assume that a general conic $C \in R^{\tilde{M}}$ satisfies $\langle C \rangle \nsubseteq X$. Since $R \cap R_2(M) \simeq \text{pr}_2^{-1}(M)$, we can assume dim $R^M \geq 6 - \dim(\text{pr}_2(W))$. We set the surface

$$
Y^M := \text{Loc}(R^M) \subset Y \cap M.
$$

Step 3. Assume dim $pr_2(W) \le 3$. Then we have dim $R^M \ge 3$, which implies that $R^M \to \mathbb{G}(2,M) = (\mathbb{P}^3)^{\vee} : C \mapsto \langle C \rangle$ is dominant. From Step 1, $S \cap M$ is a set of finite points. Thus $L \cap S = \emptyset$ for a general 2-plane $L \subset M$. Taking a general $C \in R^{\tilde{M}}$ such that $\langle C \rangle = L$, we find that $C \cap S = \emptyset$, which contradicts the condition [\(5\).](#page-4-3)

Step 4. Assume dim $pr_2(W) = 4$, that is, $pr_2(W) = \langle Y \rangle^{\vee}$. For general $(C_0, M) \in$ W, since M is general in $\langle Y \rangle^{\vee}$, $Y \cap M$ is irreducible. Then $Y^M = Y \cap M \subset M = \mathbb{P}^3$, which is a surface whose degree is equal to deg $Y = 6$.

Since $S \cap M$ is a finite set, we may assume that there exists $s \in S \cap M$ such that $s \in C$ for general $C \in R^M$. This implies that $R^M \subset R_s$. Considering $\mathcal{U}_{R^M} \to Y^M$, we find that $\dim(R^M \cap R_x) \geq 1$ for general $x \in Y^M$. Since Y^M is surface,

$$
Y^M = \text{Loc}(R^M \cap R_x) = \text{Loc}(R^M \cap R_{xs}).
$$

For general $C \in \mathbb{R}^M$, we show that $\langle C \rangle \cap Y^M$ is scheme-theoretically equal to C, as follows. Write $({\langle C \rangle \cap Y^M})_{red} = C \cup \bigcup_i E_i$ with the irreducible components E_i's. Take a general $x \in C \setminus \bigcup_i E_i$ and take a general $y \in E_1 \setminus (C \cup \langle xs \rangle)$. Taking the closure

$$
\overline{R^M \cap R_{xs}} \subset \text{Hilb}^{2t+1}(X)
$$

and consider the surjective map $\mathcal{U}_{\overline{R^M \cap R_{xs}}} \to Y^M$, we find $\tilde{C} \in \overline{R^M \cap R_{xs}}$ such that $x, y, s \in \tilde{C}$. Then $\langle \tilde{C} \rangle = \langle xyz \rangle = \langle C \rangle$, which implies that $\tilde{C} \subset Y^M \cap \langle C \rangle$. By the choice of x, it follows $C \subset \tilde{C}$. Then $C = \tilde{C}$, which implies $y \in C$, a contradiction.

Thus $(\langle C \rangle \cap Y^M)_{red} = C$. Suppose that $\langle C \rangle \cap Y^M$ is non-reduced, which means that C is a contact locus on Y^M of $\langle C \rangle$, i.e., $\gamma(C) = \langle C \rangle \in \mathbb{G}(2, M)$ for the Gauss map

$$
\gamma = \gamma_{Y^M} : Y^M \dashrightarrow \mathbb{G}(2, M)
$$

sending $x \mapsto \mathbb{T}_x Y^M$. Then dim $\gamma(Y^M) = 1$. Since $Y^M = \text{Loc}(R^M)$, $\langle C \rangle \in \gamma(Y^M)$ is a general point if so is $C \in \mathbb{R}^M$. As in [Remark 2.5,](#page-3-2) the general fiber $\gamma^{-1}(\langle C \rangle)$ is a linear variety, which contradicts $C \subset \gamma^{-1}(\langle C \rangle)$.

Thus $\langle C \rangle \cap Y^M = C$ scheme-theoretically. This contradicts deg $Y^M = 6$. \Box

3. Special point of a conic: the embedded tangent space at the point containing the conic

We use the notations of §[2.](#page-1-2) From [Lemma 2.3,](#page-3-3) we may assume $(d, \dim Y)$ = $(6, n-3)$ and the formula (3) .

Lemma 3.1. *Let* $(C, x) \in \mathcal{U}$ *. Then the following holds.*

- (a) $C \not\subset \mathbb{T}_x X$ *if and only if* $\mathbb{T}_x X \cap C = \{x\}$
- (b) $C \subset \mathbb{T}_x X$ *if and only if one of the following conditions holds:* (i) $\langle C \rangle \subset X$; (ii) $\langle C \rangle \cap X$ *is non-reduced along* C; (iii) $x \in C \cap E$ *for some irreducible component* $E \neq C$ *of* $\langle C \rangle \cap X$ *.*

Proof. (a) Since $X \subset \mathbb{P}^n$ is a hypersurface and C is a smooth conic, if $C \not\subset \mathbb{T}_x X$, then we have $\mathbb{T}_x X \cap C \subset \mathbb{T}_x X \cap (\langle C \rangle \cap C) = \mathbb{T}_x C \cap C = \{x\}.$

(b) It is sufficient to consider the case when $\langle C \rangle \not\subset X$. If $C \subset \mathbb{T}_x X$, then $\langle C \rangle \subset \mathbb{T}_x X$, and then $\langle C \rangle \cap X$ is singular at x. This means that (ii) or (iii) \Box holds.

For a smooth conic $C \subset X$, we always have a point $x \in C$ satisfying $C \subset \mathbb{T}_x X$, as follows. Since $deg(X) = 6$, if (i) and (ii) does not hold, then we have a curve $E \neq C$ in $\langle C \rangle \cap X$, and have a point $x \in C \cap E \subset \langle C \rangle = \mathbb{P}^2$ as in (iii).

Definition 3.2. We set $\mathcal{U}^* \subset \mathcal{U}$ to be an irreducible component of $\{(C, x) \in \mathcal{U} \mid \mathcal{U}\}$ $C \subset \mathbb{T}_x X$ such that $\mathcal{U}^* \to R$ is dominant, and set $R_x^* := \pi(\mathrm{ev}^{-1}(x) \cap \mathcal{U}^*)$, which consists of conics $C \subset X$ such that $x \in C \subset \mathbb{T}_x X$. Note that $\text{Loc}(R_x^*) \subset \mathbb{T}_x X$.

Lemma 3.3. $\mathfrak{U}^* \neq \mathfrak{U}$.

Proof. We consider the Gauss map $\gamma: X \to (\mathbb{P}^n)^{\vee}$. If $\mathcal{U}^* = \mathcal{U}$, then it follows from [Lemma 2.4](#page-3-4) that $Y = \text{Loc}(R_x) = \text{Loc}(R_x^{\star}) \subset \mathbb{T}_x X$ for general $x \in Y$, and then $\mathbb{T}_x X = \langle Y \rangle \subset \mathbb{P}^n$ because of [Proposition 2.6.](#page-4-0) Then $\gamma(Y) = \langle Y \rangle \in (\mathbb{P}^n)^{\vee}$, which contradicts that γ is a finite morphism as we mentioned in [Remark 2.5.](#page-3-2)

Lemma 3.4. Let $x \neq y \in Y$ *satisfy* $y \in \mathbb{T}_x X$. Then $C \subset \mathbb{T}_x X$ for any $C \in R_{xy}$.

Proof. Take $C \in R_{xy}$ and suppose $C \not\subset \mathbb{T}_x X$. Since $\langle C \rangle \not\subset \mathbb{T}_x X$, $\langle C \rangle \cap \mathbb{T}_x X$ is the line passing through x and y. On the other hand, we have $\langle C \rangle \cap \mathbb{T}_x X = \mathbb{T}_x C$, which does not intersect with any point of C except x since C is a smooth conic, a contradiction. Thus we have $C \subset \mathbb{T}_x X$.

By [Lemma 3.3,](#page-6-1) we may assume that $\mathcal{U}^* \neq \mathcal{U}$. Then dim $\mathcal{U}^* = \dim R = r$. We take the projection

$$
\mu: \mathcal{U}^\star \times_R \mathcal{U} \simeq \{ (C, x, y) \in R \times Y \times Y \mid (C, x) \in \mathcal{U}^\star, (C, y) \in \mathcal{U} \} \to \text{ev}(\mathcal{U}^\star) \times Y,
$$

such that $\mu(C, x, y) = (x, y)$, where $ev : U \rightarrow Y$ is the second projection.

Let $(C, x, y) \in \mathcal{U}^* \times_R \mathcal{U}$ be general. (Then $x, y \in C \subset \mathbb{T}_x X$.) We can take the unique irreducible component $R_{x}^{\star} \subset R_{x}^{\star}$ containing C, and take the unique irreducible component $R'_{xy} \subset R_{xy}$ containing C. (The uniqueness comes from the general choice of C .)

Lemma 3.5. In the above setting, we have $R'_{xy} \subset R_x^*$ ′ *.*

Proof. It follows from [Lemma 3.4.](#page-6-2) □

Moreover, we have the following key proposition, where for the projection, $\lim(\mu) \to \text{ev}(\mathcal{U}^*) : (x, y) \mapsto x$, we also consider the following fiber product

$$
im(\mu) \times_{ev(\mathfrak{U}^*)} im(\mu) \simeq \{ (x, y, z) \in ev(\mathfrak{U}^*) \times Y \times Y \mid (x, y), (x, z) \in im(\mu) \}.
$$

Note that, for an element (x, y, z) of the above set, there exists conics $C_1, C_2 \subset \mathbb{T}_x X$ such that $x, y \in C_1$ and $x, z \in C_2$. The projection $\text{im}(\mu) \times_{\text{ev}(\mathfrak{U}^*)} \text{im}(\mu) \to Y \times Y$ is dominant.

Proposition 3.6. *Assume* $(d, \dim Y) = (6, n-3)$ *and the formula* [\(3\)](#page-3-0)*. Then the following holds.*

- (a) $Loc(R_x^{\star})$ $\mathcal{L}(S) = \text{Loc}(R'_{xy})$ and the dimension is $n - 4$ for general $(C, x, y) \in$ $\mathfrak{U}^\star \times_R \overset{\circ}{\mathfrak{U}}$.
- (b) \langle Loc(R_x^* $\langle f \rangle$ *is an* $(n-3)$ -plane of \mathbb{P}^n for general $(C, x) \in \mathcal{U}^*$.
- (c) *The projection*

$$
\mathcal{U}^\star \times_R \mathcal{U} \times_R \mathcal{U} \to \operatorname{im}(\mu) \times_{\mathrm{ev}(\mathcal{U}^\star)} \operatorname{im}(\mu)
$$

defined by $(C, x, y, z) \mapsto (x, y, z)$ *is dominant.*

(d) Let $(C, x, y, z) \in \mathcal{U}^* \times_R \mathcal{U} \times_R \mathcal{U}$ be general (here, $x, y, z \in C \subset \mathbb{T}_x X$). Then $Loc(R_x^{\star}$ \mathcal{L}') = Loc(R'_{yz}), where $R'_{yz} \subset R_{yz}$ *is the unique irreducible component containing* C*.*

From (a) and (b), we have that $Loc(R_x^{\star})$ \prime) is a hypersurface of \langle Loc (R_x^{\star}) $\langle \rangle$ = \mathbb{P}^{n-3} . Moreover, as a corollary, later we will show that $Loc(R_x^{\star})$ ′) is a quadric hypersurface, and also will see our problem is reduced to the case of $n = 6$ [\(Corollary 3.9,](#page-9-0) [Remark 3.11\)](#page-10-0).

In order to prove this proposition, we show the following two lemmas.

Lemma 3.7. $ev(\mathcal{U}^*) \subset Y$ *is of dimension* $\geq n-5$ *. Hence* $Loc(R_x^*) \neq Y$ *for* $general \ x \in \text{ev}(\mathfrak{U}^{\star}).$

Proof. Let $(C, x, y) \in \mathcal{U}^* \times_R \mathcal{U}$ be general such that $\langle C \rangle \not\subset X$. Considering the morphism μ , $R_x^{\star'} \cap R_y \subset R_{xy}$ is of dimension $\geqslant r+1-\dim \mathrm{ev}(\mathfrak{U}^*)-\dim Y$. From [Lemma 2.1,](#page-2-2) $r+2-\dim$ ev $(\mathcal{U}^*)-\dim Y \leqslant \dim Y$. Hence ev $(\mathcal{U}^*) \geqslant r+2-2\dim(Y)$, where the right hand side is $\geq n-5$.

Since $n \geq 6$, ev(U^{*}) has positive dimension. On the other hand, we have $\#\{x \in$ $ev(\mathcal{U}^*) | \text{Loc}(R_x^*) = Y \} < \infty$ as follows. If $\text{Loc}(R_x^*) = Y$, then $Y \subset \mathbb{T}_x X$ and then $\langle Y \rangle = \mathbb{T}_x X$ because of [Proposition 2.6.](#page-4-0) Hence, by the finiteness of the Gauss map of smooth X, we have the assertion.

Lemma 3.8. Let $S \subset \mathbb{P}^n$ be a non-linear projective variety. Assume Cone_x $S =$ Cone_y S for general $x, y \in S$. Then $\langle S \rangle \subset \mathbb{P}^n$ is a $(\dim(S) + 1)$ *-plane.*

Proof. Take a general point $x \in S$ and consider the linear projection $\pi_x : \mathbb{P}^n \to$ \mathbb{P}^{n-1} from x. Then we have

$$
\pi_x(S) = \pi_x(\text{Cone}_x S) = \pi_x(\text{Cone}_y S) = \text{Cone}_{\pi_x(y)}(\pi_x(S))
$$

for general $y \in S$. Hence $\pi_x(S) = \text{Cone}_{y'}(\pi_x(S))$ for general $y' \in \pi_x(S)$. This means that $\pi_x(S)$ is a $(\dim(S))$ -plane. Hence $Cone_x S$ is a $(\dim(S) + 1)$ -plane, which implies the assertion.

Proof of [Proposition 3.6.](#page-7-0) (a) We may assume $\langle C \rangle \not\subset X$. From [Lemma 2.1](#page-2-2) and the formula [\(4\),](#page-3-1) we have dim $Loc(R'_{xy}) \geq n-4$. On the other hand, [Lemma 3.7](#page-8-0) implies $Loc(R_x^{\star})$ \mathcal{O}') ≤ n – 4. Hence [Lemma 3.5](#page-7-1) implies Loc $(R'_{xy}) =$ Loc (R'_x) ′) and the dimension is $n-4$.

(b) From (a), for general $y \in Loc(R_x^{\star})$ '), it follows $Loc(R_x^{\star})$ \mathcal{L} = Loc(R'_{xy}), where the right hand side is the closure of $\bigcup_{C \in R'_{xy}}^{\infty} C \subset \mathbb{P}^n$. Since C one_x $C = \langle C \rangle =$ $Cone_{\nu} C,$

$$
P_x := \text{Cone}_x \operatorname{Loc}(R_x^{\star}) = \overline{\bigcup_{C \in R_{xy}'} \langle C \rangle} = \text{Cone}_y \operatorname{Loc}(R_x^{\star}).
$$

For two general points $y_1, y_2 \in \text{Loc}(R_x^*$ '), we have $\text{Cone}_{y_1} \text{Loc}(R_x^*$ $') = P_x =$ Cone_{y2} Loc (R_x^{\star}) ′). Hence the assertion follows from [Lemma 3.8.](#page-8-1)

(c) Since $\text{im}(\mu) \times_{ev(\mathfrak{U}^*)} \text{im}(\mu)$ is irreducible, it is sufficient to show that, for general $x \in \text{ev}(\mathcal{U}^*)$ and for general $y, z \in \text{Loc}(R_x^*)$ $'$), there exists C such that $(C, x, y, z) \in \mathcal{U}^\star \times_R \mathcal{U} \times_R \mathcal{U}.$

First we can take a general $(C_0, x, y) \in \mathcal{U}^\star \times_R \mathcal{U}$ with some $C_0 \in R^\star_x$ ′ . From (a), we have $Loc(R_x^{\star})$ $\mathcal{L}(Y) = \text{Loc}(R'_{xy})$. Since $z \in \text{Loc}(R^*_x)$ ′) is general, we have a general $C \in R'_{xy}$ such that $z \in C$.

(d) Consider the projection

$$
\operatorname{pr}_{34}:\mathcal{U}^{\star}\times_R\mathcal{U}\times_R\mathcal{U}\to Y\times Y
$$

sending $(C, x, y, z) \mapsto (y, z)$. Let F_{yz} be an irreducible component of the fiber of pr_{34} at a general $(y, z) \in Y \times Y$. We identify F_{yz} and its image in \mathcal{U}^* under the projection pr_{12} : $(C, x, y, z) \mapsto (C, x)$.

Let us consider

$$
\bigcup_{(C,x)\in F_{yz}: \text{ general}} \text{Loc}(R_x^{\star\prime}) \quad \subset \bigcup_{(C,x)\in F_{yz}: \text{ general}} \langle \text{Loc}(R_x^{\star\prime}) \rangle.
$$

Suppose that the closure of the left hand side is equal to Y . Then (b) implies

$$
Coney Y = Coney \overline{\bigcup_{(C,x)} Loc(R_x^{*'})} = \overline{\bigcup_{(C,x)} \langle Loc(R_x^{*'}) \rangle}.
$$

In the same way, $\text{Cone}_z Y = \overline{\bigcup_{(C,x)} \langle \text{Loc}(R_x^*)\rangle}$ $\overline{\langle \rangle}$. Hence Cone_y $Y = \text{Cone}_z Y$. Since y, z ∈ Y are general, [Lemma 3.8](#page-8-1) implies that $\langle Y \rangle \subset \mathbb{P}^n$ is an $(n-2)$ -plane, which contradicts [Proposition 2.6.](#page-4-0)

Hence the closure of $\bigcup_{(C,x)\in F_{yz}: \text{ general } } \text{Loc}(R^{\star}_x)$ ') is not equal to Y. Since $Loc(R_x^{\star})$ ′) is of codimension 1 in Y, it means that $L := \text{Loc}(R_x^*$ ′) is constant for general $(C, x) \in F_{yz}$. Then $Loc(R'_{yz}) = \overline{\bigcup_{(C,x)\in F_{yz}:\text{ general }C}} \subset L$. Since dim $Loc(R'_{yz}) \geq$ $n-4$, the assertion follows.

Corollary 3.9. *Assume* $(d, \dim Y) = (6, n-3)$ *and the formula* [\(3\)](#page-3-0). *Then the following holds.*

- (a) For general $(C, x) \in \mathcal{U}^*$, Loc $(R_x^*$ \langle \mathcal{L} α \langle \mathcal{L} α $\langle R_x^* \rangle$ $\langle \rangle$) = \mathbb{P}^{n-3} *is a quadric hypersurface.*
- (b) *The projection to the second factor*

$$
q: \{ (C, H) \in R \times (\mathbb{P}^n)^{\vee} \mid C \subset H \} \to (\mathbb{P}^n)^{\vee}
$$

is dominant. (In particular, a general fiber of q is of dimension $r - 3$ *.)*

Proof. (a) Let $y, z \in \text{Loc}(R_x^*$ ') be general. Let $M \subset \langle \text{Loc}(R_x^* \mid R_{\text{max}}) \rangle$ $\langle \rangle$ be a general 2-plane such that $y, z \in M$.

From [Proposition 3.6\(](#page-7-0)c), (C_0, x, y, z) is general in $\mathcal{U}^* \times_R \mathcal{U} \times_R \mathcal{U}$ with some C_0 . From [Proposition 3.6\(](#page-7-0)d), it holds that $Loc(R'_{yz}) = Loc(R'_x)$ ′). We consider

$$
R'_{yz} \to \langle yz \rangle^* : C \mapsto \langle C \rangle,
$$

which is generically finite, where $\langle yz \rangle^* \subset \mathbb{G}(2,\langle \text{Loc}(R^*_{x}) \rangle)$ $\langle \rangle$) is the set of 2-planes containing the line $\langle yz \rangle$. Since $\dim R'_{yz} \geqslant n-5 = \dim \langle yz \rangle^*$, this morphism is dominant.

Thus we can take a general $\tilde{M} \in \langle yz \rangle^* \subset \mathbb{G}(2,\langle \text{Loc}(R^*_x))$ $\langle \rangle$) near M such that $\tilde{M} = \langle \tilde{C} \rangle$ for some $\tilde{C} \in R'_{yz}$. By generality, $\tilde{M} \cap \text{Loc}(R_x^*$ ′) is irreducible. Hence $\tilde{C} = \tilde{M} \cap \text{Loc}(R^\star_x)$ '), which means that $\deg(\text{Loc}(R_x^{\star}))$ $^{\prime})) = 2.$

(b) For \tilde{M} in (a) above, we can take a hyperplane $\tilde{H} \subset \mathbb{P}^n$ containing \tilde{M} as a general element of $(\mathbb{P}^n)^{\vee}$, and then $(\tilde{C}, \tilde{H}) \in R \times (\mathbb{P}^n)^{\vee}$, which means that q is dominant.

Considering the projection to the first factor, we find that the dimension of the left hand side of q is $r + n - 3$. Thus a general fiber of q is of dimension $r - 3$. \Box

Remark 3.10. It is known that if a smooth hypersurface in \mathbb{P}^n of degree > 2 contains an *m*-dimensional quadric hypersurface, then $m \leq (n-1)/2$. Thus [Corollary 3.9\(](#page-9-0)a) implies $n \leq 7$.

Remark 3.11. By induction on n , it is sufficient to show [Theorem 1.1](#page-1-1) in the case of $n = 6$. We see the details in the following. Let $n > 6$ and let $R \subset R_2(X)$ be an irreducible component.

Assume $(d, \dim Y) = (6, n-3)$ and the formula [\(3\),](#page-3-0) i.e., $r := \dim R$ is greater than the expected one. Since q is dominant as in [Corollary 3.9\(](#page-9-0)b), a fiber of q at general $H \in (\mathbb{P}^n)^{\vee}$, which identified with $R \cap R_2(X \cap H)$, is of dimension $r-3$. Again since q is dominant, we may take a conic $C \subset H$ as a general member of R. Then we may take an irreducible component R' of $R_2(X \cap H)$ containing C such that dim $R' \geq r - 3 \geq 3n - 16$. Since C satisfies $\langle C \rangle \not\subset \mathbb{T}_x X$, we have $\langle C \rangle \not\subset \mathbb{T}_x X \cap H = \mathbb{T}_x (X \cap H)$. Then a general $\tilde{C} \in R'$ satisfies $\langle \tilde{C} \rangle \not\subset \mathbb{T}_x (X \cap H)$. Since H is general, $X \cap H \subset H = \mathbb{P}^{n-1}$ is smooth. Once [Theorem 1.1](#page-1-1) is proved for $n-1$, we have a contradiction since dim R' must be equal to the expected

dimension $3(n-1) - 14$.

Using the above results and notations, we now prove the main theorem.

Proof of [Theorem 1.1.](#page-1-1) Let $R \neq \emptyset$ be an irreducible component of $R_2(X)$ satisfy-ing the condition [\(1\),](#page-1-0) and assume that $r := \dim R$ is greater than the expected dimension. From [Lemma 2.3,](#page-3-3) [Corollary 3.9\(](#page-9-0)b), and [Remark 3.11,](#page-10-0) we may assume $n = d = 6$ and dim $Y = 3$. Then $r \geq 5$.

Claim 3.12. $deg(Y) = 3$ *or* 4 *and* $dim(Y) \ge 5$ *.*

Proof. From [Proposition 2.6,](#page-4-0) we have $\dim(Y) \geq 5$. In particular, $\deg(Y) \geq 3$. Let $H \subset \mathbb{P}^6$ be a general hyperplane, and set $Y' = Y \cap H$. It follows from [Corollary 3.9\(](#page-9-0)b) again that $R \cap R_2(Y')$ is of dimension ≥ 2 . It is classically known that, if a surface Y' has a 2-dimensional family of conics, then Y' is projectively equivalent to either the Veronese surface $\mathbb{P}^2 \hookrightarrow \mathbb{P}^5$ or its image under linear pro-jections (see [\[13,](#page-14-3) p. 130, p. 157]). Hence $deg(Y) = deg(Y') \leq 4$.

Our first goal is to show that $\langle Y \rangle \subset \mathbb{P}^6$ is of dimension 5. Set $Q_{yz} \subset Y$ to be the surface $Loc(R'_{yz})$ for general $(C, y, z) \in \mathcal{U} \times_R \mathcal{U}$.

Claim 3.13. Q_{yz} *is a quadric surface, and* $Q_{yz} = Q_{\tilde{y}z}$ *holds for general* $(C, \tilde{y}) \in$ $\mathfrak{U}_{R'_{yz}}.$

Proof. Since $(\tilde{C}, y, z, \tilde{y})$ is general in $\mathcal{U} \times_R \mathcal{U} \times_R \mathcal{U}$, R'_{yz} (resp. $R'_{\tilde{yz}}$) is the unique irreducible component of R_{yz} (resp. $R_{\tilde{y}z}$) containing \tilde{C} . We can take $x \in \tilde{C}$ such that (\tilde{C}, x) is general in \mathcal{U}^* . Then [Proposition 3.6\(](#page-7-0)d) implies $Q_{yz} = \text{Loc}(R_x^*)$ $') =$ $Q_{\tilde{u}z}$, which is a quadric as in [Corollary 3.9.](#page-9-0)

For general $y, z_1, z_2 \in Y$ (with some C_i such that (C_i, y, z_i) is general in $\mathfrak{U} \times_R \mathfrak{U}$), we write

$$
K_y^{z_1z_2}:=\mathbb{T}_yQ_{yz_1}\cap \mathbb{T}_yQ_{yz_2}\subset \mathbb{T}_yY=\mathbb{P}^3,
$$

whose dimension is ≥ 1 . Then $y \in K_y^{z_1 z_2} \subset \langle Q_{yz_1} \rangle \cap \langle Q_{yz_2} \rangle$. (Here we do *not* know " $z_i \in K_y^{z_1 z_2}$ ".)

Claim 3.14. $\dim(\langle Q_{yz_1} \rangle \cap \langle Q_{yz_2} \rangle) = 1$ *. Hence* $K_y^{z_1z_2} = \langle Q_{yz_1} \rangle \cap \langle Q_{yz_2} \rangle$ *.*

Proof. Suppose $\dim \langle Q_{yz_1} \rangle \cap \langle Q_{yz_2} \rangle \geq 2$ for general $y, z_1, z_2 \in Y$. First we take general points $y_0, z_1, z_2 \in Y$. By generality, $z_1 \notin \langle Q_{y_0 z_2} \rangle$. For general $y \in Q_{y_0 z_2}$, we have $\langle Q_{yz_2}\rangle = \langle Q_{y_0z_2}\rangle$ because of [Claim 3.13.](#page-11-0) Consider an open subset $Y^{\circ} \subset Y$ containing y_0 such that $\dim \langle Q_{yz_1} \rangle \cap \langle Q_{yz_2} \rangle \geq 2$ for $y \in Y^{\circ}$.

Since $Y = \overline{\bigcup_{y \in Q_{y_0 z_2} \cap Y^{\circ}} Q_{y z_1}}$, we have

$$
\text{Cone}_{z_1} Y = \overline{\bigcup_{y \in Q_{y_0 z_2} \cap Y^{\circ}} \langle Q_{y z_1} \rangle},
$$

where the right hand side contains $Q_{y_0z_2}$; in fact, it contains the 3-plane $\langle Q_{y_0z_2} \rangle$ since each $\langle Q_{yz_1} \rangle$ satisfies $\dim(\langle Q_{yz_1} \rangle \cap \langle Q_{y_0z_2} \rangle) \geq 2$. Hence $\text{Cone}_{z_1} Y = \text{Cone}_{z_1} \langle Q_{y_0z_2} \rangle$, which is a 4-plane, a contradiction to $\dim(Y) \geq 5$.

Claim 3.15. $K_y^{z_1z_2} \subset Q_{yz_1}$. Hence there exists an irreducible component $K_y^{z_1}$ of $Q_{yz_1} \cap \mathbb{T}_y Q_{yz_1} \subset \langle Q_{yz_1} \rangle$ such that $K_y^{z_1} = K_y^{z_1z_2}$ for general $z_2 \in Y$.

Proof. Suppose $K_y^{z_1z_2} \not\subset Q_{yz_1}$. Let $z, w \in Y$ be general such that $w \notin Q_{yz}$. For general $\tilde{y} \in Q_{yz}$, we have $Q_{\tilde{y}z} = Q_{yz}$.

Since Y is the closure of $\bigcup_{y \in Q_{yz}:\text{ general } Q_{\tilde{y}w}} Q_{\tilde{y}w}$, we have that $\text{Cone}_w Y$ is the closure of $\bigcup_{\tilde{y}\in Q_{yz}:\text{general}}\langle Q_{\tilde{y}w}\rangle$. Since $\tilde{y}\in K_{\tilde{y}}^{zw}$, we have $Q_{yz}\subset \overline{\bigcup_{\tilde{y}\in Q_{yz}:\text{general}}K_{\tilde{y}}^{zw}}$. Moreover, since Q_{yz} is codimension one in $\langle Q_{yz} \rangle$, and since $K_{\tilde{y}}^{zw} \not\subset Q_{yz}$ and $K_{\tilde{y}}^{zw} \subset \langle Q_{yz} \rangle$, we have

$$
\langle Q_{yz} \rangle \subset \overline{\bigcup_{\tilde{y} \in Q_{yz}: \text{ general}}} K_{\tilde{y}}^{zw}.
$$

Since the right hand side is contained in $\overline{\bigcup_{\tilde{y}\in Q_{yz}:\text{general}}\langle Q_{\tilde{y}w}\rangle}$, it holds $\langle Q_{yz}\rangle \subset$ Conew Y. Then $Cone_w\langle Q_{yz}\rangle \subset \text{Cone}_w Y$, where the left hand side is a 4-plane and the right hand side is of dimension 4. It holds that $Cone_w Y$ is a 4-plane, a contradiction.

Note that, since $K_j^{z_1z_2}$ is contained in $Q_{yz_1} \cap \mathbb{T}_y Q_{yz_1} \subset \langle Q_{yz_1} \rangle$, there exists an irreducible component of $Q_{yz_1} \cap T_y Q_{yz_1}$ which is equal to $K_{y}^{z_1z_2}$ for general $z_2 \in Y$. Hence the latter statement holds.

Claim 3.16. $\langle Y \rangle \subset \mathbb{P}^6$ *is of dimension* 5*.*

Proof. Let $y, z_1 \in Y$ be general. Since Y is the closure of $\bigcup_{z_2 \in Y : \text{general}} Q_{yz_2}$, it follows that Coney Y is the closure of $\bigcup_{z_2 \in Y : \text{general}} \langle Q_{yz_2} \rangle$. Since $K_{y}^{z_1} = K_{y}^{z_1z_2} \subset$ $\langle Q_{yz_2} \rangle$ for general $z_2 \in Y$, we have that $\widehat{\text{Cone}}_y Y$ is a cone with vertex $K^{z_1}_y = \mathbb{P}^1$. Let $\tilde{y}, \tilde{z}_1 \in Y$ be general. We may assume $\tilde{y} \notin K_{y}^{z_1}$ and $y \notin K_{\tilde{y}}^{\tilde{z}_1}$. We show the statement in the following two steps.

Step 1. Suppose that two lines $K_{\tilde{y}}^{z_1}$ and $K_{\tilde{y}}^{\tilde{z_1}}$ intersect at a point v. Then, for general $s, t \in Y$, the line K_s^t also intersects with each of $K_y^{z_1}$ and $K_{\tilde{y}}^{\tilde{z_1}}$. If $v \notin K_s^t$, then $s \in K_s^t \subset \langle K_{\tilde{y}}^{z_1}, K_{\tilde{y}}^{\tilde{z_1}} \rangle$; hence we have $Y \subset \langle K_{\tilde{y}}^{z_1}, K_{\tilde{y}}^{\tilde{z_1}} \rangle = \mathbb{P}^2$, a contradiction. If $v \in K_s^t$, then since $s \in Y$ is general and $\overline{s}v = K_s^t \subset Y$, it follows that Y is a

cone with vertex v; hence $Y \subset \mathbb{T}_v X$, which implies $\langle Y \rangle = \mathbb{T}_v X = \mathbb{P}^5$.

Step 2. Suppose $K_{y}^{z_1} \cap K_{\tilde{y}}^{\tilde{z}_1} = \emptyset$. We have $K_{y}^{z_1} = K_{y}^{z_1\tilde{y}} \subset Q_{y\tilde{y}} \subset \langle Q_{y\tilde{y}} \rangle$. In the same way, $K_{\tilde{y}}^{\tilde{z_1}} \subset Q_{y\tilde{y}}$. Since $K_{\tilde{y}}^{\tilde{z_1}} = \langle Q_{\tilde{y}\tilde{z}_1} \rangle \cap \langle Q_{y\tilde{y}} \rangle$, we have

$$
K_{y}^{z_{1}} \cap \langle Q_{\tilde{y}\tilde{z}_{1}} \rangle = K_{y}^{z_{1}} \cap K_{\tilde{y}}^{\tilde{z_{1}}} = \emptyset.
$$

For the linear projection $\pi_y : \mathbb{P}^6 \dashrightarrow \mathbb{P}^5$, we consider $\pi_y(Y) = \pi_y(\text{Cone}_y Y)$, a cone with vertex $w := \pi_y(K_y^{z_1})$. Since $w \notin \pi_y(\langle Q_{\tilde{y}z_1} \rangle)$ and since $\pi_y(Q_{\tilde{y}z_1})$ is of codimension 1 in $\pi_y(Y)$, it follows that $\pi_y(Y)$ is a cone of the quadric $\pi_y(Q_{\tilde{y}\tilde{z}_1})$ with vertex w. Then $\langle \pi_y(Y) \rangle = \mathbb{P}^4$. Since $y \in Y$ is general, it follows that Y is a 3-fold of degree 3 in $\langle Y \rangle = \mathbb{P}^5$.

Let us complete the proof. By the above claim, $\langle Y \rangle = \mathbb{P}^5$. Take $X' := X \cap \langle Y \rangle$. Since the Gauss map $\gamma = \gamma_X : X \to (\mathbb{P}^6)^{\vee}$ is finite, X' is singular at most finitely many points (X' is singular at x if and only if $\gamma(x) = \langle Y \rangle$ in $(\mathbb{P}^6)^{\vee}$). In particular, X' is irreducible. (This is because, if $X' = X'_1 \cup X'_2 \subset \langle Y \rangle = \mathbb{P}^5$, then $X'_1 \cap X'_2 \subset \text{Sing } X'.$) Here $Y \subset X' \subset \langle Y \rangle$.

Take a general hyperplane $M \subset \langle Y \rangle = \mathbb{P}^5$ such that $X'' := X' \cap M$ is smooth. Then we have

$$
Y \cap M \subset X'' \subset M = \mathbb{P}^4
$$

where $Y \cap M$ is a surface of degree ≤ 4 as in [Claim 3.12](#page-10-1) and X'' is a smooth 3-fold of degree 6 in \mathbb{P}^4 . This is a contradiction since $Pic(\mathbb{P}^4) \simeq \mathbb{Z} \to Pic(X'') : \mathcal{O}_{\mathbb{P}}(1) \to$ $\mathcal{O}_{\mathbb{P}}(1)|_{X''}$ is isomorphic due to the Lefschetz theorem.

Example 3.17. Let $X \subset \mathbb{P}^7$ be the Fermat hypersurface of degree 6. Then $\mathbb{P}^3 \subset$ X. Thus $R_2(\mathbb{P}^3) \subset R_2(X)$ is of dimension 8, and the expected dimension of $R_2(X)$ is $3n-2d-2=7$. For a general hyperplane $\mathbb{P}^6 \subset \mathbb{P}^7$, we have $\mathbb{P}^2 \subset X_1 := X \cap \mathbb{P}^6$. Then $R_2(\mathbb{P}^2) \subset R_2(X_1)$ is of dimension 5 and the expected dimension of $R_2(X_1)$ is 4.

In these examples, each $C \in R_2(\mathbb{P}^3)$ (resp. $C \in R_2(\mathbb{P}^2)$) satisfies $\langle C \rangle \subset \mathbb{P}^3 \subset X$ (resp. $\langle C \rangle = \mathbb{P}^2 \subset X_1$).

Example 3.18. Let $X \subset \mathbb{P}^{10}$ be the smooth hypersurface of degree 10 defined by the following polynomial,

$$
f := x_0^8(x_0^2 + x_1^2 + x_2^2) + \sum_{i=1}^5 x_i^{10} - \sum_{i=1}^5 x_{i+5}^{10}.
$$

Then the expected dimension of $R_2(X)$ is 8. We consider the 5-plane

$$
M := \bigcap_{i=1}^{5} (x_i - x_{i+5} = 0) = \mathbb{P}^5 \subset \mathbb{P}^{10},
$$

and take $Y \subset X \cap M$, to be the zero set of $x_0^2 + x_1^2 + x_2^2$ in M. Since Y is a cone of the conic $(x_0^2 + x_1^2 + x_2^2 = 0) \subset \mathbb{P}^2$, we have a birational map $R_2(Y) \to \mathbb{G}(2, \mathbb{P}^5)$: $C \mapsto \langle C \rangle$. In particular, dim $R_2(Y) = 9$. Thus, for an irreducible component $R \subset R_2(X)$ containing $R_2(Y)$, the dimension of R is greater that the expected dimension.

REFERENCES

- [1] W. Barth and A. Van de Ven. Fano varieties of lines on hypersurfaces. Arch. Math. (Basel) 31 (1978/79), 96–104.
- [2] R. Beheshti, Lines on projective hypersurfaces. J. Reine Angew. Math. 592 (2006), 1–21.
- [3] R. Beheshti, Hypersurfaces with too many rational curves, Math. Ann. 360 (2014), 753–768.
- [4] R. Behesti and N. M. Kumar, Spaces of rational curves in complete intersections, Compositio Mathematica 149 (2013), 1041–1060.
- [5] T. D. Browning and P. Vishe, Rational curves on smooth hypersurfaces of low degree, Algebra and Number Theory 11 (2017), 1657-1675.
- [6] A. Collino, J. P. Murre, G. E. Welters, On the family of conics lying on a quartic threefold, Rend. Sem. Mat. Univ. Politec. Torino 38 (1980), 151–181.
- [7] K. Furukawa, Rational curves on hypersurfaces, J. reine angew. Math. 665 (2012), 157-188.
- [8] J. Harris, M. Roth, and J. Starr. Rational curves on hypersurfaces of low degree. J. Reine Angew. Math. 571 (2004), 73–106.
- [9] J. Harris and J. Starr. Rational curves on hypersurfaces of low degree. II. Compos. Math. 141 (2005), 35–92.
- [10] J. Kollár, Rational curves on algebraic varieties, Ergeb. Math. Grenzgeb. (3) 32, Springer-Verlag, Berlin 1996.
- [11] J. M. Landsberg and C. Robles, Fubini's theorem in codimension two. J. Reine Angew. Math. **631** (2009), 221-235.
- [12] E. Riedl and D. Yang, Kontsevich spaces of rational curves on Fano hypersurfaces, arXiv:1409.3802.
- [13] J. G. Semple and L. Roth, Introduction to algebraic geometry, Clarendon Press, Oxford, 1949.
- [14] J. Starr. The Kodaira dimension of spaces of rational curves on low degree hypersurfaces, arXiv:math/0305432.
- [15] F. L. Zak, Tangents and secants of algebraic varieties. Transl. Math. Monographs 127, Amer. Math. Soc., Providence, 1993.

Graduate School of Mathematical Sciences, the University of Tokyo, Tokyo, Japan

E-mail address: katu@ms.u-tokyo.ac.jp