

Categorical Liveness Checking by Corecursive Algebras

Natsuki Urabe* Masaki Hara

Dept. Computer Science, The University of Tokyo, Japan
{urabenatsuki, qnighy}@is.s.u-tokyo.ac.jp

*JSPS Research Fellow

Ichiro Hasuo

National Institute of Informatics, Japan
i.hasuo@acm.org


Abstract—*Final coalgebras* as “categorical greatest fixed points” play a central role in the theory of coalgebras. Somewhat analogously, most proof methods studied therein have focused on *greatest* fixed-point properties like safety and bisimilarity. Here we make a step towards categorical proof methods for *least* fixed-point properties over dynamical systems modeled as coalgebras. Concretely, we seek a categorical axiomatization of well-known proof methods for liveness, namely *ranking functions* (in nondeterministic settings) and *ranking supermartingales* (in probabilistic ones). We find an answer in a suitable combination of coalgebraic simulation (studied previously by the authors) and *corecursive algebra* as a classifier for (non-)well-foundedness.

I. INTRODUCTION


A. Backgrounds

Verification of liveness, much like that of safety, is a prototypical problem that underlines verification of more complex alternating fixed-point specifications. Liveness means that something “good” eventually occurs, while safety means that anything “bad” never occurs.

1) *Ranking Functions*: As an example, suppose that we are given a transition system as in the figure (1). Here x_1 is an *accepting* state that represents a good event. The reachability problem—a typical example of liveness checking problems—asks the following: “Does there exist a path from the initial state x_0 to x_1 ?” The answer is yes: x_1 is reachable by the path $\rightarrow x_0 \rightarrow x_2 \rightarrow x_3 \rightarrow x_1$. Note that the path does not refer to the states x_4 and x_5 .




In the example above, we assumed that the system is controlled in an *angelic* manner: we can choose the next state to eventually reach a good state. However, real-world systems often contain *demonic* branching, too, where the next state is chosen to avoid a good state. Such a system can be modeled as a *two-player game* played by angelic and demonic players. The figure (2) illustrates an example. At the state x_2 the next move is chosen by the demonic player. The answer to the reachability problem is again yes: no matter if x_3 or x_4 is chosen as the successor of x_2 , the angelic player can force reaching x_1 (by $\rightarrow x_3 \rightarrow x_1$ and $\rightarrow x_4 \rightarrow x_5 \rightarrow x_3 \rightarrow x_1$).



Numerous methods are known for such liveness checking problems (e.g. [1], [2], [3]). A well-known method is the one

using a *ranking function* [4]. For a two-player game, a ranking function is typically defined as a function $b : X \rightarrow \mathbb{N}_\infty$, from the state space X to the set $\mathbb{N}_\infty = \mathbb{N} \cup \{\infty\}$, that satisfies the following conditions: (i) for each non-accepting state x of the angelic player, there exists a successor state x' such that $b(x) \geq b(x') + 1$; and (ii) for each non-accepting state y of the demonic player, we have $b(y) \geq b(y') + 1$ for each successor state y' of y . It is known that *soundness* holds: existence of a ranking function b such that $b(x) < \infty$ implies that, regardless of the demonic player’s choice, the angelic player can construct a path $x = x_0 \rightarrow x_1 \rightarrow \dots$ that eventually reaches an accepting state. The well-foundedness of \mathbb{N} is crucial here: we have $b(x_i) > b(x_{i+1})$ for each i before an accepting state is reached; and an infinite descending chain is impossible in \mathbb{N} . For example, in the two-player game in (2), the ranking function $b = [x_0 \mapsto 5, x_1 \mapsto 0, x_2 \mapsto 4, x_3 \mapsto 1, x_4 \mapsto 3, x_5 \mapsto 2]$ ensures that x_1 is reachable from x_0 . Intuitively, the value $b(x)$ bounds the number of steps from x to an accepting state.

2) *Ranking Supermartingales*: One can consider liveness checking problems also for probabilistic systems. A typical example is the *almost-sure reachability problem*: let us consider the *probabilistic transition system* (PTS) as in the figure (3). In the almost-sure reachability problem, we want to know if the accepting state x_1 is reached with probability 1. In the PTS in (3), the answer is yes, though there exists a path that does not visit any accepting state at all (namely $\rightarrow x_0 \rightarrow x_0 \rightarrow \dots$, but this occurs with probability 0).



A notion analogous to that of ranking function is also known for probabilistic systems, namely *ranking supermartingales* [5], [6]. For a fixed positive real $\varepsilon > 0$, an (ε -additive) ranking supermartingale is a function $b' : X \rightarrow [0, \infty]$, from the state space X to the set $[0, \infty]$ of extended non-negative real numbers, that satisfies the following condition.

$$\forall x \in X \setminus \text{Acc}. b'(x) \geq \left(\sum_{x' \in X} \text{Prob}(x \rightarrow x') \cdot b'(x') \right) + \varepsilon$$

Here $\text{Prob}(x \rightarrow x')$ denotes the probability with which the system makes a transition from x to x' . This means that for each state $x \in X$, the expected value of b' decreases by at least ε after a transition. The existence of a ranking supermartingale b' such that $b'(x) < \infty$ implies that the expected value of the number of steps from x to an accepting state is finite

functor F	$c : X \rightarrow FX$ represents
$(_)^\Sigma \times \{0, 1\}$	deterministic automaton
$\mathcal{P}^2(_) \times \{0, 1\}$	two-player game
$\mathcal{D}(_) \times \{0, 1\}$	probabilistic transition system (PTS)

Fig. 1. Coalgebraic representations of transition systems. Here \mathcal{P} and \mathcal{D} denote the powerset and the distribution functors respectively (Def. II.15).

(specifically it is no bigger than $b'(x)/\varepsilon$). From this it easily follows that an accepting state is visited almost surely.

3) *Coalgebras and Algebras*: This paper aims to understand, in the categorical terms of (co)algebra, essences of liveness checking methods like ranking functions and ranking supermartingales. Coalgebras are commonly used for modeling state-based dynamics in the categorical language (see e.g. [7], [8]). Formally, for an endofunctor F over a category \mathbb{C} , an F -coalgebra is an arrow c of the type $c : X \rightarrow FX$. We can regard X as a state space, F as a specification of the branching type, and $c : X \rightarrow FX$ as a transition function. By changing the functor F we can represent various kinds of transition types (see Fig. 1). It is also known that, using coalgebras, we can generalize various automata-theoretic notions and techniques (such as behavioral equivalence [9], bisimulation [10] and simulation [11]) to various systems (e.g. nondeterministic, probabilistic, and weighted ones).

A dual notion, i.e. an arrow of type $a : FX \rightarrow X$, is known as an F -algebra. In this paper, it is used to capture *properties* (or *predicates*) over a system represented as a coalgebra.

B. Contributions

We contribute a categorical axiomatization of “ranking functions” that is behind the well-known methods that we have sketched. It combines: *corecursive algebras* as value domains (that are, like \mathbb{N}_∞ , suited to detect well-foundedness) and lax homomorphisms (like in *coalgebraic simulations* [11], [12]). Based on the axiomatization we develop a general theory; our main result is *soundness*, i.e. that existence of a categorical ranking function indeed witnesses liveness (identified with a least fixed-point property). We also exploit our general theory and derive two new notions of “ranking functions” as instances. The two concrete definitions are new to the best of our knowledge.

We shall now briefly sketch our general theory, illustrating key notions and the backgrounds from which we derive them.

1) *Corecursive Algebras for (Non-)Well-Foundedness*: In the (conventional) definition of a ranking function $b : X \rightarrow \mathbb{N}_\infty$, well-foundedness of $\mathbb{N} = \mathbb{N}_\infty \setminus \{\infty\}$ plays an important role as it ensures that no path can continue infinitely (without hitting an accepting state). Similarly, for a ranking supermartingale, it is crucial that $[0, \infty) = [0, \infty) \setminus \{\infty\}$ has no infinite sequence that decreases everywhere at least by $\varepsilon > 0$. In unifying the two notions, we need to categorically capture well-foundedness.

Our answer comprises suitable use of *corecursive algebras* [13]. An F -algebra $r : FR \rightarrow R$ is said to be corecursive if from an arbitrary coalgebra $c : X \rightarrow FX$ there exists a unique

$$FX \xrightarrow{F\langle c \rangle_r} FR \xrightarrow{r} R$$

coalgebra-algebra homomorphism $\langle c \rangle_r$ (see the diagram). Corecursive algebras have been previously used to describe general structured corecursion [13] (see also Rem. II.17). Our use of them in this paper seems novel: r being corecursive means that the function $\Phi_{c,r} : f \mapsto r \circ Ff \circ c$ has a unique fixed point; in particular its least and greatest fixed points coincide¹; we find this feature of corecursive algebras suited for their use as categorical “classifiers” for (non-)well-foundedness.

2) *Modalities and Least Fixed-Point Properties*: Liveness properties such as reachability and termination are all instances of *least fixed-point properties*: once a proper *modality* \heartsuit_σ is fixed, the property in question is described by a least fixed-point formula $\mu u. \heartsuit_\sigma u$. The way we categorically formulate these constructs, as shown below, is nowadays standard (see e.g. [17], [18]). As the base category \mathbb{C} we use **Sets** in this paper (although extensions e.g. to **Meas** would not be hard).

- We fix a domain $\Omega \in \mathbb{C}$ of *truth values* (e.g. $\Omega = \{0, 1\}$), and a *property* over $X \in \mathbb{C}$ is an arrow $u : X \rightarrow \Omega$.
- A (state-based, dynamical) system is a coalgebra $c : X \rightarrow FX$ for a suitable functor $F : \mathbb{C} \rightarrow \mathbb{C}$.
- A modality \heartsuit_σ is interpreted as an F -algebra² $\sigma : F\Omega \rightarrow \Omega$ over Ω (see Example III.4 and Prop. IV.2 for examples).
- Assuming some syntax is given, we should be able to derive the interpretation $\llbracket \heartsuit_\sigma \varphi \rrbracket_c$ of a modal formula from $\llbracket \varphi \rrbracket_c$. In the current (purely semantical) framework this goes as follows. Given a property $u : X \rightarrow \Omega$, we define the property $\Phi_{c,\sigma}(u) : X \rightarrow \Omega$ by the composite

$$\Phi_{c,\sigma}(u) = (X \xrightarrow{c} FX \xrightarrow{F^u} F\Omega \xrightarrow{\sigma} \Omega) .$$

- Assuming a suitable order structure \sqsubseteq_Ω on Ω and additional monotonicity requirements, the correspondence $\Phi_{c,\sigma} : \Omega^X \rightarrow \Omega^X$ has the least fixed point. It is denoted by $\llbracket \mu \sigma \rrbracket_c : X \rightarrow \Omega$; intuitively it is the interpretation $\llbracket \mu u. \heartsuit_\sigma u \rrbracket_c$ of the formula $\mu u. \heartsuit_\sigma u$ in the system c .

Concrete examples are in §III-B. Another standard categorical modeling of a modality (see e.g. [19]) is by a *predicate lifting*, i.e. a natural transformation $\sigma_X : \Omega^X \Rightarrow \Omega^{FX}$. It corresponds to our modeling via the Yoneda lemma; see e.g. [17].

3) *Ranking Functions, Categorically*: $FX \xrightarrow{F\llbracket \mu \sigma \rrbracket_c} F\Omega$ (4)

Our modeling is summarized on the right: the liveness property $\llbracket \mu \sigma \rrbracket_c$ in question is the *least* arrow (with respect to the order on Ω) that makes the square commute (note the subscript $=_\mu$).

The liveness checking problem is then formulated as follows: given an arrow $h : X \rightarrow \Omega$, we would like to decide if

¹Examples abound in computer science—especially in domain theory—where similar coincidences play important roles. They include: limit-colimit coincidence [14] and initial algebra-final coalgebra coincidence [15], [16].

² We use the same functor F for coalgebras (systems) and algebras (modalities). This characterization is used in [17], [18] and also found in many coalgebraic modal logic papers (e.g. [19]). However, for some examples, it comes more natural to use functors F and G together with a natural transformation $\alpha : F \Rightarrow G$, and to model a system and a modality as $c : X \rightarrow FX$ and $\sigma : G\Omega \rightarrow \Omega$ respectively. This modeling induces our current one as σ and α together induce an F -algebra $F\Omega \xrightarrow{\alpha\sigma} G\Omega \xrightarrow{\sigma} \Omega$ (cf. §L).

$h \sqsubseteq_{\Omega} \llbracket \mu\sigma \rrbracket_c$ holds. Here \sqsubseteq_{Ω} denotes the pointwise extension of the order on Ω . For example, let's say we want to check the assertion that a specific state $x_0 \in X$ satisfies the liveness property $\llbracket \mu\sigma \rrbracket_c$. In this case we would define the above “assertion” $h: X \rightarrow \Omega$, where $\Omega = \{0 \sqsubseteq_{\Omega} 1\}$, by: $h(x_0) = 1$ and $h(x) = 0$ for all $x \neq x_0$.

Our categorical framework of ranking function-based verification goes as follows.

$$\begin{array}{ccccc} FX & \xrightarrow{Fb} & FR & \xrightarrow{Fq} & F\Omega \\ \uparrow c & \sqsubseteq & \downarrow r & \sqsubseteq & \downarrow \sigma \\ X & \xrightarrow{b} & R & \xrightarrow{q} & \Omega \end{array} \quad (5)$$

(A curved arrow labeled h points from X to Ω , and a curved arrow labeled \sqsubseteq points from R to Ω .)

- We fix a *ranking domain*—the value domain for ranking functions—to be an algebra $r: FR \rightarrow R$ together with a lax homomorphism $q: R \rightarrow \Omega$ (from r to σ , in the right square in (5)). The latter q identifies $r: FR \rightarrow R$ as a “refinement” of the modality $\sigma: F\Omega \rightarrow \Omega$. A crucial requirement is that r is *corecursive*, making it suited for detecting well-foundedness.
- A (categorical) *ranking arrow* for a coalgebra $c: X \rightarrow FX$ is then defined to be a lax homomorphism $b: X \rightarrow R$, in the sense shown in the left square in (5).
- Our soundness theorem says: given an assertion $h: X \rightarrow \Omega$, in order to establish $h \sqsubseteq \llbracket \mu\sigma \rrbracket_c$, it suffices to find a ranking arrow $b: X \rightarrow R$ such that $h \sqsubseteq q \circ b$ (see (5)).

This way the problem of verifying a least fixed-point property is reduced to finding a witness b . Note that the requirement on the ranking arrow b —namely $b \sqsubseteq r \circ Fb \circ c$ —is *local* (it only involves one-step transitions) and hence easy to check.

Our main technical contribution is a proof for soundness. There we argue in terms of inequalities between arrows—much like in [11], [12]—relying on fundamental order-theoretic results on fixed points (Knaster–Tarski and Cousot–Cousot, see §II-D). Corecursiveness of r is crucial there.

4) *Concrete Examples*: Ranking functions for two-player games (§I-A) are easily seen to be an instance of our categorical notion, for suitable F, Ω and R .

Our second example in §I-A—additive ranking supermartingales in the probabilistic setting—is itself not an example, however. Analyzing its reason we are led to a few variations of the definition, among which some seem new. We discuss these variations: their relationship, advantages and disadvantages.

C. Organization of this Paper

In §II we introduce preliminaries on: our running examples (two-player games and PTSs); liveness checking methods for them; (co)algebras; and least and greatest fixed points. Our main contribution is in §III, where our categorical developments are accompanied (for illustration) by concrete examples from two-player games. The entailments of our general framework in the probabilistic setting are described in §IV. Finally in §V we conclude.

Some details and proofs are deferred to the appendices.

II. PRELIMINARIES

In this paper $[0, \infty)$ and $[0, \infty]$ denote the sets $\{a \in \mathbb{R} \mid a \geq 0\}$ and $\{a \in \mathbb{R} \mid a \geq 0\} \cup \{\infty\}$ respectively. We extend

the ordinary order \leq over \mathbb{R} to $[0, \infty]$ by regarding ∞ as the greatest element. We write \mathbb{N}_{∞} for $\mathbb{N} \cup \{\infty\}$. For a function $\varphi: X \rightarrow [0, 1]$, its *support* $\{x \in X \mid \varphi(x) > 0\}$ is denoted by $\text{supp}(\varphi)$.

A. Two-player Games and Ranking Functions

Our *two-player games* are played by an angelic player max and a demonic player min.

Definition II.1 (two-player game). A *(two-player) game structure* is a triple $G = (X_{\max}, X_{\min}, \tau)$ of a set X_{\max} of states of the player max, a set X_{\min} of states of the player min, and a transition relation $\tau \subseteq X_{\max} \times X_{\min} \cup X_{\min} \times X_{\max}$.

A *strategy of the player max* is a partial function $\alpha: X_{\max} \times (X_{\min} \times X_{\max})^* \rightarrow X_{\min}$ such that for each $x_0 y_1 x_1 \dots y_n x_n \in X_{\max} \times (X_{\min} \times X_{\max})^*$, if $\alpha(x_0 y_1 x_1 \dots y_n x_n)$ is defined then $(x_n, \alpha(x_0 y_1 x_1 \dots y_n x_n)) \in \tau$. A *strategy of the player min* is a partial function $\beta: X_{\max} \times (X_{\min} \times X_{\max})^* \times X_{\min} \rightarrow X_{\max}$ such that for each $x_0 y_1 x_1 \dots y_{n-1} x_{n-1} y_n \in X_{\max} \times (X_{\min} \times X_{\max})^* \times X_{\min}$, if $\beta(x_0 y_1 x_1 \dots y_{n-1} x_{n-1} y_n)$ is defined then $(y_n, \beta(x_0 y_1 x_1 \dots y_{n-1} x_{n-1} y_n)) \in \tau$.

For $x \in X_{\max}$ and a pair of strategies α and β of the players max and min, respectively, the *run induced by α and β from x* is a possibly infinite sequence $\rho^{\alpha, \beta, x} = x_0 y_1 x_1 y_2 x_2 \dots$ that is an element of the set

$$\begin{aligned} & (X_{\max} \times (X_{\min} \times X_{\max})^* \times \{\perp_{\max}\}) \\ & \cup (X_{\max} \times (X_{\min} \times X_{\max})^* \times X_{\min} \times \{\perp_{\min}\}) \\ & \cup (X_{\max} \times X_{\min})^{\omega} \end{aligned}$$

and is inductively defined as follows: for $n = 0$, $x_0 = x$; and for $n > 0$,

$$\begin{aligned} y_n &= \begin{cases} \perp_{\max} & (\alpha(x_0 y_1 x_1 \dots y_{n-1} x_{n-1}) \text{ is undefined}) \\ \alpha(x_0 y_1 x_1 \dots y_{n-1} x_{n-1}) & (\text{otherwise}), \text{ and} \end{cases} \\ x_n &= \begin{cases} \perp_{\min} & (\beta(x_0 y_1 x_1 \dots x_{n-1} y_n) \text{ is undefined}) \\ \beta(x_0 y_1 x_1 \dots x_{n-1} y_n) & (\text{otherwise}). \end{cases} \end{aligned}$$

The symbol \perp_{\max} (resp. \perp_{\min}) represents the end of the run at max's (resp. min's) turn: it means the player got stuck.

Once an initial state x and strategies α and β of the players max and min are fixed, a run $\rho^{\alpha, \beta, x}$ is determined. There are different ways to determine the “winner” of a run, including: max wins if min gets stuck; max wins if he does not get stuck; max wins if some specified states are visited infinitely many times (the *Büchi condition*), etc. In this paper where we focus on liveness, we choose the following (rather simple) winning condition: the player max wins if an *accepting state* is reached or the player min gets stuck. Studies of more complex conditions (like the Büchi condition) are left as future work.


Definition II.2 (reaching set). Let $G = (X_{\max}, X_{\min}, \tau)$ be a two-player game structure. We fix a set $\text{Acc} \subseteq X_{\max}$ of *accepting states*. A run $\rho^{\alpha, \beta, x} = x_0 y_1 x_1 \dots$ on $(X_{\max}, X_{\min}, \tau)$ is *winning* with respect to Acc for the player max if

- $x_n \in \text{Acc}$ for some n ; or
- $\rho^{\alpha, \beta, x}$ is a finite sequence whose last letter is \perp_{\min} .

We define the *reaching set* $\text{Reach}_{G, \text{Acc}} \subseteq X_{\max}$ by:

$$\text{Reach}_{G, \text{Acc}} = \left\{ x \in X \mid \begin{array}{l} \exists \alpha : \text{strategy of max.} \\ \forall \beta : \text{strategy of min.} \\ \rho^{\alpha, \beta, x} \text{ is winning wrt. Acc} \end{array} \right\}. \quad (6)$$

Example II.3. We define a game structure $G = (X_{\max}, X_{\min}, \tau)$ by $X_{\max} = \{x_0, x_1, x_2, x_3, x_4\}$, $X_{\min} = \{y_0, y_1, y_2, y_3\}$ and $\tau = \{(x_0, y_0), (x_0, y_1), (x_1, y_1), (x_1, y_2), (x_3, y_3), (y_0, x_2), (y_1, x_1), (y_1, x_2), (y_2, x_3), (y_2, x_4)\}$. Let $\text{Acc} = \{x_2\}$. The situation is shown on the right. Then the reaching set is $\text{Reach}_{G, \text{Acc}} = \{x_0, x_2, x_3\}$.



Ranking functions. Suppose that we are given a game structure $G = (X_{\max}, X_{\min}, \tau)$ and a set $\text{Acc} \subseteq X_{\max}$ of accepting states, and want to prove that a state x is included in $\text{Reach}_{G, \text{Acc}}$. A *ranking function* is a standard proof method in such a setting. There are variations in the definition of ranking function [4], [20], [21]: in this paper we use the following.

Definition II.4 (ranking function). Let $G = (X_{\max}, X_{\min}, \tau)$ be a game structure and $\text{Acc} \subseteq X_{\max}$. We fix an ordinal \mathfrak{z} and let $\text{Ord}_{\leq \mathfrak{z}} = \{n \mid n \leq \mathfrak{z}\}$ be the set of ordinals smaller than or equal to \mathfrak{z} . A function $b : X_{\max} \rightarrow \text{Ord}_{\leq \mathfrak{z}}$ is called a *ranking function* (for G and Acc) if it satisfies

$$\min_{y: (x, y) \in \tau} \sup_{x': (y, x') \in \tau} b(x') \hat{+} 1 \leq b(x)$$

for each $x \in X_{\max} \setminus \text{Acc}$, where $b(x') \hat{+} 1 = \min\{b(x') + 1, \mathfrak{z}\}$ denotes addition truncated at \mathfrak{z} .

The following well-known theorem states soundness, i.e. that a ranking function witnesses reachability.

Theorem II.5 (soundness, see e.g. [4]). *Let \mathfrak{z} be an ordinal, and let $b : X \rightarrow \text{Ord}_{\leq \mathfrak{z}}$ be a ranking function for G and Acc . Then $b(x) < \mathfrak{z}$ (i.e. $b(x) \neq \mathfrak{z}$) implies $x \in \text{Reach}_{G, \text{Acc}}$. \square*

Example II.6. For the game in Example II.3, let $\mathfrak{z} = \omega$ and define a function $b : X \rightarrow \text{Ord}_{\leq \mathfrak{z}}$ by $b(x_0) = 1$, $b(x_2) = 0$ and $b(x_1) = b(x_3) = b(x_4) = \omega$. Then b is a ranking function. Hence by Thm. II.5, we have $x_0 \in \text{Reach}_{G, \text{Acc}}$.

Completeness (the converse of Thm. II.5) does not hold. A counterexample is given later in Example III.17.

Remark II.7. A strategy $\alpha : X_{\max} \times (X_{\min} \times X_{\max})^* \rightarrow X_{\min}$ is said to be *positional* if its outcome depends only on the last state of the input, i.e. $x_n = x'_n$ implies $\alpha(x_0 y_1 \dots y_n x_n) = \alpha(x'_0 y'_1 \dots y'_n x'_n)$. It is known that a positional strategy suffices as long as we consider reaching sets, i.e. the set $\text{Reach}_{G, \text{Acc}}$ in Def. II.2 is unchanged if we replace “ $\exists \alpha : \text{strategy of max}$ ” in (6) with “ $\exists \alpha : \text{positional strategy of max}$ ” (see e.g. [22]).

A ranking function allows us to synthesize such a positional strategy. Let $x \in X_{\max}$ and $b : X_{\max} \rightarrow \text{Ord}_{\leq \mathfrak{z}}$ be a ranking function s.t. $b(x) < \infty$. We define a strategy α for max by

$$\alpha(x_0 y_1 \dots y_n x_n) = \arg \min_{y: (x_n, y) \in \tau} \sup_{x': (y, x') \in \tau} b(x').$$

Then it is a positional strategy such that for each strategy β of min, the run $\rho^{\alpha, \beta, x}$ is winning wrt. Acc for max.

B. Probabilistic Transition Systems and Ranking Supermartingales

Definition II.8 (PTS). A *probabilistic transition system* (PTS) is a pair $M = (X, \tau)$ of a set X and a transition function $\tau : X \rightarrow \mathcal{D}X$. Here $\mathcal{D}X = \{d : X \rightarrow [0, 1] \mid \sum_{x \in X} d(x) = 1\}$ is the set of probability distributions over X .

Definition II.9 (reachability probability). Let $M = (X, \tau)$ be a PTS. We fix a set $\text{Acc} \subseteq X$ of *accepting states*. For each $x \in X$ and $n \in \mathbb{N}$, we define a value $f_n(x) \in [0, 1]$ by:

$$f_n(x) = \sum \left\{ \prod_{i=0}^{k-1} \tau(x_i)(x_{i+1}) \mid \begin{array}{l} k \leq n-1, x_0, \dots, x_k \in X, \\ x_0 = x, x_i \notin \text{Acc} (\forall i \in [0, k-1]), \\ \text{and } x_k \in \text{Acc} \end{array} \right\}.$$


Note that if $x \in \text{Acc}$ then $f_n(x) = 1$. As the sequence $(f_n(x))_{n \in \mathbb{N}}$ is increasing for each $x \in X$, we can define a function $f : X \rightarrow [0, 1]$ by:

$$f(x) = \lim_{n \rightarrow \infty} f_n(x).$$

The function f is called the *reachability probability function* with respect to M and Acc , and is denoted by $\text{Reach}_{M, \text{Acc}}$.

Here the value $f_n(x) \in [0, 1]$ is the probability with which an accepting state is reached within n steps from x .

Example II.10. We define a PTS $M = (X, \tau)$ by $X = \{x_0, x_1, x_2, x_3\}$, and $\tau(x_0) = [x_1 \mapsto \frac{1}{2}, x_3 \mapsto \frac{1}{2}]$, $\tau(x_1) = [x_1 \mapsto \frac{1}{2}, x_2 \mapsto \frac{1}{2}]$, $\tau(x_2) = [x_2 \mapsto 1]$, and $\tau(x_3) = [x_3 \mapsto 1]$. Let $\text{Acc} = \{x_2\}$. The situation is as shown on the right.



Then $\text{Reach}_{M, \text{Acc}} : X \rightarrow [0, 1]$ assigns $\frac{1}{2}$ to x_0 , 1 to x_1 and x_2 , and 0 to x_3 .

Let us consider the *almost-sure reachability problem* for PTS. Given a PTS $M = (X, \tau)$, a set $\text{Acc} \subseteq X$ of accepting states and an (initial) state $x \in X$, we want to prove that $\text{Reach}_{M, \text{Acc}}(x) = 1$. For this problem, a ranking function-like notion called *ranking supermartingale* [5] is known. There are several variations in its definition. We follow the definition in [6]; a variation can be found in [23].

Definition II.11 (ε -additive ranking supermartingale). Let $M = (X, \tau)$ be a PTS and $\text{Acc} \subseteq X$ be the set of accepting states. Let $\varepsilon > 0$ be a real number. A function $b' : X \rightarrow [0, \infty]$ is an ε -*additive ranking supermartingale* (for M and Acc) if

$$\left(\sum_{x' \in \text{supp}(\tau(x))} \tau(x)(x') \cdot b'(x') \right) + \varepsilon \leq b'(x)$$

holds for each $x \in X \setminus \text{Acc}$.

Intuitively an ε -additive ranking supermartingale b' bounds the expected number of steps to accepting states: specifically it is no bigger than $b'(x)/\varepsilon$.

Theorem II.12 ([6]). *Let $b' : X \rightarrow [0, \infty]$ be an ε -additive ranking supermartingale for M and Acc . Then $b'(x) < \infty$ implies $\text{Reach}_{M, \text{Acc}}(x) = 1$. \square*

Example II.13. For the PTS in Example II.10, we define $b' : X \rightarrow [0, \infty]$ by $b'(x_0) = \infty$, $b'(x_1) = 2$, $b'(x_2) = 0$ and $b'(x_3) = \infty$. Then b' is a 1-additive ranking supermartingale. Hence by Thm. II.12, we have $\text{Reach}_{M, \text{Acc}}(x_1) = 1$.

C. Categorical Preliminaries

We assume that readers are familiar with basic categorical notions. For more details, see e.g. [24], [7].

Definition II.14 ((co)algebra). Let $F : \mathbb{C} \rightarrow \mathbb{C}$ be an endofunctor on a category \mathbb{C} . An F -coalgebra is a pair (X, c) of an object X in \mathbb{C} and an arrow c of the type $c : X \rightarrow FX$. An F -algebra is a pair (A, a) of an object A in \mathbb{C} and an arrow a of the type $a : FA \rightarrow A$.

In this paper we exclusively use the category **Sets** of sets and functions as the base category \mathbb{C} (although extensions e.g. to **Meas** would not be hard). We would be interested in endofunctors composed by the following.

Definition II.15 (\mathcal{P} , \mathcal{D} and $(_) \times C$). The powerset functor $\mathcal{P} : \mathbf{Sets} \rightarrow \mathbf{Sets}$ is such that:

- for each $X \in \mathbf{Sets}$, $\mathcal{P}X = \{A \subseteq X\}$; and
- for each $f : X \rightarrow Y$ and $A \in \mathcal{P}X$, $(\mathcal{P}f)(A) = \{y \in Y \mid \exists x \in A. f(x) = y\}$.

The (discrete) distribution functor $\mathcal{D} : \mathbf{Sets} \rightarrow \mathbf{Sets}$ is:

- for each $X \in \mathbf{Sets}$, $\mathcal{D}X = \{d : X \rightarrow [0, 1] \mid \sum_{x \in X} d(x) = 1\}$; and
- for each $f : X \rightarrow Y$, $\delta \in \mathcal{D}X$ and $y \in Y$, $(\mathcal{D}f)(\delta)(y) = \sum_{x \in f^{-1}(y)} \delta(x)$.

For $C \in \mathbf{Sets}$, the functor $(_) \times C : \mathbf{Sets} \rightarrow \mathbf{Sets}$ is:

- for $X \in \mathbf{Sets}$, $X \times C = \{(x, c) \mid x \in X, c \in C\}$; and
- for each $f : X \rightarrow Y$, $x \in X$ and $c \in C$, $(f \times C)(x, c) = (f(x), c)$.

We combine these functors for modeling transition types of various kinds of systems (Fig. 1). For two-player games we use the functor $F_g = \mathcal{P}^2(_) \times \{0, 1\}$. It works as follows.

- For each set X , $F_g X = \{\Gamma \subseteq \mathcal{P}X\} \times \{0, 1\}$.
- For each function $f : X \rightarrow Y$, $F_g f : F_g X \rightarrow F_g Y$ is defined by $F_g f(\Gamma, t) = (\{\{f(x) \mid x \in A\} \mid A \in \Gamma\}, t)$.

The correspondence between F_g -coalgebras and two-player games will be spelled out in Def. III.1.

The key idea in this paper is to use a *corecursive algebra* as a classifier for (non-)well-foundedness.

Definition II.16 (corecursive algebra, [13]). $F X \xrightarrow{Ff} F R$
An F -algebra $r : F R \rightarrow R$ is *corecursive* if $\begin{array}{c} \uparrow_c = r \downarrow \\ X \xrightarrow{f} R \end{array}$
given an arbitrary coalgebra $c : X \rightarrow F X$, there exists a unique arrow $f : X \rightarrow R$ such that $f = r \circ F f \circ c$.

Remark II.17. The connection between corecursive algebras and (non-)well-foundedness has been hinted by some existing results. For example, for set functors preserving monos and

inverse image diagrams, *recursive coalgebras*—the categorical dual of corecursive algebras used for general structured recursion in [25]—are known to coincide with *well-founded coalgebras* [26], where well-foundedness is categorically modeled in terms of “inductive components.” For more general categories, it is known that if a functor preserves monos then well-foundedness implies recursiveness, but its converse does not necessarily hold [27]. The dual of this result, between corecursive and *anti-founded* algebras, is pursued in [13] but with limited success.

In [28] the notion of *co-founded part* of an algebra is introduced, with a main theorem that the co-founded part of an injectively structured corecursive algebra carries a final coalgebra. The result is used for characterizing a final coalgebra as that of suitable modal formulas. Despite its name, co-founded parts have little to do with our current view of corecursive algebras here as well-foundedness classifiers.

Discussions on other works on corecursive algebra are found in §L in the appendix.

D. Verification of Least/Greatest Fixed-Point Properties

The following results are fundamental in the studies of fixed-point specifications.

Theorem II.18. *Let L be a complete lattice, and $f : L \rightarrow L$ be a monotone function.*

- 1) (Knaster–Tarski) *The set of prefixed points (i.e. those $l \in L$ such that $f(l) \sqsubseteq l$) forms a complete lattice. Moreover its least element is (not only a prefixed but) a fixed point, that is, the least fixed point μf .*
- 2) (Cousot–Cousot [29]) *Consider the (transfinite) sequence $\perp \sqsubseteq f(\perp) \sqsubseteq \dots \sqsubseteq f^\alpha(\perp) \sqsubseteq \dots$ where, for a limit ordinal α , we define $f^\alpha(\perp) = \bigsqcup_{\beta < \alpha} f^\beta(\perp)$. The sequence eventually stabilizes and its limit is the least fixed point μf .* \square

For the greatest fixed point νf we have the dual results. From these four results—Knaster–Tarski and Cousot–Cousot, for μ and ν —we derive the following four “proof principles.”

Corollary II.19. *Under the conditions of Thm. II.18:*

- (KT μ) $f(l) \sqsubseteq l$ implies $\mu f \sqsubseteq l$.
- (KT ν) $l \sqsubseteq f(l)$ implies $l \sqsubseteq \nu f$.
- (CC μ) $f^\alpha(\perp) \sqsubseteq \mu f$ for each ordinal α .
- (CC ν) $\nu f \sqsubseteq f^\alpha(\top)$ for each ordinal α . \square

Among these four, however, only two are applicable in verification: our goal is to show that an assertion h is *below* a fixed point (see §I-B3); the rules (KT ν) and (CC μ) are for *under-approximation* and thus serve our goal; but the other two are for *over-approximating* the fixed point in question.

It is these order-theoretic principles behind (namely CC and KT) that cause the difference between the proof methods for liveness (lfp’s) and safety (gfp’s). The role of ordinals α —equivalence classes of well-ordered sets—in (CC μ) can be discerned in the definitions of ranking functions/supermartingales. These proof methods for liveness are

in a sharp contrast with those for safety, in which finding an *invariant* (i.e. a post-fixed point l in (KT ν)) suffices.

The basic idea behind the current contribution—liveness checking by combination of coalgebraic simulation and corecursive algebra—can be laid out as follows. For verification it is convenient if we can rely on *certificates* whose constraints are locally checkable. Their examples include invariants, various notions of (bi)simulation and a general notion of coalgebraic simulation; they are all postfixed points in a suitable sense. They should thus be able to witness only gfp’s (not lfp’s) in view of Cor. II.19. Here we leverage the lfp-gfp coincidence in corecursive algebras to make coalgebraic simulations witness lfp’s too. The lfp-gfp coincidence might seem a serious restriction but it is a common phenomenon in many “interesting” structures in computer science (as we discussed at the end of §I-B3)

III. CATEGORICAL RANKING FUNCTIONS

Here we present our general categorical framework for ranking function-based liveness checking.

A. Running Example: Two-Player Games

In this section, in order to provide abstract notions with intuitions, we use two-player games (§II-A) as a running example. We use the functor $F_g = \mathcal{P}^2(_) \times \{0, 1\} : \mathbf{Sets} \rightarrow \mathbf{Sets}$ to model them as coalgebras (§II-C, here g stands for “game”).

Definition III.1. Given a F_g -coalgebra $c : X \rightarrow F_g X$, we define a game structure $G^c = (X_{\max}^c, X_{\min}^c, \tau^c)$ and a set $\text{Acc}^c \subseteq X_{\max}^c$ of accepting states as follows: $X_{\max}^c = X$, $X_{\min}^c = \mathcal{P}X$, $\tau^c = \{(x, A) \mid x \in X, A \in c_1(x)\} \cup \{(A, x') \mid A \in \mathcal{P}X, x' \in A\}$ and $\text{Acc}^c = \{x \in X \mid c_2(x) = 1\}$. Here we write $c(x) = (c_1(x), c_2(x)) \in \mathcal{P}^2 X \times \{0, 1\}$ for every x .

Conversely, given a game structure $G = (X_{\max}, X_{\min}, \tau)$ and a set $\text{Acc} \subseteq X_{\max}$, we define an F_g -coalgebra $c^{G, \text{Acc}} : X \rightarrow F_g X$ as follows: $X = X_{\max}$ and $c^{G, \text{Acc}}(x) = (\{\{x' \in X_{\max} \mid (y, x') \in \tau \mid y \in X_{\min}, (x, y) \in \tau\}, t)$ where t is 1 if $x \in \text{Acc}$ and 0 otherwise.

The above two transformations constitute an embedding-projection pair: games and F_g -coalgebra are almost equivalent; the former have additional freedom (in the choice of the set X_{\min}) that is however inessential.

Throughout the rest of this section, each categorical notion is accompanied by a concrete example in terms of two-player games. For readability, the details of these examples (they are all straightforward) are deferred to §A in the appendix. The other running example (PTSs) will be discussed later in §IV.

B. Modalities and Least Fixed-Point Properties, Categorically

Towards a categorical framework in which a soundness theorem is proved on the categorical level of abstraction, we need categorical modeling of modalities and least fixed-point properties. Our modeling here follows [17], [18]; it has been sketched in §I-B2.

The following function is heavily used in our developments.

Definition III.2 ($\Phi_{c,a}$). Let $F : \mathbf{Sets} \rightarrow \mathbf{Sets}$, $c : X \rightarrow FX$ be a coalgebra and $a : FA \rightarrow A$ be an algebra. We define a function $\Phi_{c,a} : A^X \rightarrow A^X$ by $\Phi_{c,a}(f) = a \circ Ff \circ c$, that is,

$$\left(X \xrightarrow{f} A \right) \xrightarrow{\Phi_{c,a}} \left(\begin{array}{ccc} FX & \xrightarrow{Ff} & FA \\ \uparrow c & & a \downarrow \\ X & & A \end{array} \right).$$

Then corecursiveness (Def. II.16) is rephrased as follows: $r : FR \rightarrow R$ is corecursive if and only if the function $\Phi_{c,r}$ has a unique fixed point for each $c : X \rightarrow FX$.

Our categorical modeling of *modality* is as follows.

Definition III.3 (a truth-value domain and an F -modality). A *truth-value domain* is a poset $(\Omega, \sqsubseteq_\Omega)$. If the order is clear from the context we simply write Ω . For a functor $F : \mathbf{Sets} \rightarrow \mathbf{Sets}$, an F -*modality* over the truth-value domain Ω is an F -algebra $\sigma : F\Omega \rightarrow \Omega$.

Example III.4. For two-player games (i.e. F_g -coalgebras) a natural truth-value domain is given by $(\{0, 1\}, \leq)$ where 1 stands for “true.” On top of this domain a natural F_g -modality $\sigma_g : F_g\{0, 1\} \rightarrow \{0, 1\}$ is given as follows.

$$\sigma_g(\Gamma, t) = \begin{cases} 1 & (t = 1) \\ \max_{A \in \Gamma} \min_{a \in A} a & (\text{otherwise}) \end{cases}$$

Here, in $(\Gamma, t) \in F_g X = \mathcal{P}^2 X \times \{0, 1\}$, $t \in \{0, 1\}$ indicates if the current state is accepting or not ($t = 1$ if yes). The second case in the above definition of $\sigma_g(\Gamma, t)$ reflects the intention that, in $\Gamma \in \mathcal{P}(\mathcal{P}X)$, the first \mathcal{P} is for the angelic player \max ’s choice while the second \mathcal{P} is for the demonic \min ’s.

Using an F -modality σ , liveness is categorically characterized as a least fixed-point property.

Definition III.5 ($\llbracket \mu \sigma \rrbracket_c$). Let $(\Omega, \sqsubseteq_\Omega)$ be a truth-value domain and $\sigma : F\Omega \rightarrow \Omega$ be a modality. We say that σ has *least fixed points* if for each $c : X \rightarrow FX$, the least fixed point of $\Phi_{c,\sigma} : \Omega^X \rightarrow \Omega^X$ (Def. III.2)—with respect to the pointwise extension of the order \sqsubseteq_Ω —exists. The least fixed point is called the (*coalgebraic*) *least fixed-point property* in c specified by σ , and is denoted by $\llbracket \mu \sigma \rrbracket_c : X \rightarrow \Omega$.

Example III.6. The F_g -modality $\sigma_g : F_g\{0, 1\} \rightarrow \{0, 1\}$ in Example III.4 has least fixed points (this follows from Prop. III.8 later). For each coalgebra $c : X \rightarrow F_g X$, the least fixed-point property $\llbracket \mu \sigma_g \rrbracket_c : X \rightarrow \{0, 1\}$ is concretely described by:

$$\llbracket \mu \sigma_g \rrbracket_c(x) = \begin{cases} 1 & (x \in \text{Reach}_{G^c, \text{Acc}^c}) \\ 0 & (\text{otherwise}). \end{cases}$$

Conversely, for each pair of a game structure G and a set Acc , $\text{Reach}_{G, \text{Acc}}$ is described by $\llbracket \mu \sigma_g \rrbracket_{c^{G, \text{Acc}}}$. See Prop. A.2 for a proof. This way we characterize reachability in two-player games in categorical terms.

C. Ranking Domains and Ranking Arrows

As we described in §I-B3, we understand *liveness checking* as the task of determining if $h \sqsubseteq \llbracket \mu\sigma \rrbracket_c$, for a given assertion $h : X \rightarrow \Omega$. Here we introduce our categorical machinery for providing witnesses to such satisfaction of liveness.

For simplicity of arguments we assume the following.

Assumption III.7. Let $F : \mathbf{Sets} \rightarrow \mathbf{Sets}$. We assume that a truth-value domain $(\Omega, \sqsubseteq_\Omega)$ and an F -modality $\sigma : F\Omega \rightarrow \Omega$ over Ω satisfy the following conditions.

- 1) The poset $(\Omega, \sqsubseteq_\Omega)$ is a complete lattice.
- 2) For each F -coalgebra $c : X \rightarrow FX$, the function $\Phi_{c,\sigma} : \Omega^X \rightarrow \Omega^X$ in Def. III.5 is monotone with respect to the pointwise extension of \sqsubseteq_Ω .

These assumptions are mild. For example, Cond. 2 is satisfied if: $F\Omega$ has an order structure; $\sigma : F\Omega \rightarrow \Omega$ is monotone; and the action $F_{X,\Omega} : \Omega^X \rightarrow (F\Omega)^{FX}$ of F on arrows is monotone, too. Cond. 1 in the above implies that Ω^X is a complete lattice. Thus we can construct a transfinite sequence $\perp_\Omega \sqsubseteq \Phi_{c,\sigma}(\perp_\Omega) \sqsubseteq \dots \sqsubseteq \Phi_{c,\sigma}^\alpha(\perp_\Omega) \sqsubseteq \dots$ as in Thm. II.18.2, to obtain the least fixed point of $\Phi_{c,\sigma}$ as its limit.

Proposition III.8. Under the conditions in Asm. III.7, σ has least fixed points (in the sense of Def. III.5). \square

Example III.9. The data F_g, σ_g for two-player games satisfy the assumptions: see Prop. A.3 (in the appendix) for a proof.

We are ready to introduce the key notions.

Definition III.10 (ranking domains). We assume Asm. III.7. Let $r : FR \rightarrow R$ be an F -algebra, $q : R \rightarrow \Omega$ be an arrow, and \sqsubseteq_R be a partial order on R . Note that for each set X , the order \sqsubseteq_Ω (resp. \sqsubseteq_R) extends to the one between functions $X \rightarrow \Omega$ (resp. $X \rightarrow R$) in a pointwise manner.

A triple (r, q, \sqsubseteq_R) is called a *ranking domain* for F and σ if the following conditions are satisfied.

- 1) We have $q \circ r \sqsubseteq_\Omega \sigma \circ Fq$ between arrows $FR \rightarrow \Omega$ (the square on the right in (7)).
- 2) The same conditions as in Asm. III.7 hold for r , i.e.
 - a) the poset (R, \sqsubseteq_R) is a complete lattice; and
 - b) for each $c : X \rightarrow FX$, the function $\Phi_{c,r} : R^X \rightarrow R^X$ (Def. III.2) is monotone.
- 3) The function $q : R \rightarrow \Omega$ is monotone (i.e. $a \sqsubseteq_R b \Rightarrow q(a) \sqsubseteq_\Omega q(b)$), strict (i.e. $q(\perp_R) = \perp_\Omega$) and continuous (i.e. for each subset $K \subseteq R$, we have $q(\bigsqcup_{a \in K} a) = \bigsqcup_{a \in K} q(a)$).
- 4) The algebra $r : FR \rightarrow R$ is corecursive.

Cond. 2 in the definition ensures that the least fixed point of $\Phi_{c,r}$ arises from the approximation sequence in Thm. II.18.2. Cond. 3 ensures that this least fixed point is preserved by q . In particular we insist on strictness—this is much like in domain theory [30]. The most significant in Def. III.10 is the

corecursiveness of r (Cond. 4): it makes r a refinement of σ that is suited for detecting well-foundedness.

Definition III.11 (ranking arrows). Let (r, q, \sqsubseteq_R) be a ranking domain for F and σ ; and $c : X \rightarrow FX$ be a coalgebra. An arrow $b : X \rightarrow R$ is called a (*coalgebraic*) *ranking arrow* for c with respect to (r, q, \sqsubseteq_R) if it satisfies $b \sqsubseteq_R \Phi_{c,r}(b) = r \circ Fb \circ c$ (the square on the left in (7)).

Now we give a soundness theorem for (categorical) ranking arrows. This is the main theorem of this paper; its proof demonstrates the role of the corecursiveness assumption.

Theorem III.12 (soundness). Let (r, q, \sqsubseteq_R) be a ranking domain. Let $c : X \rightarrow FX$ be an F -coalgebra and $b : X \rightarrow R$ be a ranking arrow for c (i.e. $b \sqsubseteq r \circ Fb \circ c$). Then we have:

$$q \circ b \sqsubseteq_\Omega \llbracket \mu\sigma \rrbracket_c.$$

$$\begin{array}{ccccc} FX & \xrightarrow{Fb} & FR & \xrightarrow{Fq} & F\Omega \\ \uparrow c & \sqcup & \downarrow r & \sqsubseteq & \downarrow \sigma \\ X & \xrightarrow{b} & R & \xrightarrow{q} & \Omega \end{array}$$

$\llbracket \mu\sigma \rrbracket_c$

Thus for liveness checking (i.e. for proving $h \sqsubseteq \llbracket \mu\sigma \rrbracket_c$) it suffices to find a ranking arrow b such that $h \sqsubseteq q \circ b$. In the proof of the theorem we use the following generalization of Thm. II.18.2. It starts from a post-fixed point l (not from \perp).

Lemma III.13. Assume the conditions in Thm. II.18, and let l be a post-fixed point of f , i.e. $l \sqsubseteq f(l)$. Then we can define a transfinite sequence $l \sqsubseteq f(l) \sqsubseteq \dots \sqsubseteq f^\alpha(l) \sqsubseteq \dots$ in a similar manner to Thm. II.18.2. The sequence eventually stabilizes and its limit is a (not necessarily least) fixed point of f . \square

Proof (Thm. III.12). By Cond. 2a in Def. III.10, the poset (R^X, \sqsubseteq_R) is a complete lattice. Moreover, by its definition, $b : X \rightarrow R$ is a post-fixed point of $\Phi_{c,r}$. Hence together with Cond. 2b, we can construct a transfinite sequence $b \sqsubseteq_R \Phi_{c,r}(b) \sqsubseteq_R \dots \sqsubseteq_R \Phi_{c,r}^\alpha(b) \sqsubseteq_R \dots$ as in Lem. III.13. By Lem. III.13, there exists an ordinal m such that $\Phi_{c,r}^m(b)$ is a fixed point of $\Phi_{c,r}$. By its definition, we have $b \sqsubseteq_R \Phi_{c,r}^m(b)$.

Note here that r is assumed to be a corecursive algebra (Cond. 4 in Def. III.10). Hence $\Phi_{c,\sigma}$ has a unique fixed point; it is denoted by $\llbracket c \rrbracket_r : X \rightarrow R$. Then we have:

$$b \sqsubseteq_R \Phi_{c,r}^m(b) = \llbracket c \rrbracket_r. \quad (8)$$

By Cond. 2a in Def. III.10, R^X is a complete lattice. Hence we can also define $\Phi_{c,r}^\alpha(\perp_R) : X \rightarrow R$ for each ordinal α (here \perp_R denotes the least element in R^X), and by Thm. II.18.2, there exists

m' such that $\Phi_{c,r}^{m'}(\perp_R)$ is also a fixed point of $\Phi_{c,r}$. Hence $\Phi_{c,r}^{m'}(\perp_R) = \llbracket c \rrbracket_r$.

We shall now prove that $q \circ \Phi_{c,r}^\alpha(\perp_R) \sqsubseteq \llbracket \mu\sigma \rrbracket_c$ holds for each ordinal α . This is by transfinite induction on α .

For $\alpha = 0$, we have:

$$\begin{aligned} q \circ \Phi_{c,r}^0(\perp_R) &= q \circ \perp_R && \text{(by definition)} \\ &= \perp_\Omega && \text{(by Cond. 3 in Def. III.10)} \\ &\sqsubseteq_\Omega \llbracket \mu\sigma \rrbracket_c. \end{aligned}$$

For a successor ordinal $\alpha + 1$, we have:

$$\begin{aligned}
q \circ \Phi_{c,r}^{\alpha+1}(\perp_R) &= q \circ r \circ F(\Phi_{c,r}^\alpha(\perp_R)) \circ c && \text{(by definition)} \\
\sqsubseteq_\Omega \sigma \circ F(q \circ \Phi_{c,r}^\alpha(\perp_R)) \circ c &&& \text{(by Cond. 1 in Def. III.10)} \\
\sqsubseteq_\Omega \sigma \circ F(\llbracket \mu\sigma \rrbracket_c) \circ c &&& \text{(by IH and Asm. III.7.2)} \\
= \llbracket \mu\sigma \rrbracket_c &&& (\llbracket \mu\sigma \rrbracket_c \text{ is a fixed point})
\end{aligned}$$

For a limit ordinal ι , we have:

$$\begin{aligned}
q \circ \Phi_{c,r}^\iota(\perp_R) &= q(\bigsqcup_{\alpha < \iota} \Phi_{c,r}^\alpha(\perp_R)) && \text{(by definition)} \\
= \bigsqcup_{\alpha < \iota} (q \circ \Phi_{c,r}^\alpha(\perp_R)) &&& \text{(by Cond. 3 in Def. III.10)} \\
\sqsubseteq_\Omega \llbracket \mu\sigma \rrbracket_c &&& \text{(by IH)}.
\end{aligned}$$

As $\Phi_{c,r}^{\iota'}(\perp_R) = \langle c \rangle_r$, the last fact yields $q \circ \langle c \rangle_r \sqsubseteq \llbracket \mu\sigma \rrbracket_c$. Combining with (8) and the monotonicity of q , we obtain $q \circ b \sqsubseteq q \circ \langle c \rangle_r \sqsubseteq \llbracket \mu\sigma \rrbracket_c$. This concludes the proof. \square

Remark III.14. Note that the requirement on ranking arrows— $b \sqsubseteq_R r \circ Fb \circ c$ (Def. III.11)—is a *local* one: it only involves one-step transitions by c and hence is easy to check.

The condition asserts that b is a suitable *post-fixed point*. In view of the order-theoretic foundations in §II-D this might seem strange: we are using an invariant-like construct b to witness a *least* fixed point, not a greatest. We are allowed to do so thanks to the corecursiveness of $r : FR \rightarrow R$ —here the least and greatest fixed points for $\Phi_{c,r}$ coincide. It is also crucial that q preserves least fixed points, being strict and continuous.

Example III.15. For two-player games as F_g -coalgebras, we can define a ranking domain

$$(r_{g,\mathfrak{z}} : F_g \mathbf{Ord}_{\leq \mathfrak{z}} \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}, q_{g,\mathfrak{z}} : \mathbf{Ord}_{\leq \mathfrak{z}} \rightarrow \{0, 1\}, \sqsubseteq_{\mathbf{Ord}})$$

as follows.

- 1) $\mathbf{Ord}_{\leq \mathfrak{z}} = \{\alpha \mid \alpha \text{ is an ordinal s.t. } \alpha \leq \mathfrak{z}\}$, and

$$r_{g,\mathfrak{z}}(\Gamma, t) = \begin{cases} 0 & (t = 1) \\ \min_{A \in \Gamma} \sup_{a \in A} (a \hat{+} 1) & \text{(otherwise)}. \end{cases}$$

- 2) $q_{g,\mathfrak{z}}(\mathfrak{z}) = 0$, and $q_{g,\mathfrak{z}}(\alpha) = 1$ for any α such that $\alpha < \mathfrak{z}$;

- 3) $\alpha \sqsubseteq_{\mathbf{Ord}} b \stackrel{\text{def}}{\iff} \alpha \geq b$ (note the directions of inequalities).

Recall that $\alpha \hat{+} 1$ denotes $\min\{\alpha + 1, \mathfrak{z}\}$. The triple $(r_{g,\mathfrak{z}}, q_{g,\mathfrak{z}}, \sqsubseteq_{\mathbf{Ord}})$ is indeed a ranking domain (see Prop. A.4 in the appendix). One can think of the above data as a classifier for (non-)well-foundedness: all the ordinals $\alpha < \mathfrak{z}$ are for “well-founded” and the maximum ordinal \mathfrak{z} is for “non-well-founded.” Observe that the map $q_{g,\mathfrak{z}}$ acts accordingly.

We indeed have the following correspondences.

- a) $b : X \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}$ is a (categorical) ranking arrow (Def. III.11) for an F_g -coalgebra $c : X \rightarrow F_g X$ wrt. $(r_{g,\mathfrak{z}}, q_{g,\mathfrak{z}}, \sqsubseteq_{\mathbf{Ord}})$ iff b is a ranking function for G^c and Acc^c (in the conventional sense of Def. II.4).

- b) $b : X \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}$ is a ranking function for a game structure G and a set Acc iff b is a ranking arrow for $c^{G, \text{Acc}}$ wrt. $(r_{g,\mathfrak{z}}, q_{g,\mathfrak{z}}, \sqsubseteq_{\mathbf{Ord}})$.
- c) $b(x) < \mathfrak{z}$ iff $q_{g,\mathfrak{z}} \circ b(x) = 1$.

Here recall the correspondence in Def. III.1. A formal statement and its proof are found in Prop. A.7.

Combined with the characterization in Example III.6, we conclude that the conventional soundness result (Thm. II.5) is an instance of our categorical soundness (Thm. III.12).

Remark III.16. Assume the conditions in Thm. III.12. As $r : FR \rightarrow R$ is a corecursive algebra, there exists a unique arrow $\langle c \rangle_r : X \rightarrow R$ such that $\langle c \rangle_r = r \circ F \langle c \rangle_r \circ c$. Since $\langle c \rangle_r$ is obviously the greatest fixed point of $\Phi_{c,\sigma}$, by the Knaster–Tarski theorem (cf. Thm. II.18), for each (categorical) ranking arrow $b : X \rightarrow R$ we have $b \sqsubseteq_R \langle c \rangle_r$. This means that if $h \sqsubseteq_\Omega q \circ b$ then $h \sqsubseteq_\Omega q \circ \langle c \rangle_r$. Therefore we can say that the unique arrow $\langle c \rangle_r : X \rightarrow R$ is the “optimal” ranking arrow in the sense that if $\langle c \rangle_r$ cannot prove liveness then no (categorical) ranking arrow can prove liveness using R and q , either. For two-player games, the optimal ranking arrow is given by the function assigning a state x the minimum number of steps from x to Acc .

We note that the converse of Thm. III.12 (i.e. *completeness*) does not necessarily hold. In other words, it is possible that there exists no ranking arrow $b : X \rightarrow R$ such that $q \circ b = \llbracket \mu\sigma \rrbracket_c$. Here is a counterexample.

Example III.17. We define an F_g -coalgebra $c : X \rightarrow F_g X$ by

$X = \{x_\alpha \mid \alpha \leq \omega\}$, $c(x_0) = (\emptyset, 1)$ and $c(x_\alpha) = (\{\{x_\beta \mid \beta < \alpha\}, 0)$ for each $\alpha > 0$.


Note that in the corresponding game structure G_c , all the choices are made by the player min. Then we have $\llbracket \mu\sigma_g \rrbracket_c(x_\omega) = 1$ because of well-foundedness of ω . However, the unique arrow $\langle c \rangle_{r_{g,\omega}} : X \rightarrow \mathbf{Ord}_{\leq \omega}$ such that $\langle c \rangle_{r_{g,\omega}} = \Phi_{c,\sigma_g}(\langle c \rangle_{r_{g,\omega}})$ assigns, to each state x_α , the ordinal α . This means that $q_{g,\omega} \circ \langle c \rangle_{r_{g,\omega}}(x_\omega) = q_{g,\omega}(\omega) = 0$. Thus $q_{g,\omega} \circ \langle c \rangle_{r_{g,\omega}} < \llbracket \mu\sigma_g \rrbracket_c$.

Similarly, for every ordinal number \mathfrak{z} , we can construct an F_g -coalgebra whose reachability cannot be proved by the ranking domain $r_{g,\mathfrak{z}} : F_g \mathbf{Ord}_{\leq \mathfrak{z}} \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}$.

By cardinality arguments we can show that sort of “completeness” holds in the example above, in the following sense: for every F_g -coalgebra c there exists an ordinal \mathfrak{z} such that the reachability of c is provable by the ranking domain $r_{g,\mathfrak{z}}$. However, in this paper we use the term “completeness” in a different sense in which we fix the domain R of ranking functions in advance.

Here is a categorical sufficient condition for completeness.

Proposition III.18 (a sufficient condition for completeness). Let (r, q, \sqsubseteq_R) be a ranking domain, $c : X \rightarrow F_g X$ be an F_g -coalgebra, and $\langle c \rangle_r : X \rightarrow R$ be the unique arrow such



that $(\llbracket c \rrbracket)_r = r \circ F(\llbracket c \rrbracket)_r \circ c$. Assume that we have the equality

$$q \circ r = \sigma \circ Fq, \quad (9)$$

instead of an inequality, in the square on the right. Then we have $q \circ (\llbracket c \rrbracket)_r = \llbracket \mu\sigma \rrbracket_c$. \square

Intuitively, the equality (9) means that r approximates the modality σ in an adequate way. The result implies that, in case $h: X \rightarrow \Omega$ satisfies $h \sqsubseteq \llbracket \mu\sigma \rrbracket_c$, the latter inequality can always be witnessed by some ranking arrow (namely $(\llbracket c \rrbracket)_r$). This is completeness of the proof method of categorical ranking arrows. An example of a complete ranking domain will be given in §IV-C.

IV. CATEGORICAL RANKING ARROWS FOR PROBABILISTIC TRANSITION SYSTEMS

We shall now investigate what our categorical framework in §III entails in the probabilistic setting of §II-B. It turns out that the well-known definition of ranking supermartingale (ε -additive ones in Def. II.11) is *not* an instance. Here we study some variations of the definition of ranking supermartingale; two among them (*distribution-valued* and *non-counting* ones, that are new to our knowledge) exhibit the nice categorical properties in §III. We also discuss some relationships between those variations, showing that the soundness of ε -additive ranking supermartingales (Def. II.11) can nevertheless be proved via the categorical arguments in §III.

A. Probabilistic Transition Systems as Coalgebras

To represent a PTS as a coalgebra, we use the functor $F_p: \mathbf{Sets} \rightarrow \mathbf{Sets}$ (p stands for “probability”) defined as follows.

Definition IV.1 (F_p). We let $F_p = \mathcal{D}(_) \times \{0, 1\}$, where \mathcal{D} is the (discrete) distribution functor in Def. II.15.

For an F_p -coalgebra $c: X \rightarrow F_p X$, we define a PTS $M^c = (X^c, \tau^c)$ and a set $\text{Acc}^c \subseteq X^c$ of accepting states by: $X^c = X$, $\tau^c(x) = c_1(x)$ for each $x \in X^c$, and $\text{Acc}^c = \{x \in X \mid c_2(x) = 1\}$. Here we write $c(x) = (c_1(x), c_2(x)) \in \mathcal{D}X \times \{0, 1\}$ for every x .

Conversely, for a PTS $M = (X, \tau)$ and a set $\text{Acc} \subseteq X$, we define an F_p -coalgebra $c^{M, \text{Acc}}: X \rightarrow F_p X$ by $c^{M, \text{Acc}}(x) = (\tau(x), t)$ where t is 1 if $x \in \text{Acc}$ and 0 otherwise.

These correspondences are indeed bijective. Analogously to Example III.6, we characterize reachability probabilities of a PTS as a coalgebraic least fixed-point property.

Proposition IV.2. We define an F_p -modality $\sigma_p: F_p[0, 1] \rightarrow [0, 1]$ over the truth-value domain $([0, 1], \leq)$ as follows. Here $[0, 1]$ is the unit interval and \leq is the usual order on it.

$$\sigma_p(\delta, t) = \begin{cases} 1 & (t = 1) \\ \sum_{a \in \text{supp}(\delta)} a \cdot \delta(a) & (\text{otherwise}). \end{cases}$$

Note here that $\delta \in \mathcal{D}[0, 1]$ and $t \in \{0, 1\}$.

Then σ_p satisfies Asm. III.7 and thus has least fixed points (in the sense of Def. III.5). The lfp property $\llbracket \mu\sigma_p \rrbracket_c$ coincides with the reachability probability function $\text{Reach}_{M^c, \text{Acc}^c}: X \rightarrow [0, 1]$ of the corresponding PTS (Def. II.9). \square

$$\begin{array}{ccccc} FX & \xrightarrow{F \llbracket \mu\sigma \rrbracket_c} & FR & \xrightarrow{Fq} & F\Omega \\ \uparrow c & \xrightarrow{F(\llbracket c \rrbracket)_r} & \downarrow r & \xrightarrow{=} & \downarrow \sigma \\ X & \xrightarrow{(\llbracket c \rrbracket)_r} & R & \xrightarrow{q} & \Omega \\ & & \llbracket \mu\sigma \rrbracket_c & & \end{array}$$

B. Known Variations: ε -Additive and α -Multiplicative Ranking Supermartingales

The definition of ranking supermartingale that we have reviewed (ε -additive ones in Def. II.11) is not an instance of our categorical notion (Def. III.10). Specifically, its value domain (the interval $[0, \infty]$ with a suitable F_p -algebraic structure) fails to be corecursive. As a result, soundness of additive ranking supermartingale (Thm. II.12) cannot be directly proved using our categorical soundness theorem (Thm. III.12).

The following is an attempt to define a ranking domain for ε -additive supermartingales. Let us fix a real number $\varepsilon > 0$ and define an F_p -algebra $r'_{p, \varepsilon}: F_p[0, \infty] \rightarrow [0, \infty]$, an arrow $q'_p: [0, \infty] \rightarrow [0, 1]$ and a partial order $\sqsubseteq_{[0, \infty]}$ over $[0, \infty]$ as follows.

1) For each $(\psi, t) \in F_p[0, \infty] = \mathcal{D}[0, \infty] \times \{0, 1\}$,

$$r'_{p, \varepsilon}(\psi, t) = \begin{cases} 0 & (t = 1) \\ (\sum_{a \in \text{supp}(\psi)} a \cdot \psi(a)) + \varepsilon & (\text{otherwise}). \end{cases}$$

2) $q'_p(\infty) = 0$ and $q'_p(a) = 1$ if $a < \infty$.

3) $a \sqsubseteq_{[0, \infty]} b \stackrel{\text{def}}{\iff} a \geq b$ (note the direction).

Proposition IV.3. In this setting, for each $c: X \rightarrow F_p X$ and $b': X \rightarrow [0, \infty]$, we have the following.


- b' is an ε -additive ranking supermartingale (Def. II.11) iff b' satisfies $b' \sqsubseteq_{[0, \infty]} r'_{p, \varepsilon} \circ b' \circ c$.
- $b'(x) \neq \infty$ iff $q'_p \circ b'(x) = 1$. \square

Therefore the triple $(r'_{p, \varepsilon}, q'_p, \sqsubseteq_{[0, \infty]})$ is suited for accommodating ε -additive supermartingales in our categorical framework. Unfortunately it is not a ranking domain (Def. III.10).

Proposition IV.4. The triple $(r'_{p, \varepsilon}, q'_p, \sqsubseteq_{[0, \infty]})$ introduced above satisfies the conditions of a ranking domain (Def. III.10), except for Cond. 4. \square

Example IV.5. We define a coalgebra


$c: X \rightarrow F_p X$ by: $X = \{x_0, x_1, x_2, x_3\}$, $c(x_0) = ([x_1 \mapsto \frac{1}{2}, x_2 \mapsto \frac{1}{2}], 0)$, $c(x_1) = ([x_1 \mapsto \frac{1}{2}, x_3 \mapsto \frac{1}{2}], 0)$, $c(x_2) = ([x_3 \mapsto 1], 0)$ and $c(x_3) = ([x_3 \mapsto 1], 1)$. The corresponding PTS is depicted on the right.



Let $\varepsilon > 0$. For this coalgebra, we define arrows $b_1, b_2: X \rightarrow [0, \infty]$ by: $b_1(x_0) = \frac{5}{2}\varepsilon$, $b_1(x_1) = 2\varepsilon$, $b_1(x_2) = \varepsilon$ and $b_1(x_3) = 0$; and $b_2(x_0) = \infty$, $b_2(x_1) = \infty$, $b_2(x_2) = \varepsilon$ and $b_2(x_3) = 0$. Both of these qualify as coalgebra-algebra homomorphisms from c to $r'_{p, \varepsilon}$. Therefore $r'_{p, \varepsilon}$ is not corecursive.

It is well-known that ε -additive supermartingales (Def. II.11) witness *positive almost-sure reachability* [31], that is, the expected number of steps to accepting states is finite. This is a property strictly stronger than almost-sure reachability (see Example IV.6). It follows that ε -additive supermartingales are not complete against almost-sure reachability.

Example IV.6. We define $c : X \rightarrow F_p X$ by $X = \{x\} \cup \{x_{i,j} \mid i, j \in \mathbb{N}, 1 \leq i, 1 \leq j \leq 2^i\}$; and $c(x) = (\delta, 0)$, where $\delta(x_{i,j})$ is $\frac{1}{2^i}$ if $j = 0$ and 0 otherwise, $c(x_{i,j}) = ([x_{i,j+1} \mapsto 1], 0)$ for each i and $j < 2^i$, and $c(x_{i,2^i}) = ([x_{i,2^i} \mapsto 1], 1)$. The corresponding



PTS M^c and the set Acc^c of accepting states are shown on the right. This system when run from x is clearly almost-sure terminating; however the expected number of steps to accepting states is infinite.

Another known variation of ranking supermartingales is given by *multiplicative ranking supermartingales* [23]. Let $\alpha \in (0, 1)$. A function $b : X \rightarrow [0, \infty]$ is an α -multiplicative ranking supermartingale if we have

$$\forall x \in X \setminus \text{Acc}. \sum_{x' \in \text{supp}(\tau(x))} \tau(x)(x') \cdot b(x') \leq \alpha \cdot b(x),$$

and moreover there exists $\delta > 0$ such that $b(x) \geq \delta$ for each $x \in X \setminus \text{Acc}$. For this multiplicative variation, results analogous to Prop. IV.3 and Prop. IV.4 hold (see §G in the appendix for the proofs).

C. Distribution-Valued Ranking Functions

Let us turn to possible instantiations of our categorical framework in the current probabilistic setting. The first uses \mathcal{DN}_∞ , the set of distributions over extended natural numbers, as a ranking domain (instead of $[0, \infty]$). In what follows, given a probability distribution γ over \mathbb{N}_∞ and $a, b \in \mathbb{N}$, $\gamma([a, b])$ denotes $\sum_{i=a}^b \gamma(i)$ and $\gamma([a, \infty))$ denotes $\sum_{i=a}^\infty \gamma(i)$.

Proposition IV.7. Recall that $\mathbb{N}_\infty = \mathbb{N} \cup \{\infty\}$ and \mathcal{DN}_∞ collects all the distributions over \mathbb{N}_∞ . We define an F_p -algebra $r_p : F_p \mathcal{DN}_\infty \rightarrow \mathcal{DN}_\infty$, a function $q_p : \mathcal{DN}_\infty \rightarrow [0, 1]$ and a partial order $\sqsubseteq_{\mathcal{DN}_\infty}$ over \mathcal{DN}_∞ as follows.

- 1) For each $(\Gamma, t) \in F_p \mathcal{DN}_\infty = \mathcal{D}^2 \mathbb{N}_\infty \times \{0, 1\}$,

$$r_p(\Gamma, t)(a) = \begin{cases} 1 & (t = 1, a = 0) \\ 0 & (t = 1, a > 0 \text{ or } t = 0, a = 0) \\ \sum_{\gamma \in \text{supp}(\Gamma)} \Gamma(\gamma) \cdot \gamma(a-1) & (t = 0, a > 0). \end{cases}$$

- 2) $q_p(\varphi) = \varphi([0, \infty))$.
- 3) $\varphi \sqsubseteq_{\mathcal{DN}_\infty} \varphi' \stackrel{\text{def}}{\iff} \forall a \in \mathbb{N}. \varphi([0, a]) \leq \varphi'([0, a])$.

Then the triple $(r_p, q_p, \sqsubseteq_{\mathcal{DN}_\infty})$ is a ranking domain. Moreover we have $q_p \circ r_p = \sigma_p \circ F_p q_p$ (cf. (9) in Prop. III.18). \square

Intuitively, the value $b(x)([0, a]) \in [0, 1]$ under-approximates the probability with which an accepting state is reached from x within a steps. The definition of $\sqsubseteq_{\mathcal{DN}_\infty}$, which is much like in probabilistic powerdomains (see e.g. [32]), also reflects this intuition. Here the Dirac distribution δ_0 (resp. δ_∞) is the greatest (resp. least) element.

The definition of ranking arrow (Def. III.11) instantiates to the following—as one sees by straightforward calculation—when we fix a ranking domain to be the one in Prop. IV.7.

Definition IV.8 (distribution-valued ranking function). Let $M = (X, \tau)$ be a PTS and $\text{Acc} \subseteq X$. A *distribution-valued ranking function* is $b : X \rightarrow \mathcal{DN}_\infty$ such that:

$$\sum_{x' \in \text{supp}(\tau)} \tau(x)(x') \cdot b(x')([0, a-1]) \geq b(x)([0, a])$$

for each $x \in X \setminus \text{Acc}$ and $a \in \mathbb{N}_\infty$. Here we let $b(x')([0, -1]) = 0$.

By Thm. III.12 (soundness) we have the following: given a PTS $c : X \rightarrow F_p X$ and an “assertion” $h : X \rightarrow [0, 1]$, if there exists a distribution-valued ranking function $b : X \rightarrow \mathcal{DN}_\infty$ such that $h \leq q_p \circ b$, then we can conclude that $h \leq \text{Reach}_{M^c, \text{Acc}^c}$. Here $\text{Reach}_{M^c, \text{Acc}^c}$ is given by reachability probabilities and coincides with $\llbracket \mu \sigma_p \rrbracket_c$ (Prop. IV.2).

We note that *quantitative verification* is possible using distribution-valued ranking functions. For example an assertion $h : X \rightarrow [0, 1]$ can be such that $h(x) = 1/2$; by finding a suitable arrow b we conclude $h \leq \text{Reach}_{M^c, \text{Acc}^c}$, that is, that the reachability probability from x is at least $1/2$.³ Such a quantitative assertion cannot be verified using ε -additive supermartingales: as in Thm. II.12 they only witness the *qualitative* property of (positive) almost-sure reachability.

Example IV.9. We define a PTS $M =$

(X, τ) by $X = \{x_0, x_1, x_2\}$, $\tau(x_0) = [x_0 \mapsto \frac{1}{3}, x_1 \mapsto \frac{1}{3}, x_2 \mapsto \frac{1}{3}]$, $\tau(x_1) = [x_1 \mapsto 1]$ and $\tau(x_2) = [x_2 \mapsto 1]$. Let $\text{Acc} = \{x_1\}$. The function $b : X \rightarrow \mathcal{DN}_\infty$ defined by $b(x_0) = [i \mapsto 1/3^{i+1}, \infty \mapsto 1/2]$, $b(x_1) = [0 \mapsto 1]$ and $b(x_2) = [\infty \mapsto 1]$ is a distribution-valued ranking function. This allows us to conclude $\text{Reach}_{M, \text{Acc}}(x_0) \geq (1 - \frac{1}{2}) = \frac{1}{2}$.

Finally we exhibit *completeness* of distribution-valued ranking functions. This is an immediate consequence of the categorical result (Prop. III.18) and Prop. IV.7.

Proposition IV.10. For each F_p -coalgebra $c : X \rightarrow F_p X$, there exists a (categorical) ranking arrow $b : X \rightarrow \mathcal{DN}_\infty$ such that $q_p \circ b = \llbracket \mu \sigma_p \rrbracket_c$. \square

Example IV.11. In Example IV.6, the function $b : X \rightarrow \mathcal{DN}_\infty$ defined by

$$b(x)(n) = \begin{cases} 1/2^i & (n = 2^i, i > 0) \\ 0 & (\text{otherwise}) \end{cases} \quad \text{and} \quad b(x_{i,j}) = \delta_{2^i - j}$$

is a distribution-valued ranking function (here $\delta_{2^i - j}$ denotes a Dirac distribution). We have $b(x)([0, \infty)) = 1$ and thus successfully verify almost-sure reachability from x .

In §IV-B we have argued that ε -additive ranking supermartingales (Def. II.11) is not an instance of our categorical ranking arrow. It is nevertheless possible to prove its soundness (Thm. II.5) using the categorical framework—specifically by showing that an ε -additive ranking supermartingale gives rise to a distribution-valued ranking function (Def. IV.8). Details are found in §I in the appendix.

³The problem solved here is more precisely the *threshold reachability checking* problem; and ranking function-based proof methods for the problem would be viable options especially when the state space X is infinite.

Similarly our framework can prove soundness of α -multiplicative ranking supermartingales (see §J for the details).

D. γ -Scaled Non-Counting Ranking Supermartingales

The notion of distribution-valued ranking function exhibits pleasant properties like completeness and quantitative assertion checking. A major drawback, however, is the complexity of its value domain \mathcal{DN}_∞ .

In many realistic verification scenarios a ranking function/supermartingale b would be *synthesized* as follows: the function b is expressed in a predetermined template $b_{\vec{p}}$ (such as polynomials up-to a certain degree) in which some parameters \vec{p} occur; the requirements on $b_{\vec{p}}$ translate to constraints on \vec{p} ; and one relies on some optimization solver (for SAT, LP, SDP, etc.) to solve the constraints. It significantly increases the complexity of the workflow if the value $b(x)$ is a distribution in \mathcal{DN}_∞ instead of an (extended) real number in $[0, \infty]$.

Here we present another probabilistic instantiation of the categorical framework. It takes values in the unit interval $[0, 1]$.

Proposition IV.12. *We fix a real number $\gamma \in [0, 1)$. We define an algebra $r_{\text{nc}, \gamma} : F_{\mathbb{P}}[0, 1] \rightarrow [0, 1]$ (here nc stands for “non-counting”) as follows.*

$$r_{\text{nc}, \gamma}(\varphi, t) = \begin{cases} 1 & (t = 1) \\ \gamma \cdot \sum_{a \in \text{supp}(\varphi)} a \cdot \varphi(a) & (\text{otherwise}). \end{cases}$$

We further define $q_{\text{nc}} : [0, 1] \rightarrow [0, 1]$ by $q_{\text{nc}}(a) = a$. Then $(r_{\text{nc}, \gamma}, q_{\text{nc}}, \leq)$, where \leq is the usual on $[0, 1]$, is a ranking domain with respect to the modality $\sigma_{\mathbb{P}}$ (Prop. IV.2). \square

Thus it makes sense to consider ranking arrows (Def. III.11) with respect to the ranking domain $(r_{\text{nc}, \gamma}, q_{\text{nc}}, \leq)$. By straightforward calculation, their definition unravels as follows.

Definition IV.13 (γ -scaled non-counting ranking supermartingale). Let $M = (X, \tau)$ be a PTS and $\text{Acc} \subseteq X$. We fix a real number γ such that $0 \leq \gamma < 1$. A γ -scaled non-counting ranking supermartingale is a function $b : X \rightarrow [0, 1]$ such that for each $x \in X \setminus \text{Acc}$ we have:

$$\gamma \cdot \sum_{x' \in X} \tau(x)(x') \cdot b(x') \geq b(x).$$

This notion of supermartingale seems new. The intuition is that $b(x)$ is a lower bound for the reachability probability. Note that, unlike for the other variations of supermartingales in this section (where we *over*-approximate the number of steps), reachability probabilities should be *under*-approximated.

We obtain the following soundness result as an instance of Thm. III.12. Here a non-counting ranking supermartingale b itself gives lower bounds for reachability probabilities since $q_{\text{nc}} : [0, 1] \rightarrow [0, 1]$ is the identity map.

Corollary IV.14. *Let $M = (X, \tau)$ be a PTS and $b : X \rightarrow [0, 1]$ be a γ -scaled non-counting ranking supermartingale. Then we have $\text{Reach}_{\text{Acc}}(x) \geq b(x)$. \square*

Example IV.15. Consider the PTS M and Acc in Example IV.9. For each $\gamma \in [0, 1)$, we define $b_\gamma : X \rightarrow [0, 1]$ by $b_\gamma(x_0) = \frac{\gamma}{3-\gamma}$, $b_\gamma(x_1) = 1$ and $b_\gamma(x_2) = 0$. Then b_γ is seen

to be a γ -scaled non-counting ranking supermartingale. By Cor. IV.14 we have $\text{Reach}_{\text{Acc}}(x_0) \geq \frac{\gamma}{3-\gamma}$; as this holds for any $\gamma \in [0, 1)$, by letting $\gamma \rightarrow 1$, we conclude $\text{Reach}_{\text{Acc}}(x_0) \geq \frac{1}{2}$.

We note that in general a scaling factor $\gamma \in [0, 1)$ results in suboptimality of under-approximation of reachability properties: in the last example $b_\gamma(x_0) = \frac{\gamma}{3-\gamma}$ is smaller than the reachability probability $1/2$. Such suboptimality is an issue especially when we aim at *qualitative* verification of almost-sure reachability. In the last example we exercised an *asymptotic* argument in which we think of γ as a free variable and take the limit under $\gamma \rightarrow 1$. This strategy can be employed for almost-sure reachability checking.

Finally, the following example demonstrates that application of non-counting supermartingales is not limited to *positive* almost-sure reachability.

Example IV.16. We define a PTS $M = (X, \tau)$ and $\text{Acc} \subseteq X$ as in Example IV.6. For each γ , if we define $b_\gamma : X \rightarrow [0, 1]$ by $b_\gamma(x) = \sum_{i=1}^{\infty} \frac{\gamma^{2^i}}{2^i}$ and $b_\gamma(x_{i,j}) = \gamma^{2^i - j}$ for each i and j , then it is a γ -scaled non-counting supermartingale. Hence we have $\text{Reach}_{M, \text{Acc}}(x) \geq \lim_{\gamma \rightarrow 1} b_\gamma(x) = 1$.

Let us summarize the section. We presented four variations of ranking supermartingales: ε -additive ones, α -multiplicative ones, distribution-valued ranking functions and γ -scaled non-counting ranking supermartingales. The former two are known in the literature while the latter two seem to be new. The known notions are not instances of our generic definition, but their soundness can be derived via our generic theory (see the end of §IV-C). Among the (seemingly) new notions, distribution-valued ranking functions enjoy nice properties like completeness (Prop. IV.10) and support of quantitative reasoning (see Example IV.9)—at the cost of their complexity (they take as values distributions in \mathcal{DN}_∞). Non-counting ranking supermartingales (whose values are simply real numbers) are advantageous in quantitative reasoning (see Example IV.15) and non-positive almost-sure termination, but the scaling factor γ in it leads to suboptimal approximation. Typically one needs to rely on asymptotic arguments to obtain sharp bounds.

V. CONCLUSIONS AND FUTURE WORK

We have given a categorical account for liveness checking: we identify the essence of ranking function-based proof methods as the combination of corecursive algebras (as value domains) and lax homomorphisms; and for our notion of ranking arrow a soundness theorem has been presented. Our leading examples have been two-player games and probabilistic transition systems; in the course of studying them we were led to (seemingly) new variations of ranking martingales.

Besides the concrete examples of “ranking functions” in this paper, we wish to derive yet other concrete examples from our categorical modeling, so that they provide novel proof methods for various liveness properties. A possible direction towards this goal is discussed in §L in the appendix, motivated by categorical closure properties of corecursive algebras. Some abstract categorical questions remain open, too, such as

characterization of the *non-well-founded part*—that represents failure of liveness properties—of a corecursive algebra. We are also interested in the relationship between these (expected) results/examples and *productivity* for coinductive datatypes in functional programming [35]. Intuitively, the latter is a property that any finite prefix of a coinductively defined data is obtained in finite time.

We have used two-player games (systems with angelic and demonic transitions) and PTSs (systems with probabilistic transitions) as leading examples. A natural direction of future work is to consider stochastic games, which involve angelic, demonic and probabilistic transitions [36].

In this paper we have focused on least fixed-point properties. Extension to nested fixed-point specifications—*persistence*, *recurrence*, and general fixed-point formulas—is important future work. There we will need to categorically axiomatize *progress measures* for parity games ([37]; see also [38]). Possibly relevant to this direction is our recent coalgebraic modeling of Büchi and parity acceptance conditions [39].

Practical implications of the proposed framework (and concrete “ranking functions” derived thereby) shall be investigated, too. We are especially interested in cyber-physical applications in which state spaces are inherently infinite but often allow succinct symbolic presentations (e.g. by polynomials). The work closely related to this direction is [23].

A categorical account on martingales is also found in recent [40], where a connection between two classic results—Kolmogorov’s extension theorem and Doob’s martingale convergence theorem—is established in categorical terms. The relationship between this work and ours shall be pursued, possibly centered around the notion of final sequence.

ACKNOWLEDGMENT

Thanks are due to Eugenia Sironi and anonymous referees for their useful comments. The authors are supported by ER-ATO HASUO Metamathematics for Systems Design Project (No. JPMJER1603), JST, and Grants-in-Aid No. 15KT0012 & 15K11984, JSPS. Natsuki Urabe is supported by Grant-in-Aid for JSPS Fellows (No. 16J08157).

REFERENCES

- [1] V. Schuppan and A. Biere, “Liveness checking as safety checking for infinite state spaces,” *Electr. Notes Theor. Comput. Sci.*, vol. 149, no. 1, pp. 79–96, 2006. [Online]. Available: <http://dx.doi.org/10.1016/j.entcs.2005.11.018>
- [2] K. Claessen and N. Sörensson, “A liveness checking algorithm that counts,” in *Formal Methods in Computer-Aided Design, FMCAD 2012, Cambridge, UK, October 22-25, 2012*, G. Cabodi and S. Singh, Eds. IEEE, 2012, pp. 52–59. [Online]. Available: <http://ieeexplore.ieee.org/xpl/articleDetails.jsp?arnumber=6462555>
- [3] P. K. Nalla, R. K. Gajavelly, H. Mony, J. Baumgartner, and R. Kanzelman, “Effective liveness verification using a transformation-based framework,” in *2014 27th International Conference on VLSI Design and 2014 13th International Conference on Embedded Systems, Mumbai, India, January 5-9, 2014*. IEEE Computer Society, 2014, pp. 74–79. [Online]. Available: <http://dx.doi.org/10.1109/VLSID.2014.20>
- [4] R. W. Floyd, “Assigning meanings to programs,” *Proceedings of Symposium on Applied Mathematics*, vol. 19, pp. 19–32, 1967.
- [5] A. Chakarov and S. Sankaranarayanan, “Probabilistic program analysis with martingales,” in *Computer Aided Verification - 25th International Conference, CAV 2013, Saint Petersburg, Russia, July 13-19, 2013. Proceedings*, ser. Lecture Notes in Computer Science, N. Sharygina and H. Veith, Eds., vol. 8044. Springer, 2013, pp. 511–526. [Online]. Available: http://dx.doi.org/10.1007/978-3-642-39799-8_34
- [6] L. M. F. Fioriti and H. Hermanns, “Probabilistic termination: Soundness, completeness, and compositionality,” in *Proceedings of the 42nd Annual ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages, POPL 2015, Mumbai, India, January 15-17, 2015*, S. K. Rajamani and D. Walker, Eds. ACM, 2015, pp. 489–501. [Online]. Available: <http://doi.acm.org/10.1145/2676726.2677001>
- [7] B. Jacobs, *Introduction to Coalgebra: Towards Mathematics of States and Observation*, ser. Cambridge Tracts in Theoretical Computer Science. Cambridge University Press, 2016.
- [8] J. J. M. Rutten, “Universal coalgebra: a theory of systems,” *Theor. Comput. Sci.*, vol. 249, pp. 3–80, 2000.
- [9] A. Sokolova, “Probabilistic systems coalgebraically: A survey,” *Theor. Comput. Sci.*, vol. 412, no. 38, pp. 5095–5110, 2011. [Online]. Available: <http://dx.doi.org/10.1016/j.tcs.2011.05.008>
- [10] P. Aczel and N. P. Mendler, “A final coalgebra theorem,” in *Category Theory and Computer Science, Manchester, UK, September 5-8, 1989, Proceedings*, ser. Lecture Notes in Computer Science, D. H. Pitt, D. E. Rydeheard, P. Dybjer, A. M. Pitts, and A. Poigné, Eds., vol. 389. Springer, 1989, pp. 357–365.
- [11] I. Hasuo, “Generic forward and backward simulations,” in *CONCUR*, ser. Lecture Notes in Computer Science, C. Baier and H. Hermanns, Eds., vol. 4137. Springer, 2006, pp. 406–420.
- [12] N. Urabe and I. Hasuo, “Coalgebraic infinite traces and Kleisli simulations,” in *Algebra and Coalgebra in Computer Science - 6th International Conference, CALCO 2015, Nijmegen, Netherlands, June 24-26, 2015. Proceedings*, 2015.
- [13] V. Capretta, T. Uustalu, and V. Vene, “Corecursive algebras: A study of general structured corecursion,” in *Formal Methods: Foundations and Applications, 12th Brazilian Symposium on Formal Methods, SBMF 2009, Gramado, Brazil, August 19-21, 2009, Revised Selected Papers*, ser. Lecture Notes in Computer Science, M. V. M. Oliveira and J. Woodcock, Eds., vol. 5902. Springer, 2009, pp. 84–100. [Online]. Available: http://dx.doi.org/10.1007/978-3-642-10452-7_7
- [14] M. B. Smyth and G. D. Plotkin, “The category theoretic solution of recursive domain equations,” *SIAM Journ. Comput.*, vol. 11, pp. 761–783, 1982.
- [15] P. J. Freyd, “Algebraically complete categories,” in *Como Conference on Category Theory*, ser. Lect. Notes Math., A. Carboni, M. C. Pedicchio, and G. Rosolini, Eds., no. 1488. Springer, Berlin, 1991, pp. 95–104.
- [16] I. Hasuo, B. Jacobs, and A. Sokolova, “Generic trace semantics via coinduction,” *Logical Methods in Computer Science*, vol. 3, no. 4, 2007.
- [17] I. Hasuo, “Generic weakest precondition semantics from monads enriched with order,” *Theor. Comput. Sci.*, vol. 604, pp. 2–29, 2015. [Online]. Available: <http://dx.doi.org/10.1016/j.tcs.2015.03.047>
- [18] W. Hino, H. Kobayashi, I. Hasuo, and B. Jacobs, “Healthiness from duality,” in *Proceedings of the 31st Annual ACM/IEEE Symposium on Logic in Computer Science, LICS '16, New York, NY, USA, July 5-8, 2016*, M. Grohe, E. Koskinen, and N. Shankar, Eds. ACM, 2016, pp. 682–691. [Online]. Available: <http://doi.acm.org/10.1145/2933575.2935319>
- [19] L. Schröder and D. Pattinson, “PSPACE bounds for rank-1 modal logics,” *ACM Trans. Comput. Log.*, vol. 10, no. 2, 2009.
- [20] C. Urban and A. Miné, “An abstract domain to infer ordinal-valued ranking functions,” in *Programming Languages and Systems - 23rd European Symposium on Programming, ESOP 2014, Held as Part of the European Joint Conferences on Theory and Practice of Software, ETAPS 2014, Grenoble, France, April 5-13, 2014, Proceedings*, ser. Lecture Notes in Computer Science, Z. Shao, Ed., vol. 8410. Springer, 2014, pp. 412–431. [Online]. Available: http://dx.doi.org/10.1007/978-3-642-54833-8_22
- [21] Z. Manna and A. Pnueli, “Adequate proof principles for invariance and liveness properties of concurrent programs,” *Sci. Comput. Program.*, vol. 4, no. 3, pp. 257–289, 1984. [Online]. Available: [http://dx.doi.org/10.1016/0167-6423\(84\)90003-0](http://dx.doi.org/10.1016/0167-6423(84)90003-0)
- [22] E. Grädel and I. Walukiewicz, “Positional determinacy of games with infinitely many priorities,” *Logical Methods in Computer Science*, vol. 2, no. 4, 2006. [Online]. Available: [http://dx.doi.org/10.2168/LMCS-2\(4:6\)2006](http://dx.doi.org/10.2168/LMCS-2(4:6)2006)

- [23] A. N. Chakarov, “Deductive verification of infinite-state stochastic systems using martingales,” Ph.D. dissertation, University of Colorado, 2016.
- [24] S. Mac Lane, *Categories for the working mathematician*, 2nd ed. Springer-Verlag New York, 1998, vol. 5.
- [25] G. Osius, “Categorical set theory: A characterization of the category of sets,” *Journal of Pure and Applied Algebra*, vol. 4, no. 1, pp. 79 – 119, 1974. [Online]. Available: <http://www.sciencedirect.com/science/article/pii/0022404974900322>
- [26] P. Taylor, *Practical foundations of mathematics*, ser. Cambridge studies in advanced mathematics. Cambridge, New York (N. Y.), Melbourne: Cambridge University Press, 1999. [Online]. Available: <http://opac.inria.fr/record=b1095522>
- [27] J. Adámek, S. Milius, L. S. Moss, and L. Sousa, “Well-pointed coalgebras,” *Logical Methods in Computer Science*, vol. 9, no. 3, 2013. [Online]. Available: [http://dx.doi.org/10.2168/LMCS-9\(3:2\)2013](http://dx.doi.org/10.2168/LMCS-9(3:2)2013)
- [28] P. B. Levy, “Final coalgebras from corecursive algebras,” in *6th Conference on Algebra and Coalgebra in Computer Science, CALCO 2015, June 24-26, 2015, Nijmegen, The Netherlands*, ser. LIPIcs, L. S. Moss and P. Sobocinski, Eds., vol. 35. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2015, pp. 221–237. [Online]. Available: <http://dx.doi.org/10.4230/LIPIcs.CALCO.2015.221>
- [29] P. Cousot and R. Cousot, “Constructive versions of Tarski’s fixed point theorems,” *Pacific Journal of Mathematics*, vol. 82, no. 1, pp. 43–57, 1979.
- [30] G. Plotkin, “Notes on domains (pisa notes),” 1983, available at <http://www.dcs.ed.ac.uk/home/gdp/publications/>.
- [31] O. Bournez and F. Garnier, “Proving positive almost-sure termination,” in *Term Rewriting and Applications, 16th International Conference, RTA 2005, Nara, Japan, April 19-21, 2005, Proceedings*, ser. Lecture Notes in Computer Science, J. Giesl, Ed., vol. 3467. Springer, 2005, pp. 323–337. [Online]. Available: http://dx.doi.org/10.1007/978-3-540-32033-3_24
- [32] N. Saheb-Djahromi, “Cpo’s of measures for nondeterminism,” *Theor. Comput. Sci.*, vol. 12, pp. 19–37, 1980. [Online]. Available: [http://dx.doi.org/10.1016/0304-3975\(80\)90003-1](http://dx.doi.org/10.1016/0304-3975(80)90003-1)
- [33] J. Adámek, M. Haddadi, and S. Milius, “Corecursive algebras, corecursive monads and bloom monads,” *Logical Methods in Computer Science*, vol. 10, no. 3, 2014. [Online]. Available: [http://dx.doi.org/10.2168/LMCS-10\(3:19\)2014](http://dx.doi.org/10.2168/LMCS-10(3:19)2014)
- [34] V. Capretta, T. Uustalu, and V. Vene, “Recursive coalgebras from comonads,” *Inf. Comput.*, vol. 204, no. 4, pp. 437–468, 2006. [Online]. Available: <http://dx.doi.org/10.1016/j.ic.2005.08.005>
- [35] A. Abel and B. Pientka, “Well-founded recursion with copatterns and sized types,” *J. Funct. Program.*, vol. 26, p. e2, 2016. [Online]. Available: <http://dx.doi.org/10.1017/S0956796816000022>
- [36] L. S. Shapley, “Stochastic games,” *Proceedings of the National Academy of Sciences*, vol. 39, no. 10, pp. 1095–1100, 1953. [Online]. Available: <http://www.pnas.org/content/39/10/1095.short>
- [37] M. Jurdzinski, “Small progress measures for solving parity games,” in *STACS*, ser. Lecture Notes in Computer Science, H. Reichel and S. Tison, Eds., vol. 1770. Springer, 2000, pp. 290–301.
- [38] I. Hasuo, S. Shimizu, and C. Cirstea, “Lattice-theoretic progress measures and coalgebraic model checking,” in *Proceedings of the 43rd Annual ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages, POPL 2016, St. Petersburg, FL, USA, January 20 - 22, 2016*, R. Bodik and R. Majumdar, Eds. ACM, 2016, pp. 718–732. [Online]. Available: <http://doi.acm.org/10.1145/2837614.2837673>
- [39] N. Urabe, S. Shimizu, and I. Hasuo, “Coalgebraic trace semantics for Büchi and parity automata,” in *27th International Conference on Concurrency Theory, CONCUR 2016, August 23-26, 2016, Québec City, Canada*, ser. LIPIcs, J. Desharnais and R. Jagadeesan, Eds., vol. 59. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2016, pp. 24:1–24:15. [Online]. Available: <http://dx.doi.org/10.4230/LIPIcs.CONCUR.2016.24>
- [40] D. Kozen, “Kolmogorov extension, martingale convergence, and compositionality of processes,” in *Proc. 31st ACM-IEEE Symp. Logic in Computer Science (LICS 2016)*, N. Shankar, Ed. New York City: ACM, July 2016, pp. 692–699.

A. Formal Discussions for Two-Player Games

Definition A.1 (σ_g). Let $(\{0, 1\}, \leq)$ be a truth-value domain where \leq denotes the usual order. We define an F_g -modality $\sigma_g : F_g\{0, 1\} \rightarrow \{0, 1\}$ by

$$\sigma_g(\Gamma, t) = \begin{cases} 1 & (t = 1) \\ \max_{A \in \Gamma} \min_{a \in A} a & (\text{otherwise}). \end{cases}$$

Proposition A.2. We define an F_g -modality $\sigma_g : F_g\{0, 1\} \rightarrow \{0, 1\}$ as in Def. A.1.

- 1) The modality σ_g has least fixed points, and for each coalgebra $c : X \rightarrow F_g X$, the corresponding least fixed-point property $\llbracket \mu \sigma_g \rrbracket_c : X \rightarrow \{0, 1\}$ is given as follows:

$$\llbracket \mu \sigma_g \rrbracket_c(x) = \begin{cases} 1 & (x \in \text{Reach}_{G^c, \text{Acc}^c}) \\ 0 & (\text{otherwise}). \end{cases}$$

- 2) For a game structure $G = (X_{\max}, X_{\min}, \tau)$ and a set $\text{Acc} \subseteq X_{\max}$, we have:

$$\llbracket \mu \sigma_g \rrbracket_{c_{G, \text{Acc}}}(x) = \begin{cases} 1 & (x \in \text{Reach}_{G, \text{Acc}}) \\ 0 & (\text{otherwise}). \end{cases}$$

Proof. We first prove (1). We define $f : X \rightarrow \{0, 1\}$ by

$$f(x) = \begin{cases} 1 & (x \in \text{Reach}_{G^c, \text{Acc}^c}) \\ 0 & (\text{otherwise}). \end{cases}$$

By definition of the least fixed-point property, it suffices to show that f is the least fixed point of the function $\Phi_{c, \sigma_g} : \{0, 1\}^X \rightarrow \{0, 1\}^X$ in Def. III.2.

We first show that f is a fixed point of Φ_{c, σ_g} . For each $x \in X$, we have:

$$\begin{aligned} & \Phi_{c, \sigma_g}(f)(x) = 1 \\ & \Leftrightarrow (\sigma_g \circ F_g f \circ c)(x) = 1 && (\text{by def. of } \Phi_{c, \sigma_g}) \\ & \Leftrightarrow c_2(x) = 1, \text{ or } \exists A \in c_1(x). \forall x' \in A. f(x') = 1 \\ & && (\text{by def. of } \sigma_g \text{ and } F_g) \\ & \Leftrightarrow c_2(x) = 1, \text{ or} \\ & \quad \exists A \in c_1(x). \forall x' \in A. \\ & \quad \quad \exists \alpha' : \text{a strategy of max. } \forall \beta' : \text{a strategy of min.} \\ & \quad \quad \rho^{\alpha', \beta', x'} \text{ is winning for max} && (\text{by def. of } f) \\ & \Leftrightarrow c_2(x) = 1, \text{ or} \\ & \quad \exists A \in c_1(x). \\ & \quad \quad \exists (\alpha'_{x'})_{x' \in A} : \text{a family of strategies of max.} \\ & \quad \quad \forall x' \in A. \forall \beta' : \text{a strategy of min.} \\ & \quad \quad \rho^{\alpha'_{x'}, \beta', x'} \text{ is winning for max} \\ & \Leftrightarrow x \in \text{Acc}^c, \text{ or} \\ & \quad \exists y \in X_{\min}^c \text{ s.t. } (x, y) \in \tau^c. \\ & \quad \exists (\alpha'_{x'})_{x' \in X_{\max}^c \text{ s.t. } (y, x') \in \tau^c} : \text{a family of strategies of max.} \\ & \quad \quad \forall x' \in X_{\max}^c \text{ s.t. } (y, x') \in \tau^c. \forall \beta' : \text{a strategy of min.} \\ & \quad \quad \rho^{\alpha'_{x'}, \beta', x'} \text{ is winning for max} \end{aligned}$$

$\Leftrightarrow \exists \alpha : \text{a strategy of max. } \forall \beta : \text{a strategy of min.}$

$\rho^{\alpha, \beta, x}$ is winning for max (by def. of $\rho^{\alpha, \beta, x}$)

$\Leftrightarrow f(x) = 1$ (by def. of f).

Hence f is a fixed point of Φ_{c, σ_g} .

It remains to show that $f : X \rightarrow \{0, 1\}$ is the least fixed point with respect to the pointwise extension of \leq . Let $f' : X \rightarrow \{0, 1\}$ be a fixed point of Φ_{c, σ_g} . To prove $f \leq f'$, it suffices to prove that $f'(x) = 0$ implies $f(x) = 0$ for each $x \in X$.

For each $x' \in X$, we have:

$$\begin{aligned} & f'(x') = 0 \\ & \Leftrightarrow \Phi_{c, \sigma_g}(f')(x') = 0 && (f' \text{ is a fixed point of } \Phi_{c, \sigma_g}) \\ & \Leftrightarrow (\sigma_g \circ F_g f' \circ c)(x') = 0 && (\text{by def. of } \Phi_{c, \sigma_g}) \\ & \Leftrightarrow c_2(x') = 0, \text{ and } \forall A \in c_1(x'). \exists x'' \in A. f'(x'') = 0 \\ & && (\text{by def. of } \sigma_g \text{ and } F_g) \\ & \Leftrightarrow x' \notin \text{Acc}^c, \text{ and} \\ & \quad \forall y \in X_{\min}^c \text{ s.t. } (x', y) \in \tau^c. \\ & \quad \quad \exists x'' \in X_{\max}^c \text{ s.t. } (y, x'') \in \tau^c. f'(x'') = 0 \\ & && (\text{by def. of } G^c \text{ and } \text{Acc}^c). \end{aligned}$$

This means that if $f'(x') = 0$, then $c_2(x') = 0$ and for each $A \in c_1(x')$ there exists $x'' \in A$ such that $f'(x'') = 0$. Hence for each $x \in X$ such that $f'(x) = 0$, we can define a strategy β_x of the player min so that for each strategy α of the player max, the resulting run $\rho^{\alpha, \beta_x, x}$ from x is not winning for max.

Therefore by the definition of f , we have $f(x) = 0$. This concludes the proof.

The item (2) is proved in a similar way. \square

1) Details for Example III.9:

Proposition A.3. The modality $\sigma_g : F_g\{0, 1\} \rightarrow \{0, 1\}$ in Def. A.1 satisfies Asm. III.7.

Proof. It is easy to see that Cond. 1 is satisfied. By monotonicity of the functions max and min that are used in σ_g , Cond. 2 is satisfied. \square

2) Details of Example III.15:

Proposition A.4. We define an F_g -modality σ_g as in Def. A.1, and fix an ordinal \mathfrak{z} . We define an F_g -algebra $r_{g, \mathfrak{z}} : F_g \mathbf{Ord}_{\leq \mathfrak{z}} \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}$, a function $q_{g, \mathfrak{z}} : F_g \mathbf{Ord}_{\leq \mathfrak{z}} \rightarrow \{0, 1\}$ and a partial order $\sqsubseteq_{\mathbf{Ord}_{\leq \mathfrak{z}}}$ over $\mathbf{Ord}_{\leq \mathfrak{z}}$ as follows.

- 1) $\mathbf{Ord}_{\leq \mathfrak{z}} = \{\mathfrak{a} \mid \mathfrak{a} \text{ is an ordinal s.t. } \mathfrak{a} \leq \mathfrak{z}\}$, and $r_{g, \mathfrak{z}} : F_g \mathbf{Ord}_{\leq \mathfrak{z}} \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}$ is defined by

$$r_{g, \mathfrak{z}}(\Gamma, t) = \begin{cases} 0 & (t = 1) \\ \min_{A \in \Gamma} \sup_{a \in A} (\mathfrak{a} \hat{+} 1) & (\text{otherwise}). \end{cases}$$

Here $\mathfrak{a} \hat{+} 1$ denotes $\min\{\mathfrak{a} + 1, \mathfrak{z}\}$.

- 2) $q_{g, \mathfrak{z}} : \mathbf{Ord}_{\leq \mathfrak{z}} \rightarrow \{0, 1\}$ is defined by

$$q_{g, \mathfrak{z}}(\mathfrak{a}) = \begin{cases} 0 & (\mathfrak{a} = \mathfrak{z}) \\ 1 & (\text{otherwise}). \end{cases}$$

3) For $\mathbf{a}, \mathbf{b} \in \mathbf{Ord}_{\leq \mathfrak{z}}$, $\mathbf{a} \sqsubseteq_{\mathbf{Ord}} \mathbf{b} \stackrel{\text{def.}}{\Leftrightarrow} \mathbf{a} \geq \mathbf{b}$ (note the directions).

Then the triple $(r_{\mathfrak{g}, \mathfrak{z}}, q_{\mathfrak{g}, \mathfrak{z}}, \sqsubseteq_{\mathbf{Ord}})$ is a ranking domain.

The most difficult part of the proof is to prove that r is a corecursive algebra (Cond. 4 in Def. III.10). We prove it separately. To this end, we first prove the following sublemma.

Lemma A.5. *The algebra $r_{\mathfrak{g}, \mathfrak{z}} : F_{\mathfrak{g}} \mathbf{Ord}_{\leq \mathfrak{z}} \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}$ satisfies Cond. 2 in Def. III.10.*

Proof. It is easy to see that Cond. 2a is satisfied.

We prove that Cond. 2b is satisfied. Let $b_1, b_2 : X \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}$ and assume that $b_1 \sqsubseteq_{\mathbf{Ord}} b_2$. Let $x \in X$. As 0 is the greatest element in $(\mathbf{Ord}_{\leq \mathfrak{z}}, \sqsubseteq_{\mathbf{Ord}})$, if $\Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}(b_2)(x) = 0$ then we have $\Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}(b_1)(x) \sqsubseteq_{\mathbf{Ord}} \Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}(b_2)(x)$.

Assume that $\Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}(b_2)(x) = n \in \mathbf{Ord}_{\leq \mathfrak{z}} \setminus \{0\}$. Let $c(x) = (\Gamma, t) \in F_{\mathfrak{g}} X = \mathcal{P}^2 X \times \{0, 1\}$. Then by definition of $r_{\mathfrak{g}, \mathfrak{z}}$ and $F_{\mathfrak{g}}$, $t = 0$ and moreover for all $A \in \Gamma$ there exists $x \in A$ such that $b_2(x) \geq n$. As $b_1 \sqsubseteq_{\mathbf{Ord}} b_2$, $b_2(x) \geq n$ implies $b_1(x) \geq n$. Hence we have $r_{\mathfrak{g}, \mathfrak{z}} \circ F_{\mathfrak{g}} b_1(\Gamma, t) \geq n$, and this implies $\Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}(b_1)(x) \sqsubseteq_{\mathbf{Ord}} \Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}(b_2)(x)$. Therefore Cond. 2b is satisfied. \square

Lemma A.6. *The algebra $r_{\mathfrak{g}, \mathfrak{z}} : F_{\mathfrak{g}} \mathbf{Ord}_{\leq \mathfrak{z}} \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}$ is corecursive.*

Proof. Let $c : X \rightarrow F_{\mathfrak{g}} X$ be an $F_{\mathfrak{g}}$ -coalgebra. It suffices to show that $\Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}$ (Def. III.2) has a unique fixed point.

By Lem. A.5, the poset $(\mathbf{Ord}_{\leq \mathfrak{z}}^X, \sqsubseteq_{\mathbf{Ord}})$ (here $\sqsubseteq_{\mathbf{Ord}}$ denotes the pointwise extension of $\sqsubseteq_{\mathbf{Ord}}$ over $\mathbf{Ord}_{\leq \mathfrak{z}}$) is a complete lattice. Therefore we can construct a transfinite sequence

$$\begin{aligned} \perp_{\mathbf{Ord}_{\leq \mathfrak{z}}} \sqsubseteq_{\mathbf{Ord}} \Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}(\perp_{\mathbf{Ord}_{\leq \mathfrak{z}}}) \sqsubseteq_{\mathbf{Ord}} \cdots \\ \sqsubseteq_{\mathbf{Ord}} \Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}^{\alpha}(\perp_{\mathbf{Ord}_{\leq \mathfrak{z}}}) \sqsubseteq_{\mathbf{Ord}} \cdots \end{aligned}$$

as in Thm. II.18.2. By Thm. II.18.2, there exists an ordinal \mathbf{n} such that $\Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}^{\mathbf{n}}(\perp_{\mathbf{Ord}_{\leq \mathfrak{z}}})$ is the least fixed point of $\Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}$.

It remains to show that this is the unique fixed point. Let $f_1, f_2 : X \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}$ be fixed points of $\Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}$. We prove $f_1(x) = \mathbf{a} \Leftrightarrow f_2(x) = \mathbf{a}$ for each $x \in X$ and $\mathbf{a} < \mathfrak{z}$ by transfinite induction on \mathbf{a} .

For the base case, we have:

$$\begin{aligned} f_1(x) = 0 \\ \Leftrightarrow r_{\mathfrak{g}, \mathfrak{z}} \circ F_{\mathfrak{g}} f_1 \circ c(x) = 0 \quad (f_1 \text{ is a fixed point of } \Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}) \\ \Leftrightarrow c_2(x) = 1 \quad (\text{by def. of } r_{\mathfrak{g}, \mathfrak{z}} \text{ and } F_{\mathfrak{g}}) \\ \Leftrightarrow r_{\mathfrak{g}, \mathfrak{z}} \circ F_{\mathfrak{g}} f_2 \circ c(x) = 0 \quad (\text{by def. of } F_{\mathfrak{g}} \text{ and } r_{\mathfrak{g}, \mathfrak{z}}) \\ \Leftrightarrow f_2(x) = 0 \quad (f_2 \text{ is a fixed point of } \Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}). \end{aligned}$$

Let $0 < \mathbf{a} < \mathfrak{z}$ and assume $f(x) = \mathbf{a}'$ iff $g(x) = \mathbf{a}'$ for each $x \in X$ and $\mathbf{a}' < \mathbf{a}$. Note here that by $\mathbf{a} < \mathfrak{z}$, we have

$\mathbf{a} \hat{+} 1 = \mathbf{a} + 1$. Hence we have:

$$\begin{aligned} f_1(x) = \mathbf{a} \\ \Leftrightarrow r_{\mathfrak{g}, \mathfrak{z}} \circ F_{\mathfrak{g}} f_1 \circ c(x) = \mathbf{a} \quad (f_1 \text{ is a fixed point of } \Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}) \\ \Leftrightarrow c_2(x) = 0 \text{ and } \min_{A \in c_1(x)} \sup_{x' \in A} f_1(x') = \mathbf{a} - 1 \\ \quad (\text{by def. of } r_{\mathfrak{g}, \mathfrak{z}} \text{ and } F_{\mathfrak{g}}) \\ \Leftrightarrow c_2(x) = 0 \text{ and } \min_{A \in c_1(x)} \sup_{x' \in A} f_2(x') = \mathbf{a} - 1 \quad (\text{by IH}) \\ \Leftrightarrow \Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}(f_2)(x) = \mathbf{a} \quad (\text{by def. of } \Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}) \\ \Leftrightarrow f_2(x) = \mathbf{a} \quad (f_2 \text{ is a fixed point of } \Phi_{c, r_{\mathfrak{g}, \mathfrak{z}}}). \end{aligned}$$

Hence we have $f_1(x) = \mathbf{a}$ iff $f_2(x) = \mathbf{a}$ for each $x \in X$ and $\mathbf{a} < \mathfrak{z}$. This immediately implies that $f_1(x) = \mathfrak{z}$ iff $f_2(x) = \mathfrak{z}$ for each $x \in X$. \square

Proof (Prop. A.4). We prove that the conditions in Def. III.10 are satisfied.

Cond. 1: Let $(\Gamma, t) \in F_{\mathfrak{g}} \mathbf{Ord}_{\leq \mathfrak{z}} = \mathcal{P}^2 r_{\mathfrak{g}, \mathfrak{z}} \times \{0, 1\}$ and assume that $\sigma_{\mathfrak{g}} \circ F_{\mathfrak{g}} q_{\mathfrak{g}, \mathfrak{z}}(\Gamma, t) = 0$. Then by the definitions of $\sigma_{\mathfrak{g}}$, $q_{\mathfrak{g}, \mathfrak{z}}$ and $F_{\mathfrak{g}}$, we have $t = 0$ and $\forall A \in \Gamma. \mathfrak{z} \in A$. Hence by the definition of $r_{\mathfrak{g}, \mathfrak{z}}$, we have $r_{\mathfrak{g}, \mathfrak{z}}(\Gamma, t) = \mathfrak{z}$. Therefore by the definition of $q_{\mathfrak{g}, \mathfrak{z}}$, we have $q_{\mathfrak{g}, \mathfrak{z}} \circ r_{\mathfrak{g}, \mathfrak{z}}(\Gamma, t) = 0$. Hence Cond. 1 is satisfied.

Cond. 2: Already proved in Lem. A.5

Cond. 3: By the definition of $q_{\mathfrak{g}, \mathfrak{z}} : R \rightarrow \{0, 1\}$, for $a_1, a_2 \in \mathbf{Ord}_{\leq \mathfrak{z}}$, $a_1 \leq a_2$ implies $q_{\mathfrak{g}, \mathfrak{z}}(a_1) \geq q_{\mathfrak{g}, \mathfrak{z}}(a_2)$ (here each \leq denotes the usual order). Therefore $q_{\mathfrak{g}, \mathfrak{z}}$ is monotone.

By its definition, $q_{\mathfrak{g}, \mathfrak{z}}(\mathfrak{z}) = 0$ and hence $q_{\mathfrak{g}, \mathfrak{z}}$ is strict.

For a subset $K \subseteq \mathbf{Ord}_{\leq \mathfrak{z}}$, it is easy to see that $\bigwedge_{a \in K} (q_{\mathfrak{g}, \mathfrak{z}}(a) \leq q_{\mathfrak{g}, \mathfrak{z}}(\bigwedge_{a \in K} a))$ is satisfied. We prove the opposite direction. Assume that $q_{\mathfrak{g}, \mathfrak{z}}(\bigwedge_{a \in K} a) = 1$. Then by the definition of $q_{\mathfrak{g}, \mathfrak{z}}$, $\bigwedge_{a \in K} a = \mathbf{n}$ for some $\mathbf{n} < \mathfrak{z}$. By the definition of $\sqsubseteq_{\mathbf{Ord}}$, this implies $a = \mathbf{n}$ for some $a \in K$. Therefore we have $q_{\mathfrak{g}, \mathfrak{z}}(a) = 1$, and this implies $\bigwedge_{a \in K} (q_{\mathfrak{g}, \mathfrak{z}}(a) = 1)$. Hence $q_{\mathfrak{g}, \mathfrak{z}}(\bigwedge_{a \in K} a) \leq \bigwedge_{a \in K} (q_{\mathfrak{g}, \mathfrak{z}}(a))$ holds, and $q_{\mathfrak{g}, \mathfrak{z}}$ is continuous.

Cond. 4: Already proved in Lem. A.6.

Hence $(r_{\mathfrak{g}, \mathfrak{z}}, q_{\mathfrak{g}, \mathfrak{z}}, \sqsubseteq_{\mathbf{Ord}})$ is a ranking domain. \square

Proposition A.7. *Let $(r_{\mathfrak{g}, \mathfrak{z}}, q_{\mathfrak{g}, \mathfrak{z}}, \sqsubseteq_{\mathbf{Ord}})$ be the ranking domain in Prop. A.4. For each function $b : X \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}$, we have the followings.*

- 1) Let $c : X \rightarrow F_{\mathfrak{g}} X$ be an $F_{\mathfrak{g}}$ -coalgebra. A function $b : X \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}$ is a ranking arrow (Def. III.11) for c wrt. $(r_{\mathfrak{g}, \mathfrak{z}}, q_{\mathfrak{g}, \mathfrak{z}}, \sqsubseteq_{\mathbf{Ord}})$ if and only if b is a ranking function (Def. II.4) for G^c and Acc^c .
- 2) Let $G = (X_{\max}, x_{\min}, \tau)$ be a game structure and $\text{Acc} \subseteq X_{\max}$ be a set of accepting states. A function $b : X_{\max} \rightarrow \mathbf{Ord}_{\leq \mathfrak{z}}$ is a ranking function for G and Acc if and only if b is a ranking arrow for $c^{G, \text{Acc}}$ wrt. $(r_{\mathfrak{g}, \mathfrak{z}}, q_{\mathfrak{g}, \mathfrak{z}}, \sqsubseteq_{\mathbf{Ord}})$.
- 3) $b(x) < \mathfrak{z}$ iff $q_{\mathfrak{g}, \mathfrak{z}} \circ b(x) = 1$.

Proof.

1: For $x \in X$ such that $c(x) = (\Gamma, t)$ we write $c_1(x)$ and $c_2(x)$ for Γ and t respectively. We have:

$$\begin{aligned}
b : X &\rightarrow \mathbf{Ord}_{\leq 3} \text{ is a ranking arrow for } c \\
&\Leftrightarrow \forall x \in X. b(x) \sqsubseteq_{\mathbf{Ord}} r_{g,3} \circ F_g b \circ c(x) \quad (\text{by Def. III.11}) \\
&\Leftrightarrow \forall x \in X. r_{g,3} \circ F_g b \circ c(x) \leq b(x) \\
&\quad (\text{by the definition of } \sqsubseteq_{\mathbf{Ord}}) \\
&\Leftrightarrow \forall x \in X. (c_2(x) = 0 \Rightarrow \min_{A \in c_1(x)} \sup_{x' \in A} b(x') \hat{+} 1 \leq b(x)) \\
&\quad (\text{by the definitions of } r_{g,3} \text{ and } F_g) \\
&\Leftrightarrow \forall x \in X_{\max}^c. \\
&\quad (x \notin \text{Acc}^c \Rightarrow \min_{y:(x,y) \in \tau^c} \sup_{x':(y,x') \in \tau^c} b(x') \hat{+} 1 \leq b(x)) \\
&\quad (\text{by the definitions of } G^c \text{ and } \text{Acc}^c) \\
&\Leftrightarrow b : X \rightarrow \mathbf{Ord}_{\leq 3} \text{ is a ranking function for } G^c \text{ and } \text{Acc}^c \\
&\quad (\text{by Def. II.4}).
\end{aligned}$$

Hence Cond. 1 is satisfied.

2: This is proved in a similar manner to Cond. 1.

3: Immediate from the definition of $q_{g,3}$. \square

B. Proof of Lem. III.13

The basic idea of the proof is similar to the proof of Thm. II.18.2 in [29].

Proof. Let m be an ordinal such that $|L| < |m|$. As $|\{f^\alpha(l) \in L \mid \alpha \leq m\}| \leq |L|$, there exist ordinals α, α' such that $f^\alpha(l) = f^{\alpha'}(l)$. Without loss of generality, we assume $\alpha < \alpha'$.

By monotonicity of f and that $l \sqsubseteq f(l)$, we have

$$f^\alpha(l) = f^{\alpha+1}(l) = \dots = f^{\alpha'}(l).$$

This concludes the proof. \square

C. Proof of Prop. III.18

Proof. By $\langle c \rangle_r = r \circ F \langle c \rangle_r \circ c$ and $q \circ r = \sigma \circ Fq$, we have:

$$q \circ \langle c \rangle_r = \Phi_{c,\sigma}(q \circ \langle c \rangle_r).$$

This means that $q \circ \langle c \rangle_r$ is a fixed point of $\Phi_{c,\sigma}$. As $\llbracket \mu\sigma \rrbracket_c$ is the least fixed point of $\Phi_{c,\sigma}$ (Def. III.5), we have

$$q \circ \langle c \rangle_r \sqsupseteq \llbracket \mu\sigma \rrbracket_c.$$

Together with Thm. III.12, we have $q \circ \langle c \rangle_r = \llbracket \mu\sigma \rrbracket_c$. \square

D. Proof of Prop. IV.2

Proof. Recall that $\text{Reach}_{\text{Acc}^c}$ is defined as the function $f : X \rightarrow [0, 1]$ in Def. II.9. We first show that this f is a fixed point of Φ_{c,σ_p} . For $x \in X$, we have:

$$\begin{aligned}
&\Phi_{c,\sigma_p}(f)(x) \\
&= (\sigma_p \circ F_p f \circ c)(x) \quad (\text{by def. of } \Phi_{c,\sigma_p}) \\
&= \begin{cases} 1 & (c_2(x) = 1) \\ \sum_{x' \in \text{supp}(c_1(x))} c_1(x)(x') \cdot f(x') & (c_2(x) = 0) \end{cases} \\
&\quad (\text{by def. of } \sigma_p \text{ and } F_p) \\
&= \begin{cases} 1 & (c_2(x) = 1) \\ \sum_{x' \in \text{supp}(c_1(x))} c_1(x)(x') \cdot \lim_{n \rightarrow \infty} f_n(x') & (c_2(x) = 0) \end{cases} \\
&\quad (\text{by def. of } f) \\
&= \begin{cases} 1 & (c_2(x) = 1) \\ \lim_{n \rightarrow \infty} \sum_{x' \in \text{supp}(c_1(x))} c_1(x)(x') \cdot f_n(x') & (c_2(x) = 0) \end{cases} \\
&= \begin{cases} 1 & (c_2(x) = 1) \\ \lim_{n \rightarrow \infty} f_{n+1}(x) & (c_2(x) = 0) \end{cases} \quad (\text{by def. of } f_n) \\
&= \lim_{n \rightarrow \infty} f_n(x) \quad (\text{by def. of } f_n) \\
&= f(x) \quad (\text{by def. of } f).
\end{aligned}$$

Hence f is a fixed point of Φ_{c,σ_p} .

It remains to show that f is the least fixed point. Let $f' : X \rightarrow [0, 1]$ be a fixed point of Φ_{c,σ_p} .

We prove $f(x) \leq f'(x)$ for each $x \in X$. To this end, by the definition of f , it suffices to prove $f_n(x) \leq f'(x)$ for each $x \in X$ and $n \in \mathbb{N}$. We prove this by induction on n .

For $n = 0$, it is immediate from that $f_0(x) = 0$.

For $n > 0$, we have:

$$\begin{aligned}
&f_n(x) \\
&= \begin{cases} 1 & (c_2(x) = 1) \\ \sum_{x' \in \text{supp}(c_1(x))} c_1(x)(x') \cdot f_{n-1}(x') & (c_2(x) = 0) \end{cases} \\
&\quad (\text{by def. of } f_n) \\
&\leq \begin{cases} 1 & (c_2(x) = 1) \\ \sum_{x' \in \text{supp}(c_1(x))} c_1(x)(x') \cdot f'(x') & (c_2(x) = 0) \end{cases} \quad (\text{by IH}) \\
&= (\sigma_p \circ F_p f' \circ c)(x) \quad (\text{by def. of } F_p \text{ and } \sigma_p) \\
&= f'(x) \quad (f' \text{ is a fixed point}).
\end{aligned}$$

Hence we have $f_n(x) \leq f'(x)$ for each $x \in X$ and $n \in \mathbb{N}$, and this implies $f(x) \leq f'(x)$ for each $x \in X$. Therefore f is the least fixed point of Φ_{c,σ_p} . \square

E. Proof of Prop. IV.3

Proof.

1: For each $b' : X \rightarrow [0, \infty]$, we have:

b' is an ε -additive ranking supermartingale

$\Leftrightarrow (\forall x \in X.$

$$c_2(x) = 0 \\ \Rightarrow \sum_{x' \in \text{supp}(c_1(x))} c_1(x)(x') \cdot b'(x') + \varepsilon \leq b'(x)$$

(by Def. II.11)

$\Leftrightarrow \forall x \in X. r'_{p,\varepsilon} \circ F_p b' \circ c(x) \leq b'(x)$ (by def. of $r'_{p,\varepsilon}$)

$\Leftrightarrow b'$ is a (categorical) ranking arrow wrt. $(r'_{p,\varepsilon}, q'_p, \sqsubseteq_{[0,\infty]})$ (by Def. III.11).

Hence Cond. 1 holds.

2: Immediate from the definition of q'_p . \square

F. Proof of Prop. IV.4

Proof.

Cond. 1: Let $(\psi, t) \in F_p[0, \infty] = \mathcal{D}[0, \infty] \times \{0, 1\}$. Then we have:

$$\begin{aligned} & q'_p \circ r'_{p,\varepsilon}(\psi, t) \\ &= \begin{cases} q'_p(0) & (t = 1) \\ q'_p(\sum_{a \in \text{supp}(\psi)} a \cdot \psi(a) + \varepsilon) & (t = 0) \end{cases} \\ & \hspace{15em} \text{(by def. of } r'_{p,\varepsilon}\text{)} \\ &= \begin{cases} 1 & (t = 1 \text{ or } \sum_{a \in \text{supp}(\psi)} a \cdot \psi(a) + \varepsilon < \infty) \\ 0 & \text{(otherwise)} \end{cases} \\ & \hspace{15em} \text{(by def. of } q'_p\text{)} \\ &\leq \begin{cases} 1 & (t = 1 \text{ or } \psi([0, \infty)) = 1) \\ 0 & \text{(otherwise)} \end{cases} \\ &\leq \begin{cases} 1 & (t = 1) \\ \psi([0, \infty)) & \text{(otherwise)} \end{cases} \\ &= \begin{cases} 1 & (t = 1) \\ 1 \cdot \mathcal{D}q_p(\psi)(1) & \text{(otherwise)} \end{cases} \hspace{2em} \text{(by def. of } q'_p\text{)} \\ &= \sigma_p \circ F_p q'_p(\psi, t) \hspace{10em} \text{(by def. of } \sigma_p\text{)}. \end{aligned}$$

Hence we have $q'_p \circ r'_{p,\varepsilon} \leq \sigma_p \circ F_p q'_p$.

Cond. 2: It is easy to see that Cond. 2a is satisfied.

We prove that Cond. 2b is satisfied. Let $b_1, b_2 : X \rightarrow [0, \infty]$ and assume that $b_1 \sqsubseteq_{[0,\infty]} b_2$. For each $x \in X$ such that $c(x) = (\varphi, t) \in F_p X = \mathcal{D}X \times \{0, 1\}$, we have:

$$\begin{aligned} & \Phi_{c,r'_{p,\varepsilon}}(b_1)(x) \\ &= r'_{p,\varepsilon} \circ F_p b_1(\varphi, t) \hspace{5em} \text{(by def. of } \Phi_{c,r'_{p,\varepsilon}}\text{)} \\ &= \begin{cases} 1 & (t = 1) \\ \sum_{x \in \text{supp}(\varphi)} \varphi(x) \cdot b_1(x) + \varepsilon & (t = 0) \end{cases} \\ & \hspace{15em} \text{(by def. of } r'_{p,\varepsilon}\text{)} \\ &\leq \begin{cases} 1 & (t = 1) \\ \sum_{x \in \text{supp}(\varphi)} \varphi(x) \cdot b_2(x) + \varepsilon & (t = 1) \end{cases} \end{aligned}$$

$$\begin{aligned} &= r'_{p,\varepsilon} \circ F_p b_2(\varphi, t) \hspace{10em} \text{(by def. of } r'_{p,\varepsilon}\text{)} \\ &= \Phi_{c,r'_{p,\varepsilon}}(b_2)(x) \hspace{10em} \text{(by def. of } \Phi_{c,r'_{p,\varepsilon}}\text{)}. \end{aligned}$$

Therefore Cond. 2b is satisfied.

Cond. 3: By the definition of q'_p , for each $a_1, a_2 \in [0, \infty]$, $a_1 \geq a_2$ implies $q'_p(a_1) \leq q'_p(a_2)$ (here each \leq denotes the standard order). Hence q'_p is monotone.

By the definition, we have $q'_p(\infty) = 0$. Hence q'_p is strict.

We prove that q'_p is continuous. Namely, for each subset $K \subseteq [0, \infty]$, we prove

$$\bigsqcup_{a \in K} (q'_p(a)) = q'_p\left(\bigsqcup_{a \in K} a\right).$$

It is easy to prove (LHS) \leq (RHS). We prove the opposite direction. To this end, by the definition of q'_p , it suffices to prove that if the right-hand side is 1 then the left-hand side is also 1. We can prove it as follows (here \bigsqcup denotes the supremum with respect to $\sqsubseteq_{[0,\infty]}$, and \bigwedge denotes the infimum with respect to the ordinary order over $[0, 1]$).

$$\begin{aligned} q'_p\left(\bigsqcup_{a \in K} a\right) = 1 &\Rightarrow \bigsqcup_{a \in K} a < \infty \hspace{5em} \text{(by def. of } q'_p\text{)} \\ &\Rightarrow \bigwedge_{a \in K} a < \infty \hspace{5em} \text{(by def. of } \sqsubseteq_{[0,\infty]}\text{)} \\ &\Rightarrow \exists a \in K. a < \infty \\ &\Rightarrow \exists a \in K. q'_p(a) = 1 \hspace{5em} \text{(by def. of } q'_p\text{)} \\ &\Rightarrow \bigsqcup_{a \in K} (q'_p(a)) = 1 \end{aligned}$$

Hence q'_p is continuous.

This concludes the proof. \square

G. Multiplicative Ranking Supermartingale, Categorically

Definition A.8. Let $M = (X, \tau)$ be a PTS. Let $\alpha \in (0, 1)$. A function $b : X \rightarrow [0, \infty]$ is an α -multiplicative ranking supermartingale if we have

$$\sum_{x' \in \text{supp}(\tau(x))} \tau(x)(x') \cdot b(x') \leq \alpha \cdot b(x)$$

for each $x \in X \setminus \text{Acc}$, and moreover there exists $\delta > 0$ such that $b(x) \geq \delta$ for each $x \in X \setminus \text{Acc}$.

The following is an attempt to define a ranking domain for α -multiplicative supermartingales. Let us fix real numbers $\alpha \in (0, 1)$ and $\delta > 0$, and define an F_p -algebra $r'_{\text{pm},\alpha} : F_p[\alpha\delta, \infty] \rightarrow [\alpha\delta, \infty]$, an arrow $q'_p : [0, \infty] \rightarrow [0, 1]$ and a partial order $\sqsubseteq_{[\alpha\delta,\infty]}$ over $[\alpha\delta, \infty]$ as follows.

1) For each $(\psi, t) \in F_p[\alpha\delta, \infty] = \mathcal{D}[\alpha\delta, \infty] \times \{0, 1\}$,

$$r'_{\text{pm},\alpha}(\psi, t) = \begin{cases} \alpha\delta & (t = 1) \\ \frac{1}{\alpha} \cdot \left(\sum_{a \in \text{supp}(\psi)} a \cdot \psi(a)\right) & \text{(otherwise)}. \end{cases}$$

2) $q'_p(\infty) = 0$ and $q'_p(a) = 1$ if $a < \infty$.

3) $a \sqsubseteq_{[\alpha\delta,\infty]} b \stackrel{\text{def.}}{\Leftrightarrow} a \geq b$ (note the direction).

The following proposition is analogous to Prop. IV.3.

Proposition A.9. *In this setting, for each $c : X \rightarrow F_p X$, we have the following.*

- a) *Let $b' : X \rightarrow [\alpha\delta, \infty]$ and assume $b' \sqsubseteq_{[\alpha\delta, \infty]} r'_{\text{pm}, \alpha} \circ F_p b' \circ c$. If we define $b : X \rightarrow [0, \infty]$ by $b(x) = b'(x)$, then b is an α -multiplicative ranking supermartingale.*
- b) *Let $b : X \rightarrow [0, \infty]$ be an α -multiplicative ranking supermartingale such that $b(x) \geq \delta$ if $x \in X \setminus \text{Acc}$. If we define $b' : X \rightarrow [\alpha\delta, \infty]$ by $b'(x) = \max\{\alpha\delta, b(x)\}$, then b' satisfies $b' \sqsubseteq_{[\alpha\delta, \infty]} r'_{\text{pm}, \alpha} \circ F_p b' \circ c$.*
- c) *For each $b' : X \rightarrow [0, \infty]$, we have $b'(x) \neq \infty$ iff $q'_p \circ b'(x) = 1$.*

Proof.

a): Let $x \in X \setminus \text{Acc}$. Then we have:

$$\begin{aligned} & \sum_{x' \in \text{supp}(\tau(x))} \tau(x)(x') \cdot b(x') \\ &= \sum_{x' \in \text{supp}(c_1(x))} c_1(x)(x') \cdot b'(x') \\ &= \alpha \cdot (r'_{\text{pm}, \alpha} \circ F_p b' \circ c)(x) \\ &\leq \alpha \cdot b'(x) \\ &= \alpha \cdot b(x). \end{aligned}$$

By the inequality above, we also have

$$b(x) \geq \frac{1}{\alpha} \sum_{x' \in \text{supp}(c_1(x))} c_1(x)(x') \cdot b'(x').$$

As $b'(x') \geq \alpha\delta$ holds for each $x' \in X$, we have $b(x) \geq \delta$. Therefore b is an α -multiplicative ranking supermartingale.

b): For each $x \in X$, we have:

$$\begin{aligned} & (r'_{\text{pm}, \alpha} \circ F_p b' \circ c)(x) \\ &= \begin{cases} \alpha\delta & (c_2(x) = 1) \\ \frac{1}{\alpha} \sum_{x' \in \text{supp}(c_1(x))} c_1(x)(x') \cdot b'(x') & (c_2(x) = 0) \end{cases} \\ &= \begin{cases} \alpha\delta & (x \in \text{Acc}) \\ \frac{1}{\alpha} \sum_{x' \in \text{supp}(\tau(x))} \tau(x)(x') \cdot b'(x') & (x \in X \setminus \text{Acc}) \end{cases} \\ &\leq \begin{cases} \alpha\delta & (x \in \text{Acc}) \\ \frac{1}{\alpha} \cdot \alpha b(x) & (x \in X \setminus \text{Acc}) \end{cases} \\ &\leq b'(x). \end{aligned}$$

Therefore by the definition of $\sqsubseteq_{[\alpha\delta, \infty]}$, we have $b' \sqsubseteq_{[\alpha\delta, \infty]} r'_{\text{pm}, \alpha} \circ F_p b' \circ c$.

c): Immediate from the definition of q'_p . \square

Therefore the triple $(r'_{\text{pm}, \alpha}, q'_p, \sqsubseteq_{[\alpha\delta, \infty]})$ is suited for accommodating α -multiplicative supermartingales in our categorical framework. However it is not a ranking domain. The following proposition is analogous to Prop. IV.4.

Proposition A.10. *The triple $(r'_{\text{pm}, \alpha}, q'_p, \sqsubseteq_{[0, \infty]})$ introduced above satisfies the conditions of a ranking domain (Def. III.10), except for Cond. 4.*

Proof.

Cond. 1: Let $(\psi, t) \in F_p[\alpha\delta, \infty] = \mathcal{D}[\alpha\delta, \infty] \times \{0, 1\}$. Then we have:

$$\begin{aligned} & q'_p \circ r'_{\text{pm}, \alpha}(\psi, t) \\ &= \begin{cases} q'_p(\alpha\delta) & (t = 1) \\ q'_p\left(\frac{1}{\alpha} \cdot \sum_{a \in \text{supp}(\psi)} a \cdot \psi(a)\right) & (t = 0) \end{cases} \\ & \hspace{15em} \text{(by def. of } r'_{\text{pm}, \alpha}\text{)} \\ &= \begin{cases} 1 & (t = 1 \text{ or } \frac{1}{\alpha} \cdot \sum_{a \in \text{supp}(\psi)} a \cdot \psi(a) < \infty) \\ 0 & \text{(otherwise)} \end{cases} \\ & \hspace{15em} \text{(by def. of } q'_p\text{)} \\ &\leq \begin{cases} 1 & (t = 1 \text{ or } \psi([0, \infty)) = 1) \\ 0 & \text{(otherwise)} \end{cases} \\ &\leq \begin{cases} 1 & (t = 1) \\ \psi([0, \infty)) & \text{(otherwise)} \end{cases} \\ &= \begin{cases} 1 & (t = 1) \\ 1 \cdot \mathcal{D}q_p(\psi)(1) & \text{(otherwise)} \end{cases} \hspace{2em} \text{(by def. of } q'_p\text{)} \\ &= \sigma_p \circ F_p q'_p(\psi, t) \hspace{5em} \text{(by def. of } \sigma_p\text{)}. \end{aligned}$$

Hence we have $q'_p \circ r'_{\text{pm}, \alpha} \leq \sigma_p \circ F_p q'_p$.

Cond. 2: It is easy to see that Cond. 2a is satisfied.

We prove that Cond. 2b is satisfied. Let $b_1, b_2 : X \rightarrow [\alpha\delta, \infty]$ and assume that $b_1 \sqsubseteq_{[\alpha\delta, \infty]} b_2$. For each $x \in X$ such that $c(x) = (\varphi, t) \in F_p X = \mathcal{D}X \times \{0, 1\}$, we have:

$$\begin{aligned} & \Phi_{c, r'_{\text{pm}, \alpha}}(b_1)(x) \\ &= r'_{\text{pm}, \alpha} \circ F_p b_1(\varphi, t) \hspace{5em} \text{(by def. of } \Phi_{c, r'_{\text{pm}, \alpha}}\text{)} \\ &= \begin{cases} 1 & (t = 1) \\ \frac{1}{\alpha} \cdot \sum_{x \in \text{supp}(\varphi)} \varphi(x) \cdot b_1(x) & (t = 0) \end{cases} \\ & \hspace{15em} \text{(by def. of } r'_{\text{pm}, \alpha}\text{)} \\ &\sqsubseteq_{[\alpha\delta, \infty]} \begin{cases} 1 & (t = 1) \\ \frac{1}{\alpha} \cdot \sum_{x \in \text{supp}(\varphi)} \varphi(x) \cdot b_2(x) & (t = 0) \end{cases} \\ &= r'_{\text{pm}, \alpha} \circ F_p b_2(\varphi, t) \hspace{5em} \text{(by def. of } r'_{\text{pm}, \alpha}\text{)} \\ &= \Phi_{c, r'_{\text{pm}, \alpha}}(b_2)(x) \hspace{5em} \text{(by def. of } \Phi_{c, r'_{\text{pm}, \alpha}}\text{)}. \end{aligned}$$

Therefore Cond. 2b is satisfied.

Cond. 3: It is proved in a similar manner to the proof of Prop. IV.4. \square

Example A.11. We define a coalgebra $c : X \rightarrow F_p X$ as in Example IV.5. We fix $\alpha \in (0, 1)$ and $\delta > 0$. We assume that $\alpha > 1/2$. For this coalgebra, we define arrows $b_1, b_2 : X \rightarrow [\alpha\delta, \infty]$ by: $b_1(x_0) = \frac{1}{2} \frac{1}{\alpha} \left(\frac{\alpha\delta}{2\alpha-1} + \delta \right)$, $b_1(x_1) = \frac{\alpha\delta}{2\alpha-1}$, $b_1(x_2) = \delta$ and $b_1(x_3) = \alpha\delta$; and $b_2(x_0) = \infty$, $b_2(x_1) = \infty$, $b_2(x_2) = \delta$ and $b_2(x_3) = \alpha\delta$. Both of these qualify as coalgebra-algebra homomorphisms from c to $r'_{\text{pm}, \alpha}$. Therefore $r'_{\text{pm}, \alpha}$ is not corecursive.

H. Proof of Prop. IV.7

We prove Prop. IV.7 in a similar manner to the proof of Prop. A.4: we first prove that r_p is a corecursive algebra separately. To this end, we first prove some lemmas.

Lemma A.12. *The order $\sqsubseteq_{\mathcal{DN}_\infty}$ in Prop. IV.7 is a partial order and \mathcal{DN}_∞ is a complete lattice with respect to this order.*

Sublemma A.13. *For every nondecreasing function $G : \mathbb{N} \rightarrow [0, 1]$, there exists a unique distribution φ over \mathbb{N}_∞ such that $\varphi([0, a]) = G(a)$ for each $a \in \mathbb{N}$.*

Proof. We define a distribution φ over \mathbb{N}_∞ by

$$\varphi(a) = \begin{cases} G(a) & (a = 0) \\ G(a) - G(a-1) & (0 < a < \infty) \\ 1 - \lim_{a' \rightarrow \infty} G(a') & (a = \infty). \end{cases}$$

As G is nondecreasing, $\varphi(a) \geq 0$ for each a . By its definition, we have $\sum_{a \in \mathbb{N}_\infty} \varphi(a) = 1$. Hence φ is a distribution.

Let φ' be a distribution such that $\varphi'([0, a]) = G(a)$. Then we have $\varphi(0) = G(0) = \varphi'([0, 0]) = \varphi'(0)$. Moreover for each $a \in \mathbb{N} \setminus \{0\}$, we have:

$$\varphi(a) = G(a) - G(a-1) = \varphi'([0, a]) - \varphi'([0, a-1]) = \varphi'(a).$$

Therefore we have $\varphi(a) = \varphi'(a)$ for each $a \in \mathbb{N}$, and this implies $\varphi(\infty) = \varphi'(\infty)$. Hence uniqueness is proved, and this concludes the proof. \square

Proof (Lem. A.12). We first prove that $\sqsubseteq_{\mathcal{DN}_\infty}$ is a partial order. Reflexivity and transitivity are immediate from those of the standard order \leq over $[0, 1]$. Assume that $\varphi \sqsubseteq_{\mathcal{DN}_\infty} \varphi'$ and $\varphi' \sqsubseteq_{\mathcal{DN}_\infty} \varphi$. By the definition of $\sqsubseteq_{\mathcal{DN}_\infty}$, we have $\varphi([0, a]) = \varphi'([0, a])$ for each $a \in [0, \infty]$. Then by Sublem. A.13, we have $\varphi = \varphi'$. Hence antisymmetry is also satisfied.

We prove that each subset $K \subseteq \mathcal{DN}_\infty$ has the least upper bound. We define $G : \mathbb{N} \rightarrow [0, 1]$ by $G(a) = \sup_{\varphi \in K} \varphi([0, a])$. Note that for each $\varphi \in K$, $a \leq b$ implies $\varphi([0, a]) \leq \varphi([0, b])$. Hence by the monotonicity of supremums, G is nondecreasing. Therefore by Sublem. A.13, there exists a unique distribution $\varphi^K \in \mathcal{DN}_\infty$ such that $\varphi^K([0, a]) = G(a)$ for each $a \in [0, \infty]$.

We prove that φ^K is the least upper bound of K .

Let $\varphi \in K$. For each $a \in \mathbb{N}$, we have:

$$\varphi([0, a]) \leq \sup_{\varphi' \in K} \varphi'([0, a]) = \varphi^K([0, a]).$$

Hence by the definition of $\sqsubseteq_{\mathcal{DN}_\infty}$, φ is an upper bound of K .

Let φ' be an upper bound of K . Then by the definition of $\sqsubseteq_{\mathcal{DN}_\infty}$, we have $\varphi([0, a]) \leq \varphi'([0, a])$ for each $\varphi \in K$ and $a \in \mathbb{N}$. Therefore we have $\varphi^K([0, a]) \leq \varphi'([0, a])$, and this means $\varphi^K \sqsubseteq_{\mathcal{DN}_\infty} \varphi'$ by the definition of $\sqsubseteq_{\mathcal{DN}_\infty}$. Hence φ^K is the least upper bound of K .

In a similar manner, we can prove that each $K \subseteq \mathcal{DN}_\infty$ has the greatest lower bound. \square

Lemma A.14. *The algebra $r_p : F_p \mathcal{DN}_\infty \rightarrow \mathcal{DN}_\infty$ in Prop IV.7 satisfies Cond. 2 in Def. III.10.*

Proof. It is already proved in Lem. A.12 that Cond. 2a is satisfied.

We prove that Cond. 2b is satisfied. Let $f_1, f_2 : X \rightarrow \mathcal{DN}_\infty$ and assume $f_1 \sqsubseteq_{\mathcal{DN}_\infty} f_2$. Let $x \in X$ and assume that $c(x) = (\varphi, t) \in F_p X$.

If $t = 1$ then we have:

$$r_p \circ F_p f_1(f)(\varphi, t) = r_p \circ F_p f_2(f)(\varphi, t) = \delta_0.$$

Therefore we have $\Phi_{c, \sigma_p}(f_1)(x) = \Phi_{c, \sigma_p}(f_2)(x)$.

Let $t = 0$ and $a \in \mathbb{N}$. If $a = 0$, by the definition of r_p , we have:

$$r_p \circ F_p f_1(f)(\varphi, t)([0, a]) = r_p \circ F_p f_2(f)(\varphi, t)([0, a]) = 0.$$

If $a > 0$, we have:

$$\begin{aligned} & r_p \circ F_p f_1(f)(\varphi, t)([0, a]) \\ &= \sum_{x \in \text{supp}(\varphi)} (\varphi(x) \cdot f_1(x)([0, a-1])) \quad (\text{by def. of } r_p) \\ &\leq \sum_{x \in \text{supp}(\varphi)} (\varphi(x) \cdot f_2(x)([0, a-1])) \quad (\text{by } f_1 \sqsubseteq_{\mathcal{DN}_\infty} f_2) \\ &= r_p \circ F_p f_2(\varphi, t)([0, a]) \quad (\text{by def. of } r_p). \end{aligned}$$

Therefore by the definition of $\sqsubseteq_{\mathcal{DN}_\infty}$, we have $r_p \circ F_p f_1(\varphi, t) \sqsubseteq_{\mathcal{DN}_\infty} r_p \circ F_p f_2(\varphi, t)$. This means that Φ_{c, r_p} is monotone. This concludes the proof. \square

Lemma A.15. *The F_p -algebra $r_p : F_p \mathcal{DN}_\infty \rightarrow \mathcal{DN}_\infty$ in Prop. IV.7 is corecursive.*

Proof. Let $c : X \rightarrow F_p X$ be an F_p -coalgebra. It suffices to show that the function $\Phi_{c, r_p} : (\mathcal{DN}_\infty)^X \rightarrow (\mathcal{DN}_\infty)^X$ (Def. III.2) has a unique fixed point.

By Lem. A.14, the poset $((\mathcal{DN}_\infty)^X, \sqsubseteq_{\mathcal{DN}_\infty})$ (here $\sqsubseteq_{\mathcal{DN}_\infty}$ denotes the pointwise extension) is a complete lattice. Therefore we can construct a transfinite sequence

$$\begin{aligned} \perp_{\mathcal{DN}_\infty} \sqsubseteq_{\mathcal{DN}_\infty} \Phi_{c, r_p}(\perp_{\mathcal{DN}_\infty}) \sqsubseteq_{\mathcal{DN}_\infty} \cdots \\ \sqsubseteq_{\mathcal{DN}_\infty} \Phi_{c, r_p}^n(\perp_{\mathcal{DN}_\infty}) \sqsubseteq_{\mathcal{DN}_\infty} \cdots \end{aligned}$$

as in Thm. II.18.2. By Thm. II.18.2, there exists an ordinal n such that $\Phi_{c, r_p}^n(\perp_{\mathcal{DN}_\infty})$ is the least fixed point of Φ_{c, r_p} .

It remains to show that this is a unique fixed point. Let $f_1, f_2 : X \rightarrow \mathcal{DN}_\infty$ be fixed points of Φ_{c, r_p} . We prove

$$f_1(x)(a) = f_2(x)(a) \quad (10)$$

for each $x \in X$ and $a \in \mathbb{N}$. For each $x \in X$ such that $(\varphi, t) = c(x)$, we write $c_1(x)$ and $c_2(x)$ for $\varphi \in \mathcal{DX}$ and $t \in \{0, 1\}$ respectively.

We first prove (10) for each $x \in X$ such that $c_2(x) = 1$. For such x , we have:

$$\begin{aligned} f_1(x)(a) &= \Phi_{c, r_p}(f_1)(x) \quad (f_1 \text{ is a fixed point}) \\ &= r_p \circ F_p f_1 \circ c(x) \\ &= \delta_0 \\ &= r_p \circ F_p f_2 \circ c(x) \\ &= \Phi_{c, r_p}(f_2)(x) \quad (f_2 \text{ is a fixed point}) \\ &= g(x). \end{aligned}$$

Hence we have $f_1(x)(a) = f_2(x)(a)$ for each a .

It remains to prove (10) for each $x \in X$ such that $c_2(x) = 0$. We prove it by the induction on a .

If $a = 0$ then by the definition of r_p , we have $f_1(x)(a) = f_2(x)(a) = 0$.

Let $a > 0$ and assume $f_1(x')(a') = f_2(x')(a')$ for each $x' \in X$ and $a' < a$. Then we have:

$$\begin{aligned}
f_1(x)(a) &= r_p \circ F_p f_1 \circ c(x)(a) && (f_1 \text{ is a fixed point}) \\
&= \sum_{x' \in \text{supp}(c_1(x))} c_1(x)(x') \cdot f_1(x')(a-1) && (\text{by def. of } r_p) \\
&= \sum_{x' \in \text{supp}(c_1(x))} c_1(x)(x') \cdot f_2(x')(a-1) && (\text{by IH}) \\
&= r_p \circ F_p f \circ c(x)(a) && (\text{by def. of } r_p) \\
&= f_2(x)(a) && (f_2 \text{ is a fixed point}).
\end{aligned}$$

Therefore we have $f_1(x)(a) = f_2(x)(a)$ for each $x \in X$ and $a \in \mathbb{N}$. Note that this implies that $f_x(x)(\infty) = f_2(x)(\infty)$ for each $x \in X$. This concludes the proof. \square

Proof (Prop. IV.7). We prove that $(r_p, q_p, \sqsubseteq_{\mathcal{DN}_\infty})$ satisfies the conditions in Def. III.10.

Cond. 1: Let $(\Gamma, t) \in F_p \mathcal{DN}_\infty = \mathcal{D}^2 \mathbb{N}_\infty \times \{0, 1\}$. If $t = 1$ then by the definitions of r_p and σ_p , we have:

$$q_p \circ r_p(\Gamma, t) = q_p(\delta_0) = 1 = \sigma_p(\mathcal{D}q_p(\Gamma), t) = \sigma_p \circ F_p q_p(\Gamma, t).$$

Assume $t = 0$. Then we have:

$$\begin{aligned}
q_p \circ r_p(\Gamma, t) &= r_p(\Gamma, 1)([0, \infty)) && (\text{by def. of } q_p) \\
&= \sum_{a=0}^{\infty} r_p(\Gamma, 1)(a) \\
&= \sum_{a=1}^{\infty} \sum_{\gamma \in \text{supp}(\Gamma)} \Gamma(\gamma) \cdot \gamma(a-1) && (\text{by def. of } r_p) \\
&= \sum_{a=0}^{\infty} \sum_{\gamma \in \text{supp}(\Gamma)} \Gamma(\gamma) \cdot \gamma(a) \\
&= \sum_{\gamma \in \text{supp}(\Gamma)} \Gamma(\gamma) \cdot \left(\sum_{a=0}^{\infty} \gamma(a) \right) \\
&= \sum_{\gamma \in \text{supp}(\Gamma)} \Gamma(\gamma) \cdot q_p(\gamma) && (\text{by def. of } q_p) \\
&= \sigma_p \circ F_p q_p(\Gamma, t) && (\text{by def. of } \sigma_p).
\end{aligned}$$

Hence Cond. 1 is satisfied. Note also that we have $q_p \circ r_p = \sigma_p \circ F_p q_p$.

Cond. 2: We have already shown in Lem. A.14.

Cond. 3: We first prove that q_p is monotone. Let $f_1, f_2 : X \rightarrow \mathcal{DN}_\infty$ and assume that $f \sqsubseteq_{\mathcal{DN}_\infty} g$. Then we have:

$$\begin{aligned}
q_p \circ f_1(x) &= f_1(x)([0, \infty)) && (\text{by def. of } q_p) \\
&= \lim_{a \rightarrow \infty} f_1(x)([0, a]) \\
&\leq \lim_{a \rightarrow \infty} f_2(x)([0, a]) && (\text{by } f_1 \sqsubseteq_{\mathcal{DN}_\infty} f_2)
\end{aligned}$$

$$\begin{aligned}
&= f_2(x)([0, \infty)) \\
&= q_p \circ f_2(x) && (\text{by def. of } q_p).
\end{aligned}$$

Hence we have $q_p \circ f_1 \leq q_p \circ f_2$, and therefore Cond. 3 was proved.

By the definition of q_p , we have $q_p(\delta_\infty) = 0$. Therefore q_p is strict.

We prove that q_p is continuous. By the definition of q_p , it suffices to prove $(\bigsqcup_{\gamma \in K} \gamma)([0, \infty)) = \sup_{\gamma \in K} (\gamma([0, \infty)))$. Note that the supremum on the left-hand side is taken with respect to $\sqsubseteq_{\mathcal{DN}_\infty}$ over \mathcal{DN}_∞ while the latter is taken with respect to the ordinary order over $[0, 1]$. We have:

$$\begin{aligned}
&\left(\bigsqcup_{\gamma \in K} \gamma \right)([0, \infty)) \\
&= \lim_{a \rightarrow \infty} \left(\bigsqcup_{\gamma \in K} \gamma \right)([0, a]) \\
&= \lim_{a \rightarrow \infty} \sup_{\gamma \in K} (\gamma([0, a])) && (\text{see the proof of Lem. A.12}) \\
&= \sup_{\gamma \in K} \lim_{a \rightarrow \infty} (\gamma([0, a])) \\
&= \sup_{\gamma \in K} (\gamma([0, \infty))).
\end{aligned}$$

Therefore q_p is continuous.

Cond. 4: We have already shown in Lem. A.15.

Hence the triple $(r_p, q_p, \sqsubseteq_{\mathcal{DN}_\infty})$ is a ranking domain. \square

I. Soundness of Additive Ranking Supermartingale, Categorically

In §IV-A, we have seen that the triple $(r'_{p,\varepsilon}, q'_p, \sqsubseteq_{[0,\infty)})$, which captures the definition of ε -additive ranking supermartingales, is not a ranking domain because $r'_{p,\varepsilon}$ is not a corecursive algebra. Hence we cannot prove soundness of ε -additive ranking supermartingales directly using our categorical framework. In this section, we show that its soundness is proved via that of distribution-valued ranking functions. More concretely, we have the following proposition.

Proposition A.16. *We define triples $(r_p, q_p, \sqsubseteq_{\mathcal{DN}_\infty})$ and $(r'_{p,\varepsilon}, q'_p, \sqsubseteq_{[0,\infty)})$ as in Prop. IV.7 and Prop. IV.3 respectively. Let $c : X \rightarrow F_p X$ be an F_p -coalgebra. Then we have:*

$$\begin{aligned}
\exists b' : X &\rightarrow [0, \infty]. \quad b' \sqsubseteq_{[0,\infty)} \Phi_{c,r'_{p,\varepsilon}}(b') \\
&\Rightarrow \exists b : X \rightarrow \mathcal{DN}_\infty. \\
&\quad b \sqsubseteq_{\mathcal{DN}_\infty} \Phi_{c,r_p}(b), \quad \text{and} \\
&\quad \forall x \in X. (q'_p \circ b'(x) = 1 \Rightarrow q_p \circ b(x) = 1).
\end{aligned}$$


This proposition is an immediate corollary of the following lemma.

Lemma A.17. *We assume the conditions in Prop. A.16. We define a function $p_p : \mathcal{DN}_\infty \rightarrow [0, \infty]$ by*

$$p_p(\gamma) = \varepsilon \cdot \sum_{a \in \text{supp}(\gamma)} a \cdot \gamma(a).$$

Then the following statements hold.

- 1) $q'_p \circ p_p \leq q_p$ (here \leq denotes the pointwise extension over $\mathcal{DN}_\infty^{([0, \infty])}$).
- 2) Let $b' : X \rightarrow [0, \infty]$ and assume $b' \sqsubseteq_{[0, \infty]} \Phi_{c, r'_{p, \varepsilon}}(b')$. As $r_p : F_p \mathcal{DN}_\infty \rightarrow \mathcal{DN}_\infty$ is corecursive (Prop. IV.7), there exists a unique function $(\llbracket c \rrbracket)_{r_p} : X \rightarrow \mathcal{DN}_\infty$ such that $(\llbracket c \rrbracket)_{r_p} = \Phi_{c, r_p}((\llbracket c \rrbracket)_{r_p})$. For this function, we have $b' \sqsubseteq_{[0, \infty]} p_p \circ (\llbracket c \rrbracket)_{r_p}$.



Sublemma A.18. For the function $p_p : \mathcal{DN}_\infty \rightarrow [0, \infty]$ in Lem. A.17, we have the following.

$$p_p \circ r_p = r'_{p, \varepsilon} \circ F_p p_p \quad (11)$$

Proof. Let $(\Gamma, t) \in F_p \mathcal{DN}_\infty = \mathcal{D}^2 \mathbb{N}_\infty \times \{0, 1\}$. If $t = 1$, then by the definitions of r_p and $r'_{p, \varepsilon}$, we have

$$p_p \circ r_p(\Gamma, t) = r'_{p, \varepsilon} \circ F_p p_p(\Gamma, t) = 0.$$

Let $t = 0$. Then we have:

$$\begin{aligned} p_p \circ r_p(\Gamma, t) &= \varepsilon \cdot \sum_{a \in \mathbb{N}_\infty \setminus \{0\}} a \cdot r_p(\Gamma, t)(a) && \text{(by def. of } p_p) \\ &= \varepsilon \cdot \sum_{a \in \mathbb{N}_\infty \setminus \{0\}} a \cdot \sum_{\gamma \in \text{supp}(\Gamma)} \Gamma(\gamma) \cdot \gamma(a-1) && \text{(by def. of } r_p) \\ &= \varepsilon \cdot \sum_{a \in \mathbb{N}_\infty} (a+1) \cdot \sum_{\gamma \in \text{supp}(\Gamma)} \Gamma(\gamma) \cdot \gamma(a) \\ &= \varepsilon \cdot \sum_{a \in \mathbb{N}_\infty} a \cdot \sum_{\gamma \in \text{supp}(\Gamma)} \Gamma(\gamma) \cdot \gamma(a) \\ &\quad + \varepsilon \cdot \sum_{a \in \mathbb{N}_\infty} \sum_{\gamma \in \text{supp}(\Gamma)} \Gamma(\gamma) \cdot \gamma(a) \\ &= \sum_{a \in \mathbb{N}_\infty} \left(a \cdot \varepsilon \cdot \sum_{\gamma \in \text{supp}(\Gamma)} \Gamma(\gamma) \cdot \gamma(a) \right) + \varepsilon \\ &= \sum_{\gamma \in \text{supp}(\Gamma)} \left(\Gamma(\gamma) \cdot \varepsilon \cdot \sum_{a \in \mathbb{N}_\infty} a \cdot \gamma(a) \right) + \varepsilon \\ &= \sum_{\gamma \in \text{supp}(\Gamma)} \left(\Gamma(\gamma) \cdot p_p(\gamma) \right) + \varepsilon && \text{(by def. of } p_p) \\ &= \sum_{b \in \text{supp}(\mathcal{D}p_p(\Gamma))} \left(b \cdot \sum_{\gamma' \in p_p^{-1}(b)} \Gamma(\gamma') \right) + \varepsilon \\ &= \sum_{b \in \text{supp}(\mathcal{D}p_p(\Gamma))} \left(b \cdot \mathcal{D}p_p(\Gamma)(b) \right) + \varepsilon \\ &= r'_{p, \varepsilon} \circ F_p p_p(\Gamma, t) && \text{(by def. of } r'_{p, \varepsilon}). \end{aligned}$$

This concludes the proof. \square


Sublemma A.19. The F_p -algebra $r'_{p, \varepsilon} : F_p [0, \infty] \rightarrow [0, \infty]$ in Prop. IV.3 satisfies the following conditions.

- 1) The poset $([0, \infty], \sqsubseteq_{[0, \infty]})$ is a complete lattice.

- 2) For each F_p -coalgebra $c : X \rightarrow F_p X$, the function $\Phi_{c, r'_{p, \varepsilon}} : [0, \infty]^X \rightarrow [0, \infty]^X$ in Def. III.5 is monotone with respect to the pointwise extension of $\sqsubseteq_{[0, \infty]}$.

Moreover the triple $(r_p : F_p \mathcal{DN}_\infty \rightarrow \mathcal{DN}_\infty, p_p : \mathcal{DN}_\infty \rightarrow [0, \infty], \sqsubseteq_{\mathcal{DN}_\infty})$ (see Prop. IV.7 and Lem. A.17) satisfies the following conditions.

- 3) We have $p_p \circ r_p \sqsubseteq_{[0, \infty]} r'_{p, \varepsilon} \circ F_p p_p$ between arrows $F_p \mathcal{DN}_\infty \rightarrow [0, \infty]$.
- 4) The following conditions are satisfied by r_p :
 - a) the poset $(\mathcal{DN}_\infty, \sqsubseteq_{\mathcal{DN}_\infty})$ has the greatest element $\top_{\mathcal{DN}_\infty}$;
 - b) the poset $(\mathcal{DN}_\infty, \sqsubseteq_{\mathcal{DN}_\infty})$ is ω^{op} -complete (i.e. for each decreasing sequence $\varphi_0 \sqsupseteq_{\mathcal{DN}_\infty} \varphi_1 \sqsupseteq_{\mathcal{DN}_\infty} \dots$ in \mathcal{DN}_∞ , the infimum $\prod_{i \in \omega} \varphi_i$ exists);
 - c) for each $c : X \rightarrow F_p X$, the function $\Phi_{c, r_p} : (\mathcal{DN}_\infty)^X \rightarrow (\mathcal{DN}_\infty)^X$ (Def. III.2) is monotone and moreover ω^{op} -continuous, i.e. for each decreasing sequence $f_0 \sqsupseteq_{\mathcal{DN}_\infty} f_1 \sqsupseteq_{\mathcal{DN}_\infty} \dots$ in $(\mathcal{DN}_\infty)^X$ wrt. the pointwise extension of $\sqsubseteq_{\mathcal{DN}_\infty}$, we have $\Phi_{c, r_p}(\prod_{i \in \omega} f_i) = \prod_{i \in \omega} \Phi_{c, r_p}(f_i)$.
- 5) The function $p_p : \mathcal{DN}_\infty \rightarrow [0, \infty]$ is monotone (i.e. $a \sqsubseteq_{\mathcal{DN}_\infty} b \Rightarrow p_p(a) \sqsubseteq_{[0, \infty]} p_p(b)$), top-preserving (i.e. $p_p(\top_{\mathcal{DN}_\infty}) = \top_{[0, \infty]}$) and ω^{op} -continuous (i.e. each decreasing sequence $\varphi_0 \sqsupseteq_{\mathcal{DN}_\infty} \varphi_1 \sqsupseteq_{\mathcal{DN}_\infty} \dots$ in \mathcal{DN}_∞ , we have $p_p(\prod_{i \in \omega} \varphi_i) = \prod_{i \in \omega} p_p(\varphi_i)$). Note that in the last equality, the infimum in the left-hand side is taken wrt. $\sqsubseteq_{\mathcal{DN}_\infty}$ while the one in the right-hand side is taken wrt. $\sqsubseteq_{[0, \infty]}$.
- 6) The algebra $r_p : F_p \mathcal{DN}_\infty \rightarrow \mathcal{DN}_\infty$ is corecursive.



Proof.

1: Easy.

2: Already proved in Prop. IV.4.

3: Immediate from Sublem. A.18.

4: It is easy to see that the Dirac distribution δ_0 concentrated at 0 is the greatest element in $(\mathcal{DN}_\infty, \sqsubseteq_{\mathcal{DN}_\infty})$. Hence Cond. 4a is satisfied. Cond. 4b is immediate from that $((\mathcal{DN}_\infty)^X, \sqsubseteq_{\mathcal{DN}_\infty})$ is a complete lattice (Lem. A.12).

We prove that Cond. 4c is satisfied. Monotonicity of Φ_{c, r_p} is already proved in Prop. IV.7. We prove that Φ_{c, r_p} is ω^{op} -continuous. To this end, it suffices to prove the following equality for each $x \in X$.

$$\Phi_{c, r_p} \left(\prod_{i \in \omega} f_i \right)(x) = \prod_{i \in \omega} (\Phi_{c, r_p}(f_i)(x)) \quad (12)$$

Note that \prod on the right-hand side denotes the infimum with respect to $\sqsubseteq_{\mathcal{DN}_\infty}$ while that on the left-hand side denotes the infimum with respect to its pointwise extension. Let $(\delta, t) = c(x)$.

If $t = 1$, then by the definition of r_p , we have $\Phi_{c,r_p}(f)(x) = \delta_0$ for each $f : X \rightarrow \mathcal{DN}_\infty$. Hence we have (12).

If $t = 0$, for each $a \in \mathbb{N}$, we have:

$$\begin{aligned}
& \left(\prod_{i \in \omega} (\Phi_{c,r_p}(f_i)(x)) \right) ([0, a]) \\
&= \inf_{i \in \omega} (\Phi_{c,r_p}(f_i)(x)([0, a])) \quad (\text{see the proof of Lem. A.12}) \\
&= \inf_{i \in \omega} \sum_{x \in \text{supp}(\delta)} \delta(x) \cdot (f_i(x)([0, a-1])) \quad (\text{by def. of } r_p) \\
&= \sum_{x \in \text{supp}(\delta)} \delta(x) \cdot \left(\inf_{i \in \omega} f_i(x)([0, a-1]) \right) \\
&\quad (\text{by the dominated convergence theorem}) \\
&= \sum_{x \in \text{supp}(\delta)} \delta(x) \cdot \left(\prod_{i \in \omega} f_i(x)([0, a-1]) \right) \\
&\quad (\text{see the proof of Lem. A.12}) \\
&= \Phi_{c,r_p} \left(\prod_{i \in \omega} f_i(x)([0, a]) \right) \quad (\text{by def. of } \Phi_{c,r_p}).
\end{aligned}$$

By Sublem. A.13, this implies (12).

5: We first prove that $p_p : \mathcal{DN}_\infty \rightarrow [0, \infty]$ is monotone.

Let $\varphi_1, \varphi_2 \in \mathcal{DN}_\infty$ and assume that $\varphi_1 \sqsubseteq_{\mathcal{DN}_\infty} \varphi_2$.

If $\varphi_2(\infty) = C > 0$, then we have $\varphi_2([0, a]) \leq 1 - C$ for each $a \in \mathbb{N}$. By the definition of $\sqsubseteq_{\mathcal{DN}_\infty}$, this implies $\varphi_1([0, a]) \leq 1 - C$ for each a , and therefore we have $\varphi_1(\infty) > 0$. Hence in this case, we have $p_p(\varphi_1) = p_p(\varphi_2) = \infty$.

Let $\varphi_2(\infty) = 0$. Then we have:

$$\begin{aligned}
& p_p(\varphi_2) \\
&= \varepsilon \cdot \sum_{a=1}^{\infty} a \cdot \varphi_2(a) \quad (\text{by def. of } p_p) \\
&= \lim_{n \rightarrow \infty} \varepsilon \cdot \sum_{a=1}^n (a \cdot \varphi_2([0, a]) - a \cdot \varphi_2([0, a-1])) \\
&= \lim_{n \rightarrow \infty} \varepsilon \cdot \left(n \cdot \varphi_2([0, n]) - \sum_{a=1}^n 1 \cdot \varphi_2([0, a-1]) \right) \\
&= \lim_{n \rightarrow \infty} \varepsilon \cdot \sum_{a=1}^n \varphi_2([a, n]) \\
&= \varepsilon \cdot \sum_{a=1}^{\infty} \varphi_2([a, \infty]) \\
&= \varepsilon \cdot \sum_{a=1}^{\infty} (1 - \varphi_2([0, a-1])) \\
&\leq \varepsilon \cdot \sum_{a=1}^{\infty} (1 - \varphi_1([0, a-1])) \quad (\text{by } \varphi_1 \sqsubseteq_{\mathcal{DN}_\infty} \varphi_2) \\
&= p_p(\varphi_1) \quad (\text{as the transformation for } \varphi_2).
\end{aligned}$$

Hence we have $p_p(\varphi_1) \sqsubseteq_{[0, \infty]} p_p(\varphi_2)$ and therefore p_p is monotone.

By the definition of p_p , $p_p(\delta_0) = 0$. Hence p_p is top-preserving.

We prove that p_p is ω^{op} -continuous. Let $\varphi_0 \sqsupseteq_{\mathcal{DN}_\infty} \varphi_1 \sqsupseteq_{\mathcal{DN}_\infty} \dots$ be a decreasing sequence in \mathcal{DN}_∞ .

Assume $\left(\prod_{i \in \omega} (\varphi_i) \right) (\infty) = C > 0$. Then for all $b \in \mathbb{N}$, there exists $i_b \in \omega$ such that $1 - \varphi_{i_b}([0, b]) \geq \frac{C}{2}$ (c.f. the proof of Lem. A.12). Therefore for each $b \in \mathbb{N}$, we have:

$$\begin{aligned}
& \prod_{i \in \omega} p_p(\varphi_i) \geq p_p(\varphi_{i_b}) \\
&= \varepsilon \cdot \sum_{a=0}^{\infty} a \cdot \varphi_{i_b}(a) \quad (\text{by def. of } p_p) \\
&\geq \varepsilon \cdot b \cdot (1 - \varphi_{i_b}([0, b])) \\
&\geq \varepsilon \cdot b \cdot \left(1 - \frac{C}{2} \right).
\end{aligned}$$

Hence we have

$$p_p \left(\prod_{i \in \omega} (\varphi_i) \right) = \infty = \prod_{i \in \omega} p_p(\varphi_i).$$

Assume $\left(\prod_{i \in \omega} (\varphi_i) \right) (\infty) = 0$. Then we have:

$$\begin{aligned}
& p_p \left(\prod_{i \in \omega} (\varphi_i) \right) \\
&= \varepsilon \cdot \sum_{a=0}^{\infty} a \cdot \left(\prod_{i \in \omega} \varphi_i \right) (a) \quad (\text{by def. of } p_p) \\
&= \varepsilon \cdot \sum_{a=1}^{\infty} \left(1 - \left(\prod_{i \in \omega} \varphi_i \right) ([0, a-1]) \right) \\
&\quad (\text{as in the proof of monotonicity of } p_p) \\
&= \varepsilon \cdot \sum_{a=1}^{\infty} \left(1 - \inf_{i \in \omega} (\varphi_i([0, a-1])) \right) \\
&\quad (\text{see the proof of Lem. A.12}) \\
&= \sup_{i \in \omega} \varepsilon \cdot \sum_{a=1}^{\infty} \left(1 - (\varphi_i([0, a-1])) \right) \\
&= \sup_{i \in \omega} p_p(\varphi_i) \quad (\text{as in the above}) \\
&= \prod_{i \in \omega} p_p(\varphi_i) \quad (\text{by def. of } \sqsubseteq_{[0, \infty]})
\end{aligned}$$

Hence we have ω^{op} -continuity.

6: It is already proved in Lem. A.15. \square

Proof (Lem. A.17).

1: For each $\varphi \in \mathcal{DN}_\infty$, we have:

$$\begin{aligned}
& q'_p \circ p_p(\varphi) \\
&= q'_p \left(\sum_{a \in \mathbb{N}} a \cdot \varphi(a) \right) \quad (\text{by def. of } p_p) \\
&= \begin{cases} 1 & (\sum_{a \in \mathbb{N}} a \cdot \varphi(a) < \infty) \\ 0 & (\text{otherwise}) \end{cases} \quad (\text{by def. of } q'_p) \\
&\leq \varphi([0, \infty)) \\
&= q_p(\varphi) \quad (\text{by def. of } q_p).
\end{aligned}$$

Hence Cond. 1 is satisfied.

2:

$$\begin{array}{ccccc}
& & F_p \llbracket \nu r'_{p,\varepsilon} \rrbracket_c & & \\
F_p X & \xrightarrow{F_p \llbracket \nu r'_{p,\varepsilon} \rrbracket_c} & F_p \mathcal{DN}_\infty & \xrightarrow{F_p p} & F_p[0, \infty] \\
\uparrow c & \xrightarrow{F_p \langle c \rangle_{r_p}} & = & \downarrow r_p & \Downarrow \exists & \downarrow r'_{p,\varepsilon} \\
X & \xrightarrow{\langle c \rangle_{r_p}} & \mathcal{DN}_\infty & \xrightarrow{p} & [0, \infty] \\
& \xrightarrow{\llbracket \nu r'_{p,\varepsilon} \rrbracket_c} & & & \\
& & b' & &
\end{array}$$

By Sublem. A.19.1–2, the F_p -modality $r'_{p,\varepsilon} : F_p[0, \infty] \rightarrow [0, \infty]$ satisfy dual conditions of Asm. III.7. Hence using the dual statement of Prop. III.8, we can show that $\Phi_{c,r'_{p,\varepsilon}} : [0, \infty]^X \rightarrow [0, \infty]^X$ has the greatest fixed point $\llbracket \nu r'_{p,\varepsilon} \rrbracket_c : X \rightarrow [0, \infty]$ with respect to the pointwise extension of $\sqsubseteq_{[0, \infty]}$. By the Knaster-Tarski theorem, we have

$$b' \sqsubseteq_{[0, \infty]} \llbracket \nu r'_{p,\varepsilon} \rrbracket_c. \quad (13)$$

By Sublem. A.19.3–6, the triple $(r_p : F_p \mathcal{DN}_\infty \rightarrow \mathcal{DN}_\infty, p_p : \mathcal{DN}_\infty \rightarrow [0, \infty], \sqsubseteq_{[0, \infty]})$ satisfies the dual conditions of the axioms of a ranking domain (Def. III.10), except that the length of a transfinite sequence in $(\mathcal{DN}_\infty)^X$ is restricted to ω . Note here that $\langle c \rangle_{r_p} : X \rightarrow \mathcal{DN}_\infty$ satisfies $\langle c \rangle_{r_p} \sqsubseteq_{\mathcal{DN}_\infty} \Phi_{c,r_p}(\langle c \rangle_{r_p})$ by its definition. Therefore in a similar manner to the proof of Thm. III.12, we can prove

$$\llbracket \nu r'_{p,\varepsilon} \rrbracket_c \sqsubseteq_{[0, \infty]} p_p \circ \langle c \rangle_{r_p}. \quad (14)$$

By (14) and (13), we have $b' \sqsubseteq_{[0, \infty]} p_p \circ \langle c \rangle_{r_p}$. \square

Proof (Prop. A.16). Immediate from Lem. A.17. \square

We can now prove the soundness of additive ranking supermartingale (Thm. II.5) using our categorical framework as follows.

$\exists b'. b'$ is an ε -additive ranking supermartingale and $b'(x) \neq \infty$

$$\stackrel{\text{Prop. IV.3}}{\Leftrightarrow} \exists b'. b' \sqsubseteq_{[0, \infty]} \Phi_{c,r'_{p,\varepsilon}}(b') \text{ and } q'_p \circ b'(x) = 1$$

$$\stackrel{\text{Prop. A.16}}{\Rightarrow} \exists b. b \text{ is a ranking arrow wrt. } (r_p, q_p, \sqsubseteq_{\mathcal{DN}_\infty}), \text{ and } q_p \circ b(x) = 1$$

$$\stackrel{\text{Thm. III.12}}{\Rightarrow} \llbracket \mu \sigma_p \rrbracket_{c,G,Acc}(x) = 1$$

$$\stackrel{\text{Prop. IV.2}}{\Leftrightarrow} \text{Reach}_{M,Acc}(x) = 1$$

J. Soundness of Multiplicative Ranking Supermartingale, Categorically

Proposition A.20. We define triples $(r_p, q_p, \sqsubseteq_{\mathcal{DN}_\infty})$ and $(r'_{p_m,\alpha}, q'_p, \sqsubseteq_{[\alpha\delta, \infty]})$ as in Prop. IV.7 and Prop. A.9 respectively. Let $c : X \rightarrow F_p X$ be an F_p -coalgebra. Then we have:

$$\begin{aligned} & \exists b'' : X \rightarrow [\alpha\delta, \infty]. b'' \sqsubseteq_{[\alpha\delta, \infty]} \Phi_{c,r'_{p_m,\alpha}}(b'') \\ & \Rightarrow \exists b : X \rightarrow \mathcal{DN}_\infty. \end{aligned}$$

$$\begin{aligned} & b \sqsubseteq_{\mathcal{DN}_\infty} \Phi_{c,r_p}(b), \text{ and} \\ & \forall x \in X. (q'_p \circ b'(x) = 1 \Rightarrow q_p \circ b(x) = 1). \end{aligned}$$

This proposition is an immediate corollary of Prop. A.16 and the following lemma.

Lemma A.21. We define triples $(r'_{p,\varepsilon}, q'_p, \sqsubseteq_{[0, \infty]})$ and $(r'_{p_m,\alpha}, q'_p, \sqsubseteq_{[\alpha\delta, \infty]})$ as in Prop. IV.3 and Prop. A.9 respectively. We define a function $p'_p : [\alpha\delta, \infty] \rightarrow [0, \infty]$ as follows.

$$p'_p(a) = \varepsilon \cdot \left(\left(\log_{\frac{1}{\alpha}} \frac{a}{\delta} \right) + 1 \right)$$

Let $c : X \rightarrow F_p X$ be an F_p -coalgebra. Then we have:

$$\begin{aligned} & \exists b'' : X \rightarrow [\alpha\delta, \infty]. b'' \sqsubseteq_{[\alpha\delta, \infty]} \Phi_{c,r'_{p_m,\alpha}}(b'') \\ & \Rightarrow p'_p \circ b'' \sqsubseteq_{[0, \infty]} \Phi_{c,r'_{p,\varepsilon}}(p'_p \circ b'') \end{aligned}$$

Sublemma A.22. For the function $F[\alpha\delta, \infty] \xrightarrow{Fp'_p} F[0, \infty]$ $p'_p : [\alpha\delta, \infty] \rightarrow [0, \infty]$ in Lem. A.21, $r'_{p_m,\alpha} \downarrow \sqsubseteq r'_{p,\varepsilon} \downarrow$ we have $p'_p \circ r'_{p_m,\alpha} \sqsubseteq_{[0, \infty]} r'_{p,\varepsilon} \circ F_p p'_p$. $[\alpha\delta, \infty] \xrightarrow{p'_p} [0, \infty]$

Proof. Let $(\psi, t) \in F_p[\alpha\delta, \infty] = \mathcal{D}[\alpha\delta, \infty] \times \{0, 1\}$. By the definition of $\sqsubseteq_{[0, \infty]}$, it suffices to prove $p'_p \circ r'_{p_m,\alpha}(\psi, t) \geq r'_{p,\varepsilon} \circ F_p p'_p(\psi, t)$.

If $t = 1$, then we have:

$$\begin{aligned} & p'_p \circ r'_{p_m,\alpha}(\psi, t) \\ & = p'_p(\alpha\delta) \quad (\text{by def. of } r'_{p_m,\alpha}) \\ & = 0 \quad (\text{by def. of } p'_p) \\ & = r'_{p,\varepsilon}(\mathcal{D}p'_p(\psi), t) \quad (\text{by def. of } r'_{p,\varepsilon}) \\ & = r'_{p,\varepsilon} \circ F_p p'_p(\psi, t). \end{aligned}$$

Let $t = 0$. Note that $\sum_{a \in \text{supp}(\psi)} \psi(a) = 1$. Hence we have:

$$\begin{aligned} & p'_p \circ r'_{p_m,\alpha}(\psi, t) \\ & = p'_p \left(\frac{1}{\alpha} \cdot \left(\sum_{a \in \text{supp}(\psi)} a \cdot \psi(a) \right) \right) \quad (\text{by def. of } r'_{p_m,\alpha}) \\ & = \varepsilon \cdot \left(\log_{\frac{1}{\alpha}} \frac{1}{\delta} \left(\frac{1}{\alpha} \cdot \left(\sum_{a \in \text{supp}(\psi)} a \cdot \psi(a) \right) \right) + 1 \right) \quad (\text{by def. of } p'_p) \\ & = \varepsilon \cdot \left(\left(\log_{\frac{1}{\alpha}} \left(\sum_{a \in \text{supp}(\psi)} \frac{a}{\delta} \cdot \psi(a) \right) \right) + 2 \right) \\ & \geq \varepsilon \cdot \left(\left(\sum_{a \in \text{supp}(\psi)} \left(\log_{\frac{1}{\alpha}} \frac{a}{\delta} \right) \cdot \psi(a) \right) + 2 \right) \\ & \quad (\log_{\frac{1}{\alpha}}(_) \text{ is an upward convex function}) \\ & = \left(\sum_{a \in \text{supp}(\psi)} \left(\varepsilon \cdot \left(\log_{\frac{1}{\alpha}} \frac{a}{\delta} \right) + 1 \right) \cdot \psi(a) \right) + \varepsilon \\ & = r'_{p,\varepsilon} \circ F_p p'_p(\varphi, t) \quad (\text{by def. of } p'_p \text{ and } r'_{p,\varepsilon}). \end{aligned}$$

This concludes the proof. \square

Proof (Lem. A.21).

$$\begin{array}{ccccc}
FX & \xrightarrow{Fb''} & F[\alpha\delta, \infty] & \xrightarrow{Fp'_p} & F[0, \infty] \\
\uparrow c & \sqcup & r'_{p_m,\alpha} \downarrow & \sqsubseteq & r'_{p,\varepsilon} \downarrow \\
X & \xrightarrow{b''} & [\alpha\delta, \infty] & \xrightarrow{p'_p} & [0, \infty]
\end{array}$$

Let $b'' : X \rightarrow [\alpha\delta, \infty]$ and assume that $b'' \sqsubseteq_{[\alpha\delta, \infty]} \Phi_{c, r'_{pm, \alpha}}(b'')$. Then we have:

$$\begin{aligned} & p'_p \circ b'' \\ & \sqsubseteq_{[0, \infty]} p'_p \circ r'_{pm, \alpha} \circ F_p b'' \circ c \quad (\text{by the assumption}) \\ & \sqsubseteq_{[0, \infty]} r'_{p, \varepsilon} \circ F_p p'_p \circ F_p b'' \circ c \quad (\text{by Sublem. A.22}) \\ & = \Phi_{c, r'_{p, \varepsilon}}(p'_p \circ b'') \quad (\text{by definition}). \end{aligned}$$

This concludes the proof. \square

Proof (Prop. A.20). Immediate from Lem. A.21 and Prop. A.16. \square

We can now prove the soundness of multiplicative ranking supermartingale using our categorical framework as follows.

$\exists b''. b''$ is an α -multiplicative ranking supermartingale, and $b''(x) \neq \infty$

$\stackrel{\text{Prop. A.9}}{\Leftrightarrow} \exists b'. b' \sqsubseteq_{[0, \infty]} \Phi_{c, r'_{p, \varepsilon}}(b')$ and $q'_p \circ b'(x) = 1$

$\stackrel{\text{Prop. A.16}}{\Rightarrow} \exists b. b$ is a ranking arrow wrt. $(r_p, q_p, \sqsubseteq_{\mathcal{DN}_\infty})$, and $q_p \circ b(x) = 1$

$\stackrel{\text{Thm. III.12}}{\Rightarrow} \llbracket \mu \sigma_p \rrbracket_{cG, \text{Acc}}(x) = 1$

$\stackrel{\text{Prop. IV.2}}{\Leftrightarrow} \text{Reach}_{M, \text{Acc}}(x) = 1$

K. Proof of Prop. IV.12

We prove that $r_{nc, \gamma}$ is a corecursive algebra separately.

Lemma A.23. *The algebra $r_{nc, \gamma} : F_p[0, 1] \rightarrow [0, 1]$ in Prop IV.12 satisfies Cond. 2 in Def. III.10.*

Proof. It is easy to see that Cond. 2a is satisfied.

We prove that Cond. 2b is satisfied. Let $b_1, b_2 : X \rightarrow [0, 1]$ and assume that $b_1 \leq b_2$. Let $x \in X$ and assume that $c(x) = (\delta, t) \in F_p X = \mathcal{DX} \times \{0, 1\}$. Then we have:

$$\begin{aligned} & r_{nc, \gamma} \circ F_p b_1(\delta, t) \\ & = \begin{cases} 1 & (t = 1) \\ \sum_{x \in X} \delta(x) \cdot b_1(x) & (t = 0) \end{cases} \\ & \quad (\text{by def. of } F_p \text{ and } r_{nc, \gamma}) \\ & = \begin{cases} 1 & (t = 1) \\ \sum_{x \in X} \delta(x) \cdot b_2(x) & (t = 0) \end{cases} \quad (\text{by } b_1 \leq b_2) \\ & = r_{nc, \gamma} \circ F_p b_2(\delta, t) \quad (\text{by def. of } r_{nc, \gamma} \text{ and } F_p). \end{aligned}$$

Therefore Cond. 2b is satisfied. \square

Lemma A.24. *The algebra $r_{nc, \gamma} : F_p[0, 1] \rightarrow [0, 1]$ in Prop. IV.12 is corecursive.*

Proof. Let $c : X \rightarrow F_p X$ be an F_p -coalgebra. For each $x \in X$ such that $c(x) = (\delta, t) \in F_p X$, we write $c_1(x)$ and $c_2(x)$ for $\delta \in \mathcal{D}[0, 1]$ and $t \in \{0, 1\}$ respectively. We prove that $\Phi_{c, r_{nc, \gamma}}$ has a unique fixed point.

We first show that $\Phi_{c, r_{nc, \gamma}}$ has a fixed point. By Lem. A.23, the poset $([0, 1]^X, \leq)$ (here \leq denotes the pointwise extension)

is a complete lattice. Therefore we can construct a transfinite sequence

$$\perp_{[0, 1]} \leq \Phi_{c, r_{nc, \gamma}}(\perp_{[0, 1]}) \leq \dots \leq \Phi_{c, r_{nc, \gamma}}^n(\perp_{[0, 1]}) \leq \dots$$

as in Thm. II.18.2. By Thm. II.18.2, there exists an ordinal n such that $\Phi_{c, r_{nc, \gamma}}^n(\perp_{[0, 1]})$ is the least fixed point of $\Phi_{c, r_{nc, \gamma}}$. Let $f = \Phi_{c, r_{nc, \gamma}}^n(\perp_{[0, 1]})$.

It remains to show that f is the unique fixed point of $\Phi_{c, r_{nc, \gamma}}$. Let $g : X \rightarrow [0, 1]$ be a fixed point of $\Phi_{c, r_{nc, \gamma}}$. As f is the least fixed point, we have $f(x) \leq g(x)$ for each $x \in X$. We now define $h : X \rightarrow [0, 1]$ by $h(x) = g(x) - f(x)$. Then we have:

$$\begin{aligned} & \sup_{x \in X} h(x) \\ & = \sup_{x \in X} (g(x) - f(x)) \quad (\text{by def. of } h) \\ & = \sup_{x \in X} (\Phi_{c, r_{nc, \gamma}}(g)(x) - \Phi_{c, r_{nc, \gamma}}(f)(x)) \\ & \quad (f \text{ and } g \text{ are fixed points}) \\ & \leq \sup_{x \in X} \left(\gamma \cdot \sum_{x' \in X} c_1(x)(x') \cdot g(x') \right. \\ & \quad \left. - \gamma \cdot \sum_{x' \in X} c_1(x)(x') \cdot f(x') \right) \\ & \quad (\text{by def. of } r_{nc, \gamma} \text{ and that } \\ & \quad c_2(x) = 1 \Rightarrow \Phi_{c, r}(f)(x) = \Phi_{c, r}(g)(x) = 1) \\ & = \gamma \cdot \sup_{x \in X} \sum_{x' \in X} c_1(x)(x') \cdot (g(x') - f(x')) \\ & = \gamma \cdot \sup_{x \in X} \sum_{x' \in X} c_1(x)(x') \cdot h(x') \quad (\text{by def. of } h) \\ & \leq \gamma \cdot \sup_{x \in X} \sup_{x' \in X} h(x') \quad (\text{by } \sum_{x' \in X} c_1(x)(x') \leq 1) \\ & = \gamma \cdot \sup_{x \in X} h(x) \end{aligned}$$

As $0 \leq \gamma < 1$, we have $\sup_{x \in X} h(x) = 0$, and this implies $f = g$. This concludes the proof. \square

Proof (Prop. IV.12). We prove that $(r_{nc, \gamma}, q_{nc}, \leq)$ satisfies the conditions in Def. III.10.

Cond. 1: Let $(\delta, t) \in F_p[0, 1] = \mathcal{D}[0, 1] \times \{0, 1\}$. Then we have:

$$\begin{aligned} & q_{nc} \circ r_{nc, \gamma}(\delta, t) \\ & = \begin{cases} q_{nc}(1) & (t = 1) \\ q_{nc}(\gamma \cdot \sum_{a \in \text{supp}(\delta)} a \cdot \delta(a)) & (t = 0) \end{cases} \quad (\text{by def. of } r_{nc, \gamma}) \\ & = \begin{cases} 1 & (t = 1) \\ \gamma \cdot \sum_{a \in \text{supp}(\delta)} a \cdot \delta(a) & (t = 0) \end{cases} \quad (\text{by def. of } q_{nc}) \\ & \leq \begin{cases} 1 & (t = 1) \\ \sum_{a \in \text{supp}(\delta)} a \cdot \delta(a) & (t = 0) \end{cases} \quad (\text{by } \gamma < 1) \\ & = \sigma_p(\delta, t) \quad (\text{by def. of } \sigma_p) \\ & = \sigma_p \circ F_p q_{nc}(\delta, t) \quad (\text{by def. of } q_{nc}). \end{aligned}$$

Hence Cond. 1 is satisfied.

Cond. 2: We have already proved in Lem. A.23.

Cond. 3: Immediate from $q_{nc} = \text{id}_{[0,1]}$.

Cond. 4: It is already proved in Lem. A.24.

Hence $(r_{nc,\gamma}, q_p, \leq)$ is a ranking domain. \square

L. Towards Further Examples

Now that we have a general categorical axiomatization of ranking functions (§III), we would like to exploit it in deriving further examples of “ranking functions” that are previously unknown, hoping that they will provide novel proof methods for various liveness properties. In the previous section we derived two variations of ranking supermartingales. Towards further examples, here we indicate a possible direction.

We can say that not many concrete examples are known of corecursive algebras. Nevertheless, the following “closure properties” can be used to derive new examples.

Lemma A.25. *Let $F: \mathbb{C} \rightarrow \mathbb{C}$ be a functor.*

- 1) ([33]) *Consider the well-known construction of the final sequence: $1 \xleftarrow{!} F1 \xleftarrow{F!} F^2 1 \xleftarrow{F^2!} \dots \xleftarrow{F^\alpha!} \dots$, where we define $F^\alpha 1$ for an arbitrary ordinal α (we assume enough limits and let $F^\alpha 1 = \lim_{b < \alpha} F^b 1$ for a limit ordinal α). Then, for each α , the algebra $F^\alpha! : F(F^\alpha 1) \rightarrow F^\alpha 1$ is corecursive.*
- 2) (dual of [34]) *Let $\tau: F \Rightarrow G$ be a natural transformation. Then a corecursive G -algebra $s: GR \rightarrow R$ induces a corecursive F -algebra $s \circ \tau_R: FR \rightarrow R$.* \square

Lem. A.25.1–2 together suggest the following workflow. For dynamical systems modeled as F -coalgebras, we pick a natural transformation $\tau: F \Rightarrow G$ and an ordinal α —the former *abstracts* (or *collapses*) F -behaviors into G -behaviors that are supposedly simpler. We then use the set $G^\alpha 1$ of “ G -behaviors up-to α ” as the ranking domain. The set carries a corecursive G -algebra by Lem. A.25.1; and via τ it carries a corecursive F -algebra, too (Lem. A.25.2).

An example of such “behavioral” ranking domains is in §M. The set $\text{Tree}_{\text{fin}}^\perp$ consists of: (unlabeled) trees that are possibly countably branching and of finite depth; and the special element \perp that designates non-well-foundedness. It is shown that, for the problem of universal reachability of tree automata, we can indeed use the set $\text{Tree}_{\text{fin}}^\perp$ to form a (categorical) ranking domain. Here we use the functors $F = (\prod_{n \in \mathbb{N} \cup \{\omega\}} \Sigma_n \times (_)^n) \times \{0, 1\}$ and $G = 1 + \prod_{n \in \mathbb{N} \cup \{\omega\}} (_)^n$, and $\tau: F \Rightarrow G$ collapses the elements $(t, 1) \in FX$ —i.e. accepting states—to the unique element of 1 in GX .

In fact the set $\text{Tree}_{\text{fin}}^\perp$ is not precisely the outcome $G^\omega 1$ of the workflow described above: $G^\omega 1$ contains all the finite and infinite trees (with suitable branching degrees) as separate elements; but in $\text{Tree}_{\text{fin}}^\perp$ a single element \perp stands for all the “non-well-founded” trees that contains at least one infinite branch. A categorical description of such collapse is our future work; so is a general order with which we can equip $G^\omega 1$.

M. A Ranking Domain for Tree Automata

Notation A.26. We write $\langle \rangle$ for the empty sequence in \mathbb{N}^* . For $w, w' \in \mathbb{N}^* \cup \mathbb{N}^\omega$, we write $w \preceq w'$ if w is a prefix of w' .

Definition A.27 (unlabeled tree). An *unlabeled tree* is a set $D \subseteq \mathbb{N}^*$ that satisfies the following conditions.

- 1) The empty sequence is in D , i.e. $\langle \rangle \in D$.
- 2) The set D is prefix-closed, i.e. $\forall w \in \mathbb{N}^*. \forall i \in \mathbb{N}. wi \in D \Rightarrow w \in D$.
- 3) The set D is downward-closed, i.e. $\forall w \in \mathbb{N}^*. \forall i \in \mathbb{N}. \forall j \leq i. wi \in D \Rightarrow wj \in D$.

An unlabeled tree D is said to be *finite-depth* if it satisfies the following additional condition.

- 4) The set D has no strictly increasing sequence with respect to \preceq , i.e.

$$\forall v \in \mathbb{N}^\omega. |\{w \in D \mid w \preceq v\}| < \infty.$$

We write Tree (resp. Tree_{fin}) for the set of all unlabeled trees (resp. unlabeled finite-depth trees).

For an unlabeled tree D , we define a function $\text{branch}_D: D \rightarrow \mathbb{N} \cup \{\infty\}$ as follows.

$$\text{branch}_D(w) = \max\{i \mid wi \in D\}.$$

The *prefix order* over Tree is a partial order \preceq that is defined by

$$D_1 \preceq D_2 \stackrel{\text{def.}}{\iff} \left(D_1 \subseteq D_2, \text{ and } \forall w \in D_1. \text{branch}_{D_1}(w) \in \{0, \text{branch}_{D_2}(w)\} \right).$$

For a (possibly infinite) family of unlabeled trees $(D_i \in \text{Tree})_{0 \leq i < \alpha}$ where $\alpha \leq \omega$, we define a new tree $\text{combine}(D_1, D_2, \dots) \subseteq \mathbb{N}^*$ as follows.

$$\text{combine}(D_1, D_2, \dots) = \{\langle \rangle\} \cup \{iw \mid i < \alpha, w \in D_i\}$$

Remark A.28. We note that $D \in \text{Tree}_{\text{fin}}$ is assumed to be *finite-depth*, but it is not assumed to be *finitely-branching*. For example, we have $(\{\langle \rangle\} \cup \{\langle n \rangle \mid n \in \mathbb{N}\}) \in \text{Tree}_{\text{fin}}$ where $\langle n \rangle$ denotes a sequence whose length is 1.

Definition A.29 (ranked alphabet). A *ranked alphabet* is a pair $\Sigma = (\Sigma, \lfloor _ \rfloor)$ of a set Σ and an *arity function* $\lfloor _ \rfloor: \Sigma \rightarrow \mathbb{N} \cup \{\infty\}$. For each $a \in \Sigma$, $|a| \in \mathbb{N} \cup \{\infty\}$ is called the *arity* of a . For each $n \in \mathbb{N} \cup \{\infty\}$, we write Σ_n for $\{a \in \Sigma \mid |a| = n\} \subseteq \Sigma$.

Definition A.30 (labeled tree). Let Σ be a ranked alphabet. A Σ -*labeled tree* is a pair $T = (D, l)$ of an unlabeled tree $D \subseteq \text{Tree}$ and a *labeling function* $l: D \rightarrow \Sigma$ that respects the arity, i.e.

$$\forall w \in D. l(w) = \text{branch}_D(w).$$

The unlabeled tree D is called the *domain* of T . We write Tree_Σ for the set of all Σ -labeled trees.

Definition A.31 ($F_{t,\Sigma}$). Let Σ be a ranked alphabet. We define a functor $F_{t,\Sigma}: \mathbf{Sets} \rightarrow \mathbf{Sets}$ by

$$F_{t,\Sigma} = \left(\prod_{n \in \mathbb{N} \cup \{\omega\}} \Sigma_n \times (_)^n \right) \times \{0, 1\}.$$

Notation A.32. Let $c : X \rightarrow F_{t,\Sigma}X$ be an $F_{t,\Sigma}$ -coalgebra. For each $x \in X$ where $c(x) = (\xi, t)$, we write $c_1(x)$ and $c_2(x)$ for $\xi \in \coprod_{n \in \mathbb{N} \cup \{\omega\}} \Sigma_n \times (_)^n$ and $t \in \{0, 1\}$, respectively.

Definition A.33 (run tree). Let $c : X \rightarrow F_{t,\Sigma}X$ be an $F_{t,\Sigma}$ -coalgebra. We define a new ranked alphabet $X \times \Sigma = (X \times \Sigma, _ |)$ by $|(x, a)| = |a|$. For $x \in X$, an $(X \times \Sigma)$ -labeled tree (D, l) is called a *run tree of c from x* if it satisfies the following conditions (here for each $w \in D$ where $l(w) = (x', a)$, $l_1(w)$ and $l_2(w)$ denote x' and a respectively):

- 1) $l_1(\langle \rangle) = x$; and
- 2) for each $w \in D$,

$$c_1(l_1(w)) = (l_2(w), l_1(w0), l_1(w1), \dots).$$

A run tree (D, l) is called *accepting* if it has no infinite branch labeled only with non-accepting states, i.e.

$$\neg (\exists v \in \mathbb{N}^\omega. \forall w \in \mathbb{N}^* \text{ s.t. } w \preceq v. (w \in D \wedge c_2(l(w)) = 0)).$$

Proposition A.34. For an $F_{t,\Sigma}$ -coalgebra $c : X \rightarrow F_{t,\Sigma}X$ and $x \in X$, there exists a unique run tree of c from x . \square

Definition A.35 (RunTree_c and Reach_c). By Prop. A.34, we write $\text{RunTree}_c(x)$ for the unique run tree of c from x . For an $F_{t,\Sigma}$ -coalgebra $c : X \rightarrow F_{t,\Sigma}X$, we define a function $\text{Reach}_c : X \rightarrow \{0, 1\}$ by

$$\text{Reach}_c(x) = \begin{cases} 1 & (\text{RunTree}_c(x) \text{ is accepting}) \\ 0 & (\text{otherwise}). \end{cases}$$

Definition A.36. We define an $F_{t,\Sigma}$ -modality $\sigma_t : F_{t,\Sigma}\{0, 1\} \rightarrow \{0, 1\}$ over a truth-value domain $(\{0, 1\}, \leq)$ by

$$\sigma_t((a, p_1, p_2, \dots), t) = \begin{cases} 1 & (t = 1 \text{ or } \forall i. p_i = 1) \\ 0 & (\text{otherwise}). \end{cases}$$

Proposition A.37. The $F_{t,\Sigma}$ -modality $\sigma_t : F_{t,\Sigma}\{0, 1\} \rightarrow \{0, 1\}$ in Def. A.36 has least fixed points. For an $F_{t,\Sigma}$ -coalgebra $c : X \rightarrow F_{t,\Sigma}X$, the (coalgebraic) least fixed property $\llbracket \mu \sigma_t \rrbracket_c : X \rightarrow \{0, 1\}$ is given by the function $\text{Reach}_c : X \rightarrow \{0, 1\}$ in Def. A.35.

Proof. We define $f : X \rightarrow \{0, 1\}$ by $f = \text{Reach}_c$. It suffices to prove that f is the least fixed-point of $\Phi_{c,\sigma_t} : (\text{Tree}_{\text{fin}}^\perp)^X \rightarrow (\text{Tree}_{\text{fin}}^\perp)^X$.

We first show that f is a fixed point of Φ_{c,σ_t} . Let $x \in X$ and assume that $c(x) = ((a, x_1, x_2, \dots), t) \in F_{t,\Sigma}X$. Then we

have:

$$\begin{aligned} & \sigma_t \circ F_{t,\Sigma}f((a, x_1, x_2, \dots), t) \\ &= \begin{cases} 1 & (t = 1 \text{ or } \forall i. f(x_i) = 1) \\ 0 & (\text{otherwise}) \end{cases} \quad (\text{by def. of } \sigma_t) \\ &= \begin{cases} 1 & (t = 1 \text{ or } \forall i. \text{RunTree}_c(x_i) \text{ is accepting}) \\ 0 & (\text{otherwise}) \end{cases} \\ & \quad (\text{by def. of } f) \\ &= \begin{cases} 1 & (\text{RunTree}_c(x) \text{ is accepting}) \\ 0 & (\text{otherwise}) \end{cases} \\ & \quad (\text{by def. of } \text{RunTree}_c) \\ &= f(x). \end{aligned}$$

Hence f is a fixed point of Φ_{c,σ_t} .

It remains to show that f is the least fixed point. Let $f' : X \rightarrow \{0, 1\}$ be a fixed point of Φ_{c,σ_t} . It suffices to prove that for each $x \in X$, $f'(x) = 0$ implies $f(x) = 0$.

Let $x \in X$ and assume $f'(x) = 0$. Let $(D, l) = \text{RunTree}_c(x)$. For each $i \in \mathbb{N}$, we inductively define $x_i \in X$ and $v_i \in \mathbb{N}$ so that $f'(x_i) = 0$ for each i as follows.

- For $i = 0$, we let $x_0 = x$. Then we have $f'(x_0) = f(x) = 0$.
- Let $i \geq 0$ and assume that we have defined x_j and v_{j-1} for each $j \leq i$ so that $f'(x_i) = 0$. Let $c(x_i) = ((a_i, x_{i,1}, x_{i,2}, \dots), t_i)$. As f' is a fixed point of Φ_{c,σ_t} , we have:

$$0 = f'(x_i) = (\sigma_t \circ F_{t,\Sigma}f' \circ c)(x_i). \quad (15)$$

By the definition σ_t , this means that there exists k such that $f'(x_{i,k}) = 0$. We let $v_i = k$ and $x_{i+1} = x_{i,k}$.

We can prove that x_i and v_i satisfy the followings.

- 1) $\forall i. (v_0 \dots v_{i-1}) \in D$
- 2) $\forall i. l(v_0 \dots v_{i-1}) = x_i$
- 3) $\forall i. c_2(l(v_0 \dots v_{i-1})) = 0$

Here (1) and (2) are immediate from the definition. The equation (15) implies that $c_2(l(v_0 \dots v_{i-1})) = c_2(x_i) = t_i = 0$ for each $i \in \mathbb{N}$.

This means that $\text{RunTree}_c(x)$ is not accepting (see Def. A.33), and therefore we have $f(x) = 0$. This concludes the proof. \square

Proposition A.38. The F_t -modality $\sigma_t : F_t\{0, 1\} \rightarrow \{0, 1\}$ in Def. A.36 satisfies conditions in Asm. III.7. \square

Proposition A.39. Let Σ be a ranked alphabet and $\text{Tree}_{\text{fin}}^\perp = \text{Tree}_{\text{fin}} \cup \{\perp_t\}$. We define an $F_{t,\Sigma}$ -algebra $r_t : F_{t,\Sigma}(\text{Tree}_{\text{fin}}^\perp) \rightarrow \text{Tree}_{\text{fin}}^\perp$, a function $q_t : \text{Tree}_{\text{fin}}^\perp \rightarrow \{0, 1\}$ and a partial order $\sqsubseteq_{\text{Tree}_{\text{fin}}^\perp}$ over $\text{Tree}_{\text{fin}}^\perp$ as follows.

- 1) $r_t((a, D_1, D_2, \dots), t) = \begin{cases} \langle \rangle & (t = 1) \\ \text{combine}(D_1, D_2, \dots) & (t = 0 \text{ and } \forall i. D_i \neq \perp_t) \\ \perp_t & (t = 0 \text{ and } \exists i. D_i = \perp_t) \end{cases}$

- 2) $q_b(D) = \begin{cases} 1 & (D \neq \perp_t) \\ 0 & (D = \perp_t) \end{cases}$.
- 3) $D_1 \sqsubseteq_{\text{Tree}_{\text{fin}}^{\perp}} D_2 \stackrel{\text{def.}}{\iff} D_1 = \perp_t$ or $(D_1, D_2 \in \text{Tree}_{\text{fin}} \text{ and } D_1 \succeq D_2)$ (note the direction).

Then $(r_t, q_t, \sqsubseteq_{\text{Tree}_{\text{fin}}^{\perp}})$ is a ranking domain.

Lemma A.40. *The poset $(\text{Tree}_{\text{fin}}^{\perp}, \sqsubseteq_{\text{Tree}_{\text{fin}}^{\perp}})$ is a complete lattice.*

Proof. Let $K \subseteq \text{Tree}_{\text{fin}}^{\perp}$.

K has the least upper bound. We first prove that K has the least upper bound. If $K \cap \text{Tree}_{\text{fin}} = \emptyset$, then \perp_t is the least upper bound of K .

Assume $K \cap \text{Tree}_{\text{fin}} \neq \emptyset$. If $K' = K \setminus \{\perp_t\}$ has the least upper bound, then it is also the least upper bound of K .

We define $A \subseteq \mathbb{N}^*$ as follows.

$$A = \bigcup \{D' \in \text{Tree}_{\text{fin}} \mid \forall D \in K'. D' \preceq D\}.$$

As $\{\langle \rangle\} \preceq D$ holds for each $D \in K'$ and Cond. 1–3 in Def. A.27 are preserved by the union, $A \in \text{Tree}$. We prove that A is the least upper bound of K' .

We first prove that A is an upper bound. Let $D \in K'$. We prove $D \sqsubseteq_{\text{Tree}_{\text{fin}}^{\perp}} A$. To this end, by the definition of \preceq , it suffices to prove $A \subseteq D$ and $\text{branch}_A(w) \in \{0, \text{branch}_D(w)\}$ for each $w \in A$. The former is immediate from the definition of A . The latter is satisfied because we have $\text{branch}_{D'}(w) \in \{0, \text{branch}_D(w)\}$ for each D' such that $D' \preceq D$ and $w \in D'$.

Hence A is an upper bound of K . We note that this also proves that A is finite-depth.

It is immediate from the definition of A that A is the least upper bound.

K has the greatest lower bound. We prove that K has the greatest lower bound. If $\perp_t \in K$, then $\prod K = \perp_t$. If $K = \emptyset$, then $\prod K = \{\langle \rangle\}$.

We assume $\perp_t \notin K$ and $K \neq \emptyset$. For $D_1, D_2 \in K$, if there exists $D_3 \in K$ such that $D_1 \preceq D_3$ and $D_2 \preceq D_3$ then by the definition of \preceq , we have:

$$\forall w \in D_1 \cap D_2. (\text{branch}_{D_1}(w) \neq 0 \text{ and } \text{branch}_{D_2}(w) \neq 0) \\ \Rightarrow \text{branch}_{D_1}(w) = \text{branch}_{D_2}(w).$$

Therefore if there exists $D_1, D_2 \in K$ that satisfy

$$\exists w \in D_1 \cap D_2. \text{branch}_{D_1}(w) \neq 0, \text{branch}_{D_2}(w) \neq 0 \\ \text{and } \text{branch}_{D_1}(w) \neq \text{branch}_{D_2}(w), \quad (16)$$

then there does not exist $D_3 \in K$ such that $D_1 \preceq D_3$ and $D_2 \preceq D_3$. This implies $\prod K = \perp_t$.

Assume that there does not exist $D_1, D_2 \in K$ that satisfy (16). We define $B \in \mathbb{N}^*$ by

$$B = \bigcup K.$$

It is easy to see that $B \in \text{Tree}$. It is also easy to see that if B is not finite-depth then \perp_t is the greatest lower bound of K .

Assume $B \in \text{Tree}_{\text{fin}}$. We prove that B is the greatest lower bound of K . Let $D \in K$. By the definition of B , we have $D \subseteq B$. As there exist no $D_1, D_2 \in K$ that satisfy (16), we have $\text{branch}_D(w) \in \{0, \text{branch}_B(w)\}$ for each $w \in D$. Hence we have $B \sqsubseteq_{\text{Tree}_{\text{fin}}^{\perp}} D$, and therefore B is a lower bound of K . It is immediate from its definition that B is the greatest lower bound. \square

Lemma A.41. *The triple $(r_t, q_t, \sqsubseteq_{\text{Tree}_{\text{fin}}^{\perp}})$ in Prop. A.39 satisfies Cond. 2 in Def. III.10.*

Proof. Cond. 2a is immediate from Lem. A.40.

It is easy to see that Cond. 2b is satisfied. \square

Lemma A.42. *The algebra $r_t : F_{t, \Sigma} \text{Tree}_{\text{fin}}^{\perp} \rightarrow \text{Tree}_{\text{fin}}^{\perp}$ in Prop. A.39 is a corecursive algebra.*

Proof. It suffices to show that the function $\Phi_{c, r_t} : (\text{Tree}_{\text{fin}}^{\perp})^X \rightarrow (\text{Tree}_{\text{fin}}^{\perp})^X$ has a unique fixed point.

By Lem. A.41, the set $((\text{Tree}_{\text{fin}}^{\perp})^X, \sqsubseteq_{\text{Tree}_{\text{fin}}^{\perp}})$ where $\sqsubseteq_{\text{Tree}_{\text{fin}}^{\perp}}$ denotes the pointwise extension is a complete lattice. Therefore by Thm. II.18, the function Φ_{c, r_t} , which is monotone by Lem. A.41, has the least fixed point.

It remains to show that this is the unique fixed point. Let $f, f' : X \rightarrow \text{Tree}_{\text{fin}}^{\perp}$ be fixed points of Φ_{c, r_t} .

Let $g : X \rightarrow \text{Tree}_{\text{fin}}^{\perp}$ and assume $g(x) = \perp_t$. Let $c(x) = ((a, x_0, x_1, \dots), t)$. As g is a fixed point of Φ_{c, r_t} , we have $(r_t \circ F_{t, \Sigma} g \circ c)(x) = \perp_t$. Hence we have $t = 0$ and that there exists i such that $g(x_i) = \perp_t$.

Therefore we can inductively define a sequence $x^0 x^1 x^2 \dots \in X^\omega$ so that:

- 1) $g(x^i) = \perp_t$;
- 2) $c_2(x^i) = 0$ for each $i \in \omega$; and
- 3) $c_1(x^i) = (a^i, x_1^i, x_2^i, \dots)$ then $x^{i+1} = x_n^i$ for some n .

Conversely, we can prove that existence of a sequence $x^0 x^1 x^2 \dots \in X^\omega$ that satisfies the conditions above implies $g(x) = \perp_t$.

Therefore we have: $f(x) = \perp_t$ if and only if $f'(x) = \perp_t$.

We now prove that for each $n \in \mathbb{N}$, $w \in \mathbb{N}^n$ and $x \in X$ such that $f(x) \neq \perp_t$ and $f'(x) \neq \perp_t$, $w \in f(x)$ if and only if $w \in f'(x)$ by the induction on n .

For $n = 0$, by Def. A.27, we have $\langle \rangle \in f(x)$ and $\langle \rangle \in f'(x)$ for each x .

Let $n > 0$ and assume that for each $w \in \mathbb{N}^n$ and $x \in X$ such that $f(x) \neq \perp_t$ and $f'(x) \neq \perp_t$, we have $w \in f(x)$ if and only if $w \in f'(x)$. Let $x \in X$ and assume $c(x) = ((a, x_0, x_1, \dots), t) \in (\Sigma \times (_)^m) \times \{0, 1\}$. For each $i \in \mathbb{N}$, we have:

$$\begin{aligned} iw \in f(x) & \\ \iff iw \in \Phi_{c, r_t}(f) & \quad (f \text{ is a fixed point}) \\ \iff i < m \text{ and } w \in f(x_i) & \quad (\text{by def. of } r_t) \\ \iff i < m \text{ and } w \in f'(x_i) & \quad (\text{by IH}) \\ \iff iw \in \Phi_{c, r_t}(f') & \quad (\text{by def. of } r_t) \\ \iff iw \in f'(x) & \quad (f' \text{ is a fixed point}). \end{aligned}$$

Hence we have $f(x) = f'(x)$ for each $x \in X$. This concludes the proof. \square

Proof (Prop. A.39). We prove that $(r_t, q_t, \sqsubseteq_{\text{Tree}_{\text{fin}}^\perp})$ satisfies conditions in Def. III.10.

1: Let $((a, D_0, D_1, \dots), t) \in F_{t, \Sigma} \text{Tree}_{\text{fin}}^\perp$. Then we have:

$$\begin{aligned}
& (q_t \circ r_t)((a, D_0, D_1, \dots), t) = 1 \\
& \Leftrightarrow r_t((a, D_0, D_1, \dots), t) \neq \perp_t && \text{(by def. of } q_t) \\
& \Leftrightarrow t = 1 \text{ or } \forall i. D_i \neq \perp_t && \text{(by def. of } r_t) \\
& \Leftrightarrow t = 1 \text{ or } \forall i. q_t(D_i) = 1 && \text{(by def. of } q_t) \\
& \Leftrightarrow (\sigma_t \circ F_{t, \Sigma} q_t)((a, D_0, D_1, \dots), t) = 1 && \text{(by def. of } \sigma_t).
\end{aligned}$$

Hence we have $q_t \circ r_t = \sigma \circ F_{t, \Sigma} q_t$.

2: We have already proved in Lem. A.41.

3: Monotonicity and strictness are immediate from the definition of q_t . We prove that q_t is continuous. Let $K \subseteq \text{Tree}_{\text{fin}}^\perp$. Then we have:

$$\begin{aligned}
& \bigsqcup_{D \in K} q_t(D) = 0 \\
& \Leftrightarrow \forall D \in K. q_t(D) = 0 \\
& \Leftrightarrow K = \{\perp_t\} && \text{(by def. of } q_t) \\
& \Leftrightarrow \bigsqcup K = \perp_t \\
& \Leftrightarrow q_t(\bigsqcup K) = 0 && \text{(by def. of } q_t).
\end{aligned}$$

Hence q_t is continuous.

4: We have already proved in Lem. A.42. \square

Proposition A.43 (completeness). *For an arbitrary $F_{t, \Sigma}$ -coalgebra $c : X \rightarrow F_{t, \Sigma} X$, there exists a ranking arrow $b : X \rightarrow \text{Tree}_{\text{fin}}^\perp$ with respect to the ranking domain $(r_t, q_t, \sqsubseteq_{\text{Tree}_{\text{fin}}^\perp})$ in Prop. A.39 that satisfies the following.*

$$q_t \circ b = \llbracket \mu \sigma_t \rrbracket_c$$

Proof. Immediate from $q_t \circ r_t = \sigma_t \circ F_{t, \Sigma} q_t$ (see the proof of Prop. A.39) and Prop. III.18. \square