



4.2	Quantum gates – Braiding anyons	20
4.3	Measurements – Fusing anyons	22
4.4	Possible error sources	22
<b>5</b>	<b>Topological quantum computation with superconducting nanowires</b>	<b>24</b>
5.1	Majorana zero modes in a superconducting nanowire	25
5.2	The Majorana qubit	28
5.3	Manipulating and reading out the Majorana qubit	28
5.4	Challenges with Majorana-based topological quantum computation	30
<b>6</b>	<b>Outlook</b>	<b>31</b>
	<b>References</b>	<b>32</b>

---

## 1 Introduction

Topological quantum computation is an approach to storing and manipulating quantum information that employs exotic quasiparticles, called anyons. Anyons are interesting on their own right in fundamental physics, as they generalise the statistics of the commonly known bosons and fermions. Due to this exotic statistical behaviour, they exhibit non-trivial quantum evolutions that are described by topology, i.e. they are abstracted from local geometrical details. When anyons are used to encode and process quantum information, this topological behaviour provides a much desired resilience against control errors and perturbations. To be more precise, the presence of certain kinds of anyons gives rise to a degenerate decoherence-free subspace, in which the state can only be evolved by moving the anyons adiabatically around each other. While from the first sight anyons appear to be an over-complicated method for performing quantum computation, they are profoundly linked to quantum error correction [1], the algorithmic means we have in dealing with errors during quantum computation. In a sense, anyonic quantum computers implement quantum error-correction at the hardware level, thus becoming resilient to control errors and erroneous perturbations. This has augmented topological quantum computation from a niche field of research to a methodology that permeates much of the research efforts in realising fault-tolerant quantum computation.

In this review we present a non-technical introduction to anyons and to the framework for performing fault-tolerant quantum computation with them. The emphasis will be on the key properties that define anyons and how they enable protected encoding and processing of quantum information. As anyons can emerge in numerous microscopically distinct systems, we discuss these concepts primarily at a platform-independent level of anyon models and provide an extensive list of references for the interested reader to go deeper. Our aim is to provide a clear and concise introduction to the underlying principles of topological quantum computation without expertise in condensed matter theory. When condensed matter concepts are needed, we introduce them in a heuristic level to give the reader a general understanding without referring to the mathematical details. In doing so, we aim this review to be accessible to anyone with a solid undergraduate understanding of quantum mechanics and the basics of quantum information.

While several reviews have already been written on the topic, we aim this review to serve as an



















































































- [178] M. Kjaergaard, F. Nichele, H. J. Suominen, M. P. Nowak, M. Wimmer, A. R. Akhmerov, J. A. Folk, K. Flensberg, J. Shabani, C. J. Palmström and C. M. Marcus, *Quantized conductance doubling and hard gap in a two-dimensional semiconductor-superconductor heterostructure*, Nat. Commun. **7**, 12841 (2016), doi:10.1038/ncomms12841.
- [179] N. H. Lindner, E. Berg, G. Refael and A. Stern, *Fractionalizing Majorana Fermions: Non-Abelian Statistics on the Edges of Abelian Quantum Hall States*, Phys. Rev. X **2**, 041002 (2012), doi:10.1103/PhysRevX.2.041002.
- [180] D. J. Clarke, J. Alicea and K. Shtengel, *Exotic non-Abelian anyons from conventional fractional quantum Hall states*, Nat. Comm. **4**, 1348 (2013), doi:10.1038/ncomms2340.
- [181] M. Barkeshli and J. D. Sau, *Physical Architecture for a Universal Topological Quantum Computer based on a Network of Majorana Nanowires*, <https://arxiv.org/abs/1509.07135>.
- [182] S. Plugge, L. A. Landau, E. Sela, A. Altland, K. Flensberg and R. Egger, *Roadmap to Majorana surface codes*, Phys. Rev. B **94**, 174514 (2016), doi:10.1103/PhysRevB.94.174514.
- [183] S. Hoffman, C. Schrade, J. Klinovaja and D. Loss, *Universal quantum computation with hybrid spin-Majorana qubits*, Phys. Rev. B **94**, 045316 (2016), doi:10.1103/PhysRevB.94.045316.