# A NOTE ON THE ENUMERATION OF EUCLIDEAN SELF-DUAL SKEW-CYCLIC CODES OVER FINITE FIELDS

IRWANSYAH, INTAN MUCHTADI-ALAMSYAH, AHMAD MUCHLIS, ALEAMS BARRA, AND DJOKO SUPRIJANTO

ABSTRACT. In this paper, we give the enumeration formulas for Euclidean selfdual skew-cyclic codes over finite fields when  $(n, |\theta|) = 1$  and for some cases when  $(n, |\theta|) > 1$ , where n is the length of the code and  $|\theta|$  is the order of automorphism  $\theta$ .

#### 1. INTRODUCTION

Let  $\mathbb{F}_q$  be the finite field of cardinality q, where q is a prime power. A linear code over  $\mathbb{F}_q$  with length n is a linear subspace of the  $\mathbb{F}_q$ -vector space  $\mathbb{F}_q^n$ . The two classes of interesting linear codes are cyclic codes and self-dual codes. The reason is, among other things, because these classes of codes have produced many examples of optimal codes. Therefore, some authors also studied the combination of these two classes of codes, namely self-dual cyclic codes [7] and the generalization of cyclic codes such as quasi-cyclic codes and skew-cyclic codes[8].

Skew-cyclic codes or  $\theta$ -cyclic codes over finite fields are a generalization of cyclic codes over finite fields. Here,  $\theta$  is an automorphism in the corresponding finite fields. This class of codes has many interesting properties such as it can be viewed as a left ideal or left submodule over a skew-polynomial ring, *etc*, see [3]. Moreover, these codes also give [38, 18, 11] code, an Euclidean self-dual code over  $\mathbb{F}_4$ , which improves the bound for optimal self-dual codes of length 36 with respect to Hamming distance [4].

Recently, the study of special class of skew-codes over finite fields *i.e.* Euclidean self-dual skew-codes are conducted regarding their existence and enumeration. Boucher [2] showed that, when  $q \equiv 1 \mod 4$ , there always exists a self-dual  $\theta$ -code in any dimension and that self-dual  $\theta$ -codes of a given dimension are either all  $\theta$ -cyclic or all  $\theta$ -negacyclic. Also, It is not exists when  $q \equiv 3 \mod 4$ . However, the enumeration of this class of codes not yet been done completely. So far, the enumeration was done for self-dual skew-codes over  $\mathbb{F}_4$  with length  $n = 2^s$  [5] and for self-dual skew-codes over  $\mathbb{F}_{p^2}$  [1].

In this paper, we study the enumeration of self-dual skew-cyclic or  $\theta$ -cyclic codes when  $(n, |\theta|) = 1$  and for some cases when  $(n, |\theta|) = d > 1$ , where  $|\theta|$  is the order of the automorphism  $\theta$ .

### 2. The Enumeration Formulas

Let us recall some definition which can be found in [1, 3, 7, 8]. Let T be a shift operator on  $\mathbb{F}_q^n$  and  $\theta$  is an automorphism in  $\mathbb{F}_q$ .

**Definition 1.** Let C be a linear code of length n over  $\mathbb{F}_q$ .

<sup>1991</sup> Mathematics Subject Classification. 11T71.

Key words and phrases. Skew-cyclic codes, Euclidean self-dual codes, Enumeration.

(1) If

$$T^{l}(c) = (c_{n-l}, c_{n-l+1}, \dots, c_{n-l-1}) \in C$$

whenever  $c = (c_0, c_1, \ldots, c_{n-1}) \in C$ , then C is a quasi-cyclic code with index l.

(2) If

$$T_{\theta}(c) = (\theta(c_{n-1}), \theta(c_0), \dots, \theta(c_{n-2})) \in C$$

whenever  $c = (c_0, c_1, \dots, c_{n-1}) \in C$ , then C is a  $\theta$ -cyclic code.

(3) If  $\theta$  in (2) above is an identity map, then C is a cyclic code.

For any  $c = (c_0, \ldots, c_{n-1})$  and  $c' = (c'_0, \ldots, c'_{n-1})$  in  $\mathbb{F}_q$ , we define the Euclidean product between c and c' as

$$[c, c'] = \sum_{i=0}^{n-1} c_i c'_i.$$

Let C be a code of length n over  $\mathbb{F}_q$  and  $C^{\perp} = \{c' \in \mathbb{F}_q^n : [c, c'] = 0, \forall c \in C\}$ . Then we can give the following definition.

**Definition 2.** A linear code C is called *Euclidean self-dual* if  $C = C^{\perp}$ .

Now, before we do the enumeration, we need to proof the following proposition.

**Proposition 3.** If C is a  $\theta$ -cyclic code over  $\mathbb{F}_q$  of length n, for some automorphism  $\theta$  in  $\mathbb{F}_q$ , then C is either a cyclic code or a quasi-cyclic code.

*Proof.* We consider two cases for  $|\theta|$ . First, if  $gcd(|\theta|, n) = 1$ , then there exist  $p, q \in \mathbb{Z}$ , such that pr + qn = 1 or  $p|\theta| = 1 + bn$  for some b. For any  $c \in C$ , let  $c = (c_0, c_1, \ldots, c_{n-1})$ , for some  $c_0, c_1, \ldots, c_{n-1} \in \mathbb{F}_q$ . Then we have

$$T_{\theta}^{p|\theta|}(c) = \left(\theta^{p|\theta|}(c_{n-p|\theta| \pmod{n}}), \theta^{p|\theta|}(c_{n-p|\theta|+1 \pmod{n}}), \dots, \theta^{p|\theta|}(c_{n-p|\theta|-1 \pmod{n}})\right)$$
$$= \left(\theta^{p|\theta|}(c_{n-1}), \theta^{p|\theta|}(c_{0}), \dots, \theta^{p|\theta|}(c_{n-2})\right)$$
$$= (c_{n-1}, c_{0}, \dots, c_{n-2})$$
$$= T(c),$$

which means,  $T(c) \in C$  or C is a cyclic code.

Second, if  $gcd(|\theta|, n) = s$ , for some  $s \in \mathbb{N}$ , where  $s \neq 1$ , then there exist  $p_1, p_2 \in \mathbb{Z}$ , such that  $p_1|\theta| = s + p_2n$ . Hence, we have

$$T_{\theta}^{p_{1}|\theta|}(c) = \left(\theta^{p_{1}|\theta|}(c_{n-p_{1}|\theta| \pmod{n}}), \theta^{p_{1}|\theta|}(c_{n-p_{1}|\theta|+1 \pmod{n}}), \dots, \theta^{p_{1}|\theta|}(c_{n-p_{1}|\theta|-1 \pmod{n}})\right)$$
$$= \left(\theta^{p_{1}|\theta|}(c_{n-s}), \theta^{p_{1}|\theta|}(c_{n-s+1}), \dots, \theta^{p_{1}|\theta|}(c_{n-s-1})\right)$$
$$= (c_{n-s}, c_{n-s+1}, \dots, c_{n-s-1})$$
$$= T^{s}(c)$$

which means, C is a quasi-cyclic code of index s, as we hope.

2.1. Case 1 :  $(n, |\theta|) = 1$ . If  $(n, |\theta|) = 1$ , then  $\theta$ -cyclic code is a cyclic code. It means, the number of self-dual  $\theta$ -cyclic codes of length n is less than or equal to the number of self-dual cyclic codes of length n. Recall that, based on the result in [7], Euclidean self-dual cyclic code of length n over  $\mathbb{F}_q$  exist only when q is a power of 2 and n is an even integer. Now, let  $n = 2^{v(n)}\tilde{n}$ , where  $(\tilde{n}, 2) = 1$ , and let  $\phi$  be the Euler function and  $ord_j(i)$  be the smallest integer e such that j divides  $i^e - 1$ . Also, let j be an odd positive integer and m be a positive integer. We say that the pair (j,m) is good if j divides  $(2^m)^k + 1$  for some integer  $k \geq 0$  and bad otherwise. We define the function  $\chi$  as follows.

$$\chi(j,m) = \begin{cases} 0, & \text{if } (j,m) \text{ good,} \\ 1, & \text{otherwise.} \end{cases}$$

Recall that, based on [7], the number of Euclidean self-dual cyclic codes over  $\mathbb{F}_q$  is

(2.1) 
$$(1+2^{v(n)})^{\frac{1}{2}\sum_{j\mid\tilde{n}}\chi(j,m)\phi(j)/ord_{j}(2^{m})}$$

Note that, by Proposition 3, when  $(n, |\theta|) = 1$ , then the number of Euclidean self-dual  $\theta$ -cyclic codes over  $\mathbb{F}_q$  is less than or equal to the one provided by (2.1).

Now, we make partitions of the set  $\{0, 1, \ldots, \tilde{n}\}$  into  $2^m$ -cyclotomic cosets denoted by  $C_s$ , where s is the smallest element in  $C_s$ . Let  $\mathcal{A}$  be the collection of union of  $2^m$ -cyclotomic cosets which represent the generators of the self-dual cyclic codes over  $\mathbb{F}_q$  of length n. Also, let  $\theta(\beta) = \beta^{2^r}$ , for all  $\beta \in \mathbb{F}_{2^m}$  and for some  $r \in \{1, 2, \ldots, m\}$ . Also, let  $\lambda$  be the map as follows

$$\begin{array}{rccc} \lambda_r: & \{0, 1, \dots, \tilde{n}\} & \longrightarrow & \{0, 1, \dots, \tilde{n}\} \\ & a & \longmapsto & 2^r a \pmod{\tilde{n}}. \end{array}$$

Furthermore, we let  $\Lambda_r$  be the map on  $\mathcal{A}$  induced by  $\lambda_r$  and  $\overline{\Lambda}_r$  be the number of elements A in  $\mathcal{A}$  such that  $\Lambda_r(A) \neq A$ . Then, we have the following result.

**Theorem 4.** If  $(n, |\theta|) = 1$ ,  $q = 2^m$ , for some m, and  $n = 2^{\nu(n)}\tilde{n}$ , for some odd positive integer  $\tilde{n}$ , then the number of Euclidean self-dual  $\theta$ -cyclic codes of length n over  $\mathbb{F}_q$  is

$$\left(1+2^{v(n)}\right)^{\frac{1}{2}\sum_{j\mid\bar{n}}\chi(j,m)\phi(j)/ord_j(2^m)}-\overline{\Lambda}_r.$$

*Proof.* Let  $C = \langle g(x) \rangle$ , where g(x) is a monic polynomial with minimum degree in C and, therefore, g(x) is a divisor of  $x^{\tilde{n}} - 1$ . Now, for any  $f(x) = \sum_{i=0}^{s} f_i x^i \in \mathbb{F}_{2^m}[x]$ , let

$$\theta(f) = \sum_{i=0}^{s} \theta(f_i) x^i = \sum_{i=0}^{s} f_i^{2^r} x^i.$$

Also, let  $\theta(C) = \langle \theta(g) \rangle$ . It is easy to see that  $\theta(C)$  is also a Euclidean self-dual cyclic code. Now, let  $\alpha$  be a primitive  $\tilde{n}$ -th root of unity in some extension of  $\mathbb{F}_{2^m}$ . Then, by [6, Theorem 4.2.1(vii)], we have

$$g(x) = \prod_{s} M_{\alpha^s}(x),$$

where  $M_{\alpha^s}(x)$  is the minimal polynomial of  $\alpha^s$  over  $\mathbb{F}_{2^m}$ , and s is the representatives of the  $2^m$ -cyclotomic cosets modulo  $\tilde{n}$ . Now, let

$$x^{n} - 1 = f_{1}(x) \cdots f_{s}(x)h_{1}(x)h_{1}^{*}(x) \cdots h_{t}(x)h_{t}^{*}(x),$$

where  $f_i(x)$   $(1 \leq i \leq s)$  are monic irreducible self-reciprocal polynomials over  $\mathbb{F}_{2^m}$ , while  $h_j(x)$  and its reciprocal polynomial  $h_j^*(x)$   $(1 \leq j \leq t)$  are both monic irreducible polynomials over  $\mathbb{F}_{2^m}$ . Then, by [6, Theorem 3.7.6], we have

$$f_i(x) = \prod_{k \in C_{s_i}} (x - \alpha^k),$$

and

$$h_j(x) = \prod_{k \in C_{s_j}} (x - \alpha^k)$$

Therefore, in order to make  $\theta(C) \subseteq C$ , we have to choose  $f_i(x)$ 's and  $h_j(x)$ 's such that their representatives in  $\mathcal{A}$  are fixed by  $\Lambda_r$ , because  $\theta(x-\alpha^l) = x-\alpha^{2^rl}$ . Since  $\overline{\Lambda_r}$  is the number of non-fixed-by- $\Lambda_r$ -elements in  $\mathcal{A}$ , we have the desired formula.  $\Box$ 

Let us consider the following examples.

**Example 5.** Let q = 4, r = 1, and n = 6. Therefore,  $\tilde{n} = 3$ . Also, let  $\mathbb{F}_4 = \mathbb{F}_2[\alpha]$ , where  $\alpha$  is the root of the polynomial  $x^2 + x + 1$ . Note that,  $\alpha$  is also a primitive  $3^{\text{th}}$ -root of unity. As we can check, we have three 4-cyclotomic cosets modulo 3, *i.e.*  $C_0 = \{0\}, C_1 = \{1\}, \text{ and } C_2 = \{2\}$ . Moreover,  $C_0$  represents a self-reciprocal polynomial, while  $C_1$  and  $C_2$  represent two polynomials which reciprocal to each other. Since  $n = 2\tilde{n}$ , we have

$$\mathcal{A} = \{ C_0 \cup C_1 \cup C_1, C_0 \cup C_2 \cup C_2, C_0 \cup C_1 \cup C_2 \}.$$

We can see that  $\Lambda_1(C_0) = C_0, \Lambda_1(C_1) = C_2$ , and  $\Lambda_1(C_2) = C_1$ . Therefore, we have  $\overline{\Lambda}_1 = 2$ , and the number of self-dual  $\theta$ -cyclic codes is  $|\mathcal{A}| - \overline{\Lambda}_1 = 1$ .

**Example 6.** Let q = 4, r = 1, and n = 14. Therefore,  $\tilde{n} = 7$ . Let  $\alpha \in \mathbb{F}_{4^2}$  be a primitive 7<sup>th</sup>-root of unity. We can check that we have three 4-cyclotomic cosets modulo 7. They are  $C_0 = \{0\}, C_1 = \{1, 2, 4\}$ , and  $C_3 = \{3, 5, 6\}$ . Also,  $C_0$  represents a self-reciprocal polynomial, while  $C_1$  and  $C_3$  represent two polynomials which reciprocal to each other. So, we have that

$$\mathcal{A} = \{ C_0 \cup C_1 \cup C_1, C_0 \cup C_3 \cup C_3, C_0 \cup C_1 \cup C_3 \}.$$

We also have  $\overline{\Lambda}_1 = 0$ , because  $\Lambda_1(C_i) = C_i$  for all i = 0, 1, 3. Therefore, the number of self-dual  $\theta$ -cyclic codes is  $|\mathcal{A}| = 3$ .

We have to note that  $|\mathcal{A}|$  is the number of self-dual cyclic codes of length n over  $\mathbb{F}_q$ . Therefore, the two examples above give us another way to count the number of self-dual cyclic codes using q-cyclotomic cosets of modulo  $\tilde{n}$ .

2.2. Case 2:  $(n, |\theta|) = d > 1$ . If  $(n, |\theta|) = d > 1$ , then  $\theta$ -cyclic code is a quasicyclic code of index d as stated in Proposition 3. Now, for any  $C \subseteq \mathbb{F}_q^n$ , define  $T_{\theta}(C) = \{T_{\theta}(c) : \forall c \in C\}$ . Then, we can easily prove the following lemma.

**Lemma 7.** If C is a Euclidean self-dual quasi-cyclic code of index d, then  $T_{\theta}(C)$  is also an Euclidean self-dual quasi-cyclic code of index d. Moreover,  $T_{\theta}(C)$  has the same Hamming weight enumerator polynomial as C.

The Lemma 7 above shows that the map  $T_{\theta}$  preserves Euclidean self-duality and quasi-cyclic property. Moreover, we have to note that, if C is a quasi-cyclic code such that  $T_{\theta}(C) \subseteq C$ , then C is also a  $\theta$ -cyclic code.

For any code  $C \subseteq \mathbb{F}_q^n$ , let  $\rho$  be the map as follows,

(2.2) 
$$\rho(C) = \begin{cases} 0, & \text{if} T_{\theta}(C) \subseteq C, \\ 1, & \text{otherwise.} \end{cases}$$

Let us recall structures of quasi-cyclic code described in [8]. Let n = dm,  $R = \mathbb{F}_q[Y]/(Y^m - 1)$ , and C be a quasi-cyclic code over  $\mathbb{F}_q$  of length dm with index d. Let

 $\mathbf{c} = (c_{00}, c_{01}, \dots, c_{0,l-1}, c_{10}, \dots, c_{1,l-1}, \dots, c_{m-1,0}, \dots, c_{m-1,l-1}) \in C.$ 

Then, the map  $\phi: \mathbb{F}_q^{dm} \longrightarrow R^d$  defined by

$$\phi(\mathbf{c}) = (\mathbf{c}_0(Y), \mathbf{c}_1(Y), \dots, \mathbf{c}_{l-1}(Y)) \in \mathbb{R}^d,$$

where  $\mathbf{c}_j(Y) = \sum_{i=0}^{m-1} c_{ij} Y^i \in R$ , is a one-to-one correspondence between quasicyclic codes over  $\mathbb{F}_q$  of index d and length dm and linear codes over R of length d. Moreover,

$$R^{d} = \left(\bigoplus_{i=1}^{s} G_{i}^{d}\right) \oplus \left(\bigoplus_{j=1}^{t} \left(H_{j}^{\prime d} \oplus H_{j}^{\prime \prime d}\right)\right),$$

where  $G_i = \mathbb{F}_q[Y]/(g_i), H'_j = \mathbb{F}_q[Y]/(h_j)$ , and  $H''_j = \mathbb{F}_q[Y]/(h^*_j)$ , for some selfreciprocal irreducible polynomials  $g_i$   $(1 \le i \le s)$  and irreducible reciprocal pairs  $h_j$ and  $h^*_j$   $(1 \le j \le t)$  which satisfy

$$Y^m - 1 = \delta g_1 \cdots g_s h_1 h_1^* \cdots h_t h_t^*$$

The above decomposition gives

$$C = \left(\bigoplus_{i=1}^{s} C_{i}\right) \oplus \left(\bigoplus_{j=1}^{t} \left(C_{j}^{\prime} \oplus C_{j}^{\prime\prime}\right)\right),$$

where  $C_i$  is a linear code over  $G_i$  of length d,  $C'_j$  is a linear code over  $H'_j$  of length d, and  $C''_j$  is a linear code over  $H''_j$  of length d.

Now, let

$$\rho_{G_i} = \sum_{C \text{ code over } G_i} \rho(\phi^{-1}(C)),$$

$$\rho_{H'_j,H''_j} = \sum_{\substack{C' \text{ code over } H'_j \\ C'' \text{ code over } H''_j}} \rho(\phi^{-1}(C' \oplus C'')),$$

and

$$N(d,q) = 1 + \sum_{k=1}^{d} \frac{(q^d - 1)(q^d - q) \cdots (q^d - q^{k-1})}{(q^k - 1)(q^k - q) \cdots (q^k - q^{k-1})}.$$

Then, we have the following result for d = 2.

**Proposition 8.** Let  $\theta$  be an automorphism in  $\mathbb{F}_q$  such that  $(n, |\theta|) = 2$ , where q be a prime power satisfying one of the following conditions,

- (1) q is a power of 2,
- (2)  $q = p^b$ , where p is a prime congruent to 1 mod 4, or
- (3)  $q = p^2 b$ , where p is a prime congruent to 3 mod 4.

Also, let m be an integer relatively prime to q. Suppose that  $Y^m - 1 = \delta g_1 \cdots g_s h_1 h_1^* \cdots h_t h_t^*$ as mentioned above. Suppose further that  $g_1 = Y - 1$  and, if m is even,  $g_2 = Y + 1$ . Let the degree of  $g_i$  be  $2d_i$  and the degree of  $h_j$  (also  $h_j^*$ ) be  $e_j$ . Then, the number of distinct Euclidean self-dual  $\theta$ -cyclic codes of length 2m with index 2 over  $\mathbb{F}_q$  is

$$\begin{aligned} &4\prod_{i=3}^{s}(q^{d_{i}}+1-\rho_{G_{i}})\prod_{j=1}^{t}(N(2,q^{e_{j}})-\rho_{H'_{j},H''_{j}}), & \text{if } m \text{ is even and } q \text{ is odd} \\ &2\prod_{i=2}^{s}(q^{d_{i}}+1-\rho_{G_{i}})\prod_{j=1}^{t}(N(2,q^{e_{j}})-\rho_{H'_{j},H''_{j}}), & \text{if } m \text{ is odd and } q \text{ is odd} \\ &\prod_{i=2}^{s}(q^{d_{i}}+1-\rho_{G_{i}})\prod_{j=1}^{t}(N(2,q^{e_{j}})-\rho_{H'_{j},H''_{j}}), & \text{if } m \text{ is odd and } q \text{ is even} \end{aligned}$$

*Proof.* Apply [8, Proposition 6.2] and use the fact that the map  $\rho$  counts the number of codes which are not invariant under the action of  $T_{\theta}$ .

Furthermore, using [8, Proposition 6.6, Proposition 6.9, Proposition 6.10, Proposition 6.12, and Proposition 6.13] and the fact about the map  $\rho$  as above, we have the following results.

**Proposition 9.** Suppose  $q \equiv 1 \mod 4$  and d is even, or  $q \equiv 3 \mod 4$  and  $d \equiv 0 \mod 4$ . Then the number of Euclidean self-dual  $\theta$ -cyclic codes of length 2d over  $\mathbb{F}_q$  is

$$4\prod_{i=1}^{\frac{d}{2}-1}(q^i+1-\rho_{G_i})^2.$$

**Proposition 10.** Suppose that q and d satisfy one of the following conditions,

(1)  $q \equiv 11 \mod 12$  and  $d \equiv 0 \mod 4$ ,

6

(2)  $q \equiv 2 \mod 3$  but  $q \not\equiv 11 \mod 12$  and d is even.

Then the number of distinct Euclidean self-dual  $\theta$ -cyclic codes over  $\mathbb{F}_q$  of length 3d is

$$b(q+1)\prod_{i=1}^{\frac{d}{2}-1}(q^{i}+1-\rho_{G_{i}})(q^{2i+1}+1-\rho_{H_{i}'}),$$

where b = 1 if q is even, 2 if q is odd.

**Proposition 11.** Let q and d satisfy one of the following conditions,

(1)  $q \equiv 7 \mod 12$  and  $d \equiv 0 \mod 4$ ,

(2)  $q \equiv 1 \mod 3$  but  $q \not\equiv 7 \mod 12$  and d is even.

Then the number of distinct Euclidean self-dual  $\theta$  -cyclic codes over  $\mathbb{F}_q$  of length 3d is

$$b\left(\prod_{i=1}^{\frac{d}{2}-1} (q^{i}+1-\rho_{G_{i}})\right) (N(d,q)-\rho_{H',H''}),$$

where b = 1 if q is even, 2 if q is odd.

**Proposition 12.** Let q be an odd prime power such that -1 is not a square in  $\mathbb{F}_q$  and let  $d \equiv 0 \mod 4$ . Then the number of distinct Euclidean self-dual  $\theta$ -cyclic codes over  $\mathbb{F}_q$  of length 4d is

$$4(q+1)\prod_{i=1}^{\frac{a}{2}-1}(q^{i}+1-\rho_{G_{i}})^{2}(q^{2i+1}+1-\rho_{H_{i}'}).$$

**Proposition 13.** Let d be an even integer and q be an odd prime power such that -1 is a square in  $\mathbb{F}_q$ . Then, the number of distinct Euclidean self-dual  $\theta$ -cyclic codes of length 4d over  $\mathbb{F}_q$  is

$$4\left(\prod_{i=1}^{\frac{d}{2}-1} (q^i + 1 - \rho_{G_i})^2\right) (N(d,q) - \rho_{H',H''})$$

# 3. CONCLUSION

Enumeration of skew-cyclic or  $\theta$ -cyclic codes over finite fields has been considered by Boucher and her coauthors in [5], [1]. However, the enumeration of this class of codes has not yet been done completely. In this paper, we study the enumeration of self-dual skew-cyclic or  $\theta$ -cyclic codes if  $(n, |\theta|) = 1$  and for some cases if  $(n, |\theta|) =$ d > 1, where  $|\theta|$  is the order of the automorphism  $\theta$ .

## 4. Acknowledgment

I, I.M-A, and A.B. are supported in part by Riset Unggulan Perguruan Tinggi Dikti 2016. D.S. is supported in part by Riset ITB 2016.

## References

- [1] D. Boucher, Construction and number of self-dual skew codes over  $\mathbb{F}_{p^2}$ , hal-01090922v2, 2014.
- [2] D. Boucher, A note on the existence of self-dual skew codes over finite fields, Lecture Notes in Computer Science, Proceedings in Honour of Thierry Berger 9084, pp. 228-239, 2015.
- [3] D. Boucher, W. Geiselmann, and F. Ulmer, Skew-cyclic codes, Applicable Algebra in Engineering, Communication, and Computing vol. 18 issue 4, pp. 379-389, 2007.
- [4] D. Boucher and F. Ulmer, Coding with skew polynomial rings, Journal of Symbolic Computation 44, pp. 1644-1656, 2009.
- [5] D. Boucher and F. Ulmer, Self-dual skew-codes and factorization of skew-polynomials, *Journal of Symbolic Computation* 60, pp. 47-61, 2013.
- [6] W. Huffman and V. Pless, Fundamentals of error correcting codes, Cambridge University Press, 2003.
- [7] Y. Jia, S. Ling, and C. Xing, On self-dual cyclic codes over finite fields, *IEEE Transactions on Information Theory* Vol. 57 No. 4, pp. 2243-2251, 2011.
- [8] S. Ling and P. Solé, On the algebraic structure of quasi-cyclic codes I: Finite fields, IEEE Transactions on Information Theory Vol. 47 No. 7, pp. 2751-2760, 2001.

MATHEMATICS DEPARTMENT, FACULTY OF MATHEMATICS AND NATURAL SCIENCES, UNIVERSITAS MATARAM, MATARAM, INDONESIA

E-mail address: irw@unram.ac.id

ALGEBRA RESEARCH GROUP, FACULTY OF MATHEMATICS AND NATURAL SCIENCES, INSTITUT TEKNOLOGI BANDUNG, JL. GANESHA 10, BANDUNG, 40132, INDONESIA

*E-mail address*: ntan, muchlis, barra@math.itb.ac.id

COMBINATORIAL MATHEMATICS RESEARCH GROUP, FACULTY OF MATHEMATICS AND NATURAL SCIENCES, INSTITUT TEKNOLOGI BANDUNG, JL. GANESHA 10, BANDUNG, 40132, INDONESIA *E-mail address*: djoko@math.itb.ac.id