

# Analytical Calculation of Stored Electrostatic Energy per Unit Length for an Infinite Charged Line and an Infinitely Long Cylinder in the Framework of Born-Infeld Electrostatics

S. K. Moayedi <sup>\*</sup>, M. Shafabakhsh <sup>†</sup>, F. Fathi <sup>‡</sup>

*Department of Physics, Faculty of Sciences, Arak University, Arak 38156-8-8349, Iran*

## Abstract

More than 80 years ago, Born-Infeld electrodynamics was proposed in order to remove the point charge singularity in Maxwell electrodynamics. In this work, after a brief introduction to Lagrangian formulation of Abelian Born-Infeld model in the presence of an external source, we obtain the explicit forms of Gauss's law and the energy density of an electrostatic field for Born-Infeld electrostatics. The electric field and the stored electrostatic energy per unit length for an infinite charged line and an infinitely long cylinder in Born-Infeld electrostatics are calculated. Numerical estimations in this paper show that the nonlinear corrections to Maxwell electrodynamics are considerable only for strong electric fields. We present an action functional for Abelian Born-Infeld model with an auxiliary scalar field in the presence of an external source. This action functional is a generalization of the action functional which was presented by Tseytlin in his studies on low energy dynamics of  $D$ -branes (Nucl. Phys. **B469**, 51 (1996); Int. J. Mod. Phys. A **19**, 3427 (2004)). Finally, we derive the symmetric energy-momentum tensor for Abelian Born-Infeld model with an auxiliary scalar field.

**Keywords:** Classical field theories; Classical electromagnetism; Other special classical field theories; Nonlinear or nonlocal theories and models

**PACS:** 03.50.-z, 03.50.De, 03.50.Kk, 11.10.Lm

---

<sup>\*</sup>E-mail: s-moayedi@araku.ac.ir

<sup>†</sup>E-mail: m-shafabakhsh@phd.araku.ac.ir

<sup>‡</sup>E-mail: f-fathi@phd.araku.ac.ir

# 1 Introduction

Maxwell electrodynamics is a very successful theory which describes a wide range of macroscopic phenomena in electricity and magnetism. On the other hand, in Maxwell electrodynamics, the electric field of a point charge  $q$  at the position of the point charge is singular, i.e.,

$$\mathbf{E}(\mathbf{x}) = \frac{q}{4\pi\epsilon_0|\mathbf{x}|^2} \frac{\mathbf{x}}{|\mathbf{x}|} \xrightarrow{|\mathbf{x}| \rightarrow 0} \infty.$$

Also, in Maxwell electrodynamics, the classical self-energy of a point charge is

$$U = \frac{q^2}{8\pi\epsilon_0} \int_0^\infty \frac{dr}{r^2} \rightarrow \infty.$$

More than 80 years ago, Born and Infeld proposed a nonlinear generalization of Maxwell electrodynamics [1]. In their generalization, the classical self-energy of a point charge was a finite value [1-6]. Recent studies in string theory show that the dynamics of electromagnetic fields on  $D$ -branes can be described by Born-Infeld theory [7-10]. In a paper on Born-Infeld theory [8], the concept of a BIon was introduced by Gibbons. BIon is a finite energy solution of a nonlinear theory with a distributional source. Today, many physicists believe that the dark energy in our universe can be described by a Born-Infeld type scalar field [11]. The authors of Ref. [12] have presented a non-Abelian generalization of Born-Infeld theory. In their generalization, they have found a one-parameter family of finite energy solutions in the case of the  $SU(2)$  gauge group. In 2013, Hendi [13] proposed a nonlinear generalization of Maxwell electrodynamics which is called exponential electrodynamics [14,15]. The black hole solutions of Einstein's gravity in the presence of exponential electrodynamics in a  $3+1$ -dimensional spacetime are obtained in Ref. [13]. In 2014, Gaete and Helayel-Neto introduced a new generalization of Maxwell electrodynamics which is known as logarithmic electrodynamics [14]. They proved that the classical self-energy of a point charge in logarithmic electrodynamics is a finite value. Recently, a novel generalization of Born-Infeld electrodynamics is presented by Gaete and Helayel-Neto in which the authors show that the field energy of a point-like charge is finite only for Born-Infeld like electrodynamics [15]. In Ref. [16], a nonlinear model for electrodynamics with manifestly broken gauge symmetry is proposed. In the above mentioned model for nonlinear electrodynamics, there are non-singular solitonic solutions which describe charged particles. Another interesting theory of nonlinear electrodynamics was proposed and developed by Heisenberg and his students Euler and Kockel [17-19]. They showed that classical electrodynamics must be corrected by nonlinear terms due to the vacuum polarization effects. In Ref. [20], the charged black hole solutions for Einstein-Euler-Heisenberg theory are obtained. There are three physical motivations in writing this paper. First, the exact solutions of nonlinear field equations are very important in

theoretical physics. These solutions help us to obtain a better understanding of physical reality. According to above statements, we attempt to obtain particular cylindrically symmetric solutions in Born-Infeld electrostatics. The search for spherically symmetric solutions in Born-Infeld electrostatics will be discussed in future works. Second, we want to show that the nonlinear corrections in electrodynamics are considerable only for very strong electric fields and extremely short spatial distances. Third, we hope to remove or at least modify the infinities which appear in Maxwell electrostatics. This paper is organized as follows. In Section 2, we study Lagrangian formulation of Abelian Born-Infeld model in the presence of an external source. The explicit forms of Gauss's law and the energy density of an electrostatic field for Born-Infeld electrostatics are obtained. In Section 3, we calculate the electric field together with the stored electrostatic energy per unit length for an infinite charged line and an infinitely long cylinder in Born-Infeld electrostatics. Summary and conclusions are presented in Section 4. Numerical estimations in Section 4 show that the nonlinear corrections to electric field of an infinite charged line at large radial distances are negligible for weak electric fields. There are two appendices in this paper. In Appendix A, a generalized action functional for Abelian Born-Infeld model with an auxiliary scalar field in the presence of an external source is proposed. In Appendix B, we obtain the symmetric energy-momentum tensor for Abelian Born-Infeld model with an auxiliary scalar field. We use SI units throughout this paper. The metric of spacetime has the signature  $(+, -, -, -)$ .

## 2 Lagrangian Formulation of Abelian Born-Infeld Model with an External Source

The Lagrangian density for Abelian Born-Infeld model in a  $3 + 1$ -dimensional spacetime is [1-6]

$$\mathcal{L}_{BI} = \epsilon_0 \beta^2 \left\{ 1 - \sqrt{1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu}} \right\} - J^\mu A_\mu, \quad (1)$$

where  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$  is the electromagnetic field tensor and  $J^\mu = (c\rho, \mathbf{J})$  is an external source for the Abelian field  $A^\mu = (\frac{1}{c}\phi, \mathbf{A})$ . The parameter  $\beta$  in Eq. (1) is called the nonlinearity parameter of the model. In the limit  $\beta \rightarrow \infty$ , Eq. (1) reduces to the Lagrangian density of the Maxwell field, i.e.,

$$\mathcal{L}_{BI}|_{large \beta} = \mathcal{L}_M + \mathcal{O}(\beta^{-2}), \quad (2)$$

where  $\mathcal{L}_M = -\frac{1}{4\mu_0}F_{\mu\nu}F^{\mu\nu} - J^\mu A_\mu$  is the Maxwell Lagrangian density. The Euler-Lagrange equation for the Born-Infeld field  $A^\mu$  is

$$\frac{\partial \mathcal{L}_{BI}}{\partial A_\lambda} - \partial_\rho \left( \frac{\partial \mathcal{L}_{BI}}{\partial (\partial_\rho A_\lambda)} \right) = 0. \quad (3)$$

If we substitute Lagrangian density (1) in the Euler-Lagrange equation (3), we will obtain the inhomogeneous Born-Infeld equations as follows:

$$\partial_\rho \left( \frac{F^{\rho\lambda}}{\sqrt{1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu}}} \right) = \mu_0 J^\lambda. \quad (4)$$

The electromagnetic field tensor  $F_{\mu\nu}$  satisfies the Bianchi identity:

$$\partial_\mu F_{\nu\lambda} + \partial_\nu F_{\lambda\mu} + \partial_\lambda F_{\mu\nu} = 0. \quad (5)$$

Equation (5) leads to the homogeneous Maxwell equations. In 3+1-dimensional spacetime, the components of  $F_{\mu\nu}$  can be written as follows:

$$F_{\mu\nu} = \begin{pmatrix} 0 & E_x/c & E_y/c & E_z/c \\ -E_x/c & 0 & -B_z & B_y \\ -E_y/c & B_z & 0 & -B_x \\ -E_z/c & -B_y & B_x & 0 \end{pmatrix}. \quad (6)$$

Using Eq. (6), Eqs. (4) and (5) can be written in the vector form as follows:

$$\nabla \cdot \left( \frac{\mathbf{E}(\mathbf{x}, t)}{\sqrt{1 - \frac{\mathbf{E}^2(\mathbf{x}, t) - c^2 \mathbf{B}^2(\mathbf{x}, t)}{\beta^2}}} \right) = \frac{\rho(\mathbf{x}, t)}{\epsilon_0}, \quad (7)$$

$$\nabla \times \mathbf{E}(\mathbf{x}, t) = -\frac{\partial \mathbf{B}(\mathbf{x}, t)}{\partial t}, \quad (8)$$

$$\nabla \times \left( \frac{\mathbf{B}(\mathbf{x}, t)}{\sqrt{1 - \frac{\mathbf{E}^2(\mathbf{x}, t) - c^2 \mathbf{B}^2(\mathbf{x}, t)}{\beta^2}}} \right) = \mu_0 \mathbf{J}(\mathbf{x}, t) + \frac{1}{c^2} \frac{\partial}{\partial t} \left( \frac{\mathbf{E}(\mathbf{x}, t)}{\sqrt{1 - \frac{\mathbf{E}^2(\mathbf{x}, t) - c^2 \mathbf{B}^2(\mathbf{x}, t)}{\beta^2}}} \right), \quad (9)$$

$$\nabla \cdot \mathbf{B}(\mathbf{x}, t) = 0. \quad (10)$$

The symmetric energy-momentum tensor for the Abelian Born-Infeld model in Eq. (1) has been obtained by Accioly [21] as follows:

$$T^\mu_\lambda = \frac{1}{\mu_0} \left[ \frac{F^{\mu\nu} F_{\nu\lambda}}{\Omega} + \frac{\beta^2}{c^2} (\Omega - 1) \delta^\mu_\lambda \right], \quad (11)$$

where  $\Omega := \sqrt{1 + \frac{c^2}{2\beta^2} F_{\alpha\gamma} F^{\alpha\gamma}}$ . The classical Born-Infeld equations (7)-(10) for an electrostatic field  $\mathbf{E}(\mathbf{x})$  are

$$\nabla \cdot \left( \frac{\mathbf{E}(\mathbf{x})}{\sqrt{1 - \frac{\mathbf{E}^2(\mathbf{x})}{\beta^2}}} \right) = \frac{\rho(\mathbf{x})}{\epsilon_0}, \quad (12)$$

$$\nabla \times \mathbf{E}(\mathbf{x}) = 0. \quad (13)$$

Equations (12) and (13) are fundamental equations of Born-Infeld electrostatics [2]. Using divergence theorem, the integral form of Eq. (12) can be written in the form

$$\oint_S \frac{1}{\sqrt{1 - \frac{\mathbf{E}^2(\mathbf{x})}{\beta^2}}} \mathbf{E}(\mathbf{x}) \cdot \mathbf{n} \, da = \frac{1}{\epsilon_0} \int_V \rho(\mathbf{x}) d^3x, \quad (14)$$

where  $V$  is the three-dimensional volume enclosed by a two-dimensional surface  $S$ . Equation (14) is Gauss's law in Born-Infeld electrostatics. Using Eqs. (6) and (11), the energy density of an electrostatic field in Born-Infeld theory is given by

$$u(\mathbf{x}) = \epsilon_0 \beta^2 \left( \frac{1}{\sqrt{1 - \frac{\mathbf{E}^2(\mathbf{x})}{\beta^2}}} - 1 \right). \quad (15)$$

In the limit  $\beta \rightarrow \infty$ , the modified electrostatic energy density in Eq. (15) smoothly becomes the usual electrostatic energy density in Maxwell theory, i.e.,

$$u(\mathbf{x})|_{\text{large } \beta} = \frac{1}{2} \epsilon_0 \mathbf{E}^2(\mathbf{x}) + \mathcal{O}(\beta^{-2}). \quad (16)$$

### 3 Calculation of Stored Electrostatic Energy per Unit Length for an Infinite Charged Line and an Infinitely Long Cylinder in Born-Infeld Electrostatics

#### 3.1 Infinite Charged Line

Let us consider an infinite charged line with a uniform positive linear charge density  $\lambda$  which is located on the  $z$ -axis. Now, we find an expression for the electric field  $\mathbf{E}(\mathbf{x})$  at a radial distance  $\rho$  from the  $z$ -axis. Because of the cylindrical symmetry of the problem, the suitable Gaussian surface is a circular cylinder of radius  $\rho$  and length  $L$ , coaxial with the  $z$ -axis (see Fig. 1).




Figure 1: The Gaussian surface for an infinite charged line. The cylindrical symmetry of the problem implies that  $\mathbf{E}(\mathbf{x}) = E_\rho(\rho)\hat{\mathbf{e}}_\rho$ , where  $\hat{\mathbf{e}}_\rho$  is the radial unit vector in cylindrical coordinates  $(\rho, \varphi, z)$ .

Using the cylindrical symmetry of the problem together with the modified Gauss's law in Eq. (14), the electric field for the Gaussian surface in Fig. 1 becomes

$$\mathbf{E}(\mathbf{x}) = \frac{\lambda}{2\pi\epsilon_0\rho} \frac{1}{\sqrt{1 + \left(\frac{\lambda}{2\pi\epsilon_0\beta\rho}\right)^2}} \hat{\mathbf{e}}_\rho. \quad (17)$$

In contrast with Maxwell electrostatics, the electric field  $\mathbf{E}(\mathbf{x})$  in Eq. (17) has a finite value on the  $z$ -axis, i.e.,

$$\lim_{\rho \rightarrow 0} \mathbf{E}(\mathbf{x}) = \beta \hat{\mathbf{e}}_\rho. \quad (18)$$

At large radial distances from the  $z$ -axis, the asymptotic behavior of the electric field in Eq. (17) is given by

$$\mathbf{E}(\mathbf{x}) = \frac{\lambda}{2\pi\epsilon_0\rho} \hat{\mathbf{e}}_\rho - \frac{\lambda^3}{16\pi^3\epsilon_0^3\beta^2\rho^3} \hat{\mathbf{e}}_\rho + \mathcal{O}(\rho^{-5}). \quad (19)$$

The first term on the right-hand side of Eq. (19) shows the electric field of an infinite charged line in Maxwell electrostatics, while the second and higher order terms in Eq. (19) show the effect of nonlinear corrections. By putting Eq. (17) in Eq. (15), the electrostatic

energy density for an infinite charged line in Born-Infeld electrostatics can be written as follows:

$$u(\mathbf{x}) = \epsilon_0 \beta^2 \left( \sqrt{1 + \left( \frac{\lambda}{2\pi\epsilon_0\beta\rho} \right)^2} - 1 \right). \quad (20)$$

Using Eq. (20), the stored electrostatic energy per unit length for an infinite charged line in the radial interval  $0 \leq \rho \leq \Lambda$  is given by

$$\begin{aligned} \frac{U}{L} &= \int_0^\Lambda \int_0^{2\pi} u(\mathbf{x}) \rho d\rho d\varphi \\ &= 2\pi\epsilon_0\beta^2 \left\{ \frac{\Lambda}{2} \sqrt{\Lambda^2 + \left( \frac{\lambda}{2\pi\epsilon_0\beta} \right)^2} - \frac{\Lambda^2}{2} \right. \\ &\quad \left. + \frac{1}{2} \left( \frac{\lambda}{2\pi\epsilon_0\beta} \right)^2 \ln \left( \frac{\Lambda + \sqrt{\Lambda^2 + \left( \frac{\lambda}{2\pi\epsilon_0\beta} \right)^2}}{\left( \frac{\lambda}{2\pi\epsilon_0\beta} \right)} \right) \right\}. \end{aligned} \quad (21)$$

It is necessary to note that the above value for  $\frac{U}{L}$  has an infinite value in Maxwell theory. In the limit of large  $\beta$ , the expression for  $\frac{U}{L}$  in Eq. (21) diverges logarithmically as  $\ln\beta$ . Hence, it seems that the finite regularization parameter  $\beta$  removes the logarithmic divergence in Eq. (21).

### 3.2 Infinitely Long Cylinder

In this subsection, we determine the electric field  $\mathbf{E}(\mathbf{x})$  and stored electrostatic energy per unit length for an infinitely long cylinder of radius  $R$  and uniform positive volume charge density  $\tau$ . As in the previous subsection, we assume that the Gaussian surface is a cylindrical closed surface of radius  $\rho$  and length  $L$  with a common axis with the infinitely long cylinder (see Fig. 2).

According to modified Gauss's law in Eq. (14), the electric field for the Gaussian surfaces in Fig. 2 is given by

$$\mathbf{E}(\mathbf{x}) = \begin{cases} \frac{\tau\rho}{2\epsilon_0} \frac{1}{\sqrt{1 + \left( \frac{\tau\rho}{2\epsilon_0\beta} \right)^2}} \hat{\mathbf{e}}_\rho; & \rho < R, \\ \frac{\tau R^2}{2\epsilon_0\rho} \frac{1}{\sqrt{1 + \left( \frac{\tau R^2}{2\epsilon_0\beta\rho} \right)^2}} \hat{\mathbf{e}}_\rho; & \rho > R. \end{cases} \quad (22)$$




Figure 2: The Gaussian surface for an infinitely long cylinder of radius  $R$ . Left) inside the cylinder ( $\rho < R$ ). Right) outside the cylinder ( $\rho > R$ ).

For the large values of the nonlinearity parameter  $\beta$ , the behavior of the electric field  $\mathbf{E}(\mathbf{x})$  in Eq. (22) is as follows:

$$\mathbf{E}(\mathbf{x}) = \begin{cases} \frac{\tau\rho}{2\epsilon_0} \hat{\mathbf{e}}_\rho - \frac{\tau^3\rho^3}{16\epsilon_0^3\beta^2} \hat{\mathbf{e}}_\rho + \mathcal{O}(\beta^{-4}); & \rho < R, \\ \frac{\tau R^2}{2\epsilon_0\rho} \hat{\mathbf{e}}_\rho - \frac{\tau^3 R^6}{16\epsilon_0^3\beta^2\rho^3} \hat{\mathbf{e}}_\rho + \mathcal{O}(\beta^{-4}); & \rho > R. \end{cases} \quad (23)$$

Hence, for the large values of  $\beta$ , the electric field  $\mathbf{E}(\mathbf{x})$  in Eq. (23) becomes the electric field of an infinitely long cylinder in Maxwell electrostatics. If we substitute Eq. (22) into Eq. (15), we will obtain the electrostatic energy density for an infinitely long cylinder in Born-Infeld electrostatics as follows:

$$u(\mathbf{x}) = \begin{cases} \epsilon_0\beta^2 \left( \sqrt{1 + \left( \frac{\tau\rho}{2\epsilon_0\beta} \right)^2} - 1 \right); & \rho < R, \\ \epsilon_0\beta^2 \left( \sqrt{1 + \left( \frac{\tau R^2}{2\epsilon_0\beta\rho} \right)^2} - 1 \right); & \rho > R. \end{cases} \quad (24)$$

Using Eq. (24), the stored electrostatic energy per unit length for an infinitely long cylinder in the radial interval  $0 \leq \rho \leq \Lambda$  ( $\Lambda > R$ ) is given by



$$\begin{aligned}
\frac{U}{L} &= 2\pi\epsilon_0\beta^2 \left\{ \int_0^R \left( \sqrt{1 + \left(\frac{\tau\rho}{2\epsilon_0\beta}\right)^2} - 1 \right) \rho d\rho + \int_R^\Lambda \left( \sqrt{1 + \left(\frac{\tau R^2}{2\epsilon_0\beta\rho}\right)^2} - 1 \right) \rho d\rho \right\} \\
&= 2\pi\epsilon_0\beta^2 \left\{ -\frac{\Lambda^2}{2} + \left(\frac{2\epsilon_0\beta}{\tau\sqrt{3}}\right)^2 \left[ \left(1 + \left(\frac{\tau R}{2\epsilon_0\beta}\right)^2\right)^{\frac{3}{2}} - 1 \right] + \frac{\Lambda^2}{2} \sqrt{1 + \left(\frac{\tau R^2}{2\epsilon_0\beta\Lambda}\right)^2} \right. \\
&\quad \left. - \frac{R^2}{2} \sqrt{1 + \left(\frac{\tau R}{2\epsilon_0\beta}\right)^2} + \frac{1}{2} \left(\frac{\tau R^2}{2\epsilon_0\beta}\right)^2 \ln \left( \frac{\Lambda + \Lambda \sqrt{1 + \left(\frac{\tau R^2}{2\epsilon_0\beta\Lambda}\right)^2}}{R + R \sqrt{1 + \left(\frac{\tau R}{2\epsilon_0\beta}\right)^2}} \right) \right\}. \tag{25}
\end{aligned}$$

In the limit of large  $\beta$ , the expression for  $\frac{U}{L}$  in Eq. (25) can be expanded in powers of  $\frac{1}{\beta^2}$  as follows:

$$\frac{U}{L} \Big|_{large \beta} = \frac{\pi\tau^2 R^4}{4\epsilon_0} \left( \frac{1}{4} + \ln \frac{\Lambda}{R} \right) + \mathcal{O}(\beta^{-2}). \tag{26}$$

The first term on the right-hand side of Eq. (26) shows the stored electrostatic energy per unit length for an infinitely long cylinder in the radial interval  $0 \leq \rho \leq \Lambda$  ( $\Lambda > R$ ) in Maxwell electrostatics.

## 4 Summary and Conclusions

In 1934, Born and Infeld introduced a nonlinear generalization of Maxwell electrodynamics, in which the classical self-energy of a point charge like electron became a finite value [1]. We showed that, in the limit of large  $\beta$ , the modified Gauss's law in Born-Infeld electrostatics is

$$\oint_S \left[ 1 + \frac{1}{2} \frac{\mathbf{E}^2(\mathbf{x})}{\beta^2} + \frac{3}{8} \frac{(\mathbf{E}^2(\mathbf{x}))^2}{\beta^4} + \mathcal{O}(\beta^{-6}) \right] \mathbf{E}(\mathbf{x}) \cdot \mathbf{n} \, da = \frac{1}{\epsilon_0} \int_V \rho(\mathbf{x}) d^3x. \tag{27}$$

By using the modified Gauss's law in Eq. (14), we calculated the electric field of an infinite charged line and an infinitely long cylinder in Born-Infeld electrostatics. The stored electrostatic energy per unit length for the above configurations of charge density has been calculated in the framework of Born-Infeld electrostatics. Born and Infeld attempted to determine  $\beta$  by equating the classical self-energy of the electron in their theory with its rest mass energy. They obtained the following numerical value for the nonlinearity parameter  $\beta$  [1]:

$$\beta = 1.2 \times 10^{20} \frac{V}{m}. \tag{28}$$

In 1973, Soff, Rafelski and Greiner [22] have estimated a lower bound on  $\beta$ . This lower bound on  $\beta$  is

$$\beta \geq 1.7 \times 10^{22} \frac{V}{m}. \quad (29)$$

Recent studies on photonic processes in Born-Infeld theory show that the numerical value of  $\beta$  is close to  $1.2 \times 10^{20} V/m$  in Eq. (28) [23]. In order to obtain a better understanding of nonlinear effects in Born-Infeld electrostatics, let us estimate the numerical value of the second term on the right-hand side of Eq. (19). For this purpose, we rewrite Eq. (19) as follows:

$$\mathbf{E}(\mathbf{x}) = \mathbf{E}_0(\mathbf{x}) + \Delta\mathbf{E}(\mathbf{x}) + \mathcal{O}(\rho^{-5}), \quad (30)$$

where

$$\mathbf{E}_0(\mathbf{x}) := \frac{\lambda}{2\pi\epsilon_0\rho} \hat{\mathbf{e}}_\rho, \quad (31)$$

$$\Delta\mathbf{E}(\mathbf{x}) := -\frac{\lambda^3}{16\pi^3\epsilon_0^3\beta^2\rho^3} \hat{\mathbf{e}}_\rho. \quad (32)$$

Using Eqs. (31) and (32), the ratio of  $\Delta\mathbf{E}(\mathbf{x})$  to  $\mathbf{E}_0(\mathbf{x})$  is given by

$$\frac{|\Delta\mathbf{E}(\mathbf{x})|}{|\mathbf{E}_0(\mathbf{x})|} = \frac{1}{2} \frac{\mathbf{E}_0^2(\mathbf{x})}{\beta^2}. \quad (33)$$

Let us assume the following approximate but realistic values [24]:

$$L = 1.80 m, \quad \rho = 0.10 m, \quad Q = +24 \mu C, \quad \lambda = 1.33 \times 10^{-5} \frac{C}{m}. \quad (34)$$

By putting Eqs. (28) and (34) into Eq. (33), we get

$$|\Delta\mathbf{E}(\mathbf{x})| \approx 2 \times 10^{-28} |\mathbf{E}_0(\mathbf{x})|. \quad (35)$$

Finally, if we put Eqs. (29) and (34) in Eq. (33), we obtain

$$|\Delta\mathbf{E}(\mathbf{x})| \lesssim 10^{-32} |\mathbf{E}_0(\mathbf{x})|. \quad (36)$$

In fact, as is clear from Eqs. (35) and (36), the nonlinear corrections to electric field in Eq. (19) are very small for weak electric fields. The authors of Ref. [25] have suggested a nonlinear generalization of Maxwell electrodynamics. In their generalization, the electric field of a point charge is singular at the position of the point charge but the classical self-energy of the point charge has a finite value. Recently, Kruglov [26,27] has proposed two different models for nonlinear electrodynamics. In these models, both the electric field of a point charge at the position of the point charge and the classical self-energy of the point charge have finite values. In future works, we hope to study the problems discussed in this research from the viewpoint of Refs. [25-27].

## Acknowledgments

We would like to thank the referees for their careful reading and constructive comments.

## Appendix A: A Generalized Action Functional for Abelian Born-Infeld Model with an Auxiliary Scalar Field

Let us consider the following action functional:

$$S(A, \psi) = \frac{1}{c} \int_{t_0}^t \int_{\mathbb{R}^3} \left[ \epsilon_0 \beta^2 \left( 1 - \omega_1 \psi^{\lambda_1} \left( 1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu} \right) - \omega_2 \psi^{\lambda_2} \right) - J^\mu A_\mu \right] d^4x, \quad (\text{A.1})$$

where  $\psi$  is an auxiliary scalar field and  $\omega_1$ ,  $\lambda_1$ ,  $\omega_2$  and  $\lambda_2$  are four non-zero constants. The variation of Eq. (A.1) with respect to  $\psi$  and  $A_\mu$  leads to the following classical field equations:

$$\psi = \left[ -\frac{\omega_2 \lambda_2}{\omega_1 \lambda_1} \left( 1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu} \right)^{-1} \right] \frac{1}{\lambda_1 - \lambda_2}, \quad (\text{A.2})$$

$$\partial_\mu (\psi^{\lambda_1} F^{\mu\nu}) = \frac{\mu_0}{2\omega_1} J^\nu. \quad (\text{A.3})$$

Substituting Eq. (A.2) into Eq. (A.3), we obtain the following classical field equation:

$$\partial_\mu \left\{ \left[ -\frac{\omega_2 \lambda_2}{\omega_1 \lambda_1} \left( 1 + \frac{c^2}{2\beta^2} F_{\alpha\gamma} F^{\alpha\gamma} \right)^{-1} \right] \frac{\lambda_1}{\lambda_1 - \lambda_2} F^{\mu\nu} \right\} = \frac{\mu_0}{2\omega_1} J^\nu. \quad (\text{A.4})$$

By choosing  $\lambda_1 = \lambda$ ,  $\lambda_2 = -\lambda$  and  $\omega_1 = \omega$ ,  $\omega_2 = \frac{1}{4\omega}$  ( $\omega > 0$ ), Eqs.(A.1) and (A.4) can be written as follows:

$$S(A, \psi) = \frac{1}{c} \int_{t_0}^t \int_{\mathbb{R}^3} \left[ \epsilon_0 \beta^2 \left( 1 - \omega \psi^\lambda \left( 1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu} \right) - \frac{1}{4\omega} \psi^{-\lambda} \right) - J^\mu A_\mu \right] d^4x, \quad (\text{A.5})$$

$$\partial_\mu \left( \frac{F^{\mu\nu}}{\sqrt{1 + \frac{c^2}{2\beta^2} F_{\alpha\gamma} F^{\alpha\gamma}}} \right) = \mu_0 J^\nu. \quad (\text{A.6})$$

Equation (A.5) is the generalized action functional for Abelian Born-Infeld model with an auxiliary scalar field  $\psi$ . Also, Eq. (A.6) is the inhomogeneous Born-Infeld equation

(see Eq. (4)). If we choose  $\omega = \frac{1}{2}$  and  $\lambda = 1$  in Eq. (A.5), we will obtain the following action functional:

$$S(A, \psi) = \frac{1}{c} \int_{t_0}^t \int_{\mathbb{R}^3} \left[ \epsilon_0 \beta^2 \left( 1 - \frac{\psi}{2} \left( 1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu} \right) - \frac{1}{2\psi} \right) - J^\mu A_\mu \right] d^4x. \quad (\text{A.7})$$

The above action functional was presented by Tseytlin in his studies on low energy dynamics of  $D$ -branes [28].

## Appendix B: The Symmetric Energy-Momentum Tensor for Abelian Born-Infeld Model with an Auxiliary Scalar Field

In this appendix, we want to obtain the symmetric energy-momentum tensor for Abelian Born-Infeld model with an auxiliary scalar field. According to Eq. (A.5), the Lagrangian density for Abelian Born-Infeld model with an auxiliary scalar field  $\psi$  in the absence of external source  $J^\mu$  is

$$\mathcal{L} = \epsilon_0 \beta^2 \left( 1 - \omega \psi^\lambda \left( 1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu} \right) - \frac{1}{4\omega} \psi^{-\lambda} \right). \quad (\text{B.1})$$

From Eq. (B.1), we obtain the following classical field equations:

$$\psi^\lambda = \frac{1}{2\omega} \left( 1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu} \right)^{-\frac{1}{2}}, \quad (\text{B.2})$$

$$\partial_\mu \left( \psi^\lambda F^{\mu\nu} \right) = 0. \quad (\text{B.3})$$

The canonical energy-momentum tensor for Eq. (B.1) is

$$\begin{aligned} T_\gamma^\alpha &= \frac{\partial \mathcal{L}}{\partial(\partial_\alpha A_\eta)} (\partial_\gamma A_\eta) + \frac{\partial \mathcal{L}}{\partial(\partial_\alpha \psi)} (\partial_\gamma \psi) - \delta_\gamma^\alpha \mathcal{L} \\ &= -2\epsilon_0 c^2 \omega \psi^\lambda F^{\alpha\eta} (\partial_\gamma A_\eta) - \epsilon_0 \beta^2 \delta_\gamma^\alpha \left( 1 - \omega \psi^\lambda \left( 1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu} \right) - \frac{1}{4\omega} \psi^{-\lambda} \right). \end{aligned} \quad (\text{B.4})$$

Using Eqs. (B.2) and (B.3), the canonical energy-momentum tensor  $T_\gamma^\alpha$  in Eq. (B.4) can be rewritten as follows:

$$\begin{aligned} T_\gamma^\alpha &= -\epsilon_0 c^2 \left( 1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu} \right)^{-\frac{1}{2}} F^{\alpha\eta} F_{\gamma\eta} \\ &\quad - \epsilon_0 \beta^2 \delta_\gamma^\alpha \left( 1 - \left( 1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu} \right)^{\frac{1}{2}} \right) + \partial_\eta M_{\gamma}^{\eta\alpha}, \end{aligned} \quad (\text{B.5})$$

where

$$M_{\gamma}^{\eta\alpha} := \frac{1}{\mu_0} \frac{F^{\eta\alpha}}{\sqrt{1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu}}} A_{\gamma}, \quad M^{\alpha\eta}_{\gamma} = -M^{\eta\alpha}_{\gamma}. \quad (\text{B.6})$$

After dropping the total divergence term  $\partial_{\eta} M^{\eta\alpha}_{\gamma}$  in Eq. (B.5), we get the following expression for the symmetric energy-momentum tensor:

$$T_{\gamma}^{\alpha} = -\epsilon_0 c^2 \left(1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu}\right)^{-\frac{1}{2}} F^{\alpha\eta} F_{\gamma\eta} - \epsilon_0 \beta^2 \delta_{\gamma}^{\alpha} \left(1 - \left(1 + \frac{c^2}{2\beta^2} F_{\mu\nu} F^{\mu\nu}\right)^{\frac{1}{2}}\right). \quad (\text{B.7})$$

If we use Eqs. (6) and (B.7), we will obtain the electrostatic energy density for Abelian Born-Infeld model with an auxiliary scalar field as follows:

$$u(\mathbf{x}) = T_0^0(\mathbf{x}) = \epsilon_0 \beta^2 \left( \frac{1}{\sqrt{1 - \frac{\mathbf{E}^2(\mathbf{x})}{\beta^2}}} - 1 \right). \quad (\text{B.8})$$

## References

- [1] M. Born and L. Infeld, Proc. R. Soc. London **A144**, 425 (1934).
- [2] D. Fortunato, L. Orsina, and L. Pisani, J. Math. Phys. **43**, 5698 (2002).
- [3] S. H. Mazharimousavi and M. Halilsoy, Found. Phys. **42**, 524 (2012).
- [4] L. P. G. De Assis, P. Gaete, J. A. Helayel-Neto, and S. O. Vellozo, Int. J. Theor. Phys. **51**, 477 (2012).
- [5] Z. K. Vad, Acta Physica Hungarica **63**, 353 (1988).
- [6] P. Gaete and I. Schmidt, Int. J. Mod. Phys. A **19**, 3427 (2004).
- [7] C. G. Callan and J. M. Maldacena, Nucl. Phys. **B513**, 198 (1998).
- [8] G. W. Gibbons, Nucl. Phys. **B514**, 603 (1998).
- [9] G. W. Gibbons, Rev. Mex. Fis. **49S1**, 19 (2003).
- [10] B. Zwiebach, A First Course in String Theory (Cambridge University Press, Cambridge, UK, 2009), 2nd ed..
- [11] H. Q. Lu, Int. J. Mod. Phys. D **14**, 355 (2005).
- [12] E. Serie, T. Masson, and R. Kerner, Phys. Rev. D **68**, 125003 (2003).

- [13] S. H. Hendi, *Ann. Phys. (N.Y.)* **333**, 282 (2013).
- [14] P. Gaete and J. Helayel-Neto, *Eur. Phys. J. C* **74**, 2816 (2014).
- [15] P. Gaete and J. Helayel-Neto, *Eur. Phys. J. C* **74**, 3182 (2014).
- [16] N. Riazi and M. Mohammadi, *Int. J. Theor. Phys.* **51**, 1276 (2012).
- [17] H. Euler and B. Kockel, *Naturwissenschaften* **23**, 246 (1935).
- [18] H. Euler, *Ann. Phys. (Leipzig)* **26**, 398 (1936).
- [19] W. Heisenberg and H. Euler, *Z. Phys.* **98**, 714 (1936).
- [20] R. Ruffini, Y.-B. Wu, and S.-S. Xue, *Phys. Rev. D* **88**, 085004 (2013).
- [21] A. Accioly, *Am. J. Phys.* **65** (9), 882 (1997).
- [22] G. Soff, J. Rafelski, and W. Greiner, *Phys. Rev. A* **7**, 903 (1973).
- [23] J. M. Davila, C. Schubert, and M. A. Trejo, *Int. J. Mod. Phys. A* **29**, 1450174 (2014).
- [24] D. Halliday, R. Resnick, and J. Walker, *Fundamentals of physics Extended* (Wiley, 2014), 10th ed..
- [25] C. V. Costa, D. M. Gitman, and A. E. Shabad, “Finite field-energy of a point charge in QED,” arXiv:1312.0447.
- [26] S. I. Kruglov, *Ann. Phys. (N.Y.)* **353**, 299 (2015).
- [27] S. I. Kruglov, “Nonlinear arcsin-electrodynamics,” arXiv:1410.7633.
- [28] A. A. Tseytlin, *Nucl. Phys.* **B469**, 51 (1996).