Structure and Topology of Band Structures in the 1651 Magnetic Space Groups

Haruki Watanabe,^{1,*} Hoi Chun Po,^{2,*} and Ashvin Vishwanath²

¹Department of Applied Physics, University of Tokyo, Tokyo 113-8656, Japan ²Department of Physics, Harvard University, Cambridge MA 02138

The properties of electrons in magnetically ordered crystals are of interest both from the viewpoint of realizing novel topological phases, such as magnetic Weyl semimetals, and from the applications perspective of creating energy-efficient memories. A systematic study of symmetry and topology in magnetic materials has been challenging given that there are 1651 magnetic space groups (MSGs). Here, by using an efficient representation of allowed band structures, we obtain a systematic description of several basic properties of free electrons in all MSGs in three dimensions as well as in the 528 magnetic layer groups relevant to two dimensional magnetic materials. We compute constraints on electron fillings and band connectivity compatible with insulating behavior. Also, by contrasting with atomic insulators, we identify band topology entailed by the symmetry transformation of bands, as determined by the MSG alone. We give an application of our results to identifying topological semimetals arising in periodic arrangements of hedgehog-like magnetic textures.

I. INTRODUCTION

The recent discovery of topological insulators and other topological phases [1] has revitalized the venerable subject of band theory. In addition to the explosion of understanding of different forms of band topology and how symmetries protect or prevent them, rapid progress is being made on several other fronts. For example, fundamental questions such as the connection between electron count and insulating behavior [2–9], as well as the constraints imposed by crystal symmetries on the connectivity of bands [2–4, 9–16], have been resolved. New information has also been gleaned by contrasting real-space and momentum-space descriptions in all 230 crystal space groups (SGs) of nonmagnetic materials [2–4, 7–9, 14, 15, 17].

Increasingly, attention is turning to electronic systems that combine magnetism with band topology. Here, a further panoply of novel topological phenomena is anticipated. Examples that have already been realized, to name just a few, include the quantized anomalous Hall effect in magnetic topological insulators [18, 19] and the topological Hall effect in skyrmion crystals [20–24]. Promising magnetic Weyl semimetals candidates are also now being intensively studied [25–27]. In addition to the fundamental physics interest of these novel phases, current-driven magnetic textures such as domain walls or skyrmions could be harnessed in technological applications, say in the development of energy-efficient memory devices [28].

Despite these strong motivations, the pace of discovery of magnetic topological materials has been relatively slow compared to their nonmagnetic counterparts. There are at least two reasons for this: First, magnetic materials are necessarily correlated, making the prediction of the magnetic structure and properties more challenging. Consequently the set of well characterized magnetic materials is relatively small. Second, the sheer complexity of combining magnetic structures with SGs makes an exhaustive study of the theoretical possibilities daunting. Indeed, there are a total of 1651 different magnetic space groups (MSGs), which were only tabulated in the 1950s. Also, unlike for the 230 nonmagnetic SGs [29, 30], relevant group-representation information is not always readily available.

Here, we will tackle the second problem by providing a systematic understanding of electronic band structures in the 1651 MSGs. Inspired by the recent synthesis of atomically thin magnets [31, 32], we also discuss the 528 magnetic layer groups (MLGs), relevant to two-dimensional magnetic materials. We report results for three key properties, tabulated in Appendix A and described below.

First, we determine electron fillings that can be compatible with insulating behavior in all MSGs. The electron count is a fundamental characteristic of an electronic crystal. The presence of nonsymmorphic symmetries, such as glides and screws, enforces connectivity of bands that raises the required fillings for realizing band insulators [2–4, 11]. These conditions may be useful in the search for magnetic Weyl semimetals, as one can target fillings that, while forbidding band insulators, are nonetheless consistent with nodal-point Fermi surfaces [12, 33, 34].

Next, it has long been known that representations of energy levels at high-symmetry points must connect in specific ways in obtaining a set of isolated bands [10]. This can be viewed as a refinement of the filling condition, which imposes additional constraints on band structures (BSs) [13–16]. The solutions to these constraints are most conveniently represented as a vector space (but with integer coefficients, so more accurately a 'lattice' in mathematical terminology) [13, 14, 35], and are described using only a handful of basis vectors whose precise number $d_{\rm BS}$ depends on the MSG.

Lastly, we contrast the general BSs defined in the momentum space with the subset of those obtained from 'atomic insulators' (AI), in which electrons are tightly bound to sites in the lattice furnishing orbitals with different symmetries. Significantly, we find that this produces a vector space of the *same* dimension, which in mathematical terms is summarized by the equality $d_{AI} = d_{BS}$. However, the bases for BSs and AIs do not generally coincide, and we determine the classes of band structures that cannot be reduced to any atomic limit. This leads to an obstruction to finding symmetric localized Wannier states [36, 37], and corresponds to band topology that

^{*} These two authors contributed equally to this work.

can be diagnosed without detailed information about the electronic wave-functions. We give an example of how this can be used to diagnose topological semimetals, and also an example of a nodal-line semimetal diagnosed through its filling. The latter consists of hedgehog-antihedgehog magnetic order with a lone electron at the core of these defects, leading to gapless behavior.

We note that much progress has recently been made on related problems for nonmagnetic SGs [8, 9, 14, 15, 17]. However, in the case of MSGs these problems have only been attacked in certain restricted settings, for example in generalizing the parity criterion [38] to identify Weyl semimetals, Chern insulators, and axion insulators [35, 39–42]. Here, we furnish a systematic study of all MSGs. In particular, it is important to emphasize that, although our approach has several features in common with K-theory-based classifications [13, 43–45], we do not seek to fully classify crystalline electronic phases here. As discussed in a recent insightful work [13] for two dimensional wallpaper groups, part of such a general classification includes band-structure connectivities, a problem that has been posed since the early days of band theory [10]. Utilizing a result we prove below, we construct a framework that greatly simplifies such computations, which enables an extension of previous results [8, 9, 14] to the physically important case of MSGs.

In addition, topological distinctions revealed using our symmetry-based indicators fit naturally into well-established frameworks [13, 14, 43, 45], and remain stable upon addition of trivial degrees of freedom. Yet, our approach is inherently symmetry-based, and so for crystals with low symmetry, say with only lattice translations, topological insulators cannot be diagnosed without further knowledge of the wavefunctions. A similar caveat pertains to the complementary quantum chemistry approach of Ref. 15, which elaborates on the theory developed in Ref. 2, and takes as input a specific set of orbitals at fixed locations in the crystal. While convenient for representing quantum chemistry information, it is less suited to capturing stable topological distinctions that survive the inclusion of additional trivial bands. Also, one forgoes the simplification arising from the vector space like representation of energy bands that allowed us to generate results for all MSGs, while Ref. 15 is currently confined to just the 230 SGs.

II. MAGNETIC BAND STRUCTURES

Let us begin by reviewing some background materials concerning electronic band structures arising from a magnetic material. There are in total 1651 MSGs and 528 MLGs [29, 30, 46–48]. Among the 1651 MSGs, 230 of them are identical to SGs in which only unitary spatial symmetries are considered (type I MSGs). All other MSGs have an equal number of unitary and anti-unitary elements: $\mathcal{M} = \mathcal{G} + \mathcal{A}$. The unitary part \mathcal{G} is identical to one of the 230 SGs and the antiunitary part \mathcal{A} can be generally written in the form $\tilde{\mathcal{T}}\mathcal{G}$, where $\tilde{\mathcal{T}} \equiv \mathcal{T}g_0$ is the product of a spatial operation g_0 and timereversal (TR) \mathcal{T} . When g_0 belongs to \mathcal{G} , the MSG is simply the direct product of a SG \mathcal{G} and $\mathbb{Z}_2^{\mathcal{T}}$. This leads to another 230 MSGs, one for each SG, that are called type II. When g_0 is not an element of \mathcal{G} , there are two types further differentiated by whether g_0 is a pure translation (type IV) or not (type III). For type II-IV MSGs, the little group of k and the sitesymmetry group of x may also have an anti-unitary part, in addition to the usual unitary part. Note that, in the literature (e.g. [49]), our definition of type III and IV MSGs are sometimes called type IIIa and IIIb respectively. In addition, there are two common labeling schemes for MSGs: Opechowski-Guccione (OG) and Belov-Neronova-Smirnova (BNS). In this work we follow the BNS notation, where an MSG is labeled by a pair of integers written as S.L, with S corresponding to one of the 230 SG and L an extra label to differentiate between different magnetic descendants. (Please refer to, e.g., Ref. 50, for the precise meaning of these numbers.)

Next, we introduce a general formalism for the efficient analysis of band structure properties based on representations of the little group [13, 14, 35]. Our main focus here is to address how the formalism developed for SGs can be readily applied to MSGs. We define a band structure (BS) as a set of bands isolated from others by band gaps above and below at all high-symmetry momenta, where by a high-symmetry momentum we refer to a k in the Brillouin zone at which the unitary part of the little group, \mathcal{G}_k , is strictly larger than the translation subgroup T. In 3D, they can be high-symmetry points, lines or planes. A BS can be characterized by the set of nonnegative integers $n = \{n_k^{\alpha}\}$ that count the number of times an irreducible representation (irrep) $u_{\mathbf{k}}^{\alpha}$ of $\mathcal{G}_{\mathbf{k}}$ appears. As we are interested in systems of spinful electrons, the irreps $u_{\mathbf{k}}^{\alpha}$ here are generally projective due to the spin-1/2 nature of electrons (also called 'double-valued'). Since $\{n_{k}^{\alpha}\}$ cannot be changed smoothly without gap closing or symmetry breaking, $\boldsymbol{n} = \{n_{\boldsymbol{k}}^{\alpha}\}$ serves as 'topological invariants' defined for each BS.

The integers $\boldsymbol{n} = \{n_{\boldsymbol{k}}^{\alpha}\}$ cannot be chosen freely, since symmetries demand that they satisfy a collection of compatibility relations [10, 29]. Since \mathcal{G} is a subgroup of $\mathcal{M} = \mathcal{G} + \mathcal{A}$, the full list of compatibility relation C imposed on an Msymmetric BS can be split into two sets, $C_{\mathcal{G}}$ arising from \mathcal{G} and $\tilde{\mathcal{C}}_{\mathcal{A}}$ from \mathcal{A} . Let us denote by $\{BS\}_{phys}^{\mathcal{G}}$ and $\{BS\}_{phys}$ the set of all n's satisfying $C_{\mathcal{G}}$ and C, respectively. Here, the subscript 'phys' indicates that all n_{k}^{α} 's in $n = \{n_{k}^{\alpha}\}$ are nonnegative, which is required for interpreting them as the multiplicities of irreps in a physical band structure. We will introduce another set $\{BS\}^{(G)}$ that relaxes this nonnegative condition later. Note that $\{BS\}_{phys}^{\mathcal{G}}$ and $\{BS\}_{phys}$ differ only in the imposition of $\tilde{\mathcal{C}}_{\mathcal{A}}$. In general, the anti-unitary part \mathcal{A} requires a pairing of $\boldsymbol{b} \in \{BS\}_{phys}^{\mathcal{G}}$ with another $\boldsymbol{b}' \in \{BS\}_{phys}^{\mathcal{G}}$, unless \boldsymbol{b} itself is already symmetric under $\tilde{\mathcal{T}}$. The pairing type can be easily determined using the Herring rule [29]. (In Appendix D, we provide a more elaborated review on the compatibility relations and the Herring rule.)

III. BAND TOPOLOGY

Having described some generalities about BSs, we now review how knowledge about the real space can inform band topology [8, 14]. We define the trivial class of BSs by the AIs, which are band insulators that are smoothly connected to a limit of vanishing hopping, and hence are deformable to product states in real space. Equivalently, an AI admits symmetric, exponentially localized Wannier functions.

To specify an AI, one should choose a position x in real space at which electrons are localized, and the type of the orbital put on that site. All inequivalent choices of the position x are classified by Wyckoff positions [51]. The orbital can be chosen from the (co-)irreps of the site-symmetry group of x (Appendix E). Given these choices, an \mathcal{M} -invariant AI can be constructed by placing a symmetry-related orbital on each site of the \mathcal{M} -symmetric lattice and filling them by electrons. The AI has a specific combination of irreps in the momentum space, which automatically satisfies $\mathcal{C} = \mathcal{C}_{\mathcal{G}} + \tilde{\mathcal{C}}_{\mathcal{A}}$. We list up all distinct n's corresponding to an AI by varying x and the orbital type. We list up all distinct n's corresponding to an AI by varying the position x and the orbital type and we obtain $\{AI\}_{phys}$, a subset of $\{BS\}_{phys}$. If one replaces \mathcal{M} above with \mathcal{G} , one gets the set of \mathcal{G} -symmetric AIs, $\{AI\}_{phys}^{\mathcal{G}}$.

Now we are ready to tell which elements of $\{BS\}_{phys}$ must be topologically nontrivial and which elements can be trivial. This can be judged by contrasting the elements of $\{BS\}_{phys}$ with those in $\{AI\}_{phys}$. Namely, any $\boldsymbol{b} \in \{BS\}_{phys}$ not belonging to $\{AI\}_{phys}$ necessarily features nontrivial band topology, because, by definition, there does not exist any atomic limit of the BS with the same combinations of irreps. This is a sufficient (but not necessary) condition to be topologically nontrivial — here we exclusively focus on the band topology that can be diagnosed by the set of irreps at highsymmetry momenta.

The simplest way in exploring the nontrivial elements of $\{BS\}_{phys.}$ is thus to consider the complement of $\{AI\}_{phys.}$ in $\{BS\}_{phys.}$, as in Refs. 14 and 52. However, this set has a complicated mathematical structure. To simplify the analysis, we allow for the formal *subtraction* of bands, and extend the values of n_k^{α} to any integer, including the negative ones, à *la* a K-theory analysis. $\{BS\}_{phys}$ then becomes an abelian group $\{BS\} = \mathbb{Z}^{d_{BS}}$ (known as a 'lattice' in the mathematical nomenclature) [13, 14, 35]. In other words, there are d_{BS} basis 'vectors' $\{b_i\}_{i=1}^{d_{BS}}$, and $\{BS\}$ can be expressed as $\{\sum_{i=1}^{d_{BS}} m_i b_i \mid m_i \in \mathbb{Z}\}$. Similarly, by allowing negative integers when taking superposition of AIs, we get another abelian group $\{AI\} = \mathbb{Z}^{d_{AI}}$, which is a subgroup of $\{BS\}$. The band topology we are interested in is now encoded in the quotient group

$$X_{\rm BS} \equiv \{\rm BS\}/\{\rm AI\},\tag{1}$$

dubbed the symmetry-based indicator of band topology [14]. As we will see shortly, the quotient group is always a finite abelian group and hence must be a product of the form $\prod_i \mathbb{Z}_{n_i}$.

IV. CONSTRUCTING BAND STRUCTURES FROM ATOMIC INSULATORS

To compute X_{BS} , the natural first step is to identify C_A , the extra compatibility relations enforced by the anti-unitary symmetries. Contrary to this expectation, we now show that, based on our earlier results on SGs, one can directly compute $\{BS\}$ and X_{BS} for any MSG \mathcal{M} without deriving \tilde{C}_A . This serves to demonstrate the power of the present approach: symmetry content and connectivity of BSs can be readily extracted without the large overheads mandated by the conventional approach.

To this end, we first revisit the relevant aspects of the theory for an SG \mathcal{G} . By definition, $\{AI\}^{\mathcal{G}}$ is a subgroup of $\{BS\}^{\mathcal{G}}$, and therefore, a priori, it could be the case that $d_{AI}^{\mathcal{G}} < d_{BS}^{\mathcal{G}}$ strictly. However, by an explicit computation for all the 230 SGs, Ref. 14 found that $d_{BS}^{\mathcal{G}} = d_{AI}^{\mathcal{G}}$ always holds. This statement has an important implication — every $\boldsymbol{b} \in \{BS\}^{\mathcal{G}}$ can, in fact, be expanded as $\boldsymbol{b} = \sum_{i} q_{i} \boldsymbol{a}_{i}$ with *rational* coefficients $q_{i} \in \mathbb{Q}$, using a basis $\{\boldsymbol{a}_{i}\}_{i=1}^{d_{AI}^{\mathcal{G}}}$ of $\{AI\}^{\mathcal{G}}$. In other words, full knowledge on the group $\{BS\}^{\mathcal{G}}$ can be obtained from that on $\{AI\}^{\mathcal{G}}$.

Based on this result for SGs, we will now prove the same statement, namely $d_{\rm BS} = d_{\rm AI}$, for any MSG \mathcal{M} . This result enables us to efficiently compute {BS} and $X_{\rm BS}$ for all MSGs and MLGs using only information contained in {AI}, which can be readily extracted from the tabulated Wyckoff positions [46–48].

Our proof is centered on the following observation: Recall $\mathcal{M} = \mathcal{G} + \mathcal{A}$, where \mathcal{G} is unitary and $\mathcal{A} = \tilde{\mathcal{T}}\mathcal{G}$ is the antiunitary part generated by $\tilde{\mathcal{T}} = \mathcal{T}g_0$ for some spatial symmetry g_0 . Note that for type II MSGs, $g_0 \in \mathcal{G}$ and can be chosen to be the identity, whereas for types III and IV, $g_0 \notin \mathcal{G}$. Now consider a \mathcal{G} -symmetric collection of fully-filled local orbitals in real space, which defines an AI $a \in {\text{AI}}_{\text{phys}}^{\mathcal{G}}$. This AI is not generally symmetric under \mathcal{M} , i.e., it may not be invariant under the action of $\tilde{\mathcal{T}}$. However, if we stack it together with its $\tilde{\mathcal{T}}$ -transformed copy, we will arrive at an \mathcal{M} -symmetric AI. Algebraically, this means $a + \tilde{\mathcal{T}}a \in {\text{AI}}_{\text{phys}}$.

In the momentum space, a similar symmetrization procedure can be performed on the representation content. Suppose that $\{|\mathbf{k}, i\rangle\}$ is a basis of an irrep $u_{\mathbf{k}}^{\alpha}$ of $\mathcal{G}_{\mathbf{k}}$. Then the $\tilde{\mathcal{T}}$ transformed copy, $\{\tilde{\mathcal{T}}|\mathbf{k}, i\rangle\}$, forms a basis of an irrep $u_{\mathbf{k}'}^{\alpha'}$ of $\mathcal{G}_{\mathbf{k}'}$ (Appendix D). When **b** represents a BS that contains $u_{\mathbf{k}}^{\alpha'}$, times, we denote by $\tilde{\mathcal{T}}\mathbf{b}$ a BS that contains $u_{\mathbf{k}'}^{\alpha'}$ the same number of times. For any $\mathbf{b} \in \{BS\}^{\mathcal{G}}, \mathbf{b} + \tilde{\mathcal{T}}\mathbf{b}$ satisfies the compatibility conditions $\tilde{\mathcal{C}}_{\mathcal{A}}$, and hence belongs to $\{BS\}$. We present the explicit form of $\tilde{\mathcal{T}}\mathbf{b}$ in Appendix F.

We are now ready to prove the statement. Observe any $B \in \{BS\}$ also belongs to $\{BS\}^{\mathcal{G}}$, and as $d_{BS}^{\mathcal{G}} = d_{AI}^{\mathcal{G}}$, it can be expanded in the basis of $\{AI\}^{\mathcal{G}}$:

$$\boldsymbol{B} = \sum_{i} q_i \boldsymbol{a}_i, \quad q_i \in \mathbb{Q}.$$
 (2)

Now we symmetrize both sides of Eq. (2) by $\tilde{\mathcal{T}}$. Since B is \mathcal{M} -invariant, $B + \tilde{\mathcal{T}}B = 2B$, so

$$\boldsymbol{B} = \frac{1}{2} (\boldsymbol{B} + \tilde{\mathcal{T}} \boldsymbol{B}) = \sum_{i} \frac{1}{2} q_i (\boldsymbol{a}_i + \tilde{\mathcal{T}} \boldsymbol{a}_i).$$
(3)

TABLE I. Characterization of band structures (BSs) in a magnetic space group (MSG); excerpt from Tables III-VIII.

MSG	a	$d^{\mathbf{b}}$	$X_{\rm BS}{}^{\rm c}$	$\nu_{\rm BS}$
2.4	Ι	9	(2, 2, 2, 4)	1
2.7	IV	5	(2)	2
3.4	IV	3	(2)	2
209.51	IV	3	(1)	2^*

- ^a MSG number in the Belov-Neronova-Smirnova notation, followed by a Roman numeral I, . . . , IV indicating its type.
- ^b Number of linearly independent BSs.
- ^c Symmetry-based indicator of band topology, which takes the form
- $\prod_i \mathbb{Z}_{n_i}$; denoted by the collection of positive integers (n_1, n_2, \cdots) .
- ^d For most of the MSGs, the set of physical BS fillings $\{\nu\}_{BS}$ and the set of AI fillings $\{\nu\}_{AI}$ agree with each other, and they take the form $\{\nu\}_{BS} = \{\nu\}_{AI} = \nu_{BS} \mathbb{N}$. The asterisk indicates violation to this rule,

TABLE II. Characterization of magnetic layer groups through the corresponding magnetic space group (MSG); excerpt from Table IX.

MSG ^a	$d^{\mathbf{b}}$	$X_{\rm BS}{}^{\sf b}$	$\nu_{\rm BS}{}^{\rm b}$
2.5 (1) II	5	(2)	2
2.5 (2) II	5	(2)	2
2.5 (3) II	5	(2)	2
3.4 (1) IV	3	(2)	2
3.4 (2) IV	2	(1)	2

^a The numbers in parentheses label the different ways to project the MSG down to 2D planes (Appendix G).

^b Defined as in Table I.

detailed in Table X.

As argued, $a_i + \tilde{\mathcal{T}} a_i \in \{AI\}$. (Note that linearity was invoked when we extend the argument form $\{AI\}_{phys}^{\mathcal{G}}$ to a general element of $\{AI\}^{\mathcal{G}}$.) This proves that any $B \in \{BS\}$ can be expanded in terms of $\{AI\}$ (using rational coefficients), an equivalent statement of $d_{AI} = d_{BS}$. Furthermore, it implies that the quotient group $X_{BS} = \{BS\}/\{AI\}$ does not contain any \mathbb{Z} -factor, and hence is a finite abelian group of the form $\prod_i \mathbb{Z}_{n_i}$.

To summarize, we have shown that the set of AIs and BSs are identical as far as their dimensionality goes. This is a powerful statement, since it means we could simply focus on AIs, study their symmetry representations in momentum space, and then take rational combinations to generate all BSs. We caution, however, that one has to properly rescale the entries of n when an irrep is paired with another copy of itself according to the Herring rule. For full details of the treatment, we refer the interested readers to Ref. 14.

Following this strategy, we perform the first calculation of $d_{\rm BS}$, $X_{\rm BS}$, and $\nu_{\rm BS}$ for all of the 1651 MSGs and 528 MLGs. The full list of the computation results are tabulated in Tables III–IX in Appendix A. For readers' convenience, we reproduce a few examples from these tables in Tables I and II.

V. REPRESENTATION-ENFORCED SEMIMETALS

As an application of our results, here we introduce new classes of representation-enforced semimetals (reSM). A BS is said to be a reSM when (i) there are gap closings at some generic (i.e., not high-symmetry) momenta in the Brillouin zone, and (ii) the gaplessness is mandated by the combination of irreps at high-symmetry momenta. Note that only unitary symmetries are incorporated in defining a high-symmetry momentum, and the notion of a BS only requires a continuous gap at all high-symmetry points, lines and planes. ReSMs arise when the topological band-gap closings are buried in the low-symmetry regions of the Brillouin zone, rendering them hard to diagnose in conventional electronic structure calculations, but are nonetheless readily detectable and are robust against numerical uncertainty due to their representationenforced nature. All previously known examples of reSMs concern TR-broken systems in 3D with either the inversion [35, 41] or $\overline{4}$ rotation symmetry [42]. The new classes we propose here are realized in spin-orbit coupled systems in 2D or 3D, and are associated to type III or IV MSGs.

Our example is the type IV MSG 3.4 (P_a 112) is generated by C_{2z} (the π rotation about z-axis) and $\tilde{\mathcal{T}} \equiv \mathcal{T}T_{\frac{1}{2}a_1}$ (half translation by $\frac{1}{2}a_1$ followed by TR) apart from lattice translations. (The type III MSG 13.69 can also host a Dirac semimetal through a nearly identical mechanism. See Appendix H.) According to Table I, this MSG has two classes of combinations of C_{2z} eigenvalues, as indicated by $X_{\rm BS} = \mathbb{Z}_2$. As we shall see now, the 3D band structure belonging to this nontrivial class hosts at least four Weyl points, where the Weyl points with opposite charities are maximally separated in the k_3 direction.

To understand this band topology step by step, let us first remove the lattice translation in z and take a single layer parallel to the xy plane. The MSG then reduces to the MLG p_a 112, which corresponds to the entry 3.4 (1) in Table II with $X_{\rm BS} = \mathbb{Z}_2$. The rotation C_{2z} is a symmetry at the four TR-invariant momenta of the 2D BZ, whose eigenvalues are either $\pm i$ for spinful electrons. When $k_1 = 0$, the antiunitary symmetry $\tilde{\mathcal{T}}$ with $\tilde{\mathcal{T}}^2 = (-1)^{\hat{F}} \hat{T}_{a_1} = -1$ demands the pairing of the two eigenvalues of C_{2z} . In contrast, when $k_1 = \pi, \tilde{\mathcal{T}}^2 = +1$ and no pairing is required. For this MLG, $X_{\rm BS} = \mathbb{Z}_2$ and the topological index distinguishing the two class is given by

$$\eta = \prod_{n:\text{occupied}} \prod_{\boldsymbol{k}=(\pi,0),(\pi,\pi)} \eta_{n,\boldsymbol{k}},\tag{4}$$

where $\eta_{n,k}$ is the C_{2z} -eigenvalue of the *n*-th band. $\eta = +1$ for any AI, while $\eta = -1$ for nontrivial BSs. Note that this invariant is similar to those identified in some earlier works [35, 38, 41, 42, 53].

Next, we show that the nontrivial BS with $\eta = -1$ corresponds to 2D magnetic Dirac reSMs (two-fold degeneracy at the gapless point) enabled by a strong spin-rotation symmetry breaking, as illustrated in Fig. 1. Denoting the standard Pauli matrices by $\sigma_{0,...,3}$, the combined symmetry $\tilde{T}C_{2z}$ dictates that the local 2 by 2 Hamiltonian $h(\mathbf{k})$ near a gapless point



FIG. 1. Magnetic representation-enforced semimetal. (a) An example tight-binding model for representation-enforced semimetal. There are two sub-lattices per unit cell (shaded) due to an anti-ferromagnetic order m_x , producing in total four bands. On-site potential J stands for the exchange coupling $-Jm_x \cdot c_x^{\dagger} \sigma c_x$. In addition to the standard nearest-neighbor (NN) hopping t, a spin-dependent hopping $\pm t_J \sigma_x$ is included, which can be viewed as originating from an exchange coupling to a magnetic moment in the middle of the NN bonds (Appendix H). (b) The dispersion relation at J/t = 1 and $t_J/t = 1/4$. In this case, η in Eq. (4) is +1 and the dispersion is gapped between the second and the third band. (c) The dispersion relation at J/t = 1 and $t_J/t = 3/4$. Now $\eta = -1$ and a pair of Dirac nodes exists as predicted. (d) The Fermi surface of the 3D version of the representation-enforced semimetal. The two Weyl points on the $k_3 = 0$ ($k_3 = \pi$) plane have the chirality +1 (-1), indicating a huge Fermi arc on some 2D surfaces. The signs on the TRIMs indicate the product of the C_{2z} rotation eigenvalues of occupied bands in this model.

takes the form

$$h(\boldsymbol{k}) = \sum_{i=0}^{2} g_i(\boldsymbol{k})\sigma_i,$$
(5)

i.e., the σ_3 term is missing. Since there are the same number of the tunable parameters k_1 and k_2 as the number of the relevant coefficients $(g_1(\mathbf{k}) \text{ and } g_2(\mathbf{k}))$, the gaplessness is stable against perturbations respecting the symmetries [54]. To see this more explicitly, we note that the symmetry $\tilde{T}C_{2z}$ quantizes the Berry phase $B(k_1) = \int_{-\pi}^{\pi} dk_2 A_2(k_1)$ to be 0 or π at any k_1 . The topological index $\eta = -1$ dictates that $B(0) \neq B(\pi)$, implying the presence of a gap closing somewhere in $0 < k_1 < \pi$. Using a $\mathbf{k} \cdot \mathbf{p}$ analysis, one can confirm that the bands disperse linearly, leading to a Dirac node. We note that, compared to the protection of Dirac node by an eigenvalue exchange of nonsymmorphic symmetries [34, 55–57], the present Dirac node has a more topological character.

Given this understanding of the 2D layer with MLG 3.4, let us now recover the third dimension by stacking the copies

of 2D layers with generic inter-layer couplings. The resulting band structure, fully symmetric under MSG 3.4, host in total four Weyl points; two on the $k_3 = 0$ plane and the other two on the $k_3 = \pi$ plane, as depicted in Fig. 1d. Note that the two Weyl points on the $k_3 = 0$ plane are related by C_{2z} rotation (or, equivalently in this setup, \mathcal{T}) symmetry, and so must have the same chirality. Consequently, the two Weyl points on the $k_3 = \pi$ plane have the opposite chirality. This large separation of the Weyl points with opposite chiralities dictates the presence of long Fermi arcs on the surface. Let us stress again that this nontrivial band structure is indicated by the rotation eigenvalues at TRIMs, via the formula in Eq. (4) applied to the $k_3 = 0$ plane of the 3D band structure. This illustrates how the ideas described in the present work can guide the search for ideal magnetic Weyl semimetals. We leave a full analysis of magnetic reSMs to future works.



FIG. 2. **Magnetic filling-enforced semimetals.** (a,b) Symmetries of a magnetic order can prohibit atomic insulators (AIs) at odd site fillings. (a) The magnetic-point-group symmetries of a ferromagnetic arrangement is compatible with nondegenerate local energy levels. (b) Those of the depicted hedgehog defect, however, force all the energy levels to exhibit even degeneracies, which forbids AIs when a lone electron is localized to the purple site. (c) When hedgehog and anti-hedgehog defects are arranged into a diamond lattice, the previous argument suggests that no AI is allowed whenever the site fillings at the defect cores are odd. (d) The hypothetical magnetic structure in (c) could be realizable in spinel structures if the diamond sites are occupied by atoms with odd atomic numbers (purple), and the magnetic atoms (blue) at the pyrochlore sites exhibit an all-in-all-out magnetic order. (e–h) Filling-enforced semimetals arising form the magnetically-ordered Fu-Kane-Mele model, Eq. (6). (e) When the magnetically modulated hopping t_J is weak compared to the spin-orbit coupling λ , the fermiology is governed by rings of gap-closing (circled in red), growing out from the original Dirac points at X when $t_J = 0$. (f) The positions of the rings in the Brillouin zone are shown in red. (g,h) For $t_J \ge 2\sqrt{2}\lambda$, the nodal rings become connected at the momentum W. (Thin lines indicate copies of the gapless momenta in the repeated zone scheme, included to illustrate the connectivity of the rings.)

VI. FILLING-ENFORCED SEMIMETALS

As another application of our theory, we attack the following problem: Given an MSG \mathcal{M} , what are the electron fillings ν at which band insulators are allowed? One of the primary interests in solving this problem lies in the search of topological semimetals - if a material has an electron filling incompatible with any band insulator, it must possess symmetryprotected gaplessness near the Fermi energy, provided a bandtheory description is applicable. We refer to such systems as filling-enforced (semi-)metals. Once the MSG describing the magnetic order is identified, such systems can be readily predicted using our results on $\nu_{\rm BS}$. For stoichiometric compounds with perfect crystalline order and commensurate magnetism, one can further show that spatial symmetries quantize the physically allowed fillings, which in turns allows one to further diagnose the fermiology of the filling-enforced (semi-)metals. We relegate a detailed discussion to Appendix B, and summarize the results on the predicted fermiology in Tables XI-XIV.

While a systematic analysis of filling-enforced semimetals (feSMs) can be performed using the tables mentioned above,

it is instructive to discuss a physical picture linking a realspace description of the magnetic ordering to the momentumspace obstruction to realizing any band insulator. To this end, we first contrast the effect of topologically trivial and nontrivial magnetic orders on local electronic energy levels. When the surrounding magnetic moments order ferromagnetically, as in Fig. 2a, the electronic levels are Zeeman-split and nondegenerate. In contrast, when the moments are arranged into the hedgehog defect shown in Fig. 2b, the magnetic-point-group symmetries at the defect core demand that all single-particle local energy levels exhibit even degeneracies, i.e., such a magnetic order forbids a gapped state when a lone electron is localized to the defect core.

To be compatible with lattice translations, however, there must be a balance between defects and anti-defects in each unit cell. An interesting situation arises when the positions of the defects and anti-defects are further related by symmetries prohibiting them from annihilation. As an example, consider the diamond lattice with one sublattice occupied by hedgehog defects, and the other sublattice, symmetry-related to the first by a glide mirror, occupied by anti-hedgehogs (Fig. 2c). The symmetries of this system are described by MSG 227.131 (type III). Suppose the defect core (i.e., the diamond sites) corresponds to an atom with an odd atomic number, which leads to an electron filling of $\nu = 2 \mod 4$ per primitive unit cell. By the previous argument, it should be impossible to obtain an AI by localizing all the electrons. This suggests the electron filling might lead to an obstruction to forming a band insulator and hence enforcing a (semi-)metallic behavior. Indeed, our result of $\nu_{\rm BS} = 4$ for this MSG (Table VIII) implies all band insulators are ruled out at the specified filling of $\nu = 2$ mod 4. From Table XII, we further see that this corresponds to a feSM with movable nodal features.

Curiously, the hypothetical structure depicted in Fig. 2c could be relevant for materials with the chemical formula AB_2O_4 taking the spinel structure, where the A atoms, occupying the diamond sites, are surrounded by the B atoms sitting at the pyrochlore positions. If the B atoms are magnetically ordered into the all-in-all-out configuration (Fig. 2d), it can be viewed as a lattice realization of the described hedgehog-antihedgehog lattice in MSG 227.131. This suggests the following tight-binding model as an example of the feSM described:

$$\hat{H} = \sum_{\langle ij \rangle} \hat{c}_{i}^{\dagger} \left(t \, \sigma_{0} + 4 \, t_{J} \, \boldsymbol{d}_{ij} \cdot \boldsymbol{\sigma} \right) \hat{c}_{j} + i \, 8\lambda \sum_{\langle \langle ij \rangle \rangle} \hat{c}_{i}^{\dagger} \left(\boldsymbol{d}_{li} \times \boldsymbol{d}_{lj} \right) \cdot \boldsymbol{\sigma} \, \hat{c}_{j},$$
(6)

where \hat{c}_i and \hat{c}_i^{\dagger} represent the fermion operators corresponding to an *s*-orbital localized to the diamond site *i* (spin indices are suppressed), $\langle ij \rangle$ and $\langle \langle ij \rangle \rangle$ respectively denote first- and second-nearest-neighbor bonds, and d_{ij} is the vector connecting the site *j* to *i* (the lattice constant of the conventional cell is set to 1). In the term $\propto \lambda$, we let *l* denote the common nearest neighbor of *i* and *j*. Note that t_J can be physically interpreted as a modulation of the nearest-neighbor hopping by exchange coupling with the magnetic moment at pyrochlore sites, and λ parameterizes the symmetry-allowed spin-orbit coupling between the second-nearest-neighbor sites.

When $t_J = 0$, Eq. (6) is TR-symmetric (corresponding to the type II MSG 227.129), and reduces to the undistorted Fu-Kane-Mele model defined on the diamond lattice hosting three symmetry-related Dirac points (Table XI) [58]. Unlike lattice distortions, which lead to topological insulators [58], TR-symmetry breaking described by $t_J \neq 0$ leads to a feSM. Interestingly, the Dirac points in the original Fu-Kane-Mele model are unstable towards such perturbation, and our results indicate that the semimetallic behavior is enforced by a more delicate band connectivity (Table XII). This is explicitly verified in the example band structures plotted in Figs. 2e-h, showing how the Dirac points split into (doubly degenerate) nodal rings as t_J increases from 0.

This example serves only as a particular instance of the 421 MSGs we identified to be compatible with feSMs (Appendix B). A full list of such MSGs is presented in Tables XI and XII. Armed with this list, we studied the magnetic structures listed on the Bilbao Crystallographic Server [50], and found that the experimentally-characterized magnetic materials YFe_4Ge_2 and LuFe₄Ge₂ (and related compounds) in the type III MSG 58.399 are realistic magnetic feSM candidates

[59, 60]. For these compounds, a nonmagnetic atom with an odd atomic number (Lu or Y) sits at a maximal-symmetry site of multiplicity 2, whereas their surrounding magnetic moments (Fe) ordered into a pattern symmetric under the combined symmetry of spatial inversion and TR. Such a magnetic ordering falls into our broad description of nontrivial defect lattices, and these compounds are expected to feature Dirac points, pinned to high-symmetry momenta, near the Fermi energy (Table XI).

We remark that the method described here can be readily applied to any other commensurate magnetic crystals where both the chemical formula and the MSG describing the magnetic order have been identified. In particular, a local-moment description is inessential — the filling criterion established based on our computation of $\nu_{\rm BS}$ applies equally well to systems exhibiting itinerant magnetism, as long as a band-theory description remains applicable.

VII. INDICATOR FOR ANTIFERROMAGNETIC TOPOLOGICAL INSULATOR

As the last application of our results, let us make a connection of symmetry-based indicators of MSGs, fully computed in this work, to some previously studied topological insulators in TR broken settings.

As a canonical example, we discuss the type IV MSG 2.7 generated by the inversion symmetry I and the half translation followed by TR, $\tilde{T} \equiv TT_{\frac{1}{2}a_z}$. This MSG has two classes of the combination of inversion parities as indicated by $X_{BS} = \mathbb{Z}_2$ in Table I. The topological index distinguishing the two classes is given by

$$\xi = (-1)^{\frac{1}{4}\sum_{n:\text{occupied}}\sum_{\text{TRIMs}}\xi_{n,\boldsymbol{k}}} = \pm 1 \tag{7}$$

Here, $\xi_{n,k}$ is the parity of the *n*-th occupied band. To understand this formula, note first that even- and odd-parity bands form a Kramers pair at the TRIMs with $k_z = \pi$. This is because I and $\tilde{\mathcal{T}}$ anti-commute when $k_z = \pi$. Thus, the four TRIMs with $k_z = \pi$ never contribute to the index. On the other hand, when $k_z = 0$, I and $\tilde{\mathcal{T}}$ commute, and $\tilde{\mathcal{T}}^2 = -1$. This means that the same parity eigenvalue appears twice at TRIMs with $k_z = 0$. In Fig. 3a, we illustrated an example of the parity combination with $\xi = -1$.

As demonstrated in Ref. [71], the index $\xi = -1$ implies a nontrivial band topology, the so-called 'antiferromagnetic topological insulator' that supports gapless surface Dirac modes. The anti-ferromagnetic topological insulator can be understood as a staggered stacking of Chern insulators, in which a Chern insulator with C = +1 (described by red disks in Fig. 3b) is on every z = n ($n \in \mathbb{Z}$) plane and the one with C = -1 (blue disks) is on $z = n + \frac{1}{2}$ ($n \in \mathbb{Z}$) plane. All of these Chern insulators are related by the $\tilde{\mathcal{T}}$ symmetry.

If we introduce a perturbation that breaks the \hat{T} symmetry, the C = +1 Chern insulator at z = n ($n \neq 0$) and the C = -1 Chern insulator at $z = n \mp \text{sign}(n)\frac{1}{2}$ can pair annihilate with each other in an inversion symmetric manner, leaving only the C = +1 Chern insulator on the z = 0 plane



FIG. 3. Symmetry indicators for anti-ferromagnetic topological insulators. (a) An example of the inversion parity combination of valence bands in the \mathbb{Z}_2 nontrivial phase of MSG 2.7. (b) The realization of the \mathbb{Z}_2 nontrivial phase by staggered stacking of Chern insulators. The red (blue) disks represent a Chern insulator with C = +1 (C = -1). (c) Breaking the \tilde{T} -symmetry (the half translation in z followed by the time-reversal) leads to a 'higher-order' state with a 1D equatorial chiral mode on the surface [61–70].

as shown in Fig. 3c. Such one-dimensional chiral edge state on the surface of 3D topological insulators has been studied in Refs. 61 and 62, and their counterparts in many other symmetry settings have been discovered recently [63–70]. Assuming that the inversion symmetry is unbroken, we see that this phase can be diagnosed by another symmetry-based indicator for the type-I MSG 2.4, which is simply the SG with only the inversion *I* apart from the lattice translation. According to the Table I, this MSG has $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4$ classification, and the insulator hosting the 1D chiral edge mode constructed here belongs to the class (0, 0, 0, 2) in this classification.

VIII. CONCLUSION

In this work, we revisit the old problem of assigning a global irrep label to each connected branch of a band structure [10], generalizing our recent results on time-reversal symmetric systems [14] to all 1651 magnetic space groups. Our central results are the computation of three fundamental quantities associated with magnetic band structures, which are tabulated in Appendix A: (i) $d_{\rm BS}$, which characterizes the number of independent building blocks of energy bands; (ii) $X_{\rm BS}$, which, akin to the Fu-Kane parity criterion [38], serves as a symmetry-based indicator of band topology; and (iii) $\nu_{\rm BS}$, which dictates the electron fillings at which band insulators are possible. We further demonstrate the utility of these results by applying them to the study of topological semimetals, focusing on cases where the absence of a band gap is either diagnosed through the symmetry representations at isolated high-symmetry points, or is mandated by the electron filling. In particular, we identify the exhaustive list of magnetic space groups capable of hosting filling-enforced semimetals (Appendix **B**).

Although a full database on our computation of the basis 'vectors' of the groups {BS} and {AI} is not included, we note that such a database can be readily generated [14]: Our proof of $d_{\rm BS} = d_{\rm AI}$ for all magnetic space groups implies full knowledge on {BS} is encoded in that of {AI}, and the latter can be computed by analyzing, for instance, hypothetical structures constructed by combining irreps of the site-

symmetry groups to each of the Wyckoff positions. As long as all high-symmetry momenta are included in the analysis and the doubling of any irrep (according to the Herring rule) is incorporated, {BS} can be recovered by forming linear superpositions of the entries in {AI} using rational coefficients, subjected only to the constraint that the resultant irrep multiplicities must all be integer-valued. Such calculations can be performed without deriving the compatibility relations, and this circumvents the careful convention-fixing required in both tabulating and utilizing such results. In view of this, we refrain from providing a database on them.

We close by highlighting several interesting open questions that we leave for future studies: (i) performing a comprehensive study on the identification and characterization of representation-enforced semimetals in all 2D and 3D magnetic space groups; (ii) developing a theory of quantized physical responses, or a proof of the absence thereof, for each nontrivial class in the exhaustive list of $X_{\rm BS}$ we computed; and (iii) discovering realistic topological materials based on systematic searches using the diagnostics our theory provides. Finally, we remark that some aspects of our filling criterion generalize to the interacting setting [72], similar to the corresponding relation between the results for time-reversal symmetric crystals [7, 9].

ACKNOWLEDGMENTS

HW thank Shuichi Murakami and Chen Fang for useful discussion. AV and HCP were supported by NSF DMR-1411343. AV acknowledges support from a Simons Investigator Award. HW acknowledges support from JSPS KAKENHI Grant Number JP17K17678.

- M. Franz and L. Molenkamp, eds., *Topological Insulators*, Contemporary Concepts of Condensed Matter Science, Vol. 6 (Elsevier, 2013).
- [2] L. Michel and J. Zak, Phys. Rev. B **59**, 5998 (1999).
- [3] L. Michel and J. Zak, Euro Phys. Lett. 50, 519 (2000).
- [4] L. Michel and J. Zak, Phys. Rep. 341, 377 (2001).
- [5] S. A. Parameswaran, A. M. Turner, D. P. Arovas, and A. Vishwanath, Nat. Phys. 9, 299 (2013).
- [6] R. Roy, arXiv:1212.2944.
- [7] H. Watanabe, H. C. Po, A. Vishwanath, and M. P. Zaletel, Proc. Natl. Acad. Sci. U.S.A. 112, 14551 (2015).
- [8] H. C. Po, H. Watanabe, M. P. Zaletel, and A. Vishwanath, Sci. Adv. 2 (2016).
- [9] H. Watanabe, H. C. Po, M. P. Zaletel, and A. Vishwanath, Phys. Rev. Lett. 117, 096404 (2016).
- [10] L. P. Bouckaert, R. Smoluchowski, and E. Wigner, Phys. Rev. 50, 58 (1936).
- [11] C. Herring, J. Franklin Inst. 233, 525 (1942).
- [12] B. J. Wieder, Y. Kim, A. M. Rappe, and C. L. Kane, Phys. Rev. Lett. 116, 186402 (2016).
- [13] J. Kruthoff, J. de Boer, J. van Wezel, C. L. Kane, and R.-J. Slager, Phys. Rev. X 7, 041069 (2017).
- [14] H. C. Po, A. Vishwanath, and H. Watanabe, Nat. Commun. 8, 50 (2017).
- [15] B. Bradlyn, L. Elcoro, J. Cano, M. G. Vergniory, Z. Wang, C. Felser, M. I. Aroyo, and B. A. Bernevig, Nature 547, 298 EP (2017).
- [16] M. G. Vergniory, L. Elcoro, Z. Wang, J. Cano, C. Felser, M. I. Aroyo, B. A. Bernevig, and B. Bradlyn, Phys. Rev. E 96, 023310 (2017).
- [17] B. J. Wieder et. al., arXiv:1705.01617.
- [18] R. Yu et al., Science **329**, 61 (2010).
- [19] C.-Z. Chang et. al., Science 340, 167 (2013).
- [20] J. Ye et. al., Phys. Rev. Lett. 83, 3737 (1999).
- [21] Y. Taguchi et. al., Science 291, 2573 (2001).
- [22] M. Lee, Y. Onose, Y. Tokura, and N. P. Ong, Phys. Rev. B 75, 172403 (2007).
- [23] B. Binz and A. Vishwanath, Physica B 403, 1336 (2008).
- [24] A. Neubauer et. al., Phys. Rev. Lett. 102, 186602 (2009).
- [25] S. Nakatsuji, N. Kiyohara, and T. Higo, Nature 527, 212 (2015).
- [26] A. K. Nayak et. al., Sci. Adv. 2 (2016).
- [27] H. Yang et. al., New J. Phys. 19, 015008 (2017).
- [28] A. Fert, V. Cros, and J. Sampaio, Nat Nano 8, 152 (2013).
- [29] C. J. Bradley and A. P. Cracknell, *The Mathematical Theory of Symmetry in Solids* (Oxford University Press, 1972).
- [30] M. I. Aroyo et. al., Acta Cryst. A 62, 115 (2006).
- [31] C. Gong et. al., Nature 546, 265 (2017).
- [32] B. Huang et. al., Nature 546, 270 (2017).
- [33] R. Chen, H. C. Po, J. B. Neaton, and A. Vishwanath, Nature Physics 14, 55 EP (2017).
- [34] S. M. Young and B. J. Wieder, Phys. Rev. Lett. 118, 186401 (2017).
- [35] A. M. Turner, Y. Zhang, R. S. K. Mong, and A. Vishwanath, Phys. Rev. B 85, 165120 (2012).
- [36] D. J. Thouless, J. Phys. C: Solid State Phys. 17, L325 (1984).
- [37] T. Thonhauser and D. Vanderbilt, Phys. Rev. B 74, 235111 (2006).
- [38] L. Fu and C. L. Kane, Phys. Rev. B 76, 045302 (2007).
- [39] A. M. Turner, Y. Zhang, and A. Vishwanath, Phys. Rev. B 82, 241102 (2010).

- [40] X. Wan, A. M. Turner, A. Vishwanath, and S. Y. Savrasov, Phys. Rev. B 83, 205101 (2011).
- [41] T. L. Hughes, E. Prodan, and B. A. Bernevig, Phys. Rev. B 83, 245132 (2011).
- [42] C. Fang, M. J. Gilbert, and B. A. Bernevig, Phys. Rev. B 86, 115112 (2012).
- [43] D. S. Freed and G. W. Moore, Annales Henri Poincaré 14, 1927 (2013).
- [44] N. Read, Phys. Rev. B 95, 115309 (2017).
- [45] K. Shiozaki, M. Sato, and K. Gomi, Phys. Rev. B 95, 235425 (2017).
- [46] D. B. Litvin, Acta Cryst. A 57, 729 (2001).
- [47] D. B. Litvin, Magnetic Group Tables: 1-, 2- and 3-Dimensional Magnetic Subperiodic Groups and Magnetic Space Groups (International Union of Crystallography, 2013).
- [48] H. T. Stokes, M. H. Dorian, and B. J. Campbell, ISOTROPY Software Suite, iso.byu.edu .
- [49] R.-X. Zhang and C.-X. Liu, Phys. Rev. B 91, 115317 (2015).
- [50] S. V. Gallego et. al., J. Appl. Cryst. 49, 1750 (2016).
- [51] T. Hahn, ed., *International Tables for Crystallography*, 5th ed., Vol. A: Space-group symmetry (Springer, 2006).
- [52] H. C. Po, H. Watanabe, and A. Vishwanath, arXiv:1709.06551
- [53] Y. Kim, B. J. Wieder, C. L. Kane, and A. M. Rappe, Phys. Rev. Lett. 115, 036806 (2015).
- [54] J. Ahn and B.-J. Yang, Phys. Rev. Lett. 118, 156401 (2017).
- [55] S. M. Young and C. L. Kane, Phys. Rev. Lett. 115, 126803 (2015).
- [56] J. Wang, Phys. Rev. B 95, 115138 (2017).
- [57] S. S. Tsirkin, I. Souza, and D. Vanderbilt, Phys. Rev. B 96, 045102 (2017).
- [58] L. Fu, C. L. Kane, and E. J. Mele, Phys. Rev. Lett. 98, 106803 (2007).
- [59] P. S. Papamantellos et. al., J. Mag. Mag. Mater. 236, 14 (2001).
- [60] P. S. Papamantellos, K. Buschow, and J. Rodríguez-Carvajal, J. Mag. Mag. Mater. 324, 3709 (2012).
- [61] M. Sitte, A. Rosch, E. Altman, and L. Fritz, Phys. Rev. Lett. 108, 126807 (2012).
- [62] F. Zhang, C. L. Kane, and E. J. Mele, Phys. Rev. Lett. 110, 046404 (2013).
- [63] W. A. Benalcazar, J. C. Y. Teo, and T. L. Hughes, Phys. Rev. B 89, 224503 (2014).
- [64] W. A. Benalcazar, B. A. Bernevig, and T. L. Hughes, Science 357, 61 (2017).
- [65] K. Hashimoto, X. Wu, and T. Kimura, Phys. Rev. B 95, 165443 (2017).
- [66] Z. Song, Z. Fang, and C. Fang, Phys. Rev. Lett. 119, 246402 (2017).
- [67] F. Schindler, A. M. Cook, M. G. Vergniory, Z. Wang, S. S. Parkin, B. A. Bernevig, and T. Neupert, arXiv preprint arXiv:1708.03636 (2017).
- [68] W. A. Benalcazar, B. A. Bernevig, and T. L. Hughes, ArXiv e-prints (2017), arXiv:1708.04230.
- [69] C. Fang and L. Fu, arXiv preprint arXiv:1709.01929 (2017).
- [70] J. Langbehn, Y. Peng, L. Trifunovic, F. von Oppen, and P. W. Brouwer, arXiv preprint arXiv:1708.03640 (2017).
- [71] R. S. K. Mong, A. M. Essin, and J. E. Moore, Phys. Rev. B 81, 245209 (2010).
- [72] H. Watanabe, Phys. Rev. B 97, 165117 (2018).

Appendix A: Tables for spinful electrons

Here we include the following tables for spinful electrons:

- Tables III–VIII: d, X_{BS} , and ν_{BS} for MSGs.
- Table IX: d, X_{BS} , and ν_{BS} for MLGs.
- Table X: $\{\nu\}_{BS}$ and $\{\nu\}_{AI}$ for exceptional MSGs.

Instructions on how to read the tables can be found in Tables I and II.

We remark that the results on d, $X_{\rm BS}$, and $\nu_{\rm BS}$ for types I and II MSGs, which account for 460 out of the 1651 entries, were already published in Ref. 14. For the readers' convenience, we reproduce them here alongside with the new results on the remaining 1191 type III and IV MSGs. Similarly, the data on exceptional filling patterns among the type II MSGs (i.e., for time-reversal symmetric systems) were reported in Refs. 8 and 9, but are also reproduced below.

TABLE III. Characterization of magnetic space groups (MSGs) in the triclinic family for spinful electrons.

MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MS	G	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MS	GG	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$
1.1 I	1	(1)	1	1.3	IV	1	(1)	2	2.5	II	9	(2, 2, 2, 4)	2	2.7 IV	5	(2)	2
1.2 II	1	(1)	2	2.4	Ι	9 ((2, 2, 2, 4)	1	2.6	III	1	(1)	2				

d: Rank of the band structure group {BS}

 $X_{\rm BS}$: Symmetry-based indicators of band topology

 $\nu_{\rm BS}$: Set of ν bands are symmetry-forbidden from being isolated by band gaps if $\nu \not\in \nu_{\rm BS} \mathbb{Z}$

MS	G	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MS	G	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$
3.1	Ι	5	(2)	1	7.24	Ι	1	(1)	2	10.47 IV	7	(2, 2)	2	13.70 IV	4	(2)	4
3.2	Π	1	(1)	2	7.25	Π	1	(1)	4	10.48 IV	6	(2)	2	13.71 IV	3	(2)	4
3.3	III	1	(1)	1	7.26	III	1	(1)	2	10.49 IV	8	(2)	2	13.72 IV	5	(2)	2
3.4	IV	3	(2)	2	7.27	IV	1	(1)	4	11.50 I	6	(2)	2	13.73 IV	3	(2)	4
3.5	IV	1	(1)	2	7.28	IV	1	(1)	2	11.51 II	5	(2, 2, 4)	4	13.74 IV	4	(2)	4
3.6	IV	3	(1)	2	7.29	IV	1	(1)	4	11.52 III	2	(1)	2	14.75 I	5	(2)	2
4.7	Ι	1	(1)	2	7.30	IV	1	(1)	4	11.53 III	1	(1)	2	14.76 II	5	(2, 4)	4
4.8	Π	1	(1)	4	7.31	IV	1	(1)	2	11.54 III	5	(2, 2, 4)	2	14.77 III	1	(1)	4
4.9	III	1	(1)	2	8.32	Ι	2	(1)	1	11.55 IV	3	(2)	4	14.78 III	1	(1)	4
4.10	IV	1	(1)	4	8.33	Π	1	(1)	2	11.56 IV	5	(2)	2	14.79 III	5	(2, 4)	2
4.11	IV	1	(1)	2	8.34	Ш	1	(1)	1	11.57 IV	3	(2)	4	14.80 IV	3	(2)	4
4.12	IV	1	(1)	2	8.35	IV	1	(1)	2	12.58 I	10	(2, 2)	1	14.81 IV	3	(2)	4
5.13	Ι	3	(1)	1	8.36	IV	1	(1)	2	12.59 II	7	(2, 2, 4)	2	14.82 IV	3	(2)	4
5.14	Π	1	(1)	2	9.37	Ι	1	(1)	2	12.60 III	1	(1)	2	14.83 IV	5	(2)	2
5.15	III	1	(1)	1	9.38	II	1	(1)	4	12.61 III	1	(1)	2	14.84 IV	3	(2)	4
5.16	IV	2	(1)	2	9.39	III	1	(1)	2	12.62 III	7	(2, 2, 4)	1	15.85 I	6	(2, 2)	2
5.17	IV	1	(1)	2	9.40	IV	1	(1)	2	12.63 IV	5	(2)	2	15.86 II	5	(2, 4)	4
6.18	Ι	3	(1)	1	9.41	IV	1	(1)	4	12.64 IV	5	(2)	2	15.87 III	1	(1)	2
6.19	Π	1	(1)	2	10.42	Ι	15	(2, 2, 2)	1	13.65 I	7	(2, 2)	2	15.88 III	2	(1)	2
6.20	III	1	(1)	1	10.43	II	9	(2, 2, 2, 4)	2	13.66 II	5	(2, 2, 4)	4	15.89 III	5	(2, 4)	2
6.21	IV	1	(1)	2	10.44	III	1	(1)	2	13.67 III	1	(1)	2	15.90 IV	4	(2)	2
6.22	IV	2	(1)	2	10.45	Ш	1	(1)	2	13.68 III	3	(1)	2	15.91 IV	3	(2)	4
6.23	IV	2	(1)	2	10.46	III	9	(2, 2, 2, 4)	1	13.69 III	5	(2, 2, 4)	2				

TABLE IV. Characterization of magnetic space groups (MSGs) in the monoclinic family for spinful electrons.

 \overline{d} : Rank of the band structure group {BS}

 $X_{\rm BS}$: Symmetry-based indicators of band topology

 $\nu_{\rm BS}$: Set of ν bands are symmetry-forbidden from being isolated by band gaps if $\nu \notin \nu_{\rm BS} \mathbb{Z}$

TABLE V. Characterization of magnetic space groups (MSGs) in the orthorhombic family for spinful electrons

16.1 1 1 2 2.3.5 III 3 1.1 </th <th>MSG</th> <th>d</th> <th>$X_{\rm BS}$</th> <th>$\nu_{\rm BS}$</th> <th>MSC</th> <th>3</th> <th>d</th> <th>$X_{\rm BS}$</th> <th>$\nu_{\rm BS}$</th> <th>MSG</th> <th>r</th> <th>d</th> <th>$X_{\rm BS}$</th> <th>$\nu_{\rm BS}$</th> <th>MSG</th> <th>d</th> <th>$X_{\rm BS}$</th> <th>$\nu_{\rm BS}$</th>	MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSC	3	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG	r	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$
16.2 II 1 <td>16.1 I</td> <td>1</td> <td>(1)</td> <td>2</td> <td>23.51</td> <td>III</td> <td>3</td> <td>(1)</td> <td>1</td> <td>29.101</td> <td>III</td> <td>1</td> <td>(1)</td> <td>4</td> <td>33.151 IV</td> <td>1</td> <td>(1)</td> <td>4</td>	16.1 I	1	(1)	2	23.51	III	3	(1)	1	29.101	III	1	(1)	4	33.151 IV	1	(1)	4
16.3 II 5 2 29.103 II 1 1 4 33.153 IV 1 1 4 16.4 IV 1 1 2 29.105 IV 1 1 33.155 IV 1 1 4 16.5 IV 1 1 2 29.105 IV 1 1 4 33.155 IV 1 1 4 16.6 IV 1 1 1 2 29.105 IV 1 1 3 1 1 2 29.107 IV 1 1 4 34.158 II 1 1 2 21.007 IV 1 1 4 34.158 II 1	16.2 II	1	(1)	2	23.52	IV	1	(1)	2	29.102	III	1	(1)	4	33.152 IV	1	(1)	4
164 IV 1 (1) 2 24.54 II 1 (1) 4 23.154 IV 1 (1) 4 165 IV 1 (1) 4 24.55 III 2 12.016 IV 1 (1) 4 34.155 II 1 (1) 2 29.106 IV 1 (1) 4 34.157 II 1 (1) 2 29.107 IV 1 (1) 4 34.157 II (1) 2 17.7 I 5 (2) 2 25.57 II (1) 2 29.100 IV 1 (1) 4 34.157 II (1) 2 14.10 1 10.12 2 34.161 IV 1 (1) 4 34.161 IV 1 1 4 34.161 IV 1 1 4 34.161 IV 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	16.3 III	5	(2)	1	24.53	Ι	4	(1)	2	29.103	III	1	(1)	4	33.153 IV	1	(1)	4
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	16.4 IV	1	(1)	2	24.54	Π	1	(1)	4	29.104	IV	1	(1)	4	33.154 IV	1	(1)	4
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	16.5 IV	1	(1)	4	24.55	III	2	(1)	2	29.105	IV	1	(1)	8	33.155 IV	1	(1)	4
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	16.6 IV	1	(1)	4	24.56	IV	3	(1)	2	29.106	IV	1	(1)	4	34.156 I	3	(1)	2
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	17.7 I	5	(1)	2	25.57	I	1	(1)	2	29.107	IV	1	(1)	4	34.157 II	1	(1)	4
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	17.8 II	1	(-) (1)	4	25.58	П	1	(1)	2	29.108	IV	1	(-) (1)	4	34.158 III	1	(-) (1)	2
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	17.9 III	1	(-) (1)	2	25.59	Ш	3	(1)	1	29,109	IV	1	(-) (1)	4	34.159 III	3	(-) (1)	2
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	17.10 III	3	(2)	2	25.60	Ш	5	(2)	1	29.110	IV	1	(1)	4	34.160 IV	$\frac{1}{2}$	(1)	4
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	17.11 IV	2	(-) (1)	4	25.61	IV	1	(-) (1)	4	30.111	I	3	(1)	2	34.161 IV	1	(1)	4
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	17.12 IV	5	(1)	2	25.62	IV	1	(1)	2	30 112	п	1	(1)	4	34 162 IV	2	(1)	4
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	17.12 IV	1	(1)	2	25.62	IV	1	(1)	4	30 113	ш	1	(1)	2	34 163 IV	3	(1)	1
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	17.13 IV	т २	(1)	4	25.05	IV	1	(1)	1	30 114	ш	1	(1)	2	34.164 IV	2	(2)	т 9
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	17.14 IV	2 2	(1)	1	25.65	IV	1	(1)	1	30 115	ш	1 2	(1)	2	35 165 I	2	(1)	2
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	17.15 IV	ე ე	(1)	- -	25.05	T	1	(1)	4 9	30.115	III IV	ე ე	(1)	4	35.165 I	1	(1)	2
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	10.10 I 19.17 П	1	(1)	4	20.00	п	1	(1)	4	30.117		2	(1)	4	35.100 II	1	(1)	∠ 1
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	10.17 II 10.10 III	1 9	(1)	4	20.07	ш	1	(1)	4 9	20 119		ა 1	(2)	4	25 169 III		(1)	1
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	10.10 III 10.10 III	ა 1	(2)	2	20.08	111 111	1	(1)	2	20,110		1	(1)	4	25 160 IV	4	(2)	1
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	18.19 III 18.20 IV	1	(1)		20.09	111 111	ა 1	(1)	2	20,120		ა ი	(1)	4	25 170 IV	1	(1)	4
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	18.20 IV	1	(1)	4	20.70		1	(1)	4	30.120		2	(1)	4	25.170 IV	2	(1)	2
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	18.21 IV	1	(1)	4	26.71	IV	1	(1)	4	30.121	11	2	(1)	4	35.171 IV	2	(1)	4
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	18.22 IV	2	(1)	4	26.72	IV	1	(1)	4	30.122	1	2	(1)	4	36.172 I	1	(1)	2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	18.23 IV	3	(1)	2	26.73	IV	1	(1)	2	31.123	l n	2	(1)	2	36.173 II	1	(1)	4
19.25 1 1 1 1 1 1 1 1 1 2 36.175 III 2 1 2 19.26 II 1 1 4 26.76 IV 1 1 4 31.126 III 2 36.175 III 1 1 2 19.27 III 1 1 4 26.77 IV 1 1 4 31.128 IV 2 36.175 IV 1 1 4 19.29 IV 1 1 4 27.79 II 1 1 4 31.129 IV 1 1 4 36.175 IV 1 1 4 36.175 IV 1 1 4 36.175 IV 1 1 4 31.128 IV 1 1 4 36.178 IV 1 1 4 31.130 IV 1 1 4 31.131 IV 1 1 4 31.132 IV 2 37.183 III 1	18.24 IV	3	(1)	2	26.74	IV	1	(1)	2	31.124	11	1	(1)	4	36.174 III		(1)	2
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	19.25 I	1	(1)	4	26.75	IV	1	(1)	4	31.125	111 	1	(1)	2	36.175 III	2	(1)	2
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	19.26 II	1	(1)	8	26.76	IV	1	(1)	4	31.126		2	(1)	2	36.176 III	1	(1)	2
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	19.27 III	1	(1)	4	26.77	IV	1	(1)	4	31.127	Ш	1	(1)	2	36.177 IV	1	(1)	2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	19.28 IV	1	(1)	4	27.78	Ι	1	(1)	2	31.128	IV	2	(1)	4	36.178 IV	1	(1)	4
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	19.29 IV	1	(1)	4	27.79	Π	1	(1)	4	31.129	IV	1	(1)	4	36.179 IV	1	(1)	2
20.31 I 3 (1) 2 27.81 III 5 (2) 2 31.131 IV 1 (1) 4 37.181 II 1 (1) 4 20.32 II 1 (1) 4 27.82 IV 1 (1) 2 31.132 IV 2 (1) 2 37.182 III 1 (1) 2 20.33 III 1 (1) 2 27.83 IV 1 (1) 4 31.134 IV 2 (1) 2 37.184 IV 1 (1) 2 20.35 IV 3 (1) 2 27.85 IV 1 (1) 4 32.135 I 3 (1) 2 37.185 IV 2 (1) 2 20.35 IV 3 (1) 2 28.87 I 4 (1) 2 32.135 II 1 (1) 2 38.187 I 1 (1) 2 28.87 II 1 1 1	19.30 IV	1	(1)	4	27.80	III	1	(1)	2	31.130	IV	1	(1)	4	37.180 I	2	(1)	2
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	20.31 I	3	(1)	2	27.81	III	5	(2)	2	31.131	IV	1	(1)	4	37.181 II	1	(1)	4
20.33 III 1 (1) 2 27.83 IV 1 (1) 4 31.133 IV 2 (1) 4 37.183 III 4 (2) 2 20.34 III 2 (1) 2 27.85 IV 1 (1) 4 31.134 IV 2 (1) 2 37.185 IV 2 (1) 2 20.35 IV 3 (1) 2 27.85 IV 1 (1) 4 32.135 I 3 (1) 2 37.186 IV 2 (1) 2 20.35 IV 3 (1) 2 28.87 I 4 (1) 2 32.135 II 1 (1) 2 38.187 I 1 (1) 2 38.187 I 1 (1) 2 38.187 I 1 (1) 2 38.189 III 1 (1) 2 38.189 III 1 (1) 2 38.189 III 1 (1) 2 38.189	20.32 II	1	(1)	4	27.82	IV	1	(1)	2	31.132	IV	2	(1)	2	37.182 III	1	(1)	2
20.34 III 2 27.84 IV 1 (1) 4 31.134 IV 2 37.184 IV 1 (1) 2 20.35 IV 3 (1) 2 27.85 IV 1 (1) 4 32.135 I 3 (1) 2 37.184 IV 1 (1) 4 20.36 IV 1 (1) 4 32.136 II 1 (1) 4 37.186 IV 2 (2) 4 20.36 IV 1 (1) 4 32.136 II 1 (1) 4 37.186 IV 2 (1) 2 20.37 IV 3 (1) 2 28.87 I 4 (1) 2 32.137 III 1 (1) 2 38.187 I 1 (1) 2 21.39 II 1 1 2 28.88 III 1 (1) 2 32.140 IV 2 (1) 1 1 1 1	20.33 III	1	(1)	2	27.83	IV	1	(1)	4	31.133	IV	2	(1)	4	37.183 III	4	(2)	2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	20.34 III	2	(1)	2	27.84	IV	1	(1)	4	31.134	IV	2	(1)	2	37.184 IV	1	(1)	2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	20.35 IV	3	(1)	2	27.85	IV	1	(1)	4	32.135	Ι	3	(1)	2	37.185 IV	2	(2)	4
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	20.36 IV	1	(1)	4	27.86	IV	1	(1)	4	32.136	II	1	(1)	4	37.186 IV	2	(1)	2
21.38 I 2 (1) 2 28.88 II 1 (1) 4 32.138 III 3 (2) 2 38.188 II 1 (1) 2 21.39 II 1 (1) 2 28.89 III 1 (1) 2 32.139 IV 1 (1) 4 38.189 III 2 (1) 1 21.40 III 4 (2) 1 28.90 III 2 (1) 2 32.139 IV 1 (1) 4 38.189 III 2 (1) 1 21.40 III 3 (1) 1 28.90 III 2 32.140 IV 2 38.191 III 3 (1) 1 21.41 IV 1 (1) 2 28.92 IV 4 (1) 2 32.143 IV 3 10 1 11 1) 2 38.191 IV 1 (1) 2 21.43 IV 2 (1) 2	20.37 IV	3	(1)	2	28.87	Ι	4	(1)	2	32.137	III	1	(1)	2	38.187 I	1	(1)	2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	21.38 I	2	(1)	2	28.88	Π	1	(1)	4	32.138	III	3	(2)	2	38.188 II	1	(1)	2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	21.39 II	1	(1)	2	28.89	III	1	(1)	2	32.139	IV	1	(1)	4	38.189 III	2	(1)	1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	21.40 III	4	(2)	1	28.90	III	2	(1)	2	32.140	IV	2	(1)	4	38.190 III	3	(1)	1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	21.41 III	3	(1)	1	28.91	III	3	(2)	2	32.141	IV	3	(1)	2	38.191 III	3	(1)	1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	21.42 IV	1	(1)	2	28.92	IV	4	(1)	2	32.142	IV	2	(1)	4	38.192 IV	1	(1)	2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	21.43 IV	2	(1)	2	28.93	IV	2	(1)	4	32.143	IV	3	(1)	4	38.193 IV	1	(1)	2
22.45 I 1 (1) 2 28.95 IV 2 (1) 4 33.145 II 1 (1) 8 39.195 I 1 (1) 2 22.46 II 1 (1) 2 28.96 IV 4 (1) 4 33.145 III 1 (1) 8 39.195 I 1 (1) 2 22.46 II 1 (1) 2 28.96 IV 4 (1) 4 33.146 III 1 (1) 4 39.196 II 1 (1) 4 22.47 III 3 (1) 1 28.97 IV 3 (1) 2 33.147 III 1 (1) 4 39.197 III 2 (1) 2 22.48 IV 1 (1) 2 28.98 IV 3 (1) 4 33.148 III 1 (1) 4 39.198 III 1 (1) 2 23.49 I 1 (1) 2 29.99 I 1 (1) 4 33.149 IV 1 (1) 4 39.199 III 3 (2) 2 23.49 I 1 <t< td=""><td>21.44 IV</td><td>2</td><td>(1)</td><td>2</td><td>28.94</td><td>IV</td><td>1</td><td>(1)</td><td>4</td><td>33.144</td><td>I</td><td>1</td><td>(1)</td><td>4</td><td>38.194 IV</td><td>1</td><td>(1)</td><td>4</td></t<>	21.44 IV	2	(1)	2	28.94	IV	1	(1)	4	33.144	I	1	(1)	4	38.194 IV	1	(1)	4
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	22.45 I	1	(1)	2	28.95	IV	2	(1)	4	33.145	II	1	(1)	8	39.195 I	1	(1)	2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	22.46 II	1	(1)	2	28.96	IV	4	(1)	4	33.146	III	1	(1)	4	39.196 II	1	(1)	4
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	22.47 III	3	(1)	1	28.97	IV	3	(1)	2	33.147	III	1	(1)	4	39.197 III	$ _2$	(1)	2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	22.48 IV	1	(1)	2	28.98	IV	3	(1)	4	33.148	III	1	(1)	4	39.198 III	1	(1)	2
	23.49 I	1	(1)	2	29.99	Ι	1	(1)	4	33.149	IV	1	(1)	4	39.199 III	3	(2)	2
25.50 II [1 (1) 2 [[29.100 II] 1 (1) 8 [[33.150 IV]] (1) 8 [[39.200 IV] 1 (1) 4	23.50 II	1	(1)	2	29.100	IJ	1	(1)	8	33.150	IV	1	(1)	8	39.200 IV	1	(1)	4

(Continued from the previous page)

39.201 IV	1	(1)	2	47.251	III	1	(1)	2	51.301	IV	4	(2)	4	54.351	IV	3	(2)	4
39.202 IV	1	(1)	2	47.252	III	15	(2, 2, 2)	1	51.302	IV	4	(2, 2)	4	54.352	IV	3	(2)	8
40.203 I	3	(1)	2	47.253	III	1	(1)	2	51.303	IV	5	(2)	4	55.353	Ι	7	(2, 2)	2
40.204 II	1	(1)	4	47.254	IV	5	(2, 2, 4)	4	51.304	IV	4	(2)	4	55.354	Π	5	(2, 4)	4
40.205 III	1	(1)	2	47.255	IV	5	(2, 4)	4	52.305	Ι	5	(2)	4	55.355	III	1	(1)	4
40.206 III	2	(1)	2	47.256	IV	5	(4)	4	52.306	Π	3	(4)	8	55.356	III	1	(1)	4
40.207 III	2	(1)	2	48.257	Ι	3	(2)	4	52.307	III	2	(1)	4	55.357	III	9	(2, 2, 2)	2
40.208 IV	3	(1)	2	48.258	Π	3	(2, 4)	4	52.308	Ш	1	(1)	4	55.358	Ш	5	(2)	2
40.209 IV	1	(1)	4	48.259	Ш	3	(1)	2	52.309	Ш	2	(1)	4	55.359	Ш	1	(1)	4
40.210 IV	3	(1)	2	48.260	Ш	5	(2)	2	52.310	Ш	4	(2)	4	55.360	IV	4	(2, 2)	4
41.211 I	2	(1)	2	48.261	Ш	1	(1)	4	52.311	Ш	4	(2)	4	55.361	IV	3	(2)	4
41.212 II	1	(1)	4	48.262	IV	2	(2)	4	52.312	Ш	3	(2)	4	55.362	IV	4	(2)	4
41.213 III	1	(1)	2	48.263	IV	3	(2)	4	52.313	ш	3	(1)	4	55.363	IV	7	(2,2)	2
41.214 III	1	(1)	2	48.264	IV	2	(2)	4	52.314	IV	3	(2)	4	55.364	IV	5	(2)	4
41.215 III	2	(2)	$\frac{1}{2}$	49.265	I	6	(2)	$\frac{1}{2}$	52.315	IV	4	(2)	4	56.365	I	3	(2)	4
41.216 IV	$\frac{1}{2}$	(1)	4	49.266	П	5	(2, 2, 4)	4	52.316	IV	3	(2)	8	56.366	П	3	(4)	8
41.217 IV	1	(1)	4	49.267	Ш	3	(1)	2	52.317	IV	4	(2)	4	56.367	ш	1	(1)	4
41 218 IV	2	(1)	2	49 268	ш	1	(1)	$\frac{1}{2}$	52.318	IV	3	(2)	4	56 368	ш	1	(1)	4
42.219 I	1	(1)	$\frac{-}{2}$	49.269	Ш	10	(2, 2)	$\frac{-}{2}$	52.319	IV	4	(2)	4	56.369	Ш	5	(2, 2)	4
42.220 II	1	(1)	$\frac{2}{2}$	49 270	ш	7	(2, 2) (2, 2)	$\frac{1}{2}$	52.320	IV	4	(2)	4	56 370	ш	3	(2, 2)	4
42.221 III	2	(1)	1	49 271	ш	1	(2, 2) (1)	4	53 321	T	9	(2)	2	56 371	ш	3	(1)	4
42.222 III	3	(2)	1	49.272	IV	3	(2)	4	53.322	П	5	(2, 4)	4	56.372	IV	$\frac{3}{2}$	(2)	8
42.223 IV	1	(1)	2	49 273	IV	6	(2)	2	53 323	ш	1	(1)	4	56 373	IV	2	(2)	4
43 224 I	2	(1)	2	49 274	IV	4	(2) (2)	4	53 324	ш	2	(1)	4	56 374	IV	3	(2)	4
43.221 I	1	(1)	4	49 275	IV	3	(2) (2)	4	53 325	ш	1	(1)	4	56 375	IV	3	(2)	4
43 226 III	1	(1)	2	49 276	IV	4	(2)	4	53 326	ш	5	(2)	2	56 376	IV	2	(2)	4
43 227 III	2	(1)	2	50 277	T	3	(2) (2)	4	53 327	ш	8	(2)	2	57 377	T	4	(2)	4
43 228 IV	1	(1)	4	50.277	п	3	(2, 4)	4	53 328	ш	6	(2, 2) (2, 2)	$\frac{2}{2}$	57 378	п	3	(2, 4)	8
44 229 T	1	(1)	2	50.279	ш	3	(2, 1) (1)	$\frac{1}{2}$	53 329	ш	2	(2, 2) (1)	4	57 379	ш	3	(2, 1) (1)	4
44 230 II	1	(1)	$\frac{2}{2}$	50.279	ш	3	(1) (1)	$\frac{2}{2}$	53 330	IV	5	(1) (2)	4	57 380	ш	1	(1)	4
44 231 III	2	(1)	1	50.281	ш	5	(2, 2)	$\frac{1}{2}$	53 331	IV	4	(2)	4	57 381	ш	1	(1)	4
44 232 III	3	(1)	1	50.282	ш	5	(2, 2)	2	53 332	IV	6	(2)	4	57 382	ш	4	(1) (2)	4
44 233 IV	1	(1)	4	50.202	ш	1	(1)	4	53 333	IV	5	(2)	4	57 383	ш	4	(2, 2)	4
44 234 IV	1	(1)	2	50.203	IV	3	(1) (2)	4	53 334	IV	8	(2)	2	57 384	ш	3	(2, 2) (2)	4
45 235 I	1	(1)	$\frac{2}{2}$	50 285	IV	2	(2)	4	53 335	IV	6	(2)	4	57 385	ш	2	(1)	4
45 236 II	1	(1)	4	50.205	IV	2	(2) (2)	8	53 336	IV	5	(2)	4	57 386	IV	2	(1) (2)	8
45 237 III	1	(1)	2	50.200	IV	3	(2) (2)	4	54 337	T	4	(2)	4	57 387	IV	3	(2)	4
45 238 III	3	(1) (2)	2	50.207	IV	2	(2) (2)	8	54 338	п	3	(2)	8	57 388	IV	4	(2)	4
45 239 IV	1	(1)	$\frac{2}{2}$	51 289	T	7	(2, 2)	$\frac{1}{2}$	54 339	ш	1	(2, 1) (1)	4	57 389	IV	4	(2)	4
45 240 IV	1	(1)	4	51 290	п	5	(2, 2) (2, 2, 4)	4	54 340	ш	2	(1)	4	57 390	IV	3	(2)	8
46 241 I	2	(1)	2	51.290	ш	1	(2, 2, 1) (1)	2	54 341	ш	1	(1)	1	57 391	IV	2	(2)	1
46.241 I	1	(1) (1)	7	51.201	ш	2	(1) (1)	$\frac{2}{2}$	54 342	ш	5	(1)	1	57 392	IV	2	(2)	-
46 243 III	1	(1)	2	51 293	ш	1	(1) (1)	4	54 343	ш	3	(2, 2)	4	58 393	T	8	(2)	2
46 244 III	2	(1) (1)	$\frac{2}{2}$	51 294	ш	7	(1)	2	54 344	ш		(2)	1	58 394	п	5	(2) (4)	4
46 245 III	2	(1) (1)	$\frac{2}{2}$	51 295	ш	6	(2, 2)	$\frac{2}{2}$	54 345	ш		(2, 2) (1)	1	58 395	ш	1	(1)	-
46.245 III	1	(1) (1)	2	51 296	ш	0	(2)	$\frac{2}{2}$	54 346	IV	1	(1)	4	58 306	ш	1	(1) (1)	4
46 247 IV	1 9	(1)	4 2	51.290	ш	3	(2, 2, 2) (1)	2	54 3/7	IV	1 1	$\binom{2}{(2)}$		58 307	ш	8	(1) (9-9)	т 9
46 248 IV	2	(1)	4	51 202	IV IV	7	(1)	$\frac{4}{2}$	54 3/9	IV	2	$\binom{2}{(2)}$		58 309	ш	5	(2, 2)	⊿ ?
47 240 IV	0	(1) (2) 2) 1)	+ 0	51 200	т v IV	2 1	(2,2)	1	54 240	IV	0 9	$\binom{2}{2}$		58 200	ш	1	(2) (1)	∠ ∕
+7.249 I	9	(2, 2, 2, 4) (2, 2, 2, 4)	2	51 200	IV IV	ы Л	(2)	4	54.349	IV IV		$\binom{2}{2}$		58 100	III IV		(1)	-± ∕I
+1.230 11	J	(2, 2, 2, 4)	4	51.300	1 V	4	(2, 2)	4	54.550	1 1	4	(4)	4	50.400	1 V	4	(4)	<u>4</u>

(Continued from the previous page)

58.401	IV	4	(2)	4	62.442	Π	3	(4)	8	65.483	III	1	(1)	2	69.524	III	9	(2,2)	1
58.402	IV	5	(2)	4	62.443	III	1	(1)	4	65.484	III	1	(1)	2	69.525	III	1	(1)	2
58.403	IV	5	(2)	4	62.444	Π	1	(1)	4	65.485	Ш	12	(2, 2, 2)	1	69.526	IV	5	(4)	2
58.404	IV	8	(2)	2	62.445	Π	2	(1)	4	65.486	Ш	10	(2, 2)	1	70.527	Ι	3	(2)	4
59.405	Ι	3	(2)	4	62.446	Ш	3	(2)	4	65.487	III	1	(1)	2	70.528	II	3	(4)	4
59.406	Π	3	(2, 4)	4	62.447	III	3	(2)	4	65.488	IV	4	(4)	4	70.529	III	2	(1)	2
59.407	Ш	2	(1)	2	62.448	III	4	(2)	4	65.489	IV	6	(2, 4)	2	70.530	III	4	(2)	2
59.408	III	1	(1)	4	62.449	III	1	(1)	4	65.490	IV	5	(4)	4	70.531	III	1	(1)	4
59.409	III	5	(2, 2)	2	62.450	IV	2	(2)	4	66.491	Ι	7	(2)	2	70.532	IV	2	(2)	4
59.410	III	4	(2)	2	62.451	IV	3	(2)	4	66.492	Π	5	(2, 4)	4	71.533	Ι	6	(4)	2
59.411	III	3	(1)	2	62.452	IV	2	(2)	8	66.493	III	2	(1)	2	71.534	II	6	(2, 4)	2
59.412	IV	3	(2)	4	62.453	IV	2	(2)	4	66.494	III	1	(1)	2	71.535	III	1	(1)	2
59.413	IV	2	(2)	8	62.454	IV	3	(2)	4	66.495	III	9	(2, 2)	2	71.536	III	9	(2, 2)	1
59.414	IV	2	(2)	8	62.455	IV	2	(2)	8	66.496	III	6	(2, 2)	2	71.537	III	1	(1)	2
59.415	IV	3	(2)	4	62.456	IV	3	(2)	4	66.497	III	1	(1)	4	71.538	IV	4	(4)	4
59.416	IV	2	(2)	4	63.457	Ι	5	(2)	2	66.498	IV	5	(2)	2	72.539	Ι	4	(2)	2
60.417	Ι	4	(2)	4	63.458	Π	4	(2, 4)	4	66.499	IV	4	(2)	4	72.540	II	4	(2, 4)	4
60.418	Π	3	(4)	8	63.459	III	2	(1)	2	66.500	IV	6	(2)	2	72.541	III	2	(1)	2
60.419	III	1	(1)	4	63.460	III	1	(1)	4	67.501	Ι	5	(2)	2	72.542	III	1	(1)	2
60.420	III	2	(1)	4	63.461	III	1	(1)	2	67.502	Π	5	(2, 2, 4)	4	72.543	III	7	(2, 2)	2
60.421	III	1	(1)	4	63.462	III	5	(2)	2	67.503	III	1	(1)	2	72.544	III	5	(2, 2)	2
60.422	III	3	(2)	4	63.463	III	6	(2, 2)	2	67.504	III	2	(1)	2	72.545	III	1	(1)	4
60.423	Π	3	(2)	4	63.464	Π	5	(2, 2)	2	67.505	Ш	7	(2, 2)	2	72.546	IV	4	(2)	2
60.424	III	4	(2, 2)	4	63.465	III	2	(1)	2	67.506	III	8	(2, 2)	2	72.547	IV	3	(2)	4
60.425	III	2	(1)	4	63.466	IV	5	(2)	2	67.507	III	2	(1)	2	73.548	Ι	3	(2)	4
60.426	IV	3	(2)	8	63.467	IV	3	(2)	4	67.508	IV	3	(2)	4	73.549	II	3	(2, 4)	8
60.427	IV	2	(2)	8	63.468	IV	4	(2)	4	67.509	IV	5	(2)	2	73.550	III	1	(1)	4
60.428	IV	3	(2)	4	64.469	Ι	5	(2)	2	67.510	IV	3	(2)	4	73.551	III	4	(2, 2)	4
60.429	IV	2	(2)	8	64.470	Π	4	(2, 4)	4	68.511	Ι	3	(2)	4	73.552	III	4	(1)	4
60.430	IV	3	(2)	4	64.471	III	1	(1)	4	68.512	Π	3	(2, 4)	4	73.553	IV	3	(2)	4
60.431	IV	4	(2)	4	64.472	III	1	(1)	4	68.513	Ш	2	(1)	2	74.554	Ι	7	(2)	2
60.432	IV	4	(2)	4	64.473	Ш	1	(1)	4	68.514	III	2	(1)	2	74.555	II	5	(2, 4)	4
61.433	Ι	3	(2)	4	64.474	III	4	(2)	2	68.515	III	5	(2, 2)	2	74.556	III	1	(1)	2
61.434	Π	3	(4)	8	64.475	III	6	(2, 2)	2	68.516	Ш	4	(2, 2)	2	74.557	ш	1	(1)	4
61.435	III	1	(1)	8	64.476	III	5	(2, 2)	2	68.517	III	2	(1)	4	74.558	Ш	6	(2, 2)	2
61.436	III	3	(2)	4	64.477	III	2	(1)	4	68.518	IV	3	(2)	4	74.559	III	7	(2, 2)	2
61.437	III	1	(1)	8	64.478	IV	4	(2)	4	68.519	IV	3	(2)	4	74.560	III	2	(1)	2
61.438	IV	2	(2)	8	64.479	IV	3	(2)	4	68.520	IV	3	(2)	4	74.561	IV	5	(2)	4
61.439	IV	3	(2)	4	64.480	IV	5	(2)	2	69.521	Ι	6	(4)	2	74.562	IV	5	(2)	2
61.440	IV	2	(2)	8	65.481	Ι	8	(2, 4)	2	69.522	II	6	(2, 2, 4)	2					
62.441	Ι	3	(2)	4	65.482	II	$\overline{7}$	(2, 2, 4)	2	69.523	III	1	(1)	2					

				-			U		, , ,			U						
MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSC	ł	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSC	3	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG	r	d	$X_{\rm BS}$	$\nu_{\rm BS}$
75.1 I	8	(4)	1	84.51	Ι	13	(2, 4)	2	90.101	IV	4	(1)	2	97.151	Ι	2	(1)	2
75.2 II	3	(1)	2	84.52	II	7	(2, 4)	4	90.102	IV	4	(1)	2	97.152	Π	2	(1)	2
75.3 III	2	(1)	2	84.53	III	7	(2)	2	91.103	Ι	4	(1)	4	97.153	III	1	(1)	2
75.4 IV	3	(1)	2	84.54	III	1	(1)	4	91.104	Π	1	(1)	8	97.154	III	5	(2)	1
75.5 IV	4	(2)	2	84.55	III	5	(2)	2	91.105	III	3	(1)	4	97.155	III	1	(1)	2
75.6 IV	4	(1)	2	84.56	IV	6	(4)	4	91.106	III	1	(1)	4	97.156	IV	2	(1)	2
76.7 I	1	(1)	4	84.57	IV	6	(4)	4	91.107	III	2	(1)	4	98.157	Ι	2	(1)	4
76.8 II	1	(1)	8	84.58	IV	7	(4)	4	91.108	IV	4	(1)	4	98.158	Π	1	(1)	4
76.9 III	1	(1)	4	85.59	Ι	11	(2, 4)	2	91.109	IV	2	(1)	8	98.159	III	3	(1)	2
76.10 IV	1	(1)	4	85.60	II	6	(2, 4)	4	91.110	IV	3	(1)	4	98.160	III	2	(1)	2
76.11 IV	1	(1)	8	85.61	III	3	(2)	4	92.111	Ι	2	(1)	4	98.161	III	1	(1)	4
76.12 IV	1	(1)	4	85.62	III	4	(1)	2	92.112	II	1	(1)	8	98.162	IV	2	(1)	4
77.13 I	4	(2)	2	85.63	Ш	6	(2)	2	92.113	III	1	(1)	4	99.163	Ι	3	(1)	2
77.14 II	1	(1)	4	85.64	IV	5	(2)	4	92.114	III	1	(1)	4	99.164	Π	3	(1)	2
77.15 III	2	(1)	2	85.65	IV	6	(2)	4	92.115	III	2	(1)	4	99.165	III	2	(1)	2
77.16 IV	1	(1)	4	85.66	IV	5	(2)	4	92.116	IV	2	(1)	4	99.166	III	1	(1)	2
77.17 IV	2	(2)	4	86.67	Ι	9	(2,2)	2	92.117	IV	1	(1)	8	99.167	Ш	8	(4)	1
77.18 IV	2	(1)	4	86.68	П	5	(2, 4)	4	92.118	IV	2	(1)	4	99.168	IV	3	(1)	4
78.19 I	1	(1)	4	86.69	Ш	3	(2)	4	93.119	I	1	(1)	4	99.169	IV	$\frac{1}{2}$	(1)	4
78.20 II	1	(1)	8	86.70	Ш	2	(-) (1)	4	93.120	П	1	(-) (1)	4	99.170	IV	$\frac{1}{2}$	(1)	4
78.21 III	1	(1)	4	86.71	Ш	-	(2)	2	93.121	Ш	1	(1)	2	100.171	T	4	(1)	2
78 22 IV	1	(1)	4	86.72	IV	4	(-)	4	93 122	ш	4	(2)	2	100 172	п	2	(1)	4
78.22 IV	1	(1)	8	86 73	IV	5	(2)	4	93 122	ш	2	(2) (1)	2	100.172	ш	1	(1)	1
78.24 IV	1	(1)	4	86 74	IV	4	(2)	4	93 124	IV	1	(1)	1	100.173	ш	2	(1)	2
70.24 IV	5	(1) (2)	1	87 75	T	т 16	(2)	т 1	93 125	IV	1	(1)	1	100.174	III	5	(1)	2
79.25 T	5	(2) (1)	2	87.76	п	0	(4, 4) (2.8)	1 2	03 126	IV	1	(1)	4	100.175	IV	2	(4)	1
79.20 H	1	(1) (1)	2	87.70	ш	5	(2, 0)	2	04 127	T	1 9	(1)	4	100.170		1	(1)	+ 0
70.27 III	1 2	(1) (1)	2	87.78	ш	2	(4)	2	0/ 128	п	1	(1)	4	100.177	IV	4	(1)	1
80.20 T	2	(1) (1)	2	87.70	ш	5	(1)	2	0/ 120	ш	1 2	(1)	2	101 170	T	1	(1)	4
80.20 II	1	(1) (1)	4	87.80		0	$\binom{2}{4}$	2	04 130	ш	2	(1)	2	101.179	п	1	(1)	4
80.30 II 80.21 III	1	(1) (1)	4 9	07.00	T	0	(4)	⊿ ົ	04 121	m	1	(2)	4	101.100	ш	1	(1)	4
80.31 III 80.32 IV	1	(1) (1)	4	00.01	п	5	(2, 2)	4	04 122	III IV	1	(1)	4	101.101	111 111	1	(1)	4 9
00.32 IV	1 1 1	(1)	4	00.02	ш ш	0 9	(4)	4	94.132		1 9	(1)	4	101.102	111		(1)	2
от.ээ т 91 24 т	12	(2, 2, 4)	1	00.03	ш	ა 1	(2)	4	94.155		2	(1)	4	101.105		4	(2)	2
01.34 II 01.25 III	0	(2)	2	00.04	ш	1 C	(1)	4 0	94.134	1 V		(1)	4	101.104		1	(1)	4
01.55 III 01.26 IV	4	(1)	∠ ົ	00.05		4	$\binom{2}{2}$	4	95.155	п	4	(1)	4	101.105		1	(1)	4
01.30 IV	5	(2)	2	00.00	1 V T	4	(2)	4	95.150	ш	1	(1)	0	101.100	1 V	1	(1)	0
01.37 IV	0 C	(2, 2)	2	09.07	п	ა ი	(1)	2	95.157	111 111	ა 1	(1)	4	102.187	п	1	(1)	4
01.30 IV	11	(2)	2 1	09.00	ш	ა 1	(1)	2	95.150	111 111	1	(1)	4	102.188	ш	1	(1)	4
62.39 I	11	(2, 2, 2)	1	09.09	ш	1	(1)	1	95.159			(1)	4	102.189	111	1	(1)	4
82.40 II	5	(2)	2	89.90	ш	8	(4)	1	95.140	IV	4	(1)	4	102.190	III	2	(1)	2
82.41 III	1	(1)	2	89.91	III	2	(1)	2	95.141	IV	2	(1)	8	102.191	III	3	(2)	2
82.42 IV	5	(2)	2	89.92	IV	3	(1)	2	95.142	IV	3	(1)	4	102.192	IV		(1)	8
83.43 1	24	(4, 4, 4)	1	89.93	IV	2	(1)	4	96.143	1	2	(1)	4	102.193	IV	2	(1)	4
83.44 II	13	(2, 4, 8)	2	89.94	IV	2	(1)	4	96.144	11	1	(1)	8	102.194	IV	2	(1)	4
83.45 III	8	(2, 4)	2	90.95	I	4	(1)	2	96.145	III	1	(1)	4	103.195	1	3	(1)	2
83.46 III	3	(1)	2	90.96	II	2	(1)	4	96.146	III	1	(1)	4	103.196	II	3	(1)	4
83.47 III	5	(2)	2	90.97	III	2	(1)	2	96.147	III	2	(1)	4	103.197	III	2	(1)	4
83.48 IV	11	(4)	2	90.98	Ш	5	(4)	2	96.148	ÍV	2	(1)	4	103.198	III	1	(1)	4
83.49 IV	12	(4, 4)	2	90.99	III	1	(1)	4	96.149	IV	1	(1)	8	103.199	III	8	(4)	2
83.50 IV	12	(4)	2	90.100	IV	2	(1)	4	96.150	IV	2	(1)	4	103.200	IV	3	(1)	2

TABLE VI. Characterization of magnetic space groups (MSGs) in the tetragonal family for spinful electrons

(Continued from the previous page)

103.201 IV	2(1) 4	111.251	Ι	5	(2)	2	117.301	III	1	(1)	4	124.351	Ι	11	(4)	2
103.202 IV	2(1) 4	111.252	Π	5	(2)	2	117.302	III	2	(1)	2	124.352	Π	8	(2, 8)	4
104.203 I	4 (1) 2	111.253	Ш	2	(1)	2	117.303	III	7	(2, 2, 4)	2	124.353	III	3	(1)	2
104.204 II	2(1) 4	111.254	Ш	1	(1)	2	117.304	IV	3	(2)	4	124.354	III	5	(4)	4
104.205 III	1(1) 4	111.255	ш	12	(2, 2, 4)	1	117.305	IV	6	(2)	2	124.355	ш	4	(4)	4
104.206 III	2(1) 4	111.256	IV	3	(2)	4	117.306	IV	4	(2)	4	124.356	ш	3	(2)	4
104.207 III	5(2)2	111.257	IV	3	(2)	4	118,307	T	6	(2)	$\frac{1}{2}$	124.357	ш	16	(-)	2
104 208 IV	2(1) 4	111 258	IV	3	(2)	4	118 308	п	3	(2)	4	124 358	ш	3	(2)	4
104.209 IV	$\frac{2}{4}$ (2) 4	112 259	T	5	(2)	2	118 309	ш	1	(1)	4	124.359	ш	3	(1)	4
104.209 IV	$\frac{1}{4}(1)$	112.259	п	3	(2) (2)	1	118 310	ш	2	(1)	2	124 360	IV	11	(1)	2
105.211 I	1(1) 2 1(1) 4	112.200	ш	2	(1)	2	118 311	ш	7	(2, 2, 2)	2	124.361	IV	6	(1)	1
105.211 I	1(1) + 1(1) + 1(1)	112.201	ш	1	(1)	4	118 312	IV	י 2	(2, 2, 2)	1	124.301		7	(4)	1
105.212 II	1(1) + 0(1) =	112.202	ш	2	(1)	4 9	110.512			(2)	4	124.502	T	6	(4)	4
105.213 III	$\frac{2}{1}$ (1) $\frac{2}{1}$	112.203	m N/		(2, 2, 2)	2	110.313	11	4	(2)	4	125.303	п	C C	$\binom{2}{2}$	4
105.214 III	1(1) 4	112.204		0 9	(2)		110.314	1 V T		(2)	2	125.304	ш ш	4	(2, 4)	4
105.215 III	4(2) 2	112.205)) ဂ	(2)	4	119.515	і п		(2)	2	125.505	ш	4	(1)	4
105.216 IV	1(1)4	112.200	1	3	(2)	4	119.310	ш ш	0 1	(2)	2	125.300	111 111	ა ი	(2)	4
105.217 IV	1(1)8	113.207	і п	0	(2)	2	119.517	111 111	1	(1)	2	125.307	111 111	ა ი	(2)	4
105.218 IV	1(1)4	113.268	ш	3	(2)	4	119.318	111	1	(1)	2	125.368	ш	3 11	(2)	4
106.219 I	2(1) 4	113.269	ш		(1)	4	119.319	III III	11	(2, 2, 2)	1	125.369	ш	11	(2, 4)	2
106.220 II	1(1)8	113.270	ш	2	(1)	2	119.320	IV	3	(2)	4	125.370	III III	6	(2)	2
106.221 III	1(1)4	113.271	III	17	(2, 2, 4)	2	120.321	1	5	(2)	2	125.371	III	2	(1)	4
106.222 III	2(1) 4	113.272	IV	3	(2)	4	120.322	Ш	3	(2)	4	125.372	IV	4	(2)	4
106.223 III	3(2)4	113.273	IV	6	(2)	2	120.323	Ш	1	(1)	4	125.373	IV	6	(2)	4
106.224 IV	1 (1) 4	113.274	IV	4	(2)	4	120.324	III	1	(1)	2	125.374	IV	4	(2)	8
106.225 IV	2(1) 4	114.275	Ι	6	(2)	2	120.325	III	7	(2, 2, 2)	2	126.375	Ι	5	(2)	4
106.226 IV	2(1) 4	114.276	Π	3	(2)	4	120.326	IV	5	(2)	2	126.376	Π	4	(4)	4
107.227 I	2(1) 2	114.277	Ш	1	(1)	4	121.327	Ι	5	(2)	2	126.377	III	4	(1)	2
107.228 II	2(1) 2	114.278	Ш	2	(1)	4	121.328	Π	4	(2)	2	126.378	III	2	(2)	4
107.229 III	1(1)2	114.279	Ш	7	(2, 2, 2)	2	121.329	Ш	1	(1)	2	126.379	Ш	2	(2)	4
107.230 III	1 (1) 2	114.280	IV	3	(2)	4	121.330	III	1	(1)	2	126.380	III	3	(2)	4
107.231 III	5(2)1	114.281	IV	4	(2)	4	121.331	Ш	9	(2, 2, 2)	1	126.381	Ш	8	(2, 2)	2
107.232 IV	2(1) 4	114.282	IV	6	(2)	2	121.332	IV	4	(2)	2	126.382	III	4	(2)	4
108.233 I	2(1) 2	115.283	Ι	5	(2)	2	122.333	Ι	7	(2)	2	126.383	III	2	(1)	4
108.234 II	2(1) 4	115.284	Π	5	(2)	2	122.334	Π	3	(2)	4	126.384	IV	5	(2)	4
108.235 III	1 (1) 4	115.285	III	2	(1)	2	122.335	III	1	(1)	4	126.385	IV	4	(2)	8
108.236 III	1 (1) 2	115.286	Ш	1	(1)	2	122.336	III	2	(1)	4	126.386	IV	5	(2)	4
108.237 III	5(4)2	115.287	Ш	12	(2, 2, 4)	1	122.337	III	6	(2, 2)	2	127.387	Ι	12	(4, 4)	2
108.238 IV	2(1) 2	115.288	IV	3	(2)	4	122.338	IV	4	(2)	4	127.388	II	8	(4, 8)	4
109.239 I	1 (1) 4	115.289	IV	3	(2)	4	123.339	Ι	13	(2, 4, 8)	2	127.389	III	2	(1)	4
109.240 II	1 (1) 4	115.290	IV	3	(2)	4	123.340	Π	13	(2, 4, 8)	2	127.390	III	5	(2, 4)	4
109.241 III	2(1) 2	116.291	Ι	5	(2)	2	123.341	III	3	(1)	2	127.391	III	6	(2, 4)	2
109.242 III	1 (1) 4	116.292	II	3	(2)	4	123.342	III	8	(2, 4)	2	127.392	III	3	(2)	4
109.243 III	2(1) 2	116.293	III	2	(1)	4	123.343	III	7	(2, 2, 4)	2	127.393	III	15	(4, 4, 4)	2
109.244 IV	1 (1) 8	116.294	III	1	(1)	2	123.344	III	5	(2)	2	127.394	III	3	(2)	4
110.245 I	1 (1) 4	116.295	III	8	(2, 2, 2)	2	123.345	III	24	(4, 4, 4)	1	127.395	III	2	(1)	4
110.246 II	1 (1) 8	116.296	IV	5	(2)	2	123.346	III	5	(2)	2	127.396	IV	6	(4)	4
110.247 III	2(1) 4	116.297	IV	3	(2)	4	123.347	III	3	(1)	2	127.397	IV	12	(4, 4)	2
110.248 III	1 (1) 4	116.298	IV	3	(2)	4	123.348	IV	8	(2, 8)	4	127.398	IV	8	(4)	4
110.249 III	2(2) 4	117.299	Ι	6	(2)	2	123.349	IV	8	(4, 8)	4	128.399	Ι	12	(4)	2
110.250 IV	1 (1) 4	117.300	II	3	(2)	4	123.350	IV	7	(8)	4	128.400	II	7	(8)	4

(Continued from the previous page)

128.401 III	2	(1)	4	131.444 I	V	6	(2, 4)	4	135.487	III	4	(4)	4	138.530 T	V	3	(2)	8
128.402 III	4	(4)	4	131.445 T	v	4	(4)	8	135.488	III	3	(2)	4	139.531	[]	9	(8)	2
128.403 III	5	(4)	4	131.446 I	V	5	(4)	4	135.489	III	8	(2, 4)	4	139.532 I	I	9	(2, 8)	2
128.404 III	3	(2)	4	132.447	I	6	(4)	4	135.490	III	3	(2)	4	139.533 I	п	2	(1)	2
128.405 III	13	(4, 4)	2	132.448	I	6	(2, 4)	4	135.491	III	1	(1)	8	139.534 I	I	5	(4)	2
128.406 III	3	(2)	4	132.449 I	Π	1	(1)	4	135.492	IV	4	(4)	4	139.535 I	I	5	(4)	2
128.407 III	2	(1)	4	132.450 I	Π	5	(4)	4	135.493	IV	6	(4)	4	139.536 I	П	5	(2)	2
128.408 IV	7	(4)	4	132.451 I	II	5	(2)	2	135.494	IV	6	(4)	4	139.537 I	I	16	(4, 4)	1
128.409 IV	8	(4)	4	132.452 I	II	3	(2)	4	136.495	Ι	7	(4)	4	139.538 I	I	4	(2)	2
128.410 IV	12	(4)	2	132.453 I	Π	12	(2, 4)	2	136.496	Π	5	(4)	4	139.539 I	I	2	(1)	2
129.411 I	6	(2)	4	132.454 I	Π	5	(2)	2	136.497	Ш	1	(1)	4	139.540 Г	V	7	(8)	4
129.412 II	6	(2, 4)	4	132.455 I	Π	1	(1)	4	136.498	III	4	(4)	4	140.541	[]	8	(4)	2
129.413 III	2	(1)	4	132.456 I	V	6	(4)	4	136.499	III	6	(2)	2	140.542 I	I	7	(2, 8)	4
129.414 III	3	(2)	4	132.457 I	V	4	(4)	4	136.500	III	3	(2)	4	140.543 I	I	2	(1)	2
129.415 III	3	(2)	4	132.458 I	V	4	(4)	8	136.501	III	9	(2, 4)	2	140.544 I	I	4	(4)	4
129.416 III	6	(2)	2	133.459	I	4	(2)	4	136.502	III	3	(2)	4	140.545 II	I	4	(4)	2
129.417 III	11	(2, 4)	2	133.460 1	Ι	3	(4)	8	136.503	III	1	(1)	4	140.546 I	I	3	(2)	4
129.418 III	3	(2)	4	133.461 I	Π	2	(1)	4	136.504	IV	4	(4)	8	140.547 I	I	13	(4, 4)	2
129.419 III	4	(1)	2	133.462 I	Π	2	(2)	4	136.505	IV	6	(4)	4	140.548 I	I	4	(2)	2
129.420 IV	4	(2)	8	133.463 I	Π	2	(2)	4	136.506	IV	7	(4)	4	140.549 I	I	2	(1)	4
129.421 IV	6	(2)	4	133.464 I	Π	3	(2)	4	137.507	Ι	4	(2)	4	140.550 Г	V	8	(4)	2
129.422 IV	4	(2)	4	133.465 I	Π	6	(2, 2)	4	137.508	Π	4	(4)	4	141.551	[]	6	(2)	4
130.423 I	5	(2)	4	133.466 I	Π	4	(2)	4	137.509	Ш	1	(1)	4	141.552 I	I	5	(4)	4
130.424 II	4	(4)	8	133.467 I	Π	1	(1)	8	137.510	III	2	(2)	4	141.553 I	I	1	(1)	4
130.425 III	2	(1)	4	133.468 I	V	4	(2)	4	137.511	III	2	(2)	4	141.554 I	I	3	(2)	4
130.426 III	2	(2)	8	133.469 I	V	4	(2)	4	137.512	III	6	(2)	2	141.555 II	I	4	(2)	4
130.427 III	2	(2)	4	133.470 I	V	3	(2)	8	137.513	III	8	(2, 2)	2	141.556 I	I	6	(2)	2
130.428 III	4	(2)	4	134.471	I	5	(2)	4	137.514	III	3	(2)	4	141.557 I	I	8	(2,2)	2
130.429 III	8	(2, 4)	4	134.472 1	Ι	5	(2, 4)	4	137.515	III	2	(1)	4	141.558 I	I	3	(2)	4
130.430 III	3	(2)	4	134.473 I	Π	2	(1)	4	137.516	IV	3	(2)	8	141.559 I	I	1	(1)	4
130.431 III	4	(1)	4	134.474 I	Ш	3	(2)	4	137.517	IV	3	(2)	8	141.560 T	۷I	4	(2)	8
130.432 IV	5	(2)	4	134.475 I	П	3	(2)	4	137.518	IV	4	(2)	4	142.561		4	(2)	4
130.433 IV	4	(2)	4	134.476 I	Ш	3	(2)	4	138.519	I	5	(2)	4	142.562 I	I	3	(4)	8
130.434 IV	5	(2)	4	134.477 I	П	9	(2,2)	2	138.520	Π	4	(4)	8	142.563 I	Π	1	(1)	4
131.435 I	7	(2, 4)	4	134.478 I	Ш	6	(2)	2	138.521	Ш	1	(1)	4	142.564 I		2	(2)	8
131.436 II	7	(2, 4)	4	134.479 1		1	(1)	4	138.522	III	3	(2)	4	142.565		2	(2)	4
131.437 III	1	(1)	4	134.480 1	V	3	(2)	8	138.523	III	3	(2)	4	142.566 1		5	(2)	4
131.438 III	17	(2)	2	134.481 1	V	4	(2)	4	138.524	III	4	(2)	4	142.567 I		5	(2,2)	4
131.439 III		(2,4)	4	134.482 1	V	4	(2)	4	138.525	III	7	(2,2)	4	142.568 I		3	(2)	4
131.440 III	5	(2)	2	135.483		6	(4)	4	138.526	Ш	3	(2)	4	142.569 I		2	(1)	8
131.441 III	13	(2,4)	2	135.484		4	(4)	8	138.527		$ ^{2}_{4}$	(1)	4	142.570 T	v	4	(2)	4
131.442 III	3	(2)	4	135.485 I		1	(1)	4	138.528	IV	4	(2)	4					
131.443 III	1	(1)	4	135.486 I	П	4	(2)	4	138.529	١V	5	(2)	4					

$\frac{\text{MSG}}{d} \frac{d}{X_{\text{BS}}} \frac{\lambda_{\text{BS}}}{\nu_{\text{BS}}} \frac{\text{MSG}}{d} \frac{d}{X_{\text{BS}}} \frac{\lambda_{\text{BS}}}{\nu_{\text{BS}}}$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
$143.1 I 7 (3) 1 \ \ 156.51 III 7 (3) 1 \ \ 160.51 III 7 (3) 1 IIII 7 (3) 1 III 7 (3) III 7 (3) III 7 (3) IIII 7 (3) IIIII 7 (3) IIII 7 (3) IIII 7 (3) IIIII 7 (3) IIII 7 (3) IIIII 7 (3) IIIII 7 (3) IIII 7 (3) IIIII 7 (3) IIIIII 7 (3) IIIII 7 (3) I$	$66.101 \text{ III} \begin{vmatrix} 11 & (2,4) & 1 \\ 177.151 \text{ III} \begin{vmatrix} 5 & (1) & 1 \\ 1 & 1 \end{vmatrix}$
143.2 II 4 (1) 2 156.52 IV 4 (1) 2 16	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
143.3 IV 4 (1) 2 157.53 I 5 (1) 1 16	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
144.4 I 1 (1) 3 157.54 II 3 (1) 2 16	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
144.5 II 1 (1) 6 157.55 III 5 (3) 1 16	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
144.6 IV 1 (1) 6 157.56 IV 3 (1) 2 16	57.106 III 3 (1) 2 178.156 II 1 (1) 12
145.7 I 1 (1) 3 158.57 I 4 (1) 2 16	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
145.8 II 1 (1) 6 158.58 II 4 (1) 4 16	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
$145.9 \text{ IV} \begin{vmatrix} 1 & (1) & 6 \end{vmatrix} \\ 158.59 \text{ III} \begin{vmatrix} 7 & (3) & 2 \end{vmatrix} \\ 1600000000000000000000000000000000000$	58.109 I 9 (6) 1 178.159 III 1 (1) 6
146.10 I 3 (1) 1 158.60 IV 4 (1) 2 16	$58.110 \text{ II} \begin{vmatrix} 4 & (1) & 2 \end{vmatrix} 178.160 \text{ IV} \begin{vmatrix} 3 & (1) & 6 \end{vmatrix}$
146.11 II 2 (1) 2 159.61 I 4 (1) 2 166	$58.111 \text{ III} \begin{vmatrix} 4 & (1) & 1 \end{vmatrix} 179.161 \text{ I} \begin{vmatrix} 3 & (1) & 6 \end{vmatrix}$
$146.12 \text{ IV} \begin{array}{ c c c c c c c c c c c c c c c c c c c$	[58.112 IV] 4 (1) 2 179.162 II 1 (1) 12
147.13 I 13 (2,12) 1 159.63 III 5 (3) 2 16	59.113 I 1 (1) 6 179.163 III 2 (1) 6
147.14 II 9 $(2,4)$ 2 159.64 IV 3 (1) 2 16	$59.114 \text{ II} \begin{vmatrix} 1 & (1) & 12 \end{vmatrix} 179.164 \text{ III} \begin{vmatrix} 2 & (1) & 6 \end{vmatrix}$
147.15 III 4 (1) 2 160.65 I 3 (1) 1 16	$59.115 \text{ III} \begin{vmatrix} 1 & (1) & 6 \end{vmatrix} 179.165 \text{ III} \begin{vmatrix} 1 & (1) & 6 \end{vmatrix}$
147.16 IV 7 (2) 2 160.66 II 2 (1) 2 16	$59.116 \text{ IV} \begin{vmatrix} 1 & (1) & 6 \end{vmatrix} 179.166 \text{ IV} \begin{vmatrix} 3 & (1) & 6 \end{vmatrix}$
148.17 I 11 (2,4) 1 160.67 III 3 (1) 1 17	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
148.18 II 8 $(2,4)$ 2 160.68 IV 2 (1) 2 17	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
148.19 III 2 (1) 2 161.69 I 2 (1) 2 17	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
148.20 IV 6 (2) 2 161.70 II 2 (1) 4 17	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
149.21 I 6 (1) 1 161.71 III 3 (1) 2 17	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
149.22 II 4 (1) 2 161.72 IV 2 (1) 2 17	71.122 II 1 (1) 6 180.172 IV 1 (1) 6
149.23 III 7 (3) 1 162.73 I 12 (2) 1 17	71.123 III 1 (1) 3 181.173 I 1 (1) 6
149.24 IV 5 (1) 2 162.74 II 9 (2,4) 2 17	71.124 IV 1 (1) 6 181.174 II 1 (1) 6
150.25 I 6 (1) 1 162.75 III 4 (1) 2 17	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
150.26 II 3 (1) 2 162.76 III 3 (1) 2 17	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
150.27 III 5 (3) 1 162.77 III 13 (2,12) 1 17	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
150.28 IV 4 (1) 2 162.78 IV 7 (2) 2 17	2.128 IV 1 (1) 6 181.178 IV 1 (1) 6
151.29 I 3 (1) 3 163.79 I 8 (2) 2 17	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
151.30 II 1 (1) 6 163.80 II 6 (4) 4 17	73.130 II 3 (1) 4 182.180 II 3 (1) 4
151.31 III 1 (1) 3 163.81 III 4 (1) 2 17	$\textbf{73.131 III} \begin{vmatrix} 4 & (1) & 2 \end{vmatrix} \textbf{182.181 III} \begin{vmatrix} 4 & (1) & 2 \end{vmatrix}$
151.32 IV 2 (1) 6 163.82 III 4 (1) 2 17	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
152.33 I 3 (1) 3 163.83 III 9 (12) 2 174	74.133 I 21 (3,3,3) 1 182.183 III 5 (3) 2
152.34 II 1 (1) 6 163.84 IV 7 (2) 2 174	24.134 II 10 (3,3) 2 182.184 IV 5 (1) 2
152.35 III 1 (1) 3 164.85 I 12 (2) 1 174	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
152.36 IV 2 (1) 6 164.86 II 9 (2,4) 2 174	74.136 IV 11 (3) 2 183.186 II 4 (1) 2
153.37 I 3 (1) 3 164.87 III 3 (1) 2 17	75.137 I 27 (6,6,6) 1 183.187 III 5 (1) 1
153.38 II 1 (1) 6 164.88 III 4 (1) 2 17	75.138 II 14 (6,12) 2 183.188 III 4 (1) 1
153.39 III 1 (1) 3 164.89 III 13 (2,12) 1 17	25.139 III 10 (3,3) 2 183.189 III 9 (6) 1
153.40 IV 2 (1) 6 164.90 IV 7 (2) 2 17	25.140 III 4 (1) 2 183.190 IV 4 (1) 4
154.41 I 3 (1) 3 165.91 I 8 (2) 2 17	25.141 III 9 (2,4) 1 184.191 I 4 (1) 2
154.42 II 1 (1) 6 165.92 II 6 (4) 4 17	25.142 IV 13 (6) 2 184.192 II 4 (1) 4
154.43 III 1 (1) 3 165.93 III 3 (1) 2 17	26.143 I 16 (3,6) 2 184.193 III 4 (1) 2
154.44 IV 2 (1) 6 165.94 III 5 (1) 2 17	76.144 II 9 (12) 4 184.194 III 3 (1) 2
155.45 I 4 (1) 1 165.95 III 9 (12) 2 17	76.145 III 11 (3) 2 184.195 III 9 (6) 2
155.46 II 2 (1) 2 165.96 IV 7 (2) 2 17	76.146 III 3 (1) 2 184.196 IV 4 (1) 2
155.47 III 3 (1) 1 166.97 I 11 (2) 1 17	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
155.48 IV 3 (1) 2 166.98 II 8 (2,4) 2 17	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
156.49 I 5 (1) 1 166.99 III 2 (1) 2 17	77.149 I 4 (1) 2 185.199 III 5 (1) 2
156.50 II 4 (1) 2 166.100 III 2 (1) 2 17	$7.150 \text{ II} \begin{vmatrix} 4 & (1) & 2 \end{vmatrix} 185.200 \text{ III} \begin{vmatrix} 3 & (1) & 2 \end{vmatrix}$

TABLE VII. Characterization of magnetic space groups (MSGs) in the hexagonal family for spinful electrons

(Continued from the previous page)

																		_
185.201 III	5	(3)	2	188.219	III	14	(3, 3)	2	191.237	III	7	(3, 3)	2	193.255	III	3	(1)	2
185.202 IV	3	(1)	2	188.220	IV	12	(3)	2	191.238	III	12	(2)	1	193.256	ш	7	(3)	4
186.203 I	3	(1)	2	189.221	Ι	10	(3, 3)	2	191.239	III	12	(2)	1	193.257	III	8	(3)	2
186.204 II	3	(1)	4	189.222	II	7	(3,3)	2	191.240	III	27	(6, 6, 6)	1	193.258	ш	8	(2)	2
186.205 III	4	(1)	2	189.223	III	4	(1)	1	191.241	III	4	(1)	2	193.259	ш	8	(2)	2
186.206 III	4	(1)	2	189.224	III	5	(1)	1	191.242	IV	9	(12)	4	193.260	ш	14	(3, 6)	2
186.207 III	5	(3)	2	189.225	III	15	(3, 3, 3)	1	192.243	Ι	13	(6)	2	193.261	ш	4	(1)	2
186.208 IV	3	(1)	2	189.226	IV	6	(3)	2	192.244	Π	9	(12)	4	193.262	IV	10	(6)	2
187.209 I	10	(3,3)	2	190.227	I	12	(3)	2	192.245	III	4	(1)	2	194.263	I	10	(6)	2
187.210 II	10	(3,3)	2	190.228	II	6	(3)	4	192.246	III	8	(3)	2	194.264	п	9	(12)	4
187.211 III	6	(1)	1	190.229	III	3	(1)	2	192.247	III	7	(3)	2	194.265	ш	3	(1)	2
187.212 III	5	(1)	1	190.230	III	4	(1)	2	192.248	III	7	(2)	2	194.266	ш	11	(3)	2
187.213 III	21	(3, 3, 3)	1	190.231	III	12	(3,3)	2	192.249	III	7	(2)	2	194.267	ш	6	(3)	4
187.214 IV	7	(3)	2	190.232	IV	9	(3)	2	192.250	Ш	18	(6, 6)	2	194.268	ш	8	(2)	2
188.215 I	12	(3)	2	191.233	I	14	(6, 12)	2	192.251	III	4	(1)	4	194.269	ш	8	(2)	2
188.216 II	7	(3)	4	191.234	II	14	(6, 12)	2	192.252	IV	13	(6)	2	194.270	ш	16	(3, 6)	2
188.217 III	5	(1)	2	191.235	III	4	(1)	2	193.253	Ι	10	(6)	2	194.271	ш	4	(1)	2
188.218 III	4	(1)	2	191.236	III	10	(3,3)	2	193.254	II	8	(12)	4	194.272	IV	9	(6)	2

MSG	ł	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSC	ì	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG	ŕ	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG		d	$X_{\rm BS}$	$\nu_{\rm BS}$
195.1	Ι	3	(1)	2	206.39	III	3	(1)	4*	216.77	IV	4	(2)	4	224.115	IV	6	(2)	4
195.2	II	2	(1)	2	207.40	Ι	4	(1)	2	217.78	Ι	6	(2)	2	225.116	Ι	11	(8)	2
195.3	IV	2	(1)	4	207.41	Π	4	(1)	2	217.79	Π	5	(2)	2	225.117	Π	11	(8)	2
196.4	I	3	(1)	2	207.42	III	3	(1)	2	217.80	Ш	3	(1)	2	225.118	ш	5	(2)	2
196.5	II	2	(1)	2	207.43	IV	3	(1)	4	218.81	Ι	6	(2)	2^*	225.119	III	10	(4)	2
196.6	IV	2	(1)	2^{*}	208.44	Ι	2	(1)	4	218.82	II	4	(2)	4	225.120	III	3	(1)	2
197.7	Ι	3	(1)	2	208.45	II	2	(1)	4	218.83	III	3	(1)	4	225.121	IV	8	(8)	4
197.8	п	2	(1)	2	208.46	III	3	(1)	2^*	218.84	IV	4	(2)	4	226.122	Ι	10	(4)	2^*
198.9	I	3	(1)	4	208.47	IV	2	(1)	4	219.85	Ι	6	(2)	2^{*}	226.123	II	8	(8)	4
198.10	II	2	(1)	8	209.48	Ι	3	(1)	2	219.86	Π	4	(2)	4	226.124	III	5	(2)	2^*
198.11	IV	2	(1)	4*	209.49	II	3	(1)	2	219.87	III	3	(1)	4	226.125	Ш	$\overline{7}$	(4)	4
199.12	Ι	4	(1)	2^{*}	209.50	III	3	(1)	2	219.88	IV	6	(2)	2^{*}	226.126	Ш	3	(1)	4
199.13	II	2	(1)	4^*	209.51	IV	3	(1)	2^*	220.89	Ι	7	(2)	2^*	226.127	IV	10	(4)	2^*
200.14	Ι	11	(2, 4)	2	210.52	I	3	(1)	4	220.90	II	4	(2)	4^*	227.128	Ι	9	(2)	4
200.15	Π	8	(2, 4)	2	210.53	II	2	(1)	4	220.91	III	3	(1)	4*	227.129	II	8	(4)	4
200.16	III	2	(1)	2	210.54	Ш	3	(1)	4	221.92	Ι	14	(4, 8)	2	227.130	III	4	(2)	4
200.17	IV	6	(4)	4	210.55	IV	3	(1)	4^*	221.93	II	14	(4, 8)	2	227.131	III	9	(2)	4
201.18	Ι	9	(2)	4	211.56	Ι	3	(1)	2	221.94	III	6	(2)	2	227.132	III	2	(1)	4
201.19	Π	6	(2, 4)	4	211.57	Π	3	(1)	2	221.95	III	11	(2, 4)	2	227.133	IV	6	(2)	8
201.20	ш	2	(1)	4	211.58	Ш	3	(1)	2	221.96	III	4	(1)	2	228.134	Ι	7	(2)	4^*
201.21	IV	5	(2)	4	212.59	Ι	3	(1)	4	221.97	IV	8	(8)	4	228.135	Π	5	(4)	8
202.22	Ι	10	(4)	2	212.60	Π	2	(1)	8	222.98	Ι	8	(2)	4	228.136	III	4	(2)	4^*
202.23	II	7	(4)	2	212.61	Ш	3	(1)	4	222.99	II	6	(4)	4	228.137	III	6	(2)	8
202.24	ш	2	(1)	2	212.62	IV	3	(1)	4^*	222.100	III	4	(2)	4	228.138	Ш	3	(1)	8
202.25	IV	6	(4)	2^{*}	213.63	Ι	3	(1)	4	222.101	III	6	(2)	4	228.139	IV	7	(2)	4^*
203.26	Ι	9	(2)	4	213.64	Π	2	(1)	8	222.102	III	3	(1)	4	229.140	Ι	11	(8)	2
203.27	Π	6	(4)	4	213.65	III	3	(1)	4	222.103	IV	8	(2)	4	229.141	Π	11	(2, 8)	2
203.28	ш	2	(1)	4	213.66	IV	3	(1)	4^*	223.104	Ι	8	(4)	4	229.142	Ш	6	(2)	2
203.29	IV	5	(2)	4^{*}	214.67	Ι	2	(1)	4	223.105	Π	7	(4)	4	229.143	Ш	10	(4)	2
204.30	I	10	(4)	2	214.68	Π	2	(1)	4^*	223.106	III	6	(2)	2^{*}	229.144	Ш	3	(1)	2
204.31	Π	7	(2, 4)	2	214.69	Ш	4	(1)	2^*	223.107	III	7	(4)	4	230.145	Ι	7	(2)	4^*
204.32	ш	2	(1)	2	215.70	Ι	6	(2)	2	223.108	III	2	(1)	4	230.146	Π	5	(4)	8^*
205.33	Ι	9	(2)	4	215.71	Π	6	(2)	2	223.109	IV	7	(4)	4	230.147	Ш	5	(2)	4^*
205.34	Π	6	(4)	8	215.72	Ш	3	(1)	2	224.110	Ι	8	(2)	4	230.148	Ш	6	(2)	4^*
205.35	ш	2	(1)	8	215.73	IV	4	(2)	4	224.111	Π	8	(2, 4)	4	230.149	III	2	(1)	8
205.36	IV	5	(2)	8	216.74	Ι	6	(2)	2	224.112	III	4	(2)	4					
206.37	I	9	(2)	4	216.75	II	6	(2)	2	224.113	III	9	(2)	4					
206.38	II	6	(2, 4)	8	216.76	Ш	3	(1)	2	224.114	III	2	(1)	4					

TABLE VIII. Characterization of magnetic space groups (MSGs) in the cubic family for spinful electrons

 $\frac{1}{d: \text{Rank of the band structure group {BS}}}{X_{BS}: \text{Symmetry-based indicators of band topology}} \nu_{BS}: \text{Set of } \nu \text{ bands are symmetry-forbidden from being isolated by band gaps if } \nu \notin \nu_{BS} \mathbb{Z}$ *: Exhibiting exceptional filling pattern; see Table X

MSG $X_{\rm BS} \ \nu_{\rm BS}$ MSG MSG $d X_{\rm BS} \nu_{\rm BS}$ d $d X_{\rm BS} \nu_{\rm BS}$ MSG $d X_{\rm BS} \nu_{\rm BS}$ 16.5 1.1 (1) Ι 1 (1) 1 6.19(1) Π 1 (1) 2 11.52 (1) III 2 (1)2 IV 1 (1) 4 1.1(2)(1)6.19(2) II 2 11.52 (2) III 2 (1) 2 17.7 (1) 4 (1) 2 Ι 1 1 (1)Ι 1 2 1.1(3)Ι 1 (1)1 6.19 (3) II 1 (1)2 11.53 (1) III 1 (1) 2 17.7 (2) Ι 4 (1)1.2(1)Π 1 6.20(1) III 11.53 (2) III 1 (1) 2 17.8 (1) II 4 (1)2 1 (1)1 1 (1)1.2(2)II 1 (1)2 6.20(2) III 1 (1)1 11.54 (1) III 3 (2) 2 17.8 (2) II 1 (1)4 1.2 (3) II 1 6.20(3) III 11.54 (2) III 3 17.9 (1) III (1)2 1 (1)1 (2)2 1 (1)2 2 11.55 IV 2 (1) 17.9 (2) III 1 (1) 2 1.3 (1) IV 1 (1)2 6.21 (1) IV 1 (1)4 11.56 (1) IV 3 (1) 17.10 (1) III 3 (2) 2 1.3 (2) IV 1 (1)2 6.21 (2) IV 1 (1)2 2 2.4(1)Ι 5 (2)1 6.22(1) IV 2 (1)2 11.56 (2) IV 3 (1) 2 17.10 (2) III 2 (1) 2 2.4(2)5 6.22 (2) IV 2 2 11.57 IV 2 (1) 17.11 IV 2 (1) 4 Ι (2)1 (1)4 2.4 (3) Ι 5 (2)1 6.23 IV 2 (1)2 12.58 I 6 (1)17.12 (1) IV 4 (1)2 1 2.5 (1) II 2 12.59 II 4 2 17.12 (2) IV 4 2 5 (2)2 7.24(1) Ι 1 (1)(2)(1)2 2.5 (2) Π 5 (2)2 7.24(2) Ι 1 (1)2 12.60 III 1 (1)2 17.13 IV 3 (1) 3 (1) 2 2.5 (3) II 5 (2)2 7.25(1) II III 1 2 18.16 I 1 (1)4 12.61 (1)7.25 (2) II III 4 18.17 1 (1) 4 2.6 (1) III 1 (1)2 1 (1)4 12.62 (2)1 Π 2.6 (2) III 1 (1)2 7.26(1) III 1 (1)2 12.64 IV 3 (1)2 18.18 III 3 (2) 2 2.6 (3) III 1 2 7.26 (2) III 2 13.65 (1) I 2 18.19 Ш 2 (1)1 (1)4 (1)1 (1)2.7 (1) IV 3 (1)2 7.27 IV 1 (1)4 13.65 (2) I 5 (2)2 18.20 IV 2 (1)4 2.7 (2) IV 3 (1) 2 7.28(1) IV 2 13.66 (1) II 3 18.23 IV 3 (1) 2 1 (1)(2)4 2 (1) 3.1 (1) (1) 7.28 (2) IV 13.66 (2) II 3 21.38 2 Ι 3 1 1 (1)2 (2) 4 Ι 5 IV 13.67 (1) III 1 (1) 21.39 Π 1 2 3.1 (2) Ι (2)1 7.29 1 (1)4 2 (1)3.1 (3) I 3 (1) IV (1) 2 13.67 (2) III 1 (1) 21.40 III 4 (2) 1 7.31 1 2 1 3.2 (1) II | 1 (1)2 8.32 Ι 2 (1)1 13.68 (1) III 2 (1) 2 21.41 III 2 (1) 1 3.2 (2) II 1 (1)2 8.33 Π 1 (1)2 13.68 (2) III 3 (1) 2 21.43 IV 2 (1)2 III 13.69 (1) III 3 3.2 (3) II 1 (1)2 8.34 1 (1)1 (2)2 25.57 (1) I 1 (1)2 IV 2 2 3.3 (1) III 1 (1)1 (1)13.69 (2) III 3 (2) 2 25.57 (2) I 1 1 8.36 (1)3.3 (2) III 1 (1) 10.42 (1) I 9 (1)13.70 IV 3 (1) 4 25.57 (3) I 2 1 1 1 (1)3.3 (3) III 1 (1)1 10.42 (2) I 10 (2, 2) 1 13.71 IV 2 (1) 4 25.58 (1) II 1 (1) 2 10.42 (3) I 2 3.4 (1) IV 3 (2)2 9 (1)1 13.72 (1) IV 3 (1)2 25.58 (2) II 1 (1) 3.4 (2) IV 2 (1)2 10.43 (1) II 5 (2)2 13.72 (2) IV 3 (1)2 25.58 (3) II 1 (1)2 3.5 (1) IV (1)2 10.43 (2) II 5 (2)2 13.73 IV 2 (1) 4 25.59 (1) III 3 (1) 1 1 3.5 (2) IV 10.43 (3) II 2 14.75 I 2 25.59 (2) III 2 (1) 1 (1)2 5 (2) 3 (1)1 2 (1) 10.44 (1) III 2 II 3 25.59 (3) III 3 (1) 3.6 IV 2 1 14.76 (2)4 (1)1 25.60 (1) III 3 (1) 4.7 (1) Ι 1 (1) 2 10.44 (2) III (1)2 14.77 III 1 (1)1 4 1 4.7 (2) Ι 1 (1)2 10.44 (3) III 1 (1)2 14.78 III 1 (1)4 25.60 (2) III 3 (1) 1 10.45 (1) III 2 14.79 III 3 25.60 (3) III 5 (2) 4.8 (1) II 1 (1) 4 1 (1)(2)2 1 4.8 (2) II | 1 (1)4 10.45 (2) III 1 (1)2 14.81 IV 2 (1)4 25.61 (1) IV 1 (1) 4 4.9 (1) III 1 10.45 (3) III IV 2 (1) 25.61 (2) IV 1 (1) 4 (1)2 1 (1)2 14.82 4 4.9 (2) III 1 10.46 (1) III IV 3 (1) 25.62 (1) IV 1 (1) 2 (1)2 5 (2)14.83 2 1 4.10 IV 1 (1)4 10.46 (2) III 5 (2)1 16.1 (1) I 1 (1) 2 25.62 (2) IV 1 (1) 2 4.11 (1) IV 1 (1) 10.46 (3) III (2) 16.1 (2) I 2 25.63 IV 1 (1) 4 2 5 1 1 (1) 10.47 (1) IV 4.11 (2) IV 1 (1)2 5 (2)2 16.1 (3) I 1 (1)2 25.64 IV 1 (1) 4 4.12 IV 1 (1)2 10.47 (2) IV 4 (1)2 16.2 (1) II 1 (1) 2 26.66 (1) I 1 (1) 2 10.48 (1) IV 5.13 Ι 2 (1)1 4 (1)2 16.2 (2) II 1 (1)2 26.66 (2) I 1 (1)2 5.14 II 1 (1)2 10.48 (2) IV 4 2 16.2 (3) II 1 (1) 2 26.67 (1) II 1 (1) 4 (1)III | 1 (1)5.15 10.49 IV 5 2 16.3 (1) III 3 (1) 26.67 (2) II 1 (1) 4 1 (1)1 5.17 IV (1)2 11.50(1) I 4 2 16.3 (2) III 3 (1) 26.68 (1) III 1 (1) 2 1 (1)1 6.18 (1) I 3 (1) 1 11.50(2) I 4 (1)2 16.3 (3) III 5 (2) 1 26.68 (2) III 1 (1) 2 (2)26.69 (1) III 2 (1) 2 6.18 (2) I 2 (1)1 11.51 (1) II 3 4 16.4 (1) IV 1 (1) 2 6.18 (3) I 3 (1) 1 11.51 (2) II 3 (2)4 16.4 (2) IV 1 (1) 2 26.69 (2) III 3 (1) 2

TABLE IX. Characterization of the projections of magnetic space groups (MSGs) which correspond to magnetic layer group, assuming spinful electrons.

(Continued from the previous page)

26.70 (1) III	1 ((1) 2	2	30.118	IV	1	(1)	4	47.255	IV	3	(2)	4	51.302 IV 3 (2)	4
26.70 (2) III	1 ((1) 2	2	30.119	IV	2	(1)	2	49.265 (1)	Ι	4	(1)	2	51.303 IV 3 (1)	4
26.71 IV	1 ((1) 4	ı	31.123	Ι	2	(1)	2	49.265 (2)	Ι	4	(1)	2	53.321 I 6 (1)	2
26.72 IV	1 ((1) 4	ı	31.124	Π	1	(1)	4	49.266 (1)	Π	3	(2)	4	53.322 II 3 (2)	4
26.73 (1) IV	1 ((1) 2	2	31.125	ш	1	(1)	2	49.266 (2)	Π	3	(2)	4	53.323 III 1 (1)	4
26.73 (2) IV	1 ((1) 2	2	31.126	ш	2	(1)	2	49.267 (1)	III	3	(1)	2	53.324 III 2 (1)	4
26.74 IV	1 ((1) 2	2	31.127	Ш	1	(1)	2	49.267 (2)	III	2	(1)	2	53.325 III 1 (1)	4
26.75 IV	1 ((1) 4	ı	31.128	IV	2	(1)	4	49.268 (1)	III	1	(1)	2	53.326 III 3 (1)	2
27.78 (1) I	1 ((1) 2	2	31.130	IV	1	(1)	4	49.268 (2)	III	1	(1)	2	53.327 III 5 (1)	2
27.78 (2) I	1 ((1) 2	2	31.132	IV	2	(1)	2	49.269 (1)	III	6	(1)	2	53.328 III 4 (2)	2
27.79 (1) II	1 ((1) 4	۱I	32.135	Ι	3	(1)	2	49.269 (2)	III	6	(1)	2	53.329 III 2 (1)	4
27.79 (2) II	1 ((1) 4	l	32.136	Π	1	(1)	4	49.270 (1)	III	4	(1)	2	53.330 IV 4 (1)	4
27.80 (1) III	1 ((1) 2	2	32.137	III	1	(1)	2	49.270 (2)	III	5	(2)	2	53.332 IV 4 (1)	4
27.80 (2) III	1 ((1) 2	2	32.138	ш	3	(2)	2	49.271 (1)	III	1	(1)	4	53.334 IV 5 (1)	2
27.81 (1) III	3 ((1) 2	2	32.140	IV	2	(1)	4	49.271 (2)	III	1	(1)	4	54.337 I 3 (1)	4
27.81 (2) III	3 ((1) 2	2	32.141	IV	3	(1)	2	49.272	IV	2	(1)	4	54.338 II 2 (2)	8
27.82 (1) IV	1 ((1) 2	2	35.165	Ι	2	(1)	2	49.273 (1)	IV	4	(1)	2	54.339 III 1 (1)	4
27.82 (2) IV	1 ((1) 2	2	35.166	Π	1	(1)	2	49.273 (2)	IV	4	(1)	2	54.340 III 2 (1)	4
27.83 IV	1 ((1) 4	ı	35.167	Ш	2	(1)	1	49.274	IV	3	(1)	4	54.341 III 1 (1)	4
27.85 IV	1 ((1) 4	۱I	35.168	ш	4	(2)	1	50.277	Ι	2	(1)	4	54.342 III 3 (1)	4
28.87 (1) I	3 ((1) 2	2	35.170	IV	2	(1)	2	50.278	Π	2	(2)	4	54.343 III 2 (1)	4
28.87 (2) I	4 ((1) 2	2	38.187	Ι	1	(1)	2	50.279	III	2	(1)	2	54.344 III 3 (2)	4
28.88 (1) II	1 ((1) 4	۱I	38.188	Π	1	(1)	2	50.280	III	3	(1)	2	54.345 III 3 (1)	4
28.88 (2) II	1 ((1) 4	۱I	38.189	ш	2	(1)	1	50.281	III	4	(2)	2	54.346 IV 3 (1)	4
28.89 (1) III	1 ((1) 2	2	38.190	ш	2	(1)	1	50.282	III	3	(1)	2	54.348 IV 2 (1)	4
28.89 (2) III	1 ((1) 2	2	38.191	ш	2	(1)	1	50.283	III	1	(1)	4	54.350 IV 3 (1)	4
28.90 (1) III	2 ((1) 2	2	38.193	IV	1	(1)	2	50.284	IV	2	(1)	4	55.353 I 5 (2)	2
28.90 (2) III	2 ((1) 2	2	39.195	Ι	1	(1)	2	50.287	IV	2	(1)	4	55.354 II 3 (2)	4
28.91 (1) III	2 ((1) 2	2	39.196	Π	1	(1)	4	51.289 (1)	Ι	5	(2)	2	55.355 III 1 (1)	4
28.91 (2) III	3 ((2) 2	2	39.197	Ш	2	(1)	2	51.289 (2)	Ι	4	(1)	2	55.356 III 1 (1)	4
28.92 (1) IV	3 ((1) 2	2	39.198	ш	1	(1)	2	51.290 (1)	Π	3	(2)	4	55.357 III 6 (2, 2)) 2
28.92 (2) IV	4 ((1) 2	2	39.199	ш	2	(1)	2	51.290 (2)	Π	3	(2)	4	55.358 III 3 (1)	2
28.93 IV	2 ((1) 4	1	39.201	IV	1	(1)	2	51.291 (1)	III	1	(1)	2	55.359 III 1 (1)	4
28.94 IV	1 ((1) 4	L	47.249 (1)	Ι	5	(2)	2	51.291 (2)	III	1	(1)	2	55.360 IV 3 (2)	4
28.96 IV	3 ((1) 4	۱I	47.249 (2)	Ι	5	(2)	2	51.292 (1)	III	2	(1)	2	55.363 IV 5 (2)	2
28.97 IV	3 ((1) 2	2	47.249 (3)	Ι	5	(2)	2	51.292 (2)	III	2	(1)	2	57.377 I 3 (1)	4
29.99 I	1 ((1) 4	ı	47.250 (1)	Π	5	(2)	2	51.293 (1)	III	1	(1)	4	57.378 II 2 (2)	8
29.100 II	1 ((1) 8	3	47.250 (2)	II	5	(2)	2	51.293 (2)	III	1	(1)	4	57.379 III 3 (1)	4
29.101 III	1 ((1) 4	۱I	47.250 (3)	II	5	(2)	2	51.294 (1)	III	4	(1)	2	57.380 III 1 (1)	4
29.102 III	1 ((1) 4	ı	47.251 (1)	Ш	1	(1)	2	51.294 (2)	III	5	(2)	2	57.381 III 1 (1)	4
29.103 III	1 ((1) 4	۱I	47.251 (2)	ш	1	(1)	2	51.295 (1)	III	4	(1)	2	57.382 III 3 (1)	4
29.104 IV	1 ((1) 4	۱I	47.251 (3)	ш	1	(1)	2	51.295 (2)	III	4	(1)	2	57.383 III 3 (2)	4
29.106 IV	1 ((1) 4	۱I	47.252 (1)	ш	9	(1)	1	51.296 (1)	III	6	(2, 2)	2	57.384 III 2 (1)	4
29.108 IV	1 ((1) 4	۱I	47.252 (2)	ш	9	(1)	1	51.296 (2)	III	6	(1)	2	57.385 III 2 (1)	4
30.111 I	2 ((1) 2	2	47.252 (3)	ш	10	(2, 2)	1	51.297 (1)	III	2	(1)	2	57.387 IV 2 (1)	4
30.112 II	1 ((1) 4	ł	47.253 (1)	III	1	(1)	2	51.297 (2)	III	3	(1)	2	57.388 IV 3 (1)	4
30.113 III	1 ((1) 2	2	47.253 (2)	III	1	(1)	2	51.298 (1)	IV	5	(2)	2	57.389 IV 3 (1)	4
30.114 III	1 ((1) 2	2	47.253 (3)	III	1	(1)	2	51.298 (2)	IV	4	(1)	2	59.405 I 2 (1)	4
30.115 III	2 ((1) 2	2	47.254 (1)	IV	3	(2)	4	51.299	IV	2	(1)	4	59.406 II 2 (2)	4
30.117 IV	2 ((1) 4	1	47.254 (2)	IV	3	(2)	4	51.300	IV	3	(2)	4	59.407 III 2 (1)	2

(Continued from the previous page)

59.400 III 4 (2) 90.98 III 5 (4) 2 125.365 III 4 (1) 2 166.85 I 8 (1) 1 59.411 III 5 (1) 2 125.365 III 4 164.85 II 6 (2) 2 59.415 IV 2 (1) 4 99.165 II 2 125.365 III 4 164.88 II 4 (2) 2 99.166 II 2 125.375 III 4 10 168.11 II 4 10 1 165.485 III 1 100.171 II 2 101.41 125.371 II 8 4 1 127.378 II 8 1 14 3 1 1 127.378 II 8 14 13 14 3 1 1 14 3 1 127.378 II 18 16 <th>59.408 III</th> <th>1</th> <th>(1)</th> <th>4</th> <th>90.97</th> <th>III</th> <th>2</th> <th>(1)</th> <th>2</th> <th>125.363</th> <th>Ι</th> <th>4</th> <th>(1)</th> <th>4</th> <th>162.76 II</th> <th>I 3</th> <th>(1)</th> <th>2</th>	59.408 III	1	(1)	4	90.97	III	2	(1)	2	125.363	Ι	4	(1)	4	162.76 II	I 3	(1)	2
59.410 III 3 (1) 2 90.99 III 1 (1) 2 153.65 III 4 (1) 4 (1) 1 <th1< th=""> <th1< th=""> 1</th1<></th1<>	59.409 III	4	(2)	2	90.98	III	5	(4)	2	125.364	Π	4	(2)	4	162.77 II	I 9	(6)	1
59.411 III 3 (1) 2 10.101 IV 4 (1) 2 125.367 II 2 (1) 4 164.87 III 3 (1) 2 55.415 IV 5 (2) 2 99.165 III 2 125.367 III 4 164.88 III 4 (1) 2 65.483 II 4 (2) 2 99.167 III 4 (1) 2 125.371 IV 4 (1) 2 168.100 I 4 (1) 2 65.483 III 1 (1) 2 99.167 II 4 (1) 4 127.383 II 4 164.113 II 4 (1) 4 173.33 II 4 101.17 II 4 101.17 II 2 127.397 III 4 17.130 II 4 101.17 1 4 10.1 110.175 <td< td=""><td>59.410 III</td><td>3</td><td>(1)</td><td>2</td><td>90.99</td><td>III</td><td>1</td><td>(1)</td><td>4</td><td>125.365</td><td>III</td><td>4</td><td>(1)</td><td>2</td><td>164.85 I</td><td>8</td><td>(1)</td><td>1</td></td<>	59.410 III	3	(1)	2	90.99	III	1	(1)	4	125.365	III	4	(1)	2	164.85 I	8	(1)	1
59.412 IV 2 (1) 4 96.163 1 3 (1) 2 125.368 II 2 (1) 4 164.87 III 3 (1) 2 55.415 I 5 (2) 2 99.166 III 1 (1) 2 125.375 III 4 (1) 2 164.89 III 9 (6) 1 65.485 III 4 (2) 2 99.166 III 1 125.371 II 2 (1) 4 125.371 II 4 (1) 1 14 3 1	59.411 III	3	(1)	2	90.101	IV	4	(1)	2	125.366	III	2	(1)	4	164.86 I	6	(2)	2
59.415 IV 2 (1) 2 (1) 2 (1) 2 (1) 2 (1) 2 (1) 2 (1) 2 (1) 2 (1) 2 (1) 2 (1) 2 (1) 2 (1) 2 (1) 2 (1) 1 1 1 1 1 1 1 2 100.171 1 2 (1) 4 (1) 4 (1) 4 (1) 1 <	59.412 IV	2	(1)	4	99.163	Ι	3	(1)	2	125.367	III	2	(1)	4	164.87 II	I 3	(1)	2
65.481 I 5 (2) 299.165 II (1) 2 125.379 II 4 (1) 2 164.89 II 9 (6) 1 65.483 II (1) 2 99.166 II (1) 2 125.371 III 4 (1) 2 168.10 II 4 (1) 2 65.484 III 1 (1) 2 100.171 II 4 (1) 2 127.387 II 8 4 2 174.133 I 4 (1) 2 65.487 II (1) 2 100.173 II 2 (1) 2 127.399 III 4 (1) 2 175.139 II 4 (1) 2 175.393 II 10 175.139 II 4 (1) 2 175.393 II 10 4 175.393 II 10 4 175.393 II 10 4 175.393 II 10 4 1175.391 1 1 175.491 1	59.415 IV	2	(1)	4	99.164	II	3	(1)	2	125.368	III	2	(1)	4	164.88 II	I 4	(1)	2
65.482 II 4 (1) 2 99.166 II 1 (1) 2 168.10 I 9 (6) 1 65.483 II (1) 2 99.167 II 8 (4) 1 125.371 II 2 168.110 II 4 (1) 1 65.485 III 6 (1) 1 100.172 II 4 (1) 2 127.387 II 8 (2) 1 14 (3) 2 65.485 III 3 (1) 2 100.173 II 5 14 171.135 II 4 (3) 2 177.390 II 4 (1) 2 177.391 II 14 (1) 2 177.391 II 4 (1) 2 177.391 II 177.137 II 4 (1) 2 177.391 II 1 11 11 11 11 11 11 11 11 11 11 11 11 11 11 11 11	65.481 I	5	(2)	2	99.165	III	2	(1)	2	125.369	III	8	(4)	2	164.89 II	I 9	(6)	1
65.483 III 1 1 2 99.167 III 8 4 1 125.371 III 2 1 4 1<1	65.482 II	4	(2)	2	99.166	III	1	(1)	2	125.370	III	4	(1)	2	168.109 I	9	(6)	1
65.484 III 1 10 10 10 10 1 100.171 1 4 10 2 127.378 1 8 (4) 1 174.133 II 4 (1) 2 10 127.388 II 8 (4) 2 174.133 II 4 (1) 1 100.173 II 1 127.388 II 8 (4) 2 174.134 II 7 (3) 2 65.487 II 1 2 100.173 II 2 127.390 III 4 175.138 II 9 (6) 2 67.501 II 3 (1) 2 110.251 I 3 (1) 2 127.393 III 4 (1) 2 175.14 II 6 0 2 175.14 II 4 (1) 2 127.393 III 4 (1) 2 175.14 II 4 11 2 11 2 11 2 11 2 11 2 11 1	65.483 III	1	(1)	2	99.167	III	8	(4)	1	125.371	III	2	(1)	4	168.110 I	4	(1)	2
65.485 III 8 (2, 2) 1 100.171 I 4 (1) 2 127.387 I 8 (4) 2 174.133 I 14 (3, 3) 1 65.486 III (1) 2 100.173 III 1 (1) 4 127.389 III 3 (1) 7 (3) 2 65.486 IV 4 100.173 III 2 (1) 2 127.390 III 3 (2) 4 175.137 I 18 (6, 6) 1 67.503 II 3 (1) 2 127.393 III 4 (2) 2 175.139 III 7 (3) 2 67.505 III 5 (1) 2 111.251 III 1 112.51 II 1 127.393 III 2 117.150 II 4 (1) 2 127.394 III 2 117.150 II 4 (1) 2 127.397 IV 8 (4) 1 111.11 <	65.484 III	1	(1)	2	99.169	IV	2	(1)	4	125.373	IV	4	(1)	4	168.111 II	I 4	(1)	1
65.486 III 6 (1) 1 100.172 II 2 (1) 4 127.388 II 5 (4) 4 174.134 II 7 (3) 2 65.487 IV 4 (2) 2 100.173 III (1) 2 127.390 III 4 (2) 2 175.138 II 9 (6) 2 67.501 I 3 (2) 4 100.175 II 5 (4) 2 127.390 III 4 105.138 II 9 (6) 2 67.503 II 3 (1) 2 127.394 III 4 (1) 2 177.150 II 4 (1) 2 127.395 III 4 11.1 1 10 2 111.251 III 1 112.54 III 1 112.54 III 1 129.411 I 4 11 11 1 11 11 129.411 I 4 11 11 1 129.411 II <t< td=""><td>65.485 III</td><td>8</td><td>(2, 2)</td><td>1</td><td>100.171</td><td>Ι</td><td>4</td><td>(1)</td><td>2</td><td>127.387</td><td>Ι</td><td>8</td><td>(4)</td><td>2</td><td>174.133 I</td><td>14</td><td>4 (3, 3)</td><td>1</td></t<>	65.485 III	8	(2, 2)	1	100.171	Ι	4	(1)	2	127.387	Ι	8	(4)	2	174.133 I	14	4 (3, 3)	1
65.487 III 1 (1) 2 100.173 III 1 (1) 4 127.389 III 2 107.435 III 4 (1) 1 65.487 II 3 (1) 2 100.174 III 2 (1) 2 107.138 II 9 (6) 2 67.502 II 3 (2) 4 101.171 V 4 (1) 2 127.392 II 2 10<.41	65.486 III	6	(1)	1	100.172	II	2	(1)	4	127.388	Π	5	(4)	4	174.134 I	7	(3)	2
65.489 IV 4 (2) 2 100.174 III 2 (1) 2 127.390 III 3 (2) 4 175.137 I I8 (6, 6) 1 67.503 III 3 (2) 4 100.177 IV 4 (1) 2 127.393 III 0 4 175.138 II 9 (6) 2 67.503 III 1 (1) 2 111.251 I 3 (1) 2 127.393 III 0 4 (1) 4 (1) 2 67.506 III 5 (1) 2 111.255 II 8 (4) 1 129.412 I 4 (2) 4 177.150 I 4 (1) 2 129.412 II 4 (1) 2 113.267 I 4 (1) 129.414 III 2 (1) 1 13.185 I 4 (1) 2 75.5 II 8 (4) 1 113.270 II 4	65.487 III	1	(1)	2	100.173	III	1	(1)	4	127.389	III	2	(1)	4	174.135 II	I 4	(1)	1
67.501 I 3 (1) 2 100.175 III 5 (4) 2 127.392 III 4 (1) 4 (1) 2 67.503 III 1 (1) 2 111.251 I 3 (1) 2 127.393 III 6 (1) 2 67.504 II 5 (1) 2 111.251 II 3 (1) 2 127.395 III 2 175.141 III 6 (2) 1 67.505 II 5 (1) 2 111.251 II 1 (1) 2 127.395 III 4 (1) 2 177.150 II 4 (1) 2 67.507 II 8 (1) 2 111.257 II 4 (1) 2 129.413 II 4 11 1 4 (1) 2 113.268 II 129.414 II 4 11 2 113.48 II 4 (1) 2 183.185 II 4 (1)<	65.489 IV	4	(2)	2	100.174	III	2	(1)	2	127.390	III	3	(2)	4	175.137 I	18	8 (6, 6)	1
67.502 II 3 (2) 4 100.177 IV 4 (1) 2 127.392 II 2 10 4 175.139 II 7 (3) 2 67.503 III 2 (11.252 II 3 (1) 2 127.393 III 2 (1) 4 175.141 III 6 (2) 1 67.505 III 2 111.253 III 1 (1) 2 127.397 IV 8 (4) 1 177.145 III 4 (1) 2 67.507 III 2 111.257 IV 2 (1) 4 (2) 4 177.153 II 4 (1) 2 75.3 II 8 (1) 2 113.268 II 2 (1) 4 133.185 I 4 (1) 2 129.415 III 2 14 4 11 2 14 183.185 I 4 11 1 1 1 183.185 I 4	67.501 I	3	(1)	2	100.175	III	5	(4)	2	127.391	III	4	(2)	2	175.138 I	9	(6)	2
67.503 III 1 11.251 I 3 (1) 2 127.393 III 10 (4, 4) 2 175.140 III 4 (1) 2 67.504 III 2 111.253 II 2 10 2 177.149 I 4 (1) 2 67.506 II 5 (1) 2 111.254 II 1 1 2 127.397 IV 8 (4) 2 177.150 II 4 (1) 2 67.507 II 2 111.257 IV 2 (1) 4 129.411 II 4 (1) 2 177.150 II 4 (1) 2 177.157 II 4 (1) 2 112.471 II 4 (1) 2 113.268 II 2 (1) 4 183.185 I 4 (1) 2 113.271 II 2 129.411 III 8 4 11 1 1 1 1 1 1 1 1	67.502 II	3	(2)	4	100.177	IV	4	(1)	2	127.392	Ш	2	(1)	4	175.139 II	I 7	(3)	2
67.504 III 2 (1) 2 111.252 II 3 (1) 2 127.394 III 2 (1) 4 175.141 III 6 (2) 1 67.505 III 5 (1) 2 111.253 III 2 (1) 2 (1) 4 177.149 I 4 (1) 2 67.506 III 2 111.255 II 8 (4) 1 129.411 II 4 (1) 4 177.150 II 4 (1) 1 11.1257 II 2 11.257 II 4 129.413 III 2 (1) 4 177.153 III 5 (1) 2 113.267 II 4 10 2 113.267 III 2 11.257 III 3 11 2 11 2 11	67.503 III	1	(1)	2	111.251	Ι	3	(1)	2	127.393	Ш	10	(4, 4)	2	175.140 II	I 4	(1)	2
67.505 III 5 (2) 2 111.253 III 2 (1) 2 (1) 4 (1) 2 67.506 III 5 (1) 2 111.253 III 1 (1) 2 (12,3)7 IV 8 (4) 2 177.150 II 4 (1) 2 67.507 II 2 111.257 II 8 (4) 1 129.411 II 4 (1) 1 177.155 III 4 (1) 1 177.157 III 6 (1) 1 13.267 I 4 129.411 III 2 (1) 4 177.153 III 9 (6) 1 75.7 II 3 (1) 2 113.267 II 4 10 2 183.187 III 5 (1) 1 13.267 II 4 129.417 III 8 4 1 13.273 II 4 10 2 183.187 II 5 (1) 1 13.273 II	67.504 III	2	(1)	2	111.252	Π	3	(1)	2	127.394	III	2	(1)	4	175.141 II	I 6	(2)	1
67.506 III 5 (1) 2 111.254 III 1 (1) 2 127.397 IV 8 (4) 2 177.150 II 4 (1) 2 67.507 III 2 (1) 2 111.255 III 8 (4) 1 129.411 I 4 (2) 4 177.151 III 4 (1) 1 75.0 IV 3 (1) 2 113.268 II 2 (1) 4 129.414 III 2 (1) 4 183.186 I 4 (1) 2 75.5 IV 4 (2) 2 113.270 III 2 (1) 2 129.416 III 4 11 5 (1) 1 81.33 I 8 (4) 1 113.271 III 5 (4) 2 129.416 III 4 11 15 (1) 1 133.188 III 4 (1) 2 183.48 II 3 11 2	67.505 III	5	(2)	2	111.253	III	2	(1)	2	127.395	III	2	(1)	4	177.149 I	4	(1)	2
67.507 III 2 111.255 III 8 (4) 1 129.411 I 4 (1) 4 (17.151 III 4 (1) 1 67.509 IV 3 (1) 2 111.257 IV 2 (1) 4 129.412 II 4 (2) 4 177.153 III 5 (1) 1 75.3 II 2 (1) 2 113.268 II 2 (1) 4 129.414 III 2 (1) 4 183.186 I 4 (1) 2 75.5 IV 4 (2) 2 113.270 III 2 129.417 III 8 4 (1) 1 1 1 1 1 1 1 1 13 1 3 (1) 2 129.417 III 8 4 2 183.188 III 4 (1) 1 1 1 1 1 1 1 1 1 1 1 1 1 1	67.506 III	5	(1)	2	111.254	III	1	(1)	2	127.397	IV	8	(4)	2	177.150 I	4	(1)	2
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	67.507 III	2	(1)	2	111.255	III	8	(4)	1	129.411	Ι	4	(1)	4	177.151 II	I 4	(1)	1
75.1I8(4)1 113.267 I4(1)2 129.413 III2(1)4 177.153 III9(6)175.2II3(1)2 113.268 II2(1)4 129.414 III2(1)4 183.185 I4(1)275.3III2(1)2 113.269 III1(1)4 129.415 III2(1)4 183.186 II4(1)281.33I8(4)1 113.271 III5(4)2 129.416 III4(1)2 183.187 III5(1)181.33II3(1)2 113.273 IV4(1)2 129.417 III8(4)2 183.188 III4(1)181.35III2(1)2 113.273 IV4(1)2 129.417 III8(4)2 183.188 III4(1)281.37IV4(2)2 115.285 III2 129.411 IV4(1)4 187.210 II7(3)283.43II8(4)2 115.286 III1 111.22 143.2 II4(1)2 187.212 III5(1)183.44II8(4)2 <td< td=""><td>67.509 IV</td><td>3</td><td>(1)</td><td>2</td><td>111.257</td><td>IV</td><td>2</td><td>(1)</td><td>4</td><td>129.412</td><td>Π</td><td>4</td><td>(2)</td><td>4</td><td>177.152 II</td><td>I 5</td><td>(1)</td><td>1</td></td<>	67.509 IV	3	(1)	2	111.257	IV	2	(1)	4	129.412	Π	4	(2)	4	177.152 II	I 5	(1)	1
75.2II3(1)2113.268II2(1)4129.414III2(1)4183.185I4(1)275.3III2(1)2113.269III1(1)4129.415III2(1)4183.186II4(1)275.5IV4(2)2113.270III2(1)2129.416III4(1)2183.187III5(1)181.33I8(4)1113.273IV4(1)2129.418III2(1)4183.189III9(6)181.35III2(1)2115.283I3(1)2129.419III4(1)2187.209I7(3)281.37IV4(2)2115.284II3(1)2129.419III4(1)4187.210II7(3)283.43I16(4,4)1115.285III2(1)2143.1I7(3)1187.211III5(1)183.44II8(4)2115.287III8(4)1147.13I9(6)1187.213III14(3,3)183.45III5(1)2117.209II	75.1 I	8	(4)	1	113.267	Ι	4	(1)	2	129.413	III	2	(1)	4	177.153 II	I 9	(6)	1
75.3III2(1)2113.269III1(1)4129.415III2(1)4183.186II4(1)275.5IV4(2)2113.270III2(1)2129.416III4(1)2183.187III5(1)181.33I8(4)1113.271III5(4)2129.417III8(4)2183.188III4(1)181.35III2(1)2113.273IV4(1)2129.418III2(1)4183.189III9(6)181.35III2(1)2115.283I3(1)2129.419III4(1)4187.209I7(3)283.43I16(4,4)1115.285III2(1)2143.1I7(3)1187.211III5(1)183.44II8(4)2115.287III8(4)1147.13I9(6)1187.213II4(3)183.45III3(1)2115.287III8(4)1147.13I9(6)1187.213II4(3)183.46III3(1)2117.209I<	75.2 II	3	(1)	2	113.268	II	2	(1)	4	129.414	III	2	(1)	4	183.185 I	4	(1)	2
75.5IV4(2)2113.270III2(1)2129.416III4(1)2183.187III5(1)181.33I8(4)1113.271III5(4)2129.417III8(4)2183.188III4(1)181.35III2(1)2115.283I3(1)2129.418III2(1)4187.209I7(3)281.37IV4(2)2115.284II3(1)2129.419III4(1)2187.209I7(3)283.43I16(4, 4)1115.285III2(1)2143.2II4(1)2187.212III5(1)183.44II8(4)2115.286III1(1)2143.2II4(1)2187.213III14(3)3183.45III5(2)2115.287III8(4)1147.13I9(6)1187.213III14(3)3283.47III3(1)2117.209I4(1)2147.15III4(1)2189.223II4(1)<1	75.3 III	2	(1)	2	113.269	III	1	(1)	4	129.415	III	2	(1)	4	183.186 I	4	(1)	2
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	75.5 IV	4	(2)	2	113.270	III	2	(1)	2	129.416	III	4	(1)	2	183.187 II	I 5	(1)	1
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	81.33 I	8	(4)	1	113.271	Ш	5	(4)	2	129.417	Ш	8	(4)	2	183.188 II	I 4	(1)	1
81.35 III 2 (1) 2 115.283 I 3 (1) 2 129.419 III 4 (1) 2 187.209 I 7 (3) 2 81.37 IV 4 (2) 2 115.284 II 3 (1) 2 129.421 IV 4 (1) 4 187.210 II 7 (3) 2 83.43 I 16 (4, 4) 1 115.285 III 2 (1) 2 143.1 I 7 (3) 1 187.212 III 5 (1) 1 83.45 III 5 (2) 2 115.287 III 8 (4) 1 147.13 I 9 (6) 1 187.213 III 14 (3) 3 2 83.45 III 3 (1) 2 117.300 II 2 147.15 III 4 (1) 1 189.223 III 4 (1) 1 189.225 III 10 (3)	81.34 II	3	(1)	2	113.273	IV	4	(1)	2	129.418	III	2	(1)	4	183.189 II	I 9	(6)	1
81.37IV4(2)2115.284II3(1)2129.421IV4(1)4187.210II7(3)283.43I16(4,4)1115.285III2(1)2143.1I7(3)1187.211III5(1)183.44II8(4)2115.286III1(1)2143.2II4(1)2187.212III5(1)183.45III5(2)2115.287III8(4)1147.13I9(6)1187.213III14(3,3)183.46III3(1)2117.299I4(1)2147.15III4(1)2189.222II5(3)283.47III3(1)2117.300II2(1)4149.21I5(1)1189.223III4(1)185.59I8(4)2117.300III2149.23III7(3)1189.225III10(3,3)185.60II4(1)2117.300III2150.25I5(1)1191.233I9(6)285.61III2117.305IV4(1)2150.26II3	81.35 III	$ ^2$	(1)	2	115.283	l n	3	(1)	2	129.419	III	4	(1)	2	187.209 I	17	(3)	2
83.43116(4, 4)1115.2851112(1)2143.117(3)1187.2111115(1)183.44II8(4)2115.286III1(1)2143.2II4(1)2187.212III5(1)183.45III5(2)2115.287III8(4)1147.13I9(6)1187.213III14(3, 3)183.46III3(1)2115.289IV2(1)4147.14II6(2)2189.221I7(3)283.47III3(1)2117.300II2(1)4149.21I5(1)1189.223III4(1)185.59I8(4)2117.301III1(1)4149.22II4(1)2189.224III4(1)185.60II4(2)4117.302III2150.25I5(1)1191.233I9(6)285.61III2117.305IV4(1)2150.25I5(1)1191.234II9(6)285.62III4(1)2123.339I8(4)2150.27 <td>81.37 IV</td> <td>4</td> <td>(2)</td> <td>2</td> <td>115.284</td> <td>11 </td> <td>3</td> <td>(1)</td> <td>2</td> <td>129.421</td> <td>IV</td> <td>4</td> <td>(1)</td> <td>4</td> <td>187.210 1</td> <td>. 7</td> <td>(3)</td> <td>2</td>	81.37 IV	4	(2)	2	115.284	11 	3	(1)	2	129.421	IV	4	(1)	4	187.210 1	. 7	(3)	2
83.44 II 8 (4) 2 115.286 III 1 (1) 2 143.2 II 4 (1) 2 187.212 III 5 (1) 1 83.45 III 5 (2) 2 115.287 III 8 (4) 1 147.13 I 9 (6) 1 187.213 III 14 (3, 3) 1 83.46 III 3 (1) 2 115.289 IV 2 (1) 4 147.15 III 6 (2) 2 189.221 I 7 (3) 2 83.47 III 3 (1) 2 117.300 II 2 (1) 4 149.21 I 5 (1) 1 189.223 III 4 (1) 1 85.60 II 4 (2) 4 117.302 III 2 150.25 I 5 (1) 1 191.233 I 9 (6) 2 85.61 III 2 123.339 I<	83.43 I) (4, 4)	1	115.285	III 	2	(1)	2	143.1	1		(3)	1	187.211 11	נן גן	(1)	1
83.45III5(2)2115.287III8(4)1147.1319(6)1187.213III14(3, 3)183.46III3(1)2115.289IV2(1)4147.14II6(2)2189.221I7(3)283.47III3(1)2117.299I4(1)2147.15III4(1)2189.222II5(3)283.49IV8(4)2117.300II2(1)4149.21I5(1)1189.223III4(1)185.59I8(4)2117.300III2(1)2149.23III7(3)1189.225III4(1)185.60II4(2)4117.302III2150.25I5(1)1191.233I9(6)285.61III4(1)2123.339I8(4)2150.26II31191.235III4(1)285.65IV4(1)4123.340II8(4)2156.49I5(1)1191.236III7(3)289.87I3(1)2123.341III3(1)215	83.44 II	8	(4)	2	115.286	111 111		(1)	2	143.2	П	4	(1)	2	187.212 II		(1)	1
83.46III3(1)2115.289IV2(1)4147.14II6(2)2189.221I7(3)283.47III3(1)2117.299I4(1)2147.15III4(1)2189.222II5(3)283.49IV8(4)2117.300II2(1)4149.21I5(1)1189.223III4(1)185.59I8(4)2117.301III1(1)4149.22II4(1)2189.225III4(1)185.60II4(2)4117.302III2(1)2149.23III7(3)1189.225III4(1)185.61III2(1)4117.305IV4(1)2150.25I5(1)1191.233I9(6)285.62III4(1)2123.339I8(4)2150.27III5(3)1191.235III4(1)285.65IV4(1)4123.340II8(4)2156.49I5(1)1191.237III5(3)289.87I3(1)2123.342III5 </td <td>83.45 III</td> <td>0</td> <td>(2)</td> <td>2</td> <td>115.287</td> <td>III III</td> <td>8</td> <td>(4)</td> <td>1</td> <td>147.13</td> <td>1</td> <td>9</td> <td>(6)</td> <td>1</td> <td>187.213 11</td> <td></td> <td>+ (3, 3)</td> <td>1</td>	83.45 III	0	(2)	2	115.287	III III	8	(4)	1	147.13	1	9	(6)	1	187.213 11		+ (3, 3)	1
85.47 III 5 (1) 2 I17.299 I 4 (1) 2 I47.15 III 4 (1) 2 I89.222 II 5 (5) 2 83.49 IV 8 (4) 2 117.300 II 2 (1) 4 149.21 I 5 (1) 1 189.223 III 4 (1) 1 85.59 I 8 (4) 2 117.301 III 1 (1) 4 149.22 II 4 (1) 1 189.223 III 4 (1) 1 189.225 III 4 (1) 1 10 (3, 3) 1 189.225 III 10 (3, 3) 1 191.233 I 9 (6) 2 85.65 III 4 (1) 2 123.339 I 8 (4) 2 150.25 II 5 (1) 1 191.235 III 4 (1) 2 185.65 IV 4 (1) 4 (1) 2 123.340 </td <td>83.46 III</td> <td>3</td> <td>(1)</td> <td>2</td> <td>115.289</td> <td>1</td> <td>2</td> <td>(1)</td> <td>4</td> <td>147.14</td> <td>Ш Ш</td> <td>0</td> <td>(2)</td> <td>2</td> <td>189.221</td> <td></td> <td>(3)</td> <td>2</td>	83.46 III	3	(1)	2	115.289	1	2	(1)	4	147.14	Ш Ш	0	(2)	2	189.221		(3)	2
85.49 IV 8 (4) 2 I17.300 II 2 (1) 4 149.21 I 5 (1) 1 189.225 III 4 (1) 1 85.59 I 8 (4) 2 117.301 III 1 (1) 4 149.22 II 4 (1) 2 189.224 III 4 (1) 1 85.60 II 4 (2) 4 117.302 III 2 149.23 III 7 (3) 1 189.225 III 10 (3, 3) 1 85.61 III 2 (1) 4 117.303 III 5 (4) 2 150.25 I 5 (1) 1 191.234 II 9 (6) 2 85.65 IV 4 (1) 4 123.340 II 8 (4) 2 156.49 I 5 (1) 1 191.235 III 4 (1) 2 85.65 IV 4 (1) 4	83.47 III	0	(1)	2	117.299	п	4	(1)	2	147.15	ш	4	(1)	2	189.222 I	. Э 1 4	(3)	2
85.39 1 8 (4) 2 117.301 11 1 (1) 4 149.22 11 4 (1) 1 189.224 11 4 (1) 1 85.60 II 4 (2) 4 117.302 III 2 149.23 III 7 (3) 1 189.225 III 10 (3, 3) 1 85.61 III 2 (1) 2 150.25 I 5 (1) 1 191.233 I 9 (6) 2 85.62 III 4 (1) 2 123.339 I 8 (4) 2 150.26 II 3 (1) 2 191.234 II 9 (6) 2 85.63 III 4 (1) 2 123.340 II 8 (4) 2 156.49 I 5 (1) 1 191.235 III 4 (1) 2 89.87 I 3 (1) 2 123.342 III 5 (2) 2	85.49 IV	0	(4)	2	117.300	ш	1	(1)	4	149.21	п		(1)	1	189.225 II	1 4 1 4	(1)	1
85.00 II 4 117.302 III 2 149.25 III 7 (3) 1 189.225 III 10 (3, 3) 1 85.61 III 2 (1) 4 117.303 III 5 (4) 2 150.25 I 5 (1) 1 191.233 I 9 (6) 2 85.62 III 4 (1) 2 123.339 I 8 (4) 2 150.26 II 3 (1) 2 191.234 II 9 (6) 2 85.63 III 4 (1) 2 123.339 I 8 (4) 2 150.27 III 5 (3) 1 191.235 III 4 (1) 2 85.65 IV 4 (1) 4 123.340 II 8 (4) 2 156.50 II 4 (1) 2 191.237 III 5 (3) 2 89.87 I 3 (1) 2 123.343 III	оз.39 I 95.60 П	0	(4)	4	117.301	ш	1	(1)	4	149.22	ш	4	(1)	1	109.224 II	I 4 I 1/	(1)	1
85.01 III 2 (1) 4 I17.305 III 5 (4) 2 I50.25 I 5 (1) 1 I91.235 I 9 (0) 2 85.62 III 4 (1) 2 117.305 IV 4 (1) 2 150.25 II 3 (1) 2 191.235 II 9 (6) 2 85.63 III 4 (1) 2 123.339 I 8 (4) 2 150.27 III 5 (3) 1 191.235 III 4 (1) 2 85.65 IV 4 (1) 4 123.340 II 8 (4) 2 156.49 I 5 (1) 1 191.237 III 5 (3) 2 89.87 I 3 (1) 2 123.342 III 5 (2) 2 156.51 III 7 (3) 1 191.237 III 8 (1) 1 89.88 II 3	85.61 III	4	(2)	4	117.302	ш	5	(1)	2	149.23	T	5	(3)	1	101 223 1		(5, 5)	1
85.62 III 4 (1) 2 I17.505 IV 4 (1) 2 I50.26 II 5 (1) 2 I51.254 II 5 (0) 2 85.63 III 4 (1) 2 I23.339 I 8 (4) 2 I50.27 III 5 (3) 1 191.235 III 4 (1) 2 85.65 IV 4 (1) 4 123.340 II 8 (4) 2 156.49 I 5 (1) 1 191.235 III 4 (1) 2 89.87 I 3 (1) 2 123.341 III 3 (1) 2 156.50 II 4 (1) 2 191.237 III 5 (3) 2 89.88 II 3 (1) 2 123.342 III 5 (2) 2 156.51 III 7 (3) 1 191.238 III 8 (1) 1 89.88 III 1	85.62 III		(1)	7 2	117.305	III IV		(1)	$\frac{2}{2}$	150.25	п	3	(1)	2	191.235 I 101 234 I	0	(6)	2
85.65 IV 4 (1) 2 123.357 I 8 (4) 2 156.27 III 5 (5) I 191.255 III 4 (1) 2 85.65 IV 4 (1) 4 123.340 II 8 (4) 2 156.49 I 5 (1) 1 191.235 III 7 (3) 2 89.87 I 3 (1) 2 123.341 III 3 (1) 2 156.50 II 4 (1) 2 191.237 III 5 (3) 2 89.88 II 3 (1) 2 123.342 III 5 (2) 2 156.51 III 7 (3) 1 191.238 III 8 (1) 1 89.89 III 1 (1) 2 123.343 III 4 (2) 2 157.53 I 5 (1) 1 191.240 III 18 (6, 6) 1 89.91 III 2 <td>85.63 III</td> <td></td> <td>(1)</td> <td>2</td> <td>123 330</td> <td>T</td> <td>8</td> <td>(1)</td> <td>2</td> <td>150.20</td> <td>ш</td> <td>5</td> <td>(1)</td> <td>1</td> <td>101 235 II</td> <td>ι, Γ Λ</td> <td>(0)</td> <td>2</td>	85.63 III		(1)	2	123 330	T	8	(1)	2	150.20	ш	5	(1)	1	101 235 II	ι, Γ Λ	(0)	2
89.85 I 3 (1) 2 123.340 II 3 (1) 2 156.50 II 4 (1) 2 191.235 III 7 (3) 2 89.87 I 3 (1) 2 123.341 III 3 (1) 2 156.50 II 4 (1) 2 191.237 III 5 (3) 2 89.88 II 3 (1) 2 123.342 III 5 (2) 2 156.51 III 7 (3) 1 191.236 III 8 (1) 1 89.89 III 1 (1) 2 123.343 III 4 (2) 2 157.53 I 5 (1) 1 191.239 III 8 (1) 1 89.90 III 8 (4) 1 123.345 III 16 (4, 4) 1 157.55 III 5 (3) 1 191.240 III 8 (6, 6) 1 89.93 IV <td< td=""><td>85.65 IV</td><td></td><td>(1)</td><td>4</td><td>123.339</td><td>п</td><td>8</td><td>(-,-)</td><td>$\frac{2}{2}$</td><td>156.49</td><td>T</td><td>5</td><td>(3)</td><td>1</td><td>191.235 II 191.236 II</td><td>I 7</td><td>(1)</td><td>2</td></td<>	85.65 IV		(1)	4	123.339	п	8	(-,-)	$\frac{2}{2}$	156.49	T	5	(3)	1	191.235 II 191.236 II	I 7	(1)	2
89.88 II 3 (1) 2 123.342 III 5 (2) 2 156.51 III 7 (3) 1 191.235 III 8 (1) 1 89.88 II 3 (1) 2 123.342 III 5 (2) 2 156.51 III 7 (3) 1 191.235 III 8 (1) 1 89.88 III 1 (1) 2 123.342 III 5 (2) 2 157.53 I 5 (1) 1 191.239 III 8 (1) 1 89.90 III 8 (4) 1 123.344 III 3 (1) 2 157.54 II 3 (1) 2 191.240 III 18 (6, 6) 1 89.91 III 2 (1) 2 123.345 III 16 (4, 4) 1 157.55 III 5 (3) 1 191.241 III 4 (1) 2 89.93 IV	89.87 T	3	(1)	т 2	123.340	ш	3	(1)	$\frac{2}{2}$	156.50	п		(1)	2	191.230 П 191.237 П	1 5	(3)	2
89.89 III 1 (1) 2 123.343 III 4 (2) 2 157.53 I 5 (1) 1 191.256 III 6 (1) 1 89.89 III 1 (1) 2 123.343 III 4 (2) 2 157.53 I 5 (1) 1 191.236 III 8 (1) 1 89.90 III 8 (4) 1 123.344 III 3 (1) 2 157.54 II 3 (1) 2 191.240 III 18 (6, 6) 1 89.91 III 2 (1) 2 123.345 III 16 (4, 4) 1 157.55 III 5 (3) 1 191.241 III 4 (1) 2 89.93 IV 2 (1) 4 123.346 III 3 (1) 2 162.73 I 8 (1) 1 191.241 III 4 (1) 2 162.74 II 6 <	89.88 П	3	(1)	2	123.341	ш	5	(1) (2)	$\frac{2}{2}$	156.50	ш	7	(1)	1	191.237 II 191.238 II		(3)	1
89.90 III 8 (4) 1 123.344 III 3 (1) 2 157.54 II 3 (1) 2 191.240 III 18 (6, 6) 1 89.90 III 2 (1) 2 123.344 III 3 (1) 2 191.240 III 18 (6, 6) 1 89.91 III 2 (1) 2 123.345 III 16 (4, 4) 1 157.55 III 5 (3) 1 191.240 III 14 (1) 2 89.93 IV 2 (1) 4 123.346 III 3 (1) 2 162.73 I 8 (1) 1 90.95 I 4 (1) 2 123.347 III 3 (1) 2 162.74 II 6 (2) 2 1	89.89 III	1	(1)	2	123.342	ш	4	(2)	$\frac{2}{2}$	157.53	T	5	(3)	1	191 230 II		(1)	1
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	89.90 III	8	(4)	1	123.344	Ш	3	(2) (1)	2	157.54	П	3	(1)	2	191.240 П		3 (6.6)	1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	89.91 III	2	(1)	2	123,345	Ш	16	(4, 4)	1	157.55	ш	5	(3)	1	191.241 П		(1)	2
90.95 I 4 (1) 2 123.347 III 3 (1) 2 162.74 II 6 (2) 2	89.93 IV	$\left \frac{2}{2} \right $	(1)	4	123.346	Ш	3	(1)	2	162.73	I	8	(1)	1		'	(1)	-
	90.95 I	$\begin{vmatrix} -4 \end{vmatrix}$	(1)	2	123.347	Ш	3	(1)	2	162.74	I	6	(2)	2				
90.96 II 2 (1) 4 23.349 IV 5 (4) 4 162.75 III 4 (1) 2	90.96 П	$ _2$	(1)	4	123.349	IV	5	(4)	4	162.75	Ш	4	(1)	2				

24

TABLE X. Magnetic space groups (MSGs) for which spinful electrons exhibit exceptional filling patterns

MSG	$\{\nu\}_{AI}$	$\{\nu\}_{\rm BS}$	MSG	$\{\nu\}_{AI}$	$\{\nu\}_{\rm BS}$	MSG	$\{\nu\}_{\mathrm{AI}}$	$\{\nu\}_{\rm BS}$	MSG	$\{\nu\}_{AI}$	$\{\nu\}_{\rm BS}$
196.6 IV	$2\mathbb{N}\setminus\{2\}$	$2\mathbb{N}$	209.51 IV	$2\mathbb{N}\setminus\{2\}$	$2\mathbb{N}$	219.88 IV	$2\mathbb{N}\setminus\{2\}$	$2\mathbb{N}$	228.134 I	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$
198.11 IV	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$	210.55 IV	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$	220.89 I	$2\mathbb{N}\setminus\{2,4,10\}$	$2\mathbb{N}\setminus\{2\}$	228.136 III	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$
199.12 I	$2\mathbb{N}\setminus\{2\}$	-	212.62 IV	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$	220.90 II	$4\mathbb{N}\setminus\{4,8,20\}$	$4\mathbb{N}\setminus\{4\}$	228.139 IV	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$
199.13 II	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$	213.66 IV	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$	220.91 III	$4\mathbb{N}\setminus\{4\}$	-	230.145 I	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$
202.25 IV	$2\mathbb{N}\setminus\{2\}$	$2\mathbb{N}$	214.68 II	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$	223.106 III	$2\mathbb{N}\setminus\{2\}$	$2\mathbb{N}$	230.146 II	$8\mathbb{N}\setminus\{8\}$	$8\mathbb{N}$
203.29 IV	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$	214.69 III	$2\mathbb{N}\setminus\{2\}$	-	226.122 I	$2\mathbb{N}\setminus\{2\}$	$2\mathbb{N}$	230.147 III	$4\mathbb{N}\setminus\{4\}$	_
206.39 III	$4\mathbb{N}\setminus\{4\}$	-	218.81 I	$2\mathbb{N}\setminus\{2\}$	$2\mathbb{N}$	226.124 III	$2\mathbb{N}\setminus\{2\}$	$2\mathbb{N}$	230.148 III	$4\mathbb{N}\setminus\{4\}$	-
208.46 III	$2\mathbb{N}\setminus\{2\}$	$2\mathbb{N}$	219.85 I	$2\mathbb{N}\setminus\{2\}$	$2\mathbb{N}$	226.127 IV	$2\mathbb{N}\setminus\{2\}$	$2\mathbb{N}$			

N: The set of natural numbers $\{\nu\}_{AI}$: Set of fillings for which physical atomic insulators are possible $\{\nu\}_{BS}$: Set of fillings for which physical band structure are possible; a dash (-) indicates $\{\nu\}_{BS} = \{\nu\}_{AI}$ for that MSG

Appendix B: Filling-enforced semimetals in stoichiometric compounds

While the electron filling ν of a physical system can, in principle, take any real value upon doping, in perfectly ordered stoichiometric crystals the realizable fillings are further constrained by crystalline symmetries. As the atoms in such a crystal form \mathcal{M} -symmetric lattices, the allowed fillings for an MSG \mathcal{M} are determined by the multiplicities of the Wyckoff positions [33]. Mathematically, this is captured by demanding $\nu \in \nu_{\rm P} \mathbb{N}$, where $\nu_{\rm P}$ is the gcd of the multiplicities of the Wyckoff positions and \mathbb{N} denotes the set of natural number. Whenever $\nu_{\rm P} < \nu_{\rm BS}$, one can find systems with a filling ν that is incompatible with a band gap, and therefore \mathcal{M} admits feSMs. We also remark that for the four 'Wyckoffmismatched' SGs identified in Ref. 8, as well as their magnetic descendants, the set of physically realizable filling does not take the simple form of $\nu_{\rm P}\mathbb{N}$. The exhaustive list of such Wyckoff-mismatched MSGs can be found in Table XXI, since the set physically realizable fillings coincides with the $\{\nu\}_{AI}$ of spinless fermions.

For TR symmetric systems described by the 230 type II MSGs, $\nu_{\rm P} < \nu_{\rm BS}$ holds generally, and the filling criterion for filling-enforced (semi-)metals simplifies into whether or not the filling is an odd multiple of $\nu_{\rm P}$. For a general MSG, however, $\nu_{\rm P} \le \nu_{\rm BS}$, and to identify magnetic settings which can host filling-enforced (semi-)metals one must first isolate the list of MSGs where $\nu_{\rm P} \ne \nu_{\rm BS}$. This is easily achieved using our theory, and we found that 647 out of the 1651 MSGs can host filling-enforced (semi-)metals (this includes the 230 type II MSGs, i.e., there are are 417 intrinsically magnetic MSGs). For all these MSGs, $\nu_{\rm BS} = 2\nu_{\rm P}$, and therefore gaplessness near the Fermi energy is enforced whenever the filling of the system is an odd multiple of $\nu_{\rm P}$.

Some of these MSGs, however, are still incompatible with feSMs due to 'band-sticking.' For instance, whenever the MSG contains the combination of spatial inversion and TR as a symmetry, each band is doubly degenerate. For such MSGs, if $\nu_{\rm P}$ is also odd, the filling-enforced gaplessness always manifests in the form of a large Fermi-surface enclosing half of the Brillouin zone, resulting in a conventional metal. A similar situation arise for band-sticking along high-symmetry line. We refer to such systems as filling-enforced metals. We found that 226 of the identified MSGs fall into this category, and they are listed in Table XIII.

This leaves behind 421 MSGs, 250 of which are intrinsically magnetic, which can potentially host feSMs. Among them, we further characterize their minimal fermiology into two types. If the irrep dimensions at any high-symmetry point is incompatible with a band gap at filling $\nu_{\rm P}$, the fermi surface of the feSM consists of pinned nodal points. These MSGs are tabulated in Table XI. If all the high-symmetry momenta can be in principle gapped, then the semimetallic behavior must be enforced by a more intricate band connectivity, resulting in nodal features that are movable, but irremovable. (Such fermiology is exemplified by that of simple nonsymmorphic SGs, as discussed in, e.g., Ref. 4.) These MSGs are tabulated in Table XII. Note that these are not mutually exclusive characterization of feSMs — certain MSGs in Table XI can also display such movable nodal features.

Lastly, we also include results specific to TR-symmetric systems (type II MSGs) in Table XIV, which describes the effect of spin-orbit coupling on the fermiology of the feSMs.

MSG	$\nu_{\rm F}$	- FS	MSG	r	$\nu_{\rm P}$	FS	MSG	$\nu_{\rm P}$	FS	MSG	ν	∕ _P F	S	MSG	$\nu_{\rm F}$	FS	MSG	$\nu_{\rm P}$	FS
1.2 II	1	2	59.415	IV	2	4	100.173 III	2	4	125.366 II	[2	4	139.540 IV	/ 2	4	195.3 IV	2	4
3.2 II	1	2	63.468	IV	2	4	101.180 II	2	4	125.368 II	[] :	2	4	140.542 II	2	4	196.4 I	1	2
5.14 II	1	2	66.492	Π	2	4	101.181 III	2	4	125.371 II	I :	2 -	4	140.544 II	1 2	4	196.5 II	1	2
6.19 II	1	2	66.497	III	2	4	101.184 IV	2	4	125.374 IV	7	4	8	140.546 II	1 2	4	197.7 I	1	2
8.33 II	1	2	67.502	Π	2	4	102.188 II	2	4	126.375 I		2	4	140.549 II	1 2	4	197.8 II	1	2
11.51 II	2	4	68.512	Π	2	4	102.189 III	2	4	126.376 II		2	4	141.551 I	2	4	201.19 II	2	4
11.57 IV	2	4	68.517	III	2	4	102.194 IV	2	4	126.378 II		2	4	141.552 II	2	4	201.20 III	2	4
13.66 II	2	4	68.520	IV	2	4	103.196 II	2	4	126.379 II	[] :	2	4	141.555 II	1 2	4	203.27 II	2	4
13.73 IV	2	4	70.528	Π	2	4	103.197 III	2	4	126.380 II	[] :	2	4	141.558 II	1 2	4	203.28 III	2	4
14.76 II	2	4	70.531	III	2	4	103.198 III	2	4	126.382 II	[] :	2	4	141.559 II	1 2	4	207.40 I	1	2
15.86 II	2	4	71.538	IV	2	4	104.204 II	2	4	126.383 II	[] :	2	4	143.2 II	1	2	207.41 II	1	2
16.1 I	1	2	72.540	Π	2	4	104.205 III	2	4	126.385 IV	7	4	8	144.5 II	3	2	207.42 III	1	2
16.2 II	1	2	72.545	III	2	4	104.206 III	2	4	127.392 II	[] :	2	4	145.8 II	3	2	207.43 IV	2	4
16.5 IV	$ _2$	4	74.555	Π	2	4	105.212 II	2	4	127.395 II		2	4	146.11 II	1	2	208.47 IV	2	4
16.6 IV	$ _2$	4	75.2	Π	1	2	105.214 III	2	4	128.403 II		2	4	149.22 II	1	2	209.48 I	1	2
18.17 II	2	4	75.3	III	1	2	105.216 IV	2	4	128.404 II	[]	2	4	150.26 II	1	2	209.49 II	1	2
21.38 I	1	2	79.26	Π	1	2	105.218 IV	2	4	128.407 II	[] :	2	4	151.30 II	3	2	209.50 III	1	2
21.39 II	1	2	79.27	III	1	2	107.232 IV	2	4	129.411 I		2	4	152.34 II	3	2	211.56 I	1	2
22.45 I	1	2	84.52	Π	2	4	108.234 II	2	4	129.420 IV	7	4	8	153.38 II	3	2	211.57 II	1	2
22.46 II	1	2	84.56	IV	2	4	108.235 III	2	4	129.421 IV		2	4	154.42 II	3	2	211.58 III	1	2
23.49 I	1	2	84.58	IV	2	4	109.240 II	2	4	130.424 II		4	8	155.46 II	1	2	218.82 II	2	4
23.50 II	1	2	85.60	Π	2	4	109.242 III	2	4	131.435 I		2	4	156.50 II	1	2	218.83 III	2	4
25.61 IV	2	4	85.65	IV	2	4	111.256 IV	2	4	131.436 II		2	4	157.54 II	1	2	220.90 II	2	4
25.63 IV	2	4	85.66	IV	2	4	112.260 II	2	4	131.439 II	[] :	2	4	160.66 II	1	2	220.91 III	2	4
25.64 IV	2	4	86.68	Π	2	4	112.262 III	2	4	131.442 II	[]	2	4	163.80 II	2	4	222.98 I	2	4
25.65 IV	2	4	86.74	IV	2	4	113.274 IV	2	4	131.443 II	[] :	2	4	165.92 II	2	4	222.99 II	2	4
26.67 II	2	4	88.82	Π	2	4	114.278 III	2	4	131.445 IV	7	4	8	167.104 II	2	4	222.100 III	2	4
27.79 II	2	4	89.87	Ι	1	2	115.288 IV	2	4	132.447 I		2	4	168.110 II	1	2	222.101 III	2	4
30.112 II	2	4	89.88	Π	1	2	115.289 IV	2	4	132.448 II		2 -	4	171.122 II	3	2	222.102 III	2	4
31.124 II	2	4	89.89	III	1	2	115.290 IV	2	4	132.450 II	[]	2	4	172.126 II	3	2	223.104 I	2	4
32.136 II	2	4	89.91	III	1	2	116.292 II	2	4	132.452 II	[] :	2	4	174.134 II	1	2	223.105 II	2	4
34.157 II	2	4	89.93	IV	2	4	116.293 III	2	4	132.455 II	[]	2 -	4	176.144 II	2	4	223.107 III	2	4
35.169 IV	2	4	89.94	IV	2	4	117.300 II	2	4	132.458 IV	7	4	8	177.149 I	1	2	223.108 III	2	4
35.171 IV	2	4	90.96	Π	2	4	117.301 III	2	4	134.471 I		2	4	177.150 II	1	2	224.110 I	2	4
36.173 II	2	4	90.99	III	2	4	118.308 II	2	4	134.472 II		2	4	180.167 I	3	2	224.111 II	2	4
37.181 II	2	4	93.126	IV	2	4	118.309 III	2	4	134.474 II	[]	2	4	180.168 II	3	2	224.112 III	2	4
43.225 II	2	4	94.128	Π	2	4	122.334 II	2	4	134.476 II	[] :	2	4	181.173 I	3	2	224.114 III	2	4
48.258 II	2	4	94.131	III	2	4	122.336 III	2	4	134.479 II	[] :	2	4	181.174 II	3	2	225.121 IV	2	4
48.261 III	2	4	97.151	Ι	1	2	124.352 II	2	4	135.484 II		4	8	183.190 IV	2 2	4	226.123 II	2	4
49.266 II	2	4	97.152	Π	1	2	124.354 III	2	4	136.495 I		2	4	184.192 II	2	4	226.125 III	2	4
49.271 III	2	4	97.153	III	1	2	124.355 III	2	4	136.500 II	[] :	2 -	4	185.198 II	2	4	226.126 III	2	4
50.278 II	2	4	97.155	III	1	2	124.356 III	2	4	136.503 II	[]	2 -	4	186.204 II	2	4	227.128 I	2	4
50.283 III	2	4	99.168	IV	2	4	124.358 III	2	4	136.504 IV	7	4	8	192.244 II	2	4	227.129 II	2	4
53.329 III	2	4	99.169	IV	2	4	124.359 III	2	4	137.507 I		2	$4 \mid$	192.251 II	1 2	4	227.130 III	2	4
55.359 III	2	4	99.170	IV	2	4	125.363 I	2	4	137.511 II	[]	2	4	195.1 I	1	2	227.132 III	2	4
58.399 III	2	4	100.172	Π	2	4	125.364 II	2	4	137.515 II	[]	2	4	195.2 II	1	2			

TABLE XI. Magnetic space groups (MSGs) which can host filling-enforced semimetals with nodal-point Fermi surfaces pinned at high-symmetry momenta.

 $\nu_{\rm P}$: The physically achievable fillings are quantized in units of $\nu_{\rm P}$

FS: Dimension of the irrep(s) dissected by the Fermi energy when the physical filling is an odd-integer-multiple of $\nu_{\rm P}$

MSG	3	ν_{P}	MSC	Ĵ	ν_{P}	MSG	ŕ	ν_{P}	MSG		ν_{P}	MSG		ν_{P}	MSG	r	ν_{P}
4.8	II	2	53.325	III	2	77.16	IV	2	102.192	IV	4	138.520	Π	4	206.38	II	4
7.25	Π	2	54.338	Π	4	77.18	IV	2	105.211	Ι	2	138.530	IV	4	208.44	Ι	2
9.38	Π	2	54.349	IV	4	78.20	Π	4	105.217	IV	4	141.553	III	2	208.45	Π	2
14.77	III	2	56.366	Π	4	80.30	Π	2	106.220	Π	4	141.554	III	2	210.52	Ι	2
14.78	III	2	57.378	Π	4	84.54	III	2	109.239	Ι	2	141.560	IV	4	210.53	Π	2
17.8	Π	2	57.390	IV	4	85.61	III	2	109.244	IV	4	142.562	Π	4	210.54	III	2
19.26	Π	4	59.405	Ι	2	86.69	III	2	110.246	Π	4	142.564	III	4	212.60	Π	4
20.32	Π	2	59.413	IV	4	86.70	III	2	119.320	IV	2	142.569	III	4	213.64	Π	4
24.54	Π	2	59.414	IV	4	88.83	III	2	120.322	Π	2	158.58	Π	2	214.67	Ι	2
28.88	Π	2	60.418	Π	4	88.84	III	2	120.323	III	2	159.62	Π	2	214.68	Π	2
29.100	Π	4	60.429	IV	4	91.104	Π	4	122.335	III	2	161.70	Π	2	216.77	IV	2
33.145	Π	4	61.434	Π	4	92.112	Π	4	125.367	III	2	169.114	Π	6	219.86	Π	2
38.194	IV	2	61.435	III	4	93.119	Ι	2	129.415	III	2	170.118	Π	6	219.87	III	2
39.196	II	2	61.437	III	4	93.120	Π	2	130.426	III	4	173.130	Π	2	224.113	III	2
40.204	Π	2	62.442	Π	4	93.124	IV	2	131.437	III	2	178.156	Π	6	227.131	III	2
41.212	Π	2	62.455	IV	4	94.127	Ι	2	132.449	III	2	179.162	Π	6	227.133	IV	4
44.233	IV	2	64.471	III	2	94.134	IV	2	133.460	Π	4	182.180	Π	2	228.135	Π	4
45.236	Π	2	64.473	III	2	95.136	Π	4	133.467	III	4	188.216	Π	2	228.137	III	4
46.242	Π	2	64.477	III	2	96.144	Π	4	133.470	IV	4	190.228	Π	2	228.138	III	4
48.257	Ι	2	68.511	Ι	2	98.157	Ι	2	134.473	III	2	198.10	Π	4	230.146	Π	4
50.277	Ι	2	70.527	Ι	2	98.158	Π	2	134.475	III	2	199.13	Π	2	230.149	III	4
50.286	IV	4	73.549	Π	4	98.161	III	2	134.480	IV	4	201.18	Ι	2			
50.288	IV	4	74.557	III	2	101.179	Ι	2	135.491	III	4	203.26	Ι	2			
52.306	II	4	76.8	Π	4	101.186	IV	4	137.516	IV	4	205.34	Π	4			
53.323	III	2	77.14	II	2	102.187	Ι	2	137.517	IV	4	205.35	III	4			

TABLE XII. Magnetic space groups (MSGs) which can host filling-enforced semimetals with movable nodal Fermi surfaces.

 $\nu_{\rm P}$: The physically achievable fillings are quantized in units of $\nu_{\rm P}$

TABLE XIII. Magnetic space groups (MSGs) which can host filling-enforced metals.

MSG	ŕ	ν_{P}	Deg	MSC	Ĵ	ν_{P}	Deg	MSG		$\nu_{\rm P}$	Deg	MSG		ν_{P}	Deg	MSG	ſ	ν_{P}	Deg	MSG	r	ν_{P}	Deg
2.5	II	1	3	59.406	II	2	1	99.166	III	1	1	123.350	IV	2	1	147.14	Π	1	3	202.22	Ι	1	1
2.6	ш	1	3	59.408	III	2	1	107.227	Ι	1	1	125.373	IV	2	1	147.15	III	1	3	202.23	Π	1	3
10.43	II	1	3	59.416	IV	2	1	107.228	Π	1	1	126.386	IV	2	1	148.18	Π	1	3	202.24	III	1	3
10.44	ш	1	3	63.458	Π	2	1	107.229	III	1	1	127.388	II	2	1	148.19	III	1	3	204.30	Ι	1	1
10.45	ш	1	3	63.460	III	2	1	107.230	Ш	1	1	127.389	III	2	1	162.74	Π	1	3	204.31	Π	1	3
12.59	II	1	3	64.470	Π	2	1	111.251	Ι	1	1	127.390	III	2	1	162.75	III	1	3	204.32	III	1	3
12.60	ш	1	3	64.472	III	2	1	111.252	Π	1	1	127.394	III	2	1	162.76	III	1	3	215.70	Ι	1	1
12.61	ш	1	3	65.481	Ι	1	1	111.253	Ш	1	1	128.400	Π	2	1	164.86	Π	1	3	215.71	Π	1	1
25.57	I	1	1	65.482	Π	1	3	111.254	Ш	1	1	128.401	III	2	1	164.87	III	1	3	215.72	III	1	1
25.58	п	1	1	65.483	Ш	1	3	111.257	IV	2	1	128.402	III	2	1	164.88	III	1	3	215.73	IV	2	1
35.165	I	1	1	65.484	Ш	1	3	111.258	IV	2	1	128.406	III	2	1	166.98	Π	1	3	216.74	Ι	1	1
35.166	п	1	1	65.487	Ш	1	3	112.266	IV	2	1	129.412	Π	2	1	166.99	III	1	3	216.75	Π	1	1
38.187	I	1	1	65.488	IV	2	1	113.268	Π	2	1	129.413	III	2	1	166.100	III	1	3	216.76	III	1	1
38.188	II	1	1	65.490	IV	2	1	113.269	III	2	1	129.414	III	2	1	175.138	Π	1	3	217.78	Ι	1	1
42.219	Ι	1	1	67.510	IV	2	1	114.276	Π	2	1	129.418	III	2	1	175.139	III	1	3	217.79	Π	1	1
42.220	п	1	1	69.521	Ι	1	1	114.277	III	2	1	129.422	IV	2	1	175.140	III	1	3	217.80	III	1	1
44.229	I	1	1	69.522	Π	1	3	115.283	Ι	1	1	131.444	IV	2	1	183.185	Ι	1	1	218.84	IV	2	1
44.230	II	1	1	69.523	III	1	3	115.284	Π	1	1	131.446	IV	2	1	183.186	Π	1	1	221.92	Ι	1	1
47.249	I	1	1	69.525	III	1	3	115.285	III	1	1	132.456	IV	2	1	187.209	Ι	1	1	221.93	Π	1	3
47.250	п	1	3	71.533	Ι	1	1	115.286	III	1	1	134.482	IV	2	1	187.210	Π	1	1	221.94	III	1	3
47.251	ш	1	3	71.534	Π	1	3	119.315	Ι	1	1	136.496	II	2	1	189.221	Ι	1	1	221.95	III	1	1
47.253	ш	1	3	71.535	ш	1	3	119.316	Π	1	1	136.497	III	2	1	189.222	Π	1	1	221.96	III	1	3
47.254	IV	2	1	71.537	III	1	3	119.317	III	1	1	136.498	III	2	1	191.233	Ι	1	1	221.97	IV	2	1
47.255	IV	2	1	81.34	Π	1	1	119.318	III	1	1	136.502	III	2	1	191.234	Π	1	3	222.103	IV	2	1
47.256	IV	2	1	81.35	III	1	1	121.327	Ι	1	1	136.506	IV	2	1	191.235	III	1	3	223.109	IV	2	1
48.264	IV	2	1	82.40	Π	1	1	121.328	Π	1	1	137.508	II	2	1	191.236	III	1	3	224.115	IV	2	1
50.287	IV	2	1	82.41	III	1	1	121.329	III	1	1	137.509	III	2	1	191.237	III	1	3	225.116	Ι	1	1
51.290	п	2	1	83.44	Π	1	3	121.330	III	1	1	137.510	III	2	1	191.241	III	1	3	225.117	Π	1	3
51.293	ш	2	1	83.45	III	1	1	123.339	Ι	1	1	137.514	III	2	1	191.242	IV	2	1	225.118	III	1	3
51.302	IV	2	1	83.46	III	1	3	123.340	Π	1	3	137.518	IV	2	1	193.254	Π	2	1	225.119	III	1	1
53.322	п	2	1	83.47	Ш	1	3	123.341	Ш	1	3	139.531	Ι	1	1	193.256	III	2	1	225.120	III	1	3
53.324	ш	2	1	87.76	Π	1	3	123.342	Ш	1	1	139.532	Π	1	3	194.264	Π	2	1	229.140	Ι	1	1
55.354	Π	2	1	87.77	Ш	1	1	123.343	Ш	1	1	139.533	III	1	3	194.267	III	2	1	229.141	Π	1	3
55.355	ш	2	1	87.78	Ш	1	3	123.344	Ш	1	3	139.534	III	1	1	200.14	Ι	1	1	229.142	III	1	3
55.356	ш	2	1	87.79	Ш	1	3	123.346	Ш	1	3	139.535	III	1	1	200.15	Π	1	3	229.143	III	1	1
58.394	II	2	1	99.163	Ι	1	1	123.347	III	1	3	139.536	III	1	3	200.16	III	1	3	229.144	III	1	3
58.395	ш	2	1	99.164	II	1	1	123.348	IV	2	1	139.538	III	1	3	200.17	IV	2	1				
58.396	Ш	2	1	99.165	Ш	1	1	123.349	IV	2	1	139.539	Ш	1	3	201.21	IV	2	1				

 $\nu_{\rm P}$: The physically achievable fillings are quantized in units of $\nu_{\rm P}$ Deg: Dimension of the manifold on which band sticking occurs, which enforces metallic behavior at fillings which are odd-integer multiples of $\nu_{\rm P}$

				0	1			<i>.</i>				2		U			· · ·
SG	ν_{P}	FS	SG	ν_{P}	FS	SG	ν_{P}	FS	SG	$\nu_{\rm P}$	FS	SG	ν_{P}	FS	SG	ν_{P}	FS
1	1	$M_{3} \rightarrow p$	40	2	$M_1 \to m$	79	1	$M_{3} \rightarrow p$	118	2	$M_1 \to p$	157	1	$M_{3} \rightarrow p$	196	1	$M_{3} \rightarrow p$
2	1	$M_3 \to M_3$	41	2	$M_1 \to m$	80	2	$\boldsymbol{p} \to \boldsymbol{m}$	119	1	$M_3 \to M_1$	158	2	$M_1 \to m$	197	1	$M_3 \to p$
3	1	$M_3 \to p$	42	1	$M_{3} \rightarrow M_{1} \\$	81	1	$M_3 \to M_1$	120	2	$M_1 \to m$	159	2	$M_1 \to m$	198	4	$\boldsymbol{p} \to \boldsymbol{m}$
4	2	$M_2 \to m$	43	2	$M_1 \to p$	82	1	$M_{3} \rightarrow M_{1} \\$	121	1	$M_3 \to M_1$	160	1	$M_{3} \rightarrow p$	199	2	$\boldsymbol{p} \to \boldsymbol{m}$
5	1	$M_3 \to p$	44	1	$M_{3} \rightarrow M_{1} \\$	83	1	$M_{\rm 3} \to M_{\rm 3}$	122	2	$M_1 \to p$	161	2	$M_1 \to m$	200	1	$M_{\rm 3} \to M_{\rm 3}$
6	1	$M_3 \to p$	45	2	$M_1 \to m$	84	2	$\boldsymbol{p} \rightarrow \boldsymbol{p}$	123	1	$M_{3} \rightarrow M_{3}$	162	1	$M_{\rm 3} \to M_{\rm 3}$	201	2	$M_1 \to p$
7	2	$M_1 \to m$	46	2	$M_1 \to m$	85	2	$M_1 \to p$	124	2	$M_1 \to p$	163	2	$M_1 \to p$	202	1	$M_3 \to M_3$
8	1	$M_3 \to p$	47	1	$M_{\rm 3} \to M_{\rm 3}$	86	2	$M_1 \to p$	125	2	$M_1 \to p$	164	1	$M_{\rm 3} \to M_{\rm 3}$	203	2	$M_1 \to p$
9	2	$M_1 \to m$	48	2	$M_1 \to p$	87	1	$M_{\rm 3} \to M_{\rm 3}$	126	2	$M_1 \to p$	165	2	$M_1 \to p$	204	1	$M_{\rm 3} \to M_{\rm 3}$
10	1	$M_3 \to M_3$	49	2	$M_1 \to p$	88	2	$M_1 \to p$	127	2	$M_2 \to M_1$	166	1	$M_{3} \rightarrow M_{3}$	205	4	$M_1 \to m$
11	2	$M_2 \to p$	50	2	$M_1 \to p$	89	1	$M_{3} \rightarrow p$	128	2	$M_2 \to M_1$	167	2	$M_1 \to p$	206	4	$\boldsymbol{p} \to \boldsymbol{m}$
12	1	$M_3 \to M_3$	51	2	$M_2 \to M_1$	90	2	$M_2 \to p$	129	2	$M_2 \to M_1$	168	1	$M_{3} \rightarrow p$	207	1	$M_3 \to p$
13	2	$M_1 \to p$	52	4	$\boldsymbol{p} \to \boldsymbol{m}$	91	4	$\boldsymbol{m} \to \boldsymbol{m}$	130	4	$\mathbf{p} \to \mathbf{p}$	169	6	$M_2 \to m$	208	2	$\boldsymbol{m} \to \boldsymbol{m}$
14	2	$M_2 \to p$	53	2	$M_2 \to M_1$	92	4	$\boldsymbol{p} \to \boldsymbol{m}$	131	2	$M_1 \to p$	170	6	$M_2 \to m$	209	1	$M_3 \to p$
15	2	$M_1 \to p$	54	4	$\boldsymbol{p} \to \boldsymbol{m}$	93	2	$\boldsymbol{m} \to \boldsymbol{m}$	132	2	$M_1 \to p$	171	3	$M_{3} \rightarrow p$	210	2	$\boldsymbol{p} \to \boldsymbol{m}$
16	1	$M_3 \to p$	55	2	$M_2 \to M_1$	94	2	$M_2 \to p$	133	4	$\boldsymbol{m} \to \boldsymbol{m}$	172	3	$M_3 \to p$	211	1	$M_3 \to p$
17	2	$M_2 \to m$	56	4	$\boldsymbol{p} \to \boldsymbol{m}$	95	4	$\boldsymbol{m} \to \boldsymbol{m}$	134	2	$M_1 \to p$	173	2	$M_2 \to m$	212	4	$\boldsymbol{p} \to \boldsymbol{m}$
18	2	$M_2 \to p$	57	4	$M_1 \to m$	96	4	$\boldsymbol{p} \rightarrow \boldsymbol{m}$	135	4	$\boldsymbol{p} \rightarrow \boldsymbol{p}$	174	1	$M_{3} \rightarrow p$	213	4	$\boldsymbol{p} \to \boldsymbol{m}$
19	4	$\boldsymbol{p} \to \boldsymbol{m}$	58	2	$M_2 \to M_1$	97	1	$M_3 \to p$	136	2	$M_2 \to M_1$	175	1	$M_3 \to M_3$	214	2	$\boldsymbol{p} \to \boldsymbol{m}$
20	2	$M_2 \to m$	59	2	$M_2 \to M_1$	98	2	$\boldsymbol{p} \to \boldsymbol{m}$	137	2	$M_2 \to M_1$	176	2	$M_2 \to p$	215	1	$M_3 \to M_1$
21	1	$M_3 \to p$	60	4	$M_1 \to m$	99	1	$M_3 \to M_1$	138	4	$\boldsymbol{p} \to \boldsymbol{m}$	177	1	$M_{3} \rightarrow p$	216	1	$M_3 \to M_1$
22	1	$M_3 \to p$	61	4	$M_1 \to m$	100	2	$M_1 \to p$	139	1	$M_3 \rightarrow M_3$	178	6	$M_2 \to m$	217	1	$M_3 \to M_1$
23	1	$M_3 \to p$	62	4	$M_1 \to m$	101	2	$M_1 \to p$	140	2	$M_1 \to p$	179	6	$M_2 \to m$	218	2	$M_1 \to p$
24	2	$\textbf{p} \rightarrow \textbf{m}$	63	2	$M_2 \to M_1$	102	2	$M_1 \to p$	141	2	$M_1 \to p$	180	3	$M_3 \to p$	219	2	$M_1 \to m$
25	1	$M_3 \to M_1$	64	2	$M_2 \to M_1$	103	2	$M_1 \to p$	142	4	$\boldsymbol{p} \rightarrow \boldsymbol{m}$	181	3	$M_3 \to p$	220	2	$M_1 \to p$
26	2	$M_2 \to p$	65	1	$M_3 \to M_3$	104	2	$M_1 \to p$	143	1	$M_3 \to p$	182	2	$M_2 \to m$	221	1	$M_3 \to M_3$
27	2	$M_1 \to p$	66	2	$M_1 \to p$	105	2	$M_1 \to p$	144	3	$M_3 \to p$	183	1	$M_3 \to M_1$	222	2	$M_1 \to p$
28	2	$M_1 \to m$	67	2	$M_1 \to p$	106	4	$\boldsymbol{m} \rightarrow \boldsymbol{m}$	145	3	$M_3 \to p$	184	2	$M_1 \to p$	223	2	$M_1 \to p$
29	4	$\textbf{p} \rightarrow \textbf{m}$	68	2	$M_1 \to p$	107	1	$M_3 \to M_1$	146	1	$M_3 ightarrow p$	185	2	$M_2 \to p$	224	2	$M_1 \to p$
30	2	$M_1 \to p$	69	1	$M_3 ightarrow M_3$	108	2	$M_1 \to p$	147	1	$M_3 ightarrow M_3$	186	2	$M_2 \to p$	225	1	$M_3 \to M_3$
31	2	$M_2 \to p$	70	2	$M_1 \to p$	109	2	$M_1 \to p$	148	1	$M_3 \to M_3$	187	1	$M_3 \to M_1$	226	2	$M_1 \to p$
32	2	$M_1 \to p$	71	1	$M_3 \to M_3$	110	4	$\boldsymbol{p} \to \boldsymbol{m}$	149	1	$M_3 \to p$	188	2	$M_1 \to m$	227	2	$M_1 \to p$
33	4	$\textbf{p} \rightarrow \textbf{m}$	72	2	$M_1 \to p$	111	1	$M_3 \to M_1$	150	1	$M_3 \to p$	189	1	$M_3 \to M_1$	228	4	$\boldsymbol{p} \to \boldsymbol{m}$
34	2	$M_1 \to p$	73	4	$\boldsymbol{p} \to \boldsymbol{m}$	112	2	$M_1 \to p$	151	3	$M_3 \to p$	190	2	$M_1 \to m$	229	1	$M_3 \to M_3$
35	1	$M_3 \to M_1$	74	2	$M_1 \to p$	113	2	$M_2 \to M_1$	152	3	$M_3 \to p$	191	1	$M_3 \to M_3$	230	4	$\boldsymbol{p} \to \boldsymbol{m}$
36	2	$M_2 \to p$	75	1	$M_3 \to p$	114	2	$M_2 \to M_1$	153	3	$M_3 \to p$	192	2	$M_1 \to p$			
37	2	$M_1 \to p$	76	4	$\boldsymbol{m} \rightarrow \boldsymbol{m}$	115	1	$M_3 \to M_1$	154	3	$M_3 \to p$	193	2	$M_2 \to M_1$			
38	1	$M_3 \to M_1$	77	2	$\boldsymbol{m} \rightarrow \boldsymbol{m}$	116	2	$M_1 \to p$	155	1	$M_3 \to p$	194	2	$M_2 \to M_1$			
39	2	$M_1 \rightarrow m$	78	4	$\boldsymbol{m} \to \boldsymbol{m}$	117	2	$M_1 \rightarrow p$	156	1	$M_3 \rightarrow p$	195	1	$M_3 ightarrow p$			

TABLE XIV. Effect of turning on spin-orbit coupling on the fermiology of time-reversal symmetric filling-enforced (semi-)metals

SG: Space group describing the spatial symmetry of a material with time-reversal symmetry

 $\nu_{\rm P}$: The physically achievable fillings are quantized in units of $\nu_{\rm P}$

FS: Fermiology for a material with a filling being an odd-integer multiple of $\nu_{\rm P}$, dubbed a filling-enforced (semi-)metal M_d : Filling-enforced metals due to band sticking on d-dimensional manifolds

p: Filling-enforced semimetals with pinned, nodal-point Fermi surfaces

m: Filling-enforced semimetals with movable nodal Fermi surfaces

Appendix C: Notations

For a MSG $\mathcal{M} = \mathcal{G} + \mathcal{A}$, $g \in \mathcal{G}$ maps the position \boldsymbol{x} and the momentum \boldsymbol{k} to

$$g(\boldsymbol{x}) \equiv p_g \boldsymbol{x} + \boldsymbol{t}_g, \ g(\boldsymbol{k}) \equiv p_g(\boldsymbol{k}), \ p_g \in O(3),$$
 (C1)

while $a \in \mathcal{A}$ acts on them as

$$a(\mathbf{x}) = p_a \mathbf{x} + \mathbf{t}_a, \quad a(\mathbf{k}) = -p_a \mathbf{k}, \ p_a \in O(3).$$
 (C2)

The little group of k is defined by

$$\mathcal{M}_{\boldsymbol{k}} = \{ m \in \mathcal{M} \mid m(\boldsymbol{k}) = \boldsymbol{k} \} = \mathcal{G}_{\boldsymbol{k}} + \mathcal{A}_{\boldsymbol{k}}, \quad (C3)$$

where \mathcal{G}_{k} is the unitary part and \mathcal{A}_{k} is the anti-unitary part:

$$\mathcal{G}_{\boldsymbol{k}} = \{ g \in \mathcal{G} \mid g(\boldsymbol{k}) = \boldsymbol{k} + \boldsymbol{G} \},$$
(C4)

$$\mathcal{A}_{\boldsymbol{k}} = \{ a \in \mathcal{A} \, | \, a(\boldsymbol{k}) = \boldsymbol{k} + \boldsymbol{G} \}$$
(C5)

and G is a reciprocal lattice vector. \mathcal{G}_k is a subgroup of \mathcal{M}_k but \mathcal{A}_k is not even a group.

Similarly, the site symmetry group of x is defined by

$$\mathcal{M}_{\boldsymbol{x}} = \{ m \in \mathcal{G} \mid m(\boldsymbol{x}) = \boldsymbol{x} \} = \mathcal{G}_{\boldsymbol{x}} + \mathcal{A}_{\boldsymbol{x}}.$$
(C6)

where $\mathcal{G}_{\boldsymbol{x}}$ is the unitary part and $\mathcal{A}_{\boldsymbol{x}}$ is the anti-unitary part:

$$\mathcal{G}_{\boldsymbol{x}} = \{ g \in \mathcal{G} \, | \, g(\boldsymbol{x}) = \boldsymbol{x} \}, \tag{C7}$$

$$\mathcal{A}_{\boldsymbol{x}} = \{ a \in \mathcal{A} \, | \, a(\boldsymbol{x}) = \boldsymbol{x} \}. \tag{C8}$$

Since we are interested in systems of spinful electrons with spin-orbit coupling, we assume a projective representation, i.e.,

$$u(g)u(g') = z_{g,g'}u(gg'),$$
 (C9)

$$u(g)u(a') = z_{g,a'}u(ga'),$$
 (C10)

$$u(a)u(g')^* = z_{a,g'}u(ag'),$$
 (C11)

$$u(a)u(a')^* = z_{a,a'}u(aa').$$
 (C12)

For example, $z_{\mathcal{T},\mathcal{T}} = -1$ for the time-reversal symmetry \mathcal{T} . For spinless electrons, all z should be set to be unity in the following discussions.

Appendix D: Compatibility relations

Here we review the compatibility relations among the integers $n = \{n_k^{\alpha}\}$ to be consistent with $\mathcal{M} = \mathcal{G} + \mathcal{A}$.

1. Basics

Suppose that $\{|\mathbf{k}, i\rangle\}$ is a basis of an irrep $u_{\mathbf{k}}^{\alpha}$ of $\mathcal{G}_{\mathbf{k}}$, i.e.,

$$\hat{h}|\boldsymbol{k},i\rangle = |\boldsymbol{k},j\rangle [u_{\boldsymbol{k}}^{\alpha}(h)]_{ji}, \quad h \in \mathcal{G}_{\boldsymbol{k}}.$$
 (D1)

Here we show that $\{\hat{g}|\boldsymbol{k},i\rangle\}$ $(g \in \mathcal{G})$ forms a basis of the irrep

$$u_{g(\mathbf{k})}^{\alpha'}(h') \equiv \rho_{h',g} u_{\mathbf{k}}^{\alpha}(g^{-1}h'g), \quad h' \in \mathcal{G}_{g(\mathbf{k})}.$$
(D2)

Here, $\rho_{h,g}$ is related to the projective factor $z_{g,g'} = \pm 1$ as

$$\rho_{h,g} = \frac{z_{h,g}}{z_{g,g^{-1}hg}} = \pm 1.$$
(D3)

To see this, note that $h' \in \mathcal{G}_{g(k)}$ can be written as $h' = ghg^{-1}$ using $h \in \mathcal{G}_k$. Then,

$$\begin{aligned} h'(\hat{g}|\boldsymbol{k},i\rangle) &= z_{h',g}(h'g)|\boldsymbol{k},i\rangle \\ &= z_{h',g}(\hat{g}h)|\boldsymbol{k},i\rangle = \frac{z_{h',g}}{z_{g,h}}\hat{g}(\hat{h}|\boldsymbol{k},i\rangle) \\ &= \rho_{h',g}\hat{g}(|\boldsymbol{k},j\rangle[u_{\boldsymbol{k}}^{\alpha}(h)]_{ji}) \\ &= (\hat{g}|\boldsymbol{k},j\rangle)\rho_{h',g}[u_{\boldsymbol{k}}^{\alpha}(g^{-1}h'g)]_{ji}. \end{aligned}$$
(D4)

Similarly, $\{\hat{a} | \boldsymbol{k}, i \rangle\}$ $(a \in \mathcal{A})$ forms a basis of the irrep

$$u_{a(\mathbf{k})}^{\alpha'}(h') \equiv \rho_{h',a} [u_{\mathbf{k}}^{\alpha}(a^{-1}h'a)]_{ji}^{*}, \quad h' \in \mathcal{G}_{g(\mathbf{k})}.$$
(D5)

where

$$\rho_{h,a} = \frac{z_{h,a}}{z_{a,a^{-1}ha}} = \pm 1.$$
 (D6)

Again writing $h' \in \mathcal{G}_{a(k)}$ as $h' = aha^{-1}$ using $h \in \mathcal{G}_k$,

$$\hat{h}'(\hat{a}|\boldsymbol{k},i\rangle) = z_{h',a}(\hat{h}'a)|\boldsymbol{k},i\rangle$$

$$= z_{h',a}(\hat{a}h)|\boldsymbol{k},i\rangle = \frac{z_{h',a}}{z_{a,h}}\hat{a}(\hat{h}|\boldsymbol{k},i\rangle)$$

$$= \rho_{h',a}\hat{a}(|\boldsymbol{k},j\rangle[u_{\boldsymbol{k}}^{\alpha}(h)]_{ji})$$

$$= (\hat{a}|\boldsymbol{k},j\rangle)\rho_{h',a}[u_{\boldsymbol{k}}^{\alpha}(a^{-1}h'a)]_{ji}^{*}.$$
(D7)

2. Compatibility relations from the unitary part $C_{\mathcal{G}}$

Here we summarize the compatibility relations arising from the unitary part \mathcal{G} of \mathcal{M} . There are three major categories:

(i) The representations used at symmetry-related momenta are related to each other. Suppose that an irrep $u_{\mathbf{k}}^{\alpha}$ of $\mathcal{G}_{\mathbf{k}}$ is used $n_{\mathbf{k}}^{\alpha}$ times. When $g \notin \mathcal{G}_{\mathbf{k}}$, \mathbf{k} and $g(\mathbf{k})$ are symmetry-related but distinct momenta. Let $\{|\mathbf{k}, i\rangle\}$ be the basis of the representation. As explained above, the g-transformed copy, $\{\hat{g}|\mathbf{k},i\rangle\}$ forms a basis of the irrep $u_{g(\mathbf{k})}^{\alpha'}$ in Eq. (D1). It implies

$$n_{g(\boldsymbol{k})}^{\alpha'} = n_{\boldsymbol{k}}^{\alpha}.$$
 (D8)

(ii) At a high-symmetry momenta slightly off from a higher-symmetry point k, the continuity of the band structure requires that

$$n_{\boldsymbol{k}+\delta(\boldsymbol{k})}^{\beta} = \sum_{\alpha} n_{\boldsymbol{k}}^{\alpha} m^{\alpha\beta}, \qquad (D9)$$

where nonnegative integers $m^{\alpha\beta}$ are the coefficients appearing in the decomposition $u_{\mathbf{k}}^{\alpha} = \sum_{\beta} c^{\alpha\beta} u_{\mathbf{k}+\delta(\mathbf{k})}^{\beta}$ regarding $u_{\mathbf{k}}^{\alpha}$ as a representation of $\mathcal{G}_{\mathbf{k}+\delta(\mathbf{k})} < \mathcal{G}_{\mathbf{k}}$.

(iii) For nonsymmorphic operations g with a 'fractional translation', such as glide reflections or screw rotations,

$$\sum_{\alpha} n_{\boldsymbol{k}}^{\alpha} \operatorname{tr} u_{\boldsymbol{k}}^{\alpha}(g) = 0, \qquad (D10)$$

again due to the continuity of the band structure.

3. Compatibility relations from the anti-unitary part \tilde{C}_{A}

The anti-unitary part A adds two more constraints.

(iv) \mathcal{A} may relate two momenta that were not related by \mathcal{G} . This occurs when the anti-unitary part \mathcal{A}_{k} is empty. Suppose that an irrep u_{k}^{α} of \mathcal{G}_{k} is used n_{k}^{α} times. Let $\{|\mathbf{k}, i\rangle\}$ be the basis of the representation. The *a*-transformed copy, $\{\hat{a}|\mathbf{k}, i\rangle\}$ forms a basis of the irrep $u_{a(\mathbf{k})}^{\alpha'}$ in Eq. (D5). It requires a constraint

$$n_{a(\boldsymbol{k})}^{\alpha'} = n_{\boldsymbol{k}}^{\alpha}.$$
 (D11)

(v) When the anti-unitary part \mathcal{A}_{k} is not empty, $u_{a(k)}^{\alpha'}$ in Eq. (D5) is also an irrep at k if $a \in \mathcal{A}_{k}$. The nature of paring due to \mathcal{A}_{k} can be judged by the Herring rule [29]:

$$\xi_{\boldsymbol{k}}^{\alpha} = \frac{1}{|\mathcal{A}_{\boldsymbol{k}}/T|} \sum_{a \in \mathcal{A}_{\boldsymbol{k}}/T} \delta_{a(\boldsymbol{k}),\boldsymbol{k}} z_{a,a} \, \chi_{\boldsymbol{k}}^{\alpha}(a^2), \quad (\text{D12})$$

where $\chi_{\mathbf{k}}^{\alpha}(h) \equiv \operatorname{tr} u_{\mathbf{k}}^{\alpha}(h)$ is the character of the irrep. When $\xi_{\mathbf{k}}^{\alpha} = 0, u_{\mathbf{k}}^{\alpha'}$ and $u_{\mathbf{k}}^{\alpha}$ are inequivalent and $n_{\mathbf{k}}^{\alpha'} = n_{\mathbf{k}}^{\alpha} (\alpha' \neq \alpha)$ is required. When $\xi_{\mathbf{k}}^{\alpha} = -1$ or $+1, u_{\mathbf{k}}^{\alpha'}$ and $u_{\mathbf{k}}^{\alpha}$ are equivalent $(\alpha' = \alpha)$. When $\xi_{\mathbf{k}}^{\alpha} = -1, \{\hat{a}|\mathbf{k},i\rangle\}$ and $\{|\mathbf{k},i\rangle\}$ are orthogonal to each other and $n_{\mathbf{k}}^{\alpha}$ must be an even integer. When $\xi_{\mathbf{k}}^{\alpha} = +1, \{\hat{a}|\mathbf{k},i\rangle\}$ and $\{|\mathbf{k},i\rangle\}$ are linearly dependent and there is no extra constraint on $n_{\mathbf{k}}^{\alpha}$.

Appendix E: Details of Atomic insulators

As explained in the main text, an AI can be specified by choosing the position x in real space at which electrons are localized and the type of orbital on the site. The type of orbital is a (co-)representation of the site symmetry group \mathcal{M}_x in Eq. (C6).

Let u_x^r (r = 1, 2, ...) be an irrep of \mathcal{G}_x . When \mathcal{A}_x is empty, u_x^r is already an irrep of \mathcal{M}_x . We put an orbital u_x^r on the site x. One can construct an AI by placing a symmetry-related copy on each site of an \mathcal{M} -symmetric lattice $\mathcal{M}x \equiv \{mx \mid m \in \mathcal{M}\}$. In this case the lattice $\mathcal{M}x$ contains twice as many number of site as $\mathcal{G}x$.

On the other hand, when \mathcal{A}_x is not empty, \mathcal{A}_x dictates a certain pairing among u_x^r 's, leading to a co-irrep of \mathcal{M}_x . The details of paring can be again judged by the Herring rule [29]:

$$\xi_{\boldsymbol{x}}^{r} = \frac{1}{|\mathcal{A}_{\boldsymbol{x}}|} \sum_{a \in \mathcal{A}_{\boldsymbol{x}}} z_{a,a} \, \chi_{\boldsymbol{x}}^{r}(a^{2}). \tag{E1}$$

When $\xi_x^r = 0$, $u_x^{r'}$ and u_k^r are inequivalent $(r' \neq r)$ and must be paired to form a co-irrep of \mathcal{M}_x . When $\xi_x^r = -1$ or +1, $u_x^{r'}$ and u_x^r are equivalent (r' = r). When $\xi_x^r = -1$, $\{\hat{a}|x,i\rangle\}$ and $\{|x,i\rangle\}$ are orthogonal to each other, and u_x^r is 'paired with itself'. Finally, when $\xi_x^r = +1$, $\{\hat{a}|x,i\rangle\}$ and $\{|x,i\rangle\}$ are linearly dependent and u_x^r alone stands a co-irrep of \mathcal{M}_x . We put a co-irrep on the site x and generate AI by placing symmetry-related copy of the orbital on each site of the lattice $\mathcal{M}x = \mathcal{G}x$.

Appendix F: The details of $\tilde{\mathcal{T}}b$

Let us write $\boldsymbol{b} = \{n_{\boldsymbol{k}}^{\alpha}\}$ and $\tilde{\mathcal{T}}\boldsymbol{b} = \{\tilde{n}_{\boldsymbol{k}}^{\alpha}\}$. By definition, \boldsymbol{b} has the representation $\bigoplus_{\alpha} n_{\boldsymbol{k}}^{\alpha} u_{\boldsymbol{k}}^{\alpha}$ of $\mathcal{G}_{\boldsymbol{k}}$ at \boldsymbol{k} . Then $a = \tilde{\mathcal{T}}$ maps it to the representation $\bigoplus_{\alpha} n_{\boldsymbol{k}}^{\alpha} u_{a(\boldsymbol{k})}^{\alpha'}$ of $\mathcal{G}_{a(\boldsymbol{k})}$ at $a(\boldsymbol{k})$ [see Eq. (D5)]. By definition of $\tilde{\mathcal{T}}\boldsymbol{b}$, this coincides with $\bigoplus_{\alpha} \tilde{n}_{a(\boldsymbol{k})}^{\alpha} u_{a(\boldsymbol{k})}^{\alpha}$, i.e.,

$$\oplus_{\alpha} \tilde{n}^{\alpha}_{a(\mathbf{k})} u^{\alpha}_{a(\mathbf{k})}(h) = \oplus_{\alpha} n^{\alpha}_{\mathbf{k}} u^{\alpha'}_{a(\mathbf{k})}(h).$$
(F1)

Solving for $\tilde{n}_{\boldsymbol{k}}^{\alpha}$, we get

$$\tilde{n}_{\boldsymbol{k}}^{\beta} = \frac{1}{|\mathcal{G}_{\boldsymbol{k}}/T|} \sum_{h \in \mathcal{G}_{\boldsymbol{k}}/T} \chi_{\boldsymbol{k}}^{\beta}(h)^* \sum_{\alpha} n_{a^{-1}(\boldsymbol{k})}^{\alpha} \chi_{\boldsymbol{k}}^{\alpha'}(h).$$
(F2)

Appendix G: Magnetic layer groups

For each MLG, there is a corresponding MSG that can be achieved by stacking of 2D layers. Conversely, if an MSG \mathcal{M} is such that every element of $m \in \mathcal{M}$ maps, say, z to either $n \pm z$ ($n \in \mathbb{Z}$) (i.e., no mixing of z with x or y or no fractional translation along z), the MSG can be reduced to an MLG by projecting down to xy plane. (In the actual calculation, it suffices to set $k_z = 0$). The projection down to yz and zx planes, if allowed, may also lead to an MSG. These three possible ways of projections lead to sometimes distinct and sometimes the identical MLGs. We present the computation results in in Appendix A in a redundant way ---for each 'parent' MSG, we list the computed numbers for all possible directions of projections among the three, regardless of their dependence/independence as MLGs. For example, for MSG 3.4, two projections down to xy and xz planes are possible and correspondingly we have two rows 3.4(1) and 3.4 (2). For this MSG, the two projections correspond to two different MLGs. On the other hand, for the type II MSG 2.5, the three ways of projection lead to an identical MLG. Even in this case the redundant presentation has an advantage: e.g. it tells us that the three \mathbb{Z}_2 factor of $X_{BS} = (\mathbb{Z}_2)^3 \times \mathbb{Z}_4$ for MSG 2.5 can actually be attributed to the thee orthogonal 2D planes, implying the presence of 'weak' phases.

Appendix H: The tight-binding model for representation-enforced semimetal

Here we present a tight-binding model that realizes the 2D reSM described by the main text. We assume the MLG p_a 112. There are two sites in a unit cell. The A-sublattice is at $\boldsymbol{x} = n_1\boldsymbol{a}_1 + n_2\boldsymbol{a}_2$ $(n_1, n_2 \in \mathbb{Z})$ and the B-sublattice is at $\boldsymbol{x} = (n_1 + \frac{1}{2})\boldsymbol{a}_1 + n_2\boldsymbol{a}_2$ $(n, m \in \mathbb{Z})$. For simplicity let us set $\boldsymbol{a}_1 = (2, 0), \boldsymbol{a}_2 = (-1, 1)$. In the following, four by four matrices are in the basis $(|A, \uparrow\rangle, |A, \downarrow\rangle, |B, \uparrow\rangle, |B, \downarrow\rangle)$. We

assume the standard symmetry representation:

$$U_{\boldsymbol{k}}(C_{2z}) = i \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & e^{-ik_1} & 0 \\ 0 & 0 & 0 & -e^{-ik_1} \end{pmatrix}, \quad (H1)$$
$$U_{\boldsymbol{k}}(\mathcal{T}T_{\frac{1}{2}\boldsymbol{a}_1}) = \begin{pmatrix} 0 & 0 & 0 & -e^{ik_1} \\ 0 & 0 & e^{ik_1} & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \quad (H2)$$

satisfying

$$U_{-k}(C_{2z})U_k(C_{2z}) = -1,$$
 (H3)

$$U_{-k}(\mathcal{T}T_{\frac{1}{2}a_1})U_k(\mathcal{T}T_{\frac{1}{2}a_1})^* = -e^{-ik_1}.$$
 (H4)

An example of the tight-binding Hamiltonian in the momentum space reads

$$h(\boldsymbol{k}) = \sum_{\mu,\nu=0,4} g_{\mu,\nu}(\boldsymbol{k}) \, \sigma_{\mu} \otimes \sigma_{\nu}, \tag{H5}$$

$$g_{1,0}(\mathbf{k}) = t[1 + \cos k_1 + \cos k_2 + \cos(k_1 + k_2)],$$
 (H6)

$$g_{2,0}(\mathbf{k}) = t[\sin k_1 - \sin k_2 + \sin(k_1 + k_2)]$$
(H7)

$$g_{1,1}(\mathbf{k}) = t_J [1 - \cos k_1 + \cos k_2 - \cos(k_1 + k_2)],$$
 (H8)

$$g_{2,1}(\mathbf{k}) = -t_J [\sin k_1 + \sin k_2 + \sin(k_1 + k_2)], \quad (H9)$$

$$g_{3,3}(k) = J.$$
 (H10)

It is easy to check the symmetry of the Hamiltonian:

$$U_{\boldsymbol{k}}(C_{2z})h(\boldsymbol{k}) = h(-\boldsymbol{k})U_{\boldsymbol{k}}(C_{2z}), \tag{H11}$$

$$U_{\boldsymbol{k}}(\mathcal{T}T_{\frac{1}{2}\boldsymbol{a}_1})h(\boldsymbol{k})^* = h(-\boldsymbol{k})U_{\boldsymbol{k}}(\mathcal{T}T_{\frac{1}{2}\boldsymbol{a}_1}). \quad (H12)$$

When $|t_J| < \frac{1}{2}J$, $\eta = +1$ and the system is a band insulator. When $|t_J| > \frac{1}{2}J$, $\eta = -1$ and the system realizes a Dirac semimetal.

The MLG corresponding to the type III MSG 13.69 (P112'/a') can also host a Dirac semimetal through an almost identical mechanism. The MLG is generated by T_{a_2} , I (the 3D inversion), and $\mathcal{T}G$ ($G \equiv T_{\frac{1}{2}a_1}IC_{2z}$ is the glide symmetry). The corresponding topological invariant η is defined by

$$\eta = \prod_{n:\text{occupied}} \prod_{\boldsymbol{k}=(0,0),(0,\pi)} \eta_{n,\boldsymbol{k}},$$
(H13)

where $\eta_{n,k}$ represents the inversion eigenvalue ± 1 .

Appendix I: Tables for spinless electrons

Here we include the following tables for spinless electrons:

- Tables XV–XX: d, X_{BS} , and ν_{BS} for MSGs.
- Table XXI: $\{\nu\}_{BS}$ and $\{\nu\}_{AI}$ for exceptional MSGs.

Similar to the remarks in Appendix A, some previously published results are reproduced below for completeness.

TABLE XV. Characterization of magnetic space groups (MSGs) in the triclinic family for spinless electrons.

MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MS	SG	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MS	SG	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$
1.1 I	1	(1)	1	1.3	IV	1	(1)	2	2.5	II	9	(2, 2, 2, 4)	1	2.7 IV	5	(2, 2, 4)	2
1.2 II	1	(1)	1	2.4	Ι	9	(2, 2, 2, 4)	1	2.6	III	1	(1)	1				

 \overline{d} : Rank of the band structure group {BS} X_{BS} : Symmetry-based indicators of band topology

 ν_{BS} : Set of ν bands are symmetry-forbidden from being isolated by band gaps if $\nu \notin \nu_{BS} \mathbb{Z}$

							-								-		_		
MS	G	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MS	G	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSC	3	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG	G	d	$X_{\rm BS}$	$\nu_{\rm BS}$
3.1	Ι	5	(2)	1	7.24	Ι	1	(1)	2	10.47	IV	9	(2, 2, 2)	2	13.70	IV	4	(2, 2)	4
3.2	Π	5	(2)	1	7.25	II	1	(1)	2	10.48	IV	10	(2, 2)	2	13.71	IV	5	(2, 2)	4
3.3	III	1	(1)	1	7.26	III	1	(1)	2	10.49	IV	8	(2, 2)	2	13.72	IV	7	(2, 2)	2
3.4	IV	3	(2)	2	7.27	IV	1	(1)	4	11.50	Ι	6	(2)	2	13.73	IV	5	(2)	2
3.5	IV	5	(2)	2	7.28	IV	1	(1)	2	11.51	Π	6	(2)	2	13.74	IV	4	(2)	4
3.6	IV	3	(1)	2	7.29	IV	1	(1)	4	11.52	Ш	2	(1)	2	14.75	Ι	5	(2)	2
4.7	Ι	1	(1)	2	7.30	IV	1	(1)	4	11.53	Ш	1	(1)	2	14.76	Π	5	(2)	2
4.8	Π	1	(1)	2	7.31	IV	1	(1)	2	11.54	Ш	5	(2, 2, 4)	2	14.77	III	1	(1)	2
4.9	III	1	(1)	2	8.32	Ι	2	(1)	1	11.55	IV	4	(2)	4	14.78	III	1	(1)	2
4.10	IV	1	(1)	4	8.33	Π	2	(1)	1	11.56	IV	6	(2)	2	14.79	III	5	(2, 4)	2
4.11	IV	1	(1)	2	8.34	III	1	(1)	1	11.57	IV	4	(2)	2	14.80	IV	3	(2)	4
4.12	IV	1	(1)	2	8.35	IV	2	(1)	2	12.58	Ι	10	(2, 2)	1	14.81	IV	3	(2)	4
5.13	Ι	3	(1)	1	8.36	IV	2	(1)	2	12.59	Π	10	(2, 2)	1	14.82	IV	3	(2)	4
5.14	Π	3	(1)	1	9.37	Ι	1	(1)	2	12.60	Ш	2	(1)	1	14.83	IV	5	(2)	2
5.15	III	1	(1)	1	9.38	II	1	(1)	2	12.61	Ш	3	(1)	1	14.84	IV	3	(2)	4
5.16	IV	2	(1)	2	9.39	III	1	(1)	2	12.62	Ш	7	(2, 2, 4)	1	15.85	Ι	6	(2, 2)	2
5.17	IV	3	(2)	2	9.40	IV	1	(1)	2	12.63	IV	6	(2, 2)	2	15.86	Π	6	(2, 2)	2
6.18	Ι	3	(1)	1	9.41	IV	1	(2)	2^*	12.64	IV	8	(2, 2)	2	15.87	III	1	(1)	2
6.19	Π	3	(1)	1	10.42	Ι	15	(2, 2, 2)	1	13.65	Ι	7	(2, 2)	2	15.88	III	2	(1)	2
6.20	III	1	(1)	1	10.43	II	15	(2, 2, 2)	1	13.66	Π	7	(2, 2)	2	15.89	III	5	(2, 4)	2
6.21	IV	3	(1)	2	10.44	III	3	(1)	1	13.67	III	1	(1)	2	15.90	IV	5	(2, 2)	2
6.22	IV	2	(1)	2	10.45	III	5	(2)	1	13.68	III	3	(2)	2	15.91	IV	4	(2, 2, 2)	2^*
6.23	IV	2	(1)	2	10.46	III	9	(2, 2, 2, 4)	1	13.69	III	5	(2, 2, 4)	2					

TABLE XVI. Characterization of magnetic space groups (MSGs) in the monoclinic family for spinless electrons.

d: Rank of the band structure group {BS}

 $X_{\rm BS}$: Symmetry-based indicators of band topology

 ν_{BS} : Set of ν bands are symmetry-forbidden from being isolated by band gaps if $\nu \notin \nu_{BS} \mathbb{Z}$

*: Exhibiting exceptional filling pattern; see Table XXI

MSG MSG $d X_{\rm BS} \nu_{\rm BS}$ MSG $d X_{\rm BS} \nu_{\rm BS}$ $d X_{\rm BS} \nu_{\rm BS}$ $d X_{\rm BS} \nu_{\rm BS}$ MSG 16.1 I 13(1)1 23.51 III 3 (1) 1 29.101 III 1 (1) 33.151 IV 1 4 (1)4 16.2 II 13(1)1 23.52 IV 5 (2) $\mathbf{2}$ 29.102 III 1 (1) 4 33.152 IV 1 (1)4 $\mathbf{2}$ 16.3 III 5(2)1 24.53 Ι |4(1)|29.103 III 1 (1) 33.153 IV 1 (1)4 4 16.4 IV 9 (2) $\mathbf{2}$ 24.54 II |4(1)|229.104 IV 1 (1) 4 33.154 IV 1 (1)4 24.55 III 2 (1) 16.5 IV 7(2) $\mathbf{2}$ $\mathbf{2}$ 29.105 IV 1 (2) 33.155 IV 1 4^{*} (1)4 $\mathbf{2}$ 16.6 IV 7(2) $\mathbf{2}$ 24.56 IV 4 (2) $\mathbf{2}$ 29.106 IV 1 (1) 34.156 I 3 4 (1)17.7 (1) $\mathbf{2}$ 25.57 Ι |9(1)|29.107 IV 1 (1) 34.157 II 3 $\mathbf{2}$ Ι 51 4 (1)25.58 II 9 (1) 29.108 IV 1 (1) 34.158 III 1 17.8 II 5(1) $\mathbf{2}$ 1 4 (1)217.9 III 1 (1) $\mathbf{2}$ 25.59 III 3 (1) 1 29.109 IV 1 (1) 4 34.159 III 3 (1) $\mathbf{2}$ 3 25.60 III 5 (2) 29.110 IV 1 (1) 34.160 IV 2 17.10 III (2)21 (1)4 4 17.11 IV 4 (2)4 25.61 IV 9 (1) 230.111 I 3 (1) 234.161 IV 3 (2)4 17.12 IV 5(1) $\mathbf{2}$ 25.62 IV 6 (1) $\mathbf{2}$ 30.112 II 3 (1) 34.162 IV 2 2(1)4 17.13 IV 4 $\mathbf{2}$ 25.63 IV 5 (1) $\mathbf{2}$ 30.113 III 1 (1) 34.163 IV 3 (1) $\mathbf{2}$ (2)4 17.14 IV 3 (2)4 25.64 IV 6 (1) $\mathbf{2}$ 30.114 III 1 (1) 2 34.164 IV 3 (1) $\mathbf{2}$ 17.15 IV 3 25.65 IV 5 (1) 30.115 III 3 (1) 35.165 I (1)4 $\mathbf{2}$ 26 (1)1 18.16 I 3 (1) $\mathbf{2}$ 26.66 Ι |3(1)|230.116 IV 2 (1) 4 35.166 II 6 (1)1 II |3(1)|18.17 II 2 $\mathbf{2}$ 30.117 IV 3 (2) 35.167 III 2 3 (1)26.67 4 (1)1 18.18 III (2) $\mathbf{2}$ 26.68 III 1 (1) $\mathbf{2}$ 30.118 IV 3 (2) 35.168 III 4 3 4(2)1 18.19 III 1 (1)226.69 III 3 (1) $\mathbf{2}$ 30.119 IV 3 (1) 35.169 IV 6 (1) $\mathbf{2}$ 218.20 IV 24 26.70 III 1 (1) $\mathbf{2}$ 30.120 IV | 2 (2)35.170 IV 5 (1) $\mathbf{2}$ (1)4 18.21 IV 3 26.71 IV 2 (1) 30.121 IV | 2 (1)35.171 IV 5 $\mathbf{2}$ (2)444(1)18.22 IV 2(1)26.72 IV 3 (1) 30.122 IV | 2 (1)36.172 I $\mathbf{2}$ 4 4 4 |2|(1)|2(1)| $\mathbf{2}$ 18.23 IV 3 (1) $\mathbf{2}$ 26.73 IV 3 (1)231.123 I 236.173 II 2 (1)18.24 IV 3 $\mathbf{2}$ 26.74 IV 3 (1) 31.124 II 2 (1) 36.174 III 1 $\mathbf{2}$ (1)22(1)19.25 I 26.75 IV 2 (1)31.125 III 1 (1) 36.175 III 2 $\mathbf{2}$ 1 (1)4 2(1)4 26.76 IV |2| (1) 36.176 III | 1 19.26 II 1 (1)31.126 III 2 (1) 2(1) $\mathbf{2}$ 4 4 26.77 IV 2 (1) 36.177 IV 2 19.27 III 1 (1)4 4 31.127 III 1 (1) $\mathbf{2}$ (1) $\mathbf{2}$ 19.28 IV 27.78 Ι |5(2)| $\mathbf{2}$ 31.128 IV 2 (1) 36.178 IV 2 1 (1)4 4 (1)4 II | 5 (2) $\mathbf{2}$ 19.29 IV 1 (1)27.79 231.129 IV 2 (1) 36.179 IV 2 (1)4 4 19.30 IV 1 27.80 III 1 (1) $\mathbf{2}$ 31.130 IV 2 (1) 37.180 I $\mathbf{2}$ (1)4 4 4 (2)20.31 I 3 (1) $\mathbf{2}$ 27.81 III 5 (2) $\mathbf{2}$ 31.131 IV | 2 (1)37.181 II 4 (2) $\mathbf{2}$ 4 27.82 IV 5 (2) 231.132 IV | 2 (1)37.182 III 1 $\mathbf{2}$ 20.32 II 3 (1) $\mathbf{2}$ $\mathbf{2}$ (1)20.33 III 1 27.83 IV 3 (2) |31.133 IV| 2 (1)37.183 III 4 (2) $\mathbf{2}$ (1)24420.34 III $\mathbf{2}$ 27.84 IV 3 (2) 31.134 IV 2 (1) 37.184 IV 4 $\mathbf{2}$ (1)22(2)4 20.35 IV 3 (1) $\mathbf{2}$ 27.85 IV 3 (2)32.135 I 3 (1) 37.185 IV 3 (2,2) 2^{*} 4 220.36 IV 3 (2)427.86 IV 3 (2) 432.136 II 3 (1) 237.186 IV 3 (2) $\mathbf{2}$ $\mathbf{2}$ $\mathbf{2}$ 32.137 III 1 (1) 38.187 I 6 20.37 IV 3 (1)28.87Ι $|4\ (1)$ $\mathbf{2}$ (1)1 32.138 III 3 (2) 38.188 II 6 21.38 I 8 (1)1 28.88 II |4|(1)| $\mathbf{2}$ 2(1)1 21.39 II 8 (1)1 28.89 III 1 (1) $\mathbf{2}$ 32.139 IV 3 (2) 4 38.189 III 2 (1)1 21.40 III 4(2)28.90 III 2 (1) $\mathbf{2}$ 32.140 IV 2 (1) 38.190 III 3 1 4 (1)1 21.41 III 3 28.91 III 3 (2) $\mathbf{2}$ 32.141 IV 3 (1) 38.191 III 3 (1)1 $\mathbf{2}$ (1)1 21.42 IV 6(2) $\mathbf{2}$ 28.92 IV 4 (1) $\mathbf{2}$ 32.142 IV 2 (1) 38.192 IV 4 (1) $\mathbf{2}$ 4 21.43 IV 7(2) $\mathbf{2}$ 28.93 IV 3 (1) 4 32.143 IV 3 (1) 4 38.193 IV 6 (1) $\mathbf{2}$ $\mathbf{2}$ 38.194 IV 4 $\mathbf{2}$ 21.44 IV 5(2)28.94 IV 4 (1) 33.144 I | 1 (1) (1)4 4 22.45 I 7(1)28.95 IV 3 (1) 33.145 II 1 (1) 39.195 I 4 $\mathbf{2}$ 1 4 4 (1)22.46 II 7(1)1 28.96 IV 4 (1) 4 33.146 III 1 (1) 39.196 II 4 (1) $\mathbf{2}$ 4 22.47 III 28.97 IV 3 (1) 33.147 III 1 (1) 39.197 III 2 (1) $\mathbf{2}$ 3 (1)1 24 22.48 IV 7(2)228.98 IV 3 (1) 4 33.148 III 1 (1) 4 39.198 III 1 (1)27 $\mathbf{2}$ 23.49 I (1)1 29.99 I | 1 (1)433.149 IV 1 (1) 4 39.199 III 3 (2)29.100 II 1 (1) 33.150 IV 1 (1) 39.200 IV 3 (1) 23.50 II 7 (1)1 48 4

TABLE XVII. Characterization of magnetic space groups (MSGs) in the orthorhombic family for spinless electrons

(Continued from the previous page)

39.201 IV	4	(1)	2	47.251	III	9	(1)	1	51.301	IV	7	(1)	4	54.351	IV	4	(2)	4
39.202 IV	3	(1)	2	47.252	III	15	(2, 2, 2)	1	51.302	IV	9	(1)	2	54.352	IV	4	(2)	8
40.203 I	3	(1)	2	47.253	III	13	(1)	1	51.303	IV	8	(1)	4	55.353	Ι	9	(1)	2
40.204 II	3	(1)	2	47.254	IV	18	(1)	2	51.304	IV	7	(1)	4	55.354	Π	9	(1)	2
40.205 III	1	(1)	2	47.255	IV	15	(1)	2	52.305	Ι	5	(2)	4	55.355	ш	3	(1)	2
40.206 III	2	(1)	2	47.256	IV	14	(1)	2	52.306	Π	5	(2)	4	55.356	III	3	(1)	2
40.207 III	2	(1)	2	48.257	Ι	9	(2)	2	52.307	III	2	(1)	4	55.357	III	9	(2, 2, 2)	2
40.208 IV	3	(1)	2	48.258	Π	9	(2)	2	52.308	III	1	(1)	4	55.358	III	5	(2)	2
40.209 IV	3	(1)	4	48.259	III	3	(1)	2	52.309	III	2	(1)	4	55.359	ш	3	(1)	2
40.210 IV	3	(1)	2	48.260	III	5	(2)	2	52.310	III	4	(2)	4	55.360	IV	6	(1)	4
41.211 I	2	(1)	2	48.261	III	7	(2)	2	52.311	III	4	(2)	4	55.361	IV	6	(1)	4
41.212 II	2	(1)	2	48.262	IV	6	(2, 2)	4	52.312	III	3	(2)	4	55.362	IV	5	(1)	4
41.213 III	1	(1)	2	48.263	IV	7	(2, 2)	4	52.313	III	3	(1)	4	55.363	IV	9	(1)	2
41.214 III	1	(1)	2	48.264	IV	8	(2)	2	52.314	IV	5	(2, 2)	4	55.364	IV	6	(1)	4
41.215 III	2	(2)	2	49.265	Ι	14	(2)	2	52.315	IV	4	(2)	4	56.365	Ι	5	(2)	4
41.216 IV	2	(1)	4	49.266	Π	14	(2)	2	52.316	IV	4	(2, 2)	8	56.366	Π	5	(2)	4
41.217 IV	2	(2)	2^*	49.267	III	4	(1)	2	52.317	IV	5	(2)	4	56.367	ш	1	(1)	4
41.218 IV	2	(1)	2	49.268	III	5	(2)	2	52.318	IV	4	(2)	4	56.368	ш	3	(2)	4
42.219 I	5	(1)	1	49.269	III	10	(2, 2)	2	52.319	IV	4	(2, 2)	4	56.369	ш	5	(2, 2)	4
42.220 II	5	(1)	1	49.270	III	7	(2, 2)	2	52.320	IV	5	(2)	4	56.370	III	3	(2)	4
42.221 III	2	(1)	1	49.271	III	9	(2)	2	53.321	Ι	9	(2)	2	56.371	ш	3	(2)	4
42.222 III	3	(2)	1	49.272	IV	9	(2)	4	53.322	Π	9	(2)	2	56.372	IV	3	(2)	8
42.223 IV	5	(1)	2	49.273	IV	14	(2)	2	53.323	Ш	3	(1)	2	56.373	IV	4	(2)	4
43.224 I	2	(1)	2	49.274	IV	9	(2)	4	53.324	Ш	3	(1)	2	56.374	IV	4	(2)	4
43.225 II	2	(1)	2	49.275	IV	8	(2)	4	53.325	Ш	2	(1)	2	56.375	IV	5	(2)	4
43.226 III	1	(1)	2	49.276	IV	8	(2)	4	53.326	III	5	(2)	2	56.376	IV	4	(2)	4
43.227 III	2	(1)	2	50.277	Ι	9	(2)	2	53.327	III	8	(2, 2)	2	57.377	Ι	5	(1)	4
43.228 IV	2	(2)	2^*	50.278	Π	9	(2)	2	53.328	III	6	(2, 2)	2	57.378	Π	5	(1)	4
44.229 I	5	(1)	1	50.279	III	3	(1)	2	53.329	III	4	(1)	2	57.379	ш	3	(1)	4
44.230 II	5	(1)	1	50.280	III	3	(1)	2	53.330	IV	7	(2)	4	57.380	III	2	(1)	4
44.231 III	2	(1)	1	50.281	III	5	(2, 2)	2	53.331	IV	6	(2)	4	57.381	III	1	(1)	4
44.232 III	3	(1)	1	50.282	III	5	(2)	2	53.332	IV	7	(2)	4	57.382	III	4	(2)	4
44.233 IV	5	(1)	2	50.283	III	7	(2)	2	53.333	IV	6	(2)	4	57.383	III	4	(2, 2)	4
44.234 IV	4	(1)	2	50.284	IV	8	(2, 2)	4	53.334	IV	9	(2)	2	57.384	III	3	(2)	4
45.235 I	3	(2)	2	50.285	IV	6	(2, 2)	4	53.335	IV	6	(2)	4	57.385	ш	2	(1)	4
45.236 II	3	(1)	2	50.286	IV	5	(2, 2)	4	53.336	IV	6	(2)	4	57.386	IV	4	(1)	8
45.237 III	1	(1)	2	50.287	IV	9	(2)	2	54.337	Ι	6	(2)	4	57.387	IV	5	(1)	4
45.238 III	3	(2)	2	50.288	IV	6	(2)	4	54.338	Π	6	(2)	4	57.388	IV	5	(1)	4
45.239 IV	3	(2)	2	51.289	Ι	12	(1)	2	54.339	III	1	(1)	4	57.389	IV	5	(1)	4
45.240 IV	2	(2)	2^*	51.290	Π	12	(1)	2	54.340	III	2	(1)	4	57.390	IV	4	(1)	4
46.241 I	3	(1)	2	51.291	Ш	3	(1)	2	54.341	Ш	3	(2)	4	57.391	IV	4	(1)	4
46.242 II	3	(1)	2	51.292	III	4	(1)	2	54.342	III	5	(2, 2)	4	57.392	IV	4	(1)	4
46.243 III	1	(1)	2	51.293	Ш	6	(1)	2	54.343	Ш	3	(2)	4	58.393	Ι	8	(2)	2
46.244 III	2	(1)	2	51.294	III	7	(2, 2)	2	54.344	III	4	(2, 2)	4	58.394	Π	8	(2)	2
46.245 III	2	(1)	2	51.295	III	6	(2)	2	54.345	III	4	(2)	4	58.395	III	2	(1)	2
46.246 IV	3	(1)	2	51.296	III	9	(2, 2, 2)	2	54.346	IV	6	(2)	4	58.396	ш	3	(1)	2
46.247 IV	3	(1)	2	51.297	III	5	(1)	2	54.347	IV	4	(2, 2)	4*	58.397	III	8	(2, 2)	2
46.248 IV	3	(1)	4	51.298	IV	12	(1)	2	54.348	IV	6	(2)	4	58.398	III	5	(2)	2
47.249 I	27	(1)	1	51.299	IV	8	(1)	4	54.349	IV	4	(2)	4	58.399	III	3	(1)	2
47.250 II	27	(1)	1	51.300	IV	9	(1)	4	54.350	IV	6	(2)	4	58.400	IV	5	(2)	4

(Continued from the previous page)

58.401	IV	6	(2)	4	62.442	II	4	(1)	4	65.483	III	6	(1)	1	69.524	III	9	(2, 2)	1
58.402	IV	5	(2)	4	62.443	III	2	(1)	4	65.484	III	6	(1)	1	69.525	III	7	(2)	1
58.403	IV	6	(2)	4	62.444	III	1	(1)	4	65.485	Ш	12	(2, 2, 2)	1	69.526	IV	14	(1)	2
58.404	IV	8	(2)	2	62.445	III	2	(1)	4	65.486	Ш	10	(2, 2)	1	70.527	Ι	6	(2)	2
59.405	Ι	7	(1)	2	62.446	Ш	3	(2)	4	65.487	III	8	(1)	1	70.528	Π	6	(2)	2
59.406	II	7	(1)	2	62.447	III	3	(2)	4	65.488	IV	12	(1)	2	70.529	III	2	(1)	2
59.407	III	2	(1)	2	62.448	III	4	(2)	4	65.489	IV	15	(1)	2	70.530	III	4	(2)	2
59.408	Ш	5	(1)	2	62.449	Ш	1	(1)	4	65.490	IV	11	(1)	2	70.531	ш	4	(1)	2
59.409	III	5	(2, 2)	2	62.450	IV	3	(1)	4	66.491	Ι	11	(2)	2	70.532	IV	5	(2, 2)	2^*
59.410	III	4	(2)	2	62.451	IV	4	(1)	4	66.492	Π	11	(2)	2	71.533	Ι	15	(1)	1
59.411	III	3	(1)	2	62.452	IV	3	(1)	8	66.493	III	3	(1)	2	71.534	Π	15	(1)	1
59.412	IV	6	(1)	4	62.453	IV	3	(1)	4	66.494	III	4	(2)	2	71.535	III	5	(1)	1
59.413	IV	6	(1)	4	62.454	IV	4	(1)	4	66.495	III	9	(2, 2)	2	71.536	III	9	(2, 2)	1
59.414	IV	5	(1)	4	62.455	IV	3	(1)	4	66.496	III	6	(2, 2)	2	71.537	III	7	(1)	1
59.415	IV	7	(1)	2	62.456	IV	4	(1)	4	66.497	Ш	6	(2)	2	71.538	IV	11	(1)	2
59.416	IV	6	(1)	2	63.457	Ι	8	(1)	2	66.498	IV	10	(2)	2	72.539	Ι	9	(1)	2
60.417	Ι	4	(2)	4	63.458	Π	8	(1)	2	66.499	IV	8	(2)	4	72.540	Π	9	(1)	2
60.418	II	4	(2)	4	63.459	Ш	3	(1)	2	66.500	IV	9	(2)	2	72.541	ш	3	(1)	2
60.419	Ш	1	(1)	4	63.460	Ш	4	(1)	2	67.501	Ι	13	(1)	2	72.542	ш	3	(2)	2
60.420	III	2	(1)	4	63.461	III	2	(1)	2	67.502	II	13	(1)	2	72.543	III	7	(2, 2)	2
60.421	III	1	(1)	4	63.462	Ш	5	(2)	2	67.503	III	4	(1)	2	72.544	III	5	(2, 2)	2
60.422	III	3	(2)	4	63.463	Ш	6	(2, 2)	2	67.504	III	5	(1)	2	72.545	III	5	(2)	2
60.423	III	3	(2)	4	63.464	Ш	5	(2, 2)	2	67.505	III	7	(2, 2)	2	72.546	IV	9	(1)	2
60.424	III	4	(2, 2)	4	63.465	Ш	3	(1)	2	67.506	III	8	(2, 2)	2	72.547	IV	7	(1)	4
60.425	III	2	(1)	4	63.466	IV	8	(1)	2	67.507	III	7	(2)	2	73.548	Ι	6	(2)	2^*
60.426	IV	3	(2, 2)	4*	63.467	IV	7	(1)	4	67.508	IV	9	(1)	4	73.549	Π	6	(1)	4
60.427	IV	3	(2)	8	63.468	IV	7	(1)	2	67.509	IV	13	(1)	2	73.550	III	2	(2)	2^*
60.428	IV	4	(2)	4	64.469	Ι	7	(1)	2	67.510	IV	9	(1)	2	73.551	III	4	(2, 2)	4
60.429	IV	3	(2)	4	64.470	Π	7	(1)	2	68.511	Ι	7	(2)	2	73.552	III	4	(2, 2)	4
60.430	IV	3	(2)	4	64.471	Ш	2	(1)	2	68.512	Π	7	(2)	2	73.553	IV	6	(2)	2^*
60.431	IV	4	(2)	4	64.472	Ш	3	(1)	2	68.513	III	2	(1)	2	74.554	Ι	10	(1)	2
60.432	IV	4	(2)	4	64.473	III	2	(1)	2	68.514	III	3	(2)	2	74.555	Π	10	(1)	2
61.433	Ι	3	(2)	4	64.474	Ш	4	(2)	2	68.515	III	5	(2, 2)	2	74.556	III	3	(1)	2
61.434	II	3	(1)	4	64.475	III	6	(2, 2)	2	68.516	III	4	(2, 2)	2	74.557	III	4	(1)	2
61.435	III	1	(1)	4	64.476	Ш	5	(2, 2)	2	68.517	III	5	(2, 2)	2	74.558	III	6	(2, 2)	2
61.436	III	3	(2)	4	64.477	III	3	(2)	2	68.518	IV	7	(2)	4	74.559	III	7	(2, 2)	2
61.437	III	1	(1)	4	64.478	IV	6	(1)	4	68.519	IV	7	(2, 2)	2^{*}	74.560	III	4	(1)	2
61.438	IV	2	(2)	4*	64.479	IV	6	(1)	4	68.520	IV	7	(2)	2	74.561	IV	8	(1)	4
61.439	IV	3	(2)	4	64.480	IV	7	(1)	2	69.521	Ι	15	(1)	1	74.562	IV	9	(1)	2
61.440	IV	2	(2)	4*	65.481	Ι	18	(1)	1	69.522	II	15	(1)	1					
62.441	Ι	4	(1)	4	65.482	Π	18	(1)	1	69.523	III	5	(1)	1					

d: Rank of the band structure group {BS} X_{BS} : Symmetry-based indicators of band topology ν_{BS} : Set of ν bands are symmetry-forbidden from being isolated by band gaps if $\nu \notin \nu_{BS} \mathbb{Z}$ *: Exhibiting exceptional filling pattern; see Table XXI

					~			арз (:									
MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSC	ť	d	X _{BS}	$\nu_{\rm BS}$	MSG	d	X _{BS}	$\nu_{\rm BS}$	MSG		d	$X_{\rm BS}$	$\nu_{\rm BS}$
75.1 1	8	(4)	1	84.51	Ι	13	(2, 4)	2	90.101 IV	7	(1)	2	97.151	I	8	(1)	1
75.2 II	6	(2)	1	84.52	П	11	(2, 2)	2	90.102 IV	6	(2)	2	97.152	П	8	(1)	1
75.3 III	4	(1)	1	84.53	III	9	(2)	2	91.103 I	4	(1)	4	97.153	Ш	5	(1)	1
75.4 IV	6	(2)	2	84.54	III	4	(2)	2	91.104 II	4	(1)	4	97.154	III	5	(2)	1
75.5 IV	5	(2)	2	84.55	III	8	(2, 2, 2)	2	91.105 III	3	(1)	4	97.155	III	5	(1)	1
75.6 IV	5	(2)	2	84.56	IV	10	(2)	2	91.106 III	1	(1)	4	97.156	IV	7	(1)	2
76.7 I	1	(1)	4	84.57	IV	8	(2, 2)	4	91.107 III	2	(1)	4	98.157	Ι	5	(1)	2
76.8 II	1	(1)	4	84.58	IV	9	(2)	2	91.108 IV	4	(1)	4	98.158	Π	5	(1)	2
76.9 III	1	(1)	4	85.59	Ι	11	(2, 4)	2	91.109 IV	3	(2)	4*	98.159	Ш	3	(1)	2
76.10 IV	1	(1)	4	85.60	Π	8	(2)	2	91.110 IV	3	(1)	4	98.160	Ш	2	(1)	2
76.11 IV	1	(2)	4^{*}	85.61	III	5	(1)	2	92.111 I	2	(1)	4	98.161	Ш	4	(1)	2
76.12 IV	1	(1)	4	85.62	III	5	(2)	2	92.112 II	2	(1)	4	98.162	IV	5	(2)	2^*
77.13 I	4	(2)	2	85.63	III	7	(2, 2, 2)	2	92.113 III	1	(1)	4	99.163	Ι	9	(1)	1
77.14 II	4	(2)	2	85.64	IV	7	(2)	4	92.114 III	1	(1)	4	99.164	Π	9	(1)	1
77.15 III	4	(2)	2	85.65	IV	8	(2)	2	92.115 III	2	(1)	4	99.165	III	5	(1)	1
77.16 IV	4	(1)	2	85.66	IV	7	(2)	2	92.116 IV	2	(1)	4	99.166	Ш	6	(1)	1
77.17 IV	3	(2, 2)	2^*	86.67	Ι	9	(2, 2)	2	92.117 IV	2	(2)	4*	99.167	Ш	8	(4)	1
77.18 IV	3	(1)	2	86.68	Π	7	(2)	2	92.118 IV	2	(1)	4	99.168	IV	9	(1)	2
78.19 I	1	(1)	4	86.69	III	5	(2)	2	93.119 I	10	(1)	2	99.169	IV	6	(1)	2
78.20 II	1	(1)	4	86.70	III	3	(2)	2	93.120 II	10	(1)	2	99.170	IV	6	(1)	2
78.21 III	1	(1)	4	86.71	III	7	(2, 2, 2, 2)	2	93.121 III	8	(2)	2	100.171	Ι	6	(1)	2
78.22 IV	1	(1)	4	86.72	IV	6	(1)	4	93.122 III	4	(2)	2	100.172	Π	5	(1)	2
78.23 IV	1	(2)	4^*	86.73	IV	7	(2, 2)	2^*	93.123 III	6	(2)	2	100.173	III	4	(1)	2
78.24 IV	1	(1)	4	86.74	IV	6	(1)	2	93.124 IV	7	(1)	2	100.174	III	3	(1)	2
79.25 I	5	(2)	1	87.75	Ι	16	(4, 4)	1	93.125 IV	7	(2)	2^{*}	100.175	III	5	(4)	2
79.26 II	4	(1)	1	87.76	II	12	(2, 2)	1	93.126 IV	6	(1)	2	100.176	IV	5	(1)	4
79.27 III	3	(1)	1	87.77	III	8	(1)	1	94.127 I	5	(1)	2	100.177	IV	6	(1)	2
79.28 IV	4	(1)	2	87.78	III	4	(2)	1	94.128 II	5	(1)	2	100.178	IV	6	(1)	4
80.29 I	2	(1)	2	87.79	III	7	(2, 2)	1	94.129 III	3	(2)	2	101.179	Ι	5	(1)	2
80.30 II	2	(1)	2	87.80	IV	11	(2, 2)	2	94.130 III	3	(2)	2	101.180	Π	5	(1)	2
80.31 III	2	(1)	2	88.81	Ι	8	(2, 2)	2	94.131 III	5	(2)	2	101.181	Ш	5	(1)	2
80.32 IV	2	(2)	2^*	88.82	II	6	(2)	2	94.132 IV	4	(1)	4	101.182	Ш	4	(2)	2
81.33 I	12	(2, 2, 4)	1	88.83	III	4	(2)	2	94.133 IV	5	(2)	2^{*}	101.183	Ш	4	(2)	2
81.34 II	8	(2, 2)	1	88.84	III	2	(1)	2	94.134 IV	4	(1)	2	101.184	IV	5	(1)	2
81.35 III	4	(1)	1	88.85	III	6	(2, 2)	2	95.135 I	4	(1)	4	101.185	IV	4	(1)	4
81.36 IV	8	(2, 2)	2	88.86	IV	5	(2, 2)	2^*	95.136 II	4	(1)	4	101.186	IV	4	(1)	4
81.37 IV	7	(2, 2, 2)	2	89.87	Ι	12	(1)	1	95.137 III	3	(1)	4	102.187	Ι	4	(1)	2
81.38 IV	7	(2, 2)	2	89.88	II	12	(1)	1	95.138 III	1	(1)	4	102.188	Π	4	(1)	2
82.39 I	11	(2, 2, 2)	1	89.89	III	8	(1)	1	95.139 III	2	(1)	4	102.189	Ш	4	(1)	2
82.40 II	7	(2)	1	89.90	III	8	(4)	1	95.140 IV	4	(1)	4	102.190	Ш	3	(2)	2
82.41 III	3	(1)	1	89.91	III	6	(1)	1	95.141 IV	3	(2)	4*	102.191	Ш	3	(2)	2
82.42 IV	7	(2, 2)	2	89.92	IV	9	(2)	2	95.142 IV	3	(1)	4	102.192	IV	4	(1)	4
83.43 I	24	(4, 4, 4)	1	89.93	IV	8	(1)	2	96.143 I	2	(1)	4	102.193	IV	4	(1)	4
83.44 II	18	(2, 2, 2)	1	89.94	IV	7	(2)	2	96.144 II	2	(1)	4	102.194	IV	4	(1)	2
83.45 III	12	(1)	1	90.95	Ι	7	(1)	2	96.145 III	1	(1)	4	103.195	Ι	6	(2)	2
83.46 III	6	(2)	1	90.96	Π	6	(1)	2	96.146 III	1	(1)	4	103.196	II	6	(2)	2
83.47 III	8	(2, 2)	1	90.97	III	3	(1)	2	96.147 III	2	(1)	4	103.197	III	4	(1)	2
83.48 IV	14	(2, 4)	2	90.98	III	5	(4)	2	96.148 IV	2	(1)	4	103.198	III	4	(1)	2
83.49 IV	15	(2, 2, 2)	2	90.99	III	5	(1)	2	96.149 IV	2	(2)	4^{*}	103.199	III	8	(4)	2
83.50 IV	13	(2, 4)	2	90.100	IV	5	(2)	4	96.150 IV	2	(1)	4	103.200	IV	6	(2)	2

TABLE XVIII. Characterization of magnetic space groups (MSGs) in the tetragonal family for spinless electrons

(Continued from the previous page)

103.201 IV 4 (1) 4	111.251 I	13	(1)	1	117.301	III	5	(1)	2	124.351 I 17	(4)	2
103.202 IV 4 (2) 4	111.252 II	13	(1)	1	117.302	III	3	(1)	2	124.352 II 15	(2)	2
104.203 I 5 (2) 2	111.253 III	5	(1)	1	117.303	III	7	(2, 2, 4)	2	124.353 III 6	(2)	2
104.204 II 4 (1) 2	111.254 III	8	(1)	1	117.304	IV	6	(2)	4	124.354 III 9	(1)	2
104.205 III 3 (1) 2	111.255 III	12	(2, 2, 4)	1	117.305	IV	9	(2)	2	124.355 III 10	(1)	2
104.206 III 3 (1) 2	111.256 IV	9	(1)	2	117.306	IV	6	(2)	4	124.356 III 8	(1)	2
104.207 III 5 (2) 2	111.257 IV	8	(1)	2	118.307	I	9	(2)	2	124.357 III 16 (4	4.4)	2
104.208 IV 4 (2) 4	111.258 IV	8	(1)	2	118.308	П	7	(-) (1)	2	124.358 III 7	(1)	2
104.209 IV 5 (2) 4	112 259 T	12	(2)	2	118 309	ш	5	(1)	2	124 359 III 9	(2)	2
104.210 IV 5 (2) 2	112.259 Т 112.260 П	10	(2) (1)	2	118 310	ш	3	(1)	$\frac{2}{2}$	124.360 IV 17	(2) (4)	2
105,211 I 6 (1) 2	112.200 H	10	(1) (2)	2	118 311	ш	7	(2, 2, 2)	$\frac{2}{2}$	124.360 IV 11	(1)	1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	112.201 III	8	(2) (1)	2	118 312	IV	6	(2, 2, 2) (2)	1	124.362 IV 11	(2) (4)	1
105.212 II 0 (1) 2 105.213 III 4 (2) 2	112.202 III	8	(1)	2	118 313	IV		$\binom{2}{2}$	1	124.302 IV II 125.363 I 13	(1)	т 9
105.215 III + (2) 2 105.214 III - 6 (1) 2	112.205 III	10	(2, 2, 2)	2	110.515	IV IV	0	$\binom{2}{2}$	-	125.303 I 13	(1)	2
105.214 III 0 (1) 2 105.215 III 4 (2) 2	112.204 IV	10	$\binom{2}{2}$	2 0*	110.314	1 V T	0	(2) (1)	1	125.304 II 13	(1)	2
105.215 III 4 (2) 2 105.216 IV (2) (1) 2	112.203 IV	7	(2)	4	119.515	і п	10	(1)	1	125.305 III 0	(1)	2
105.210 IV 0 (1) 2 105.217 IV 4 (1) 4	112.200 IV	((2)	2	119.310	11 111	10	(1)	1	125.300 III 8	(1)	2
105.217 IV 4 (1) 4	113.267 I	8	(1)	2	119.317	III	5	(1)	1	125.367 III 7	(1)	2
105.218 IV 4 (1) 2	113.268 II	6	(1)	2	119.318	III	4	(1)	1	125.368 III 8	(1)	2
106.219 1 3 (2) 2^*	113.269 III	4	(1)	2	119.319	III	11	(2, 2, 2)	1	125.369 III 11 (1	(2, 4)	2
106.220 II 3 (1) 4	113.270 III	3	(1)	2	119.320	IV	8	(1)	2	125.370 III 9	(2)	2
106.221 III 3 (2) 4	113.271 III	7	(2, 2, 4)	2	120.321	I	9	(2)	2	125.371 III 8	(1)	2
106.222 III 3 (2) 2^*	113.272 IV	6	(1)	4	120.322	Π	7	(1)	2	125.372 IV 9	(1)	4
106.223 III 3 (2) 4	113.273 IV	8	(1)	2	120.323	III	5	(1)	2	125.373 IV 13	(1)	2
106.224 IV 3 (1) 4	113.274 IV	6	(1)	2	120.324	III	3	(2)	2	125.374 IV 9	(1)	4
106.225 IV 3 (2) 2^*	114.275 I	7	(2)	2	120.325	III	7	(2, 2, 2)	2	126.375 I 10	(2)	2
106.226 IV 3 (1) 4	114.276 II	5	(1)	2	120.326	IV	9	(2)	2	126.376 II 9	(1)	2
107.227 I 6 (1) 1	114.277 III	3	(2)	2	121.327	Ι	10	(1)	1	126.377 III 5	(2)	2
107.228 II 6 (1) 1	114.278 III	3	(1)	2	121.328	Π	9	(1)	1	126.378 III 5	(1)	2
107.229 III 4 (1) 1	114.279 III	7	(2, 2, 2)	2	121.329	III	4	(1)	1	126.379 III 6	(1)	2
107.230 III 4 (1) 1	114.280 IV	5	(2)	4	121.330	III	5	(1)	1	126.380 III 7	(1)	2
107.231 III 5 (2) 1	114.281 IV	5	(2)	2^*	121.331	III	9	(2, 2, 2)	1	126.381 III 8 (2	2, 2)	2
107.232 IV 6 (1) 2	114.282 IV	7	(2)	2	121.332	IV	8	(1)	2	126.382 III 6	(2)	2
108.233 I 5 (1) 2	115.283 I	12	(1)	1	122.333	Ι	7	(2)	2	126.383 III 7	(1)	2
108.234 II 5 (1) 2	115.284 II	12	(1)	1	122.334	Π	5	(1)	2	126.384 IV 9	(2)	4
108.235 III 4 (1) 2	115.285 III	6	(1)	1	122.335	III	2	(1)	2	126.385 IV 8	(2)	4
108.236 III 3 (1) 2	115.286 III	6	(1)	1	122.336	III	3	(1)	2	126.386 IV 10	(2)	2
108.237 III 5 (4) 2	115.287 III	12	(2, 2, 4)	1	122.337	III	6	(2, 2)	2	127.387 I 18	(1)	2
108.238 IV 5 (1) 2	115.288 IV	9	(1)	2	122.338	IV	5	(2)	2^{*}	127.388 II 15	(1)	2
109.239 I 3 (1) 2	115.289 IV	8	(1)	2	123.339	Ι	27	(1)	1	127.389 III 5	(1)	2
109.240 II 3 (1) 2	115.290 IV	7	(1)	2	123.340	Π	27	(1)	1	127.390 III 12	(1)	2
109.241 III 2 (1) 2	116.291 I	10	(2)	2	123.341	ш	9	(1)	1	127.391 III 9	(1)	2
109.242 III 3 (1) 2	116 292 II	8	(1)	2	123 342	Ш	15	(1)	1	127 392 III 6	(1)	2
109.242 III 0 (1) 2 109.243 III 2 (1) 2	116 293 III	6	(1)	2	123.343	ш	18	(1)	1	127 393 III 15 (4	(1)	2
109.243 III 2 (1) 2 109.244 IV 3 (1) 4	116 294 III	1	(1) (2)	2	123.344	ш	13	(1)	1	127.393 III 10 (1	(1)	2
109.241 IV = 0 (1) 4 $110.245 \text{ I} = 2 (2) 2^{*}$	116 295 III	8	(2, 2, 2)	2	123.345	ш	24	(1)	1	127.395 III 6	(1)	2
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	116 206 IV	0	(2, 2, 2)	2	123.345	ш	124	(4, 4, 4)	1	127.395 III 0	(1)	2 1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	116 207 IV	9 7	(2)	2 1	123.340	ш	12	(1)	1	127.390 IV 11	(1) (1)	ч 9
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	116 200 11	۱ ۶	(2)	-± 1	123.347	ш IV	12	(1)	1 0	127.377 IV 10	(1) (1)	∠ ∧
$110.240 \text{ III} \left[\begin{array}{c} 2 \\ 2 \end{array} \right] \left[\begin{array}{c} 2 \end{array} \right] \left[\begin{array}{c} 2 \\ 2 \end{array} \right] \left[\begin{array}{c} 2 \end{array} \left[\begin{array}{c} 2 \end{array} \right] \left[\begin{array}{c} 2 \end{array} \right] \left[\begin{array}{c} 2 \end{array} \left[\begin{array}{c} 2 \end{array} \right] \left[\begin{array}{c} 2 \end{array} \left[\begin{array}{c} 2 \end{array} \right] \left[\begin{array}{c} 2 \end{array} \left[\end{array} \left] \left[\begin{array}{c} 2 \end{array} \right] \left[\begin{array}{c} 2 \end{array} \right] \left[\end{array} \left[\begin{array}{c} 2 \end{array} \left[\end{array} \right] \left[\end{array} \left] \left[\begin{array}{c} 2 \end{array} \right] \left[\end{array} \left] \left[\end{array} \left[\begin{array}{c} 2 \end{array} \right] \left[\end{array} \left] \left[\end{array} \left]$	117 200 T	0	(2) (2)	4 0	123.340	11	10	(1)	4 0	127.370 IV 12 128 200 I 14	(1) (4)	4 0
$110.249 \text{ III} \begin{bmatrix} 2 & (2) & 4 \\ 110.250 & W \\ 0 & 0 \end{bmatrix} = 0$	117.299 I	9	(2)	2	123.349	11	18	(1)	2	128.399 I 14	(4)	2
110.250 IV 2 (2) 2^*	117.300 II	1	(1)	2	123.350	١V	15	(1)	2	128.400 11 11	(2)	2

(Continued from the previous page)

128 401 III	4	(2)	2	131 ///	IV	15	(1)	2	135 / 87	ш	6	(1)	4	138 530	w	7	(1)	1
128.401 III 128.402 III	8	(2) (1)	$\frac{2}{2}$	131.444		11	(1)	4	135.487	ш	5	(1) (2)	4	130.550	T	18	(1)	4
128.403 III	7	(1)	$\frac{2}{2}$	131 446	IV	12	(1)	2	135 489	ш	8	(2 4)	4	139 532	п	18	(1)	1
128.404 III	5	(1)	2	132,447	T	15	(1)	2	135 490	ш	6	(2, 1)	2*	139 533	ш	6	(1)	1
128.405 III	13	$(4 \ 4)$	2	132.448	П	15	(1)	2	135 491	Ш	4	(1)	4	139 534	Ш	11	(1)	1
128.406 III	6	(1, 1)	$\frac{2}{2}$	132.449	ш	5	(1)	$\frac{2}{2}$	135.492	IV	8	(1)	4	139.535	ш	11	(1)	1
128.407 III	5	(1)	$\frac{-}{2}$	132.450	Ш	11	(1)	$\frac{-}{2}$	135.493	IV	9	(1)	4	139.536	Ш	10	(1)	1
128.408 IV	10	(4)	4	132.451	Ш	10	(2)	$\frac{-}{2}$	135.494	IV	9	(1)	4	139.537	Ш	16	(4, 4)	1
128.409 IV	11	(2)	4	132.452	Ш	9	(1)	2	136.495	I	12	(1)	2	139.538	Ш	9	(1)	1
128.410 IV	14	(4)	2	132.453	ш	12	(2, 4)	2	136.496	п	11	(1)	2	139.539	Ш	8	(1)	1
129.411 I	12	(1)	2	132.454	III	10	(2)	2	136.497	III	4	(1)	2	139.540	IV	15	(1)	2
129.412 II	12	(1)	2	132.455	III	8	(1)	2	136.498	III	10	(1)	2	140.541	I	15	(1)	2
129.413 III	6	(1)	2	132.456	IV	14	(1)	2	136.499	III	7	(2)	2	140.542	II	14	(1)	2
129.414 III	8	(1)	2	132.457	IV	11	(1)	4	136.500	Ш	6	(1)	2	140.543	Ш	5	(1)	2
129.415 III	6	(1)	2	132.458	IV	10	(1)	4	136.501	III	9	(2, 4)	2	140.544	Ш	10	(1)	2
129.416 III	8	(1)	2	133.459	Ι	9	(2)	2^{*}	136.502	III	7	(2)	2	140.545	III	8	(1)	2
129.417 III	11	(2, 4)	2	133.460	Π	8	(1)	4	136.503	III	5	(1)	2	140.546	III	7	(1)	2
129.418 III	8	(1)	2	133.461	III	3	(2)	2^{*}	136.504	IV	9	(1)	4	140.547	III	13	(4, 4)	2
129.419 III	7	(1)	2	133.462	III	5	(2)	4	136.505	IV	11	(1)	4	140.548	III	8	(1)	2
129.420 IV	9	(1)	4	133.463	III	6	(2)	2^{*}	136.506	IV	11	(1)	2	140.549	III	7	(2)	2
129.421 IV	12	(1)	2	133.464	III	7	(2)	4	137.507	Ι	8	(1)	2	140.550	IV	15	(1)	2
129.422 IV	9	(1)	2	133.465	III	6	(2, 2)	4	137.508	Π	8	(1)	2	141.551	Ι	9	(1)	2
130.423 I	8	(2)	4	133.466	III	6	(2, 2)	2^*	137.509	III	4	(1)	2	141.552	II	9	(1)	2
130.424 II	7	(2)	4	133.467	III	6	(1)	4	137.510	III	5	(2)	2	141.553	Ш	3	(1)	2
130.425 III	4	(2)	4	133.468	IV	8	(1)	4	137.511	III	5	(1)	2	141.554	Ш	5	(2)	2
130.426 III	5	(1)	4	133.469	IV	9	(2)	2^{*}	137.512	Ш	7	(2)	2	141.555	III	6	(1)	2
130.427 III	4	(1)	4	133.470	IV	7	(1)	4	137.513	Π	8	(2, 2)	2	141.556	III	7	(2)	2
130.428 III	5	(1)	4	134.471	Ι	12	(1)	2	137.514	III	7	(1)	2	141.557	III	8	(2, 2)	2
130.429 III	8	(2, 4)	4	134.472	Π	12	(1)	2	137.515	III	4	(1)	2	141.558	III	6	(1)	2
130.430 III	6	(1)	4	134.473	III	4	(1)	2	137.516	IV	7	(1)	4	141.559	III	4	(1)	2
130.431 III	6	(2)	4	134.474	III	8	(1)	2	137.517	IV	7	(1)	4	141.560	IV	7	(1)	4
130.432 IV	8	(2)	4	134.475	III	7	(2)	2	137.518	IV	8	(1)	2	142.561	Ι	7	(2)	2^*
130.433 IV	7	(1)	4	134.476	III	8	(1)	2	138.519	Ι	10	(1)	4	142.562	II	6	(1)	4
130.434 IV	8	(2)	4	134.477	III	9	(2, 2)	2	138.520	Π	9	(1)	4	142.563	III	2	(2)	2^*
131.435 I	18	(1)	2	134.478	III	9	(2, 2)	2	138.521	III	4	(1)	4	142.564	III	4	(2)	4
131.436 II	18	(1)	2	134.479	III	7	(1)	2	138.522	III	8	(1)	4	142.565	III	4	(2)	2^*
131.437 III	6	(1)	2	134.480	IV	8	(1)	4	138.523	III	5	(2)	4	142.566	III	5	(2, 2)	4
131.438 III	10	(2)	2	134.481	IV	10	(1)	4	138.524	III	6	(1)	4	142.567	III	5	(2, 2)	4
131.439 III	15	(1)	2	134.482	IV	10	(1)	2	138.525	III	7	(2, 2)	4	142.568	III	5	(2)	2^*
131.440 III	12	(2)	2	135.483	Ι	9	(1)	4	138.526	III	7	(2)	4	142.569	III	4	(2)	4
131.441 III	13	(2, 4)	2	135.484	Π	8	(1)	4	138.527	III	5	(1)	4	142.570	IV	7	(2)	2^*
131.442 III	9	(1)	2	135.485	III	3	(2)	2^{*}	138.528	IV	8	(1)	4					
131.443 III	9	(1)	2	135.486	III	7	(2)	4	138.529	IV	10	(1)	4					

 $\frac{|V| + |V|}{|V|} = \frac{|V| + |V|}{|V|} = \frac{|V| + |V|}{|V|} = \frac{|V| + |V|}{|V|} = \frac{|V|}{|V|} = \frac{|V$

							- 1	, ,				<u> </u>			
MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$	MSG	d	$X_{\rm BS}$	$\nu_{\rm BS}$
143.1 I	7	(3)	1	156.51 III	7	(3)	1	166.101 III	11	(2, 4)	1	177.151 III	5	(1)	1
143.2 II	4	(1)	1	156.52 IV	5	(1)	2	166.102 IV	7	(2)	2	177.152 III	6	(1)	1
143.3 IV	4	(1)	2	157.53 I	5	(1)	1	167.103 I	7	(2)	2	177.153 III	9	(6)	1
144.4 I	1	(1)	3	157.54 II	4	(1)	1	167.104 II	6	(2)	2	177.154 IV	8	(2)	2
144.5 II	1	(1)	3	157.55 III	5	(3)	1	167.105 III	2	(1)	2	178.155 I	3	(1)	6
144.6 IV	1	(1)	6	157.56 IV	4	(1)	2	167.106 III	3	(1)	2	178.156 II	3	(1)	6
145.7 I	1	(1)	3	158.57 I	4	(1)	2	167.107 III	7	(4)	2	178.157 III	2	(1)	6
145.8 II	1	(1)	3	158.58 II	4	(1)	2	167.108 IV	7	(2)	2	178.158 III	2	(1)	6
145.9 IV	1	(1)	6	158.59 III	7	(3)	2	168.109 I	9	(6)	1	178.159 III	1	(1)	6
146.10 I	3	(1)	1	158.60 IV	4	(1)	2	168.110 II	6	(2)	1	178.160 IV	3	(1)	6
146.11 II	2	(1)	1	159.61 I	4	(1)	2	168.111 III	4	(1)	1	179.161 I	3	(1)	6
146.12 IV	2	(1)	2	159.62 II	3	(1)	2	168.112 IV	6	(2)	2	179.162 II	3	(1)	6
147.13 I	13	(2, 12)	1	159.63 III	5	(3)	2	169.113 I	1	(1)	6	179.163 III	2	(1)	6
147.14 II	9	(2, 4)	1	159.64 IV	3	(1)	2	169.114 II	1	(1)	6	179.164 III	2	(1)	6
147.15 III	4	(1)	1	160.65 I	3	(1)	1	169.115 III	1	(1)	6	179.165 III	1	(1)	6
147.16 IV	7	(4)	2	160.66 II	3	(1)	1	169.116 IV	1	(1)	6	179.166 IV	3	(1)	6
148.17 I	11	(2, 4)	1	160.67 III	3	(1)	1	170.117 I	1	(1)	6	180.167 I	7	(1)	3
148.18 II	8	(2, 4)	1	160.68 IV	3	(1)	2	170.118 II	1	(1)	6	180.168 II	7	(1)	3
148.19 III	2	(1)	1	161.69 I	2	(1)	2	170.119 III	1	(1)	6	180.169 III	3	(1)	3
148.20 IV	6	(4)	2	161.70 II	2	(1)	2	170.120 IV	1	(1)	6	180.170 III	3	(1)	3
149.21 I	6	(1)	1	161.71 III	3	(1)	2	171.121 I	3	(2)	3	180.171 III	3	(2)	3
149.22 II	6	(1)	1	161.72 IV	2	(1)	2	171.122 II	3	(2)	3	180.172 IV	5	(2)	6
149.23 III	7	(3)	1	162.73 I	12	(2)	1	171.123 III	1	(1)	3	181.173 I	7	(1)	3
149.24 IV	5	(1)	2	162.74 II	12	(2)	1	171.124 IV	3	(2)	6	181.174 II	7	(1)	3
150.25 I	6	(1)	1	162.75 III	5	(1)	1	172.125 I	3	(2)	3	181.175 III	3	(1)	3
150.26 II	5	(1)	1	162.76 III	5	(1)	1	172.126 II	3	(2)	3	181.176 III	3	(1)	3
150.27 III	5	(3)	1	162.77 III	13	(2, 12)	1	172.127 III	1	(1)	3	181.177 III	3	(2)	3
150.28 IV	4	(1)	2	162.78 IV	8	(2)	2	172.128 IV	3	(2)	6	181.178 IV	5	(2)	6
151.29 I	3	(1)	3	163.79 I	8	(2)	2	173.129 I	5	(3)	2	182.179 I	5	(1)	2
151.30 II	3	(1)	3	163.80 II	7	(2)	2	173.130 II	3	(1)	2	182.180 II	5	(1)	2
151.31 III	1	(1)	3	163.81 III	4	(1)	2	173.131 III	4	(1)	2	182.181 III	4	(1)	2
151.32 IV	2	(1)	6	163.82 III	4	(1)	2	173.132 IV	3	(1)	2	182.182 III	5	(1)	2
152.33 I	3	(1)	3	163.83 III	9	(12)	2	174.133 I	21	(3, 3, 3)	1	182.183 III	5	(3)	2
152.34 II	3	(1)	3	163.84 IV	8	(2)	2	174.134 II	12	(1)	1	182.184 IV	5	(1)	2
152.35 III	1	(1)	3	164.85 I	12	(2)	1	174.135 III	4	(1)	1	183.185 I	8	(1)	1
152.36 IV	2	(1)	6	164.86 II	12	(2)	1	174.136 IV	11	(3)	2	183.186 II	8	(1)	1
153.37 I	3	(1)	3	164.87 III	4	(1)	1	175.137 I	27	(6, 6, 6)	1	183.187 III	5	(1)	1
153.38 II	3	(1)	3	164.88 III	6	(1)	1	175.138 II	18	(2, 2, 2)	1	183.188 III	4	(1)	1
153.39 III	1	(1)	3	164.89 III	13	(2, 12)	1	175.139 III	12	(1)	1	183.189 III	9	(6)	1
153.40 IV	2	(1)	6	164.90 IV	8	(2)	2	175.140 III	6	(2)	1	183.190 IV	8	(1)	2
154.41 I	3	(1)	3	165.91 I	8	(2)	2	175.141 III	9	(2, 4)	1	184.191 I	6	(2)	2
154.42 II	3	(1)	3	165.92 II	7	(2)	2	175.142 IV	15	(2, 6)	2	184.192 II	6	(2)	2
154.43 III	1	(1)	3	165.93 III	3	(1)	2	176.143 I	16	(3, 6)	2	184.193 III	4	(1)	2
154.44 IV	2	(1)	6	165.94 III	5	(1)	2	176.144 II	10	(2)	2	184.194 III	3	(1)	2
155.45 I	4	(1)	1	165.95 III	9	(12)	2	176.145 III	11	(3)	2	184.195 III	9	(6)	2
155.46 II	4	(1)	1	165.96 IV	8	(2)	2	176.146 III	3	(1)	2	184.196 IV	6	(2)	2
155.47 III	3	(1)	1	166.97 I	11	(2)	1	176.147 III	7	(4)	2	185.197 I	4	(1)	2
155.48 IV	3	(1)	$\frac{1}{2}$	166.98 П	11	(2)	1	176.148 IV	$ _{10}$	(2)	$\frac{-}{2}$	185.198 П	4	(1)	2
156.49 I	5	(1)	1	166.99 III	3	(1)	1	177.149 T	10	(1)	1	185.199 III	5	(1)	2
156.50 II	5	(1)	1	166.100 III	4	(1)	1	177,150 II	10	(1)	1	185,200 III	3	(1)	2
II	5	(*)	-		L.*	(*/	-	II	1-0	(*)	-		<u> </u>	(*)	

TABLE XIX. Characterization of magnetic space groups (MSGs) in the hexagonal family for spinless electrons

(Continued from the previous page)

185.201 III	5	(3)	2	188.219	III	14	(3, 3)	2	191.237	III	12	(1)	1	193.255	III	4	(1)	$\overline{2}$
185.202 IV	4	(1)	2	188.220	IV	12	(3)	2	191.238	III	12	(2)	1	193.256	ш	10	(1)	2
186.203 I	4	(1)	2	189.221	Ι	15	(1)	1	191.239	Ш	12	(2)	1	193.257	ш	9	(3)	2
186.204 II	4	(1)	2	189.222	II	12	(1)	1	191.240	III	27	(6, 6, 6)	1	193.258	ш	8	(2)	2
186.205 III	4	(1)	2	189.223	III	4	(1)	1	191.241	III	10	(1)	1	193.259	ш	8	(2)	2
186.206 III	4	(1)	2	189.224	III	5	(1)	1	191.242	IV	16	(1)	2	193.260	III	14	(3, 6)	2
186.207 III	5	(3)	2	189.225	III	15	(3,3,3)	1	192.243	Ι	17	(6)	2	193.261	III	5	(1)	2
186.208 IV	4	(1)	2	189.226	IV	9	(1)	2	192.244	Π	14	(2)	2	193.262	IV	13	(1)	2
187.209 I	15	(1)	1	190.227	Ι	12	(3)	2	192.245	III	6	(2)	2	194.263	Ι	13	(1)	2
187.210 II	15	(1)	1	190.228	Π	8	(1)	2	192.246	III	9	(1)	2	194.264	II	13	(1)	2
187.211 III	6	(1)	1	190.229	III	3	(1)	2	192.247	III	8	(1)	2	194.265	III	4	(1)	2
187.212 III	5	(1)	1	190.230	III	4	(1)	2	192.248	III	7	(2)	2	194.266	III	12	(3)	2
187.213 III	21	(3,3,3)	1	190.231	III	12	(3,3)	2	192.249	III	7	(2)	2	194.267	III	9	(1)	2
187.214 IV	10	(1)	2	190.232	IV	9	(3)	2	192.250	III	18	(6, 6)	2	194.268	ш	8	(2)	2
188.215 I	12	(3)	2	191.233	Ι	24	(1)	1	192.251	III	8	(2)	2	194.269	ш	8	(2)	2
188.216 II	9	(1)	2	191.234	II	24	(1)	1	192.252	IV	17	(6)	2	194.270	ш	16	(3, 6)	2
188.217 III	5	(1)	2	191.235	III	8	(1)	1	193.253	Ι	13	(1)	2	194.271	ш	5	(1)	2
188.218 III	4	(1)	2	191.236	III	15	(1)	1	193.254	II	12	(1)	2	194.272	IV	12	(1)	2

MSG	Τ	d	XDC	$\nu_{\rm PC}$	MSG	d	XDC	Vpc	MSG	T	d XDC	Vpc	MSG	d	XDC	Vpg
195.1 I		7	(1)	1	206.39 III	3	(1)	4	216.77 IV	7	7 (1)	2	224.115 IV	10	(1)	2
195.2 II	[6	(1)	1	207.40 I	9	(1)	1	217.78 I		9 (1)	1	225.116 I	17	(1)	1
195.3 IV	7	4	(2)	2	207.41 II	9	(1)	1	217.79 II		8 (1)	1	225.117 II	17	(1)	1
196.4 I		5	(1)	1	207.42 III	6	(1)	1	217.80 II	I	5(1)	1	225.118 III	8	(1)	1
196.5 II	[4	(1)	1	207.43 IV	6	(2)	2	218.81 I		9 (2)	2	225.119 III	13	(1)	1
196.6 IV	7	4	(2)	2	208.44 I	7	(1)	2	218.82 II		7 (1)	2	225.120 III	7	(1)	1
197.7 I		5	(1)	1	208.45 II	7	(1)	2	218.83 II	I	6 (1)	2	225.121 IV	13	(1)	2
197.8 II	[4	(1)	1	208.46 III	6	(2)	2	218.84 IV	7	6(2)	2	226.122 I	14	(1)	2
198.9 I		3	(1)	4	208.47 IV	5	(1)	2	219.85 I		8 (2)	2	226.123 II	12	(1)	2
198.10 II	[2	(1)	4	209.48 I	7	(1)	1	219.86 II		6 (1)	2	226.124 III	7	(1)	2
198.11 IV	7	2	(1)	4*	209.49 II	7	(1)	1	219.87 II	I	5(1)	2	226.125 III	10	(1)	2
199.12 I		4	(1)	2^*	209.50 III	5	(1)	1	219.88 IV	7	8 (2)	2	226.126 III	6	(2)	2
199.13 II	[3	(1)	2^*	209.51 IV	6	(1)	2	220.89 I		7(2)	2^*	226.127 IV	14	(1)	2
200.14 I		17	(1)	1	210.52 I	4	(1)	2	220.90 II		5(1)	2^*	227.128 I	11	(1)	2
200.15 II	[14	(1)	1	210.53 II	4	(1)	2	220.91 II	I	4 (1)	2^*	227.129 II	11	(1)	2
200.16 II	I	6	(1)	1	210.54 III	4	(1)	2	221.92 I	2	22(1)	1	227.130 III	6	(1)	2
200.17 IV	7	9	(1)	2	210.55 IV	4	(2)	2^{*}	221.93 II	2	22(1)	1	227.131 III	10	(2)	2
201.18 I		11	(2)	2	211.56 I	7	(1)	1	221.94 II	I 1	.0 (1)	1	227.132 III	4	(1)	2
201.19 II	[8	(2)	2	211.57 II	7	(1)	1	221.95 II	I 1	.6(1)	1	227.133 IV	8	(1)	4
201.20 II	I	4	(2)	2	211.58 III	5	(1)	1	221.96 II	I	9(1)	1	228.134 I	9	(2)	2^*
201.21 IV	7	7	(2)	2	212.59 I	3	(1)	4	221.97 IV	/ 1	(1)	2	228.135 II	7	(1)	4
202.22 I		13	(1)	1	212.60 II	3	(1)	4	222.98 I	1	(1)	2	228.136 III	5	(2)	2^*
202.23 II	[10	(1)	1	212.61 III	3	(1)	4	222.99 II		9(1)	2	228.137 III	7	(2)	4
202.24 II	I	4	(2)	1	212.62 IV	3	(1)	4*	222.100 II	I	6(1)	2	228.138 III	4	(2)	4
202.25 IV	7	9	(1)	2	213.63 I	3	(1)	4	222.101 II	I	8 (1)	2	228.139 IV	9	(2)	2^*
203.26 I		10	(2)	2	213.64 II	3	(1)	4	222.102 II	I	6(1)	2	229.140 I	17	(1)	1
203.27 II	[7	(2)	2	213.65 III	3	(1)	4	222.103 IV	/ 1	(1)	2	229.141 II	17	(1)	1
203.28 II	I	3	(1)	2	213.66 IV	3	(1)	4*	223.104 I	1	(1)	2	229.142 III	9	(1)	1
203.29 IV	/	6	(2, 2)	2^*	214.67 I	5	(1)	2^{*}	223.105 II	1	(1)	2	229.143 III	13	(1)	1
204.30 I		13	(1)	1	214.68 II	5	(1)	2^{*}	223.106 II	I	9(2)	2	229.144 III	7	(1)	1
204.31 II	[10	(1)	1	214.69 III	4	(1)	2^{*}	223.107 II	I 1	(1)	2	230.145 I	9	(2)	2^*
204.32 II	I	4	(1)	1	215.70 I	10	(1)	1	223.108 II	I	6(1)	2	230.146 II	7	(1)	4^*
205.33 I		9	(2)	4	215.71 II	10	(1)	1	223.109 IV	/ 1	(1)	2	230.147 III	5	(2)	4^*
205.34 II	[6	(1)	4	215.72 III	6	(1)	1	224.110 I	1	(1)	2	230.148 III	7	(2)	2^*
205.35 II	I	2	(1)	4	215.73 IV	7	(1)	2	224.111 II	1	(1)	2	230.149 III	4	(1)	4^*
205.36 IV	/	5	(2)	4^*	216.74 I	9	(1)	1	224.112 II	I	7(1)	2				
206.37 I		10	(2)	2^*	216.75 II	9	(1)	1	224.113 II	I 1	(1)	2				
206.38 II	[]]	$\overline{7}$	(1)	4	216.76 III	5	(1)	1	224.114 II	I	6 (1)	2				

TABLE XX. Characterization of magnetic space groups (MSGs) in the cubic family for spinless electrons

d: Rank of the band structure group {BS} X_{BS} : Symmetry-based indicators of band topology ν_{BS} : Set of ν bands are symmetry-forbidden from being isolated by band gaps if $\nu \notin \nu_{BS} \mathbb{Z}$ *: Exhibiting exceptional filling pattern; see Table XXI

MSC	ť	$\{\nu\}_{AI}$	$\{\nu\}_{BS}$	MSG	$\{\nu\}_{AI}$	$\{\nu\}_{BS}$	MSG	$\{\nu\}_{AI}$	$\{\nu\}_{BS}$	MSG	$\{\nu\}_{AI}$	$\{\nu\}_{BS}$
9.41	IV	$4\mathbb{N}$	$2\mathbb{N}$	80.32 IV	$4\mathbb{N}$	$2\mathbb{N}$	133.459 I	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$	212.62 IV	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$
15.91	IV	$4\mathbb{N}$	$2\mathbb{N}$	86.73 IV	$4\mathbb{N}$	$2\mathbb{N}$	133.461 II	$4\mathbb{N}$	$2\mathbb{N}$	213.66 IV	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$
29.105	IV	$8\mathbb{N}$	$4\mathbb{N}$	88.86 IV	$4\mathbb{N}$	$2\mathbb{N}$	133.463 II	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$	214.67 I	$2\mathbb{N}\setminus\{2\}$	—
37.185	IV	$4\mathbb{N}$	$2\mathbb{N}$	91.109 IV	$8\mathbb{N}$	$4\mathbb{N}$	133.466 II	$4\mathbb{N}$	$2\mathbb{N}$	214.68 II	$2\mathbb{N}\setminus\{2\}$	-
41.217	IV	$4\mathbb{N}$	$2\mathbb{N}$	92.117 IV	$8\mathbb{N}$	$4\mathbb{N}$	133.469 IV	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$	214.69 III	$2\mathbb{N}\setminus\{2\}$	—
43.228	IV	$4\mathbb{N}$	$2\mathbb{N}$	93.125 IV	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$	135.485 II	$4\mathbb{N}$	$2\mathbb{N}$	220.89 I	$2\mathbb{N}\setminus\{2,4,10\}$	$2\mathbb{N}\setminus\{2\}$
45.240	IV	$4\mathbb{N}$	$2\mathbb{N}$	94.133 IV	$4\mathbb{N}$	$2\mathbb{N}$	135.490 II	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$	220.90 II	$2\mathbb{N}\setminus\{2,4,10\}$	—
54.347	IV	$8\mathbb{N}$	$4\mathbb{N}$	95.141 IV	$8\mathbb{N}$	$4\mathbb{N}$	142.561 I	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$	220.91 III	$2\mathbb{N}\setminus\{2,4,10\}$	—
60.426	IV	$8\mathbb{N}$	$4\mathbb{N}$	96.149 IV	$8\mathbb{N}$	$4\mathbb{N}$	142.563 II	$4\mathbb{N}$	$2\mathbb{N}$	228.134 I	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$
61.438	IV	$8\mathbb{N}$	$4\mathbb{N}$	98.162 IV	$4\mathbb{N}$	$2\mathbb{N}$	142.565 II	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$	228.136 III	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$
61.440	IV	$8\mathbb{N}$	$4\mathbb{N}$	106.219 I	$4\mathbb{N}$	$2\mathbb{N}$	142.568 II	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$	228.139 IV	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$
68.519	IV	$4\mathbb{N}$	$2\mathbb{N}$	106.222 III	$4\mathbb{N}$	$2\mathbb{N}$	142.570 IV	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$	230.145 I	$4\mathbb{N}\setminus\{4\}$	$2\mathbb{N}\setminus\{2,4\}$
70.532	IV	$4\mathbb{N}$	$2\mathbb{N}$	106.225 IV	$4\mathbb{N}$	$2\mathbb{N}$	198.11 IV	$ 4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$	230.146 II	$4\mathbb{N}\setminus\{4\}$	—
73.548	Ι	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$	110.245 I	$4\mathbb{N}$	$2\mathbb{N}$	199.12 I	$2\mathbb{N}\setminus\{2\}$	-	230.147 III	$4\mathbb{N}\setminus\{4\}$	$4\mathbb{N}$
73.550	Ш	$4\mathbb{N}$	$2\mathbb{N}$	110.248 III	$4\mathbb{N}$	$2\mathbb{N}$	199.13 II	$2\mathbb{N}\setminus\{2\}$	-	230.148 III	$4\mathbb{N}\setminus\{4\}$	$2\mathbb{N}\setminus\{2,4,10\}$
73.553	IV	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$	110.250 IV	$4\mathbb{N}$	$2\mathbb{N}$	203.29 IV	$4\mathbb{N}$	$2\mathbb{N}$	230.149 III	$4\mathbb{N}\setminus\{4\}$	—
76.11	IV	$8\mathbb{N}$	$4\mathbb{N}$	112.265 IV	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$	205.36 IV	81	$4\mathbb{N}$			
77.17	IV	$4\mathbb{N}$	$2\mathbb{N}$	114.281 IV	$4\mathbb{N}$	$2\mathbb{N}$	206.37 I	$4\mathbb{N}$	$2\mathbb{N}\setminus\{2\}$			
78.23	IV	$8\mathbb{N}$	$4\mathbb{N}$	122.338 IV	$4\mathbb{N}$	$2\mathbb{N}$	210.55 IV	$4\mathbb{N}$	$2\mathbb{N}$			

TABLE XXI. Magnetic space groups (MSGs) for which spinless electrons exhibit exceptional filling patterns

 \mathbb{N} : The set of natural numbers

 $\{\nu\}_{AI}$: Set of fillings for which physical atomic insulators are possible $\{\nu\}_{BS}$: Set of fillings for which physical band structure are possible; a dash (-) indicates $\{\nu\}_{BS} = \{\nu\}_{AI}$ for that MSG