

Abundance ratios & ages of stellar populations in HARPS-GTO sample

E. Delgado Mena¹, M. Tsantaki², V. Zh. Adibekyan¹, S. G. Sousa^(1,3),
N. C. Santos^(1,3), J. I. González Hernández^(4,5) and G. Israelian^(4,5)

¹Instituto de Astrofísica e Ciências do Espaço,
Universidade do Porto, CAUP, Rua das Estrelas, PT4150-762 Porto, Portugal.
email: elisa.delgado@astro.up.pt

²Instituto de Radioastronomía y Astrofísica, IRyA, UNAM,
Campus Morelia, A.P. 3-72, C.P. 58089, Michoacán, Mexico

³Departamento de Física e Astronomia, Faculdade de Ciências, U. Porto, Portugal

⁴Instituto de Astrofísica de Canarias, C/ Via Lactea, s/n, 38205, La Laguna, Tenerife, Spain

⁵Departamento de Astrofísica, Universidad de La Laguna, 38206 La Laguna, Tenerife, Spain

Abstract.

In this work we present chemical abundances of heavy elements ($Z > 28$) for a homogeneous sample of 1059 stars from HARPS planet search program. We also derive ages using parallaxes from Hipparcos and Gaia DR1 to compare the results. We study the $[X/Fe]$ ratios for different populations and compare them with models of Galactic chemical evolution. We find that thick disk stars are chemically disjunct for Zn and Eu. Moreover, the high-alpha metal-rich population presents an interesting behaviour, with clear overabundances of Cu and Zn and lower abundances of Y and Ba with respect to thin disk stars. Several abundance ratios present a significant correlation with age for chemically separated thin disk stars (regardless of their metallicity) but thick disk stars do not present that behaviour. Moreover, at supersolar metallicities the trends with age tend to be weaker for several elements.

Keywords. stars: abundances - stars: fundamental parameters - Galaxy: evolution - Galaxy: disk - solar neighborhood

1. Introduction

In the era of large spectroscopic surveys such as APOGEE, Gaia-ESO Survey or RAVE, among others, the contribution of smaller samples with high-resolution and high quality spectra is of great importance to understand the Galactic Chemical Evolution (GCE). In this work we have derived abundances for Cu, Zn, Sr, Y, Zr, Ba, Ce, Nd and Eu (Delgado Mena et al. 2017) for 1111 stars within the volume-limited HARPS-GTO planet search sample in order to complement our previous works for light elements (Delgado Mena et al. 2014, Delgado Mena et al. 2015, Suárez Andrés et al. 2016, Bertrán de Lis et al. 2015), α -elements and Fe-peak elements (Adibekyan et al. 2012). The main purpose of this work is to evaluate the GCE evolution of those heavier elements and the dependence on stellar ages of different abundance ratios.

2. Stellar ages

We derive the masses, radii and ages with the PARAM v1.3 tool[†] using the PARSEC isochrones (Bressan et al. 2012) with our values for T_{eff} and $[Fe/H]$, the V magnitudes

[†] <http://stev.oapd.inaf.it/cgi-bin/param>




Figure 1. General $[X/Fe]$ ratios as a function of age for the reduced sample with reliable ages. Thin disk stars, thick disk stars and $h\alpha mr$ are depicted with dots, triangles and squares, respectively. The linear fit to all the stars just serves as eye guiding.

from the main Hipparcos catalogue (Perryman et al. 1997)) and the parallaxes from the Hipparcos new reduction (van Leeuwen 2007) or from the first release (DR1) of Gaia (Lindegren et al. 2016). We note that we added a systematic error of 0.3 mas to the formal error of the Gaia DR1 parallaxes as recommended by the Gaia collaboration. Meanwhile Hipparcos provides parallaxes for 1051 out of the 1059 stars within our sample, only 923 stars have parallaxes in GAIA DR1. Moreover, there are significant differences in many cases, leading to non-negligible differences in age. In order to have a sample with ages as reliable as possible we decided to select the Hipparcos ages with a difference less than 1 Gyr with respect to the ages derived with GAIA parallaxes and with an error in age lower than 2 Gyr. This final sample is composed by 377 stars belonging to the thin disk, thick disk and high- α metal-rich stars (hereafter $h\alpha mr$, a population with high α abundances at $[Fe/H] > -0.2$ dex discovered by Adibekyan et al. 2011).

3. Abundance ratios vs age

In Fig. 1 we can see how several elements depend on age. By combining elements that increase and decrease with age, respectively, it is possible to have steeper and more constrained trends. For example, $[Mg/Fe]$ shows a tight increasing trend with age, meanwhile $[Eu/Fe]$, $[Zn/Fe]$ and $[Al/Fe]$ also show this dependency though with more dispersion. This trend is expected since these elements are mainly formed by massive stars which started to contribute to the Galaxy chemical enrichment earlier than the lower mass stars responsible for Fe production. On the other hand, the light- s process elements Y and Sr




Figure 2. $[\text{Y}/\text{Mg}]$, $[\text{Y}/\text{Zn}]$, $[\text{Y}/\text{Al}]$ and $[\text{Sr}/\text{Mg}]$, $[\text{Sr}/\text{Zn}]$, $[\text{Sr}/\text{Al}]$ for $-0.8 < [\text{Fe}/\text{H}] < -0.2$. Symbols as in Fig 1.

show the most clear decreasing trends with age. These elements are formed by low-mass AGB stars so we can expect them to increase with time (for younger stars) due to the increasing and delayed contribution of low-mass stars as the Galaxy evolves. In Figs. 2 and 3 we show different combinations of previously mentioned elements at different metallicity regions. Previous works have explored and confirmed the tight correlation of these abundance ratios with age (e.g. da Silva et al. 2012), Nissen 2015, Spina et al. 2016) but only using solar twins or solar analogues. However, Feltzing et al. (2017) noted that $[\text{Y}/\text{Mg}]$ clock is not valid at $[\text{Fe}/\text{H}] < -0.5$ dex. In our sample we find that still at $[\text{Fe}/\text{H}] < -0.5$ dex the different abundance ratios show a correlation with age (steeper for $[\text{Sr}/\text{X}]$ than for $[\text{Y}/\text{X}]$) but this is only valid for thin disk stars. We note however that our sample of thick disk stars with reliable ages is quite small. It is also clear that the trends become flat at ages $\gtrsim 8$ Gyr. On the other hand, at higher metallicities, in the bin $-0.2 < [\text{Fe}/\text{H}] < 0.2$ dex, the abundance ratios of Y and Sr (with respect to Mg, Zn and Al) present similar slopes. Nevertheless, we remark that meanwhile $[\text{Sr}/\text{Fe}]$ presents a constant correlation with age at different metallicities, $[\text{Y}/\text{Fe}]$ becomes flatter as $[\text{Fe}/\text{H}]$ increases. Moreover, we can observe that thin disk stars present mostly no dependence on age for ages $\gtrsim 8$ Gyr but *halms* stars show a continuous dependence in the full age range for $[\text{Y}/\text{X}]$ ratios. The improvement of parallaxes from GAIA DR2 will help to determine more precise ages for our stars increasing the sample size and allowing us to better understand the behaviour of the abundance-age trends for different populations in the Galaxy.

Acknowledgements

E.D.M., V.Zh.A., N.C.S. and S.G.S. acknowledge the support from Fundação para a Ciência e a Tecnologia (FCT) through national funds and from FEDER through COMPETE2020 by the following grants




Figure 3. [Y/Mg],[Y/Zn],[Y/Al] and [Sr/Mg],[Sr/Zn],[Sr/Al] for $-0.2 < [Fe/H] < 0.2$. Symbols as in Fig 1.

UID/FIS/04434/2013 & POCI-01-0145-FEDER-007672, PTDC/FIS-AST/7073/2014 & POCI-01-0145-FEDER-016880 and PTDC/FIS-AST/1526/2014 & POCI-01-0145-FEDER-016886. E.D.M., V.Zh.A., N.C.S. and S.G.S. also acknowledge the support from FCT through Investigador FCT contracts IF/00849/2015, IF/00650/2015, IF/00169/2012/CP0150/CT0002 and IF/00028/2014/CP1215/CT0002 funded by FCT (Portugal) and POPH/FSE (EC). This research has made use of the SIMBAD database operated at CDS, Strasbourg (France).

This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement.

References

- Adibekyan, V. Z., Santos, N. C., Sousa, S. G., & Israelian, G. 2011, *A&A*, 535, L11
 Adibekyan, V. Z., Sousa, S. G., Santos, N. C., et al. 2012, *A&A*, 545, A32
 Bertran de Lis, S., Delgado Mena, E., Adibekyan, V. Z., et al. 2015, *A&A*, 576, A89
 Bressan, A., Marigo, P., Girardi, L., et al. 2012, *MNRAS*, 427, 127
 da Silva, R., Porto de Mello, G. F., Milone, A. C., et al. 2012, *A&A*, 542, A84
 Delgado Mena, E., Bertrán de Lis, S., Adibekyan, V. Z., et al. 2015, *A&A*, 576, A69
 Delgado Mena, E., Israelian, G., González Hernández, J. I., et al. 2014, *A&A*, 562, A92
 Delgado Mena, E., Tsantaki, M., Adibekyan, V. Z., et al. 2017, *arXiv:1705.04349*
 Feltzing, S., Howes, L. M., McMillan, P. J., & Stenokutė, E. 2017, *MNRAS*, 465, L109
 Lindegren, L., Lammers, U., Bastian, U., et al. 2016, *A&A*, 595, A4
 Nissen, P. E. 2015, *A&A*, 579, A52
 Perryman, M. A. C., Lindegren, L., Kovalevsky, J., et al. 1997, *A&A*, 323, L49
 Spina, L., Meléndez, J., Karakas, A. I., et al. 2016, *A&A*, 593, A125
 Suárez-Andrés, L., Israelian, G., González Hernández, J. I., et al. 2016, *A&A*, 591, A69
 van Leeuwen, F. 2007, *A&A*, 474, 653