Top Quark Rare Decays via Loop-Induced FCNC Interactions in Extended Mirror Fermion Model

P. Q. Hung[∗](#page-1-0)

Department of Physics, University of Virginia, Charlottesville, VA 22904-4714, USA

and

Center for Theoretical and Computational Physics, Hue University College of Education, Hue, Vietnam

Yu-Xiang Lin[†](#page-1-1) and Chrisna Setyo Nugroho[‡](#page-1-2)

Department of Physics, National Taiwan Normal University, Taipei 116, Taiwan

Tzu-Chiang Yuan[§](#page-1-3)

Institute of Physics, Academia Sinica, Nangang, Taipei 11529, Taiwan

and

Physics Division,

National Center for Theoretical Sciences, Hsinchu, Taiwan

(Dated: November 1, 2021)

Abstract

Flavor changing neutral current (FCNC) interactions for a top quark t decays into Xq with X represents a neutral gauge or Higgs boson, and q a up- or charm-quark are highly suppressed in the Standard Model (SM) due to the Glashow-Iliopoulos-Miami mechanism. Whilst current limits on the branching ratios of these processes have been established at the order of 10⁻⁴ from the Large Hadron Collider experiments, SM predictions are at least nine orders of magnitude below. In this work, we study some of these FCNC processes in the context of an extended mirror fermion model, originally proposed to implement the electroweak scale seesaw mechanism for non-sterile right-handed neutrinos. We show that one can probe the process $t \to Zc$ for a wide range of parameter space with branching ratios varying from 10^{-6} to 10^{-8} , comparable with various new physics models including the general two Higgs doublet model with or without flavor violations at tree level, minimal supersymmetric standard model with or without R-parity, and extra dimension model.

[∗] pqh@virginia.edu

[†] x90417@gmail.com

[‡] setyo13nugros@gmail.com

[§] tcyuan@phys.sinica.edu.tw

I. INTRODUCTION

Absence of flavor changing neutral current (FCNC) interactions in the Standard Model (SM) at tree level is quite a unique property due to the special quantum numbers of the three generations of fermions (quarks and leptons) and one Higgs doublet under the gauge group of $SU(3)_C \times SU(2)_L \times U(1)_Y$. FCNC interactions can nevertheless be induced at the quantum loop level and therefore are suppressed by the GIM mechanism [\[1\]](#page-26-0). Experimental results for various FCNC processes in the kaon, D and B meson systems are all in line with the SM expectations.

For the heavy top quark t , the story is quite different. Since there is no time for the heavy top quark to form bound states, we can discuss its free decay, like the dominant decay mode $t \to W^+b$ at tree level or its rare FCNC decays. The SM branching ratios $\mathcal{B}(t \to Xq)$ where X denotes one of the following neutral particle Z, γ, g or h in SM, and q denotes the light u or c quark, vary in the range $10^{-17} - 10^{-12}$ [\[2\]](#page-26-1), which are unobservable at the present technology. However, in many models beyond the SM, branching ratios for some of these processes of order up to 10^{-3} can be achieved. Observations of these rare top quark FCNC decays at the Large Hadron Collider (LHC) with significant larger branching ratios than the SM predictions would then be clear signals, albeit indirect, of new physics.

Indeed LHC can be considered as a top quark factory, estimated to produce $10^8 t\bar{t}$ pair with an integrated luminosity of 100 inverse femtobarn. For an updated review on top quark properties at the LHC, see [\[3\]](#page-26-2). The current limits for $t \to Zq$ [\[4–](#page-26-3)[7\]](#page-26-4) are

$$
\mathcal{B}(t \to Zu) \leq \begin{cases} 2.2 \times 10^{-4} \text{ [CMS]} ,\\ 1.7 \times 10^{-4} \text{ [ATLAS]} , \end{cases}
$$
 (1)

$$
\mathcal{B}(t \to Zc) \le \begin{cases} 4.9 \times 10^{-4} \text{ [CMS]} ,\\ 2.3 \times 10^{-4} \text{ [ATLAS]} ; \end{cases}
$$
 (2)

and for $t \to \gamma q$, we have the limits from CMS [\[8\]](#page-26-5)

$$
\mathcal{B}(t \to \gamma u) \le 1.3 \times 10^{-4},
$$

$$
\mathcal{B}(t \to \gamma c) \le 1.7 \times 10^{-3}.
$$
 (3)

Projected limits for the above as well as other FCNC processes for the top quark are expected to be improved constantly in the future at the LHC. Thus searching for or discovery of any one of these FCNC rare top decays $t \rightarrow X q$ at the LHC would be providing interesting constraints or discriminations of various new physics models in the future.

Over the years FCNC top quark decays had been studied intensively in the literature for many new physics models, like the minimal supersymmetric standard model (MSSM) with $[9]$ or without $[10, 11]$ $[10, 11]$ R-parity, flavor conserving $[12]$ or flavor violating [\[13](#page-26-10)[–16\]](#page-27-0) two Higgs doublet model (2HDM), aligned two Higgs doublet model (A2HDM) [\[17\]](#page-27-1), warped extra-dimensions [\[18,](#page-27-2) [19\]](#page-27-3), and effective Lagrangian frame-work [\[20\]](#page-27-4), *etc.* Branching ratios for FCNC top quark decays in all these models are typically many orders of magnitude above the SM and some of them may lead to detectable signals at the LHC.

In this work, we compute the FCNC decays of $t \to V q$ $(V = Z, \gamma; q = u, c)$ in an extension of mirror fermion model [\[21\]](#page-27-5) originally proposed by one of us [\[22\]](#page-27-6). In contrast with various left-right symmetric models, the model in [\[22\]](#page-27-6) did not include the gauge group $SU(2)_R$ while adding the mirror partners of the SM fermions. Despite having the same SM gauge group, the scalar sector must be enlarged. In additional to employ the bi-triplets in the Georgi-Machacek (GM) model [\[23,](#page-27-7) [24\]](#page-27-8) and a Higgs singlet to implement the electroweak scale seesaw mechanism for the non-sterile right-handed neutrino masses [\[22\]](#page-27-6), one needs to add a mirror Higgs doublet [\[25\]](#page-27-9) in the scalar sector so as to make consistency with the various signal strengths of the 125 GeV Higgs measured at the LHC. We will briefly review this class of mirror fermion model and its further extension with a horizontal A_4 symmetry in the following section.

We layout the paper as follows. In Section [II,](#page-4-0) after giving a brief highlight on some of the salient features of the model, we present the relevant interaction Lagrangian. Our calculation and analysis are presented in Section [III](#page-10-0) and [IV](#page-11-0) respectively. We finally summarize our results in Section [V.](#page-16-0) Analytical expressions for the loop functions are collected in the Appendix.

II. THE MIRROR FERMION MODEL

A. A Lightning Review

As already eluded to in the Introduction, the mirror fermion model in [\[22\]](#page-27-6) was devised to implement the so-called electroweak scale seesaw mechanism for the neutrino masses. We first list the particle content of the model in Table [I](#page-5-0) for further discussions. One special feature of this mirror model is to treat the right-handed neutrino in each generation to be non-sterile by grouping it with a new heavy mirror right-handed charged lepton into a weak doublet l_{Ri}^M , regarded as the mirror of the SM doublet l_{Li} with i labelling the generation. When the Higgs singlet ϕ_{0S} develops a small vacuum expectation value (VEV) of order 10⁵ eV, through its Yukawa couplings between the SM lepton doublets and their mirror partners, it can provide a small Dirac mass term for the neutrinos. On the other hand, when the triplet $\tilde{\chi}$ field with hypercharge $Y/2 = 1$ in Table [I](#page-5-0) develops a VEV of order $v_{\rm SM} = 246$ GeV, a Majorana mass term of electroweak scale can be generated through its Yukawa couplings among these new mirror lepton doublets. Details of this electroweak scale seesaw mechanism in the mirror fermion model can be found in [\[22\]](#page-27-6).

Note that the other triplet ξ which has zero hypercharge is grouped with $\tilde{\chi}$ to form

Fields	$(SU(3), SU(2), U(1)_Y; A_4)$	
$l_{Li} = \left(\begin{array}{c} \nu_L \ e_L \end{array} \right) \quad , \ \ l_{Ri}^M = \left(\begin{array}{c} \nu_R \ e_R \ e_R^M \end{array} \right) \, .$	$(1, 2, -\frac{1}{2}; 3)$	
e_{Ri} , e_{Li}^M	$(1, 1, -1; 3)$	
$\left q_{Li} = \left(\begin{array}{c} u_L \ d_L \end{array} \right) \right \;\;,\;\; q_{Ri}^M = \left(\begin{array}{c} u_R^M \ d_R^M \end{array} \right)_i$	$(3, 2, \frac{1}{6} ; 3)$	
u_{Ri} , u_{Li}^M	$(3, 1, \frac{2}{3} ; 3)$	
d_{Ri} , d_{Li}^M	$(3, 1, -\frac{1}{3} ; 3)$	
ϕ_{0S}	(1, 1, 0; 1)	
ϕ_{iS}	(1, 1, 0; 3)	
Φ $\ ,\ \ \Phi_M$	$(1, 2, \frac{1}{2}; 1)$	
ξ	(1, 3, 0; 1)	
$\tilde{\chi}$	(1, 3, 1; 1)	

TABLE I: The Standard Model quantum numbers of the fermion and scalar sectors in the extended mirror model together with their assignments under the horizontal A⁴ symmetry.

the bi-triplets in the GM model to maintain the custodial symmetry and therefore the ρ parameter equals unity at tree level. In [\[26\]](#page-27-10), the potential dangerous contributions from the GM triplets to the S and T oblique parameters are shown to be partially cancelled by the opposite contributions from mirror fermions such that the model is still healthy against electroweak precision tests.

Mirrors of other SM fermions, both leptons and quarks, can be introduced in the same way as listed in Table [I.](#page-5-0) Searches for these heavy mirror fermions at the LHC were presented in [\[27\]](#page-27-11) and [\[28\]](#page-27-12) for the mirror quarks and mirror leptons respectively.

In order to reproduce the signal strengths of $h_{125} \rightarrow \gamma\gamma$ and $h_{125} \rightarrow Z\gamma$ for the 125 GeV Higgs observed at the LHC, a mirror Higgs doublet Φ_M of the SM one Φ was introduced [\[25\]](#page-27-9). We note that mixing effects among the two doublets as well as with the triplet ξ must be taken into account in order to satisfy the LHC results. A global $U(1) \times U(1)$ symmetry was also enforced in the Yukawa interactions so that the SM Higgs doublet only couples to the SM fermions and the mirror Higgs doublet only couples to the mirror fermions. Thus there is no FCNC Higgs interactions at tree level in the model. Processes like $h \to \tau \mu$ [\[29\]](#page-28-0) and $t \to hq$ can only occur at the quantum loop level.

To address the issues of neutrino and charged lepton masses and mixings, the original mirror model was extended in [\[21\]](#page-27-5) by introducing a horizontal family symmetry of the tetrahedral group A_4 . The A_4 assignments of all the scalars and fermions as well as their SM quantum numbers are shown at the last column in Table [I.](#page-5-0) The lone singlet ϕ_{0S} is now accompanied with a A_4 triplet $\vec{\phi}_S = (\phi_{1S}, \phi_{2S}, \phi_{3S})$. Both ϕ_{0S} and $\vec{\phi}_S$ are electroweak singlets and they are the only fields communicating the SM sector with the mirror sector through the Yukawa couplings, which must be invariant under both gauge symmetry and A_4 . Other scalars are A_4 singlets.

Phenomenological implications of the extended mirror fermion model with the A_4 symmetry have been explored for the charged lepton flavor violating (CLFV) processes $\mu \to e\gamma$ [\[30\]](#page-28-1), $\mu - e$ conversion [\[31\]](#page-28-2) and $h_{125} \to \tau \mu$ [\[29\]](#page-28-0), as well as for the electron electric dipole moment [\[32\]](#page-28-3). Here we will explore its implication in the rare FCNC top decays. Implications of the A_4 symmetry for the quark masses and mixings will be given in [\[33\]](#page-28-4).

B. Interaction Lagrangian for Quarks and Their Mirrors

Here we will write down the interactions for the quarks and their mirrors that are relevant to the FCNC processes $t \to Vq$ that we are studying. Since the result for $t \to gq$ can be easily obtained from that of $t \to \gamma q$, we will not present detailed formulas for the former process. As for $t \to hq$, one must consider the mixing effects from the more complicated Higgs sector in the extended mirror model. We will leave it for future work.

1. Quark Yukawa Couplings with A⁴ Symmetry

Recall that the tetrahedron symmetry group A_4 has four irreducible representations $1, 1', 1''$, and 3 with the following multiplication rule

$$
3 \times 3 = 3_1(23, 31, 12) + 3_2(32, 13, 21) + 1(11 + 22 + 33) + 1'(11 + \omega^2 22 + \omega 33) + 1''(11 + \omega 22 + \omega^2 33)
$$
(4)

where $\omega = e^{2\pi i/3} = -\frac{1}{2} + i$ $\sqrt{3}$ $\frac{2}{2}$.

Using the above A_4 multiplication rules one can construct new Yukawa couplings in the leptonic sector, which are both gauge invariant and A_4 symmetric, to implement small Dirac neutrino masses in electroweak seesaw and to discuss charged lepton mixings [\[21\]](#page-27-5).

In the same vein, one can write down the following new Yukawa couplings for the quarks and their mirrors (both in the flavor basis with subscripts "0") with the scalar singlets, which are both gauge invariant and A_4 symmetric,

$$
-\mathcal{L}_{Y} \supset g_{0S}^{Q} \phi_{0S}(\overline{q_{L,0}} q_{R,0}^{M})_{1} + g_{1S}^{Q} \overline{\phi}_{S} \cdot (\overline{q_{L,0}} \times q_{R,0}^{M})_{3_{1}} + g_{2S}^{Q} \overline{\phi}_{S} \cdot (\overline{q_{L,0}} \times q_{R,0}^{M})_{3_{2}} + g_{0S}^{u} \phi_{0S}(\overline{u_{R,0}} u_{L,0}^{M})_{1} + g_{1S}^{u} \overline{\phi}_{S} \cdot (\overline{u_{R,0}} \times u_{L,0}^{M})_{3_{1}} + g_{2S}^{u} \overline{\phi}_{S} \cdot (\overline{u_{R,0}} \times u_{L,0}^{M})_{3_{2}} \qquad (5) + g_{0S}^{d} \phi_{0S}(\overline{d_{R,0}} d_{L,0}^{M})_{1} + g_{1S}^{d} \overline{\phi}_{S} \cdot (\overline{d_{R,0}} \times d_{L,0}^{M})_{3_{1}} + g_{2S}^{d} \overline{\phi}_{S} \cdot (\overline{d_{R,0}} \times d_{L,0}^{M})_{3_{2}} + \text{H.c.}
$$

where $g_{0S}^{Q,u,d}$ $g_{0S}^{Q,u,d}, g_{1S}^{Q,u,d}$ $q_{1S}^{Q,u,d}$ and $q_{2S}^{Q,u,d}$ $\mathcal{L}_{2S}^{Q,u,a}$ are in general complex coupling constants. Implications of the above Yukawa interactions on the quark mixings will be presented in [\[33\]](#page-28-4).

Next we move to the physical basis by making the following unitary transformations on the left-handed fields

$$
u_{L,0} = V_L^u u_L, \ d_{L,0} = V_L^d u_L, \ u_{L,0}^M = V_L^{u^M} u_L^M, \ d_{L,0}^M = V_L^{d^M} d_L^M,
$$

and similarly for the right-handed fields

$$
u_{R,0} = V_R^u u_R, \ d_{R,0} = V_R^d d_R, \ u_{R,0}^M = V_R^{u^M} u_R^M, \ d_{R,0}^M = V_R^{d^M} d_R^M.
$$

We can then recast the Yukawa interactions in the following form

$$
\mathcal{L}_Y \supset - \bar{u} \left(V_L^{u\dagger} M_S^Q(\phi) V_R^{uM} P_R + V_R^{u\dagger} M_S^u(\phi) V_L^{uM} P_L \right) u^M
$$

$$
- \bar{d} \left(V_L^{d\dagger} M_S^Q(\phi) V_R^{dM} P_R + V_R^{d\dagger} M_S^d(\phi) V_L^{dM} P_L \right) d^M + \text{H.c.}
$$
 (6)

with $P_{L,R} = (1 \mp \gamma_5)/2$. Here $M_S^{Q,u,d}$ $S^{Q,u,a}(\phi)$ are three field-dependent three by three matrices which can be decomposed in terms of the four scalar fields according to

$$
M_S^Q(\phi) = M^{Q,0}\phi_{0S} + M^{Q,1}\phi_{1S} + M^{Q,2}\phi_{2S} + M^{Q,3}\phi_{3S} ,\qquad (7)
$$

where

$$
M^{Q,0} = \begin{pmatrix} g_{0S}^Q & 0 & 0 \\ 0 & g_{0S}^Q & 0 \\ 0 & 0 & g_{0S}^Q \end{pmatrix} , \quad M^{Q,1} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & g_{1S}^Q \\ 0 & g_{2S}^Q & 0 \end{pmatrix} ,
$$

\n
$$
M^{Q,2} = \begin{pmatrix} 0 & 0 & g_{2S}^Q \\ 0 & 0 & 0 \\ g_{1S}^Q & 0 & 0 \end{pmatrix} , \quad M^{Q,3} = \begin{pmatrix} 0 & g_{1S}^Q & 0 \\ g_{2S}^Q & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} ,
$$
\n(8)

and similar decompositions for $M_S^u(\phi)$ and $M_S^d(\phi)$ with $M^{u,k}$ and $M^{d,k}$ obtained by the substitutions of $g_{iS}^Q \to g_{iS}^u$ and g_{iS}^d respectively in Eq. [\(8\)](#page-8-0). Introducing the following combinations of the coupling matrices $M^{Q,k}$ and $M^{q,k}$ $(k = 0, 1, 2, 3)$ with the fermion mixing matrices

$$
V_L^{q,k} \equiv V_L^{q\dagger} M^{Q,k} V_R^{q^M} ,
$$

\n
$$
V_R^{q,k} \equiv V_R^{q\dagger} M^{q,k} V_L^{q^M} ,
$$
\n(9)

we arrive at the final form of the Yukawa interactions

$$
\mathcal{L}_Y \supset -\sum_{k=0}^3 \sum_{i,j=1}^3 \bar{u}_i \left\{ V_L^{u,k}{}_{ij} P_R + V_R^{u,k}{}_{ij} P_L \right\} u_j^M \phi_{kS} + (u \leftrightarrow d) + \text{H.c.} \tag{10}
$$

We have combined the four scalars ϕ_{0S} and $\vec{\phi}_S$ into ϕ_{kS} with $k = 0, 1, 2, 3$. For the FCNC rare top decays that we are studying, only $V_{(L)}^{u,k}$ $\chi_{(L,R)}^{u,\kappa}$ are relevant.

2. Neutral Currents

We also need the neutral current interactions for the SM Z boson and photon couple to the quarks and their mirrors.

$$
\mathcal{L}_{\rm NC} \supset g Z^{\mu} J_{\mu}^{Z} + e A^{\mu} J_{\mu}^{\rm EM} \tag{11}
$$

with

$$
J_{\mu}^{Z} = \frac{1}{\cos \theta_{W}} \left[\bar{q}_{L} \gamma_{\mu} \left(T^{3} - Q_{q} \sin^{2} \theta_{W} \right) q_{L} - \bar{q}_{R} \gamma_{\mu} Q_{q} \sin^{2} \theta_{W} q_{R} \right. \\
\left. + \bar{q}_{R}^{M} \gamma_{\mu} \left(T^{3} - Q_{q} \sin^{2} \theta_{W} \right) q_{R}^{M} - \bar{q}_{L}^{M} \gamma_{\mu} Q_{q} \sin^{2} \theta_{W} q_{L}^{M} \right] \,,
$$
\n(12)

$$
J_{\mu}^{\text{EM}} = Q_q \left(\bar{q} \gamma_{\mu} q + \bar{q}^M \gamma_{\mu} q^M \right) \tag{13}
$$

The above neutral current interactions in SM and the new Yukawa couplings can induce FCNC decay $t \rightarrow Vq$ at one-loop level as depicted by the three Feynman diagrams in Fig. [1.](#page-10-1)

FIG. 1: Feynman diagrams contributing to $t \to Vq$.

III. FCNC TOP DECAYS $t \rightarrow Vq$

The effective Lagrangian for $t\to Zq$ and $t\to \gamma q$ can be expressed as

$$
\mathcal{L}_{\text{eff}} = -\bar{q}\gamma_{\mu}(C_{L}P_{L} + C_{R}P_{R})tZ^{\mu} - \frac{1}{m_{t}}\bar{q}\sigma_{\mu\nu}(A_{L}P_{L} + A_{R}P_{R})tZ^{\mu\nu}
$$

$$
-\frac{1}{m_{t}}\bar{q}\sigma_{\mu\nu}(A'_{L}P_{L} + A'_{R}P_{R})tF^{\mu\nu} + \text{H.c.}
$$
(14)

where $q = (u, c)$; $Z^{\mu\nu} = \partial^{\mu}Z^{\nu} - \partial^{\nu}Z^{\mu}$ and $F^{\mu\nu} = \partial^{\mu}A^{\nu} - \partial^{\nu}A^{\mu}$; and $A_{L,R}$, $A'_{L,R}$ and ${\cal C}_{L,R}$ are dimensionless quantities.

In terms of the dimensionless mass ratios

$$
r_q \equiv m_q/m_t \quad , \qquad r_Z \equiv m_Z/m_t \tag{15}
$$

the partial decay rate for $t\to Zq$ is given by

$$
\Gamma(t \to qZ) = \frac{1}{16\pi} \frac{1}{m_t} \lambda^{\frac{1}{2}} \left(1, r_q^2, r_Z^2\right) \left\langle \sum |\mathcal{M}|^2 \right\rangle ,\qquad (16)
$$

where
$$
\lambda(1, y, z) = (1 - (\sqrt{y} + \sqrt{z})^2)(1 - (\sqrt{y} - \sqrt{z})^2)
$$
 and
\n
$$
\left\langle \sum |\mathcal{M}|^2 \right\rangle = \frac{m_t^2}{2} \left\{ +2 \left(|C_L|^2 + |C_R|^2 \right) \left(1 + r_q^2 - r_Z^2 \right) + 4 \left(|A_L|^2 + |A_R|^2 \right) \left[2 \left(1 - r_q^2 \right)^2 - \left(1 + r_q^2 \right) r_Z^2 - r_Z^4 \right] - 16 \operatorname{Re} (C_L C_R^*) r_q - 48 \operatorname{Re} (A_L A_R^*) r_q r_Z^2 - 12 \operatorname{Re} (C_L A_L^* + C_R A_R^*) r_q \left(1 - r_q^2 + r_Z^2 \right) + 12 \operatorname{Re} (C_L A_R^* + C_R A_L^*) \left(1 - r_q^2 - r_Z^2 \right) + \frac{1}{r_Z^2} \left[\left(|C_L|^2 + |C_R|^2 \right) \left(\left(1 - r_q^2 \right)^2 - \left(1 + r_q^2 \right) r_Z^2 \right) + 4 \operatorname{Re} (C_L C_R^*) r_q r_Z^2 \right] \right\}.
$$
 (17)

For a given model, the dimensionless quantities $A_{L,R}$ and $C_{L,R}$ can be determined. In the mirror fermion model, these quantities are induced at one loop level, as depicted by the Feynman diagrams in Fig. [1.](#page-10-1) Their analytical expressions are given in the Appendix.

Similarly, for $t \to \gamma q$ we have

$$
\Gamma(t \to q\gamma) = \frac{1}{16\pi} \frac{1}{m_t} \lambda^{\frac{1}{2}} \left(1, r_q^2, 0\right) \left\langle \sum |\mathcal{M}|^2 \right\rangle \tag{18}
$$

with $\langle \sum |\mathcal{M}|^2 \rangle = 4m_t^2(1 - r_q^2)^2(|A'_L|^2 + |A'_R|^2)$. The expressions for A'_L and A'_R are also given in the Appendix.

IV. ANALYSIS

In our numerical analysis, we will make the following assumptions on the parameter space of the model.

(1) First, we will take all the unknown Yukawa couplings to be real and assume $g_{iS}^q = g_{iS}^Q$ for $q = (u, d)$ and $i = 0, 1, 2$. We will explore how our results depend on the couplings g_{iS}^Q . We note that it has been shown recently [\[34\]](#page-28-5) in the extended mirror fermion model [\[25\]](#page-27-9) the complex values of some of these Yukawa couplings, combined with the electroweak scale seesaw mechanism generating the minuscule neutrino masses, one can provide a solution to the strong CP problem without introducing axion.

(2) Since only the product $V_{\text{CKM}} = (V_L^u)^{\dagger} V_L^d$ are known experimentally, we will study the following scenarios for illustrative purpose. Scenario 1:

$$
V_L^u = V_{\text{CKM}}^{\dagger} ,
$$

\n
$$
V_R^u = V_L^{u^M} = V_R^{u^M} = 1 .
$$
\n(19)

Scenario 2:

$$
V_L^u = V_L^{u^M} = V_{\text{CKM}}^{\dagger} ,
$$

\n
$$
V_R^u = V_R^{u^M} = 1 .
$$
\n(20)

(3) For the three generation of mirror quark masses, we assume

$$
m_{q_1^M} : m_{q_2^M} : m_{q_3^M} = M : M + 10 \text{ GeV} : M + 20 \text{ GeV} , \qquad (21)
$$

and vary the common mirror quark mass M from 150 to 800 GeV. We note that mirror fermions in this class of electroweak scale mirror fermion model are expected to have masses of electroweak scale to satisfy unitarity [\[27\]](#page-27-11).

- (4) For the scalars ϕ_{kS} , their masses are necessarily small since they are link to the Dirac neutrino masses [\[21,](#page-27-5) [22\]](#page-27-6). We set their masses m_{kS} all equal 10 MeV. Our numerical results are not sensitive to this choice as long as $m_{kS} \ll m_{q_m^M}$.
- (5) For the SM parameters, we use [\[35\]](#page-28-6)

$$
m_t = 173.21 \,\text{GeV} \ , \ m_c = 1.275 \,\text{GeV} \ , \ m_u = 2.3 \,\text{MeV} \ ,
$$

\n
$$
\sin^2 \theta_W = 0.23126 \ , \ \alpha = 1/127.944 \ ,
$$

\n
$$
\Gamma_t = 1.41 \,\text{GeV} \ , \ \mathcal{B} \left(t \to W^+ b \right) = 0.957 \ .
$$
\n(22)

FIG. 2: Branching ratios of $t \to \gamma u$ (left) and $t \to \gamma c$ (right) versus the logarithmic of Yukawa couplings g_{0}^Q $_{0S}^{Q}$ and g_{1S}^{Q} with $g_{2S}^{Q} = 10^{-3}$ and $M = 150$ GeV in Scenario 1.

FIG. 3: Branching ratios of $t \to Zu$ (left) and $t \to Ze$ (right) versus the logarithmic of Yukawa couplings g_{0}^Q $_{0S}^{Q}$ and g_{1S}^{Q} with $g_{2S}^{Q} = 10^{-3}$ and $M = 150$ GeV in Scenario 1.

In Fig. [\(2\)](#page-13-0), we show the contour plots of log $\mathcal{B}(t \to \gamma u)$ (left panel) and log $\mathcal{B}(t \to \gamma u)$ γc) (right panel) on the $(\log_{10}(g_{0.9}^Q$ $_{0S}^{Q}), \log_{10}(g_{1S}^{Q})$ $\binom{Q}{1S}$) plane in the case of Scenario 1 with

 g_{2}^Q $_{2S}^{Q}$ set to be 10⁻³. Fig. [\(3\)](#page-13-1) is similar as Fig. [\(2\)](#page-13-0) but for $t → Zq$. Figs. [\(4\)](#page-14-0) and [\(5\)](#page-14-1) are the same as Figs. [\(2\)](#page-13-0) and [\(3\)](#page-13-1) respectively but for Scenario 2. The common mirror fermion mass M is set to be 150 GeV in these 4 figures.

FIG. 4: Same as Fig. [\(2\)](#page-13-0) in Scenario 2.

FIG. 5: Same as Fig. [\(3\)](#page-13-1) in Scenario 2.

Mirror quarks may be pair produced at the LHC [\[27\]](#page-27-11). Once produced the heavier mirror fermions may cascade into lighter ones by emitting an on-shell or off-shell SM W-boson, depending on the detail mass spectrum of the mirror fermions. The lightest mirror quark will then decay into its SM partner with any one of the scalar singlets ϕ_{kS} via the new Yukawa interactions which are responsible to the FCNC decays of the top quark studied here. In the mirror lepton case, the corresponding Yukawa couplings g_{iS}^L are necessarily small since they are responsible for providing small Dirac masses to the neutrinos in the electroweak scale seesaw mechanism. Assuming the lightest mirror fermion is u^M . In Fig. [\(6\)](#page-16-1), we plot the contours of the decay length of u^M in the $(M, \log_{10}(g_{0S}^Q))$ $\binom{Q}{0S}$) plane. We take all the Yukawa couplings to be the same just for illustrations. One can see that for very small Yukawa couplings $< 10^{-6}$, can the decay length reach a few mm for a displaced vertex. Search strategies for the mirror fermions would then be quite different from the usual cases, involving not merely the missing energies but displaced vertices as well [\[27\]](#page-27-11). Current experiments at the LHC have the capability to perform such kind of searches. Further studies of this issue are warranted.

Nevertheless, for the mirror quarks, there is no a priori reason that these new Yukawa couplings have to be very small except that there are stringent constraints from the mixings between SM fermions and their mirrors. The mixing angle is roughly of order $g_{iS}^Q \langle \phi_{kS} \rangle / M$. For $g_{iS}^Q \sim 1$, $\langle \phi_{kS} \rangle \sim 1$ MeV and $M \sim 500$ GeV, this mixing angle is about 2×10^{-6} . A full analysis taking into the account of the mixing effects is beyond the scope of this paper.

In Figs. [\(7\)](#page-17-0) and [\(8\)](#page-18-0), we show the scatter plots for the logarithmic of branching ratios of the 4 processes, $t \to \gamma u$ and $t \to \gamma c$ in the left panel and $t \to Zu$ and $t \to Zc$ in the right panel, versus $\log_{10}(g_{0.8}^Q)$ $\binom{Q}{0S}$ for Scenarios 1 and 2 respectively. We have set all the Yukawa couplings equal to each other in these plots. The different colors in the scatter plots represent different values of the common mirror fermion

FIG. 6: Decay length of the lightest mirror quark versus $\log_{10}(g_{0.9}^Q)$ $\binom{Q}{0S}$ assuming all the unknown Yukawa couplings equal to each other.

mass M varied from 150 to 800 GeV as indicated by the color palettes on the top of each plot. Current experimental limits of these processes are also shown in these plots by the horizontal red dashed lines, while the black dashed lines are the SM predictions. It is clear from these plots that the mirror quarks in this class of model with mass less than 800 GeV could play an important role in FCNC decays of the top quark, provided that the Yukawa couplings are of the same size as the top quark Yukawa coupling in SM. However if the Yukawa couplings are very small to allow for a displaced vertex for the lightest mirror fermion, all these FCNC top decays are beyond the reach at LHC.

V. CONCLUSION

In Table [II,](#page-19-0) we summarize our numerical results as well as those from SM and other three popular new physics models taken from [\[36\]](#page-28-7) for comparisons. The numbers in

FIG. 7: Scatter plots for the branching ratios of $t\to Vq$ in Scenario 1.

brackets in the 2HDM and MSSM columns are for 2HDM with tree level flavor violation and MSSM with R-parity violation respectively. Our results shown in the last column are taken from Figs. [\(7\)](#page-17-0) and [\(8\)](#page-18-0) for $g_{0S}^Q = y_t =$ √ $2m_t/v_{\rm SM} \sim 1$ and the mirror mass M varying from 150 to 800 GeV. On the other hand, if $|g_{0}^{Q}$ $\frac{Q}{0S}$ turns out

FIG. 8: Same as Fig. [\(7\)](#page-17-0) in Scenario 2.

to be small, of order 10[−]⁴ or less as suggested by the new solution to the strong CP problem discussed in [\[34\]](#page-28-5), all FCNC top decays in the model would be unobservable.

While the experimental results of many FCNC processes in the kaon, D and B meson systems, accumulated over the past several decades, had been mostly consis-

Process	SМ	2HDM	MSSM	Extra-Dimension Mirror Model	
		$t \rightarrow Zc$ $\left 1 \times 10^{-14} \right \le 10^{-10} (10^{-6}) \left \le 10^{-7} (10^{-6}) \right $		$\leq 10^{-5}$	$10^{-6} - 10^{-8}$
		$t{\rightarrow}Zu$ $ 7\times10^{-17} $ -	$\leq 10^{-7} (10^{-6})$		$10^{-8} - 10^{-10}$
		$t \rightarrow \gamma c \left 5 \times 10^{-14} \right \leq 10^{-9} (10^{-7}) \left \leq 10^{-8} (10^{-9}) \right $		$\leq 10^{-9}$	$10^{-7} - 10^{-9}$
	$t\rightarrow \gamma u \ \left 4\times10^{-16}\right $	$\mathcal{L}^{\text{max}}_{\text{max}}$	$\leq 10^{-8} (10^{-9})$		$10^{-8} - 10^{-11}$

TABLE II: Comparisons of theoretical predictions for the branching ratios of FCNC rare top decay $t \to Vq$ in various models. The numbers in brackets in 2HDM and MSSM are for 2HDM with tree level flavor violations and MSSM with R-parity violation respectively.

tent with the SM expectations, theoretical predictions for FCNC processes involving the top quark and/or the Higgs boson have not been challenged by high energy experiments until recent years.

In this work, we have computed the FCNC processes $t \to Zq$ and γq in a class of mirror fermion model equipped with a horizontal A_4 symmetry in the fermion and scalar sectors. We found that branching ratio for $t \to Zc$ is typically of order $10^{-8} - 10^{-6}$ as mirror quark masses are varying in the range from 800 to 150 GeV and new Yukawa couplings are of the same size of the top Yukawa coupling in SM. At 14 TeV the total cross section for $t\bar{t}$ production at the LHC is about 598 pb. With a luminosity of 300 (1000) fb⁻¹, we thus expect 180 (6) events of $t \to Zc$ for a branching ratio of 10^{-6} (10^{-8}) before any experimental cuts. For the other processes $t \to Zu$ and $t \to \gamma q$, their branching ratios are typically smaller by 1 – 2 orders of magnitude.

For the gluon mode $t \to gq$ its partial width is about 42 times larger than that of the photon mode $t \to \gamma q$. The LHC limits for the branching ratios of $t \to gu$ and $t \to gc$ are 2.0×10^{-5} and 4.1×10^{-4} respectively from CMS [\[37\]](#page-28-8), and 4.0×10^{-5} and 20×10^{-5} respectively from ATLAS [\[38\]](#page-28-9). These branching ratios are extracted from the single top production via FCNC interactions from gluon plus up- or charmquark initial states. They are still $1 - 2$ orders of magnitude above our theoretical predictions.

It is also interesting to consider FCNC processes involving both the heavy top quark and the 125 GeV Higgs, the two heaviest particles in the SM. One particular important process is $t \to hq$, which LHC has obtained the following limits [\[39,](#page-28-10) [40\]](#page-28-11)

$$
\mathcal{B}(t \to hu) \leq \begin{cases} 4.5 \times 10^{-3} \text{ [ATLAS]} ,\\ 5.5 \times 10^{-3} \text{ [CMS]} ; \end{cases}
$$
 (23)

$$
\mathcal{B}(t \to hc) \le \begin{cases} 4.6 \times 10^{-3} [\text{ATLAS}] ,\\ 4.0 \times 10^{-3} [\text{CMS}]. \end{cases}
$$
 (24)

In the mirror fermion model, the mirror Higgs as well as the GM triplets could be an imposter for the 125 GeV Higgs due to mixing effects, which must be taken into account. This work will be reported elsewhere [\[41\]](#page-28-12).

LHC has unique opportunity for probing the top quark FCNC decays in new physics models since the SM contributions are at least nine orders of magnitude below the current limits of these processes. With its high luminosity upgrade in the second phase, HL-LHC can impose powerful constraints on any underlying new physics responsible for the FCNC interactions.

ACKNOWLEDGMENTS

We would like to thank Chuan-Ren Chen for stimulating discussions. This work is supported by the Ministry of Science and Technology (MoST) of Taiwan under Grant Number MOST-104-2112-M-001-001-MY3.

APPENDIX

Form Factors for $t\to Zq$

All three Feynman diagrams (A), (B) and (C) in Fig. [\(1\)](#page-10-1) contribute to the form factors ${\cal C}_L$ and ${\cal C}_R$:

$$
C_{(L,R)} = C_{(L,R)}^A + C_{(L,R)}^B + C_{(L,R)}^C.
$$

To minimize cluttering in our expressions given below, we define

$$
a = \frac{g}{4\cos\theta_W} \left(1 - \frac{8}{3}\sin^2\theta_W\right) , \qquad b = \frac{g}{4\cos\theta_W} ,
$$

and

$$
V_{ij}^{L,k} = \left(V_L^{u,k}\right)_{ij}, \qquad V_{ij}^{R,k} = \left(V_R^{u,k}\right)_{ij},
$$

where $V_{(L)}^{u,k}$ $\chi_{(L,R)}^{u,\kappa}$ are given by Eq. [\(9\)](#page-9-0). The individual contributions from each diagrams can be computed using the automated tools FormCalc and LoopTools in FeynArts [\[42\]](#page-28-13). The results are listed as follows.

$$
C_{L}^{A} = \frac{1}{(m_{t}^{2} - m_{q}^{2})} \frac{(a+b)}{16\pi^{2}} \sum_{k=0}^{3} \sum_{m=1}^{3} \left\{ m_{q}^{2} V_{qm}^{L,k} V_{tm}^{L,k^{*}} \left[B_{0}(m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right. \right.+ B_{1}(m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \left. \right] + m_{t} m_{q} V_{qm}^{R,k} V_{tm}^{R,k^{*}} \left[B_{0}(m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right.+ B_{1}(m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \left. \right] + m_{q} m_{q_{m}^{M}} V_{qm}^{R,k} V_{tm}^{L,k^{*}} B_{0}(m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) + m_{t} m_{q_{m}^{M}} V_{qm}^{L,k} V_{tm}^{R,k^{*}} B_{0}(m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \left. \right\},
$$
\n
$$
(25)
$$

$$
C_{L}^{B} = \frac{-1}{(m_{t}^{2} - m_{q}^{2})} \frac{(a+b)}{16\pi^{2}} \sum_{k=0}^{3} \sum_{m=1}^{3} \left\{ m_{t}^{2} V_{qm}^{L,k} V_{tm}^{L,k*} \left[B_{0}(m_{t}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right. \right.\left. + B_{1}(m_{t}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right] + m_{t} m_{q} V_{qm}^{R,k} V_{tm}^{R,k*} \left[B_{0}(m_{t}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right. \right.\left. + B_{1}(m_{t}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right] + m_{q} m_{q_{m}^{M}} V_{qm}^{R,k} V_{tm}^{L,k*} B_{0}(m_{t}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) + m_{t} m_{q_{m}^{M}} V_{qm}^{L,k} V_{tm}^{R,k*} B_{0}(m_{t}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right\},
$$
\n(26)

and

$$
C_{L}^{C} = \frac{-1}{16\pi^{2}} \sum_{k=0}^{3} \sum_{m=1}^{3} \Big\{ (a+b)V_{qm}^{L,k}V_{tm}^{L,k*} \Big[\frac{1}{2} - 2C_{00}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) \Big] + m_{q_{m}^{M}}^{2}(a-b)V_{qm}^{L,k}V_{tm}^{L,k*} C_{0}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + m_{Z}^{2}(a+b)V_{qm}^{L,k}V_{tm}^{L,k*} C_{12}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + m_{t}m_{q}(a-b)V_{qm}^{L,k}V_{tm}^{R,k*} [2C_{1}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + 2C_{2}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + 2C_{12}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + C_{11}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + C_{22}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + C_{0}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + C_{0}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m
$$

$$
C_{R}^{A} = \frac{1}{(m_{t}^{2} - m_{q}^{2})} \frac{(a - b)}{16\pi^{2}} \sum_{k=0}^{3} \sum_{m=1}^{3} \left\{ m_{q}^{2} V_{qm}^{R,k} V_{tm}^{R,k*} \left[B_{0} (m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right. \right.\left. + B_{1} (m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right] + m_{t} m_{q} V_{qm}^{L,k} V_{tm}^{L,k*} \left[B_{0} (m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right.\left. + B_{1} (m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right] + m_{q} m_{q_{m}^{M}} V_{qm}^{L,k} V_{tm}^{R,k*} B_{0} (m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \left. + m_{t} m_{q_{m}^{M}} V_{qm}^{R,k} V_{tm}^{L,k*} B_{0} (m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right\},
$$
\n
$$
(28)
$$

$$
C_{R}^{B} = \frac{-1}{(m_{t}^{2} - m_{q}^{2})} \frac{(a - b)}{16\pi^{2}} \sum_{k=0}^{3} \sum_{m=1}^{3} \left\{ m_{t}^{2} V_{qm}^{R,k} V_{tm}^{R,k^{*}} \left[B_{0}(m_{t}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right. \right.\left. + B_{1}(m_{t}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right] + m_{t} m_{q} V_{qm}^{L,k} V_{tm}^{L,k^{*}} \left[B_{0}(m_{t}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right.\left. + B_{1}(m_{t}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right] + m_{q} m_{q_{m}^{M}} V_{qm}^{L,k} V_{tm}^{R,k^{*}} B_{0}(m_{t}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \left. + m_{t} m_{q_{m}^{M}} V_{qm}^{R,k} V_{tm}^{L,k^{*}} B_{0}(m_{t}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}) \right\},
$$
\n(29)

and

$$
C_{R}^{C} = \frac{-1}{16\pi^{2}} \sum_{k=0}^{3} \sum_{m=1}^{3} \Big\{ (a-b)V_{qm}^{R,k}V_{tm}^{R,k} \Big[\frac{1}{2} - 2C_{00}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{nt}^{M}}^{2}, m_{q_{nt}^{M}}^{2}) \Big] + m_{q_{m}^{M}}^{2}(a+b)V_{qm}^{R,k}V_{tm}^{R,k*} C_{0}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{nt}^{M}}^{2}, m_{q_{nt}^{M}}^{2}) + m_{Z}^{2}(a-b)V_{qm}^{R,k}V_{tm}^{R,k*} C_{12}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{nt}^{M}}^{2}, m_{q_{nt}^{M}}^{2}) + m_{t}m_{q}(a+b)V_{qm}^{L,k}V_{tm}^{L,k*} [2C_{1}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{nt}^{M}}^{2}, m_{q_{nt}^{M}}^{2}) + 2C_{2}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{nt}^{M}}^{2}, m_{q_{nt}^{M}}^{2}) + 2C_{12}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{nt}^{M}}^{2}, m_{q_{nt}^{M}}^{2}) + C_{11}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{nt}^{M}}^{2}, m_{q_{nt}^{M}}^{2}) + C_{22}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{nt}^{M}}^{2}, m_{q_{nt}^{M}}^{2}) + C_{0}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{nt}^{M}}^{2}, m_{q_{nt}^{M}}^{2}) + C_{1}(m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{
$$

Each of the above contributions $C_{L,R}^A$, $C_{L,R}^B$ and $C_{L,R}^C$ are ultraviolet divergent. However by using the divergent parts of the Passarino-Veltman (PV) functions

$$
\text{Div}[B_0] = +\Delta_{\epsilon} ,
$$
\n
$$
\text{Div}[B_1] = -\frac{1}{2}\Delta_{\epsilon} ,
$$
\n
$$
\text{Div}[C_{00}] = +\frac{1}{4}\Delta_{\epsilon} ,
$$
\n(31)

where $\Delta_{\epsilon} = 2/\epsilon - \gamma_E + \ln 4\pi$ with $\epsilon = 4 - d$ is the regulator in dimensional regularization and γ_E being the Euler's constant, one can easily verify that the divergences in the three diagrams summed up to nil leading to finite results for C_L and C_R .

Only Diagram (C) contributes to the dipole form factors A_L and A_R . They are given by

$$
A_{L} = \frac{1}{16\pi^{2}} \sum_{k=0}^{3} \sum_{m=1}^{3} \left\{ \frac{m_{t}^{2}}{2} (a-b) V_{qm}^{R,k} V_{tm}^{R,k*} \left[C_{1} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + C_{11} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + C_{12} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) \right] + \frac{m_{t} m_{q}}{2} (a+b) V_{qm}^{L,k} V_{tm}^{L,k*} \left[C_{2} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + C_{12} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}) + C_{22} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) \right] + \frac{m_{t} m_{q_{m}^{M}}}{2} V_{qm}^{R,k} V_{tm}^{L,k*} \left[(a-b) C_{1} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + (a+b) C_{2} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{ks}}^{2}, m_{q_{m}^{M}}^{2}) \right] \right\},
$$
\n(32)

and

$$
A_{R} = \frac{1}{16\pi^{2}} \sum_{k=0}^{3} \sum_{m=1}^{3} \left\{ \frac{m_{t}^{2}}{2} (a+b) V_{qm}^{L,k} V_{tm}^{L,k*} \left[C_{1} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + C_{11} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + C_{12} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) \right] + \frac{m_{t} m_{q}}{2} (a-b) V_{qm}^{R,k} V_{tm}^{R,k*} \left[C_{2} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + C_{12} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + C_{22} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) \right] + \frac{m_{t} m_{q_{m}^{M}} V_{L,k} V_{tm}^{R,k*}}{2} \left[(a+b) C_{1} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) + (a-b) C_{2} (m_{t}^{2}, m_{Z}^{2}, m_{q}^{2}, m_{\phi_{kS}}^{2}, m_{q_{m}^{M}}^{2}, m_{q_{m}^{M}}^{2}) \right] \right\}.
$$
\n(33)

Since the PV functions C_1, C_2, C_{11}, C_{12} and C_{22} do not have ultraviolet divergences, A_L and A_R are finite, as one should expect for they are the coefficients of the nonrenormalizable magnetic and electric dipole operators.

Form Factors for $t \to \gamma q$ and $t \to gq$

 A'_L and A'_R can be obtained from the above A_L and A_R respectively by replacing

$$
m_Z^2 \to 0
$$
, $a \to \frac{2}{3}e$, $b \to 0$. (34)

The decay rate for $t \to gq$ can be obtained from that of $t \to \gamma q$ simply by replacing the top quark electric charge $\frac{2}{3}e$ by the strong coupling g_s and multiply the final result by an overall color factor $(N_C^2 - 1)/2N_C$ where N_C is the number of color. Thus

$$
\frac{\Gamma(t \to gq)}{\Gamma(t \to \gamma q)} = \frac{9}{4} \cdot \frac{N_C^2 - 1}{2N_C} \cdot \frac{\alpha_s}{\alpha_{\rm em}}.
$$
\n(35)

Taking $N_C = 3$, $\alpha_s = 0.11$ and $\alpha_{em}^{-1} = 128$, this ratio is about 42. Next-to-leading order QCD corrections to the processes $t \to \gamma q$, $t \to Zq$ and $t \to gq$ can be found in [\[43,](#page-28-14) [44\]](#page-28-15). Moreover, the next-to-leading order and next-to-next-to-leading order QCD corrections for the dominant SM top quark decay mode $t \to W^+b$ had been computed in [\[45,](#page-29-0) [46\]](#page-29-1) and [\[47\]](#page-29-2) respectively.

- [1] S. L. Glashow, J. Iliopoulos and L. Maiani, Phys. Rev. D 2, 1285 (1970).
- [2] J. A. Aguilar-Saavedra, Acta Phys. Polon. B 35, 2695 (2004) [hep-ph/0409342].
- [3] For an updated top quark properties at the LHC, see the [Moriond talk](http://moriond.in2p3.fr/QCD/2017/WednesdayMorning/Soares.pdf) [\(http://moriond.in2p3.fr/QCD/2017/WednesdayMorning/Soares.pdf\)](http://moriond.in2p3.fr/QCD/2017/WednesdayMorning/Soares.pdf) by M. Soares.
- [4] [The CMS Collaboration], CMS PAS TOP-12-039.
- [5] [The CMS Collaboration], CMS PAS TOP-14-020.
- [6] G. Aad et al. [ATLAS Collaboration], Eur. Phys. J. C 76, no. 1, 12 (2016) [arXiv:1508.05796 [hep-ex]].
- [7] The ATLAS collaboration [ATLAS Collaboration], ATLAS-CONF-2017-070.
- [8] V. Khachatryan et al. [CMS Collaboration], JHEP 1604, 035 (2016) [arXiv:1511.03951 [hep-ex]].
- [9] J. J. Cao, G. Eilam, M. Frank, K. Hikasa, G. L. Liu, I. Turan and J. M. Yang, Phys. Rev. D 75, 075021 (2007) [hep-ph/0702264].
- [10] J. M. Yang, B. L. Young and X. Zhang, Phys. Rev. D 58, 055001 (1998) [hepph/9705341].
- [11] G. Eilam, A. Gemintern, T. Han, J. M. Yang and X. Zhang, Phys. Lett. B 510, 227 (2001) [hep-ph/0102037].
- [12] G. Eilam, J. L. Hewett and A. Soni, Phys. Rev. D 44, 1473 (1991) Erratum: [Phys. Rev. D 59, 039901 (1999)].
- [13] D. Atwood, L. Reina and A. Soni, Phys. Rev. D 55, 3156 (1997) [hep-ph/9609279].
- [14] S. Bejar, hep-ph/0606138.
- [15] C. Kao, H. Y. Cheng, W. S. Hou and J. Sayre, Phys. Lett. B 716, 225 (2012) [arXiv:1112.1707 [hep-ph]].
- [16] K. F. Chen, W. S. Hou, C. Kao and M. Kohda, Phys. Lett. B 725, 378 (2013) [arXiv:1304.8037 [hep-ph]].
- [17] G. Abbas, A. Celis, X. Q. Li, J. Lu and A. Pich, JHEP 1506, 005 (2015) [arXiv:1503.06423 [hep-ph]].
- [18] K. Agashe, G. Perez and A. Soni, Phys. Rev. D 75, 015002 (2007) [hep-ph/0606293].
- [19] K. Agashe and R. Contino, Phys. Rev. D 80, 075016 (2009) [arXiv:0906.1542 [hep-ph]].
- [20] H. Hesari, H. Khanpour and M. Mohammadi Najafabadi, Phys. Rev. D 92, no. 11, 113012 (2015) [arXiv:1508.07579 [hep-ph]]; H. Hesari, H. Khanpour, M. Khatiri Yanehsari and M. Mohammadi Najafabadi, Adv. High Energy Phys. 2014, 476490 (2014) [arXiv:1412.8572 [hep-ex]]; H. Khanpour, S. Khatibi, M. Khatiri Yanehsari and M. Mohammadi Najafabadi, arXiv:1408.2090 [hep-ph].
- [21] P. Q. Hung and T. Le, JHEP 1509, 001 (2015) Erratum: [JHEP 1509, 134 (2015)] [arXiv:1501.02538 [hep-ph]].
- [22] P. Q. Hung, Phys. Lett. B 649, 275 (2007) [hep-ph/0612004].
- [23] H. Georgi and M. Machacek, Nucl. Phys. B 262, 463 (1985).
- [24] M. S. Chanowitz and M. Golden, Phys. Lett. B 165, 105 (1985).
- [25] V. Hoang, P. Q. Hung and A. S. Kamat, Nucl. Phys. B 896, 611 (2015) [arXiv:1412.0343 [hep-ph]].
- [26] V. Hoang, P. Q. Hung and A. S. Kamat, Nucl. Phys. B 877, 190 (2013) [arXiv:1303.0428 [hep-ph]].
- [27] S. Chakdar, K. Ghosh, V. Hoang, P. Q. Hung and S. Nandi, Phys. Rev. D 93, no. 3, 035007 (2016) [arXiv:1508.07318 [hep-ph]].
- [28] S. Chakdar, K. Ghosh, V. Hoang, P. Q. Hung and S. Nandi, Phys. Rev. D 95, no. 1, 015014 (2017) [arXiv:1606.08502 [hep-ph]].
- [29] C. F. Chang, C. H. V. Chang, C. S. Nugroho and T. C. Yuan, Nucl. Phys. B 910, 293 (2016) [arXiv:1602.00680 [hep-ph]].
- [30] P. Q. Hung, T. Le, V. Q. Tran and T. C. Yuan, JHEP 1512, 169 (2015) [arXiv:1508.07016 [hep-ph]]; P. Q. Hung, Phys. Lett. B 659, 585 (2008) [arXiv:0711.0733 [hep-ph]].
- [31] P. Q. Hung, T. Le, V. Q. Tran and T. C. Yuan, arXiv:1701.01761 [hep-ph].
- [32] C. F. Chang, P. Q. Hung, C. S. Nugroho, V. Q. Tran and T. C. Yuan, arXiv:1702.04516 [hep-ph].
- [33] P. Q. Hung and T. Le, work in progress.
- [34] P. Q. Hung, arXiv:1710.00498 [hep-ph]; arXiv:1704.06390 [hep-ph].
- [35] C. Patrignani et al. (Particle Data Group), Chin. Phys. C, 40, 100001 (2016).
- [36] K. Agashe et al. [Top Quark Working Group], arXiv:1311.2028 [hep-ph].
- [37] V. Khachatryan et al. [CMS Collaboration], JHEP 1702, 028 (2017) [arXiv:1610.03545] [hep-ex]].
- [38] G. Aad et al. [ATLAS Collaboration], Eur. Phys. J. C 76, no. 2, 55 (2016) [arXiv:1509.00294 [hep-ex]].
- [39] G. Aad et al. [ATLAS Collaboration], JHEP 1512, 061 (2015) [arXiv:1509.06047 [hepex]].
- [40] V. Khachatryan et al. [CMS Collaboration], JHEP 1702, 079 (2017) [arXiv:1610.04857 [hep-ex]].
- [41] Y.-X. Lin and C. S. Nugroho, in preparation.
- [42] <http://www.feynarts.de/>
- [43] J. J. Zhang, C. S. Li, J. Gao, H. Zhang, Z. Li, C.-P. Yuan and T. C. Yuan, Phys. Rev. Lett. 102, 072001 (2009) [arXiv:0810.3889 [hep-ph]].
- [44] J. J. Zhang, C. S. Li, J. Gao, H. X. Zhu, C.-P. Yuan and T. C. Yuan, Phys. Rev. D 82, 073005 (2010) [arXiv:1004.0898 [hep-ph]].
- [45] M. Jezabek and J. H. Kuhn, Nucl. Phys. B **314**, 1 (1989).
- [46] C. S. Li, R. J. Oakes and T. C. Yuan, Phys. Rev. D 43, 3759 (1991).
- [47] J. Gao, C. S. Li and H. X. Zhu, Phys. Rev. Lett. 110, no. 4, 042001 (2013) [arXiv:1210.2808 [hep-ph]].