

A Distributed Algorithm for Least Square Solutions of Linear Equations

Xuan Wang, Jingqiu Zhou, Shaoshuai Mou and Martin J. Corless *

October 2, 2017

Abstract

A distributed discrete-time algorithm is proposed for multi-agent networks to achieve a common least squares solution of a group of linear equations, in which each agent only knows some of the equations and is only able to receive information from its nearby neighbors. For fixed, connected, and undirected networks, the proposed discrete-time algorithm results in each agents solution estimate to converging exponentially fast to the same least squares solution. Moreover, the convergence does not require careful choices of time-varying small step sizes.

1 Introduction

A significant amount of effort in the control community has recently been given to distributed algorithms for solving a set of linear equations over multi-agent networks, in which each agent only knows some of the equations and controls a state vector that can be looked upon as an estimate to the solution of the overall linear equations [1–4]. The key idea of these distributed algorithms is a so-called “agreement principle” [5], in which each agent limits the update of its state to satisfy its own equation while trying to reach a consensus with its nearby neighbors’ states. Different from the well-studied consensus problem [6–17], which aims to drive all agents’ states to be the same, the agreement principle allows agents to cooperatively reach the same solution to the overall set of linear equation as long solutions exist. Numerous extensions along this direction include achieving solutions with the minimum Euclidean norm [18, 19], elimination of the initialization step [20], and reduction of state vector dimension by utilizing the sparsity of the linear equations [21].

Linear equations arising from many engineering problems are, however, overdetermined, for which all the above distributed algorithms based on the agreement principle are not directly applicable. For example, in distributed parameter estimation [22], observations are subject to measurement noise that leads to no solution of the resulting equations; in power networks, the mode estimation of voltage oscillations asks for the least squares solution of linear equations resulted from the output of phasor measurement units [23, 24]; a distributed least squares solver can also be applied to the position determination of multi-agent formation control [25–27], state estimation in signal processing [28–30] and real-time data fitting of financial models [31]. One idea for dealing with the case of overdetermined linear equations is briefly discussed in [1], which however does not scale well with the network size. A common approach to achieve least squares solution is to reformulate it as a distributed optimization problem. In order to find the optimal solution in the sense of the total network, classical methods employ a centralized agent (coordinator) to collect the information in

*This work was supported by a funding from Northrop Grumman Corporation. X. Wang, J. Zhou, S. Mou and M. J. Corless are with the School of Aeronautics and Astronautics, Purdue University, West Lafayette, IN 47906 USA (e-mail: wang3156@purdue.edu, zhou745@purdue.edu, mous@purdue.edu, corless@purdue.edu). Corresponding Author: Shaoshuai Mou.

the network or assign computation tasks to other agents [28,32]. Such a structure, however, puts too much load on the central agent and has a strict requirement on the network topology. Compared with this, consensus based algorithms can solve distributed optimization problem with no requirement on a central agent [33–38]. For example, the methods based on the projection-consensus flow proposed in [29, 30, 39], are able to drive agents’ states to a neighborhood of the least square solution by introducing a sufficiently large gain. As an improvement of these methods, the exact least squares solution can be obtained by introducing a decaying weight to the local gradient [37, 38], but at the cost of losing fixed exponential convergence rate. Many other algorithms, like [34–37], have good results on both exact solution and convergence rate but require all agents to share a common, time-varying small step size that has to be carefully chosen for convergence. A similar requirement of sufficiently small step sizes has to be made when adapting classical continuous algorithms in [40–42] to achieve the least squares solution.

The major contribution of this paper comes from devising a discrete-time algorithm, which is distributed; achieves exact least square solutions; converges exponentially fast for fixed undirected connected networks; and does not involve any small or time-varying step sizes for convergence. These attributes differentiate the proposed algorithm from those in existence for achieving a least squares solution. The remainder of this paper is organized as follows. In Section II, we formulate the problem of obtaining a least squares solution in a distributed manner. Then, a discrete-time distributed algorithm is proposed in Section III and Section IV contains the main theorem which claims exponential convergence of the proposed algorithm to a least squares solution. A proof of the main theorem is contained in Section V, which is followed by numerical simulations in Section VI and conclusions in Section VII. Proofs of a lemma and a corollary are given in the Appendix.

Notation: The vector $\mathbf{1}_r$ is the vector in \mathbb{R}^r with all its components equal to 1 and I denotes an identity matrix. The transpose and kernel of any matrix M is denoted by M' , respectively. By $M > 0$ and $M \geq 0$ it is meant that the symmetric matrix M is positive definite and positive semi-definite, respectively. Finally,

$$\text{col} \{A_1, A_2, \dots, A_r\} = [A'_1 \quad A'_2 \quad \dots \quad A'_r]'$$

and $\text{diag} \{A_1, A_2, \dots, A_r\}$ is a block diagonal matrix with the i th diagonal block equal to A_i , $i = 1, 2, \dots, r$.

2 Problem Formulation

Consider a network of m agents, $i = 1, 2, \dots, m$, in which each agent can communicate with certain other agents called its *neighbors*. Suppose that each agent wishes to solve the following *least squares optimization problem*

$$\min_{x \in \mathbb{R}^n} \frac{1}{2} \sum_{i=1}^m \|A_i x - b_i\|_2^2 \quad (1)$$

for the *least squares solution* x^* , where $\|\cdot\|_2$ denotes the 2-norm, but *each agent i only knows matrices $A_i \in \mathbb{R}^{n_i \times n}$ and $b_i \in \mathbb{R}^{n_i}$.*

Suppose that at each time $t = 0, 1, \dots$, each agent i controls a state vector $x_i(t) \in \mathbb{R}^n$, which can be viewed as agent i 's estimate of x^* . The **problem** of interest in this paper is to develop a local rule for each agent i to iteratively update its state vector $x_i(t)$ by only using its neighbors' states such that all $x_i(t)$ converge exponentially fast to a least squares solution x^* .

Note that, if x^* is a least squares solution then, all least squares solutions are given by

$$x^* + r \quad (2)$$

with $A_i r = 0$ for $i = 1, \dots, m$, that is,

$$A r = 0 \quad (3)$$

where

$$A = \text{col} \{A_1, A_2, \dots, A_m\} \quad (4)$$

Thus the problem (1) has a unique solution if and only if $\ker(A) = 0$.

To proceed, we let \mathcal{N}_i denote the set of agent i 's neighbors. We assume that each agent is a neighbor of itself, that is, $i \in \mathcal{N}_i$. Neighbor relations can be described by an undirected graph \mathbb{G} with self-arcs such that there is an undirected edge connecting two different nodes i and j if and only if i and j are neighbors. In this paper we only consider the case in which \mathbb{G} is connected and fixed.

3 A Distributed Discrete-Time Update

In this section we present a distributed and discrete update algorithm for each agent to asymptotically achieve the same least squares solution x^* . We note that the problem (1) is equivalent to the following constrained optimization problem:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \sum_{i=1}^m |A_i x_i - b_i|_2^2 \\ & \text{subject to} && x_1 = x_2 = \dots = x_m \end{aligned} \quad (5)$$

To obtain an update algorithm, let W be a symmetric weighting matrix associated with the undirected graph \mathbb{G} such that its ij -th and ji -th entries, w_{ij} and w_{ji} , are positive if and only if there is an undirected edge between i and j in \mathbb{G} , and are zero, otherwise. Since each agent is a neighbor of itself one has $w_{ii} > 0$. Let D denote the diagonal matrix whose i th diagonal entry, denoted by d_i , is the i -th row sum of W , that is,

$$d_i = \sum_{j=1}^m w_{ij} = \sum_{j \in \mathcal{N}_i} w_{ij}. \quad (6)$$

Now introduce the Laplacian L matrix associated with the weighted graph:

$$L = D - W \quad (7)$$

let $\bar{L} = L \otimes I_n$ where \otimes denotes the Kronecker product and let

$$\mathbf{x} = \text{col} \{x_1, x_2, \dots, x_m\} \quad (8)$$

be the column consisting of all the state vectors. Since \mathbb{G} is connected, a vector is in the kernel of L if and only if it is a scalar multiple of $\mathbf{1}_m$ [43]. Using this property, one obtains that the constraint in (5) is equivalent to $\bar{L}\mathbf{x} = 0$. Thus problem (5) and, hence, the original problem, is equivalent to the following problem:

$$\begin{aligned} & \text{minimize} && \frac{1}{2} |\bar{A}\mathbf{x} - b|_2^2 \\ & \text{subject to} && \bar{L}\mathbf{x} = \mathbf{0} \end{aligned} \quad (9)$$

where

$$\bar{A} = \text{diag} \{A_1, A_2, \dots, A_m\} \quad (10)$$

$$b = \text{col} \{b_1, b_2, \dots, b_m\} \quad (11)$$

Note that \mathbf{x}^* solves (9) if and only if

$$\mathbf{x}^* = \mathbf{1}_m \otimes x^* \quad (12)$$

where x^* is a least squares solution to the original problem.

The linear constraint quadratic optimization problem (9) is analytically solvable by Lagrange Method [42,44]. That is, define

$$G(\mathbf{x}, \mathbf{z}) = \frac{\bar{c}}{2} (\bar{A}\mathbf{x} - b)' (\bar{A}\mathbf{x} - b) + \mathbf{z}' \bar{L}\mathbf{x} \quad (13)$$

where $\mathbf{z} = \text{col} \{z_1, z_2, \dots, z_m\} \in \mathbb{R}^{nm}$ is the so-called Lagrange multiplier and $\bar{c} > 0$ is an arbitrary positive constant introduced for the purpose of adjusting the weights between the two terms summed in $G(\mathbf{x}, \mathbf{z})$. Note that the Hessian matrix of the objective function is $\bar{A}'\bar{A} \geq 0$. Then, \mathbf{x}^* solves problem (9) if and only if there exists \mathbf{z}^* such that $\nabla_{\mathbf{x}, \mathbf{z}} G(\mathbf{x}^*, \mathbf{z}^*) = 0$. Then the problem of achieving a least square solution x^* to (1) is equivalent to finding \mathbf{x}^* and \mathbf{z}^* such that

$$\bar{c}(\bar{A}'\bar{A}\mathbf{x}^* - \bar{A}'b) + \bar{L}'\mathbf{z}^* = 0 \quad (14)$$

$$\bar{L}\mathbf{x}^* = 0 \quad (15)$$

Since W is symmetric, $L = L'$; hence (14)-(15) are equivalent to

$$\bar{c}(A'_i A_i x_i^* - A'_i b_i) + \sum_{j \in \mathcal{N}_i} w_{ij} (z_i^* - z_j^*) = 0 \quad (16)$$

$$\sum_{j \in \mathcal{N}_i} w_{ij} (x_i^* - x_j^*) = 0 \quad (17)$$

for $i = 1, 2, \dots, m$.

By introducing an additional state vector $z_i(t) \in \mathbb{R}^n$ for each agent i , one could achieve a distributed solution to (16) and (17) by the saddle-point dynamics proposed in [42,45]. Discretization of such a continuous update usually requires a careful choice of sufficiently small step size to guarantee convergence. To eliminate such a requirement, we propose a new discrete-time update as follows:

$$\begin{aligned} x_i(t+1) &= x_i(t) - \bar{c}\kappa_i [A'_i A_i x_i(t+1) - A'_i b_i] \\ &\quad - \kappa_i \sum_{j \in \mathcal{N}_i} w_{ij} [z_i(t+1) - z_j(t)] \\ &\quad - c\kappa_i \sum_{j \in \mathcal{N}_i} w_{ij} [x_i(t+1) - x_j(t)] \end{aligned} \quad (18)$$

$$z_i(t+1) = z_i(t) + \kappa_i \sum_{j \in \mathcal{N}_i} w_{ij} [x_i(t+1) - x_j(t)] \quad (19)$$

Here $c \geq 0$ is arbitrary non-negative constant and $\kappa_i > 0$, $i = 1, \dots, m$, are parameters to be chosen. As will be shown later, a simple and distributed way of choosing c, \bar{c}, κ_i for each agent is

$$c \geq 0, \quad \bar{c} > 0, \quad \kappa_i = \frac{1}{d_i}$$

Under this choice the updates (18)-(19) will be totally distributed without any designed parameters and the effectiveness for driving all $x_i(t)$ to a least square solution will be shown later in next section.

The updates (18)-(19) result from a mixed use of each agent's upcoming states $x_i(t+1), z_i(t+1)$ and its neighbors current states $x_j(t), z_j(t), j \in \mathcal{N}_i$. This enables us to derive from (18) and (19) the following update without introducing any step size:

$$\begin{bmatrix} x_i(t+1) \\ z_i(t+1) \end{bmatrix} = E_i \begin{bmatrix} x_i(t) + \kappa_i \sum_{j \in \mathcal{N}_i} w_{ij} [cx_j(t) + z_j(t)] + \bar{c}\kappa_i A'_i b_i \\ -\kappa_i \sum_{j \in \mathcal{N}_i} w_{ij} x_j(t) + z_i(t) \end{bmatrix} \quad (20)$$

where

$$E_i = \begin{bmatrix} I_n + \bar{c}\kappa_i A_i' A_i + c\kappa_i d_i I_n & \kappa_i d_i I_n \\ -\kappa_i d_i I_n & I_n \end{bmatrix}^{-1}$$

Note right away that the update (20) is distributed since each agent i only uses A_i, b_i and states of its neighbors and itself; it requires each agent to control a state vector in \mathbb{R}^{2n} whose size is independent of the underlying network, and does not involve any step size. Exponential convergence under the proposed update will be expounded on in next section.

4 Main Result

To present the main result of the paper, Theorem 1, let

$$\bar{W} = W \otimes I_n, \quad \bar{D} = D \otimes I_n, \quad \bar{\mathcal{K}} = \mathcal{K} \otimes I_n$$

where $\mathcal{K} \in \mathbb{R}^{m \times m}$ is the diagonal matrix whose ii entry is κ_i .

Theorem 1 *Suppose \mathbb{G} is undirected and connected, W is symmetric, $\bar{c}, \kappa_1, \dots, \kappa_m > 0, c \geq 0$*

$$DKD - WKW \geq 0 \tag{21}$$

and $c > 0$ if there exists a non-zero vector \mathbf{u} such that

$$\bar{A}\mathbf{u} = 0 \tag{22}$$

$$(\bar{D}\bar{\mathcal{K}}\bar{D} - \bar{W}\bar{\mathcal{K}}\bar{W})\mathbf{u} = 0 \tag{23}$$

$$\bar{L}\mathbf{u} \neq 0. \tag{24}$$

Then the proposed update (20) results in all $x_i(t)$ converging exponentially fast to the same least squares solution to $Ax = b$.

By Theorem 1, convergence of the proposed update depends on choosing parameters κ_i to satisfy (21). This can be achieved in a simple and **distributed** way as illustrated by the following corollary.

Corollary 1 *If $c \geq 0, \bar{c} > 0$ and $\kappa_i = 1/d_i$ for $i = 1, 2, \dots, m$ then the proposed update (20) results in all $x_i(t)$ converging exponentially fast to the same least squares solution to $Ax = b$.*

A proof of Corollary 1 is given in the Appendix.

5 Proof of Theorem 1

In order to prove our main result, we first re-write the update equations (18)-(19) in vector form:

$$\begin{aligned} \mathbf{x}(t+1) &= \mathbf{x}(t) - \bar{c}\bar{\mathcal{K}} [\bar{A}'\bar{A}\mathbf{x}(t+1) - \bar{A}'b] \\ &\quad - \bar{\mathcal{K}} [\bar{D}\mathbf{z}(t+1) - \bar{W}\mathbf{z}(t)] \\ &\quad - c\bar{\mathcal{K}} [\bar{D}\mathbf{x}(t+1) - \bar{W}\mathbf{x}(t)] \end{aligned} \tag{25}$$

$$\mathbf{z}(t+1) = \mathbf{z}(t) + \bar{\mathcal{K}} [\bar{D}\mathbf{x}(t+1) - \bar{W}\mathbf{x}(t)] \tag{26}$$

where

$$\mathbf{x} = \text{col} \{x_1, \dots, x_m\} \quad (27)$$

$$\mathbf{z} = \text{col} \{z_1, \dots, z_m\} \quad (28)$$

Since $\bar{\mathcal{K}}$ is invertible and $\bar{c} > 0$, all equilibrium states $\text{col} \{\mathbf{x}^e, \mathbf{z}^e\}$ of (25)-(26) are given by $\bar{c}[\bar{A}'\bar{A}\mathbf{x}^e - \bar{A}'b] + \bar{L}\mathbf{z}^e + c\bar{L}\mathbf{x}^e = 0$ and $\bar{L}\mathbf{x}^e = 0$ where

$$\bar{L} = \bar{D} - \bar{W} = L \otimes I \quad (29)$$

Thus, the equilibrium states are given by

$$\bar{c}(\bar{A}'\bar{A}\mathbf{x}^e - \bar{A}'b) + \bar{L}\mathbf{z}^e = 0 \quad (30)$$

$$\bar{L}\mathbf{x}^e = 0 \quad (31)$$

Clearly the set an equilibrium states to (25)-(26) is the same as the set of solutions to (14)-(15). So, to prove Theorem 1 we just need to show that every solution to (25)-(26) converges to an equilibrium state. To achieve this we re-write (25)-(26) compactly as

$$\mathbf{y}(t+1) = Q\mathbf{y}(t) + \mathbf{b} \quad (32)$$

where $\mathbf{y} = \text{col} \{\mathbf{x}, \mathbf{z}\}$ and

$$Q = \begin{bmatrix} I_{mn} + \bar{c}\bar{\mathcal{K}}\bar{A}'\bar{A} + c\bar{\mathcal{K}}\bar{D} & \bar{\mathcal{K}}\bar{D} \\ -\bar{\mathcal{K}}\bar{D} & I_{mn} \end{bmatrix}^{-1} \begin{bmatrix} I_{mn} + c\bar{\mathcal{K}}\bar{W} & \bar{\mathcal{K}}\bar{W} \\ -\bar{\mathcal{K}}\bar{W} & I_{mn} \end{bmatrix} \quad (33)$$

$$\mathbf{b} = \begin{bmatrix} I_{mn} + \bar{c}\bar{\mathcal{K}}\bar{A}'\bar{A} + c\bar{\mathcal{K}}\bar{D} & \bar{\mathcal{K}}\bar{D} \\ -\bar{\mathcal{K}}\bar{D} & I_{mn} \end{bmatrix}^{-1} \begin{bmatrix} \bar{c}\bar{\mathcal{K}}\bar{A}'b \\ 0 \end{bmatrix} \quad (34)$$

The equilibrium states \mathbf{y}^e of (32) are given by $(I - Q)\mathbf{y}^e = \mathbf{b}$. Let \mathbf{y}^* be any equilibrium state of (32). Then all equilibrium states \mathbf{y}^e of (32) are given by $\mathbf{y}^e = \mathbf{y}^* + \mathbf{v}$ where $\mathbf{v} = 0$ or \mathbf{v} an eigenvector of Q corresponding to eigenvalue one. The evolution of $\mathbf{e} = \mathbf{y} - \mathbf{y}^*$ is governed by

$$\mathbf{e}(t+1) = Q\mathbf{e}(t) \quad (35)$$

So, to prove Theorem 1 we now just have to show that every solution of (35) exponentially converges to zero or to an eigenvector of Q corresponding to eigenvalue one. To achieve this we need the following lemma whose proof is in the Appendix

Lemma 1 *Suppose (21) holds. Then Q has the following properties.*

- (a) *Every eigenvalue of Q has magnitude less than or equal to one and -1 is not an eigenvalue of Q .*
- (b) *If Q has a complex eigenvalue of magnitude one then $c = 0$ and there is a non-zero vector \mathbf{u} which satisfies (22) - (24) in Theorem 1.*
- (c) *One is an eigenvalue of Q and its algebraic multiplicity is equal to its geometric multiplicity. A non-zero vector $\text{col} \{\mathbf{u}, \bar{\mathbf{u}}\}$ is a eigenvector corresponding to one if and only if*

$$\bar{A}\mathbf{u} = 0, \quad \bar{L}\mathbf{u} = 0, \quad \bar{L}\bar{\mathbf{u}} = 0 \quad (36)$$

As a consequence of the hypotheses of Theorem 1, Lemma 1 tells us that every eigenvalue of Q has magnitude less than or equal to one. Also, one is the only eigenvalue of magnitude one and its

algebraic and geometric multiplicities are equal. Hence, there exists a non-singular matrix T such that

$$Q = T \begin{bmatrix} I & 0 \\ 0 & R \end{bmatrix} T^{-1} \quad (37)$$

and all the eigenvalues of R have magnitude strictly less than one. Every solution of (35) satisfies $\mathbf{e}(t) = Q^t \mathbf{e}(0)$. Since

$$Q^t = T \begin{bmatrix} I & 0 \\ 0 & R^t \end{bmatrix} T^{-1}$$

and all the eigenvalues of R have magnitude strictly less than one, it follows that $\mathbf{e}(t)$ exponentially converges to

$$\mathbf{v} = T \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} T^{-1} \mathbf{e}(0)$$

Note that

$$\begin{aligned} Q\mathbf{v} &= T \begin{bmatrix} I & 0 \\ 0 & R \end{bmatrix} T^{-1} T \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} T^{-1} \mathbf{e}(0) \\ &= T \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} T^{-1} \mathbf{e}(0) = \mathbf{v} \end{aligned}$$

that is, $\mathbf{v} = 0$ or \mathbf{v} is an eigenvector of Q corresponding to eigenvalue one. Hence every solution of (35) exponentially converges to zero or to an eigenvector of Q corresponding to eigenvalue one. ■

6 Example

Numerical simulations will be performed with the 5-node network in Fig. 1 to illustrate Theorem 1.




Figure 1: Five-node connected network \mathbb{G}

From Fig. 1 we obtain that

$$W = \begin{bmatrix} 0.9 & 1.5 & 0 & 0.6 & 0 \\ 1.5 & 0.7 & 1.8 & 0 & 0 \\ 0 & 1.8 & 1 & 2.2 & 0 \\ 0.6 & 0 & 2.2 & 0.8 & 1.4 \\ 0 & 0 & 0 & 1.4 & 0.6 \end{bmatrix}$$

Hence

$$D = \text{diag} \{3, 4, 5, 5, 2\}$$

and based on the distributed way of choosing weights in Corollary 1,

$$\mathcal{K} = \text{diag} \left\{ \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{5}, \frac{1}{2} \right\}$$

We consider solving a set of linear equations that has multiple least square solutions on network \mathbb{G} , in which agents 1, 2, 3, 4, 5 know

$$\begin{aligned} A_1 &= [1 \ 2 \ 3 \ 4], & b_1 &= 10 \\ A_2 &= [4 \ 5 \ 6 \ 7], & b_2 &= 20 \\ A_3 &= [1 \ 2 \ 3 \ 4], & b_3 &= 15 \\ A_4 &= [5 \ 6 \ 3 \ 4], & b_4 &= 17 \\ A_5 &= [4 \ 3 \ 2 \ 1], & b_5 &= 6 \end{aligned}$$

respectively. Since the least squares solution is not unique, to show the effectiveness of our method, we introduce $W(t)$ as

$$W(t) = \frac{1}{2m} \sum_{i=1}^m |A'Ax_i(t) - A'b|_2^2 + \frac{1}{2m^2} \sum_{i=1}^m \sum_{j=1}^m |x_i(t) - x_j(t)|_2^2$$

where the first term is a cost associated with the $x_i(t)$ not being a least squares solution of $Ax = b$; the second term is a cost associated with the $x_i(t)$ not achieving consensus to the same value. Numerical simulation results in Fig. 2 validates the exponential convergence of $W(t)$ for $\bar{c} = 1$ and $c = 0, 2, 4$, respectively. As a comparison, Fig. 3 validates the exponential convergence of $W(t)$




Figure 2: Example: multiple least squares solution, $\bar{c} = 1$, different c

for $c = 0$ and $\bar{c} = 0.1, 0.3, 1, 3$, respectively.

It is worth mentioning that $W(t)$ goes to 0 doesn't mean x_i converge to a constant value, to show this, we let $c = 0, \bar{c} = 1$ and use Fig. 4 to demonstrate the history of x_1 . This, along with the consensus result in Fig. 2 and 3 validates that all x_i converge to constant values.

The simulation results show that different parameters c and \bar{c} lead to different convergence rates, this because the eigenvalues of matrix Q are different. However, finding the best parameter set c and \bar{c} is not straightforward because the eigenvalues of matrix Q are also determined by the equation $Ax = b$ and the network \mathbb{G} , both information are global information that cannot be obtained by agents.




Figure 3: Example: Multiple least squares solution, $c = 0$, different \bar{c}




Figure 4: History of x_1 when $c = 0$ and $\bar{c} = 1$

7 Conclusion

We have proposed a discrete-time update for multi-agent networks, which enable each agent to achieve the same least square solutions exponentially fast when the network is undirected and connected. The exponential convergence does not rely on any time-varying and small step-size, which differs from the proposed updates in existence. Future work includes proper design of parameters c , \bar{c} and the generalization of the proposed update to networks that are directed and time-varying.

Appendix

Proof of Lemma 1

First note that due to the mixed-product property of Kronecker product, $\bar{D}\bar{K}\bar{D} = (DKD) \otimes I_n$; $\bar{W}\bar{K}\bar{W} = (WKW) \otimes I_n$. Assumption (21) in Theorem 1 is equivalent to

$$\bar{D}\bar{K}\bar{D} - \bar{W}\bar{K}\bar{W} \geq 0$$

If suppose λ is an eigenvalue of Q . Then there is a nonzero vector $\text{col}\{\mathbf{u}, \bar{\mathbf{u}}\}$ such that

$$Q \begin{bmatrix} \mathbf{u} \\ \bar{\mathbf{u}} \end{bmatrix} = \lambda \begin{bmatrix} \mathbf{u} \\ \bar{\mathbf{u}} \end{bmatrix}$$

which, recalling (33), is equivalent to

$$\begin{bmatrix} I_{mn} + c\bar{\mathcal{K}}\bar{W} & \bar{\mathcal{K}}\bar{W} \\ -\bar{\mathcal{K}}\bar{W} & I_{mn} \end{bmatrix} \begin{bmatrix} \mathbf{u} \\ \bar{\mathbf{u}} \end{bmatrix} = \lambda \begin{bmatrix} I_{mn} + \bar{c}\bar{\mathcal{K}}\bar{A}'\bar{A} + c\bar{\mathcal{K}}\bar{D} & \bar{\mathcal{K}}\bar{D} \\ -\bar{\mathcal{K}}\bar{D} & I_{mn} \end{bmatrix} \begin{bmatrix} \mathbf{u} \\ \bar{\mathbf{u}} \end{bmatrix}$$

that is,

$$\begin{aligned} [I_{mn} + c\bar{\mathcal{K}}\bar{W} - \lambda(I_{mn} + \bar{c}\bar{\mathcal{K}}\bar{A}'\bar{A} + c\bar{\mathcal{K}}\bar{D})] \mathbf{u} &= \bar{\mathcal{K}}(\lambda\bar{D} - \bar{W}) \bar{\mathbf{u}} \\ \bar{\mathcal{K}}(\lambda\bar{D} - \bar{W}) \mathbf{u} &= (\lambda - 1) \bar{\mathbf{u}} \end{aligned}$$

and, since $\bar{\mathcal{K}}$ is nonsingular, these can be written as

$$[\bar{\mathcal{K}}^{-1} + c\bar{W} - \lambda(\bar{\mathcal{K}}^{-1} + \bar{c}\bar{A}'\bar{A} + c\bar{D})] \mathbf{u} = (\lambda\bar{D} - \bar{W}) \bar{\mathbf{u}} \quad (38)$$

$$(\lambda\bar{D} - \bar{W}) \mathbf{u} = (\lambda - 1) \bar{\mathcal{K}}^{-1} \bar{\mathbf{u}} \quad (39)$$

In the case of $\lambda \neq 1$, (39) is equivalent to

$$\bar{\mathbf{u}} = \frac{1}{\lambda - 1} \bar{\mathcal{K}}(\lambda\bar{L}) \mathbf{u} \quad (40)$$

Hence $\mathbf{u} \neq 0$ and (38) is equivalent to

$$M(\lambda) \mathbf{u} = 0 \quad (41)$$

where

$$\begin{aligned} M(\lambda) &= (\lambda\bar{D} - \bar{W}) \bar{\mathcal{K}}(\lambda\bar{D} - \bar{W}) + \lambda(\lambda - 1) \bar{c}\bar{A}'\bar{A} \\ &\quad + (\lambda - 1)^2 \bar{\mathcal{K}}^{-1} + (\lambda - 1)c(\lambda\bar{D} - \bar{W}) \end{aligned} \quad (42)$$

Note that

$$M(\lambda) = \lambda^2 M_2 + \lambda M_1 + M_0$$

where

$$\begin{aligned} M_2 &= \bar{D}\bar{\mathcal{K}}\bar{D} + \bar{\mathcal{K}}^{-1} + c\bar{D} + \bar{c}\bar{A}'\bar{A} > 0 \\ M_1 &= -\bar{W}\bar{\mathcal{K}}\bar{D} - \bar{D}\bar{\mathcal{K}}\bar{W} - 2\bar{\mathcal{K}}^{-1} - c\bar{D} - c\bar{W} - \bar{c}\bar{A}'\bar{A} \\ M_0 &= \bar{W}\bar{\mathcal{K}}\bar{W} + \bar{\mathcal{K}}^{-1} + c\bar{W} > 0 \end{aligned}$$

Thus, we have shown that $\lambda \neq 1$ is an eigenvalue of Q if and only if there is a nonzero vector \mathbf{u} such that (41) holds.

In the case of $\lambda = 1$, equations (38) and (39) reduce to

$$\begin{aligned} -[\bar{c}\bar{A}'\bar{A} + c(\bar{D} - \bar{W})] \mathbf{u} &= (\bar{D} - \bar{W}) \bar{\mathbf{u}} \\ (\bar{D} - \bar{W}) \mathbf{u} &= 0 \end{aligned}$$

Recall that $\bar{D} - \bar{W} = \bar{L}$, thus $\lambda = 1$ is an eigenvalue of Q and $\text{col}\{\mathbf{u}, \bar{\mathbf{u}}\}$ is an eigenvector corresponding to 1 if and only if $\text{col}\{\mathbf{u}, \bar{\mathbf{u}}\}$ is nonzero and

$$\bar{L}\mathbf{u} = 0 \quad (43)$$

$$\bar{L}\bar{\mathbf{u}} = -\bar{c}\bar{A}'\bar{A}\mathbf{u} \quad (44)$$

Proof of (a)

If $|\lambda| > 1$, then we have shown in the previous section that λ is an eigenvalue of Q if and only if there is a nonzero vector \mathbf{u} such that (41) holds.

Suppose λ is real with $|\lambda| > 1$ and recall expression (42) for $M(\lambda)$. Observe that both $\lambda(\lambda - 1)$ and $(\lambda - 1)^2$ are positive; $\bar{A}'\bar{A}$ and $(\lambda\bar{D} - \bar{W})\bar{K}(\lambda\bar{D} - \bar{W})$ are positive semi-definite; and \bar{K}^{-1} is positive definite. Furthermore, if $\lambda > 1$ then, $\lambda - 1$ is positive and $c(\lambda\bar{D} - \bar{W})$ is positive semi-definite; if $\lambda < -1$ then, $\lambda - 1$ is negative and $c(\lambda\bar{D} - \bar{W})$ is negative semi-definite. Thus, we conclude that when λ is real with $|\lambda| > 1$, the matrix $M(\lambda)$ in (42) is positive definite. Hence there is not a non-zero vector \mathbf{u} for which $M(\lambda)\mathbf{u} = 0$ and so λ is not an eigenvalue of Q .

Suppose that λ is complex. Left-multiplying equation (41) by \mathbf{u}' yields

$$\lambda^2 c_2 + \lambda c_1 + c_0 = 0$$

where

$$c_0 = \mathbf{u}'M_0\mathbf{u}, \quad c_1 = \mathbf{u}'M_1\mathbf{u}, \quad c_2 = \mathbf{u}'M_2\mathbf{u} \quad (45)$$

Since M_0, M_1, M_2 are symmetric, c_0, c_1 and c_2 are real. Let $\lambda = p + qi$, where $i = \sqrt{-1}$ and p and q are real with $q \neq 0$. Then equating the real and imaginary parts of (45) to zero results in

$$(p^2 - q^2)c_2 + pc_1 + c_0 = 0 \quad (46)$$

$$q(2pc_2 + c_1) = 0 \quad (47)$$

Since $q \neq 0$, equation (47) implies that

$$2pc_2 + c_1 = 0 \quad (48)$$

which upon substitution into (46) yields

$$|\lambda|^2 c_2 - c_0 = 0 \quad (49)$$

Since M_2 is positive definite and $\mathbf{u} \neq 0$, we must have $c_2 > 0$. Note that $c_2 - c_0 = \mathbf{u}'(M_2 - M_0)\mathbf{u}$ and

$$M_2 - M_0 = \bar{D}\bar{K}\bar{D} - \bar{W}\bar{K}\bar{W} + c(\bar{D} - \bar{W}) + \bar{c}\bar{A}'\bar{A} \quad (50)$$

Since $\bar{D} - \bar{W} = \bar{L} \geq 0$, $\bar{A}'\bar{A} \geq 0$, $c, \bar{c} \geq 0$ and, by assumption $\bar{D}\bar{K}\bar{D} - \bar{W}\bar{K}\bar{W} \geq 0$, one has $M_2 - M_0 \geq 0$ and $c_2 \geq c_0$. Since $c_2 > 0$, it now follows from (49) that $|\lambda|^2 = c_0/c_2 \leq 1$. Hence $|\lambda| \leq 1$.

Recall that $\lambda = -1$ is an eigenvalue of Q if and only if there is a nonzero vector \mathbf{u} such that $M(-1)\mathbf{u} = 0$. From (42) we have

$$\begin{aligned} M(-1) &= (\bar{D} + \bar{W})\bar{K}(\bar{D} + \bar{W}) + 2\bar{c}\bar{A}'\bar{A} + 4\bar{K}^{-1} + 2c(\bar{D} + \bar{W}) \\ &> 0 \end{aligned}$$

Since $M(-1)$ is positive definite, $M(-1)\mathbf{u} \neq 0$ for all non-zero $\mathbf{u} = 0$. Thus -1 is not an eigenvalue of Q .

Proof of (b)

Suppose that λ is a complex eigenvalue of Q with $|\lambda| = 1$. Recalling the proof of (a), there must exist a nonzero vector \mathbf{u} such that (48) and (49) hold, that is,

$$\mathbf{u}'(2pM_2 + M_1)\mathbf{u} = 0 \quad (51)$$

$$\mathbf{u}'(M_2 - M_0)\mathbf{u} = 0 \quad (52)$$

where $\lambda = p + qi$. Recall (50). Since $c(\bar{D} - \bar{W})$ and $\bar{c}\bar{A}'\bar{A}$ are positive semi-definite, $\bar{c} > 0$ and, by assumption, $\bar{D}\bar{K}\bar{D} - \bar{W}\bar{K}\bar{W}$ is positive semi-definite, (52) implies that

$$(\bar{D}\bar{K}\bar{D} - \bar{W}\bar{K}\bar{W})\mathbf{u} = 0 \quad (53)$$

$$\bar{A}\mathbf{u} = 0 \quad (54)$$

$$c(\bar{D} - \bar{W})\mathbf{u} = 0 \quad (55)$$

Equation (51), along with (54), results in

$$\begin{aligned} & \mathbf{u}'[2p(\bar{D}\bar{K}\bar{D} + \bar{K}^{-1} + c\bar{D}) \\ & - (\bar{W}\bar{K}\bar{D} + \bar{D}\bar{K}\bar{W} + 2\bar{K}^{-1} + c\bar{D} + c\bar{W})]\mathbf{u} = 0 \end{aligned} \quad (56)$$

If $c \neq 0$ then (55) implies that $\bar{D}\mathbf{u} = \bar{W}\mathbf{u}$ and we obtain that

$$2(p-1)\mathbf{u}'[\bar{D}\bar{K}\bar{D} + \bar{K}^{-1} + c\bar{D}]\mathbf{u} = 0 \quad (57)$$

Since $1 = |\lambda|^2 = p^2 + q^2 = 1$ and $q \neq 0$, we must have $p < 1$ and (57) implies that

$$\mathbf{u}'[\bar{D}\bar{K}\bar{D} + \bar{K}^{-1} + c\bar{D}]\mathbf{u} = 0 \quad (58)$$

The matrix $\bar{D}\bar{K}\bar{D} + \bar{K}^{-1} + c\bar{D}$ is positive definite, so, (58) yields the contradiction that $\mathbf{u} = 0$; hence $c = 0$.

If $\bar{L}\mathbf{u} = 0$ then $\bar{D}\mathbf{u} = \bar{W}\mathbf{u}$ and (57) holds. Again we get the contradiction that $\mathbf{u} = 0$. Hence $\bar{L}\mathbf{u} \neq 0$. This along with (53), (54), lead to the equations (22)-(24) in Theorem 1. ■

Proof of (c)

We have seen that one is an eigenvalue for Q and $\text{col}\{\mathbf{u}, \bar{\mathbf{u}}\}$ is a corresponding eigenvector if and only if $\text{col}\{\mathbf{u}, \bar{\mathbf{u}}\}$ is nonzero and satisfies (43) and (44). Since \bar{L} is symmetric, (43) implies that $\mathbf{u}'\bar{L} = 0$. Multiplying both sides of (44) by \mathbf{u}' , it now follows that $-\bar{c}\mathbf{u}'\bar{A}'\bar{A}\mathbf{u} = 0$; since $\bar{c} > 0$, this is equivalent to $\bar{A}\mathbf{u} = 0$. Equation (43) now implies that $\bar{L}\bar{\mathbf{u}} = 0$. Thus (36) holds. In addition, since \bar{L} is singular, a nonzero solution $\text{col}\{\mathbf{u}, \bar{\mathbf{u}}\}$ to (36) exists; hence one is an eigenvalue for Q .

Now, we prove the multiplicity property of eigenvalue one by contradiction. Suppose the algebraic multiplicity of the eigenvalue one is not equal to its geometric multiplicity. Then there exists a nonzero vector $\text{col}\{\mathbf{v}, \bar{\mathbf{v}}\}$ [46] such that

$$(Q - I) \begin{bmatrix} \mathbf{v} \\ \bar{\mathbf{v}} \end{bmatrix} = \begin{bmatrix} \mathbf{u} \\ \bar{\mathbf{u}} \end{bmatrix} \quad (59)$$

where $\text{col}\{\mathbf{u}, \bar{\mathbf{u}}\}$ is an eigenvector corresponding to one. It follows from (59) and the definition of Q in (33) that

$$\begin{aligned} & (c\bar{K}\bar{W} - c\bar{K}\bar{D} - \bar{c}\bar{K}\bar{A}'\bar{A})\mathbf{v} - (\bar{K}\bar{W} - \bar{K}\bar{D})\bar{\mathbf{v}} \\ & = (I + \bar{c}\bar{K}\bar{A}'\bar{A} + c\bar{K}\bar{D})\mathbf{u} + \bar{K}\bar{D}\bar{\mathbf{u}} \\ & (\bar{K}\bar{D} - \bar{K}\bar{W})\mathbf{v} = -\bar{K}\bar{D}\mathbf{u} + \bar{\mathbf{u}} \end{aligned}$$

Left multiplying both equations by \bar{K}^{-1} and recalling that $\bar{L} = \bar{D} - \bar{W}$ yields

$$-c(\bar{L} + \bar{A}'\bar{A})\mathbf{v} + \bar{L}\bar{\mathbf{v}} = (\bar{K}^{-1} + \bar{c}\bar{A}'\bar{A} + c\bar{D})\mathbf{u} + \bar{D}\bar{\mathbf{u}} \quad (60)$$

$$\bar{L}\mathbf{v} = -\bar{D}\mathbf{u} + \bar{K}^{-1}\bar{\mathbf{u}} \quad (61)$$

Since $\text{col}\{\mathbf{u}, \bar{\mathbf{u}}\}$ is an eigenvector corresponding to one, it is nonzero and satisfies (36); hence left multiplying (60) and (61) by \mathbf{u}' and $\bar{\mathbf{u}}'$, respectively, results in

$$\begin{aligned} 0 &= \mathbf{u}' (\bar{\mathcal{K}}^{-1} + c\bar{D}) \mathbf{u} + \mathbf{u}' \bar{D} \bar{\mathbf{u}} \\ 0 &= -\bar{\mathbf{u}}' \bar{D} \mathbf{u} + \bar{\mathbf{u}}' \bar{\mathcal{K}}^{-1} \bar{\mathbf{u}} \end{aligned}$$

Thus

$$\mathbf{u}' (\bar{\mathcal{K}}^{-1} + c\bar{D}) \mathbf{u} + \bar{\mathbf{u}}' \bar{\mathcal{K}}^{-1} \bar{\mathbf{u}} = 0 \quad (62)$$

Since both $(\bar{\mathcal{K}}^{-1} + c\bar{D})$ and $\bar{\mathcal{K}}^{-1}$ are positive definite, we obtain the contradiction that $\mathbf{u} = \bar{\mathbf{u}} = 0$. Hence the algebraic multiplicity of eigenvalue one must equal its geometric multiplicity. ■

Proof of Corollary 1

We prove this corollary by showing that the hypotheses of Theorem 1 hold. Since $\kappa_i = 1/d_i$ we have $\bar{\mathcal{K}} = \bar{D}^{-1}$ and

$$\begin{aligned} \bar{D}\bar{\mathcal{K}}\bar{D} - \bar{W}\bar{\mathcal{K}}\bar{W} &= \bar{D} - \bar{W}\bar{D}^{-1}\bar{W} \\ &= \bar{D}^{\frac{1}{2}}[I - (\bar{D}^{-\frac{1}{2}}\bar{W}\bar{D}^{-\frac{1}{2}})^2]\bar{D}^{\frac{1}{2}} \end{aligned}$$

By the Gershgorin Disk Theorem, $\bar{L} = \bar{D} - \bar{W} \geq 0$ [43] and thus $I \geq \bar{D}^{-\frac{1}{2}}\bar{W}\bar{D}^{-\frac{1}{2}}$. Hence $I \geq (\bar{D}^{-\frac{1}{2}}\bar{W}\bar{D}^{-\frac{1}{2}})^2$ that is,

$$I - (\bar{D}^{-\frac{1}{2}}\bar{W}\bar{D}^{-\frac{1}{2}})^2 \geq 0$$

Thus, $\bar{D}\bar{\mathcal{K}}\bar{D} - \bar{W}\bar{\mathcal{K}}\bar{W} \geq 0$, that is, (21) holds.

We now show that equations (22)-(24) do not have a solution. If (23) holds then,

$$(\bar{D} - \bar{W}\bar{D}^{-1}\bar{W}) \mathbf{u} = 0$$

that is

$$\bar{D}^{\frac{1}{2}} \left(I + \bar{D}^{-\frac{1}{2}}\bar{W}\bar{D}^{-\frac{1}{2}} \right) \left(I - \bar{D}^{-\frac{1}{2}}\bar{W}\bar{D}^{-\frac{1}{2}} \right) \bar{D}^{\frac{1}{2}} \mathbf{u} = 0 \quad (63)$$

Recall that \mathbb{G} has self-arcs, that is, $\bar{D} + \bar{W} > 0$ [43]; hence $I_{mn} + \bar{D}^{-\frac{1}{2}}\bar{W}\bar{D}^{-\frac{1}{2}} > 0$ is nonsingular. Thus, equation (63) leads to

$$\left(I_{mn} - \bar{D}^{-\frac{1}{2}}\bar{W}\bar{D}^{-\frac{1}{2}} \right) \bar{D}^{\frac{1}{2}} \mathbf{u} = 0 \quad (64)$$

which, upon left-multiplying by $\bar{D}^{\frac{1}{2}}$ yields

$$(\bar{D} - \bar{W}) \mathbf{u} = 0 \quad (65)$$

that is (24) is not satisfied. Hence there does not exist a vector \mathbf{u} satisfying (22) - (24). Application of Theorem 1 now yields exponential convergence for any $c \geq 0$ and $\bar{c} > 0$. ■

References

- [1] S. Mou, J. Liu, and A. S. Morse, "A distributed algorithm for solving a linear algebraic equation," *IEEE Transactions on Automatic Control*, no. 11, pp. 2863–2878, 2015.

- [2] B. D. O. Anderson, S. Mou, U. R. Helmke, and A. S. Morse, “Decentralized gradient algorithm for solution of a linear equation,” *Numerical Algebra, Control and Optimization*, no. 3, pp. 319–328, 2016.
- [3] J. Lu and C. Y. Tang, “A distributed algorithm for solving positive definite linear equations over networks with membership dynamics,” *IEEE Transactions on Control of Network Systems*, 2016.
- [4] J. Wang and N. Elia, “Distributed solution of linear equations over unreliable networks,” *Proceedings of American Control Conference*, pp. 6471–6476, Jul. 2016.
- [5] S. Mou and A. S. Morse, “A fixed-neighbor, distributed algorithm for solving a linear algebraic equation,” *European Control Conference*, pp. 2269–2273, 2013.
- [6] A. Jadbabaie, J. Lin, and A. S. Morse, “Coordination of groups of mobile autonomous agents using nearest neighbor rules,” *IEEE Trans. Autom. Control*, pp. 988–1001, June 2003, also in Proc. 41st IEEE CDC, pages 2953 - 2958, 2002.
- [7] Z. Lin, B. Francis, and M. Maggiore, “State agreement for continuous-time coupled nonlinear systems,” *SIAM Journal on Control and Optimization*, vol. 46, no. 1, pp. 288–307, 2007.
- [8] M. Cao, A. S. Morse, and B. D. O. Anderson, “Agree asynchronously,” *IEEE Transactions on Automatic Control*, vol. 53, no. 8, pp. 1826–1838, 2008.
- [9] Z. Sun, S. Mou, B. D. O. Anderson, and A. S. Morse, “Formation movements in minimally rigid formation control with mismatched mutual distances,” in *Decision and Control (CDC), 2014 IEEE 53rd Annual Conference on*. IEEE, 2014, pp. 6161–6166.
- [10] M. Fruhnert and M. Corless, “Consensus control of linear second-order discrete-time systems with guaranteed rate of convergence,” in *Control Conference (ECC), 2015 European*. IEEE, 2015, pp. 1492–1497.
- [11] M. Cao, A. S. Morse, and B. D. O. Anderson, “Reaching a consensus in a dynamically changing environment: a graphical approach,” *SIAM Journal on Control and Optimization*, vol. 47, pp. 575–600, Feb. 2008.
- [12] X. Wang and Y. Hong, “Finite-time consensus for multi-agent networks with second-order agent dynamics,” *IFAC Proceedings Volumes*, vol. 41, no. 2, pp. 15 185–15 190, 2008.
- [13] K. Cai and H. Ishii, “Quantized consensus and averaging on gossip digraphs,” *IEEE Transactions on Automatic Control*, vol. 56, no. 9, pp. 2087–2100, Sept 2011.
- [14] —, “Average consensus on general strongly connected digraphs,” *Automatica*, vol. 48, no. 11, pp. 2750–2761, 2012.
- [15] M. Liu, F. Wirth, M. Corless, and R. Shorten, “On the stability and convergence of a class of consensus systems with a nonlinear input,” *Automatica*, Accepted 2017.
- [16] X. Chen, J. Liu, M. A. Belabbas, Z. Xu, and T. Baar, “Distributed evaluation and convergence of self-appraisals in social networks,” *IEEE Transactions on Automatic Control*, vol. 62, no. 1, pp. 291–304, Jan 2017.
- [17] Y. Hong, J. Hu, and L. Gao, “Tracking control for multi-agent consensus with an active leader and variable topology,” *Automatica*, vol. 42, no. 7, pp. 1177–1182, 2006.
- [18] X. Wang, S. Mou, and D. Sun, “Improvement of a distributed algorithm for solving linear equations,” *IEEE Transactions on Industrial Electronics*, 2016, doi:10.1109/TIE.2016.2636119.

- [19] P. Wang, W. Ren, and Z. Duan, "Distributed minimum weighted norm solution to linear equations associated with weighted inner product," *Proceedings of the 55th Conference on Decision and Control*, pp. 5220–5225, 2016.
- [20] L. Wang, D. Fullmer, and A. S. Morse, "A distributed algorithm with an arbitrary initialization for solving a linear algebraic equation," *Proceedings of American Control Conference*, pp. 1078–1081, 2016.
- [21] S. Mou, Z. Lin, L. Wang, D. Fullmer, and A. S. Morse, "A distributed algorithm for efficiently solving linear equations and its applications (special issue jcw)," *Systems & Control Letters*, pp. 21–27, 2016.
- [22] S. Kar, J. M. F. Moura, and K. Ramanan, "Distributed parameter estimation in sensor networks: Nonlinear observations models and imperfect communication," *IEEE Transactions on Information Theory*, no. 6, pp. 1–52, 2012.
- [23] S. Nabavi, J. Zhang, and A. Chakraborty, "Distributed optimization algorithms for wide-area oscillation monitoring in power systems using interregional pmu-pdc architectures," *IEEE Transactions on Smart Grid*, vol. 6, no. 5, pp. 2529–2538, Sept 2015.
- [24] B. Zhao, F. Lin, C. Wang, X. Zhang, M. Polis, and L. Wang, "Supervisory control of networked timed discrete event systems and its applications to power distribution networks," *IEEE Transactions on Control of Network Systems*, vol. PP, no. 99, pp. 1–1, 2015.
- [25] J. Qin, C. Yu, and B. Anderson, "On leaderless and leader-following consensus for interacting clusters of second-order multi-agent systems," *Automatica*, vol. 74, pp. 214 – 221, 2016.
- [26] Z. Lin, M. Broucke, and B. Francis, "Local control strategies for groups of mobile autonomous agents," *IEEE Trans. Autom. Control*, vol. 49, no. 4, pp. 622–629, April 2004.
- [27] Z. Sun, U. Helmke, and B. D. O. Anderson, "Rigid formation shape control in general dimensions: an invariance principle and open problems," in *2015 54th IEEE Conference on Decision and Control (CDC)*, Dec 2015, pp. 6095–6100.
- [28] T. C. Aysal and K. E. Barner, "Constrained decentralized estimation over noisy channels for sensor networks," *IEEE Transactions on Signal Processing*, vol. 56, no. 4, pp. 1398–1410, April 2008.
- [29] C. G. Lopes and A. H. Sayed, "Diffusion least-mean squares over adaptive networks: Formulation and performance analysis," *IEEE Transactions on Signal Processing*, vol. 56, no. 7, pp. 3122–3136, July 2008.
- [30] G. Mateos, I. D. Schizas, and G. B. Giannakis, "Distributed recursive least-squares for consensus-based in-network adaptive estimation," *IEEE Transactions on Signal Processing*, vol. 57, no. 11, pp. 4583–4588, Nov 2009.
- [31] F. Longstaff and E. Schwartz, "Valuing american options by simulation: a simple least-squares approach," *Review of Financial studies*, vol. 14, no. 1, pp. 113–147, 2001.
- [32] Z. Luo, "Universal decentralized estimation in a bandwidth constrained sensor network," *IEEE Transactions on Information Theory*, vol. 51, no. 6, pp. 2210–2219, June 2005.
- [33] A. Nedic and A. Ozdaglar, "Distributed sub-gradient methods for multi-agent optimization," *IEEE Transactions on Automatic Control*, no. 1, pp. 48–61, 2009.

- [34] J. C. Duchi, A. Agarwal, and M. J. Wainwright, “Dual averaging for distributed optimization: Convergence analysis and network scaling,” *IEEE Transactions on Automatic Control*, no. 3, pp. 592–606, 2012.
- [35] D. Jakovetic, J. M. F. Moura, and J. Xavier, “Fast distributed gradient methods,” *IEEE Transactions on Automatic Control*, no. 5, pp. 1131–1146, 2014.
- [36] T. Chang, A. Nedic, and A. Scaglione, “Distributed constrained optimization by consensus-based primal-dual perturbation method,” *IEEE Transactions on Automatic Control*, no. 6, pp. 1524–1538, 2014.
- [37] A. Nedic, A. Ozdaglar, and P. A. Parrilo, “Constrained consensus and optimization in multi-agent networks,” *IEEE Transactions on Automatic Control*, no. 4, pp. 922–938, 2010.
- [38] A. K. Sahu, S. Kar, J. M. F. Moura, and H. V. Poor, “Distributed constrained recursive nonlinear least-squares estimation: Algorithms and asymptotics,” *ArXiv e-prints*, Jan. 2016.
- [39] G. Shi, B. D. O. Anderson, and U. Helmke, “Network flows that solve linear equations,” *IEEE Transactions on Automatic Control*, vol. 62, no. 6, pp. 2659–2674, June 2017.
- [40] J. Wang and N. Elia, “Distributed least square with intermittent communications,” in *2012 American Control Conference (ACC)*, June 2012, pp. 6479–6484.
- [41] —, “A control perspective for centralized and distributed convex optimization,” in *2011 50th IEEE Conference on Decision and Control and European Control Conference*, Dec 2011, pp. 3800–3805.
- [42] B. Gharesifard and J. Cortés, “Distributed continuous-time convex optimization on weight-balanced digraphs,” *IEEE Trans. Autom. Control*, vol. 59, no. 3, pp. 781–786, Mar. 2014.
- [43] F. Chung, *Spectral graph theory*. American Mathematical Soc., 1997, vol. 92.
- [44] D. Bertsekas, *Nonlinear programming*. Athena scientific Belmont, 1999.
- [45] J. Wang and N. Elia, “Control approach to distributed optimization,” in *2010 48th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, Sept 2010, pp. 557–561.
- [46] R. Bronson, *Matrix methods: an introduction*. Gulf Professional Publishing, 1991.