

Quantum critical singularities in two-dimensional metallic XY ferromagnets

Chandra M. Varma

Department of Physics, University of California, Riverside, CA. 92521

W. J. Gannon and M. C. Aronson

Department of Physics and Astronomy,

Texas A&M University, College Station, TX 77843

J. A. Rodriguez-Rivera and Y. Qiu

NIST Center for Neutron Research,

National Institute of Standards and Technology, Gaithersburg, MD 20899 USA

(Dated: June 18, 2018)

Abstract

An important problem in contemporary physics concerns quantum-critical fluctuations in metals. A scaling function for the momentum, frequency, temperature and magnetic field dependence of the correlation function near a 2D-ferromagnetic quantum-critical point (QCP) is constructed, and its singularities are determined by comparing to the recent calculations of the correlation functions of the dissipative quantum XY model (DQXY). The calculations are motivated by the measured properties of the metallic compound $\text{YFe}_2\text{Al}_{10}$, which is a realization of the DQXY model in 2D. The frequency, temperature and magnetic field dependence of the scaling function as well as the singularities measured in the experiments are given by the theory without adjustable exponents. The same model is applicable to the superconductor-insulator transitions, classes of metallic AFM-QCPs, and as fluctuations of the loop-current ordered state in hole-doped cuprates. The results presented here lend credence to the solution found for the 2D-DQXY model, and its applications in understanding quantum-critical properties of diverse systems.

I. INTRODUCTION

$\text{YFe}_2\text{Al}_{10}$ is nearly tetragonal, with a divergent uniform magnetic susceptibility at low temperatures with field applied in the a-c plane, but a constant value at the same temperatures for fields applied along the b-axis¹. There is no observed anisotropy of the susceptibility within the a-c plane. These results suggest that the metal is accidentally close to a ferromagnetic quantum critical point and that the relevant model for criticality is the 2D-XY model. The specific heat divided by temperature is logarithmic in temperature. We show here that the singularity in the susceptibility and the specific heat together and the singularity in the frequency/temperature dependence of the correlations² and their contrast with the momentum dependence are consistent with the recent solution of the 2D-DQXY model.

Classical 2D FM transitions of the Berezinskii, Kosterlitz-Thouless^{3,4} variety at finite T have been found in some insulating compounds in the past⁵. $\text{YFe}_2\text{Al}_{10}$ appears to be the first metallic compound to be very near a planar ferro-magnetic quantum-transition.

II. RESPONSE FUNCTION OF A 2D XY MODEL NEAR QUANTUM CRITICALITY

The 2D-dissipative quantum XY model describes the physics of interacting quantum rotors lying in a plane and includes dissipation due to transfer of energy to other excitations. It is specified by the action given, for example, by Eq. (1) in Ref. 6. Without dissipation, the phase diagram and the correlation functions of the quantum XY model in 2D belong to the classical 3D XY universality class. But in a metal, the dissipation introduced by coupling of the fluctuations to corresponding incoherent fluctuations of the fermions, leads to a much richer phase diagram⁶⁻⁸. A theory of the phase diagram and of the quantum-critical fluctuations has been derived and tested by quantum Monte-Carlo calculations^{6,9,10}. The fluctuations in such theories present a new paradigm in quantum critical phenomena. The conventional theories of quantum-critical phenomena^{11,12} are based on anharmonic soft spin-fluctuations, which are extensions of the theory of classical dynamical critical phenomena¹³, applicable to models of the Ginzburg-Landau-Wilson type. In such theories, the frequency and momentum dependence of the correlation function are always entangled and a finite dynamical exponent z given by the dispersion of the spin-wave excitations in the presence of

dissipation relates the spatial and temporal correlations. A quite different class of correlation functions are found for the 2D-DQXY model because the critical properties are determined not by spin-wave excitations but by topological excitations in space and time.

The 2D- DQXY model can be exactly transformed^{9,14} to a model of orthogonal topological charges, warps and vortices. Warps interact with each other in (imaginary) time and are essentially local in space while the vortices interact purely in space. The correlation function of the order parameter $e^{i\theta(\mathbf{r},\tau)}$ of the 2D-DQXY model have been derived by quantum Monte-Carlo⁶ which also checks their relation to the correlation functions of warps and vortices. The model transformed to interacting topological excitations has also been solved analytically¹⁵. The correlation function is found in an extensive region of parameters in which the proliferation of warps determines the criticality to be,

$$\mathcal{C}(r, \tau) \equiv \langle e^{i\theta(\mathbf{r},\tau)} e^{-i\theta(0,0)} \rangle \approx \chi_0 \log(r_0/r) e^{(-r/\xi_r)} \frac{1}{\tau} e^{-\left(\frac{\tau}{\xi_\tau}\right)}. \quad (1)$$

The three especially note-worthy features of (1) are (i) it is separable in its r and τ dependence, (ii) its thermal Fourier transform at criticality, when $\xi_\tau \rightarrow \infty$ has the ω/T scaling⁹, introduced in critical phenomena in Ref.¹⁶ and termed "Planckian"¹⁷, and (iii) that^{6,15}

$$(\xi_r/a) = \log(\xi_\tau/\tau_c). \quad (2)$$

This means that the dynamical critical exponent z is effectively ∞ . ξ_τ has an essential singularity as a function of the dimensionless dissipation parameter α but an algebraic singularity as a function of the dimensionless parameter $\tilde{K} \equiv \sqrt{K K_\tau}$. Here K is the Josephson coupling and K_τ is the kinetic energy parameter in the quantum XY model. On the disordered side of the QCP, ξ_τ is given by,

$$\xi_\tau/\tau_c = e^{\sqrt{c\alpha_c/(\alpha_c-\alpha)}}, \text{ at constant } \tilde{K}, \text{ and } = \left(\frac{\tilde{K}_c}{\tilde{K}_c - \tilde{K}} \right)^{\nu_\tau}, \text{ at constant } \alpha; \quad \nu_\tau \approx 1/2. \quad (3)$$

c is a constant of $O(1)$ and τ_c a short-time cut-off. If the transition, as expected is driven by $(\tilde{K} - \tilde{K}_c)$, the logarithmic dependence of the spatial correlation function may lead to a very short observed correlation length unless the sample is tuned to very small values of $(\tilde{K} - \tilde{K}_c)$, and other effects, such as disorder do not change the asymptotic critical properties.

At criticality, i.e. for $\xi_\tau^{-1} = \xi_r^{-1} = 0$, the *thermal Fourier transform* of the correlation function is

$$\mathcal{C}(q, w, T) \propto \frac{1}{q^2} \tanh \frac{\omega}{2T}, \quad (4)$$

with a high frequency cut-off. For finite ξ_τ and ξ_r the infra-red singularities are cut-off and their form is given in the Appendix in Ref.(6).

III. SCALING FOR THE FERROMAGNETIC QUANTUM XY MODEL IN A FIELD

The magnetic field B_\perp in the plane couples linearly to the order parameter and serves as a cut-off to the quantum critical regime. To address the experimental results, we first present a scaling theory for the correlation function in a magnetic field, and connect the results to the calculated form, Eq. (1) derived at $B = 0$.

A novelty is to derive a scaling form of the correlation function when the spatial correlations depend logarithmically on the temporal correlation length and neither may bear power-law relations to the control parameters. Consider the response of the 2D-XY ferromagnet with a uniform field B in the easy plane at a temperature T to a small applied time and space dependent field $h(r, t)$, also in the easy plane. Follow the usual process of scaling for the correlation function on taking the derivative of logarithm of the partition function with respect to $h(r_1, \tau_1)$ and $h(r_2, \tau_2)$, $r = |r_1 - r_2|$, $\tau = \tau_1 - \tau_2$, and scale the space and time-metric together with the scaling operators in the action so as to keep the singular part of the partition function invariant. The space-metric is expanded by the correlation length ξ_r and the time-metric by ξ_τ . The renormalization group eigenvalue for B_\perp on scaling time is defined to be z_b .

$$\mathcal{C}(r, \tau, T, B_\perp) = \xi_r^{-2d} \xi_\tau^{-2} \xi_\tau^{2z_b} \chi\left(\frac{r}{\xi_r}, \frac{\tau}{\xi_\tau}, T\xi_\tau, B_\perp \xi_\tau^{z_b}\right). \quad (5)$$

The $q = 0, \omega = 0$ limit of the correlation function is found by integrating over r and τ . Divided by T , this gives the temperature and magnetic field dependence of the static uniform susceptibility, Eq. (6). The integration over the space-variable brings a factor ξ_r^d , as usual. At this point the special properties of the results in (1) may be used. Since the temporal correlation function is $\propto 1/\tau$ at criticality, integration over τ can produce at most only logarithmic corrections, which may be neglected to begin with in comparison with the rest. Also, since $\xi_r \propto \log \xi_\tau$, the space dependent prefactors may also be neglected to

logarithmic accuracy. So we get

$$\begin{aligned}\chi(T, B_\perp) &\equiv \frac{dM(T, B_\perp)}{dB_\perp} = \frac{1}{T} \langle \cos^2(\theta) \rangle = \frac{1}{T} C(q=0, \omega=0, T, B_\perp) \\ &= \frac{1}{T} \xi_\tau^{-2+2z_b} \chi(T\xi_\tau, B_\perp \xi_\tau^{z_b}).\end{aligned}\quad (6)$$

On re-scaling $T\xi_\tau \rightarrow 1$ to express ξ_τ in terms of T , one gets

$$\chi(T, B_\perp) \propto T^{(1-2z_b)} f_{1,\chi}\left(\frac{B_\perp}{T^{z_b}}\right),\quad (7)$$

or equivalently

$$\chi(T, B_\perp) \propto B_\perp^{(1-2z_b)/z_b} f_{2,\chi}(T/B_\perp^{(1/z_b)}). \quad (8)$$

On comparing (7) with the static susceptibility calculated from (1) and again neglecting logarithmic corrections, we find that the two are mutually consistent only if $z_b = 1$. Given the $1/T$ factor in (6), the correlation function has an exponent 0 which is consistent with having logarithmic corrections. Scaling cannot give the logarithmic corrections, which turn out to be important in relation to experiments, as seen below. We therefore explicitly calculate the magnetic susceptibility by the Monte-Carlo technique using the procedure of Refs. 6 for the dissipative quantum XY model.

A. Monte-Carlo Calculations:

The uniform magnetic susceptibility per unit-cell is

$$\chi(T) \equiv \frac{1}{N^2} \sum_i^{N^2} \int_0^\beta d\tau \langle M(i, \tau) M(i, 0) \rangle; \quad M(i, \tau) = \cos(\theta_i)(\tau), \quad (9)$$

where N^2 is the number of unit-cells on a lattice labelled by i . This is converted to a form suitable for quantum Montecarlo calculations on a discrete space $N \times N$ and imaginary time one-dimensional lattice τ_n of N_τ cells,

$$\chi(T) = \frac{1}{N^2 N_\tau} \sum_{i,n}^{N^2, N_\tau} \langle \cos(\theta_i, n) \cos(\theta_i, 0) \rangle. \quad (10)$$

The calculation is entirely as in the calculation of the action susceptibility, Eq. (10), of Ref. 6. The discretization and calculation procedure is also fully described there in Sec. II-C. $\tau_n = n\delta\tau = n\beta/N_\tau$. $\delta\tau = \tau_c$ is the ultra-violet (short) time cut-off. The temperature

in the calculation is controlled by N_τ^{-1} . This has been calculated on a $N^2 = 50 \times 50$ lattice and with N_τ ranging from 20 to 200. With an upper-cutoff $\tau_c^{-1} = 160K$, this effectively gives results at discrete temperatures from 20 to 0.8 K. The results for $\chi(T)$ are given in Fig. (1).

The black crosses in Fig. (1) are the result, and they compare favorably to the measured uniform susceptibility χ , also shown in Fig. 1. Motivated by the discussion above, we look for logarithmic factors multiplying T^{-1} . We find that the calculated susceptibility fits $T^{-1} \log^2(T/\omega_c)$, where $\omega_c = 1/\tau_c$ is the high energy cut-off given in Eq. (3). We also show the experimentally derived function $T^{-1.4}$, which mimics $\frac{1}{T} (\log T\tau_c)^2$ very well over the range of experimental temperatures, with $\tau_c^{-1} = 160$ K. The previously reported scaling analysis¹ is purely phenomenological, with two critical exponents that are determined by the experiments and with a spatial correlation length, discussed below, which is in qualitative conflict with experiments. In contrast, the logarithmic corrections found here leave no parameter in the theory undetermined.

The experimental results¹ for the scaling of $M(B, T)$ in YFe_2Al_{10} , previously fitted¹ to the scaling expression

$$-(d(M/B_\perp)/dT)B_\perp^{1.4} \propto F(T/B_\perp^{(1-0.4)}), \quad (11)$$

are compared to the result

$$-(d(M/B_\perp)/dT) \frac{B_\perp}{\log^2(B\tau_c)} \propto f_M\left(\frac{T}{B_\perp \log^2(B\tau_c)}\right). \quad (12)$$

in Fig. 2.

Fig. 2 shows that Eqn. 12 gives an acceptable scaling collapse, with a tiny off-set of the field $B \rightarrow B'$, $B' = B + 0.07$ T and $B_c = 100$ T, the latter in reasonable agreement with the value of $\tau_c^{-1} = 160$ K, using $g\mu_B B_c = \hbar/\tau_c$ with the Landé g-factor taken to be 2 and μ_B the Bohr magneton. We do not know the origin of the small off-set of 0.07 T required to best fit the data which spans the range up to 6 T; it may be due to impurities in the sample.

B. Scaling of the Free-energy:

The scaling for the free-energy per unit volume may be considered similarly

$$f(T, B) \propto T \xi_r^{-d} \xi_\tau^{-1} \Phi(T \xi_\tau, B_\perp \xi_\tau^{z_b}) \quad (13)$$




FIG. 1. Comparison between the uniform magnetic susceptibility $\chi = M/B$ of $\text{YFe}_2\text{Al}_{10}$ measured in a field of 0.05 T in the critical $a - c$ plane (red circles)¹ to fits to $\propto T^{-1.4}$ (blue line) and the form calculated from the correlation functions of the dissipative 2D-quantum XY Model in this paper with $\tau_c^{-1} = 160$ K, (green line) which is the approximate scale below which the divergent form appears in the experiments. Also shown are direct calculation of χ by Monte-Carlo method (black crosses), with temperature and susceptibility scaled to the experimental data. Fits are performed for $T < 20$ K with $\tau_c^{-1} = 160$ K as a fitted parameter in the theory for the 2DXY model.

This gives, using the same results as for the calculation of magnetization, that

$$f(T, B) \propto T^2 \ln^2(T\tau_c) \tilde{\Phi}\left(\frac{B_\perp}{T/\log^2(T\tau_c)}\right). \quad (14)$$

With (14), the results for $M(T, B_\perp)$ and $\chi(T, B_\perp)$ derived above from the correlation functions follow to logarithmic accuracy. The specific heat divided by T at constant B_\perp has in addition to a constant and a $\log(T)$ term a $\log^2(T)$ term with a coefficient that is 1/3 of the logarithmic term. The specific heat as a function of magnetic field B_\perp , similarly follows. Note the factor T in (13). This is un-important for classical transitions, where it is replaced near criticality by T_c but essential to keep for a transition with $T \rightarrow 0$




FIG. 2. (A) The scaling of the uniform magnetization divided by field M/B as a function of temperature T measured at different fixed fields B indicated in the figure to Eqn. 11, as shown in¹. Measurements were made for temperatures $1.8 \leq T \leq 30$ K with the field in the critical $a-c$ plane. (B) Same data as (A), which can be scaled using Eqn. 12, with a substitution of the external field B with $B' = B + 0.07$ T and $B_c = 100$ T. Colors indicating different magnetic fields are the same as (A).

C. Dynamics

Consider now the extension of the correlation function, Eq. (5) to obtain the frequency and momentum dependent magnetic response function. In the absence of the detailed Monte-Carlo calculations of the correlation function in a magnetic field, one may guess on grounds given below that the magnetic response function has the approximate scaling form,

$$\chi''(q, \omega, T, B_{\perp}) \propto \frac{\log(\sqrt{\omega^2 + (2k_B T)^2} \tau_c)}{\sqrt{\omega^2 + (2k_B T)^2}} f_{\chi}\left(\frac{\omega}{T}, qa \log(\xi_{\tau}/\tau_c), \frac{B_{\perp}}{(T/\log(\sqrt{\omega^2 + (2k_B T)^2} \tau_c))}\right) \quad (15)$$

This follows the form of the derived correlation function (1) except for the modifications necessary due to the scaling corrections due to B_{\perp} . The logarithmic term and its argument have been chosen so that it reproduces the temperature dependence of the calculated uniform magnetic susceptibility, derived by using the Kramers-Kronig relation between the imaginary part $\chi''(q, \omega, T, B_{\perp})$ and the real part at $\omega = 0$, as well as the magnetic field dependence of the magnetization derived above.

Eq. (15) may be put in various other forms as desired. It follows that for finite B_{\perp} ,

this divergence is cut-off. It is predicted that together with $\omega/2T$ scaling of the form calculated in microscopic theory to be of the form $\tanh(\omega/2T)$, with a cut-off at $\omega_c = \tau_c^{-1}$, there should be singular pre-factors. This has been tested by inelastic neutron scattering as described in Ref.², where it is found that a cutoff energy dependence of the form $(\omega^2 + (\pi T)^2)^{-1.4/2} \tanh(\omega/T)$ describes the data reasonably well. The data can be equally fitted by Eq. (15).

A comparison of Eq. (15) presented here to the energy dependence of the measured dynamical susceptibility is shown in Fig. 3 for $\omega \gg T$. The correspondence between temperature and energy revealed by a previous scaling analysis¹ and the Kramers-Kronig relation² suggests that the momentum-integrated dynamical susceptibility χ'' is a function of $(1/\omega) \log^2(\omega\tau_c)$ with a high energy cutoff ω_c . Fixing $\tau_c^{-1} = 14$ meV (≈ 160 K) gives the fit shown in Fig. 3, nearly indistinguishable from the phenomenological $E^{-1.4}$ power law behavior used previously². The correspondence of Eq. (15) to the change in the dynamics in a magnetic field may be seen in that paper.

On considering the q -dependence, one encounters an interesting discrepancy in relation to the experiments. The sample is, not surprisingly, not exactly at criticality. The dynamical measurements, both through neutron scattering and more directly through the muon spin-relaxation rate¹⁹ suggest a low temperature cut-off in the experiments of about 1 K. So $\xi_\tau^{-1} \approx 1K$. In experiments not exactly at criticality, T and B_\perp should be replaced approximately by $\sqrt{T^2 + \xi_\tau^{-2}}$ and $\sqrt{B_\perp^2 + B_x^2}$ with $(g\mu_B B_x)^2 < S^2 \gg \xi_\tau^{-2}$, where $(g\mu_B)^2 < S^2 >$ is a measure of the mean-square magnetic moment in the fluctuations. Using $\tau_c^{-1} \approx 100K$, the corresponding cut-off in the spatial correlation length ξ_r may be estimated using Eqs. (2) to be about 4 lattice constants. But the spatial correlation length in neutron scattering experiments is only about a lattice constant, although independent of temperature in accord with the theory. A possible explanation²⁰ of such extreme spatial locality while scale-invariant behavior is observed with long temporal correlation length ξ_τ/τ_c of $O(10^2)$ may lie in the crossover due to disorder in quantum-critical problems with large dynamical critical exponent z . This matter can be tested by further experiments in samples closer to criticality. Tuning closer to criticality may be difficult since using the second of (3), which is the more likely applicable, $\xi_\tau/\tau_c \approx 10^2$ implies that already $(1 - \tilde{K}/\tilde{K}_c) \approx 10^{-3}$.




FIG. 3. The dynamical susceptibility χ'' at $T = 0.1K$ as a function of energy obtained by inelastic neutron scattering using the MACS spectrometer at the National Institute For Standards and Technology¹⁸. Data (red circles) were measured at a constant fixed energy over a large area of q -space, corrected for the Fe^{2+} form factor, then integrated over four Brillouin zones that are out of the critical ($a-c$) plane (along q_K , parallel to the crystal b -axis) and over one Brillouin zone within the critical plane (along q_L , parallel to the crystal c -axis)². Measurements were made with a small bias field of 0.025 T along the crystal a -axis to suppress superconductivity in the aluminum sample holder. Fits are made to the forms given by Eq. (15) for $E = \hbar\omega \gg k_B T$, so that $\chi'' \sim E^{-1.4}$ (black line), and $(1/E) \log^2(E \tau_c)$ (green line), fixing $(\tau_c^{-1} = 14 \text{ meV} \approx 160 \text{ K})$. Error bars on data points represent one standard deviation.

IV. CONCLUDING REMARKS

These results test the theory of the 2+1 D - XY model in considerable detail. In particular, the success of the results in explaining the singularities in the properties associated with the free energy depends on the novel results of the theory that the correlation function is the product of a function in space and a function in time, and that the spatial correlations vary logarithmically as the temporal correlations. The result that at criticality the time-

dependence is proportional to $1/\tau$, i.e has the Planckian scaling ω/T , has also been crucial. As may easily be seen, these results cannot be obtained by simply putting the dynamical exponent $z \rightarrow \infty$ in the conventional dynamical critical theory. Further tests of the theory require samples in which the distance to quantum-criticality can be systematically changed, for example, by applying pressure, thereby observing a longer spatial correlation length varying logarithmically as the distance to the critical point.

Acknowledgements: CMV acknowledges with pleasure discussions with Joerg Schmalian and Alexei Tsvelik. Special thanks are due to Changtao Hou and Lijun Zhu who wrote the Monte-Carlo routines used for the results shown in Fig. (1). Part of this research was conducted at Brookhaven National Laboratory, where W. J. G and M. C. A were supported under the auspices of the US Department of Energy, Office of Basic Energy Sciences, under contract DE-AC02-98CH1886. Access to MACS was provided by the Center for High Resolution Neutron Scattering, a partnership between the National Institute of Standards and Technology and the National Science Foundation under Agreement No. DMR-1508249.

-
- ¹ L. S. Wu, M. S. Kim, K. Park, A. M. Tsvelik, and M. C. Aronson, Proceedings of the National Academy of Sciences **39**, 14088 (2014).
- ² W. J. Gannon, L. S. Wu, I. A. Zaliznyak, W. Xu, A. M. Tsvelik, J. A. Rodriguez-Rivera, Y. Qiu, and M. C. Aronson, arXiv:1712.04033 (2017).
- ³ J. Kosterlitz and D. Thouless, Journal of Physics C: Solid State Physics **6**, 1181 (1973).
- ⁴ V. Berezinskii, Zh. Eksp. Teor. Fiz. **32**, 493 (1970).
- ⁵ S. T. Bramwell, P. C. W. Holdsworth, and M. T. Hutchings, Journal of the Physical Society of Japan **64**, 3066 (1995).
- ⁶ L. Zhu, Y. Chen, and C. M. Varma, Physical Review B **91**, 205129 (2015).
- ⁷ E. B. Stiansen, I. B. Sperstad, and A. Sudbø, Physical Review B **85**, 224531 (2012).
- ⁸ A Caldeira-Leggett form of dissipation is used in Refs. 6, 9, and 10. In the present context, this can be shown to come from the coupling $\nabla\theta$, which represents the collective ferromagnetic spin-current, to the spin-current of the incoherent fermions.
- ⁹ V. Aji and C. M. Varma, Physical Review Letters **99**, 067003 (2007).
- ¹⁰ L. Zhu, C. Hou, and C. M. Varma, Physical Review B **94**, 235156 (2016).

- ¹¹ T. Moriya, Spin Fluctuations in Itinerant Electron Magnetism, Springer Series in Solid-State Sciences (Springer-Verlag, Berlin, 1985).
- ¹² J. A. Hertz, Physical Review B **14**, 1165 (1976).
- ¹³ P. C. Hohenberg and B. I. Halperin, Reviews of Modern Physics **49**, 435 (1977).
- ¹⁴ V. Aji and C. M. Varma, Physical Review B **82**, 174501 (2010).
- ¹⁵ C. Hou and C. M. Varma, Physical Review B **94**, 201101(R) (2016).
- ¹⁶ C. M. Varma, P. B. Littlewood, S. Schmitt-Rink, E. Abrahams, and A. E. Ruckenstein, Physical Review Letters **63**, 1996 (1989).
- ¹⁷ J. Zaanen, Nature **430**, 512 (2004).
- ¹⁸ J. A. Rodriguez, D. M. Adler, P. C. Brand, C. Broholm, J. C. Cook, C. Brocker, R. Hammond, Z. Huang, P. Hundertmark, J. W. Lynn, N. C. Maliszewski, J. Moyer, J. Orndorff, D. Pierce, T. D. Pike, G. Scharfstein, S. A. Smee, and R. Vilaseca, Measurement Science and Technology **19**, 034023 (2008).
- ¹⁹ K. Huang, C. Tan, J. Zhang, Z. Ding, D. E. MacLaughlin, O. O. Bernal, P.-C. Ho, C. Baines, L. S. Wu, M. C. Aronson, and L. Shu, arXiv:1801.03659 (2018).
- ²⁰ C. M. Varma, arXiv:1701.03853 (2017).