

EVERY EXPONENTIAL GROUP SUPPORTS A POSITIVE HARMONIC FUNCTION

GIDEON AMIR AND GADY KOZMA

ABSTRACT. We prove that all groups of exponential growth support non-constant positive harmonic functions. In fact, our results hold in the more general case of strongly connected, finitely supported Markov chains invariant under some transitive group of automorphisms for which the directed balls grow exponentially.

1. INTRODUCTION

Let G be a finitely generated group, and let S be a set of generators. We call a function $f : G \rightarrow \mathbb{R}$ (right) harmonic if for every $g \in G$ we have

$$f(g) = \frac{1}{|S|} \sum_{s \in S} f(gs).$$

There is considerable interest in identifying spaces of harmonic functions with various properties and relating them to properties of the group G . In particular, the spaces of *bounded* harmonic functions and of *positive* harmonic functions were studied (these are related, respectively, to the Poisson and Martin boundaries of G). Our interest here is the following question: for which G and S does a positive, non-constant harmonic function exist?

One example where this problem is relatively understood is the case of nilpotent groups. In this case, Margulis [4] reduced the question from G to $G/[G, G]$ i.e. from general nilpotent groups to abelian groups. When S is symmetric, i.e. when $s \in S \Leftrightarrow s^{-1} \in S$, any positive harmonic function on an abelian group is constant, see [2] for a proof and some results also in the non-symmetric case (Margulis' reduction is not restricted to the symmetric case). Similarly, Hebisch and Saloff-Coste showed that when G has polynomial word growth and S is symmetric the only positive harmonic functions are the constants [3, Theorem 6.2]. Since by Gromov's theorem groups with polynomial word growth are virtually nilpotent, this result is quite close to Margulis', but the proof of [3] is analytic in nature (and in particular does not use Gromov's theorem or any other algebraic structure result).

In the other direction, Bougerol and Élie showed that for a certain class of groups, close in spirit to Lie groups, exponential word growth implies that a positive non-constant harmonic function does exist [1, Theorem 1.4]. Here we show that most of the assumptions of [1] are not necessary: the result

holds for any finitely generated group and any finite set of generators. Even symmetry is not necessary. Let us state our result:

Theorem. *Let R_n be a strongly connected, finitely supported Markov chain on some set G , invariant to some transitive group of automorphisms of G . Assume the directed balls on G grow exponentially. Then there exists a non-constant positive harmonic function on G .*

We recall that a Markov chain is called strongly connected if for any two states of the chain (here this is the same as the elements of G) there is a positive probability for the chain to reach from one state to the other in a finite number of steps. It is called “finitely supported” if from any state one may reach only finitely many states in a single step (in general this might depend on the starting state, but here due to the invariance all starting states are identical). The directed ball around g of radius r , denoted by $B(g, r)$, is the set of all $h \in G$ such that for some $s \leq r$ the probability to reach h from g in s steps of the Markov chain is positive. Finally, a function $f : G \rightarrow \mathbb{R}$ is called harmonic to a Markov chain R_n on G if $f(R_n)$ is a martingale. The reader may readily verify that these definitions coincide with the definitions given above for a group, when G is a group and the Markov chain is the one moving from a $g \in G$ to each of the elements of the form gs for some $s \in S$ with probability $1/|S|$.

Question. Does the Grigorchuk group support a non-trivial positive harmonic function?

2. PROOF

It will be convenient to use graph notation, so whenever R_n is a Markov chain on some set G , we will consider G as a directed graph: the elements of G will be called “vertices”, and a couple $x, y \in G$ will be called a (directed) edge if the Markov chain can move from x to y in a single step (we denote $x \rightarrow y$). A path will be a sequence of vertices x_1, x_2, \dots such that $x_i \rightarrow x_{i+1}$ and a geodesic is a path which is shortest possible between its end vertices (equivalently, between any two vertices in it).

Let R_n be a Markov chain on some G . For a set S of vertices, an $a \in S$ and a vertex $x \in \partial S$ (i.e. $x \notin S$ but $\exists y \in S$ such that $y \rightarrow x$) let $\mu_S(a, x)$ be the probability that R_n , when started from a , exits S at x . If a and b are two vertices we denote

$$\epsilon(S) = \epsilon(S; a, b) = \max_x \frac{|\mu_S(a, x) - \mu_S(b, x)|}{\mu_S(a, x)}$$

where the maximum is taken over all $x \in \partial S$ such that $\mu_S(a, x) > 0$.

Lemma 1. *Let R_n be a strongly connected Markov chain on some G . Then there exists a non-trivial positive harmonic function on G if and only if there exist vertices a and b such that $\epsilon(S; a, b) \not\rightarrow 0$ as $S \rightarrow G$.*

(we say that sets of vertices S converge to G if every vertex of G is eventually contained in S)

Proof. Let us start with the “if” part, for which we assume a and b exist and we need to construct a positive harmonic function f . In more details, our assumption is that there exist an $\epsilon > 0$, a sequence S_n and vertices $x_n \in \partial S_n$ such that

$$|\mu_{S_n}(a, x_n) - \mu_{S_n}(b, x_n)| > \epsilon \mu_{S_n}(a, x_n).$$

We now define functions f_n using

$$f_n(v) = \frac{\mu_{S_n}(v, x_n)}{\mu_{S_n}(a, x_n)}$$

(extend f_n to outside S_n arbitrarily).

We wish to take a subsequential limit of the f_n and for this we need to get an a priori bound. Here is where we use strong connectedness. Indeed, if v and w are vertices, and if γ is some path from v to w which R may take with probability $p > 0$, then for every n sufficiently large such that $\gamma \subset S_n$ we have $\mu(v, x_n) > p\mu(w, x_n)$ and hence $f_n(v) > pf_n(w)$. Since $f_n(a) = 1$ we get that for every v there exists a constant $p(v)$ such that $|f_n(v)| \leq p(v)$ for all n sufficiently large. This means that we may take a subsequence n_k such that $f_{n_k}(v)$ converges for every v . The resulting limit is positive, harmonic, and nontrivial since $f_{n_k}(a) = 1$ but $|f_{n_k}(b) - 1| \geq \epsilon$ and both facts are preserved in the limit.

We move to the “only if” part. We thus assume that for all a and $b \in S$ $\epsilon(S) \rightarrow 0$. Let f be some positive harmonic function. Let a and b be vertices, and S some finite set containing both. Since $f(R_n)$ is a martingale (by definition), the optional stopping theorem tells us that

$$f(a) = \sum_{x \in \partial S} \mu(a, x) f(x).$$

Using this also for b and subtracting gives

$$\begin{aligned} |f(a) - f(b)| &= \left| \sum_{x \in \partial S} (\mu(a, x) - \mu(b, x)) f(x) \right| \\ &\leq \sum_x |\mu(a, x) - \mu(b, x)| f(x) \\ &\leq \sum_x \epsilon(S) \mu(a, x) f(x) = \epsilon(S) f(a). \end{aligned}$$

Taking $S \rightarrow G$ and using the assumption we get that $f(a) = f(b)$. Since a and b were arbitrary, the function is constant. \square

Lemma 2. *Let R_n be a Markov chain on some G . Let $A \subset B$ be two finite subsets of vertices and let $a, b \in A$. Then*

$$\epsilon(B) \leq \epsilon(A).$$

Proof. Let $x \in \partial B$. Using the strong Markov property at the exit time from A we may write

$$\mu_B(a, x) = \sum_{y \in \partial A} \mu_A(a, y) \mu_B(y, x).$$

Writing the analogous equality for b and subtracting we get

$$\begin{aligned} |\mu_B(a, x) - \mu_B(b, x)| &\leq \sum_{y \in \partial A} |\mu_A(a, y) - \mu_A(b, y)| \mu_B(y, x) \\ &\leq \epsilon(A) \sum_{y \in \partial A} \mu_A(a, y) \mu_B(y, x) = \epsilon(A) \mu_B(a, x). \end{aligned}$$

Taking the maximum over $x \in \partial B$ proves the lemma. \square

Proof of the theorem. Assume such a function does not exist. By lemma 1 we know that for all a and b $\epsilon(B(a, r); a, b) \rightarrow 0$ where $B(a, r)$ is the directed ball of radius r centered at a , or in other words, all vertices which can be reached from a by a path of length less than r which the walker may take with positive probability (we use here strong connectedness to claim that $B(a, r) \rightarrow G$). Let $\delta > 0$ be some parameter. Since R_n has finite support there exists some r_0 sufficiently large such that $\epsilon(B(a, s); a, b) < \delta$ for all $s > r_0$ and all $a \rightarrow b$ (the value of a does not matter, from transitivity).

Let now $x \in \partial B(a, r)$. Let γ be a geodesic from a to x i.e. $a = \gamma_0 \rightarrow \gamma_1 \rightarrow \dots \rightarrow \gamma_r = x$. Write

$$\mu_{B(a, r)}(a, x) = \prod_{i=0}^{r-1} \frac{\mu_{B(a, r)}(\gamma_i, x)}{\mu_{B(a, r)}(\gamma_{i+1}, x)}.$$

(with the convention that $\mu_{B(a, r)}(x, x) = 1$). Write also

$$\left| \frac{\mu_{B(a, r)}(\gamma_i, x)}{\mu_{B(a, r)}(\gamma_{i+1}, x)} - 1 \right| \leq \epsilon(B(a, r); \gamma_i, \gamma_{i+1}). \quad (1)$$

Now, $B(\gamma_i, r - i) \subset B(a, r)$ and hence by lemma 2

$$\epsilon(B(a, r); \gamma_i, \gamma_{i+1}) \leq \epsilon(B(\gamma_i, r - i); \gamma_i, \gamma_{i+1}). \quad (2)$$

Let φ be an automorphism of G taking γ_i to a and preserving the Markov chain R , and let $s_i = \varphi(\gamma_{i+1})$. Then

$$\epsilon(B(\gamma_i, r - i); \gamma_i, \gamma_{i+1}) = \epsilon(B(a, r - i); a, s_i). \quad (3)$$

Finally, if $i < r - r_0$ then

$$\left| \frac{\mu_{B(a, r)}(\gamma_i, x)}{\mu_{B(a, r)}(\gamma_{i+1}, x)} - 1 \right| \stackrel{(1,2,3)}{\leq} \epsilon(B(a, r - i); a, s_i) \leq \delta.$$

For $i \geq r - r_0$ we estimate simply $\mu_{B(a, r)}(\gamma_{i+1}, x) \geq p \mu_{B(a, r)}(\gamma_i, x)$ where p is the minimum non-zero probability that a step of R may take ($p > 0$ because R is finitely supported and invariant). We get

$$\mu_{B(a, r)}(x) \geq (1 - \delta)^{r - r_0} p^{r_0}.$$

Since $x \in \partial B(a, r)$ was arbitrary we get that

$$|\partial B(a, r)| \leq (1 - \delta)^{r_0 - r} p^{-r_0}.$$

Summing over r gives

$$|B(a, s)| = \sum_{r=0}^{s-1} |\partial B(1, r)| \leq \sum_{r=0}^{s-1} (1 - \delta)^{r_0 - r} p^{-r_0} \leq \frac{p^{-r_0}}{\delta} (1 - \delta)^{-s}.$$

Since δ was an arbitrary positive number, we reach a contradiction to our assumption that directed balls grow exponentially. \square

Acknowledgements. While performing this research, G.A. was supported by the Israel Science Foundation grant #575/16 and by GIF grant #I-1363-304.6/2016. G.K. was supported by the Israel Science Foundation grant #1369/15 and by the Jesselson Foundation.

REFERENCES

- [1] Philippe Bougerol and Laure Élie, *Existence of positive harmonic functions on groups and on covering manifolds*. Ann. Inst. H. Poincaré Probab. Statist. 31:1 (1995), 59–80. Available at: numdam.org/31_1_59_0
- [2] Gustave Choquet and Jacques Deny, *Sur l'équation de convolution $\mu = \mu * \sigma$* [French: On the convolution equation $\mu = \mu * \sigma$]. C. R. Hebd. Acad. Sci. 250 (1960), 799–801. Available at: gallica.bnf.fr/f825
- [3] Waldemar Hebisch and Laurent Saloff-Coste, *Gaussian estimates for Markov chains and random walks on groups*. Ann. Probab. 21:2 (1993), 673–709. jstor.org/2244671
- [4] Gregory A. Margulis, *Positive harmonic functions on nilpotent groups*. Dokl. Akad. Nauk SSSR 166 (1966), 1054–1057 (Russian); English translation in: Soviet Math. Dokl. 7 (1966), 241–244.

BAR-ILAN UNIVERSITY, RAMAT GAN 52900, ISRAEL.
E-mail address: gidi.amir@gmail.com

THE WEIZMANN INSTITUTE OF SCIENCE, REHOVOT 76100, ISRAEL
E-mail address: gady.kozma@weizmann.ac.il