The Thirring quantum cellular automaton

Alessandro Bisio,* Giacomo Mauro D'Ariano,† Paolo Perinotti,‡ and Alessandro Tosini§

Dipartimento di Fisica dell'Università di Pavia, via Bassi 6, 27100 Pavia and

Istituto Nazionale di Fisica Nucleare, Gruppo IV, via Bassi 6, 27100 Pavia

We analytically diagonalize a discrete-time on-site interacting fermionic cellular automaton in the two-particle sector. Important features of the solutions sensibly differ from those of analogous Hamiltonian models. In particular, we found a wider variety of scattering processes, we have bound states for every value of the total momentum, and there exist bound states also in the free case, where the coupling constant is null.

PACS numbers: 71.10.Fd, 03.67.Ac

Quantum cellular automata and quantum walks constitute an increasingly attractive arena for research in many body systems [1–3], quantum computation [4–7], and foundations of quantum field theory [8–12].

The notion of quantum cellular automaton introduced by Feynman [13] as a universal quantum simulator, was mathematically formalized in Ref. [14, 15]. In the case of non-interacting theories the evolution of field operators is linear, and its simulation through quantum cellular automata reduces to simulation of a single particle through a quantum walk [16–18]. The interacting case is largely unexplored, and was mainly approached by extending the quantum walk formalism, introducing decoherence [19], or a classical external field [10, 20–22]. A notable exception is Ref. [23], where bound states in interacting quantum walks are studied.

In the present paper we study a one-dimensional massive Fermionic cellular automaton with a four-Fermion on-site interaction. The main result consists in the complete analytical solution in the two-particle sector. The linear part of the evolution corresponds to a onedimensional Dirac walk [10], with an interaction having the most general on-site, number-preserving form. The same kind of interaction characterizes the most studied integrable quantum systems [24–27] such as Hubbard's [28] and Thirring's [29] models. For this reason we call the present model Thirring quantum cellular automaton.

Despite the similarities, the present cellular automaton differs from the above models mainly in the discreteness of time evolution. This feature produces non-trivial differences in the dynamical solutions of the model, in particular a wider spectrum of scattering states, and the existence of bound states for every value of the total momentum. As a consequence of the departure of the present discrete-time evolution from the usual Hamiltonian paradigm, we are not allowed to borrow the common Bethe ansatz technique straightforwardly.

We start defining a quantum walk for interacting particles on the lattice \mathbb{Z} , assuming the particle statistics to be Fermionic. First we introduce the walk W for a free two-component Fermionic field ψ defined at any lattice point $x \in \mathbb{Z}$ and at any discrete time $t \in \mathbb{Z}$

$$\begin{split} \psi(x,t+1) &= W\psi(x,t), \qquad \psi(x,t) = \begin{pmatrix} \psi_{\uparrow}(x,t) \\ \psi_{\downarrow}(x,t) \end{pmatrix} \\ W &= \begin{pmatrix} \nu T_x^{\dagger} & -i\mu \\ -i\mu & \nu T_x \end{pmatrix}, \quad \nu,\mu > 0, \ \nu^2 + \mu^2 = 1, \end{split}$$

where T_x is the translation operator $T_x\phi(x) = \phi(x+1)$ and ψ_{\uparrow} and ψ_{\downarrow} denote the two components of the field. In the one-particle sector the above walk is a unitary operator W over the Hilbert space $\mathcal{H} = \mathbb{C}^2 \otimes \ell^2(\mathbb{Z})$ for which we will use the factorized orthonormal basis $|a\rangle|x\rangle$, with $a \in \{\uparrow, \downarrow\}$.

Notice that the walk evolution is local, with the field at time t and at site x depending only on the field at sites $x \pm 1$ at time t - 1 (first-neighbouring scheme). Moreover, since W commutes with translations along the lattice, the walk can be diagonalized in the momentum space. In the Fourier representation the operator W is expressed in terms of the momentum $p \in (-\pi, \pi]$ $(|p\rangle := (2\pi)^{-1/2} \sum_{x} e^{-ipx} |x\rangle)$ as follows

$$W = \int dp W(p) \otimes |p\rangle \langle p|, \quad W(p) = \begin{pmatrix} \nu e^{ip} & -i\mu \\ -i\mu & \nu e^{-ip} \end{pmatrix}$$
$$W(p) \mathbf{v}_p^s = e^{-is\omega(p)} \mathbf{v}_p^s, \quad \mathbf{v}_p^s := \frac{1}{|N_s|} \begin{pmatrix} -i\mu \\ g_s(p) \end{pmatrix}, \quad (1)$$
$$\omega(p) := \operatorname{Arccos}(\nu \cos p), \qquad s \in \{+, -\},$$

with $g_s(p) = -i(s\sin\omega(p) + \nu\sin p), |N_s|^2 = \mu^2 + |g_s|^2.$

The function $\omega(p)$ is the walk dispersion relation. In Ref. [30] it is shown that for small p this discrete dynamics recovers that of a free Dirac field of mass μ .

A *N*-particle walk can then be described taking $\mathcal{H}_N = \mathcal{H}^{\otimes N}$ as the Hilbert space of the system and $W_N := W^{\otimes N}$ as the operator providing the evolution. Within this scenario we introduce a coupling between particles defining the dynamical time step via a unitary operator of the form $W_N V_{int}$. Here we consider the following interacting term

$$V_{int} := V(\chi) = e^{i\chi n_{\uparrow}(x)n_{\downarrow}(x)}.$$

^{*} alessandro.bisio@unipv.it

[†] dariano@unipv.it

[‡] paolo.perinotti@unipv.it

[§] alessandro.tosini@unipv.it

where $n_a(x) = \psi_a^{\dagger}(x)\psi_a(x)$ is the number operator at site x and with internal state $a \in \{\uparrow,\downarrow\}$. This corresponds to an on-site coupling, namely the action of $V(\chi)$ is non trivial if and only if two Fermions lie at the same site of the lattice. Moreover, since $V(\chi)$ commutes with the total number operator $n = \sum_x (\psi_{\uparrow}^{\dagger}(x)\psi_{\uparrow}(x)+\psi_{\downarrow}^{\dagger}(x)\psi_{\downarrow}(x))$, the dynamics preserves the number of particles. This is the most general possible coupling with the above properties for the case of a Fermionic walk [31]. In this paper we restrict to the N = 2 case and therefore we focus on the walk

$$U_2 := W_2 V_2(\chi), \qquad V_2(\chi) := e^{i\chi \delta_{y,0}(1 - \delta_{\alpha_1,\alpha_2})}, \qquad (2)$$

which is written in the center of mass basis $|a_1, a_2\rangle |y\rangle |w\rangle$ for the two-particles Hilbert space $\mathcal{H}_2 = \mathbb{C}^4 \otimes \ell^2(\mathbb{Z})$, with $a_1, a_2 \in \{\uparrow, \downarrow\}, y = x_1 - x_2$ and $w = x_1 + x_2$ the relative and the center of mass coordinate respectively. Denoting by $k = \frac{1}{2}(p_1 - p_2)$ the (half) relative momentum and by $p = \frac{1}{2}(p_1 + p_2)$ the (half) total momentum, the free walk W_2 in the momentum representation is written as follows

$$W_{2} = \int dk \, dp \, W_{2}(p,k) \otimes |k\rangle \langle k| \otimes |p\rangle \langle p|, \qquad (3)$$
$$W_{2}(p,k) \mathbf{v}^{sr} = e^{-i\omega_{sr}(p,k)} \mathbf{v}^{sr}, \quad \mathbf{v}^{sr} := \mathbf{v}^{s} \otimes \mathbf{v}^{r}$$

$$\omega_{sr}(p,k) := s\omega(p+k) + r\omega(p-k), \quad s, r \in \{+, -\},$$

where the eigenvectors of $W_2(p,k) := W(p+k) \otimes W(p-k)$ are easily computed as the tensor product of the singleparticle ones in Eq. (1).

Since the interacting dynamics U_2 commutes with translation T_w in the center of mass coordinate w, it is convenient to write the walk in the hybrid basis $|a_1, a_2\rangle |y\rangle |p\rangle$, in the following block-diagonal form

$$\begin{split} U_2 &= \int \mathrm{d}p \, U_2(\chi,p) \otimes |p\rangle \langle p|, \quad U_2(\chi,p) := W_2(p) \tilde{V}_2(\chi), \\ W_2(p) &:= \mu \nu \begin{pmatrix} \frac{\nu}{\mu} e^{i2p} & -ie^{ip}T_y & -ie^{ip}T_y^{\dagger} & -\frac{\mu}{\nu} \\ -ie^{ip}T_y & \frac{\nu}{\mu}T_y^2 & -\frac{\mu}{\nu} & -ie^{-ip}T_y \\ -ie^{ip}T_y^{\dagger} & -\frac{\mu}{\nu} & \frac{\nu}{\mu}T_y^{\dagger 2} & -ie^{-ip}T_y^{\dagger} \\ -\frac{\mu}{\nu} & -ie^{-ip}T_y & -ie^{-ip}T_y^{\dagger} & \frac{\nu}{\mu}e^{-i2p} \end{pmatrix} \\ \tilde{V}_2(\chi) &:= \begin{pmatrix} I & 0 & 0 & 0 \\ 0 & e^{i\chi\delta_{y,0}}I & 0 & 0 \\ 0 & 0 & 0 & I \end{pmatrix}, \end{split}$$

with T_y the translation in the relative coordinate y.

The first step is to solve the linear difference equation

$$U_{2}(\chi, p) \mathbf{f}_{p,\omega,\chi} = e^{i\omega} \mathbf{f}_{p,\omega,\chi}, \qquad (4)$$
$$\mathbf{f}_{p,\omega,\chi} : \mathbb{Z} \to \mathbb{C}^{4}, \quad \omega \in \mathbb{C},$$

for any possible value of χ and p. Among all the possible solutions of Equation (4) we will then choose those ones which are eigenvectors (or generalized eigenvectors) of $U_2(\chi, p)$ considered as an operator on the Hilbert space $\mathbb{C}^4 \otimes \ell^2(\mathbb{Z})$. Since the interacting particles are Fermions,



FIG. 1. (Colors online) Top: k_R and k_I represent the real and the imaginary part of the relative momentum k in the twofermion state. The highlighted regions collect the values of $k \in \mathbb{C}$ providing a real value of the quasi-energy ω . Bottom: the disjoint subregions of the unit circle are the images under $k \mapsto e^{i\omega_{rs}(p,k)}$ of the disjoint regions in the top figure, for fixed values of the total momentum p = 0.55 and mass $\mu = 0.8$. Ω_f coincides with the continuous spectrum of $U_2(\chi, p)$ (see Eq. (2)). The discrete spectrum lies in the other regions, and for a fixed value of the coupling constant χ it consists of a single point. Varying the value of χ the unit circle is covered, and the boundary points of the arcs depend on p.

we are only interested in the solutions that are antisymmetric under the exchange of the two particles, i.e.

$$\mathbf{f}_{p,\omega,\chi}(y) = -E\mathbf{f}_{p,\omega,\chi}(-y),$$

where E is represented as $E = \frac{1}{2} \sum_{i=0}^{3} \sigma_i \otimes \sigma_i$ (with $\sigma_0 = I$, and σ_i , i = 1, 2, 3, the Pauli matrices).

In the following, in order to lighten the notation, we will omit the explicit dependence of the solutions from p, ω, χ and we will write $\mathbf{f}(y)$ for $\mathbf{f}_{p,\omega,\chi}(y)$.

Since the interacting term acts only at the origin, for y > 0 Eq. (4) becomes a linear recurrence relation with constant coefficient whose most general solutions [32] are of two forms: $\mathbf{f}_{\infty}(y)$ or $\mathbf{f}(y)$, given by

$$\mathbf{f}_{\infty}(y) = (\zeta_{\infty}, 0, 0, \zeta'_{\infty})^T \delta_{y,1} \quad \zeta_{\infty}, \zeta'_{\infty} \in \mathbb{C} \quad y > 0,$$
(5)

$$\mathbf{f}(y) = \sum_{s,r=\pm} \int_{\mathsf{S}} \mathrm{d}k \, e^{-iky} g_{\omega}(s,r,k) \mathbf{v}_k^{sr} \qquad y > 0 \tag{6}$$

$$k = k_R + ik_I, \ \mathsf{S} := \{k \in \mathbb{C} \mid k_R \in (-\pi, \pi]\},\$$
$$g_{\omega}(s, r, k) \in \mathbb{C} \text{ s.t. } e^{-i\omega} \neq e^{-i\omega_{sr}(p, k)} \Rightarrow g_{\omega}(s, r, k) = 0,$$

where the function $\omega_{rs}(p,k)$ and the vectors $\mathbf{v}_k^{sr} := \mathbf{v}_{p,k}^{sr}$ have been defined in Eq. (3), and for complex argument Arccos is the principal value of the arccosine function. Let us first consider the functions given by Eq. (5). A necessary condition for a function obeying Eq. (5) to be a (proper or improper) eigenvector of $U_2(\chi, p)$ is that $\omega_{sr}(p,k) \in \mathbb{R}$. In order to analyse this condition, it is useful to introduce the following sets (see Fig. 1):

$$\Gamma_f := \{k \in \mathsf{S} | k_I = 0\},
\Gamma_z := \{k \in \mathsf{S} | k_R = z\frac{\pi}{2}\}, \quad z = 0, \pm 1, 2,$$
(7)

$$\Omega_f^{sr} := \{ \exp(-i\omega_{sr}(p,k)) \mid k \in \Gamma_f \},$$

$$\Omega_f^{sr} = \{ \exp(-i\omega_{sr}(p,k)) \mid k \in \Gamma_f \},$$
(8)

$$\Omega_z^{sr} := \{ \exp(-i\omega_{sr}(p,k)) \mid k \in \Gamma_z, \ (-1)^z = sr \}.$$

Reminding that $\omega(x+\pi) = \pi - \omega(x)$, one can verify that

$$\Omega_{f}^{++} = \Omega_{f}^{--}, \quad \Omega_{f}^{+-} = \Omega_{f}^{-+}, \quad \Omega_{0}^{++} = \Omega_{2}^{--}, \\ \Omega_{\pi}^{++} = \Omega_{0}^{--}, \quad \Omega_{-1}^{+-} = \Omega_{1}^{-+}, \quad \Omega_{1}^{+-} = \Omega_{-1}^{-+}.$$
(9)

The following technical result, proved in Appendix A, marks the first important difference from the Hamiltonian integrable models, relying in the degeneracy of twoparticle levels. The degeneracy in the Hamiltonian case is two, corresponding to the intuitive one-dimensional picture where either a classical elastic bounce or a quantum tunnelling where the particles cross each other occur. On the other hand, in the discrete case the degeneration is four, allowing also for scattering events where hopping to a distant region in the Brillouin zone can occur. This phenomenon is due to periodicity of the quasi-energy spectrum, which is where the failure of the Bethe ansatz lurks.

Lemma 1 Let $\omega_{sr}(p,k)$ be defined as in Eq. (3) and let us assume $p \neq z\frac{\pi}{2}$ ($z \in \mathbb{Z}$). Then we have:

1.
$$\omega_{\pm\pm}(p,k) \in \mathbb{R} \implies k \in \Gamma_{\rm f} \cup \Gamma_0 \cup \Gamma_2$$
$$\omega_{\pm\mp}(p,k) \in \mathbb{R} \implies k \in \Gamma_{\rm f} \cup \Gamma_{-1} \cup \Gamma_1$$
(10)

- The six sets Ω⁺⁺_f, Ω⁺⁻_f, Ω⁺⁺₀, Ω⁺⁺₂, Ω⁺⁻₋₁, Ω⁺⁻₁ are disjoint and their union is the whole unit circle except for the points e^{±i2p}.
- 3. For any $\omega \in \mathbb{R}$ such that $e^{-i\omega} \neq e^{\pm i2p}$ the equation $e^{-i\omega} = e^{-i\omega_{sr}(p,k)}$ has four distinct solutions. If the triple (+,+,k) is a solution then also (+,+,-k), $(-,-,\pi-k)$ and $(-,-,k-\pi)$ are solutions. If the triple (+,-,k) is a solution then (-,+,-k), $(+,-,\pi-k)$ and $(-,+,k-\pi)$ are solutions.

By Lemma 1, Eq. (5) yields two classes of solutions:

$$\begin{aligned} \mathbf{f}_{k}^{\pm}(y) & \text{with } k \in \Gamma_{f} \cup \Gamma_{0} \cup \Gamma_{2}, \\ \mathbf{f}_{k}^{-}(y) & \text{with } k \in \Gamma_{f} \cup \Gamma_{-1} \cup \Gamma_{1}, \end{aligned}$$
(11)
$$\mathbf{f}_{k}^{\pm}(y) = \begin{cases} [\alpha_{\pm} \mathbf{v}_{k}^{\pm\pm} - (-1)^{y} \delta_{\pm} \mathbf{v}_{k-\pi}^{-\mp}] e^{-iyk} + \\ -[\beta_{\pm} \mathbf{v}_{-k}^{\pm\pm} - (-1)^{y} \gamma_{\pm} \mathbf{v}_{\pi-k}^{\pm-}] e^{iyk} & y > 0 \\ (0, \eta_{\pm}, -\eta_{\pm}, 0)^{T} & y = 0 \\ \text{antisymmetrized} & y < 0, \end{cases} \end{aligned}$$

where $\alpha_{\pm}, \beta_{\pm}, \ldots$, are complex coefficients which depend on p, k, m, χ . We now determine these coefficients by requiring that Eq. (4) is satisfied. Because of the locality of the evolution, this constraint needs to be verified only for y = 0, 1, 2. A tedious albeit straightforward calculation allows one to bring Eq. (11) into the following form after suitable reparametrization

$$\mathbf{f}_{k}^{\pm}(y) = c_{1}\mathbf{f}_{k}^{\pm,f}(y) + c_{2}\mathbf{f}_{k}^{\pm,i}(y), \qquad (12)$$

$$\mathbf{f}_{k}^{\pm,j}(y) = [\mathbf{v}_{k}^{\pm,\pm} + (-1)^{y} \mathbf{v}_{k}^{\pm,\pm}]e^{-iyk} + \\ - [\mathbf{v}_{-k}^{\pm,\pm} + (-1)^{y} \mathbf{v}_{\pi-k}^{\pm,-}]e^{iyk}, \\ \mathbf{f}_{k}^{\pm,i}(y) = \begin{cases} e^{-i\delta_{y,0}\chi} \{ [\mathbf{v}_{k}^{\pm,\pm} - (-1)^{y} \mathbf{v}_{k-\pi}^{\pm,\pm}]e^{-iyk} + \\ -T_{\pm} [\mathbf{v}_{-k}^{\pm,\pm} - (-1)^{y} \mathbf{v}_{\pi-k}^{\pm,\pm}]e^{iyk} \} \\ \text{antisymmetrized} & y < 0, \end{cases}$$

$$T_{\pm} := \frac{g_{\pm}(p+k) + e^{-i\chi}g_{\pm}(p-k)}{g_{\pm}(p-k) + e^{-i\chi}g_{\pm}(p+k)}, \qquad c_1, c_2 \in \mathbb{C}.$$

The next step of the analysis is to identify, among the set of functions of Eq. (12), those ones which correspond to eigenvectors or generalized eigenvectors of $U_2(\chi, p)$.

For $k \in \Gamma_f$, Eq. (12) gives the generalized eigenvector of $U_2(\chi, p)$ corresponding to the continuous spectrum $\sigma_c = \Omega_f^{++} \cup \Omega_f^{+-}$. Since $U_2 - W_2$ is a finite rank operator, the continuous spectrum of the two-particle interacting case is the same as that of the free walk (see Theorem IV 5.35 of Ref.[33]). From Eq. (12) we have that the solutions of the kind $\mathbf{f}_k^{\pm,f}$ are generalized eigenvectors of the free theory which are also generalized eigenvectors of the interacting theory. This is easily understood since $\mathbf{f}_k^{\pm,f}(0) = \mathbf{0}$, and therefore those eigensolutions do not feel the presence of the interaction—which is localized at y = 0. On the other hand, we can interpret the solution of the kind $\mathbf{f}_k^{\pm,i}$ as a scattering of plane waves with the T_{\pm} playing the role of transmission coefficients.

For $k \notin \Gamma_f$, necessary conditions for \mathbf{f}_k^{\pm} to be a (proper or generalized) eigenvector of $U_2(\chi, p)$ are that $k_I = \Im(k) < 0, c_1 = 0, c_2 \neq 0$ and $T_{\pm} = 0$ (otherwise \mathbf{f}_k^{\pm} is exponentially divergent). In appendix B we prove the following result:

Lemma 2 Let T_{\pm} defined as in Eq. (12) and let us assume $p \neq z\frac{\pi}{2}$. If $e^{i\chi} \notin \{e^{\pm i2p}, 1, -1\}$, then there exists a unique $k \in \Gamma_0 \cup \Gamma_{-1} \cup \Gamma_1 \cup \Gamma_2$ with $k_I < 0$ such that either $T_+ = 0$ or $T_- = 0$. On the other hand, if $e^{i\chi} \in \{e^{\pm i2p}, 1, -1\}$ then $T_+ \neq 0$ and $T_- \neq 0$ for all $k \in \Gamma_0 \cup \Gamma_{-1} \cup \Gamma_1 \cup \Gamma_2$ with $k_I < 0$.

The above result tells us that, for $e^{i\chi} \notin \{e^{\pm i2p}, 1, -1\}$, the two-particles interacting evolution $U_2(\chi, p)$ has one proper eigenvector whose corresponding eigenvalue constitutes the discrete spectrum of $U_2(\chi, p)$. This eigenstate is easily interpreted as a bound state of two particles.

We now consider the functions given by Eq. (6) which lead to the antisymmetric functions

$$\mathbf{f}_{\infty}(y) = \begin{cases} (\zeta_{\infty}, 0, 0, \zeta_{\infty}')^T \delta_{y,1} & y > 0, \\ (0, \eta_{\infty}, -\eta_{\infty}, 0)^T & y = 0, \\ (-\zeta_{\infty}, 0, 0, -\zeta_{\infty}')^T \delta_{y,-1} & y < 0. \end{cases}$$
(13)

$$\mathbf{f}_{\pm\infty}(y) = \begin{cases} ie^{\pm ip} (-\frac{1\pm 1}{2}, 0, 0, -\frac{1\pm 1}{2})^T \delta_{y,1} & y > 0, \\ (0, \frac{\mu}{\nu}, -\frac{\mu}{\nu}, 0)^T & y = 0, \\ ie^{\pm ip} (\frac{1\pm 1}{2}, 0, 0, -\frac{-1\pm 1}{2})^T \delta_{y,-1} & y < 0, \end{cases}$$

$$U_2(\chi, p)\mathbf{f}_{\pm\infty} = e^{\pm i2p}\mathbf{f}_{\pm\infty} \text{ for } e^{i\chi} = e^{\pm i2p}.$$
 (14)

Eq. (14) provides the proper eigenstate of $U_2(\chi, p)$ for the cases $e^{i\chi} = e^{\pm i2p}$ which were missing in Lemma 2.

In Fig. 2 we plot the spectrum of $U_2(\chi, p)$ as a function of p for different values of the χ . We can then write, for $p \neq z\frac{\pi}{2}$, the spectral resolution of $U_2(\chi, p)$, i.e.

$$U_{2}(\chi,p) = \sum_{\substack{s=\pm, \\ j=f,i}} \int_{-\pi}^{\pi} dk \, e^{-i\omega_{+s}(p,k)} |\phi_{p,\chi}^{s,j}(k)\rangle \langle \phi_{p,\chi}^{s,j}(k)| + e^{-i\tilde{\omega}} |\varphi_{p,\chi}\rangle \langle \varphi_{p,\chi}|,$$

where we defined

$$\begin{split} \langle y | \phi_{p,\chi}^{s,j}(k) \rangle &:= N_{p,\chi,s,j,k} \, \mathbf{f}_{k}^{s,j}(y), \\ \langle y | \varphi_{p,\chi} \rangle &:= \begin{cases} M_{p,\chi,\tilde{k}} \, \mathbf{f}_{k}^{+,i}(y), & e^{i\chi} \neq e^{\pm i2p}, \ T_{+}(\tilde{k}) = 0 \\ M_{p,\chi,\tilde{k}} \, \mathbf{f}_{k}^{-,i}(y), & e^{i\chi} \neq e^{\pm i2p}, \ T_{-}(\tilde{k}) = 0 \\ M_{p}^{\pm} \, \mathbf{f}_{\pm\infty}(y), & e^{i\chi} = e^{\pm i2p}, \end{cases} \\ \tilde{\omega} &:= \begin{cases} \omega_{++}(p,\tilde{k}), & e^{i\chi} \neq e^{\pm i2p}, \ T_{+}(\tilde{k}) = 0, \\ \omega_{+-}(p,\tilde{k}), & e^{i\chi} \neq e^{\pm i2p}, \ T_{-}(\tilde{k}) = 0, \\ \pm 2p, & e^{i\chi} = e^{\pm i2p}, \end{cases} \end{split}$$

and N, M are normalization factors such that

$$\begin{aligned} \langle \phi_{p,\chi}^{s,j}(k) | \phi_{p,\chi}^{s',j'}(k') \rangle &= \delta_{s,s'} \delta_{j,j'} \delta(k-k') \\ \langle \varphi_{p,\chi} | \varphi_{p,\chi} \rangle &= 1 \end{aligned}$$



FIG. 2. Spectrum of the 2 particle automaton of Eq. (2): In red and yellow are depicted the continuous spectrum bands; in black the discrete band for different values of the coupling: $\chi_1 = -\frac{\pi}{5}, \chi_2 = -\frac{\pi}{2}, \chi_3 = -\frac{4\pi}{5}, \chi_4 = \frac{4\pi}{5}, \chi_5 = \frac{\pi}{2}, \chi_6 = \frac{\pi}{5}.$

We conclude our analysis with the discussion of the cases $p = z\frac{\pi}{2}$ starting from p = 0. We have $\omega_{\pm\pm}(0,k) = \pm 2\omega(k)$, with $\omega(k) \in (-\pi,\pi]$ and $\omega(k) \neq 0$, iff $k \in \Gamma_f \cup \Gamma_0 \cup \Gamma_2$. On the other hand $\omega_{\pm\mp}(0,k) = 0$ for all $k \in \mathbb{C}$, and thus $\omega_{\pm\mp}(0,k) \neq \omega_{\pm\pm}(0,k')$ for all values of k,k'. Therefore the previous analysis still holds for $e^{-i\omega} \neq 1$

and, by setting p = 0, the solutions \mathbf{f}_k^+ of Eq. (12) are (proper and improper) eigenvectors of $U_2(\chi, 0)$. Thus, the spectrum of $U_2(\chi, 0)$ decomposes into a continuous spectrum, which is the arc of the unit circle wich contains -1 and has $e^{\pm 2i\omega(0)}$ as extremes, and a point spectrum made of two distinct points: $e^{-i2\omega(\tilde{k})}$ (where \tilde{k} is the solution of $T_+ = 0$ when p = 0) and 1. Since $U_2(\chi, 0)$ is unitary, if $e^{-i\omega}$ belongs to the point spectrum then it is a proper eigenvalue of $U_2(\chi, 0)$. Let us denote with $P_0^$ the projection on the eigenspace of the eigenvalue 1, and by P_p^- the following projection

Now, since $\lim_{p\to 0} ||U_2(\chi, p) - U_2(\chi, 0)|| = 0$, and 1 is a separated part of the spectrum of $U_2(\chi, 0)$, then $\lim_{p\to 0} ||P_p^- - P_0^-|| = 0$ (see Theorem IV 3.16 of Ref.[33]). We have then that

$$\begin{split} P_0^- &= \sum_{\substack{n \in \mathbb{Z} \\ j=f,i}} |\psi_0^-(n,j)\rangle \langle \psi_0^-(n,j)|, \\ |\psi_0^-(n,j)\rangle &:= \int_{-\pi}^{\pi} g_n(k) |\phi_{0,\chi}^{-,j}(k)\rangle \end{split}$$

where $g_n(k)$ is an orthonormal basis for $L^2(-\pi,\pi]$. The cases $p = \pi, \pm \frac{\pi}{2}$ can be analysed in the same way. The eigenspace corresponding to the eigenvalue 1 is thus a separable Hilbert space of stationary bound states. This result marks an important departure from the behaviour of analogous Hamiltonian models. Remarkably, it occurs even in the non-interacting case $\chi = 0$.

The diagonalisation of $U_2(\chi, p)$ summarized by the following proposition

Proposition 1 Let $U_2(\chi, p)$ be defined as in Equation (3). Then its spectral resolution is

$$\begin{split} U_{2}(\chi,p) &= \\ &= \begin{cases} \sum_{\substack{s=\pm, \\ j=f,i}} U_{p,\chi}^{s,j} + e^{-i\tilde{\omega}} P_{p,\chi} & p \neq z\frac{\pi}{2}, \\ \sum_{\substack{j=f,i \\ j=f,i}} U_{z\pi,\chi}^{+,j} + P_{z\pi}^{-} + e^{-i\tilde{\omega}} P_{z\pi,\chi} & p = 0, \pi, \\ \sum_{\substack{j=f,i \\ j=f,i}} U_{\pm\frac{\pi}{2},\chi}^{-,j} - P_{\pm\frac{\pi}{2}}^{+} + e^{-i\tilde{\omega}} P_{\pm\frac{\pi}{2},\chi} & p = \pm\frac{\pi}{2}, \end{cases} \\ U_{p,\chi}^{s,j} &:= \int_{-\pi}^{\pi} dk \, e^{-i\omega_{+s}(p,k)} |\phi_{p,\chi}^{s,j}(k)\rangle \langle \phi_{p,\chi}^{s,j}(k)|, \\ P_{p,\chi} &= |\varphi_{p,\chi}\rangle \langle \varphi_{p,\chi}|. \end{split}$$

We diagonalized an on-site interacting fermionic cellular automaton in the two-particle sector. Differently from analogous Hamiltonian models, (i) bound states exist for every value of the total momentum, (ii) there are four classes of scattering solutions instead of two, (iii) the bound states exist also in the free case.

- C. M. Dawson, J. Eisert, and T. J. Osborne, Phys. Rev. Lett. 100, 130501 (2008).
- [2] B. M. Boghosian and W. Taylor, Phys. Rev. E 57, 54 (1998).
- [3] J. I. Cirac, D. Perez-Garcia, N. Schuch, and F. Verstraete, Journal of Statistical Mechanics: Theory and Experiment 2017, 083105 (2017).
- [4] A. M. Childs, Phys. Rev. Lett. **102**, 180501 (2009).
- [5] N. B. Lovett, S. Cooper, M. Everitt, M. Trevers, and V. Kendon, Phys. Rev. A 81, 042330 (2010).
- [6] A. Schreiber, K. N. Cassemiro, V. Potocek, A. Gabris, P. J. Mosley, E. Andersson, I. Jex, and C. Silberhorn, Phys. Rev. Lett. **104**, 050502 (2010).
- [7] L. Sansoni, F. Sciarrino, G. Vallone, P. Mataloni, A. Crespi, R. Ramponi, and R. Osellame, Phys. Rev. Lett. 108, 010502 (2012).
- [8] I. Bialynicki-Birula, Physical Review D 49, 6920 (1994).
- [9] D. Meyer, Journal of Statistical Physics 85, 551 (1996).
- [10] A. Bisio, G. M. D'Ariano, and A. Tosini, Phys. Rev. A 88, 032301 (2013).
- [11] P. Arrighi, S. Facchini, and M. Forets, Quantum Information Processing 15, 3467 (2016).
- [12] G. M. D'Ariano and P. Perinotti, Phys. Rev. A 90, 062106 (2014).
- [13] R. Feynman, International journal of theoretical physics 21, 467 (1982).
- [14] B. Schumacher and R. Werner, Arxiv preprint quantph/0405174 (2004).
- [15] D. Gross, V. Nesme, H. Vogts, and R. Werner, Communications in Mathematical Physics , 1 (2012).
- [16] A. Ambainis, E. Bach, A. Nayak, A. Vishwanath, and J. Watrous, in *Proceedings of the thirty-third annual* ACM symposium on Theory of computing (ACM, 2001) pp. 37–49.
- [17] Y. Aharonov, L. Davidovich, and N. Zagury, Phys. Rev. A 48, 1687 (1993).
- [18] G. Grossing and A. Zeilinger, Complex Systems 2, 197 (1988).
- [19] Y. Shikano, K. Chisaki, E. Segawa, and N. Konno, Phys. Rev. A 81, 062129 (2010).
- [20] P. Arnault and F. Debbasch, Phys. Rev. A 93, 052301 (2016).
- [21] P. Arnault, G. Di Molfetta, M. Brachet, and F. Debbasch, Phys. Rev. A 94, 012335 (2016).
- [22] D. A. Meyer, Phys. Rev. E 55, 5261 (1997).
- [23] A. Ahlbrecht, A. Alberti, D. Meschede, V. B. Scholz, A. H. Werner, and R. F. Werner, New Journal of Physics 14, 073050 (2012).
- [24] E. H. Lieb and F. Y. Wu, Physical Review Letters 20, 1445 (1968).
- [25] S. Coleman, Phys. Rev. D 11, 2088 (1975).
- [26] V. E. Korepin, Theoretical and Mathematical Physics 41, 953 (1979).
- [27] F. H. Essler, H. Frahm, F. Göhmann, A. Klümper, and V. E. Korepin, *The one-dimensional Hubbard model* (Cambridge University Press, 2005).
- [28] Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences 276, 238 (1963).
- [29] W. E. Thirring, Annals of Physics 3, 91 (1958).
- [30] A. Bisio, G. M. D'Ariano, and A. Tosini, Annals of

Physics **354**, 244 (2015).

- [31] S. Östlund and E. Mele, Phys. Rev. B 44, 12413 (1991).
- [32] P. Cull, M. Flahive, and R. Robson, *Difference equa*tions: from rabbits to chaos (Springer Science & Business Media, 2005).
- [33] T. Kato, Perturbation theory for linear operators, Vol. 132 (Springer Science & Business Media, 2013).

Appendix A: Proof of Lemma 1

1. Proof of item 1

Let us define $\hat{\omega}_{\pm} + i\tilde{\omega}_{\pm} := \omega(p \pm k)$. Since $\omega(z^*) = \omega^*(z)$, we have that both $\hat{\omega}_{\pm}$ and $\tilde{\omega}_{\pm}$ are real. Then we have

$$\Im(\omega_{rs}(p,k)) = 0 \iff r\tilde{\omega}_{+} = -s\tilde{\omega}_{-} \implies \cosh\tilde{\omega}_{+} = \cosh\tilde{\omega}_{-} =: \cosh\tilde{\omega}.$$
(A1)

Reminding that $\cos \omega (p \pm k) = \nu \cos(p \pm k)$, Eq. (A1) implies that

$$\cos^2 \hat{\omega}_{\pm} \cosh^2 \tilde{\omega} = \nu^2 \cos^2(p \pm k_R) \cosh^2 k_I$$

$$\sin^2 \hat{\omega}_{\pm} \sinh^2 \tilde{\omega} = \nu^2 \sin^2(p \pm k_R) \sinh^2 k_I.$$
 (A2)

From the above relations we find that

$$\cos^2(p \pm k_R) \frac{\cosh^2 k_I}{\cosh^2 \tilde{\omega}} + \sin^2(p \pm k) \frac{\sinh^2 k_I}{\sinh^2 \tilde{\omega}} = \frac{1}{\nu^2}$$
(A3)

which gives

$$\left[\sin^2(p+k_R) - \sin^2(p-k_R)\right] \left(\frac{\sinh^2 k_I}{\sinh^2 \tilde{\omega}} - \frac{\cosh^2 k_I}{\cosh^2 \tilde{\omega}}\right) = 0$$

Now, since $\frac{\sinh^2 k_I}{\sinh^2 \tilde{\omega}} - \frac{\cosh^2 k_I}{\cosh^2 \tilde{\omega}} = 0$ implies $\frac{\sinh^2 k_I}{\sinh^2 \tilde{\omega}} = \frac{\cosh^2 k_I}{\cosh^2 \tilde{\omega}} = 1$, which is not compatible with Eq. (A3), it must be $\sin^2(p+k_R) = \sin^2(p-k_R)$, which gives

$$k_R = \frac{z}{2}\pi \lor p = \frac{z}{2}\pi, \qquad z \in \mathbb{Z}.$$

By explicit computation one obtains

$$\Im(\omega_{\pm\pm}(p,k)) = 0 \land k_I \neq 0 \implies k_R = 0, \pi \lor p = \pm \frac{\pi}{2},$$

$$\Im(\omega_{\pm\mp}(p,k)) = 0 \land k_I \neq 0 \implies k_R = \pm \frac{\pi}{2}, \lor p = 0, \pi,$$

which proves the first item of Lemma 1.

2. Proof of item 2

Let us consider the case in which $p \in (0, \frac{\pi}{2})$. The function $k \mapsto \omega_{++}(p, k)$ is smooth and periodic with period 2π and therefore it ranges between its maximum and minimal values. The maximum and minimum values are found by setting $\partial_k \omega_{++}(p, k) = 0$. By explicit computation one obtains

$$\frac{\sin(p+k)}{\sqrt{1-\nu^2\cos^2(p+k)}} = \frac{\sin(p-k)}{\sqrt{1-\nu^2\cos^2(p-k)}},$$

which implies, for $p \neq z\frac{\pi}{2}$, that $k = 0, \pi$. we have than that $\omega_{++}(p,k)$ ranges between $2\omega(p)$ and $2\pi - 2\omega(p)$. By noticing that $\omega_{++}(p,\frac{\pi}{2}) = \pi$ we have that Ω_f^{++} is the arc which connects $e^{i2\omega(p)}$ and $e^{-i2\omega(p)}$ and which includes -1 (see Fig. 1). With the same procedure we find that Ω_f^{+-} is the arc connecting $e^{i(2\omega(p+\frac{\pi}{2})-\pi)}$ and $e^{-i(2\omega(p+\frac{\pi}{2})-\pi)}$ which includes 1 (see Fig. 1). We now verify that Ω_f^{++} and Ω_f^{+-} are disjoint. Since $\omega(p) < \frac{\pi}{2}$, $\omega(p+\frac{\pi}{2}) > \frac{\pi}{2}$ we have $e^{-i\omega} \in \Omega_f^{++}$ iff $\omega \mod 2\pi \in (-\pi, -2\omega(p)] \cup [2\omega(p), \pi]$ and $e^{-i\omega} \in \Omega_f^{+-}$ iff $\omega \mod 2\pi \in [\pi - 2\omega(p+\frac{\pi}{2}), 2\omega(p+\frac{\pi}{2}) - \pi,]$. Then, from the inequality $|\frac{\mathrm{d}}{\mathrm{d}x}\omega(x)| < 1$, $\forall x \in \mathbb{R}$, we have

$$\omega(p+\frac{\pi}{2}) - \omega(p) = \int_p^{p+\frac{\pi}{2}} \frac{\mathrm{d}}{\mathrm{d}x} \omega(x) < \int_p^{p+\frac{\pi}{2}} \frac{\mathrm{d}}{\mathrm{d}x} < \frac{\pi}{2},$$

which implies that the sets $(-\pi, -2\omega(p)] \cup [2\omega(p), \pi]$ and $[\pi - 2\omega(p + \frac{\pi}{2}), 2\omega(p + \frac{\pi}{2}) - \pi]$ are disjoint.

Let us now consider the set Ω_0^{++} . For $\pi \neq 0, \pi$, the function $\mathbb{R} \ni k_I \mapsto \omega_{++}(p, ik_I) = \omega(p+ik_I) + \omega(p-ik_I)$ is smooth. Therefore, the extremal points of its range occur either in its stationary points or at its limiting values for $k_I \to \pm \infty$. By setting $\partial_{k_I} \omega_{++}(p, ik_I) = 0$ we obtain

$$\frac{\sin(p+ik_I)}{\sqrt{1-\nu^2\cos^2(p+ik_I)}} = \frac{\sin(p-ik_I)}{\sqrt{1-\nu^2\cos^2(p-ik_I)}} \Longrightarrow$$
$$\sin^2(p+ik_I) = \sin^2(p-ik_I) \Longrightarrow$$
$$\sin(p+ik_I) = \pm\sin(p-ik_I) \Longrightarrow k_I = 0,$$

where we used the hypothesis $p \neq z\frac{\pi}{2}$. When $k_I = 0$ we clearly have $\omega_{++}(p,0) = 2\omega(p)$. Let us now compute $\lim_{k_I \to +\infty} \omega_{++}(p,ik_I)$. Since $\omega_{++}(p,ik_I)$ is an even function of k_I the limit $k_I \to +\infty$ and $k_I \to -\infty$ coincide. We have

$$\begin{split} \omega_{++}(p,ik_I) &= \omega(p+ik_I) + \omega(p-ik_I) = \\ &= 2\Re\,\omega(p+ik_I) = 2\Re\operatorname{Arccos}(\nu\cos(p+ik_I)) = \\ &= 2\Re\operatorname{Arccos}(\nu\cos p\cosh k_I - i\sin p\cosh k_I) = \\ &= 2\operatorname{Arccos} 2^{-1}\left(\sqrt{(1+\cos p\cosh k_I)^2 + \sin^2 p\cosh^2 k_I} - \frac{\sqrt{(1-\cos p\cosh k_I)^2 + \sin^2 p\cosh^2 k_I}}{\sqrt{(1-\cos p\cosh k_I)^2 + \sin^2 p\cosh^2 k_I}}\right) \xrightarrow{k_I \to +\infty} \\ &\to 2\operatorname{Arccos}\cos p = 2|p|. \end{split}$$

Since we are assuming $p \in (0, \frac{\pi}{2})$ we have that

$$\frac{\mathrm{d}}{\mathrm{d}p}(\omega(p) - p) = \frac{\mathrm{d}}{\mathrm{d}p}\omega(p) - 1 < 0,$$
$$\omega(0) > 0 \text{ and } \omega(\frac{\pi}{2}) = \frac{\pi}{2},$$

which imply $\omega(p)-p > 0$ for $p \in (0, \frac{\pi}{2})$. Similarly one can show $\omega(p+\frac{\pi}{2})-\frac{\pi}{2} < p$ for $p \in (0, \frac{\pi}{2})$. From $\omega(p+\frac{\pi}{2})-\frac{\pi}{2} we have that <math>e^{-i\omega} \in \Omega_0^{++}$ iff $\omega \mod 2\pi \in (-2\omega(p),), -2p$. Moreover we have that $e^{-i\omega} \in \Omega_2^{++}$ iff $e^{i\omega} \in \Omega_0^{++}$ iff $\omega \mod 2\pi \in (2p, 2\omega(p))$. This proves that, for $p \in (0, \frac{\pi}{2}) \ \Omega_0^{++}, \ \Omega_f^{+-} \ \text{and} \ \Omega_2^{++}$ are disjoint sets (see Fig. 1). Following the same derivation it is easy to show that $e^{-i\omega} \in \Omega_1^{+-}$ iff $\omega \mod 2\pi \in (-2p, \pi - 2\omega(p+1))$

 $(\frac{\pi}{2}))$ and $e^{-i\omega} \in \Omega_{-1}^{+-}$ iff $\omega \mod 2\pi \in (2\omega(p+\frac{\pi}{2})-\pi, 2p)$ which proves item 2 of Lemma 1 for $p \in (0, \frac{\pi}{2})$ (see Fig. 1). The same line of derivation can be followed for the cases $p \in (-\frac{\pi}{2}, 0), p \in (\frac{\pi}{2}, \pi)$ and $p \in (-\pi, -\frac{\pi}{2})$ thus completing the proof.

3. Proof of item 3

Let us consider a value $e^{-i\omega} \neq e^{\pm i2p}$. From item 2 of Lemma 1 we have that the sets $\Omega_f^{++}, \Omega_f^{+-}, \Omega_0^{++}, \Omega_2^{++}, \Omega_{-1}^{+-}, \Omega_1^{+-}$ cover the whole unit circle except the points $e^{\pm i2p}$ and therefore $e^{-i\omega}$ must belong to one of those sets. We prove the thesis for the case $e^{-i\omega} \in \Omega_f^{++}$, and the remaining cases can be proved in the same way. If $e^{-i\omega} \in \Omega_f^{++}$, then there exists $k \in \Gamma_f$ such that $\omega_{++}(p,k) = \omega$ By direct computation one verify that mod 2π . also $\omega_{++}(p,-k) = \omega_{--}(p,k-\pi) = \omega_{--}(p,\pi-k)$ mod $2\pi = \omega$ In order to prove that these are the only admissible solutions we must check that $k' \neq \pm k$ implies $\omega_{++}(p,k') \neq \omega \mod 2\pi$ By contradiction let us suppose that there exists $k' \neq \pm k$ such that $\omega_{++}(p,k') = \omega$ mod 2π . This clearly implies $\omega_{++}(p,k') = \omega_{++}(p,k')$ since the range of ω_{++} is smaller than 2π . Let us consider the case 0 < k' < k. Since ω_{++} is smooth, there must exists k'' such that k' < k'' < k and $\left[\frac{\mathrm{d}}{\mathrm{d}k}\omega_{++}\right](p,k') = 0$. By direct computation one proves that this is impossible. The generalization to the cases $-k < k' < 0, \ k < k' < \pi$, $-\pi < k' < k$ is straightforward. The analysis of the cases $k' = 0, \pi$ is easily done by direct computation.

Appendix B: Proof of Lemma 2

In order to prove the lemma it is convenient to introduce the following function from the negative half line $k_I \in (-\infty, 0]$ to the unit circle S^1 :

$$G_{z}: \mathbb{R}^{-} \to S^{1}, \quad G_{z}(k_{I}) = -\frac{A_{z}(p,k)}{A_{z}^{*}(p,k)}, \qquad j = 0, 2, \pm 1,$$

$$A_{0}(p,k) = \sin(\omega(p-ik_{I})) + \nu \sin(p-ik_{I}),$$

$$A_{2}(p,k) = \sin(\omega(p-ik_{I})) - \nu \sin(p-ik_{I}),$$

$$A_{\pm 1}(p,k) = \sin(\omega(p \mp \frac{\pi}{2} - ik_{I})) + \nu \sin(p \mp \frac{\pi}{2} - ik_{I}).$$

The above function allows to study the constraint $T_{\pm}(p,k,\chi) = 0$ through the following five properties

(a)
$$k \in \Gamma_0 : T_+(p,k,\chi) = 0 \iff G_0(k_I) = e^{i\chi},$$

 $k \in \Gamma_2 : T_+(p,k,\chi) = 0 \iff G_2(k_I) = e^{i\chi},$
 $k \in \Gamma_{\pm 1} : T_-(p,k,\chi) = 0 \iff G_{\pm 1}(k_I) = e^{i\chi},$

(b)
$$G_0(0) = G_2(0) = -1, \qquad G_{\pm 1}(0) = 1,$$

(c)
$$\forall k_I, \ G_0(k_I) \neq 1, G_2(k_I) \neq 1, \ G_{\pm 1}(k_I) \neq -1,$$

(d)
$$\forall k_I, \quad \frac{d}{dk_I} G_z(k_I) \neq 0, \quad z = 0, 2, \pm 1,$$

(e)
$$\lim_{\substack{k_I \to -\infty}} G_0(k_I) = e^{i2|p|},$$

$$\lim_{\substack{k_I \to -\infty}} G_2(k_I) = e^{-i2|p|},$$

$$\left(e^{\pm i2p} \quad n \in (-\frac{\pi}{2}, \frac{\pi}{2})\right)$$

$$\lim_{k_I \to -\infty} G_{\pm 1}(k_I) = \begin{cases} e^{\pm i2p}, & p \in (-\frac{\pi}{2}, \frac{\pi}{2}) \\ e^{\pm i2p}, & p \in (-\pi, -\frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi) \end{cases}.$$

The statement of the lemma is now proved noticing that the functions $G_z(k_I)$, $z = 0, 2, \pm 1$, from the negative half line $k_I \in (-\infty, 0]$ to the unit circle S^1 , are injective and their ranges are

- Range of G_0 : the smallest arc having -1 and $e^{i2|p|}$ as its extremal points,
- Range of G_2 : the smallest arc having -1 and $e^{-i2|p|}$ as its extremal points,
- Range of $G_{\pm 1}$: the smallest arc having 1 and (depending on the value of p) $e^{\pm i 2|p|}$ as its extremal points.

Here we provide the proofs of items 1-5 for the case $k \in \Gamma_0$. The proof for the other three cases $k \in \Gamma_2$ and $k \in \Gamma_{\pm 1}$ is almost identical.

a. Proof of item (a) for $k \in \Gamma_0$

If $k \in \Gamma_0$ it is $k = ik_I$, and starting from the definition of Eq. (12) we can rewrite $T_+(p, k, \chi)$ as follows

$$T_{+} = \frac{A_{0}^{*}(p,k) + e^{-i\chi}A_{0}(p,k)}{e^{-i\chi}A_{0}^{*}(p,k) + A_{0}(p,k)}$$

Let us replace $A_0(p, k)$ with A and $T_+(p, k, \chi)$ with T_+ in order to lighten the notation. We have that $T_+ = 0 \iff A^* + e^{-i\chi}A = 0 \land e^{-i\chi}A^* + A \neq 0$. First we observe that $A \neq 0$, indeed $A = 0 \iff \sin(\omega(p - ik)) =$ $-n\sin(p - ik) \implies \sin^2(\omega(p - ik)) = n^2\sin^2(p - ik) \implies$ $\nu^2 = 1$, which is not an admissible value. Accordingly, a straightforward computation shows that $T_+ = 0 \iff$ $A^* + e^{-i\chi}A = 0 \land e^{-i\chi}A^* + A = 0$ iff $\chi = m\pi \ (m \in \mathbb{Z})$. However, $T_+(p, k, m\pi) = (-1)^m \neq 0$ and we conclude that $T_+ = 0$ if and only if $A^* + e^{-i\chi}A = 0$ which proves item (a).

b. Proof of item (b) for $k \in \Gamma_0$

Notice that $G_0(k_I) = -1 \implies A = A^*$ which implies $\Im[A] = 0 \implies \Im[A^2] = 0$ (where we replaced $A_0(p,k)$ with A in order to lighten the notation). Since we have $A^2 = 1 - \nu^2 + 2\nu \sin(p - ik_I)A$ the condition $\Im[A] = 0 \land \Im[A^2] = 0$ implies $\Im \sin(p - ik_I) = 0$ that is $\cos(p)\sinh(k_I) = 0$ and then $k_I = 0 \lor p = \frac{\pi}{2} + m\pi$ $(m \in \mathbb{Z})$ which proves item (b).

c. Proof of item (c) for $k \in \Gamma_0$

We have $G_0(k_I) = 1 \implies A = -A^*$ which implies $\Re[A] = 0 \implies \Im[A^2] = 0$ (where we replaced $A_0(p,k)$) with A in order to lighten the notation). Since it is $A^2 = 1 - \nu^2 + 2\nu \sin(p - ik_I)A$ the condition $\Re[A] = 0 \land \Im[A^2] = 0$ implies $\Re \sin(p - ik_I) = 0$ that is $\sin(p) \cosh(k_I) = 0$. Since the last equality is satisfied only for $p = m\pi$ $(m \in \mathbb{Z})$, which are not admissible values of p, item (c) is proved.

d. Proof of item (d) for $k \in \Gamma_0$

We prove that $\frac{d}{dk_I}G_0(k_I) = 0 \implies p = m\frac{\pi}{2} \ (m \in \mathbb{Z})$, which are not admissible values of p. Again we replace $A_0(p,k)$ with A in order to lighten the notation. Consider

$$\frac{d}{dk_I}G_0(k_I) = \frac{A'A^* - A{A'}^*}{(A^*)^2}$$

where $A' = \partial_{k_I} A$ and $A^{*'} := \partial_{k_I} A^* = A'^*$. Then, reminding that $A \neq 0$ (see the proof of item (a)) and noticing that

$$A'A^* - AA'^* = -i|1 + \omega'_-|^2 \sin(\omega_- + \omega_+),$$

$$\omega_{\pm} := \omega(p \pm ik), \quad \omega'(x) := \frac{d}{dx}\omega(x),$$

(this can be verified rewriting A as $A = \sin(\omega(p-ik))(1+\omega'(p-ik)))$ one has

$$\frac{d}{dk_I}G_0(k_I) = 0 \iff 1 + \omega'_- = 0 \lor \sin(\omega_- + \omega_+) = 0.$$

Let us investigate the two possible cases. In the first case $1 + \omega'_{-} = 0$ it must be

$$\omega'_{-}^{2} = \frac{\nu^{2} \sin^{2}(p - ik_{I})}{\sin^{2}(\omega(p - ik_{I}))} = 1 \implies \nu = 1,$$

which is not admissible. Let us now consider the case $\sin(\omega_- + \omega_+) = 0$, that is $\omega_- + \omega_+ = m\pi$. We have for m even $\cos \omega_- = \cos \omega_+ \implies \sin p = 0 \lor k_I = 0$. On the other hand, if m is odd we have $\cos \omega_- = -\cos \omega_+ \implies \cos p = 0$. Item (d) is thus proved.

e. Proof of item (e) for $k \in \Gamma_0$

For convenience in the following we replace $A_0(p,k)$ with A. First we rewrite the function G_0 as follows

$$G_0(k_I) = -\frac{Z}{Z^*},$$

$$Z := -iA = e^{-i\omega(p-ik_I)} - \nu e^{i(p-ik_I)}$$

Reminding that in this case it is $k \in \Gamma_0$ it is $k = k_R + ik_I$ with $k_R = 0$, from Appendix C we have the expressions of $e^{-i\omega(p-ik_I)}$ for $k_I \to -\infty$:

$$p > 0 \implies e^{-i\omega(p-ik_I)} = \frac{1}{\nu} e^{-ip} e^{k_I},$$
$$p < 0 \implies e^{-i\omega(p-ik_I)} = \nu e^{ip} e^{-k_I} - \frac{\mu^2}{\nu} e^{-ip} e^{k_I}.$$

from which item (e) follows.

Appendix C: Asymptotic behaviour of the walk eigenvalues

The one-particle Dirac walk in momentum space is defined through the matrix valued function of Eq. (1). Since $U(p) \in \mathbb{SU}(2)$, its eigenvalues are $e^{-i\omega(p)}$ and $e^{i\omega(p)}$ where $\omega(p)$ is the solution of the equation $\cos \omega = \nu \cos p$ with positive value. Then we write

$$\omega: (-\pi,\pi] \to [0,\pi] \quad p \mapsto \omega(p) = \arccos(\nu \cos p) \ge 0.$$

For our purposes it is convenient to consider the analytic continuation of U(p) to the subset $S := \{p \in \mathbb{C} \mid \Re(p) = p_R \in (-\pi, \pi], \Im(p) = p_I \leq 0\}$ of the complex plane. The eigenvalues of U(p), with $p \in S$, are $e^{-i\omega(p)}$ and $e^{i\omega(p)}$ where now $\omega(p) = \operatorname{Arccos}(\nu \cos p)$, and Arccos denotes the principal value of the multivalued analytic function arccos. We notice that $\operatorname{Arg}(e^{i\omega(p)}) = \Re(\omega(p)) = \Re(\Delta(p)) = \Re(\operatorname{Arccos}(\nu \cos p) \in [0, \pi].$

In the two-particles case we introduced the center of mass coordinates p, k, representing respectively the total and the relative momentum. While p is always real, k can have an imaginary part. Let us study the eigenvalues of U(p-k) in the limit $k_I \to -\infty$. We have

$$\begin{split} U(p-k) &= \\ \nu e^{i(p-k_R)} e^{-k_I} \begin{pmatrix} e^{2k_I} & -i\frac{\mu}{\nu} e^{-i(p-k_R)} e^{k_I} \\ -i\frac{\mu}{\nu} e^{-i(p-k_R)} e^{k_I} & e^{-i2(p-k_R)} \end{pmatrix}, \end{split}$$

and denoting with λ'_1, λ'_2 the two eigenvalues of $\nu^{-1}e^{-i(p-k_R)}e^{k_I}D(p-k)$ we have for $k_I \to -\infty$

$$\lambda_1' = 1 - \frac{\mu^2}{\nu^2} e^{-2i(p-k_R)} e^{2k_I}, \quad \lambda_2' = \frac{1}{\nu^2} e^{-2i(p-k_R)} e^{2k_I}.$$

Accordingly, for $k_I \to -\infty$, the eigenvalues λ_1 and λ_2 of D(p-k) are

$$\lambda_{1} = \nu e^{i(p-k_{R})} e^{-k_{I}} - \frac{\mu^{2}}{\nu} e^{-i(p-k_{R})} e^{k_{I}},$$

$$\lambda_{2} = \frac{1}{\nu} e^{-i(p-k_{R})} e^{k_{I}},$$

and noticing that $\lim_{k_I \to -\infty} \operatorname{Arg}(\lambda_1) = p - k_R$, and $\lim_{k_I \to -\infty} \operatorname{Arg}(\lambda_2) = -(p - k_R)$ we get

$$p - k_R > 0 \implies e^{-i\omega(p-k)} = \lambda_2, \ e^{i\omega(p-k)} = \lambda_1,$$

$$p - k_R < 0 \implies e^{-i\omega(p-k)} = \lambda_1, \ e^{i\omega(p-k)} = \lambda_2.$$