Interaction of Non-Abelian Tensor Gauge Fields

George Savvidy

Institute of Nuclear and Particle Physics
Demokritos National Research Centre
Ag. Paraskevi, Athens, Greece

Abstract

The non-Abelian tensor gauge fields take values in extended Poincaré algebra. In order to define the invariant Lagrangian we introduce a vector variable in two alternative ways: through the transversal representation of the extended Poincaré algebra and through the path integral over the auxiliary vector field with the U(1) Abelian action. We demonstrate that path integral formulation allows to fix the unitary gauge, derive scattering amplitudes in spinor representation and to define the gauge invariant Lagrangian in a curved space-time.

1. The concept of local gauge invariance formulated by Yang and Mills [1] allows to define the non-Abelian gauge fields, to derive the Lagrangian, the dynamical field equations and to develop a universal point of view on matter interactions [2, 3, 4]. It is appealing to extend the gauge principle so that it will define the interaction of non-Abelian tensor gauge fields as well¹.

The recently proposed generalisation of Yang-Mills theory [26, 27, 28, 29, 34, 35, 36] is based on the extension of the Poincaré algebra $L_G(\mathcal{P})$ with additional generators $L_a^{\lambda_1...\lambda_n}$. The invariant Lagrangian is defined in terms of a composite gauge field $\mathcal{A}_{\mu}^a(x,e_{\perp})$ and the field strength tensor $\mathcal{G}_{\mu\nu}^a(x,e_{\perp})$ depending on the space-time coordinates x_{μ} and the transversal space-like vector variable e_{\perp}^{λ} [30, 31, 32, 29, 34]. The variable e_{\perp}^{λ} belongs to irreducible transversal representation $L_a^{\lambda_1...\lambda_n} = L_a e_{\perp}^{\lambda_1}...e_{\perp}^{\lambda_n}$ of the algebra $L_G(\mathcal{P})$ and fulfils the equations: $e_{\perp}^2 = -1$, $(n \cdot e_{\perp}) = 0$, $n^2 = 0$. These equations are reminiscent of the Abelian gauge field equations. We shall elevate this fact into a guiding principle postulating that the dynamics of the vector variables $e_{\lambda}(x)$ is governed by the Abelian U(1) action and the Lagrangian of the tensor gauge fields is

$$\mathcal{L} = -\frac{1}{4} \langle \mathcal{G}^{a}_{\mu\nu}(x,e) \mathcal{G}^{a\mu\nu}(x,e) \rangle = -\frac{1}{4} \int \mathcal{G}^{a}_{\mu\nu}(x,e) \mathcal{G}^{a\mu\nu}(x,e) e^{-\frac{i}{4} \int F^{2}(e)d^{4}x} \mathcal{D}e_{\lambda} , \qquad (1)$$

where $F_{\rho\lambda} = \partial_{\rho}e_{\lambda} - \partial_{\lambda}e_{\rho}$ and $\langle ... \rangle = \int ... \mathcal{D}e_{\lambda}$ denotes a path integral over $e_{\lambda}(x)$. The advantage of the above formulation lies in the fact that the dynamics of the auxiliary variable $e_{\lambda}(x)$ is identical to the photon dynamics and therefore naturally describes the space-like transversal polarisations e_{\perp} appearing earlier in $L_a^{\lambda_1...\lambda_n}$ representation of the $L_G(\mathcal{P})$ algebra. The expansion of the integrand $\mathcal{G}^a_{\mu\nu}(x,e)\mathcal{G}^{a\mu\nu}(x,e)$ over the auxiliary variable e_{\perp}^{λ} expresses the Lagrangian \mathcal{L} in terms of vacuum expectation values

$$\mathcal{L} = \sum_{s=0}^{\infty} \frac{1}{s!} \mathcal{L}_{\lambda_1 \dots \lambda_s} \left\langle e_{\perp}^{\lambda_1} \dots e_{\perp}^{\lambda_s} \right\rangle . \tag{2}$$

The $L_G(\mathcal{P})$ generators have the form $L_a^{\lambda_1...\lambda_n} = L_a e_{\perp}^{\lambda_1}...e_{\perp}^{\lambda_n}$ and the vacuum expectation values $\langle L_a^{\lambda_1...\lambda_n} | L_b^{\lambda_{n+1}...\lambda_s} \rangle = \delta_{ab} \langle e_{\perp}^{\lambda_1}...e_{\perp}^{\lambda_n} e_{\perp}^{\lambda_{n+1}}...e_{\perp}^{\lambda_s} \rangle$ play the role of the Killing metric. The Killing metric contracts only the space-like components and guarantee the absence of negative norm time-like components of the tensor gauge fields in the Lagrangian \mathcal{L} .

¹The research in high spin field theories has long and rich history. One should mention the early works of Majorana [5], Fierz [6], Pauli [7], Dirac [8], Ginsburg and Tamm [9], Weinberg [10], Minkowski [11], Nambu [12],Schwinger [13], Ramond [14], Singh and Hagen [15], Fronsdal [16], Brink et.al. [17, 18], Berends et.al. [19], Fradkin[20], Vasiliev [21], Sagnotti, Sezgin and Sundel [22], Metsaev [23], Manvelyan et.al. [24] and many other works (see also the references in [25]).

2. The non-Abelian tensor gauge fields $A^a_{\mu\lambda_1...\lambda_s}(x)$, s=0,1,2,... are totally symmetric in indices $\lambda_1...\lambda_s$ and are the components of the field $\mathcal{A}_{\mu}(x,e)$ [26, 27, 28]:

$$\mathcal{A}_{\mu}(x,e) = \sum_{s=0}^{\infty} \frac{1}{s!} A^{a}_{\mu\lambda_{1}...\lambda_{s}}(x) L_{a}e^{\lambda_{1}}...e^{\lambda_{s}}$$

$$\tag{3}$$

taking values in the extended Poincaré algebra. The $L_G(\mathcal{P})$ algebra [28, 29, 34, 35] is:

$$[P^{\mu}, P^{\nu}] = 0,$$

$$[M^{\mu\nu}, P^{\lambda}] = \eta^{\nu\lambda} P^{\mu} - \eta^{\mu\lambda} P^{\nu},$$

$$[M^{\mu\nu}, M^{\lambda\rho}] = \eta^{\mu\rho} M^{\nu\lambda} - \eta^{\mu\lambda} M^{\nu\rho} + \eta^{\nu\lambda} M^{\mu\rho} - \eta^{\nu\rho} M^{\mu\lambda},$$

$$[P^{\mu}, L_a^{\lambda_1 \dots \lambda_s}] = 0,$$

$$[M^{\mu\nu}, L_a^{\lambda_1 \dots \lambda_s}] = \eta^{\nu\lambda_1} L_a^{\mu\lambda_2 \dots \lambda_s} - \eta^{\mu\lambda_1} L_a^{\nu\lambda_2 \dots \lambda_s} + \dots + \eta^{\nu\lambda_s} L_a^{\mu\lambda_1 \dots \lambda_{s-1}} - \eta^{\mu\lambda_s} L_a^{\nu\lambda_1 \dots \lambda_{s-1}},$$

$$[L_a^{\lambda_1 \dots \lambda_k}, L_b^{\lambda_{k+1} \dots \lambda_s}] = i f_{abc} L_c^{\lambda_1 \dots \lambda_s}.$$

$$(4)$$

It incorporates the Poincaré and internal algebra L_G with structure constants f_{abc} . The generators $L_a^{\lambda_1...\lambda_s}$ carry the *internal charges and spins*. There is no conflict with the Coleman-Mandula theorem [33] because the theorem applies to the symmetries that act on S-matrix elements and not to symmetries which acts on the gauge field $\mathcal{A}_{\mu}(x,e)$. The algebra (4) is invariant with respect to the "gauge" transformations [29, 34, 35]:

$$L_a^{\lambda_1...\lambda_s} \to L_a^{\lambda_1...\lambda_s} + \sum_1 \chi \ P^{\lambda_1} L_a^{\lambda_2...\lambda_s} + \sum_2 \chi \ P^{\lambda_1} P^{\lambda_2} L_a^{\lambda_3...\lambda_s} + \dots + \chi^s \ P^{\lambda_1} \dots P^{\lambda_s} L_a$$

$$M^{\mu\nu} \to M^{\mu\nu}, \quad P^{\lambda} \to P^{\lambda}, \tag{5}$$

where the sums $\Sigma_1, \Sigma_2, ...$ are over all inequivalent index permutations and χ is a scalar. The invariance retains if one consider any translationally invariant vector n^{λ} instead of P^{λ} . The representations of $L_G(\mathcal{P})$ are therefore defined modulo longitudinal terms proportional to the momentum P^{λ} . The algebra $L_G(\mathcal{P})$ has representation in terms of the differential operators:

$$P^{\mu} = k^{\mu},$$

$$M^{\mu\nu} = i(k^{\mu} \frac{\partial}{\partial k_{\nu}} - k^{\nu} \frac{\partial}{\partial k_{\mu}}) + i(e^{\mu} \frac{\partial}{\partial e_{\nu}} - e^{\nu} \frac{\partial}{\partial e_{\mu}}),$$

$$L_{a}^{\lambda_{1}...\lambda_{s}} = e^{\lambda_{1}}...e^{\lambda_{s}} \otimes L_{a}.$$
(6)

The irreducible transversal representation is defined by invariant equations [30, 31, 32]:

$$n^2 = 0, \quad (n \cdot e_\perp) = 0, \quad e_\perp^2 = -1,$$
 (7)

where n_{λ} is an arbitrary light-like vector². These equations have the solution

$$e^{\mu}_{\perp} = \chi n^{\mu} + e^{i\varphi} e^{\mu}_{+} + e^{-i\varphi} e^{\mu}_{-},$$
 (8)

where e^{μ}_{\pm} are helicity polarisation vectors, the variables χ and φ are defined on the cylinder $\varphi \in S^1, \chi \in R^1$. The representation of the generators $L_a^{\lambda_1...\lambda_s}$ therefore is:

$$L_a^{\perp \lambda_1 \dots \lambda_s} = \prod_{n=1}^s (\chi n^{\lambda_n} + e^{i\varphi} e_+^{\lambda_n} + e^{-i\varphi} e_-^{\lambda_n}) \otimes L_a$$
 (9)

and it is transversal because of (7):

$$n_{\lambda_1} L_a^{\perp \lambda_1 \dots \lambda_s} = 0, \quad s = 1, 2, \dots$$
 (10)

Opening the brackets in (9) and separating terms containing the variables n^{λ_i} we shall get the first term of the form $\prod_{n=1}^{s} (e^{i\varphi}e_+^{\lambda_n} + e^{-i\varphi}e_-^{\lambda_n})$ which represents the helicity generators $(L_{a+\cdots+}^{\lambda_1\dots\lambda_s}, \dots, L_{a-\cdots-}^{\lambda_1\dots\lambda_s})$ with helicity spectrum $h = (+s, +s-2, \dots, -s+2, -s)$. The rest of the terms contain n^{λ} and correspond to the transformation (5) of L_a^{\perp} $\lambda_1\dots\lambda_s$.

Substituting the representation (9) of the $L_a^{\perp \lambda_1 \dots \lambda_s}$ into the expansion (3) and collecting terms in front of the helicity generators $(L_{a+\dots+}^{\lambda_1 \dots \lambda_s}, \dots, L_{a-\dots-}^{\lambda_1 \dots \lambda_s})$ we shall get (see Appendix for details)

$$\mathcal{A}_{\mu}(x,e) = \sum_{s=0}^{\infty} \frac{1}{s!} \left(A_{\mu+\cdots+} + \dots + A_{\mu-\cdots-} \right), \tag{11}$$

where s is the number of helicity indices and $A_{\mu+\cdots} = A^a_{\mu\lambda_1\dots\lambda_s} L^{\lambda_1\dots\lambda_s}_{a+\cdots}$. It follows from (11) that the time components $\lambda_i = 0, i = 1, ..., s$ of the tensor gauge fields have been gauged away. The rest of the time components $(A_{0+\cdots+}, ..., A_{0-\cdots-})$, when the first index $\mu = 0$ is equal to zero, will be gauged away by the gauge transformation [1, 28, 29]:

$$\mathcal{A}'_{\mu}(x,e) = U(\xi)\mathcal{A}_{\mu}(x,e)U^{-1}(\xi) - \frac{i}{g}\partial_{\mu}U(\xi)\ U^{-1}(\xi),\tag{12}$$

where $U(\xi) = \exp(i\xi(x,e))$ and the parameter $\xi(x,e)$ depends on the vector variable [28, 29]:

$$\xi(x,e) = \sum_{s} \frac{1}{s!} \, \xi^{a}_{\lambda_{1} \dots \lambda_{s}}(x) \, L^{\lambda_{1} \dots \lambda_{s}}_{a}.$$

 $^{^{2}}$ A vector variable, in addition to the space-time coordinate x, was introduced earlier by Yukawa [30], Fierz [31] and Wigner[32].

The components $\xi^a_{\lambda_1...\lambda_s}(x)$ are totally symmetric. The commutator of covariant derivatives $\nabla^{ab}_{\mu} = (\partial_{\mu} - ig\mathcal{A}_{\mu}(x,e))^{ab}$ defines the field strength tensor

$$\mathcal{G}_{\mu\nu}(x,e) = \partial_{\mu}\mathcal{A}_{\nu}(x,e) - \partial_{\nu}\mathcal{A}_{\mu}(x,e) - ig[\mathcal{A}_{\mu}(x,e) \,\mathcal{A}_{\nu}(x,e)]
= \sum_{s} \frac{1}{s!} G^{a}_{\mu\nu,\lambda_{1}...\lambda_{s}}(x) L^{\lambda_{1}...\lambda_{s}}_{a},$$
(13)

which transforms homogeneously $\mathcal{G}'_{\mu\nu}(x,e) = U(\xi)\mathcal{G}_{\mu\nu}(x,e)U^{-1}(\xi)$ and the gauge invariant Lagrangian is defined as [26, 27, 28]

$$\mathcal{L}(x) = \langle \mathcal{L}(x, e) \rangle = -\frac{1}{4} \langle \mathcal{G}^{a}_{\mu\nu}(x, e) \mathcal{G}^{a\mu\nu}(x, e) \rangle. \tag{14}$$

Using the $L_G(\mathcal{P})$ generators (9) one can calculate the Killing metric [29, 34, 36]:

$$L_{G}: \qquad \langle L_{a}|L_{b}\rangle = \delta_{ab}, \tag{15}$$

$$L_{P}: \qquad \langle P^{\mu}|P^{\nu}\rangle = 0$$

$$\langle M_{\mu\nu}|P_{\lambda}\rangle = 0$$

$$\langle M^{\mu\nu}|M^{\lambda\rho}\rangle = \eta^{\mu\lambda}\eta^{\nu\rho} - \eta^{\mu\rho}\eta^{\nu\lambda}$$

$$L_{G}(P): \qquad \langle P^{\mu}|L_{a}^{\perp \lambda_{1}...\lambda_{s}}\rangle = 0, \tag{2}$$

$$\langle M^{\mu\nu}|L_{a}^{\perp \lambda_{1}...\lambda_{s}}\rangle = 0, \tag{3}$$

$$\langle L_{a}^{\perp \lambda_{1}...\lambda_{n}}|L_{b}^{\perp \lambda_{n+1}...\lambda_{2s+1}}\rangle = 0, \qquad s = 0, 1, 2, 3, ...$$

$$\langle L_{a}^{\perp \lambda_{1}...\lambda_{n}}|L_{b}^{\perp \lambda_{n+1}...\lambda_{2s}}\rangle = \delta_{ab} \ s! \ (\bar{\eta}^{\lambda_{1}\lambda_{2}}\bar{\eta}^{\lambda_{3}\lambda_{4}}...\bar{\eta}^{\lambda_{2s-1}\lambda_{2s}} + \text{perm}),$$

where $\bar{\eta}^{\lambda_1 \lambda_2}$ is the projector into the two-dimensional plane transversal to the light-like vector n^{λ} [13]:

$$\bar{\eta}^{\lambda_1 \lambda_2} = -\eta^{\lambda_1 \lambda_2} + \frac{n^{\lambda_1} \bar{n}^{\lambda_2} + \bar{n}^{\lambda_1} n^{\lambda_2}}{n \bar{n}} = \sum_{i=+} e_i^{\lambda_1} e_i^{\lambda_2}, \qquad n_{\lambda_1} \bar{\eta}^{\lambda_1 \lambda_2} = n_{\lambda_2} \bar{\eta}^{\lambda_1 \lambda_2} = 0, \qquad (16)$$

where $n = (n_0, \vec{n})$ and $\bar{n}^{\mu} = (n_0, -\vec{n})$ and the Killing metric is transversal (7):

$$n_{\lambda_i} \langle L_a^{\perp \lambda_1 \dots \lambda_n}; L_b^{\perp \lambda_{n+1} \dots \lambda_{2s}} \rangle = 0, \quad i = 1, 2, \dots 2s.$$
 (17)

Expanding the $\mathcal{L}(x,e)$ in (14) over the vector variable e^{λ} we shall get [26, 27, 28]:

$$\mathcal{L}(x) = \langle \mathcal{L}(x, e) \rangle = \sum_{s=0}^{\infty} \frac{1}{s!} \mathcal{L}_{\lambda_1 \dots \lambda_s}(x) \langle e^{\lambda_1} \dots e^{\lambda_s} \rangle . \tag{18}$$

In particular, the term quadratic in e^{λ} is

$$\mathcal{L}_2 = \mathcal{L}_{\lambda_1 \lambda_2} \langle e^{\lambda_1} e^{\lambda_2} \rangle = -\frac{1}{4} (G^a_{\mu\nu,\lambda_1} G^{a\mu\nu,}_{\lambda_2} + G^a_{\mu\nu,\lambda_1 \lambda_2} G^{a\mu\nu}) \bar{\eta}^{\lambda_1 \lambda_2}$$
(19)

or using the bar symbol over the indices indicating the transversal Killing metric contraction (16) we get

$$\mathcal{L}_2 = -\frac{1}{4} G^a_{\mu\nu,\bar{\lambda}} G^{a\mu\nu,\bar{\lambda}} - \frac{1}{4} G^a_{\mu\nu,\bar{\lambda}} \bar{A} G^{a\mu\nu}. \tag{20}$$

3. In the transversal representation (9) of the $L_G(\mathcal{P})$ algebra the tensor gauge fields (11) are partially gauged and the residual gauge transformation of (12), (48) takes the following form [26, 27, 28]

$$A'_{\mu} = A_{\mu} + \nabla_{\mu}\xi,$$

$$A'_{\mu\pm} = A_{\mu\pm} + \partial_{\mu}\xi_{\pm} - ig[A_{\mu}, \xi_{\pm}] - ig[A_{\mu\pm}, \xi],$$

$$A'_{\mu\pm\pm} = A_{\mu\pm\pm} + \nabla_{\mu}\xi_{\pm\pm} - ig([A_{\mu\pm}, \xi_{\pm}] + [A_{\mu\pm}, \xi_{\pm}] + [A_{\mu\pm\pm}, \xi]),$$

$$A'_{\mu+-} = A_{\mu+-} + \nabla_{\mu}\xi_{+-} - ig([A_{\mu+}, \xi_{-}] + [A_{\mu-}, \xi_{+}] + [A_{\mu+-}, \xi]),$$
......

where $A_{\mu\pm} = A^a_{\mu\lambda} L^{\perp\lambda}_{a\pm}$, $\xi_{\pm} = \xi^b_{\lambda} L^{\perp\lambda}_{b\pm}$, $A_{\mu\pm\pm} = A^a_{\mu\nu\lambda} L^{\perp\nu\lambda}_{a\pm\pm}$, $\xi_{\pm\pm} = \xi^b_{\nu\lambda} L^{\perp\nu\lambda}_{b\pm\pm}$ and so on. Using the residual gauge transformation (21) we can impose the Lorentz gauge on the rest of the field components:

$$\partial^{\mu} A'_{\mu+\dots} = 0. \tag{22}$$

Indeed, we shall impose the standard Lorentz gauge $\partial^{\mu}A_{\mu}^{'}=0$ by solving the equation

$$\partial^{\mu} A_{\mu} + \partial^{\mu} \nabla_{\mu} \xi = 0, \quad \xi = -(\partial^{\mu} \nabla_{\mu})^{-1} \partial^{\nu} A_{\nu} . \tag{23}$$

Imposing a similar gauge $\partial^{\mu}A'_{\mu\pm}=0$ on the tensor gauge field we get

$$\partial^{\mu} A_{\mu\pm} + \partial^{\mu} \nabla_{\mu} \xi_{\pm} - ig \partial^{\mu} [A_{\mu\pm}, \xi] = 0, \tag{24}$$

which can be solved by using the gauge parameters ξ_{\pm}

$$\xi_{\pm} = -(\partial^{\mu} \nabla_{\mu})^{-1} \partial^{\nu} (A_{\nu \pm} - ig[A_{\nu \pm}, \xi]), \tag{25}$$

The gauge condition $\partial^{\mu}A_{\mu\pm\pm}^{\prime}=0$ on the rank-3 gauge field can also be resolved by

$$\xi_{\pm\pm} = -(\partial^{\mu}\nabla_{\mu})^{-1}\partial^{\nu}(A_{\nu\pm\pm} - 2ig[A_{\nu\pm}, \xi_{\pm}] - ig[A_{\nu\pm\pm}, \xi]),$$

$$\xi_{+-} = -(\partial^{\mu}\nabla_{\mu})^{-1}\partial^{\nu}(A_{\nu+-} - ig[A_{\nu+}, \xi_{-}] - ig[A_{\nu-}, \xi_{+}] - ig[A_{\nu+-}, \xi]).$$
 (26)

Thus the two-stage gauge fixing procedure (7) - (11) and (22) - (26)

$$A_{\mu+\cdots} = A^a_{\mu\lambda_1\cdots\lambda_s} \ L^{\lambda_1\cdots\lambda_s}_{a+\cdots} \quad \text{and} \quad \partial^{\mu} A'_{\mu+\cdots} = 0$$
 (27)

allows to completely exclude the time-like components of the tensor gauge fields leaving only pure transversal helicity polarisations.

4. As it was demonstrated above, the vector variable e^{λ} has the properties which are reminiscent of the Abelian gauge field. We shall elevate this fact into a guiding principle postulating that the dynamics of the vector variables $e^{\lambda}(x)$ is governed by the Abelian U(1) action. We shall consider now the Killing metric (15) as a correlation function of the Abelian U(1) gauge field $e^{\lambda}(x)$ defined in terms of the Feynman path integral:

$$\langle Te^{\lambda_1}(x_1)...e^{\lambda_s}(x_s)\rangle = \int e^{i\int (-\frac{1}{4}F_{\alpha\beta}(e)F^{\alpha\beta}(e)+J_{\alpha}e^{\alpha})}e^{\lambda_1}(x_1)...e^{\lambda_s}(x_s)\mathcal{D}e_{\lambda}(x), \tag{28}$$

where the external current is conserved: $\partial^{\alpha} J_{\alpha} = 0$ [13]. The action for the vector variable $e^{\lambda}(x)$ in the axial gauge has the form

$$\int \left\{ -\frac{1}{4} F_{\alpha\beta}(e) F^{\alpha\beta}(e) + \lambda^2 e^{\alpha}(x) n_{\alpha} n_{\beta} e^{\beta}(x) \right\} d^4 x = \frac{1}{2} \int e^{\alpha}(x) H_{\alpha\beta} e^{\beta}(x) d^4 x, \tag{29}$$

where $H_{\alpha\beta} = \eta_{\alpha\beta}\partial^2 - \partial_\alpha\partial_\beta + \lambda^2 n_\alpha n_\beta$, $H_{\alpha\beta} = -\eta_{\alpha\beta}k^2 + k_\alpha k_\beta + \lambda^2 n_\alpha n_\beta$. The vector variable propagator $H_{\alpha\beta}(k)\Delta^{\beta\gamma}(k) = i\delta^{\gamma}_{\alpha}$ has the following expression:

$$\Delta^{\alpha\beta}(k) = \left(-\eta^{\alpha\beta} + \frac{k^2 - \lambda^2 n^2}{\lambda^2 (k \cdot n)^2} k^{\alpha} k^{\beta} + \frac{k^{\alpha} n^{\beta} + n^{\alpha} k^{\beta}}{(k \cdot n)}\right) \frac{i}{k^2 + i\epsilon}.$$

In the limit $\lambda^2 \to \infty$

$$\Delta^{\alpha\beta}(k) = i\left(-\eta^{\alpha\beta} - \frac{n^2}{(k\cdot n)^2}k^{\alpha}k^{\beta} + \frac{k^{\alpha}n^{\beta} + n^{\alpha}k^{\beta}}{(k\cdot n)}\right)\frac{i}{k^2 + i\epsilon}$$

the propagator is explicitly transversal

$$n_{\alpha} \Delta^{\alpha\beta}(k) = 0. (30)$$

In the light-cone gauge, when $n^2 = 0$, we shall get

$$\Delta^{\alpha\beta}(x-y) = \langle 0|Te^{\alpha}(x)e^{\beta}(y)|0\rangle = \int \frac{d^dk}{(2\pi)^d} \left(-\eta^{\alpha\beta} + \frac{k^{\alpha}n^{\beta} + n^{\alpha}k^{\beta}}{(k \cdot n)}\right) \frac{i}{k^2 + i\epsilon} e^{-ik \cdot (x-y)}$$
(31)

and the correlation function at the coincident points $x \to y$ is:

$$\langle 0|Te^{\alpha}(x)e^{\beta}(x)|0\rangle = \int \frac{d^dk}{(2\pi)^d} \left(-\eta^{\alpha\beta} + \frac{k^{\alpha}n^{\beta} + n^{\alpha}k^{\beta}}{(k \cdot n)}\right) \frac{i}{k^2 + i\epsilon}.$$

The above integral depends on the covariant expression of the form

$$\langle 0|Te^{\alpha}(x)e^{\beta}(x)|0\rangle = -A\eta^{\alpha\beta} + B(n^{\alpha}\bar{n}^{\beta} + \bar{n}^{\alpha}n^{\beta}), \tag{32}$$

where $n^{\alpha}=(n_0,\vec{n})$, $\bar{n}^{\alpha}=(n_0,-\vec{n})$ and $n^2=\bar{n}^2=n_0^2-\vec{n}^2=0$. Calculating the trace I and contraction with n_{α} , we shall get two equations: $-4A+2B(n\cdot\bar{n})=I$, $-A+B(n\cdot\bar{n})=0$, therefore $B=A/(n\cdot\bar{n})$ and

$$A = \int \frac{d^d k}{(2\pi)^d} \frac{i}{k^2 + i\epsilon} \ . \tag{33}$$

These lead us to the expression for the vector variable correlation function

$$\frac{1}{A}\langle 0|Te^{\alpha}(x)e^{\beta}(x)|0\rangle = -\eta^{\alpha\beta} + \frac{n^{\alpha}\bar{n}^{\beta} + \bar{n}^{\alpha}n^{\beta}}{(n\cdot\bar{n})},\tag{34}$$

which is transversal to the vector n_{α} (30). Because $n^2 = 0$, the matrix (34) has two eigenvectors e_{\pm} in the space-like directions:

$$n^2 = \bar{n}^2 = 0$$
, $(n \cdot e_{\pm}) = 0$ $(\bar{n} \cdot e_{\pm}) = 0$, $e_{\pm}^2 = 0$, $(e_{+} \cdot e_{-}) = 1$ (35)

and

$$\bar{\eta}^{\alpha\beta} = \frac{1}{A} \langle 0 | T e^{\alpha}(x) e^{\beta}(x) | 0 \rangle = -\eta^{\alpha\beta} + \frac{n^{\alpha} \bar{n}^{\beta} + \bar{n}^{\alpha} n^{\beta}}{(n \cdot \bar{n})} = \sum_{i=+}^{\infty} e_i^{\alpha} e_i^{\beta}, \tag{36}$$

Using the Wick theorem we shall get the metric (15). Thus the Killing metric, expressed in terms of the Feynman path integral (28) and in terms of traces of the $L_G(\mathcal{P})$ generators (15) in the transversal representation (9) coincide.

The conclusion is that the Lagrangian (1), (14) for the tensor gauge fields can be defined in flat space-time in two equivalent ways: either by using transversal representation of the $L_G(\mathcal{P})$ generators (15) or by postulating that the dynamics of the vector variable is governed by the Abelian action (28). In both cases the vector variable has two transversal polarisations and the corresponding Killing metric contracts transversal space-like components of the tensor gauge fields. Being equivalent in flat space-time they are not equivalent in curved. The advantage of the path integral formulation lies in the fact that it opens a prospect to define the gauge invariant Lagrangian in curved space-time.

4. The Lagrangian (1), (14) defines the propagation of free tensor gauge bosons and their interactions. The high-spin bosons interact through the triple and quartic interaction vertices and a dimensionless coupling constant. In order to calculate scattering amplitudes

it is convenient to use spinor representation of amplitudes developed in [37, 38, 39, 40]. A scattering amplitude $M_n = M_n(\lambda_1, \tilde{\lambda}_1, h_1; ...; \lambda_n, \tilde{\lambda}_n, h_n)$ for the massless particles of momenta p_i and polarisation tensors ε_i (i = 1, ..., n) can be represented in terms of spinors:

$$k_{a\dot{a}} = \lambda_a \tilde{\lambda}_{\dot{a}}, \quad e_{a\dot{a}}^+ = \frac{\mu_a \tilde{\lambda}_{\dot{a}}}{\langle \mu, \lambda \rangle}, \quad e_{a\dot{a}}^- = \frac{\lambda_a \tilde{\mu}_{\dot{a}}}{[\lambda, \mu]}, \tag{37}$$

where

$$\lambda_a = (\sqrt{k^+}, \frac{k_x + ik_y}{\sqrt{k^+}}), \quad \tilde{\lambda}_{\dot{a}} = (\sqrt{k^+}, \frac{k_x - ik_y}{\sqrt{k^+}}), \quad k^+ = k_t + k_z ,$$

and μ_a is a reference spinor. In Yang-Mills theory the interaction vertex VVV is

$$\mathcal{V}_{\alpha\beta\gamma}^{abc}(k,p,q) = -igf^{abc}F_{\alpha\beta\gamma}(k,p,q) = -igf^{abc}[\eta_{\alpha\beta}(p-k)_{\gamma} + \eta_{\alpha\gamma}(k-q)_{\beta} + \eta_{\beta\gamma}(q-p)_{\alpha}].$$
(38)

In (1), (14) the interaction vertex VTT, the tensor-vector-tensor vertex is [28]

$$\mathcal{V}_{\alpha\dot{\alpha}\beta\gamma\dot{\gamma}}^{abc}(k,p,q) = -igf^{abc}[\eta_{\alpha\beta}(p-k)_{\gamma} + \eta_{\alpha\gamma}(k-q)_{\beta} + \eta_{\beta\gamma}(q-p)_{\alpha}]\bar{\eta}_{\dot{\alpha}\dot{\gamma}}, \qquad (39)$$

where the indices $(a, \alpha \dot{\alpha}, k)$ belong to the tensor gauge boson, (b, β, p) to the vector gauge boson and $(c, \gamma \dot{\gamma}, q)$ to the second tensor gauge boson. In massless theory the momentum conservation $\delta(k+p+q)$ equation has the solution $k=(\omega,0,0,r), \quad p=(\omega,0,0,r), \quad q=(-2\omega,0,0,-2r)$ ($\omega^2=r^2$) that can be deformed by a complex parameter z [38, 37, 39, 40, 41], $k=(\omega,z,iz,r), \quad p=(\omega,-z,-iz,r), \quad q=(-2\omega,0,0,-2r)$ and the corresponding polarisation vectors

$$e_k^+ = \frac{1}{\sqrt{2}}(\frac{z}{\omega}, 1, -i, -\frac{z}{r}), \quad e_p^+ = \frac{1}{\sqrt{2}}(-\frac{z}{\omega}, 1, -i, \frac{z}{r}), \quad e_q^- = \frac{1}{\sqrt{2}}(0, 1, i, 0)$$

are orthogonal to the momenta $k \cdot e_k^+ = 0$, $p \cdot e_p^+ = 0$, $q \cdot e_q^- = 0$. Computing the matrix element of the YM vertex VVV (38) we shall get:

$$M_{YM}(+1, +1, -1) = F_{\alpha\beta\gamma}(k, p, q) \ e_k^{+\alpha} e_p^{+\beta} e_q^{-\gamma} =$$

$$= (e_k^+ \cdot e_p^+) \ (p - k \cdot e_q^-) + (e_p^+ \cdot e_q^-) \ (q - p \cdot e_k^+) + (e_q^- \cdot e_k^+) \ (k - q \cdot e_p^+) =$$

$$= 2(e_k^+ \cdot e_p^+) \ (p \cdot e_q^-) + 2(e_p^+ \cdot e_q^-) \ (q \cdot e_k^+) + 2(e_q^- \cdot e_k^+) \ (k \cdot e_p^+). \tag{40}$$

In a spinor representation (37) the amplitude (40) is

$$M_{YM}(+1,+1,-1) = \frac{[1,2]^4}{[1,2][2,3][3,1]},$$

where the reference spinors have been chosen as $\mu^{(k)} = \lambda^{(p)}$, $\mu^{(p)} = \lambda^{(q)}$ and $\mu^{(q)} = \lambda^{(p)}$. For the VTT amplitude $M_{GYM} = \varepsilon^{\alpha\dot{\alpha}}(k)e^{\beta}(p)\varepsilon^{\gamma\gamma'}(q)$ $\mathcal{V}_{\alpha\dot{\alpha}\beta\gamma\dot{\gamma}}(k,p,q)$ $\delta(k+p+q)$ we shall get

$$M_{GYM}(+2,+1,-2) = 2\left(e_k^+ \cdot e_p^+\right) \left(p \cdot e_q^-\right) + \left(e_p^+ \cdot e_q^-\right) \left(q \cdot e_k^+\right) + \left(e_q^- \cdot e_k^+\right) \left(k \cdot e_p^+\right) \left(e_k^+ \cdot e_q^-\right). \tag{41}$$

The last term has the following form:

$$(e_k^+ \cdot e_q^-) = \frac{\langle \mu^{(k)}, \lambda^{(q)} \rangle [\lambda^{(k)}, \mu^{(q)}]}{\langle \mu^{(k)}, \lambda^{(k)} \rangle [\lambda^{(q)}, \mu^{(q)}]} = \left(\frac{[1, 2]}{[2, 3]}\right)^2 , \tag{42}$$

and in spinor representation we shall get

$$M_{GYM}(+2,+1,-2) = \frac{[1,2]^4}{[1,2][2,3][3,1]} \left(\frac{[1,2]}{[2,3]}\right)^2. \tag{43}$$

The reference spinors for both tensor bosons are equal to each other: $\mu^{(k)} = \mu^{(q)}$. Considering the high spin tensors one can find

$$M_{GYM}(+s,+1,-s) = \frac{[1,2]^4}{[1,2][2,3][3,1]} \left(\frac{[1,2]}{[2,3]}\right)^{2s-2},$$
(44)

which reduces to the YM amplitude when s = 1. The TTT-amplitude for particles of helicities (h_1, h_2, h_3) has the following general form [17, 18, 41, 43]:

$$M_3 = gf^{abc} < 1, 2 >^{-2h_1 - 2h_2 - 1} < 2, 3 >^{2h_1 + 1} < 3, 1 >^{2h_2 + 1}, \quad h_3 = -1 - h_1 - h_2,$$

$$M_3 = gf^{abc} [1, 2]^{2h_1 + 2h_2 - 1} [2, 3]^{-2h_1 + 1} [3, 1]^{-2h_2 + 1}, \quad h_3 = 1 - h_1 - h_2.$$

$$(45)$$

In particular, considering the interaction between bosons of helicities (+s, +1, -s) one can get convinced that the general expression (45) is in full agreement with the direct calculation (44).

5. The advantage of the path integral formulation of the Lagrangian (1), (14) is that it can be defined in a background gravitational field as well [26, 27, 28]:

$$\mathcal{L} = -\frac{1}{4} \int \mathcal{G}^{a}_{\mu\nu}(x,e) \mathcal{G}^{a\mu\nu}(x,e) e^{-i\frac{1}{4} \int F_{\alpha\beta}(e)F^{\alpha\beta}(e)\sqrt{-g}d^4x} \mathcal{D}e_{\lambda}(x), \tag{46}$$

where the field strength tensors are well defined in terms of ordinary derivatives:

$$F_{\alpha\beta}(e) = e_{\beta;\alpha} - e_{\alpha;\beta} = \partial_{\alpha}e_{\beta} - \partial_{\beta}e_{\alpha}$$

$$\mathcal{G}_{\mu\nu}(x,e) = \mathcal{A}_{\nu;\mu}(x,e) - \mathcal{A}_{\mu;\nu}(x,e) - ig[\mathcal{A}_{\mu}(x,e) \mathcal{A}_{\nu}(x,e)]$$

$$= \partial_{\mu}\mathcal{A}_{\nu}(x,e) - \partial_{\nu}\mathcal{A}_{\mu}(x,e) - ig[\mathcal{A}_{\mu}(x,e) \mathcal{A}_{\nu}(x,e)]. \tag{47}$$

It follows therefore that the extended gauge invariance, (12) and (48), still holds.

6. We calculated the scattering amplitudes of non-Abelian tensor gauge bosons at tree level approximation in [43], as well as their one-loop contribution into the Callan-Symanzik beta function [44]. This contribution is negative and corresponds to an asymptotically free theory. The review article [45] contains the details of the calculations.

I would like to thank the CERN Theory Division, where part of this work was completed, for kind hospitality.

1 Appendix

The explicit expression for the gauge fields transformation is [26, 27, 28]:

and the components of the field strength tensor are [26, 27, 28]:

$$G^{a}_{\mu\nu} = \partial_{\mu}A^{a}_{\nu} - \partial_{\nu}A^{a}_{\mu} + gf^{abc} A^{b}_{\mu} A^{c}_{\nu},$$

$$G^{a}_{\mu\nu,\lambda} = \partial_{\mu}A^{a}_{\nu\lambda} - \partial_{\nu}A^{a}_{\mu\lambda} + gf^{abc} (A^{b}_{\mu} A^{c}_{\nu\lambda} + A^{b}_{\mu\lambda} A^{c}_{\nu}),$$

$$G^{a}_{\mu\nu,\lambda\rho} = \partial_{\mu}A^{a}_{\nu\lambda\rho} - \partial_{\nu}A^{a}_{\mu\lambda\rho} + gf^{abc} (A^{b}_{\mu} A^{c}_{\nu\lambda\rho} + A^{b}_{\mu\lambda} A^{c}_{\nu\rho} + A^{b}_{\mu\rho} A^{c}_{\nu\lambda} + A^{b}_{\mu\lambda\rho} A^{c}_{\nu}),$$

$$(49)$$

The transversal tensor gauge fields $A_{\mu+\cdots}$ appear in the expansion (11):

$$\mathcal{A}(x,e) = A_{\mu} + A_{\mu\lambda_{1}}\chi n^{\lambda_{1}} + \frac{1}{2}A_{\mu\lambda_{1}\lambda_{2}}(\chi^{2}n^{\lambda_{1}}n^{\lambda_{2}} + e_{+}^{\lambda_{1}}e_{-}^{\lambda_{2}} + e_{-}^{\lambda_{1}}e_{+}^{\lambda_{2}}) + \dots
+ A_{\mu\lambda_{1}}e^{i\varphi}e_{+}^{\lambda_{1}} + \frac{1}{2}A_{\mu\lambda_{1}\lambda_{2}}e^{i\varphi}(e_{+}^{\lambda_{1}}\chi n^{\lambda_{2}} + \chi n^{\lambda_{1}}e_{+}^{\lambda_{2}}) + \dots
+ A_{\mu\lambda_{1}}e^{-i\varphi}e_{-}^{\lambda_{1}} + \frac{1}{2}A_{\mu\lambda_{1}\lambda_{2}}e^{-i\varphi}(e_{-}^{\lambda_{1}}\chi n^{\lambda_{2}} + \chi n^{\lambda_{1}}e_{-}^{\lambda_{2}}) + \dots
+ \frac{1}{2}A_{\mu\lambda_{1}\lambda_{2}}e^{2i\varphi}e_{+}^{\lambda_{1}}e_{+}^{\lambda_{2}} + \frac{1}{2}A_{\mu\lambda_{1}\lambda_{2}}e^{-2i\varphi}e_{-}^{\lambda_{1}}e_{-}^{\lambda_{2}} + \dots
= > A_{\mu} + A_{\mu+} + A_{\mu-} + \frac{1}{2}A_{\mu++} + \frac{1}{2}A_{\mu--} \quad \dots$$
(50)

The phase factor $e^{ih\varphi}$ defines the helicity h of the generators $L_{a+\cdots-}^{\lambda_1\dots\lambda_s}=e^{ih\varphi}e_+^{\lambda_1}\dots e_-^{\lambda_s}\otimes L_a$.

References

- [1] C.N.Yang and R.L.Mills. Conservation of Isotopic Spin and Isotopic Gauge Invariance. Phys. Rev. **96** (1954) 191
- [2] S.S.Chern. *Topics in Defferential Geometry*, Ch. III "Theory of Connections" (The Institute for Advanced Study, Princeton, 1951)
- [3] H. Weyl, *Electron and Gravitation.*, Z. Phys. **56** (1929) 330
- [4] E. Cartan, Sur les variétés à connexion affine et la théorie de la relativité généralisée. Annales Sci. Ecole Norm. Sup. 40 (1923) 325; 41 (1924) 1; 42 (1925) 17
- [5] E.Majorana. Teoria Relativistica di Particelle con Momento Intrinseco Arbitrario, Nuovo Cimento 9 (1932) 335
- [6] M. Fierz. Uber die relativistische Theorie kräftefreier Teilchen mit beliebigem Spin, Helv. Phys. Acta. 12 (1939) 3.
- [7] M. Fierz and W. Pauli. On Relativistic Wave Equations for Particles of Arbitrary Spin in an Electromagnetic Field, Proc. Roy. Soc. A173 (1939) 211.
- [8] P.A.M.Dirac. Relativistic wave equations, Proc. Roy. Soc. A155 (1936) 447;
 Unitary Representation of the Lorentz Group, Proc. Roy. Soc. A183 (1944) 284.
- [9] V. L. Ginzburg and I. E. Tamm, To the theory of spin, ZETP 17 (1947) 227
- [10] S. Weinberg, Feynman Rules For Any Spin, Phys. Rev. 133 (1964) B1318.
- [11] P. Minkowski. Versuch einer konsistenten Theorie eines Spin-2 Mesons, Helv. Phys. Acta. 32 (1966) 477
- [12] Y. Nambu, Relativistic groups and infinite-component fields, In *N. Svartholm, Elementary Particle Theory. Proceedings Of The Nobel Symposium Held 1968 At Lerum, Sweden*, Stockholm 1968, 105-117
- [13] J.Schwinger, Particles, Sourses, and Fields (Addison-Wesley, Reading, MA, 1970)
- [14] P. Ramond, Dual Theory for Free Fermions, Phys. Rev. D 3 (1971) 2415.

- [15] L. P. S. Singh and C. R. Hagen, Lagrangian formulation for arbitrary spin. I. The boson case, Phys. Rev. D9 (1974) 898
- [16] C.Fronsdal, Massless fields with integer spin, Phys.Rev. **D18** (1978) 3624
- [17] A. K. Bengtsson, I. Bengtsson and L. Brink, "Cubic Interaction Terms For Arbitrary Spin," Nucl. Phys. B **227** (1983) 31.
- [18] A. K. Bengtsson, I. Bengtsson and L. Brink, "Cubic Interaction Terms For Arbitrarily Extended Supermultiplets," Nucl. Phys. B 227 (1983) 41.
- [19] F. A. Berends, G. J. H Burgers and H. Van Dam, On the Theoretical problems in Constructing Interactions Involving Higher-Spin Massless Particles, Nucl. Phys. B 260 (1985) 295.
- [20] E.S. Fradkin and M.A. Vasiliev, Dokl. Acad. Nauk. 29, 1100 (1986); Ann. of Phys. 177, 63 (1987).
- [21] M.A.Vasiliev et. al. Nonlinear Higher Spin Theories in Various Domensions hep-th/0503128
- [22] A. Sagnotti, E. Sezgin and P. Sundell, On higher spins with a strong Sp(2,R) condition, arXiv:hep-th/0501156.
- [23] R. R. Metsaev, Cubic interaction vertices of massive and massless higher spin fields, Nucl. Phys. B 759 (2006) 147
- [24] R. Manvelyan, K. Mkrtchyan and W. Ruhl, General trilinear interaction for arbitrary even higher spin gauge fields, Nucl. Phys. B 836 (2010) 204
- [25] S. Guttenberg and G. Savvidy, Schwinger-Fronsdal Theory of Abelian Tensor Gauge Fields, SIGMA 4 (2008) 061 [arXiv:0804.0522 [hep-th]].
- [26] G. Savvidy, Non-Abelian tensor gauge fields: Generalization of Yang-Mills theory, Phys. Lett. B 625 (2005) 341
- [27] G. Savvidy, Non-abelian tensor gauge fields. I, Int. J. Mod. Phys. A 21 (2006) 4931;
- [28] G. Savvidy, Non-abelian tensor gauge fields. II, Int. J. Mod. Phys. A 21 (2006) 4959;

- [29] G. Savvidy, Extension of the Poincaré Group and Non-Abelian Tensor Gauge Fields, Int. J. Mod. Phys. A **25** (2010) 5765 [arXiv:1006.3005 [hep-th]].
- [30] H.Yukawa, Quantum Theory of Non-Local Fields. Part I. Free Fields, Phys. Rev. 77 (1950) 219
- [31] M. Fierz, Non-Local Fields, Phys. Rev. 78 (1950) 184
- [32] E. Wigner, Invariant Quantum Mechanical Equations of Motion, in Theoretical Physics ed. A.Salam (International Atomic Energy, Vienna, 1963) p 59
- [33] S. R. Coleman and J. Mandula, All possible symmetries of the S matrix, Phys. Rev. 159 (1967) 1251.
- [34] G. Savvidy, Non-Abelian Tensor Gauge Fields, Proc. Steklov Inst. Math. 272 (2011) 201 [arXiv:1004.4456 [hep-th]].
- [35] I. Antoniadis, L. Brink and G. Savvidy, Extensions of the Poincaré group, J. Math. Phys. **52** (2011) 072303 [arXiv:1103.2456 [hep-th]].
- [36] G. Savvidy, Invariant scalar product on extended Poincaré algebra, J. Phys. A 47 (2014) 5, 055204 [arXiv:1308.2695 [hep-th]].
- [37] S. J. Parke and T. R. Taylor, An Amplitude for n Gluon Scattering, Phys. Rev. Lett. 56 (1986) 2459.
- [38] L. J. Dixon, Calculating scattering amplitudes efficiently, arXiv:hep-ph/9601359.
- [39] E. Witten, Perturbative gauge theory as a string theory in twistor space, Commun. Math. Phys. **252** (2004) 189 [arXiv:hep-th/0312171].
- [40] R. Britto, F. Cachazo, B. Feng and E. Witten, Direct proof of tree-level recursion relation in Yang-Mills theory, Phys. Rev. Lett. 94 (2005) 181602 [arXiv:hep-th/0501052].
- [41] P. Benincasa and F. Cachazo, Consistency Conditions on the S-Matrix of Massless Particles, arXiv:0705.4305 [hep-th].
- [42] N. Arkani-Hamed, T. C. Huang and Y. t. Huang, Scattering Amplitudes For All Masses and Spins, arXiv:1709.04891 [hep-th].

- [43] G. Georgiou and G. Savvidy, Production of non-Abelian tensor gauge bosons. Tree amplitudes and BCFW recursion relation, Int. J. Mod. Phys. A 26 (2011) 2537 [arXiv:1007.3756 [hep-th]].
- [44] G. Savvidy, Asymptotic freedom of non-Abelian tensor gauge fields, Phys. Lett. B **732** (2014) 150.
- [45] G. Savvidy, Generalisation of the Yang-Mills Theory, Int. J. Mod. Phys. A **31** (2016) no.01, 1630003 doi:10.1142/S0217751X16300039, 10.1142/9789814725569_0015 [arXiv:1511.00274 [hep-th]].