

Using the CIFIST grid of CO5BOLD 3D model atmospheres to study the effects of stellar granulation on photometric colours.

I. Grids of 3D corrections in the UBVRI, 2MASS, Hipparcos, Gaia, and SDSS systems.

P. Bonifacio¹, E. Caffau¹, H.-G. Ludwig^{2,1}, M. Steffen^{3,1}, F. Castelli⁴, A. J. Gallagher^{5,1}, A. Kučinskas⁶, D. Prakashavičius⁶, R. Cayrel¹, B. Freytag⁷, B. Plez⁸, and D. Homeier²

¹ GEPI, Observatoire de Paris, PSL Research University, CNRS, Place Jules Janssen, 92195 Meudon, France

² Zentrum für Astronomie der Universität Heidelberg, Landessternwarte, Königstuhl 12, 69117 Heidelberg, Germany

³ Leibniz-Institut für Astrophysik Potsdam, An der Sternwarte 16, D-14482 Potsdam, Germany

⁴ Istituto Nazionale di Astrofisica, Osservatorio Astronomico di Trieste, Via Tiepolo 11, I-34131 Trieste, Italy

⁵ Max Planck Institut für Astronomie, D-69117 Heidelberg, Germany

⁶ Astronomical Observatory, Institute of Theoretical Physics and Astronomy, Vilnius University, Saulėtekio al. 3, Vilnius LT-10257, Lithuania

⁷ Department of Physics and Astronomy at Uppsala University, Regementsvägen 1, Box 516, 75120, Uppsala, Sweden

⁸ LUPM, Université de Montpellier, CNRS, UMR 5299, 34095 Montpellier Cedex 05, France

Received ...; Accepted ...

ABSTRACT

Context. The atmospheres of cool stars are temporally and spatially inhomogeneous due to the effects of convection. The influence of this inhomogeneity, referred to as granulation, on colours has never been investigated over a large range of effective temperatures and gravities.

Aims. We aim to study, in a quantitative way, the impact of granulation on colours.

Methods. We use the CIFIST (Cosmological Impact of the First Stars) grid of CO⁵BOLD (COnservative COde for the COmputation of COmpressible COnvection in a BOx of L Dimensions, L=2,3) hydrodynamical models to compute emerging fluxes. These in turn are used to compute theoretical colours in the *UBVRI*, 2MASS, Hipparcos, Gaia and SDSS systems. Every CO⁵BOLD model has a corresponding one dimensional (1D) plane-parallel LHD (Lagrangian HydroDynamics) model computed for the same atmospheric parameters, which we used to define a "3D correction" that can be applied to colours computed from fluxes computed from any 1D model atmosphere code. As an example, we illustrate these corrections applied to colours computed from ATLAS models.

Results. The 3D corrections on colours are generally small, of the order of a few hundredths of a magnitude, yet they are far from negligible. We find that ignoring granulation effects can lead to underestimation of T_{eff} by up to 200 K and overestimation of gravity by up to 0.5 dex, when using colours as diagnostics. We have identified a major shortcoming in how scattering is treated in the current version of the CIFIST grid, which could lead to offsets of the order 0.01 mag, especially for colours involving blue and UV bands. We have investigated the Gaia and Hipparcos photometric systems and found that the $(G - H_p)$, $(BP - RP)$ diagram is immune to the effects of granulation. In addition, we point to the potential of the RVS photometry as a metallicity diagnostic.

Conclusions. Our investigation shows that the effects of granulation should not be neglected if one wants to use colours as diagnostics of the stellar parameters of F,G,K stars. A limitation is that scattering is treated as true absorption in our current computations, thus our 3D corrections are likely an upper limit to the true effect. We are already computing the next generation of the CIFIST grid, using an approximate treatment of scattering.

Key words. Hydrodynamics - Stars: atmospheres

1. Introduction

The use of theoretical colours computed from model atmospheres is widespread, however, so far all available grids of theoretical colours (e.g. Bessell et al. 1998; Castelli 1999; Öneħag et al. 2009; Casagrande & Vandenberg 2014) are based on one dimensional (1D) static model atmospheres. Extensive computations of fluxes from the three-dimensional (3D) models of the STAGGER grid (Magic et al. 2013) have been done by Magic et al. (2015), however, so-far only the limb-darkening coefficients have been published. In a series of two papers, we wish to ex-

plore the parameter space covered by the CIFIST (Cosmological Impact of the First Stars) grid of 3D models (Ludwig et al. 2009, and Ludwig et al., in preparation) that has been computed with the CO⁵BOLD (COnservative COde for the COmputation of COmpressible COnvection in a BOx of L Dimensions, L=2,3) code (Freytag et al. 2012). This effort expands on the pioneering computations of colours from CO⁵BOLD models presented earlier by Kučinskas et al. (2005) and Kučinskas et al. (2009).

For comparison with the CO⁵BOLD models, we computed two grids of 1D model atmospheres. The first was computed using the LHD (Lagrangian HydroDynamics) 1D model atmosphere code (Caffau & Ludwig 2007) assuming a mixing length

Table 1. 2MASS in-band fluxes and corresponding magnitudes from Cohen et al. (2003)

Star	FJ	J	FH	H	FK_S	K_S
	Wcm^{-2}	mag	Wcm^{-2}	mag	Wcm^{-2}	mag
	$\times 10^{-14}$		$\times 10^{-14}$		$\times 10^{-14}$	
Vega	5.076	0.001	2.857	-0.005	1.121	0.001
0 mag	5.082	0.000	2.843	0.000	1.122	0.000

parameter $\alpha_{\text{LHD}} = 1.0$ ¹. The value 1.0 is closest to the value used in the Castelli & Kurucz (2003) grid of ATLAS models; therefore we chose this value here. The effects of different choices of the mixing length parameter in the reference 1D model are further explored in Paper II of the series (Kučinskas et al. 2018). The second grid was computed using ATLAS (Kurucz 2005) assuming a mixing length parameter 1.25, and the Castelli & Kurucz (2003) Opacity Distribution Functions (ODFs) with 2 km s^{-1} microturbulence. The choice of mixing length and microturbulent velocity were made to be consistent with the Castelli & Kurucz (2003) grid of ATLAS models. Both grids of models were specifically computed to match atmospheric parameters of the CO⁵BOLD models in the CIFIST grid.

The purpose of the present paper (Paper I of the series) is to present the computational methods employed and compare the fluxes and colours computed from 1D static model atmospheres to those computed from 3D hydrodynamical model atmospheres. We discuss in detail the effects of the treatment of scattering both in the computation of model atmospheres and in the computation of emerging fluxes. For the *UBVRI*, 2MASS, Hipparcos, Gaia and SDSS *ugriz* systems we provide grids of “3D corrections” that can be applied to colours computed from 1D model atmospheres. We also provide a corresponding grid of ATLAS 3D-corrected colours.

In the second paper of the series (Kučinskas et al. 2018), we select ten model atmospheres for two different metallicities (0.0 and -2.0) that cover the main phases of stellar evolution (main sequence, turn off, sub giant, red giant) and explore the effects of granulation in the corresponding parameter space.

There are some differences of approach to the computation of colours in the two papers, which are explained in detail in Kučinskas et al. (2018). In both papers, we adopt a strictly differential approach, discussing only 3D-1D differences, which makes the above-mentioned differences irrelevant.

2. Computation of synthetic colours

2.1. Photometric systems and zero-points

The problem of computing synthetic colours has been thoroughly addressed in many excellent papers (e.g. Bessell et al. 1998; Castelli 1999; Önehag et al. 2009; Castelli & Kurucz 2006; Bessell & Murphy 2012; Casagrande & Vandenberg 2014). Our own approach to the problem has been extensively described in Bonifacio et al. (2017). We refer the reader to that paper for justification on our choices of zero points for the synthetic photometry and bolometric magnitudes. In this paper, we investigate the *UBVRI* and Hipparcos-Tycho systems, using the bandpasses defined by Bessell & Murphy (2012), the 2MASS system (Cohen et al. 2003) and the Gaia system, as defined by

¹ Associated to each model in the CIFIST grid we always compute three LHD models with $\alpha_{\text{LHD}} = 0.5, 1.0, 2.0$.

the pre-launch bandpasses². The above are all “Vega” systems, and we tie the synthetic photometry to Vega using the CALSPEC flux of VEGA³ and assuming all magnitudes for Vega to be 0.03, except for the 2MASS magnitudes where we assume for Vega $J=0.001$, $H=-0.005$ and $K=0.001$, in order to be consistent with the zero-magnitude fluxes of Cohen et al. (2003, see table 1). As explained in Bonifacio et al. (2017), we did not define the zero points of the “Vega” systems by using a model atmosphere of Vega, since such a model cannot be computed with CO⁵BOLD, but we assumed a radius of $1 R_{\odot}$ and a distance of 10 pc for each model atmosphere and applied to this flux the same zero points adopted for the observed fluxes. We also investigated the SDSS *ugriz* system, which is AB type (Oke & Gunn 1983), tied to the standards defined in Fukugita et al. (1996). In the SDSS catalogue, what are reported are not magnitudes, but “luptitudes” defined in Lupton et al. (1999). A “luptitude” is defined as

$$m = -\frac{2.5}{\ln(10)} [\text{asinh}((f/f_0)/(2b)) + \ln(b)] \quad (1)$$

where f is the flux of the object, f_0 is the flux of the 0 magnitude object, and b is a constant, called the softening parameter. The softening parameter determines the magnitude of an object with zero flux, that is, a non-detection in the survey; this is the magnitude limit of the survey. It is important to keep in mind that objects with magnitudes equal or very close to this limit have essentially undefined colours and therefore should not be compared to theoretical colours. Other SDSS-like systems that are becoming increasingly popular and are used in many telescopes may use magnitudes rather than luptitudes. In order to be able to compare our computed colours directly to the SDSS catalogue, we computed luptitudes rather than magnitudes to derive SDSS colours. We note however that the difference in 3D correction between colours computed using luptitudes and magnitudes is always less than or equal to 10^{-5} mag. So for any practical purpose the 3D corrections provided in Tables A.13 to A.20 can also be applied to any SDSS-like system.

2.2. Computation of emerging fluxes

As described in greater detail in Bonifacio et al. (2017) we used the NLTE3D code to compute the emerging fluxes from the CO⁵BOLD models. Each 3D model is formed by a time series of 3D structures, usually of the order of several hundreds, that we call snapshots. Consecutive snapshots are often correlated, and including correlated events does not improve the statistic. As already mentioned, computations with 3D structures, whether they be fluxes or spectrum synthesis, are time consuming. We then select a representative number of statistically independent snapshots able to represent the model. Usually we select about 20 snapshots from a model, because we think this is a good compromise between the computation time for the fluxes or spectrum synthesis, and the representativeness of the selection for the model. This number has been found to be representative in a detailed investigation on the solar model (Caffau 2009) and we assume this to be the case for any CO⁵BOLD model. The snapshot selection is made so that its statistical properties are the same as those of the total ensemble of computed snapshots; we refer to Caffau & Ludwig (2007) for further details on this.

² <http://www.cosmos.esa.int/documents/29201/302420/normalisedPassbands.txt/a65b04bd-4060-44fa-be36-91975f2bd58a>

³ ftp://ftp.stsci.edu/cdbs/current_calspec/alpha_lyr_stis_008.fits

For each model, we averaged the emerging flux from the selected snapshots.

We relied on the Castelli & Kurucz (2003) opacity distribution functions (ODFs) in order to take into account the line opacity. We used the “LITTLE” ODFs, with 1212 frequency points, rather than the “BIG” ODFs that only have 328 frequency points. The continuum opacities were computed using the `iondis` and `opal` routines of the `Linfor3D` spectrum synthesis code (Steffen et al. 2017; Gallagher et al. 2017).

This approach is inconsistent, since the models have been computed using MARCS continuum and line opacities, while the fluxes have been computed assuming `iondis` and `opal` continuum opacities and ATLAS line opacities. Since we rely mainly on a differential approach (3D-1D), we do not expect this inconsistency to be a major shortcoming, as we explain below.

2.3. The treatment of scattering

Scattering opacity has been treated as true absorption and this, in our view is the main physical limitation of our computation. The treatment of scattering enters on two occasions in our problem: first in the computation of the model, where its treatment may influence the temperature structure of the model, then in the computation of the emerging flux from the model. The choice of treating scattering as true absorption is made for computational reasons. A fully consistent treatment of scattering requires that the mean intensity J_ν be evaluated at each point in the computational box during the computation. Collet et al. (2011) questioned the soundness of treating scattering as true absorption and showed that it may lead to a mean temperature structure that is substantially warmer than that obtained when explicitly treating the scattering. On the other hand, they also noticed that a better and computationally less expensive approximation is to treat scattering as true absorption in the optically thick layers and ignore the scattering opacity in the optically thin layers as described in Hayek et al. (2010). In this approximation one does not really treat scattering while computing the radiative transfer, but uses an appropriate opacity table. Ludwig & Steffen (2012) computed a CO⁵BOLD model using the Hayek et al. (2010) approximation and agreed qualitatively with the conclusions of Collet et al. (2011), although in their case the influence of the treatment of scattering was clearly less important than in the example studied by Collet et al. (2011). In the Ludwig & Steffen (2012) study, the difference in mean temperature between the “true absorption” and “approximate scattering” models is about 100 K at $\log(\tau)=-4$, while in the Collet et al. (2011) study, this difference is about five times larger. The two models used by Ludwig & Steffen (2012) and Collet et al. (2011) have very similar atmospheric parameters, therefore we attribute this difference between the two studies to differences between the two codes used to compute the hydrodynamical simulations. We have used this model, with $T_{\text{eff}}=5000$ K, $\log g=2.5$, and metallicity -3.0 , and another model, with the same atmospheric parameters, but computed treating scattering as true absorption, to assess the influence of scattering on the computed fluxes and colours. This particular model was chosen because it was in this case that Collet et al. (2011) obtained the largest effects of scattering. Therefore these results may be considered an upper limit to the effects of scattering.

NLTE3D can treat scattering in detail (no approximations) in the computation of the flux from a given model, but the computation is much more intensive. We limited the comparison to a single snapshot. In order to check that a single snapshot is representative, we computed the colours for each of the 20 snapshots




Fig. 1. Ratio of the fluxes of one snapshot of the CO⁵BOLD models d3t50g25mm30n01 (scattering treated as true absorption) and d3t50g25mm30n02 (scattering in the Hayek et al. (2010) approximation) to the fluxes computed from their corresponding LHD models under different assumptions. The solid line (black) corresponds to the d3t50g25mm30n02 model, and the flux has been computed fully taking into account scattering. The dashed line (red) corresponds to the d3t50g25mm30n01 model and the flux has been computed treating scattering as true absorption. The dotted line (blue) corresponds again to the d3t50g25mm30n01 model, but in this case scattering has been properly taken into account in the flux computation.

that are our selection for this model. The snapshot-to-snapshot difference in colours was always below 0.01 mag. As we shall see below, the differences due to the different assumptions on scattering are always larger than this. Thus our comparison is meaningful.

In Fig. 1, we show the ratio of the flux computed from the CO⁵BOLD models to that of their corresponding LHD models. The dotted line corresponds to the model for which scattering has been treated as true absorption, but scattering has been taken into account in the flux computation. The dashed line corresponds to the same model; however in this case, consistently with what was done for the model computation, also in the flux computation scattering has been treated as true absorption. Finally the solid line corresponds to the model for which scattering has been treated in the approximation of Hayek et al. (2010) and scattering has been fully taken into account in the flux computation. The same approximations apply to both the 1D and 3D models. We present this in the UV-visible region where the differences due to the different treatment of scattering are largest. The first thing that is obvious is that the treatment of scattering mostly affects the spectral regions characterised by strong lines. Treating scattering as true absorption results in a flux from the CO⁵BOLD model that can be 1.2 to almost 1.6 times larger than that in the corresponding LHD model. This is not so in the case where scattering is treated using the Hayek et al. (2010) approximation in the model, and fully treated in the flux computation. In this case, the flux from the CO⁵BOLD model is only about

5% larger than that from the LHD model, but the flux ratio varies rather smoothly with wavelength, unlike what is seen when treating scattering as true absorption. The “hybrid” case, in which we treat scattering in the flux computation using a model which has been computed treating scattering as true absorption, is interesting. In the current version of the CIFIST grid (Ludwig et al. 2009) all models have been computed treating scattering as true absorption. A new version of the grid is currently being computed in which scattering is treated in the Hayek et al. (2010) approximation. In the following, for simplicity, we refer to the case where scattering has been treated as true absorption both in the model and in the flux calculations as true absorption; we refer to the case in which the model has been computed treating scattering in the Hayek et al. (2010) approximation and the flux has been computed fully treating scattering as “scattering”; we refer to the case in which the model has been computed treating scattering as true absorption but the flux has been computed fully taking into account scattering, as “hybrid”.

Is there anything to be gained in treating scattering in the flux computation, from the current version of the CIFIST grid? Figure 1 suggests that this is not the case. Although in this hybrid case the run of the 3D/1D ratio has a shape that is very similar to that of the scattering case, it is typically 10% smaller than what is expected in the scattering case. We are ultimately interested in colours and therefore we computed the colour-corrections (3D-1D) for all three cases shown in Fig. 1. Assuming that the scattering computations are the ones closest to the truth we find the difference in 3D correction between the scattering case and the true absorption case is on average -0.013 mag while this difference is -0.009 mag for the hybrid case. Obviously this is largest, in absolute value, for the bluest colours. For $U - B$ we find a difference in the 3D correction of -0.024 for the true absorption case and -0.18 for the hybrid case, while for the $V - K$ colour we find -0.018 for the true absorption case and -0.014 for the hybrid case. In the wavelength range 500 nm to 2500 nm (not shown in Fig. 1) the ratio of the scattering 3D/1D flux ratio to the true absorption 3D/1D flux ratio varies from 0.95 to 1.017 with a mean value of 1.003.

From this test, on one model we decided to use the fluxes computed from the CIFIST grid treating scattering as true absorption. Although the hybrid approach does present a flux that has a similar shape to the consistent scattering approach, we cannot convincingly conclude that this improvement is worth the considerable computational effort needed to perform hybrid computations of the whole grid, considering that the differences in 3D corrections are of only a few thousandths of a magnitude.

All fluxes from the ATLAS models were computed with the ATLAS code, where true absorption was assumed for the continuum scattering and no line scattering was considered. These were also the assumptions adopted in the computation of the ATLAS models.

3. Results

We adopted a differential approach: for each colour, magnitude, or bolometric correction, we computed the “3D correction” as colour (CO⁵BOLD) - colour (LHD). These corrections are provided in the Tables A.1 to A.20. As an example of how these corrections have to be used, we also provide the ATLAS 3D-corrected colours. For each CO⁵BOLD model we computed an ATLAS model with exactly the same parameters and its emerging flux. From this flux, we computed colours and the 3D corrected colour is then simply colour (ATLAS) + “3D correction”.




Fig. 2. A $u - g$ vs. $g - z$ colour-colour diagram for solar metallicity (bottom panel) and -2.0 (upper panel). The solid lines are the lines of constant temperature and constant surface gravity defined by the Castelli & Kurucz (2003) grid. The black dots correspond to our ATLAS models, that sample points within the Castelli & Kurucz (2003) grid that correspond to the parameters of the models in the CIFIST grid. The Castelli & Kurucz (2003) grid has steps of 250 K in T_{eff} and 0.5 dex in $\log g$. The red squares are the “3D-corrected” colours.

Clearly, this same procedure can be applied to colours computed from any grid of 1D model atmospheres.

The ATLAS models computed for this paper are almost identical to those of the Castelli & Kurucz (2003) grid. The only difference is that we used updated ODFs for which the contribution of the H₂O lines has been revised. This has only a minor effect on models cooler than 4500 K. Our computed colours from our




Fig. 3. The 3D–1D corrections for the $g - z$ and $u - g$ colours for two metallicities: 0.0 and -3.0 .

ATLAS models precisely fall on the theoretical curves defined in the Castelli & Kurucz (2003) grid, and may be slightly off for some colours when the temperature decreases due to H_2O opacity.

Figure 2 depicts colour-colour diagrams for $g - z$ and $u - g$ bands for metallicities at $[\text{M}/\text{H}]=0.0$ and -2.0 . If the metallicity is known (e.g. for stars in a cluster) such a diagram can be used as a diagnostic in T_{eff} and $\log g$. While the corrections are generally small around the solar chemical composition, they increase with decreasing metallicity. At low metallicity, the 3D-corrected $g - z$ tends to be larger (i.e. redder), therefore, hotter temperatures will be found from observed $g-z$ colours. At the hot end of our model grid, this change is rather large, of the order of 200 K. At solar metallicity the situation is similar for the hotter models, however for the cooler models ($T_{\text{eff}} < 4500$ K) the situation reverses and the 3D-corrected $g - z$ is smaller (bluer); thus, for a given observed $g - z$, one would infer a cooler T_{eff} , although by only about 50 K. A similar behaviour is observed for the $u - g$ colour at low metallicity; the 3D-corrected $u - g$ is larger, thus, for an observed $u - g$, one infers a lower gravity, by almost 0.5 dex. At solar metallicity the situation is different. For the hotter

models, there is hardly any change in $u - g$, but below 5500 K the 3D-corrected $u - g$ is smaller, implying gravities that are larger, by about 0.1 dex. The situation is illustrated in Fig.3, where the 3D-1D corrections for metallicities 0.0 and -3.0 are plotted as a function of T_{eff} . To guide the eye, the corrections relative to models with the same surface gravity are connected by a solid line and a different colour is used for each surface gravity as given in the legend.

In Fig. 4 we show the $U - B$ versus $V - K$ diagram, which is morphologically similar to the $u - g$ versus $g - z$, and thus the effect of granulation is similar. However, the use of the longer baseline colour $V - K$ makes another effect apparent. At all metallicities, the 3D-corrected colours are redder for the hotter models and bluer for the cooler models. This results in a more compressed scale in $V - K$. It is also apparent that the difference in inferred temperature is smaller in this case, less than 100 K for all gravities and metallicities.

In Fig. 5 we compare two colours that are often used as temperature indicators: $V - I$ and $J - K$. There is a tight correlation between the two colours, up to about 4250 K when the relation splits up and the two colours respond differently to changes in




Fig. 4. A $U - B$ vs. $V - K$ colour-colour diagram for solar metallicity (bottom panel) and -2.0 (upper panel). The meaning of the lines and symbols is the same as in Fig. 2.

gravity. The “3D-corrected” colours show exactly the same behaviour, except that the relation is offset by 0.05 mag.

Let us turn our attention to the Gaia photometry. We should warn the reader that the Gaia colours have been computed using the pre-launch transmission curves. This is useful to gain an understanding of the general properties of this photometry. Detailed comparison with the observations will be possible when the post-launch transmission curves become available, after the Gaia Data Release 2. We notice that the wide band used for the Gaia photometry, G , is sufficiently different from the analogue band of Hipparcos, H_p , that the colour $G-H_p$ is tightly correlated with effective temperature. Even more interestingly, it is very tightly correlated with the colour that can be formed from the Blue Prism (BP) and Red Prism (RP) spectra, $BP-RP$. In




Fig. 5. A $J - K$ vs. $V - I$ colour-colour diagram for solar metallicity (bottom panel) and -2.0 (upper panel). The meaning of the lines and symbols is the same as in Fig. 2.

Fig. 6 we show how this colour-colour diagram provides a very tight correlation, essentially putting all the stars of any metallicity or gravity on the same relation. It is interesting to note that the 3D corrections do not destroy this correlation, since all the points move along the curve. This should be a very robust correlation and should prove useful to determine the effective pass-bands of the Gaia system and reddening and to detect binary systems. Any star that is an outlier with respect to such a correlation should be considered as a potential binary system. The $G-H_p$ correlates with any temperature-dependent colour, in particular with any colour that measures the slope of the Paschen continuum. With other colours, however, the relation splits due to gravity and metallicity dependencies. As an example, we show its correlation with the Tycho $B_T - V_T$ colour in Fig. 7. In this case the




Fig. 6. The colour $G-H_p$ as a function of the $BP-RP$ colour. Colours computed from models of different metallicity are shown with a different colour: black for $[M/H]=0.0$, red for $[M/H]=-1.0$, green for $[M/H]=-2.0$ and blue for $[M/H]=-3.0$. Open circles are the colours computed from ATLAS models, while filled squares are the corresponding 3D-corrected colours. We stress that the Gaia bands are the pre-launch bands.




Fig. 7. The colour $G-H_p$ as a function of the $BT-VT$ colour. Symbols are the same as in Fig. 6. The dashed lines connect the points from the Castelli & Kurucz (2003) grid.

3D corrections induce relations that are distinctly different from what is deduced from the 1D model atmospheres. Because of the splitting seen due to surface gravity and metallicity, such a colour-colour diagram has a clear diagnostic potential, however the theoretical colours must be properly modelled.

Finally let us turn our attention to the potential of the RVS colour as a metallicity diagnostic. At the faint end of the sensi-




Fig. 8. The CaT colour as a function of the $BP-RP$ colour for models with $\log g=4.0$. Filled circles are the colours from the ATLAS models and filled squares are the 3D-corrected colours.

tivity of the RVS spectrograph (Katz et al. 2004), in the interval $G=15$ to $G=17$, it is likely that the spectra will be too noisy to allow a metallicity determination. Yet, since the spectra will be flux-calibrated one can derive an RVS magnitude, that is, a narrow-band filter centered on the IR $Ca\ II$ triplet. In ground-based observations, such a filter is not very useful, since the signal of the $Ca\ II$ lines is severely diluted by the signal coming from the atmospheric OH emission lines. Data taken from space, however, like Gaia, do not suffer from such a limitation. In Fig. 8 we show the behaviour of the CaT colour as a function of the $BP-RP$ colour. We show several metallicities for $\log g=4.0$ and it is clear that even a precision of 0.02 mag in this colour should allow an estimate of the metallicity with a precision of 0.5 dex, down to a metallicity of -3.0 , at least. The colour is not strongly sensitive to gravity. However Gaia will also provide the parallaxes, thus the surface gravity will be known. It is interesting to see that the 3D corrections move the points along the curve defined by the 1D models, so that the metallicity sensitivity of the colour remains the same. The metallicity information from the RVS magnitude will be complementary to that coming from the full spectral energy distribution, derived from the prism spectra, for stars for which both will be available.

4. Conclusions

We have used the CIFIST grid of CO⁵BOLD models to investigate the effects of granulation on fluxes and colours of stars of spectral type F, G, and K. Our investigation is exploratory, since we realise that treating scattering as true absorption, as done in the current version of the CIFIST models, leads to incorrect results in the computed emerging fluxes, especially in the blue and UV regions. The influence of this assumption can be quantified in terms of colours and is of the order of 0.02 magnitudes for colours involving blue or UV bands. While the next generation of the CIFIST grid is being computed using the approximate treatment of scattering suggested by Hayek et al. (2010), we be-

lieve it is still interesting to publish the results obtained from the current version of the CIFIST grid.

The effects of granulation on colours are generally small, of a few hundredths of a magnitude; they cannot, however be considered negligible. On some colours, such as $g - z$, the effect can translate to a temperature estimate that is hotter by 200 K.

We publish tables with 3D corrections that can be applied to colours computed from any 1D model atmosphere. For $T_{\text{eff}} \geq 5000\text{K}$, the corrections are smooth enough, as a function of atmospheric parameters, that it is possible to interpolate the corrections between grid points; thus the coarseness of the CIFIST grid should not be a major limitation. However at the cool end there are still far too few models to allow a reliable interpolation.

We have investigated the effects on the Gaia photometric system, although only the pre-launch transmission curves are available. We confirm that a tight correlation between the $G - H_p$ colour and the BP-RP colour should exist, and that this correlation should be the same also after granulation effects are taken into account. We have also investigated the potential of the $(G - RVS) - (BP - RP)$ colour as a metallicity diagnostic and we confirm that it is indeed a good indicator, even after 3D corrections have been taken into account, provided this colour can be measured with a precision of 0.02 mag or better. If one is only interested in selecting stars that are more metal-poor than -3.0 , the colour is still very powerful even if the precision is of the order of 0.05 mag.

Acknowledgements. We are grateful to the referee, Santi Cassisi, whose report helped us to improve the paper. This project has been supported by foundation MERAC and Observatoire de Paris. We acknowledge financial support from CNRS Institut National de Sciences de l'Univers Programme National de Cosmologie et Galaxies and Programme National de Physique Stellaire. This work was granted access to the HPC resources of MesoPSL financed by the Région Île de France and the project Equip@Meso (reference ANR-10-EQPX-29-01) of the programme Investissements d'Avenir supervised by the Agence Nationale pour la Recherche. HGL and DH acknowledge financial support by the Sonderforschungsbereich SFB 881 "The Milky Way System" (subproject A4) of the German Research Foundation (DFG). This research was supported by a grant (MIP-089/2015) from the Research Council of Lithuania.

References

- Bessell, M. S., Castelli, F., & Plez, B. 1998, *A&A*, 333, 231
- Bessell, M., & Murphy, S. 2012, *PASP*, 124, 140
- Bonifacio, P., Caffau, E., Ludwig, H.-G., et al. 2017 *Memorie della Società Astronomica Italiana*, 88, 90
- Caffau, E. 2009, Thèse de Doctorat en Astronomie et Astrophysique, Observatoire de Paris, Ecole Doctorale Astronomie et Astrophysique d'Île-de-France.
- Caffau, E., & Ludwig, H.-G. 2007, *A&A*, 467, L11
- Casagrande, L., & Vandenberg, D. A. 2014, *MNRAS*, 444, 392
- Castelli, F. 1999, *A&A*, 346, 564
- Castelli, F. & Kurucz, R. L. 2003, in *IAU Symposium*, ed. N. Piskunov, W. W. Weiss, & D. F. Gray, 20P, arXiv:astro-ph/0405087v1
- Castelli, F., & Kurucz, R. L. 2006, *A&A*, 454, 333
- Cohen, M., Wheaton, W. A., & Megeath, S. T. 2003, *AJ*, 126, 1090
- Collet, R., Hayek, W., Asplund, M., et al. 2011, *A&A*, 528, A32
- Freytag, B., Steffen, M., Ludwig, H.-G., et al. 2012, *J. Comp. Phys.*, 231, 919
- Fukugita, M., Ichikawa, T., Gunn, J. E., et al. 1996, *AJ*, 111, 1748
- Gallagher, A. J., Steffen, M., Caffau, E., et al. 2017, *Mem. Soc. Astron. Italiana*, 88, 82
- Hayek, W., Asplund, M., Carlsson, M., et al. 2010, *A&A*, 517, A49
- Katz, D., Munari, U., Cropper, M., et al. 2004, *MNRAS*, 354, 1223
- Kučinskas, A., Hauschildt, P.H., Ludwig, H.-G., et al. 2005, *A&A*, 442, 281
- Kučinskas, A., Ludwig, H.-G., Caffau, E., & Steffen, M. 2009, *Mem. Soc. Astron. Italiana*, 80, 723
- Kučinskas, A., Klevas, J., Ludwig, H.-G., Bonifacio, P., Steffen, M., & Caffau, E., 2018, *A&A* in press, arXiv:1802.00073.
- Kurucz, R. L. 2005, *Memorie della Società Astronomica Italiana Supplementi*, 8, 14
- Ludwig, H.-G., & Steffen, M. 2012, *Astrophysics and Space Science Proceedings*, 26, 125
- Ludwig, H.-G., Caffau, E., Steffen, M., et al. 2009, *Mem. Soc. Astron. Italiana*, 80, 711
- Lupton, R. H., Gunn, J. E., & Szalay, A. S. 1999, *AJ*, 118, 1406
- Magic, Z., Collet, R., Asplund, M., et al. 2013, *A&A*, 557, A26
- Magic, Z., Chiavassa, A., Collet, R., & Asplund, M. 2015, *A&A*, 573, A90
- Önehag, A., Gustafsson, B., Eriksson, K., & Edvardsson, B. 2009, *A&A*, 498, 527
- Oke, J. B., & Gunn, J. E. 1983, *ApJ*, 266, 713
- Steffen, M., Ludwig, H.-G., Wedemeyer-Böhm, S., & Gallagher, A. J. 2017, *Linfor3D User Manual* <http://www.aip.de/Members/msteffen/linfor3d>

Appendix A: Tables of colours and colour corrections.

In this Appendix we present the 3D-1D colour corrections for colours in the Johnson-Cousins, 2MASS, Hipparcos, Gaia and SDSS systems. Bolometric corrections are also provided for the Gaia G magnitude. For each colour, we provide the colour from the ATLAS model, and the "3D-corrected" colour, that is, the ATLAS colour to which we added the corresponding 3D-1D corrections. The 3D-corrected colours can be used directly to interpret observed colours. The 3D corrections can also be used to correct colours computed from any 1D model atmosphere.

Table A.1. Colours and corrections for the Johnson-Cousins system for metallicity $[M/H]=0.0$. In columns labelled ‘ATLAS_c’ the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$U-B$ ATLAS	$U-B$ ATLAS _c	Δ_{U-B} 3D-1D	$B-V$ ATLAS	$B-V$ ATLAS _c	Δ_{B-V} 3D-1D	$V-R$ ATLAS	$V-R$ ATLAS _c	Δ_{V-R} 3D-1D	$V-I$ ATLAS	$V-I$ ATLAS _c	Δ_{V-I} 3D-1D
3635	1.00	+1.880	+1.792	-0.089	+1.671	+1.583	-0.014	+0.944	+0.956	+0.012	+2.062	+2.090	+0.028
3813	4.00	+1.123	+1.073	-0.050	+1.306	+1.256	+0.032	+0.878	+0.872	-0.006	+1.888	+1.861	-0.027
4018	1.50	+1.688	+1.580	-0.108	+1.541	+1.433	-0.034	+0.783	+0.764	-0.019	+1.495	+1.476	-0.019
3964	4.50	+1.095	+1.035	-0.060	+1.314	+1.255	-0.015	+0.844	+0.832	-0.012	+1.687	+1.689	+0.002
4501	2.00	+1.069	+1.011	-0.058	+1.268	+1.210	-0.027	+0.608	+0.601	-0.007	+1.149	+1.140	-0.009
4477	2.50	+1.004	+0.979	-0.025	+1.235	+1.210	-0.012	+0.618	+0.615	-0.003	+1.169	+1.167	-0.003
4478	4.00	+0.854	+0.833	-0.021	+1.136	+1.115	-0.008	+0.645	+0.640	-0.005	+1.203	+1.207	+0.005
4509	4.50	+0.879	+0.874	-0.005	+1.127	+1.122	-0.003	+0.658	+0.659	+0.001	+1.212	+1.231	+0.019
4582	3.20	+0.834	+0.813	-0.021	+1.143	+1.122	-0.008	+0.595	+0.593	-0.002	+1.123	+1.124	+0.001
4777	3.20	+0.679	+0.662	-0.017	+1.067	+1.049	-0.008	+0.540	+0.540	+0.000	+1.026	+1.029	+0.003
4968	2.50	+0.594	+0.561	-0.033	+1.019	+0.986	-0.020	+0.496	+0.494	-0.002	+0.945	+0.946	+0.001
5038	3.00	+0.503	+0.485	-0.019	+0.969	+0.950	-0.014	+0.481	+0.481	-0.001	+0.921	+0.922	+0.001
4923	3.50	+0.558	+0.543	-0.014	+0.998	+0.983	-0.009	+0.508	+0.507	-0.000	+0.967	+0.969	+0.002
4954	4.00	+0.529	+0.522	-0.007	+0.971	+0.965	-0.004	+0.507	+0.509	+0.002	+0.962	+0.966	+0.004
4982	4.50	+0.520	+0.523	+0.003	+0.951	+0.954	+0.001	+0.510	+0.516	+0.006	+0.961	+0.970	+0.010
5432	3.50	+0.247	+0.239	-0.008	+0.798	+0.790	-0.012	+0.411	+0.410	-0.001	+0.794	+0.795	+0.001
5475	4.00	+0.209	+0.212	+0.003	+0.781	+0.784	-0.005	+0.407	+0.408	+0.001	+0.784	+0.788	+0.004
5488	4.50	+0.202	+0.207	+0.006	+0.779	+0.784	+0.003	+0.409	+0.414	+0.005	+0.784	+0.794	+0.010
5774	4.40	+0.068	+0.077	+0.009	+0.680	+0.689	+0.003	+0.364	+0.368	+0.004	+0.703	+0.712	+0.009
5884	3.50	+0.082	+0.091	+0.009	+0.632	+0.641	-0.012	+0.342	+0.341	-0.001	+0.665	+0.667	+0.002
5928	4.00	+0.034	+0.043	+0.010	+0.624	+0.633	-0.004	+0.339	+0.340	+0.001	+0.659	+0.664	+0.005
5865	4.50	+0.031	+0.045	+0.014	+0.652	+0.666	+0.003	+0.352	+0.355	+0.004	+0.681	+0.688	+0.007
6228	4.00	-0.027	-0.015	+0.012	+0.531	+0.543	-0.003	+0.299	+0.300	+0.001	+0.584	+0.590	+0.006
6233	4.50	-0.069	-0.054	+0.014	+0.541	+0.555	-0.000	+0.303	+0.305	+0.002	+0.591	+0.595	+0.005
6484	4.00	-0.049	-0.036	+0.013	+0.461	+0.474	-0.003	+0.267	+0.267	+0.000	+0.522	+0.526	+0.004
6456	4.50	-0.101	-0.085	+0.016	+0.481	+0.497	+0.001	+0.276	+0.278	+0.002	+0.539	+0.544	+0.005
6726	4.30	-0.091	-0.073	+0.018	+0.411	+0.429	-0.004	+0.242	+0.239	-0.003	+0.473	+0.470	-0.003

Table A.2. Colours and corrections for the Johnson-Cousins system for metallicity $[M/H]=-1.0$. In columns labelled ‘ATLAS_c’ the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$U-B$ ATLAS	$U-B$ ATLAS _c	Δ_{U-B} 3D-1D	$B-V$ ATLAS	$B-V$ ATLAS _c	Δ_{B-V} 3D-1D	$V-R$ ATLAS	$V-R$ ATLAS _c	Δ_{V-R} 3D-1D	$V-I$ ATLAS	$V-I$ ATLAS _c	Δ_{V-I} 3D-1D
3790	4.00	+0.882	+0.843	-0.038	+1.326	+1.288	-0.012	+0.873	+0.859	-0.013	+1.744	+1.739	-0.005
4040	1.50	+1.068	+0.996	-0.073	+1.339	+1.266	-0.031	+0.736	+0.718	-0.018	+1.424	+1.403	-0.021
4001	4.50	+0.800	+0.768	-0.032	+1.268	+1.236	-0.020	+0.818	+0.806	-0.012	+1.555	+1.560	+0.006
4485	2.00	+0.553	+0.526	-0.027	+1.069	+1.042	-0.021	+0.575	+0.571	-0.004	+1.137	+1.132	-0.005
4490	2.50	+0.484	+0.479	-0.004	+1.036	+1.031	-0.005	+0.576	+0.577	+0.001	+1.136	+1.139	+0.003
4525	4.00	+0.417	+0.440	+0.023	+0.978	+1.000	+0.014	+0.595	+0.609	+0.013	+1.145	+1.167	+0.022
4499	4.50	+0.469	+0.471	+0.003	+0.994	+0.997	+0.003	+0.624	+0.628	+0.004	+1.180	+1.193	+0.013
4754	2.50	+0.309	+0.307	-0.002	+0.927	+0.925	-0.010	+0.507	+0.509	+0.002	+1.013	+1.016	+0.004
4993	2.50	+0.179	+0.191	+0.011	+0.830	+0.842	-0.008	+0.459	+0.462	+0.003	+0.921	+0.928	+0.007
4930	3.50	+0.156	+0.170	+0.014	+0.836	+0.850	+0.003	+0.475	+0.482	+0.008	+0.947	+0.958	+0.011
4986	4.00	+0.127	+0.154	+0.027	+0.814	+0.841	+0.013	+0.469	+0.483	+0.014	+0.931	+0.950	+0.019
5061	4.50	+0.102	+0.125	+0.022	+0.790	+0.812	+0.008	+0.460	+0.472	+0.012	+0.910	+0.923	+0.014
5481	3.50	-0.067	-0.035	+0.032	+0.649	+0.680	+0.006	+0.381	+0.389	+0.008	+0.770	+0.784	+0.014
5532	4.00	-0.103	-0.075	+0.027	+0.639	+0.666	+0.004	+0.376	+0.383	+0.007	+0.759	+0.769	+0.010
5473	4.50	-0.087	-0.059	+0.027	+0.665	+0.692	+0.006	+0.388	+0.397	+0.009	+0.779	+0.791	+0.012
5772	4.40	-0.176	-0.150	+0.026	+0.576	+0.602	+0.006	+0.346	+0.354	+0.008	+0.700	+0.712	+0.012
5890	3.50	-0.133	-0.092	+0.041	+0.523	+0.564	+0.007	+0.325	+0.332	+0.007	+0.661	+0.676	+0.015
5850	4.00	-0.171	-0.137	+0.034	+0.545	+0.579	+0.003	+0.334	+0.338	+0.004	+0.677	+0.684	+0.007
5923	4.50	-0.210	-0.171	+0.039	+0.537	+0.576	+0.013	+0.328	+0.338	+0.009	+0.665	+0.679	+0.014
6210	3.50	-0.135	-0.102	+0.033	+0.441	+0.474	-0.005	+0.284	+0.284	-0.000	+0.581	+0.583	+0.001
6261	4.00	-0.196	-0.158	+0.038	+0.447	+0.485	+0.003	+0.286	+0.287	+0.001	+0.585	+0.587	+0.001
6236	4.00	-0.196	-0.165	+0.030	+0.442	+0.472	+0.004	+0.284	+0.288	+0.004	+0.579	+0.588	+0.009
6238	4.50	-0.247	-0.209	+0.037	+0.460	+0.497	+0.011	+0.291	+0.298	+0.006	+0.594	+0.604	+0.010
6503	4.00	-0.184	-0.153	+0.031	+0.392	+0.423	+0.002	+0.255	+0.256	+0.001	+0.524	+0.526	+0.003
6456	4.50	-0.251	-0.211	+0.039	+0.414	+0.454	+0.013	+0.267	+0.274	+0.007	+0.546	+0.558	+0.012
6730	4.30	-0.208	-0.189	+0.019	+0.359	+0.378	+0.003	+0.234	+0.236	+0.001	+0.481	+0.485	+0.004

Table A.3. Colours and corrections for the Johnson-Cousins system for metallicity $[M/H]=-2.0$. In columns labelled ‘ATLAS_c’ the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$U-B$ ATLAS	$U-B$ ATLAS _c	Δ_{U-B} 3D-1D	$B-V$ ATLAS	$B-V$ ATLAS _c	Δ_{B-V} 3D-1D	$V-R$ ATLAS	$V-R$ ATLAS _c	Δ_{V-R} 3D-1D	$V-I$ ATLAS	$V-I$ ATLAS _c	Δ_{V-I} 3D-1D
4001	1.50	+0.895	+0.857	-0.038	+1.336	+1.298	-0.015	+0.733	+0.729	-0.004	+1.440	+1.433	-0.007
4000	4.50	+0.657	+0.654	-0.003	+1.255	+1.252	-0.001	+0.785	+0.784	-0.001	+1.490	+1.491	+0.000
4506	1.50	+0.440	+0.417	-0.023	+1.050	+1.027	-0.039	+0.568	+0.563	-0.004	+1.143	+1.139	-0.004
4506	2.00	+0.301	+0.300	-0.000	+0.989	+0.988	-0.011	+0.557	+0.564	+0.007	+1.126	+1.138	+0.013
4478	2.50	+0.344	+0.350	+0.006	+1.006	+1.012	+0.001	+0.555	+0.565	+0.010	+1.123	+1.136	+0.013
4504	4.00	+0.272	+0.283	+0.011	+0.960	+0.971	+0.015	+0.571	+0.582	+0.011	+1.133	+1.147	+0.015
4539	4.50	+0.280	+0.290	+0.010	+0.959	+0.969	+0.012	+0.576	+0.584	+0.008	+1.134	+1.146	+0.012
4743	2.50	+0.129	+0.132	+0.004	+0.871	+0.875	-0.014	+0.497	+0.501	+0.005	+1.013	+1.020	+0.006
4778	2.00	+0.164	+0.206	+0.042	+0.872	+0.914	-0.014	+0.495	+0.490	-0.005	+1.008	+0.998	-0.010
5024	2.50	-0.012	+0.017	+0.029	+0.745	+0.774	-0.012	+0.444	+0.448	+0.004	+0.912	+0.919	+0.007
4976	3.50	-0.030	-0.005	+0.026	+0.772	+0.798	+0.004	+0.454	+0.465	+0.012	+0.928	+0.943	+0.015
4955	4.00	-0.010	+0.022	+0.031	+0.789	+0.820	+0.010	+0.462	+0.477	+0.015	+0.940	+0.959	+0.019
5013	4.50	-0.019	+0.006	+0.025	+0.778	+0.803	+0.007	+0.458	+0.470	+0.012	+0.927	+0.941	+0.014
5502	2.50	-0.112	-0.076	+0.036	+0.563	+0.599	-0.014	+0.367	+0.366	-0.000	+0.760	+0.763	+0.002
5505	3.50	-0.209	-0.168	+0.041	+0.579	+0.620	+0.005	+0.371	+0.377	+0.006	+0.769	+0.777	+0.008
5472	4.00	-0.220	-0.177	+0.042	+0.604	+0.646	+0.016	+0.379	+0.392	+0.013	+0.782	+0.801	+0.019
5479	4.50	-0.221	-0.181	+0.040	+0.618	+0.658	+0.012	+0.383	+0.394	+0.012	+0.786	+0.800	+0.014
5772	3.70	-0.253	-0.201	+0.052	+0.509	+0.561	+0.015	+0.337	+0.345	+0.008	+0.701	+0.712	+0.012
5734	4.40	-0.285	-0.244	+0.041	+0.539	+0.580	+0.018	+0.347	+0.357	+0.010	+0.718	+0.732	+0.014
5861	3.50	-0.236	-0.189	+0.047	+0.482	+0.530	+0.007	+0.324	+0.329	+0.004	+0.675	+0.682	+0.007
5856	4.00	-0.282	-0.233	+0.049	+0.496	+0.544	+0.016	+0.329	+0.337	+0.008	+0.683	+0.695	+0.012
5923	4.50	-0.316	-0.269	+0.047	+0.494	+0.541	+0.021	+0.325	+0.335	+0.011	+0.674	+0.688	+0.015
6287	3.50	-0.206	-0.168	+0.038	+0.392	+0.430	+0.002	+0.271	+0.274	+0.002	+0.569	+0.574	+0.005
6206	4.00	-0.281	-0.246	+0.036	+0.421	+0.457	+0.008	+0.287	+0.290	+0.003	+0.600	+0.606	+0.006
6323	4.50	-0.331	-0.290	+0.042	+0.412	+0.453	+0.017	+0.279	+0.287	+0.007	+0.583	+0.594	+0.011
6534	4.00	-0.249	-0.215	+0.034	+0.362	+0.396	+0.006	+0.250	+0.252	+0.003	+0.525	+0.529	+0.005
6533	4.50	-0.319	-0.281	+0.038	+0.376	+0.415	+0.015	+0.257	+0.263	+0.006	+0.538	+0.547	+0.009

Table A.4. Colours and corrections for the Johnson-Cousins system for metallicity $[M/H]=-3.0$. In columns labelled ‘ATLAS_c’ the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$U-B$ ATLAS	$U-B$ ATLAS _c	Δ_{U-B} 3D-1D	$B-V$ ATLAS	$B-V$ ATLAS _c	Δ_{B-V} 3D-1D	$V-R$ ATLAS	$V-R$ ATLAS _c	Δ_{V-R} 3D-1D	$V-I$ ATLAS	$V-I$ ATLAS _c	Δ_{V-I} 3D-1D
3990	1.50	+0.962	+0.980	+0.018	+1.437	+1.456	-0.010	+0.769	+0.771	+0.002	+1.515	+1.518	+0.003
4503	1.50	+0.373	+0.338	-0.035	+1.039	+1.004	-0.050	+0.583	+0.576	-0.006	+1.177	+1.171	-0.007
4488	2.00	+0.302	+0.298	-0.005	+1.011	+1.006	-0.016	+0.571	+0.577	+0.006	+1.159	+1.170	+0.011
4523	2.50	+0.214	+0.207	-0.006	+0.965	+0.958	-0.012	+0.553	+0.560	+0.007	+1.126	+1.136	+0.011
4494	4.00	+0.193	+0.201	+0.008	+0.973	+0.981	+0.005	+0.572	+0.578	+0.005	+1.157	+1.164	+0.007
4522	4.50	+0.186	+0.193	+0.007	+0.967	+0.974	+0.003	+0.571	+0.575	+0.004	+1.152	+1.156	+0.004
4739	2.50	+0.049	+0.060	+0.010	+0.837	+0.848	-0.007	+0.500	+0.508	+0.008	+1.027	+1.042	+0.015
4891	1.80	+0.049	+0.067	+0.018	+0.769	+0.787	-0.010	+0.473	+0.478	+0.005	+0.975	+0.985	+0.010
4884	2.00	+0.023	+0.055	+0.033	+0.767	+0.800	-0.010	+0.473	+0.475	+0.003	+0.974	+0.979	+0.005
5018	2.50	-0.089	-0.049	+0.041	+0.698	+0.739	+0.005	+0.443	+0.453	+0.010	+0.919	+0.937	+0.018
4978	3.50	-0.113	-0.087	+0.026	+0.734	+0.760	+0.013	+0.455	+0.467	+0.013	+0.938	+0.959	+0.020
5160	4.00	-0.186	-0.157	+0.030	+0.676	+0.706	+0.018	+0.427	+0.439	+0.012	+0.883	+0.902	+0.019
4992	4.50	-0.102	-0.083	+0.019	+0.759	+0.778	+0.007	+0.466	+0.473	+0.007	+0.954	+0.964	+0.010
5501	2.50	-0.167	-0.120	+0.047	+0.536	+0.583	-0.007	+0.363	+0.366	+0.003	+0.762	+0.769	+0.007
5535	3.50	-0.272	-0.227	+0.045	+0.542	+0.588	+0.015	+0.365	+0.374	+0.009	+0.764	+0.778	+0.014
5476	4.00	-0.282	-0.241	+0.041	+0.572	+0.613	+0.028	+0.377	+0.393	+0.016	+0.786	+0.813	+0.027
5487	4.50	-0.285	-0.256	+0.029	+0.586	+0.616	+0.021	+0.381	+0.392	+0.011	+0.791	+0.809	+0.018
5789	3.70	-0.303	-0.261	+0.041	+0.486	+0.527	+0.013	+0.333	+0.340	+0.007	+0.699	+0.712	+0.013
5873	3.50	-0.281	-0.239	+0.042	+0.463	+0.506	+0.007	+0.321	+0.325	+0.004	+0.675	+0.683	+0.008
5846	4.00	-0.328	-0.285	+0.043	+0.481	+0.523	+0.017	+0.329	+0.337	+0.008	+0.689	+0.703	+0.013
5920	4.50	-0.361	-0.326	+0.035	+0.478	+0.513	+0.020	+0.324	+0.334	+0.009	+0.679	+0.694	+0.015
6306	3.50	-0.237	-0.204	+0.033	+0.379	+0.412	+0.002	+0.268	+0.270	+0.002	+0.568	+0.573	+0.005
6269	4.00	-0.312	-0.277	+0.035	+0.400	+0.435	+0.011	+0.279	+0.284	+0.004	+0.589	+0.597	+0.008
6242	4.00	-0.315	-0.282	+0.032	+0.404	+0.437	+0.008	+0.282	+0.285	+0.003	+0.595	+0.601	+0.006
6273	4.50	-0.368	-0.333	+0.035	+0.411	+0.446	+0.019	+0.284	+0.292	+0.008	+0.598	+0.611	+0.013
6408	4.00	-0.296	-0.273	+0.023	+0.376	+0.399	+0.009	+0.264	+0.267	+0.004	+0.557	+0.565	+0.008
6550	4.50	-0.348	-0.317	+0.030	+0.366	+0.397	+0.013	+0.255	+0.259	+0.004	+0.538	+0.545	+0.007

Table A.5. Colours and corrections for the 2MASS and Hipparcos-Tycho systems for metallicity $[M/H]=0.0$. In columns labelled ‘ATLAS_c’ the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$J - K$	$J - K$	Δ_{J-K}	$V - K$	$V - K$	Δ_{V-K}	$V - H_p$	$V - H_p$	Δ_{V-H_p}	$B_T - V_T$	$B_T - V_T$	$\Delta_{B_T-V_T}$
		ATLAS	ATLAS _c	3D-1D	ATLAS	ATLAS _c	3D-1D	ATLAS	ATLAS _c	3D-1D	ATLAS	ATLAS _c	3D-1D
3635	1.00	+1.146	+1.134	-0.012	+4.567	+4.637	+0.070	-0.179	-0.172	+0.007	+1.857	+1.840	-0.017
3813	4.00	+0.958	+0.960	+0.002	+4.026	+3.964	-0.061	-0.150	-0.159	-0.009	+1.346	+1.385	+0.039
4018	1.50	+0.949	+0.933	-0.017	+3.513	+3.463	-0.050	-0.217	-0.218	-0.001	+1.782	+1.739	-0.043
3964	4.50	+0.889	+0.870	-0.020	+3.627	+3.594	-0.033	-0.162	-0.164	-0.002	+1.382	+1.364	-0.018
4501	2.00	+0.725	+0.714	-0.011	+2.717	+2.681	-0.036	-0.225	-0.222	+0.003	+1.469	+1.430	-0.039
4477	2.50	+0.742	+0.733	-0.009	+2.762	+2.741	-0.021	-0.215	-0.214	+0.001	+1.419	+1.402	-0.017
4478	4.00	+0.756	+0.727	-0.028	+2.802	+2.762	-0.040	-0.184	-0.185	-0.001	+1.273	+1.261	-0.013
4509	4.50	+0.741	+0.723	-0.018	+2.781	+2.777	-0.004	-0.178	-0.177	+0.001	+1.257	+1.249	-0.008
4582	3.20	+0.707	+0.697	-0.010	+2.636	+2.619	-0.018	-0.202	-0.201	+0.001	+1.301	+1.288	-0.013
4777	3.20	+0.634	+0.628	-0.007	+2.396	+2.385	-0.011	-0.201	-0.200	+0.001	+1.213	+1.200	-0.013
4968	2.50	+0.564	+0.556	-0.008	+2.182	+2.162	-0.020	-0.203	-0.200	+0.003	+1.154	+1.123	-0.031
5038	3.00	+0.547	+0.541	-0.007	+2.120	+2.104	-0.015	-0.195	-0.193	+0.003	+1.091	+1.070	-0.022
4923	3.50	+0.587	+0.577	-0.010	+2.242	+2.223	-0.019	-0.194	-0.192	+0.002	+1.129	+1.114	-0.015
4954	4.00	+0.581	+0.572	-0.009	+2.219	+2.205	-0.014	-0.187	-0.186	+0.001	+1.098	+1.089	-0.009
4982	4.50	+0.574	+0.568	-0.006	+2.201	+2.199	-0.002	-0.180	-0.179	+0.001	+1.074	+1.072	-0.003
5432	3.50	+0.445	+0.439	-0.007	+1.781	+1.765	-0.016	-0.169	-0.167	+0.002	+0.879	+0.861	-0.018
5475	4.00	+0.438	+0.433	-0.005	+1.753	+1.745	-0.008	-0.165	-0.164	+0.001	+0.861	+0.852	-0.009
5488	4.50	+0.436	+0.434	-0.003	+1.748	+1.750	+0.002	-0.163	-0.163	+0.000	+0.861	+0.861	+0.000
5774	4.40	+0.374	+0.373	-0.001	+1.540	+1.543	+0.003	-0.147	-0.147	-0.001	+0.736	+0.738	+0.002
5884	3.50	+0.348	+0.344	-0.003	+1.453	+1.441	-0.012	-0.139	-0.136	+0.002	+0.674	+0.658	-0.017
5928	4.00	+0.342	+0.340	-0.002	+1.433	+1.428	-0.005	-0.136	-0.136	+0.001	+0.665	+0.658	-0.007
5865	4.50	+0.356	+0.354	-0.003	+1.480	+1.478	-0.002	-0.141	-0.142	-0.001	+0.702	+0.703	+0.001
6228	4.00	+0.289	+0.290	+0.001	+1.247	+1.247	-0.000	-0.117	-0.117	+0.000	+0.554	+0.549	-0.005
6233	4.50	+0.291	+0.290	-0.002	+1.255	+1.251	-0.004	-0.119	-0.119	-0.000	+0.566	+0.565	-0.001
6484	4.00	+0.249	+0.253	+0.004	+1.101	+1.103	+0.003	-0.102	-0.101	+0.000	+0.473	+0.469	-0.004
6456	4.50	+0.257	+0.257	+0.001	+1.130	+1.130	+0.000	-0.106	-0.106	-0.001	+0.496	+0.496	+0.000
6726	4.30	+0.216	+0.222	+0.005	+0.981	+0.982	+0.001	-0.090	-0.089	+0.001	+0.417	+0.414	-0.003

Table A.6. Colours and corrections for the 2MASS and Hipparcos-Tycho systems for metallicity $[M/H]=-1.0$. In columns labelled ‘ATLAS_c’ the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$J - K$	$J - K$	Δ_{J-K}	$V - K$	$V - K$	Δ_{V-K}	$V - H_p$	$V - H_p$	Δ_{V-H_p}	$B_T - V_T$	$B_T - V_T$	$\Delta_{B_T-V_T}$
		ATLAS	ATLAS _c	3D-1D	ATLAS	ATLAS _c	3D-1D	ATLAS	ATLAS _c	3D-1D	ATLAS	ATLAS _c	3D-1D
3790	4.00	+0.859	+0.842	-0.017	+3.683	+3.649	-0.034	-0.157	-0.160	-0.003	+1.360	+1.350	-0.011
4040	1.50	+0.968	+0.940	-0.028	+3.430	+3.358	-0.072	-0.199	-0.200	-0.001	+1.498	+1.460	-0.039
4001	4.50	+0.820	+0.802	-0.018	+3.345	+3.329	-0.017	-0.163	-0.163	-0.001	+1.322	+1.298	-0.023
4485	2.00	+0.756	+0.742	-0.014	+2.734	+2.700	-0.034	-0.195	-0.192	+0.003	+1.182	+1.153	-0.029
4490	2.50	+0.758	+0.741	-0.017	+2.733	+2.707	-0.026	-0.186	-0.185	+0.001	+1.138	+1.128	-0.010
4525	4.00	+0.743	+0.728	-0.015	+2.710	+2.712	+0.002	-0.165	-0.165	+0.000	+1.057	+1.068	+0.011
4499	4.50	+0.745	+0.717	-0.028	+2.750	+2.720	-0.030	-0.160	-0.160	+0.000	+1.068	+1.069	+0.000
4754	2.50	+0.656	+0.642	-0.014	+2.414	+2.390	-0.024	-0.180	-0.178	+0.002	+1.009	+0.993	-0.016
4993	2.50	+0.578	+0.565	-0.013	+2.168	+2.148	-0.020	-0.169	-0.167	+0.002	+0.890	+0.877	-0.013
4930	3.50	+0.602	+0.593	-0.009	+2.238	+2.231	-0.007	-0.165	-0.164	+0.001	+0.899	+0.900	+0.001
4986	4.00	+0.585	+0.580	-0.005	+2.186	+2.196	+0.010	-0.160	-0.160	-0.000	+0.874	+0.886	+0.012
5061	4.50	+0.561	+0.556	-0.005	+2.117	+2.120	+0.003	-0.154	-0.153	+0.001	+0.847	+0.853	+0.007
5481	3.50	+0.454	+0.446	-0.008	+1.767	+1.763	-0.003	-0.137	-0.138	-0.001	+0.671	+0.676	+0.005
5532	4.00	+0.444	+0.438	-0.007	+1.733	+1.728	-0.005	-0.134	-0.135	-0.000	+0.661	+0.665	+0.004
5473	4.50	+0.457	+0.453	-0.004	+1.777	+1.778	+0.001	-0.138	-0.138	+0.000	+0.695	+0.701	+0.006
5772	4.40	+0.395	+0.385	-0.010	+1.572	+1.562	-0.011	-0.122	-0.123	-0.001	+0.586	+0.591	+0.005
5890	3.50	+0.369	+0.366	-0.003	+1.486	+1.489	+0.004	-0.112	-0.113	-0.002	+0.524	+0.532	+0.008
5850	4.00	+0.379	+0.371	-0.008	+1.520	+1.507	-0.013	-0.116	-0.116	-0.001	+0.549	+0.552	+0.003
5923	4.50	+0.367	+0.365	-0.002	+1.479	+1.486	+0.007	-0.114	-0.116	-0.003	+0.541	+0.557	+0.016
6210	3.50	+0.310	+0.311	+0.001	+1.288	+1.282	-0.006	-0.094	-0.093	+0.001	+0.434	+0.430	-0.004
6261	4.00	+0.311	+0.310	-0.001	+1.288	+1.281	-0.007	-0.095	-0.096	-0.001	+0.440	+0.446	+0.006
6236	4.00	+0.306	+0.304	-0.003	+1.274	+1.271	-0.003	-0.094	-0.095	-0.001	+0.433	+0.438	+0.005
6238	4.50	+0.313	+0.310	-0.003	+1.298	+1.297	-0.001	-0.097	-0.100	-0.003	+0.453	+0.467	+0.014
6503	4.00	+0.268	+0.268	+0.000	+1.139	+1.135	-0.004	-0.083	-0.083	-0.001	+0.381	+0.384	+0.004
6456	4.50	+0.279	+0.279	-0.000	+1.181	+1.186	+0.005	-0.087	-0.091	-0.004	+0.403	+0.420	+0.017
6730	4.30	+0.237	+0.242	+0.005	+1.030	+1.036	+0.005	-0.075	-0.076	-0.001	+0.346	+0.351	+0.004

Table A.7. Colours and corrections for the 2MASS and Hipparcos-Tycho systems for metallicity $[M/H]=-2.0$. In columns labelled ‘ATLAS_c’ the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$J-K$	$J-K$	Δ_{J-K}	$V-K$	$V-K$	Δ_{V-K}	$V-H_p$	$V-H_p$	Δ_{V-H_p}	B_T-V_T	B_T-V_T	$\Delta_{B_T-V_T}$
		ATLAS	ATLAS _c	3D-1D	ATLAS	ATLAS _c	3D-1D	ATLAS	ATLAS _c	3D-1D	ATLAS	ATLAS _c	3D-1D
4001	1.50	+0.985	+0.938	-0.047	+3.479	+3.395	-0.084	-0.202	-0.201	+0.001	+1.481	+1.460	-0.022
4000	4.50	+0.785	+0.773	-0.011	+3.216	+3.201	-0.015	-0.170	-0.170	-0.000	+1.330	+1.329	-0.001
4506	1.50	+0.740	+0.737	-0.003	+2.711	+2.690	-0.021	-0.197	-0.190	+0.007	+1.136	+1.086	-0.050
4506	2.00	+0.758	+0.747	-0.012	+2.723	+2.711	-0.012	-0.184	-0.180	+0.004	+1.063	+1.045	-0.017
4478	2.50	+0.745	+0.725	-0.020	+2.697	+2.676	-0.021	-0.190	-0.187	+0.003	+1.083	+1.080	-0.003
4504	4.00	+0.719	+0.693	-0.026	+2.661	+2.635	-0.025	-0.171	-0.172	-0.001	+1.025	+1.040	+0.015
4539	4.50	+0.686	+0.666	-0.020	+2.602	+2.584	-0.017	-0.170	-0.170	-0.000	+1.021	+1.034	+0.013
4743	2.50	+0.661	+0.645	-0.015	+2.420	+2.395	-0.025	-0.172	-0.168	+0.004	+0.918	+0.899	-0.019
4778	2.00	+0.646	+0.650	+0.004	+2.389	+2.385	-0.004	-0.174	-0.171	+0.003	+0.917	+0.906	-0.011
5024	2.50	+0.576	+0.563	-0.013	+2.150	+2.129	-0.021	-0.153	-0.149	+0.004	+0.764	+0.749	-0.014
4976	3.50	+0.585	+0.580	-0.006	+2.183	+2.187	+0.003	-0.157	-0.155	+0.002	+0.800	+0.804	+0.003
4955	4.00	+0.584	+0.578	-0.005	+2.193	+2.203	+0.010	-0.158	-0.157	+0.002	+0.822	+0.832	+0.011
5013	4.50	+0.559	+0.551	-0.008	+2.132	+2.132	-0.000	-0.156	-0.154	+0.002	+0.810	+0.816	+0.007
5502	2.50	+0.457	+0.454	-0.003	+1.763	+1.751	-0.012	-0.119	-0.115	+0.004	+0.553	+0.539	-0.013
5505	3.50	+0.462	+0.452	-0.010	+1.776	+1.762	-0.015	-0.121	-0.122	-0.000	+0.570	+0.577	+0.008
5472	4.00	+0.466	+0.461	-0.005	+1.796	+1.804	+0.007	-0.126	-0.128	-0.002	+0.599	+0.618	+0.019
5479	4.50	+0.459	+0.455	-0.004	+1.784	+1.789	+0.005	-0.129	-0.130	-0.000	+0.616	+0.632	+0.016
5772	3.70	+0.409	+0.400	-0.009	+1.601	+1.592	-0.009	-0.107	-0.110	-0.003	+0.491	+0.511	+0.020
5734	4.40	+0.414	+0.409	-0.005	+1.626	+1.628	+0.002	-0.113	-0.116	-0.003	+0.524	+0.548	+0.024
5861	3.50	+0.390	+0.384	-0.006	+1.540	+1.529	-0.011	-0.101	-0.103	-0.002	+0.463	+0.474	+0.011
5856	4.00	+0.394	+0.387	-0.007	+1.552	+1.546	-0.006	-0.104	-0.107	-0.003	+0.476	+0.498	+0.022
5923	4.50	+0.381	+0.374	-0.007	+1.513	+1.512	-0.002	-0.104	-0.108	-0.004	+0.474	+0.501	+0.027
6287	3.50	+0.312	+0.313	+0.001	+1.277	+1.276	-0.001	-0.082	-0.083	-0.000	+0.371	+0.376	+0.005
6206	4.00	+0.331	+0.331	-0.000	+1.343	+1.340	-0.003	-0.088	-0.090	-0.002	+0.398	+0.410	+0.012
6323	4.50	+0.315	+0.310	-0.005	+1.290	+1.285	-0.006	-0.086	-0.090	-0.004	+0.387	+0.409	+0.022
6534	4.00	+0.277	+0.277	+0.000	+1.157	+1.153	-0.004	-0.075	-0.077	-0.002	+0.340	+0.350	+0.010
6533	4.50	+0.282	+0.279	-0.003	+1.178	+1.173	-0.004	-0.078	-0.082	-0.004	+0.352	+0.371	+0.019

Table A.8. Colours and corrections for the 2MASS and Hipparcos-Tycho systems for metallicity $[M/H]=-3.0$. In columns labelled ‘ATLAS_c’ the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$J-K$	$J-K$	Δ_{J-K}	$V-K$	$V-K$	Δ_{V-K}	$V-H_p$	$V-H_p$	Δ_{V-H_p}	B_T-V_T	B_T-V_T	$\Delta_{B_T-V_T}$
		ATLAS	ATLAS _c	3D-1D	ATLAS	ATLAS _c	3D-1D	ATLAS	ATLAS _c	3D-1D	ATLAS	ATLAS _c	3D-1D
3990	1.50	+0.940	+0.922	-0.018	+3.506	+3.469	-0.038	-0.212	-0.210	+0.002	+1.588	+1.576	-0.012
4503	1.50	+0.725	+0.727	+0.002	+2.728	+2.709	-0.019	-0.195	-0.186	+0.008	+1.096	+1.033	-0.063
4488	2.00	+0.729	+0.723	-0.006	+2.714	+2.708	-0.006	-0.190	-0.185	+0.005	+1.066	+1.043	-0.022
4523	2.50	+0.711	+0.695	-0.015	+2.644	+2.626	-0.019	-0.184	-0.179	+0.005	+1.012	+0.994	-0.018
4494	4.00	+0.660	+0.654	-0.006	+2.589	+2.585	-0.004	-0.180	-0.180	+0.001	+1.015	+1.020	+0.005
4522	4.50	+0.634	+0.630	-0.004	+2.538	+2.535	-0.003	-0.179	-0.179	+0.000	+1.008	+1.012	+0.004
4739	2.50	+0.651	+0.644	-0.007	+2.421	+2.418	-0.003	-0.167	-0.163	+0.004	+0.856	+0.846	-0.010
4891	1.80	+0.611	+0.604	-0.007	+2.287	+2.275	-0.012	-0.157	-0.153	+0.004	+0.775	+0.763	-0.012
4884	2.00	+0.614	+0.597	-0.017	+2.292	+2.262	-0.030	-0.156	-0.153	+0.003	+0.773	+0.761	-0.012
5018	2.50	+0.581	+0.572	-0.009	+2.167	+2.164	-0.002	-0.143	-0.143	+0.001	+0.693	+0.699	+0.006
4978	3.50	+0.575	+0.570	-0.006	+2.178	+2.187	+0.009	-0.150	-0.150	-0.000	+0.734	+0.749	+0.015
5160	4.00	+0.527	+0.520	-0.007	+2.025	+2.030	+0.005	-0.140	-0.142	-0.002	+0.667	+0.688	+0.022
4992	4.50	+0.543	+0.540	-0.003	+2.134	+2.137	+0.003	-0.154	-0.154	-0.000	+0.762	+0.770	+0.008
5501	2.50	+0.464	+0.461	-0.003	+1.779	+1.773	-0.007	-0.113	-0.111	+0.002	+0.515	+0.509	-0.005
5535	3.50	+0.463	+0.453	-0.011	+1.773	+1.765	-0.009	-0.113	-0.116	-0.003	+0.518	+0.536	+0.019
5476	4.00	+0.471	+0.463	-0.007	+1.809	+1.822	+0.012	-0.120	-0.125	-0.005	+0.549	+0.582	+0.033
5487	4.50	+0.460	+0.451	-0.008	+1.792	+1.793	+0.001	-0.123	-0.127	-0.004	+0.565	+0.590	+0.025
5789	3.70	+0.413	+0.407	-0.007	+1.608	+1.603	-0.005	-0.102	-0.105	-0.003	+0.458	+0.474	+0.017
5873	3.50	+0.395	+0.389	-0.006	+1.550	+1.540	-0.010	-0.097	-0.099	-0.002	+0.436	+0.446	+0.010
5846	4.00	+0.403	+0.393	-0.010	+1.576	+1.566	-0.009	-0.101	-0.105	-0.004	+0.452	+0.473	+0.021
5920	4.50	+0.389	+0.376	-0.012	+1.533	+1.521	-0.012	-0.100	-0.105	-0.005	+0.448	+0.472	+0.024
6306	3.50	+0.314	+0.317	+0.002	+1.281	+1.282	+0.001	-0.079	-0.080	-0.001	+0.353	+0.357	+0.004
6269	4.00	+0.327	+0.324	-0.003	+1.324	+1.318	-0.006	-0.083	-0.086	-0.003	+0.371	+0.384	+0.014
6242	4.00	+0.332	+0.328	-0.004	+1.339	+1.331	-0.009	-0.084	-0.087	-0.002	+0.375	+0.386	+0.011
6273	4.50	+0.330	+0.323	-0.007	+1.335	+1.328	-0.006	-0.086	-0.091	-0.005	+0.380	+0.402	+0.022
6408	4.00	+0.303	+0.306	+0.003	+1.244	+1.248	+0.003	-0.078	-0.081	-0.002	+0.348	+0.359	+0.011

Table A.9. Colours and corrections for the Gaia system for metallicity $[M/H]=0.0$. In columns $ATLAS_c$ the 3D correction has been added to the $ATLAS$ colour. The CaT colour is defined as $(G - RVS) - (BP - RP)$. Bol_G is the bolometric correction in the G-band.

T_{eff}	$\log g$	$G - H_p$ ATLAS	$G - H_p$ ATLAS _c	Δ_{G-H_p} 3D-1D	$BP - RP$ ATLAS	$BP - RP$ ATLAS _c	Δ_{BP-RP} 3D-1D	CaT ATLAS	CaT ATLAS _c	Δ_{CaT} 3D-1D	Bol_G ATLAS	Bol_G ATLAS _c	Δ_{Bol_G} 3D-1D
3635	1.00	-0.767	-0.780	-0.013	+2.247	+2.277	+0.030	-0.494	-0.511	-0.017	-1.153	-1.272	-0.119
3813	4.00	-0.661	-0.655	+0.007	+2.022	+2.006	-0.016	-0.383	-0.377	+0.006	-0.887	-0.857	+0.029
4018	1.50	-0.541	-0.531	+0.010	+1.752	+1.730	-0.021	-0.372	-0.359	+0.013	-0.650	-0.628	+0.022
3964	4.50	-0.587	-0.586	+0.001	+1.852	+1.850	-0.001	-0.359	-0.349	+0.010	-0.695	-0.687	+0.008
4501	2.00	-0.399	-0.394	+0.006	+1.415	+1.401	-0.014	-0.275	-0.269	+0.006	-0.338	-0.322	+0.016
4477	2.50	-0.402	-0.400	+0.002	+1.422	+1.417	-0.005	-0.270	-0.267	+0.003	-0.353	-0.346	+0.007
4478	4.00	-0.402	-0.402	+0.000	+1.419	+1.419	+0.000	-0.265	-0.259	+0.006	-0.361	-0.332	+0.029
4509	4.50	-0.404	-0.409	-0.005	+1.422	+1.435	+0.013	-0.274	-0.270	+0.004	-0.350	-0.358	-0.008
4582	3.20	-0.377	-0.377	+0.001	+1.359	+1.357	-0.001	-0.249	-0.246	+0.003	-0.307	-0.304	+0.004
4777	3.20	-0.339	-0.339	-0.000	+1.259	+1.259	+0.001	-0.222	-0.221	+0.001	-0.230	-0.229	+0.002
4968	2.50	-0.310	-0.308	+0.002	+1.180	+1.175	-0.005	-0.210	-0.206	+0.004	-0.167	-0.161	+0.006
5038	3.00	-0.297	-0.296	+0.001	+1.144	+1.142	-0.002	-0.196	-0.193	+0.003	-0.150	-0.148	+0.003
4923	3.50	-0.313	-0.313	+0.000	+1.190	+1.189	-0.001	-0.204	-0.202	+0.002	-0.184	-0.182	+0.002
4954	4.00	-0.309	-0.309	-0.001	+1.176	+1.178	+0.002	-0.200	-0.201	-0.000	-0.177	-0.176	+0.001
4982	4.50	-0.306	-0.309	-0.003	+1.168	+1.176	+0.008	-0.201	-0.204	-0.003	-0.170	-0.174	-0.004
5432	3.50	-0.241	-0.240	+0.001	+0.984	+0.981	-0.002	-0.155	-0.151	+0.004	-0.071	-0.073	-0.001
5475	4.00	-0.236	-0.237	-0.001	+0.969	+0.971	+0.002	-0.150	-0.148	+0.002	-0.067	-0.069	-0.002
5488	4.50	-0.236	-0.239	-0.003	+0.967	+0.976	+0.010	-0.150	-0.152	-0.001	-0.065	-0.074	-0.008
5774	4.40	-0.203	-0.206	-0.003	+0.865	+0.874	+0.009	-0.125	-0.126	-0.000	-0.032	-0.039	-0.008
5884	3.50	-0.187	-0.187	+0.001	+0.819	+0.817	-0.002	-0.116	-0.111	+0.005	-0.014	-0.028	-0.014
5928	4.00	-0.185	-0.186	-0.001	+0.809	+0.812	+0.003	-0.112	-0.110	+0.002	-0.016	-0.022	-0.006
5865	4.50	-0.194	-0.196	-0.003	+0.836	+0.843	+0.008	-0.119	-0.119	-0.000	-0.024	-0.031	-0.007
6228	4.00	-0.155	-0.156	-0.001	+0.712	+0.716	+0.004	-0.090	-0.087	+0.004	+0.004	-0.014	-0.018
6233	4.50	-0.158	-0.159	-0.001	+0.719	+0.723	+0.005	-0.091	-0.090	+0.001	-0.002	-0.009	-0.007
6484	4.00	-0.132	-0.133	-0.001	+0.633	+0.636	+0.003	-0.074	-0.071	+0.004	+0.015	-0.002	-0.017
6456	4.50	-0.138	-0.140	-0.002	+0.652	+0.658	+0.005	-0.077	-0.075	+0.002	+0.005	-0.003	-0.008
6726	4.30	-0.115	-0.113	+0.001	+0.571	+0.567	-0.004	-0.062	-0.059	+0.002	+0.014	+0.010	-0.004

Table A.10. Colours and corrections for the Gaia system for metallicity $[M/H]=-1.0$. In columns $ATLAS_c$ the 3D correction has been added to the $ATLAS$ colour. The CaT colour is defined as $(G - RVS) - (BP - RP)$. Bol_G is the bolometric correction in the G-band.

T_{eff}	$\log g$	$G - H_p$ ATLAS	$G - H_p$ $ATLAS_c$	Δ_{G-H_p} 3D-1D	$BP - RP$ ATLAS	$BP - RP$ $ATLAS_c$	Δ_{BP-RP} 3D-1D	CaT ATLAS	CaT $ATLAS_c$	Δ_{CaT} 3D-1D	Bol_G ATLAS	Bol_G $ATLAS_c$	Δ_{Bol_G} 3D-1D
3790	4.00	-0.612	-0.608	+0.004	+1.905	+1.899	-0.006	-0.368	-0.358	+0.010	-0.717	-0.705	+0.013
4040	1.50	-0.499	-0.488	+0.011	+1.659	+1.635	-0.024	-0.327	-0.314	+0.013	-0.618	-0.570	+0.048
4001	4.50	-0.539	-0.537	+0.001	+1.738	+1.737	-0.001	-0.343	-0.331	+0.012	-0.568	-0.566	+0.001
4485	2.00	-0.374	-0.370	+0.004	+1.356	+1.347	-0.009	-0.230	-0.224	+0.006	-0.353	-0.342	+0.011
4490	2.50	-0.370	-0.370	-0.001	+1.345	+1.346	+0.001	-0.221	-0.221	+0.000	-0.352	-0.346	+0.007
4525	4.00	-0.366	-0.375	-0.009	+1.330	+1.350	+0.021	-0.219	-0.226	-0.007	-0.338	-0.344	-0.007
4499	4.50	-0.379	-0.383	-0.004	+1.361	+1.371	+0.010	-0.234	-0.235	-0.001	-0.347	-0.328	+0.019
4754	2.50	-0.319	-0.319	-0.000	+1.211	+1.211	+0.000	-0.185	-0.183	+0.002	-0.250	-0.247	+0.004
4993	2.50	-0.281	-0.282	-0.001	+1.103	+1.107	+0.004	-0.158	-0.157	+0.001	-0.182	-0.182	-0.001
4930	3.50	-0.289	-0.293	-0.004	+1.125	+1.135	+0.010	-0.159	-0.162	-0.004	-0.202	-0.205	-0.003
4986	4.00	-0.281	-0.289	-0.008	+1.102	+1.122	+0.020	-0.154	-0.162	-0.008	-0.187	-0.202	-0.015
5061	4.50	-0.272	-0.277	-0.005	+1.075	+1.088	+0.013	-0.150	-0.157	-0.007	-0.168	-0.173	-0.005
5481	3.50	-0.216	-0.222	-0.005	+0.911	+0.926	+0.015	-0.106	-0.109	-0.003	-0.101	-0.116	-0.015
5532	4.00	-0.212	-0.216	-0.004	+0.897	+0.908	+0.011	-0.103	-0.106	-0.003	-0.098	-0.104	-0.007
5473	4.50	-0.221	-0.225	-0.004	+0.922	+0.934	+0.012	-0.111	-0.116	-0.005	-0.103	-0.111	-0.008
5772	4.40	-0.189	-0.194	-0.004	+0.822	+0.836	+0.013	-0.086	-0.089	-0.003	-0.079	-0.078	+0.001
5890	3.50	-0.173	-0.178	-0.005	+0.773	+0.790	+0.017	-0.073	-0.075	-0.002	-0.063	-0.091	-0.027
5850	4.00	-0.180	-0.182	-0.003	+0.793	+0.801	+0.008	-0.077	-0.079	-0.002	-0.073	-0.073	-0.000
5923	4.50	-0.175	-0.181	-0.006	+0.777	+0.795	+0.018	-0.075	-0.081	-0.006	-0.072	-0.093	-0.021
6210	3.50	-0.143	-0.143	+0.000	+0.675	+0.675	+0.000	-0.053	-0.051	+0.002	-0.045	-0.052	-0.007
6261	4.00	-0.145	-0.146	-0.001	+0.678	+0.681	+0.003	-0.053	-0.054	-0.002	-0.057	-0.075	-0.018
6236	4.00	-0.143	-0.146	-0.003	+0.671	+0.681	+0.010	-0.051	-0.052	-0.001	-0.056	-0.051	+0.005
6238	4.50	-0.148	-0.152	-0.004	+0.687	+0.701	+0.014	-0.054	-0.059	-0.004	-0.065	-0.074	-0.009
6503	4.00	-0.124	-0.124	-0.001	+0.604	+0.608	+0.004	-0.040	-0.041	-0.001	-0.049	-0.057	-0.008
6456	4.50	-0.131	-0.136	-0.005	+0.629	+0.645	+0.016	-0.043	-0.047	-0.004	-0.063	-0.081	-0.018
6730	4.30	-0.110	-0.111	-0.001	+0.554	+0.558	+0.005	-0.032	-0.032	-0.000	-0.055	-0.069	-0.014

Table A.11. Colours and corrections for the Gaia system for metallicity $[M/H]=-2.0$. In columns $ATLAS_c$ the 3D correction has been added to the $ATLAS$ colour. The CaT colour is defined as $(G - RVS) - (BP - RP)$. Bol_G is the bolometric correction in the G-band.

T_{eff}	$\log g$	$G - H_p$ ATLAS	$G - H_p$ ATLAS _c	Δ_{G-H_p} 3D-1D	$BP - RP$ ATLAS	$BP - RP$ ATLAS _c	Δ_{BP-RP} 3D-1D	CaT ATLAS	CaT ATLAS _c	Δ_{CaT} 3D-1D	Bol_G ATLAS	Bol_G ATLAS _c	Δ_{Bol_G} 3D-1D
4001	1.50	-0.505	-0.501	+0.004	+1.677	+1.668	-0.009	-0.317	-0.314	+0.003	-0.646	-0.600	+0.046
4000	4.50	-0.515	-0.515	+0.000	+1.689	+1.689	+0.000	-0.327	-0.326	+0.001	-0.522	-0.518	+0.003
4506	1.50	-0.375	-0.370	+0.006	+1.361	+1.347	-0.014	-0.216	-0.207	+0.009	-0.352	-0.348	+0.005
4506	2.00	-0.362	-0.365	-0.003	+1.326	+1.333	+0.007	-0.196	-0.197	-0.001	-0.358	-0.362	-0.004
4478	2.50	-0.363	-0.368	-0.004	+1.330	+1.340	+0.010	-0.201	-0.206	-0.005	-0.349	-0.346	+0.002
4504	4.00	-0.360	-0.367	-0.007	+1.318	+1.334	+0.016	-0.198	-0.206	-0.008	-0.332	-0.329	+0.003
4539	4.50	-0.361	-0.366	-0.005	+1.319	+1.331	+0.013	-0.202	-0.208	-0.006	-0.310	-0.308	+0.003
4743	2.50	-0.313	-0.314	-0.001	+1.196	+1.197	+0.001	-0.159	-0.157	+0.001	-0.263	-0.258	+0.005
4778	2.00	-0.312	-0.307	+0.005	+1.193	+1.180	-0.012	-0.160	-0.158	+0.003	-0.252	-0.258	-0.007
5024	2.50	-0.268	-0.269	-0.001	+1.068	+1.071	+0.003	-0.122	-0.121	+0.001	-0.193	-0.195	-0.002
4976	3.50	-0.276	-0.281	-0.005	+1.088	+1.102	+0.014	-0.129	-0.135	-0.006	-0.203	-0.214	-0.011
4955	4.00	-0.281	-0.288	-0.007	+1.103	+1.122	+0.018	-0.136	-0.144	-0.009	-0.203	-0.218	-0.014
5013	4.50	-0.277	-0.281	-0.005	+1.089	+1.101	+0.013	-0.134	-0.142	-0.007	-0.186	-0.193	-0.006
5502	2.50	-0.204	-0.203	+0.001	+0.876	+0.874	-0.002	-0.071	-0.069	+0.003	-0.118	-0.125	-0.007
5505	3.50	-0.208	-0.211	-0.003	+0.884	+0.893	+0.009	-0.072	-0.077	-0.004	-0.134	-0.140	-0.005
5472	4.00	-0.215	-0.222	-0.007	+0.903	+0.925	+0.022	-0.079	-0.088	-0.008	-0.137	-0.163	-0.026
5479	4.50	-0.218	-0.223	-0.005	+0.911	+0.927	+0.016	-0.084	-0.093	-0.009	-0.132	-0.144	-0.012
5772	3.70	-0.182	-0.187	-0.005	+0.799	+0.816	+0.017	-0.053	-0.061	-0.008	-0.118	-0.122	-0.004
5734	4.40	-0.190	-0.196	-0.006	+0.823	+0.842	+0.020	-0.060	-0.070	-0.009	-0.122	-0.139	-0.017
5861	3.50	-0.172	-0.175	-0.003	+0.768	+0.778	+0.010	-0.047	-0.052	-0.005	-0.110	-0.116	-0.006
5856	4.00	-0.176	-0.181	-0.005	+0.778	+0.795	+0.017	-0.049	-0.057	-0.008	-0.118	-0.128	-0.010
5923	4.50	-0.173	-0.179	-0.006	+0.767	+0.788	+0.021	-0.048	-0.059	-0.011	-0.117	-0.131	-0.014
6287	3.50	-0.135	-0.136	-0.002	+0.645	+0.651	+0.006	-0.028	-0.030	-0.002	-0.085	-0.096	-0.011
6206	4.00	-0.146	-0.148	-0.002	+0.679	+0.688	+0.009	-0.032	-0.035	-0.004	-0.105	-0.117	-0.012
6323	4.50	-0.140	-0.145	-0.005	+0.658	+0.675	+0.017	-0.028	-0.036	-0.008	-0.114	-0.123	-0.009
6534	4.00	-0.120	-0.122	-0.002	+0.592	+0.600	+0.007	-0.021	-0.024	-0.003	-0.094	-0.105	-0.011
6533	4.50	-0.125	-0.129	-0.004	+0.605	+0.620	+0.015	-0.020	-0.027	-0.007	-0.111	-0.122	-0.011

Table A.12. Colours and corrections for the Gaia system for metallicity $[M/H]=-3.0$. In columns $ATLAS_c$ the 3D correction has been added to the $ATLAS$ colour. The CaT colour is defined as $(G - RVS) - (BP - RP)$. Bol_G is the bolometric correction in the G-band.

T_{eff}	$\log g$	$G - H_p$ ATLAS	$G - H_p$ $ATLAS_c$	Δ_{G-H_p} 3D-1D	$BP - RP$ ATLAS	$BP - RP$ $ATLAS_c$	Δ_{BP-RP} 3D-1D	CaT ATLAS	CaT $ATLAS_c$	Δ_{CaT} 3D-1D	Bol_G ATLAS	Bol_G $ATLAS_c$	Δ_{Bol_G} 3D-1D
3990	1.50	-0.544	-0.544	+0.000	+1.766	+1.766	+0.000	-0.347	-0.347	+0.001	-0.661	-0.640	+0.021
4503	1.50	-0.387	-0.379	+0.008	+1.388	+1.370	-0.018	-0.211	-0.200	+0.011	-0.361	-0.359	+0.003
4488	2.00	-0.377	-0.379	-0.002	+1.364	+1.369	+0.005	-0.201	-0.201	+0.000	-0.358	-0.362	-0.005
4523	2.50	-0.361	-0.363	-0.002	+1.322	+1.326	+0.005	-0.185	-0.187	-0.001	-0.336	-0.338	-0.003
4494	4.00	-0.372	-0.375	-0.003	+1.348	+1.355	+0.006	-0.195	-0.198	-0.004	-0.317	-0.319	-0.002
4522	4.50	-0.370	-0.371	-0.002	+1.342	+1.346	+0.004	-0.195	-0.197	-0.003	-0.300	-0.300	+0.001
4739	2.50	-0.315	-0.319	-0.004	+1.198	+1.208	+0.010	-0.145	-0.147	-0.002	-0.269	-0.282	-0.012
4891	1.80	-0.292	-0.294	-0.002	+1.133	+1.139	+0.006	-0.126	-0.126	-0.000	-0.230	-0.237	-0.007
4884	2.00	-0.291	-0.292	-0.000	+1.132	+1.133	+0.002	-0.125	-0.125	-0.001	-0.233	-0.230	+0.003
5018	2.50	-0.266	-0.272	-0.006	+1.059	+1.076	+0.018	-0.102	-0.108	-0.005	-0.208	-0.229	-0.021
4978	3.50	-0.276	-0.283	-0.008	+1.085	+1.107	+0.021	-0.113	-0.121	-0.008	-0.212	-0.233	-0.020
5160	4.00	-0.253	-0.261	-0.008	+1.017	+1.040	+0.023	-0.095	-0.105	-0.009	-0.183	-0.201	-0.018
4992	4.50	-0.284	-0.288	-0.004	+1.107	+1.117	+0.010	-0.123	-0.128	-0.005	-0.199	-0.207	-0.009
5501	2.50	-0.202	-0.203	-0.001	+0.867	+0.872	+0.005	-0.057	-0.057	-0.000	-0.136	-0.150	-0.014
5535	3.50	-0.203	-0.209	-0.006	+0.866	+0.885	+0.019	-0.055	-0.063	-0.008	-0.154	-0.162	-0.008
5476	4.00	-0.213	-0.224	-0.011	+0.895	+0.928	+0.034	-0.063	-0.075	-0.012	-0.158	-0.191	-0.033
5487	4.50	-0.216	-0.224	-0.008	+0.904	+0.927	+0.023	-0.068	-0.077	-0.009	-0.152	-0.164	-0.012
5789	3.70	-0.179	-0.184	-0.005	+0.788	+0.805	+0.017	-0.040	-0.047	-0.007	-0.140	-0.154	-0.014
5873	3.50	-0.170	-0.173	-0.003	+0.761	+0.771	+0.011	-0.036	-0.041	-0.005	-0.131	-0.141	-0.010
5846	4.00	-0.176	-0.181	-0.006	+0.777	+0.796	+0.019	-0.038	-0.046	-0.008	-0.141	-0.151	-0.009
5920	4.50	-0.172	-0.179	-0.007	+0.765	+0.786	+0.021	-0.037	-0.046	-0.009	-0.141	-0.144	-0.003
6306	3.50	-0.133	-0.134	-0.002	+0.638	+0.644	+0.006	-0.020	-0.022	-0.001	-0.104	-0.118	-0.013
6269	4.00	-0.140	-0.144	-0.003	+0.660	+0.672	+0.012	-0.021	-0.026	-0.005	-0.125	-0.136	-0.011
6242	4.00	-0.142	-0.145	-0.003	+0.668	+0.676	+0.009	-0.022	-0.026	-0.004	-0.126	-0.134	-0.008
6273	4.50	-0.144	-0.150	-0.006	+0.669	+0.689	+0.020	-0.021	-0.029	-0.008	-0.136	-0.148	-0.011
6408	4.00	-0.130	-0.133	-0.003	+0.625	+0.635	+0.011	-0.017	-0.020	-0.003	-0.120	-0.137	-0.017
6550	4.50	-0.124	-0.127	-0.004	+0.601	+0.613	+0.012	-0.013	-0.019	-0.005	-0.132	-0.138	-0.006

We show in Fig.A.1 the bolometric corrections to Gaia G magnitude as a function of effective temperature. This Figure supersedes and replaces the analogous Figure 2 of Bonifacio et al. (2017), in which a large correction was predicted for the hottest solar metallicity models. Those corrections were in fact spurious and due to the use of low resolution LHD models. It can be appreciated as for solar-type solar-metallicity stars most of the stellar flux is emitted in the G bandpass, leading to essentially zero bolometric corrections. For cooler stars a larger part of the flux is emitted in the near infra-red, outside the G bandpass and thus the bolometric correction becomes non-negligible. In a similar manner we note that for metal-poor stars, even of solar-type, the bolometric correction is small, but non-negligible. This is due to the fact that because of the lower opacity more flux is emitted in the ultra-violet, outside the G bandpass.

Table A.13. Colours and corrections for SDSS $u-g$, $g-z$, $g-r$ and $g-i$ for metallicity $[M/H]=0.0$. In columns ATLAS_c the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$u-g$ ATLAS	$u-g$ ATLAS _c	Δ_{u-g} 3D-1D	$g-z$ ATLAS	$g-z$ ATLAS _c	Δ_{g-z} 3D-1D	$g-r$ ATLAS	$g-r$ ATLAS _c	Δ_{g-r} 3D-1D	$g-i$ ATLAS	$g-i$ ATLAS _c	Δ_{g-i} 3D-1D
3635	1.00	+3.578	+3.479	-0.099	+2.869	+2.906	+0.037	+1.492	+1.485	-0.007	+2.340	+2.355	+0.014
3813	4.00	+2.579	+2.542	-0.037	+2.484	+2.468	-0.016	+1.294	+1.312	+0.017	+2.031	+2.025	-0.007
4018	1.50	+3.324	+3.197	-0.127	+2.075	+2.030	-0.045	+1.322	+1.280	-0.042	+1.781	+1.737	-0.044
3964	4.50	+2.548	+2.483	-0.066	+2.233	+2.226	-0.008	+1.301	+1.275	-0.025	+1.879	+1.864	-0.015
4501	2.00	+2.575	+2.502	-0.073	+1.470	+1.445	-0.025	+0.998	+0.978	-0.020	+1.297	+1.275	-0.022
4477	2.50	+2.487	+2.457	-0.030	+1.485	+1.475	-0.010	+0.997	+0.987	-0.010	+1.306	+1.297	-0.009
4478	4.00	+2.264	+2.239	-0.025	+1.490	+1.489	-0.001	+0.995	+0.983	-0.011	+1.311	+1.306	-0.004
4509	4.50	+2.279	+2.271	-0.008	+1.499	+1.520	+0.021	+1.010	+1.009	-0.001	+1.325	+1.338	+0.012
4582	3.20	+2.267	+2.241	-0.025	+1.379	+1.376	-0.003	+0.932	+0.926	-0.006	+1.219	+1.216	-0.004
4777	3.20	+2.080	+2.059	-0.022	+1.210	+1.210	+0.001	+0.834	+0.832	-0.003	+1.080	+1.081	+0.000
4968	2.50	+1.977	+1.935	-0.043	+1.077	+1.069	-0.008	+0.764	+0.753	-0.011	+0.976	+0.968	-0.008
5038	3.00	+1.860	+1.835	-0.025	+1.021	+1.017	-0.004	+0.726	+0.719	-0.007	+0.928	+0.924	-0.004
4923	3.50	+1.926	+1.906	-0.019	+1.098	+1.095	-0.002	+0.768	+0.764	-0.004	+0.989	+0.987	-0.002
4954	4.00	+1.882	+1.872	-0.010	+1.077	+1.080	+0.003	+0.756	+0.757	+0.001	+0.973	+0.976	+0.004
4982	4.50	+1.860	+1.861	+0.001	+1.065	+1.077	+0.013	+0.751	+0.760	+0.009	+0.965	+0.978	+0.013
5432	3.50	+1.523	+1.510	-0.012	+0.771	+0.768	-0.003	+0.574	+0.568	-0.006	+0.722	+0.718	-0.004
5475	4.00	+1.475	+1.476	+0.001	+0.749	+0.752	+0.003	+0.562	+0.561	-0.001	+0.704	+0.707	+0.002
5488	4.50	+1.464	+1.470	+0.006	+0.746	+0.760	+0.014	+0.562	+0.570	+0.008	+0.703	+0.716	+0.013
5774	4.40	+1.287	+1.297	+0.010	+0.594	+0.607	+0.013	+0.470	+0.477	+0.007	+0.578	+0.589	+0.011
5884	3.50	+1.289	+1.295	+0.006	+0.524	+0.522	-0.002	+0.425	+0.418	-0.007	+0.517	+0.514	-0.004
5928	4.00	+1.232	+1.241	+0.009	+0.510	+0.515	+0.004	+0.419	+0.418	-0.000	+0.508	+0.511	+0.003
5865	4.50	+1.237	+1.253	+0.015	+0.550	+0.561	+0.011	+0.445	+0.450	+0.006	+0.542	+0.552	+0.010
6228	4.00	+1.136	+1.149	+0.012	+0.368	+0.374	+0.007	+0.334	+0.333	-0.001	+0.390	+0.394	+0.004
6233	4.50	+1.092	+1.108	+0.015	+0.379	+0.385	+0.006	+0.343	+0.345	+0.002	+0.402	+0.407	+0.005
6484	4.00	+1.091	+1.104	+0.013	+0.254	+0.259	+0.005	+0.267	+0.265	-0.002	+0.295	+0.297	+0.002
6456	4.50	+1.038	+1.056	+0.018	+0.283	+0.291	+0.007	+0.287	+0.289	+0.002	+0.323	+0.328	+0.005
6726	4.30	+1.029	+1.047	+0.019	+0.165	+0.159	-0.006	+0.218	+0.212	-0.006	+0.224	+0.218	-0.007

Table A.14. Colours and corrections for SDSS $u-g$, $g-z$, $g-r$ and $g-i$ for metallicity $[M/H]=-1.0$. In columns ATLAS_c the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$u-g$ ATLAS	$u-g$ ATLAS _c	Δ_{u-g} 3D-1D	$g-z$ ATLAS	$g-z$ ATLAS _c	Δ_{g-z} 3D-1D	$g-r$ ATLAS	$g-r$ ATLAS _c	Δ_{g-r} 3D-1D	$g-i$ ATLAS	$g-i$ ATLAS _c	Δ_{g-i} 3D-1D
3790	4.00	+2.311	+2.272	-0.039	+2.334	+2.318	-0.016	+1.346	+1.320	-0.026	+1.951	+1.930	-0.021
4040	1.50	+2.575	+2.488	-0.086	+1.890	+1.845	-0.046	+1.171	+1.134	-0.037	+1.615	+1.573	-0.042
4001	4.50	+2.208	+2.169	-0.039	+2.052	+2.046	-0.006	+1.270	+1.242	-0.027	+1.758	+1.741	-0.017
4485	2.00	+1.938	+1.903	-0.035	+1.370	+1.353	-0.016	+0.865	+0.851	-0.013	+1.186	+1.171	-0.014
4490	2.50	+1.847	+1.841	-0.006	+1.355	+1.357	+0.002	+0.853	+0.853	+0.000	+1.171	+1.174	+0.002
4525	4.00	+1.735	+1.760	+0.025	+1.344	+1.380	+0.036	+0.860	+0.884	+0.024	+1.168	+1.201	+0.032
4499	4.50	+1.784	+1.786	+0.002	+1.401	+1.417	+0.016	+0.907	+0.914	+0.006	+1.222	+1.236	+0.014
4754	2.50	+1.627	+1.622	-0.006	+1.137	+1.137	-0.000	+0.730	+0.728	-0.002	+0.994	+0.995	+0.001
4993	2.50	+1.456	+1.466	+0.010	+0.969	+0.973	+0.005	+0.635	+0.636	+0.001	+0.858	+0.863	+0.005
4930	3.50	+1.427	+1.442	+0.015	+1.007	+1.022	+0.015	+0.655	+0.667	+0.012	+0.888	+0.903	+0.015
4986	4.00	+1.385	+1.416	+0.031	+0.973	+1.004	+0.031	+0.639	+0.663	+0.024	+0.862	+0.892	+0.030
5061	4.50	+1.348	+1.372	+0.023	+0.931	+0.952	+0.021	+0.620	+0.640	+0.020	+0.831	+0.853	+0.022
5481	3.50	+1.125	+1.162	+0.036	+0.682	+0.702	+0.021	+0.473	+0.485	+0.012	+0.625	+0.645	+0.019
5532	4.00	+1.082	+1.112	+0.030	+0.662	+0.676	+0.014	+0.463	+0.473	+0.010	+0.610	+0.625	+0.014
5473	4.50	+1.106	+1.136	+0.030	+0.699	+0.715	+0.017	+0.488	+0.502	+0.014	+0.642	+0.660	+0.017
5772	4.40	+0.980	+1.009	+0.029	+0.552	+0.569	+0.017	+0.404	+0.416	+0.012	+0.523	+0.541	+0.017
5890	3.50	+1.012	+1.060	+0.047	+0.478	+0.501	+0.022	+0.357	+0.368	+0.011	+0.459	+0.478	+0.019
5850	4.00	+0.976	+1.014	+0.038	+0.508	+0.518	+0.009	+0.376	+0.382	+0.006	+0.486	+0.495	+0.009
5923	4.50	+0.929	+0.976	+0.046	+0.487	+0.510	+0.023	+0.367	+0.384	+0.016	+0.471	+0.493	+0.022
6210	3.50	+0.986	+1.021	+0.035	+0.333	+0.332	-0.001	+0.276	+0.272	-0.004	+0.341	+0.338	-0.002
6261	4.00	+0.919	+0.961	+0.043	+0.340	+0.341	+0.002	+0.282	+0.283	+0.001	+0.349	+0.351	+0.002
6236	4.00	+0.917	+0.952	+0.035	+0.329	+0.342	+0.012	+0.276	+0.282	+0.006	+0.340	+0.351	+0.011
6238	4.50	+0.865	+0.910	+0.045	+0.355	+0.371	+0.017	+0.293	+0.305	+0.012	+0.363	+0.379	+0.016
6503	4.00	+0.916	+0.951	+0.035	+0.231	+0.234	+0.003	+0.223	+0.223	+0.001	+0.261	+0.263	+0.003
6456	4.50	+0.847	+0.895	+0.048	+0.269	+0.290	+0.020	+0.246	+0.259	+0.013	+0.294	+0.313	+0.018
6730	4.30	+0.880	+0.901	+0.022	+0.157	+0.163	+0.006	+0.185	+0.187	+0.002	+0.203	+0.207	+0.004

Table A.15. Colours and corrections for SDSS $u-g$, $g-z$, $g-r$ and $g-i$ for metallicity $[M/H]=-2.0$. In columns ATLAS_c the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$u-g$ ATLAS	$u-g$ ATLAS _c	Δ_{u-g} 3D-1D	$g-z$ ATLAS	$g-z$ ATLAS _c	Δ_{g-z} 3D-1D	$g-r$ ATLAS	$g-r$ ATLAS _c	Δ_{g-r} 3D-1D	$g-i$ ATLAS	$g-i$ ATLAS _c	Δ_{g-i} 3D-1D
4001	1.50	+2.411	+2.364	-0.047	+1.912	+1.894	-0.019	+1.158	+1.148	-0.010	+1.623	+1.610	-0.013
4000	4.50	+2.076	+2.072	-0.003	+1.951	+1.950	-0.001	+1.209	+1.207	-0.002	+1.676	+1.675	-0.001
4506	1.50	+1.824	+1.784	-0.040	+1.375	+1.351	-0.024	+0.851	+0.828	-0.024	+1.185	+1.162	-0.023
4506	2.00	+1.644	+1.639	-0.006	+1.327	+1.338	+0.011	+0.810	+0.814	+0.004	+1.135	+1.145	+0.010
4478	2.50	+1.701	+1.706	+0.005	+1.330	+1.346	+0.016	+0.817	+0.830	+0.013	+1.142	+1.159	+0.017
4504	4.00	+1.589	+1.605	+0.016	+1.320	+1.345	+0.025	+0.818	+0.839	+0.021	+1.137	+1.163	+0.026
4539	4.50	+1.594	+1.608	+0.014	+1.320	+1.339	+0.019	+0.825	+0.841	+0.016	+1.142	+1.162	+0.020
4743	2.50	+1.417	+1.415	-0.002	+1.122	+1.122	+0.001	+0.696	+0.695	-0.000	+0.970	+0.972	+0.002
4778	2.00	+1.457	+1.500	+0.043	+1.114	+1.094	-0.020	+0.695	+0.682	-0.013	+0.967	+0.948	-0.018
5024	2.50	+1.219	+1.246	+0.027	+0.927	+0.929	+0.002	+0.585	+0.584	-0.001	+0.811	+0.814	+0.002
4976	3.50	+1.206	+1.233	+0.027	+0.959	+0.979	+0.020	+0.607	+0.622	+0.015	+0.839	+0.859	+0.020
4955	4.00	+1.233	+1.267	+0.034	+0.981	+1.009	+0.028	+0.623	+0.646	+0.023	+0.860	+0.887	+0.028
5013	4.50	+1.217	+1.243	+0.026	+0.958	+0.977	+0.019	+0.614	+0.633	+0.018	+0.844	+0.865	+0.021
5502	2.50	+1.048	+1.084	+0.036	+0.640	+0.635	-0.005	+0.421	+0.413	-0.008	+0.576	+0.570	-0.006
5505	3.50	+0.943	+0.989	+0.046	+0.657	+0.667	+0.010	+0.434	+0.442	+0.008	+0.592	+0.603	+0.011
5472	4.00	+0.939	+0.990	+0.051	+0.685	+0.715	+0.030	+0.454	+0.475	+0.021	+0.618	+0.646	+0.028
5479	4.50	+0.942	+0.988	+0.046	+0.695	+0.716	+0.021	+0.464	+0.483	+0.019	+0.629	+0.651	+0.022
5772	3.70	+0.872	+0.934	+0.062	+0.532	+0.551	+0.020	+0.366	+0.381	+0.016	+0.491	+0.512	+0.020
5734	4.40	+0.845	+0.897	+0.051	+0.568	+0.592	+0.024	+0.391	+0.410	+0.019	+0.524	+0.548	+0.024
5861	3.50	+0.883	+0.938	+0.055	+0.486	+0.495	+0.010	+0.340	+0.347	+0.007	+0.453	+0.463	+0.010
5856	4.00	+0.836	+0.895	+0.059	+0.502	+0.522	+0.020	+0.351	+0.367	+0.016	+0.468	+0.489	+0.021
5923	4.50	+0.796	+0.855	+0.058	+0.486	+0.512	+0.026	+0.346	+0.368	+0.021	+0.459	+0.485	+0.026
6287	3.50	+0.892	+0.934	+0.042	+0.301	+0.307	+0.006	+0.241	+0.243	+0.002	+0.305	+0.310	+0.005
6206	4.00	+0.814	+0.856	+0.042	+0.354	+0.364	+0.009	+0.272	+0.278	+0.006	+0.350	+0.359	+0.009
6323	4.50	+0.755	+0.806	+0.051	+0.326	+0.344	+0.019	+0.260	+0.275	+0.015	+0.330	+0.349	+0.019
6534	4.00	+0.834	+0.873	+0.039	+0.225	+0.232	+0.007	+0.205	+0.209	+0.004	+0.247	+0.254	+0.007
6533	4.50	+0.758	+0.805	+0.047	+0.247	+0.263	+0.016	+0.219	+0.232	+0.013	+0.267	+0.284	+0.017




Fig. A.1. Bolometric corrections in the Gaia G band for four sets of models: $\log g = 2.5$ (right panel) and $\log g = 4.5$ (left panel) at $[M/H]=0.0$ and $[M/H]=-2.0$. Black symbols are computed from our ATLAS models, the lines are the bolometric corrections computed from the Castelli & Kurucz (2003) grid. The solid line refers to models of solar metallicity, the dashed line to models with metallicity -2.0 . The red symbols are the values to which we added the 3D correction.

Table A.16. Colours and corrections for SDSS $u-g$, $g-z$, $g-r$ and $g-i$ for metallicity $[M/H]=-3.0$. In columns ATLAS_c the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$u-g$ ATLAS	$u-g$ ATLAS _c	Δ_{u-g} 3D-1D	$g-z$ ATLAS	$g-z$ ATLAS _c	Δ_{g-z} 3D-1D	$g-r$ ATLAS	$g-r$ ATLAS _c	Δ_{g-r} 3D-1D	$g-i$ ATLAS	$g-i$ ATLAS _c	Δ_{g-i} 3D-1D
3990	1.50	+2.543	+2.560	+0.018	+2.054	+2.050	-0.004	+1.241	+1.237	-0.004	+1.748	+1.746	-0.002
4503	1.50	+1.748	+1.691	-0.058	+1.421	+1.389	-0.032	+0.868	+0.837	-0.031	+1.220	+1.189	-0.032
4488	2.00	+1.660	+1.647	-0.013	+1.385	+1.391	+0.006	+0.841	+0.841	-0.001	+1.186	+1.191	+0.005
4523	2.50	+1.544	+1.530	-0.014	+1.320	+1.326	+0.006	+0.800	+0.803	+0.003	+1.130	+1.136	+0.007
4494	4.00	+1.516	+1.526	+0.009	+1.361	+1.371	+0.010	+0.826	+0.835	+0.009	+1.169	+1.180	+0.011
4522	4.50	+1.506	+1.513	+0.008	+1.349	+1.355	+0.006	+0.823	+0.829	+0.006	+1.163	+1.170	+0.007
4739	2.50	+1.316	+1.324	+0.008	+1.131	+1.145	+0.014	+0.689	+0.694	+0.005	+0.972	+0.984	+0.012
4891	1.80	+1.294	+1.310	+0.016	+1.029	+1.035	+0.007	+0.631	+0.630	-0.001	+0.889	+0.894	+0.005
4884	2.00	+1.263	+1.296	+0.034	+1.028	+1.027	-0.001	+0.629	+0.626	-0.003	+0.887	+0.888	+0.001
5018	2.50	+1.114	+1.160	+0.047	+0.921	+0.945	+0.023	+0.566	+0.578	+0.012	+0.798	+0.819	+0.021
4978	3.50	+1.099	+1.131	+0.032	+0.962	+0.992	+0.030	+0.594	+0.614	+0.020	+0.835	+0.864	+0.028
5160	4.00	+0.997	+1.036	+0.038	+0.860	+0.891	+0.031	+0.540	+0.562	+0.021	+0.755	+0.784	+0.029
4992	4.50	+1.118	+1.141	+0.022	+0.990	+1.005	+0.015	+0.619	+0.630	+0.011	+0.867	+0.881	+0.014
5501	2.50	+0.978	+1.027	+0.050	+0.635	+0.639	+0.004	+0.407	+0.405	-0.002	+0.565	+0.567	+0.002
5535	3.50	+0.860	+0.915	+0.055	+0.638	+0.661	+0.023	+0.412	+0.428	+0.016	+0.570	+0.593	+0.022
5476	4.00	+0.858	+0.912	+0.054	+0.681	+0.726	+0.045	+0.440	+0.469	+0.030	+0.608	+0.649	+0.042
5487	4.50	+0.858	+0.897	+0.039	+0.693	+0.723	+0.031	+0.451	+0.473	+0.022	+0.621	+0.651	+0.030
5789	3.70	+0.809	+0.858	+0.050	+0.523	+0.544	+0.020	+0.352	+0.365	+0.013	+0.479	+0.499	+0.019
5873	3.50	+0.826	+0.875	+0.049	+0.481	+0.492	+0.011	+0.329	+0.336	+0.007	+0.444	+0.456	+0.011
5846	4.00	+0.779	+0.831	+0.052	+0.507	+0.529	+0.022	+0.346	+0.361	+0.016	+0.468	+0.490	+0.022
5920	4.50	+0.741	+0.786	+0.045	+0.490	+0.516	+0.026	+0.340	+0.360	+0.020	+0.457	+0.483	+0.026
6306	3.50	+0.853	+0.890	+0.037	+0.296	+0.302	+0.007	+0.233	+0.234	+0.001	+0.297	+0.302	+0.005
6269	4.00	+0.773	+0.815	+0.042	+0.332	+0.346	+0.013	+0.255	+0.264	+0.009	+0.329	+0.342	+0.013
6242	4.00	+0.771	+0.810	+0.038	+0.343	+0.352	+0.009	+0.260	+0.267	+0.006	+0.338	+0.347	+0.009
6273	4.50	+0.713	+0.758	+0.044	+0.349	+0.372	+0.023	+0.266	+0.283	+0.017	+0.344	+0.367	+0.023
6408	4.00	+0.784	+0.812	+0.028	+0.278	+0.291	+0.013	+0.227	+0.234	+0.007	+0.286	+0.297	+0.011
6550	4.50	+0.723	+0.761	+0.038	+0.245	+0.259	+0.014	+0.214	+0.224	+0.010	+0.263	+0.277	+0.014

Table A.17. Colours and corrections for SDSS $r - i$, $i - z$, $u - i$ and $u - z$ for metallicity $[M/H]=0.0$. In columns ATLAS_c the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$r - i$ ATLAS	$r - i$ ATLAS _c	Δ_{r-i} 3D-1D	$i - z$ ATLAS	$i - z$ ATLAS _c	Δ_{i-z} 3D-1D	$u - i$ ATLAS	$u - i$ ATLAS _c	Δ_{u-i} 3D-1D	$u - z$ ATLAS	$u - z$ ATLAS _c	Δ_{u-z} 3D-1D
3635	1.00	+0.848	+0.869	+0.021	+0.529	+0.551	+0.023	+5.918	+5.833	-0.084	+6.446	+6.385	-0.062
3813	4.00	+0.737	+0.713	-0.024	+0.452	+0.443	-0.009	+4.611	+4.567	-0.044	+5.063	+5.010	-0.053
4018	1.50	+0.459	+0.457	-0.002	+0.293	+0.292	-0.001	+5.105	+4.935	-0.171	+5.399	+5.227	-0.172
3964	4.50	+0.578	+0.588	+0.010	+0.354	+0.362	+0.008	+4.427	+4.346	-0.081	+4.782	+4.708	-0.073
4501	2.00	+0.299	+0.298	-0.002	+0.173	+0.170	-0.003	+3.872	+3.778	-0.094	+4.045	+3.948	-0.098
4477	2.50	+0.309	+0.309	+0.001	+0.179	+0.178	-0.001	+3.793	+3.754	-0.039	+3.972	+3.932	-0.040
4478	4.00	+0.316	+0.323	+0.007	+0.179	+0.183	+0.004	+3.574	+3.545	-0.029	+3.753	+3.728	-0.026
4509	4.50	+0.315	+0.328	+0.013	+0.173	+0.182	+0.009	+3.605	+3.609	+0.004	+3.778	+3.791	+0.013
4582	3.20	+0.287	+0.290	+0.003	+0.160	+0.161	+0.001	+3.486	+3.457	-0.029	+3.646	+3.618	-0.028
4777	3.20	+0.246	+0.249	+0.003	+0.129	+0.130	+0.000	+3.161	+3.139	-0.021	+3.290	+3.269	-0.021
4968	2.50	+0.212	+0.215	+0.003	+0.101	+0.101	-0.000	+2.953	+2.903	-0.051	+3.054	+3.004	-0.051
5038	3.00	+0.202	+0.204	+0.003	+0.093	+0.093	+0.000	+2.788	+2.759	-0.030	+2.882	+2.852	-0.030
4923	3.50	+0.221	+0.223	+0.002	+0.109	+0.108	-0.001	+2.915	+2.893	-0.021	+3.023	+3.001	-0.022
4954	4.00	+0.217	+0.219	+0.002	+0.104	+0.104	-0.001	+2.854	+2.848	-0.006	+2.958	+2.951	-0.007
4982	4.50	+0.214	+0.218	+0.004	+0.100	+0.100	+0.000	+2.825	+2.838	+0.013	+2.925	+2.938	+0.013
5432	3.50	+0.148	+0.150	+0.002	+0.049	+0.049	+0.000	+2.245	+2.229	-0.016	+2.294	+2.278	-0.016
5475	4.00	+0.143	+0.146	+0.003	+0.045	+0.045	+0.001	+2.180	+2.183	+0.003	+2.224	+2.228	+0.004
5488	4.50	+0.141	+0.146	+0.005	+0.043	+0.044	+0.001	+2.167	+2.186	+0.019	+2.210	+2.230	+0.020
5774	4.40	+0.108	+0.112	+0.005	+0.016	+0.018	+0.002	+1.864	+1.886	+0.022	+1.880	+1.904	+0.024
5884	3.50	+0.092	+0.096	+0.003	+0.006	+0.008	+0.002	+1.807	+1.809	+0.002	+1.813	+1.817	+0.004
5928	4.00	+0.089	+0.093	+0.003	+0.003	+0.004	+0.001	+1.740	+1.752	+0.012	+1.743	+1.756	+0.014
5865	4.50	+0.098	+0.102	+0.004	+0.008	+0.009	+0.001	+1.780	+1.805	+0.025	+1.787	+1.814	+0.026
6228	4.00	+0.056	+0.061	+0.005	-0.022	-0.020	+0.003	+1.526	+1.543	+0.016	+1.504	+1.523	+0.019
6233	4.50	+0.059	+0.062	+0.003	-0.023	-0.021	+0.001	+1.494	+1.514	+0.021	+1.471	+1.493	+0.022
6484	4.00	+0.028	+0.032	+0.004	-0.042	-0.039	+0.003	+1.386	+1.401	+0.015	+1.345	+1.362	+0.018
6456	4.50	+0.036	+0.039	+0.003	-0.040	-0.038	+0.002	+1.361	+1.384	+0.023	+1.321	+1.346	+0.025
6726	4.30	+0.006	+0.006	-0.000	-0.059	-0.058	+0.001	+1.253	+1.265	+0.012	+1.194	+1.207	+0.013

Table A.18. Colours and corrections for SDSS $r - i$, $i - z$, $u - i$ and $u - z$ for metallicity $[M/H]=-1.0$. In columns ATLAS_c the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$r - i$ ATLAS	$r - i$ ATLAS _c	Δ_{r-i} 3D-1D	$i - z$ ATLAS	$i - z$ ATLAS _c	Δ_{i-z} 3D-1D	$u - i$ ATLAS	$u - i$ ATLAS _c	Δ_{u-i} 3D-1D	$u - z$ ATLAS	$u - z$ ATLAS _c	Δ_{u-z} 3D-1D
3790	4.00	+0.605	+0.610	+0.005	+0.383	+0.388	+0.005	+4.262	+4.201	-0.060	+4.645	+4.589	-0.056
4040	1.50	+0.444	+0.439	-0.005	+0.276	+0.272	-0.004	+4.189	+4.061	-0.128	+4.465	+4.333	-0.132
4001	4.50	+0.488	+0.499	+0.011	+0.294	+0.305	+0.011	+3.966	+3.911	-0.055	+4.260	+4.216	-0.044
4485	2.00	+0.321	+0.320	-0.001	+0.184	+0.182	-0.002	+3.124	+3.074	-0.050	+3.308	+3.256	-0.052
4490	2.50	+0.318	+0.320	+0.002	+0.184	+0.183	-0.001	+3.018	+3.014	-0.004	+3.202	+3.198	-0.005
4525	4.00	+0.308	+0.316	+0.008	+0.176	+0.180	+0.003	+2.903	+2.960	+0.058	+3.079	+3.140	+0.061
4499	4.50	+0.315	+0.322	+0.007	+0.179	+0.181	+0.002	+3.006	+3.022	+0.016	+3.185	+3.203	+0.018
4754	2.50	+0.265	+0.267	+0.002	+0.143	+0.142	-0.001	+2.622	+2.617	-0.005	+2.764	+2.759	-0.006
4993	2.50	+0.223	+0.227	+0.004	+0.110	+0.110	+0.000	+2.314	+2.329	+0.014	+2.425	+2.439	+0.015
4930	3.50	+0.233	+0.237	+0.004	+0.119	+0.118	-0.000	+2.315	+2.345	+0.031	+2.434	+2.464	+0.030
4986	4.00	+0.223	+0.229	+0.006	+0.111	+0.112	+0.001	+2.247	+2.308	+0.061	+2.358	+2.420	+0.062
5061	4.50	+0.211	+0.213	+0.002	+0.100	+0.099	-0.001	+2.179	+2.225	+0.045	+2.280	+2.324	+0.044
5481	3.50	+0.153	+0.159	+0.007	+0.056	+0.058	+0.002	+1.751	+1.806	+0.055	+1.807	+1.864	+0.057
5532	4.00	+0.147	+0.151	+0.004	+0.051	+0.051	-0.000	+1.692	+1.737	+0.045	+1.743	+1.788	+0.045
5473	4.50	+0.155	+0.158	+0.003	+0.056	+0.056	-0.001	+1.749	+1.796	+0.047	+1.805	+1.851	+0.047
5772	4.40	+0.120	+0.124	+0.005	+0.029	+0.029	-0.000	+1.503	+1.550	+0.047	+1.532	+1.578	+0.046
5890	3.50	+0.102	+0.110	+0.008	+0.019	+0.022	+0.003	+1.472	+1.538	+0.066	+1.491	+1.561	+0.070
5850	4.00	+0.110	+0.113	+0.003	+0.022	+0.022	-0.000	+1.462	+1.509	+0.047	+1.485	+1.531	+0.047
5923	4.50	+0.103	+0.109	+0.006	+0.016	+0.017	+0.001	+1.400	+1.468	+0.069	+1.416	+1.485	+0.070
6210	3.50	+0.065	+0.067	+0.002	-0.007	-0.006	+0.001	+1.327	+1.360	+0.032	+1.320	+1.354	+0.034
6261	4.00	+0.067	+0.068	+0.001	-0.009	-0.009	-0.001	+1.267	+1.312	+0.045	+1.258	+1.303	+0.044
6236	4.00	+0.064	+0.069	+0.005	-0.011	-0.009	+0.002	+1.258	+1.303	+0.045	+1.247	+1.294	+0.047
6238	4.50	+0.070	+0.075	+0.004	-0.009	-0.008	+0.001	+1.228	+1.289	+0.061	+1.220	+1.281	+0.061
6503	4.00	+0.038	+0.040	+0.002	-0.029	-0.029	+0.001	+1.176	+1.214	+0.038	+1.147	+1.185	+0.038
6456	4.50	+0.048	+0.054	+0.005	-0.025	-0.023	+0.002	+1.141	+1.207	+0.066	+1.116	+1.184	+0.068
6730	4.30	+0.018	+0.021	+0.003	-0.046	-0.044	+0.002	+1.083	+1.108	+0.026	+1.037	+1.065	+0.028

Table A.19. Colours and corrections for SDSS $r - i$, $i - z$, $u - i$ and $u - z$ for metallicity $[M/H]=-2.0$. In columns ATLAS_c the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$r - i$ ATLAS	$r - i$ ATLAS _c	Δ_{r-i} 3D-1D	$i - z$ ATLAS	$i - z$ ATLAS _c	Δ_{i-z} 3D-1D	$u - i$ ATLAS	$u - i$ ATLAS _c	Δ_{u-i} 3D-1D	$u - z$ ATLAS	$u - z$ ATLAS _c	Δ_{u-z} 3D-1D
4001	1.50	+0.464	+0.462	-0.002	+0.290	+0.284	-0.006	+4.033	+3.974	-0.060	+4.323	+4.257	-0.066
4000	4.50	+0.467	+0.468	+0.001	+0.275	+0.275	+0.000	+3.752	+3.747	-0.004	+4.026	+4.022	-0.004
4506	1.50	+0.334	+0.334	+0.000	+0.190	+0.190	-0.001	+3.009	+2.945	-0.063	+3.199	+3.135	-0.064
4506	2.00	+0.325	+0.331	+0.006	+0.192	+0.192	+0.001	+2.780	+2.784	+0.004	+2.971	+2.976	+0.005
4478	2.50	+0.325	+0.329	+0.004	+0.188	+0.187	-0.001	+2.843	+2.865	+0.022	+3.031	+3.052	+0.020
4504	4.00	+0.319	+0.324	+0.005	+0.183	+0.182	-0.001	+2.726	+2.768	+0.041	+2.910	+2.950	+0.040
4539	4.50	+0.317	+0.321	+0.004	+0.177	+0.177	-0.000	+2.737	+2.770	+0.034	+2.914	+2.947	+0.033
4743	2.50	+0.274	+0.276	+0.002	+0.152	+0.150	-0.001	+2.387	+2.387	-0.000	+2.539	+2.537	-0.002
4778	2.00	+0.271	+0.266	-0.005	+0.148	+0.146	-0.002	+2.424	+2.448	+0.025	+2.571	+2.594	+0.023
5024	2.50	+0.226	+0.230	+0.003	+0.116	+0.116	-0.000	+2.030	+2.060	+0.030	+2.146	+2.175	+0.029
4976	3.50	+0.233	+0.237	+0.005	+0.120	+0.120	+0.000	+2.045	+2.092	+0.047	+2.165	+2.212	+0.047
4955	4.00	+0.237	+0.242	+0.005	+0.121	+0.121	-0.000	+2.092	+2.154	+0.062	+2.214	+2.276	+0.062
5013	4.50	+0.230	+0.232	+0.002	+0.113	+0.112	-0.002	+2.061	+2.108	+0.047	+2.175	+2.220	+0.045
5502	2.50	+0.154	+0.157	+0.003	+0.064	+0.065	+0.001	+1.624	+1.654	+0.030	+1.688	+1.719	+0.031
5505	3.50	+0.158	+0.161	+0.003	+0.065	+0.064	-0.001	+1.535	+1.592	+0.058	+1.600	+1.656	+0.057
5472	4.00	+0.164	+0.171	+0.007	+0.067	+0.069	+0.001	+1.557	+1.636	+0.079	+1.624	+1.704	+0.080
5479	4.50	+0.165	+0.168	+0.003	+0.066	+0.065	-0.001	+1.571	+1.639	+0.068	+1.636	+1.704	+0.067
5772	3.70	+0.126	+0.130	+0.005	+0.040	+0.040	-0.001	+1.364	+1.446	+0.082	+1.404	+1.486	+0.082
5734	4.40	+0.133	+0.138	+0.005	+0.044	+0.043	-0.000	+1.369	+1.445	+0.076	+1.413	+1.488	+0.076
5861	3.50	+0.114	+0.117	+0.003	+0.032	+0.032	-0.001	+1.336	+1.401	+0.065	+1.369	+1.433	+0.064
5856	4.00	+0.117	+0.122	+0.005	+0.033	+0.033	-0.001	+1.304	+1.384	+0.080	+1.337	+1.417	+0.079
5923	4.50	+0.112	+0.117	+0.005	+0.028	+0.027	-0.001	+1.255	+1.340	+0.085	+1.283	+1.367	+0.084
6287	3.50	+0.063	+0.066	+0.003	-0.004	-0.003	+0.001	+1.197	+1.244	+0.047	+1.193	+1.241	+0.048
6206	4.00	+0.078	+0.081	+0.003	+0.004	+0.004	+0.000	+1.165	+1.215	+0.051	+1.169	+1.220	+0.051
6323	4.50	+0.070	+0.074	+0.004	-0.004	-0.005	-0.001	+1.085	+1.155	+0.070	+1.080	+1.150	+0.070
6534	4.00	+0.042	+0.044	+0.003	-0.022	-0.022	+0.000	+1.081	+1.127	+0.046	+1.059	+1.105	+0.046
6533	4.50	+0.048	+0.052	+0.004	-0.020	-0.021	-0.000	+1.026	+1.089	+0.064	+1.005	+1.069	+0.063

Table A.20. Colours and corrections for SDSS $r - i$, $i - z$, $u - i$ and $u - z$ for metallicity $[M/H]=-3.0$. In columns ATLAS_c the 3D correction has been added to the ATLAS colour.

T_{eff}	$\log g$	$r - i$ ATLAS	$r - i$ ATLAS _c	Δ_{r-i} 3D-1D	$i - z$ ATLAS	$i - z$ ATLAS _c	Δ_{i-z} 3D-1D	$u - i$ ATLAS	$u - i$ ATLAS _c	Δ_{u-i} 3D-1D	$u - z$ ATLAS	$u - z$ ATLAS _c	Δ_{u-z} 3D-1D
3990	1.50	+0.507	+0.509	+0.002	+0.306	+0.304	-0.002	+4.290	+4.306	+0.016	+4.596	+4.611	+0.014
4503	1.50	+0.352	+0.352	-0.000	+0.201	+0.200	-0.000	+2.969	+2.879	-0.089	+3.169	+3.080	-0.090
4488	2.00	+0.344	+0.350	+0.006	+0.199	+0.200	+0.001	+2.846	+2.838	-0.008	+3.045	+3.038	-0.007
4523	2.50	+0.329	+0.334	+0.004	+0.190	+0.190	-0.000	+2.674	+2.666	-0.007	+2.864	+2.856	-0.008
4494	4.00	+0.343	+0.345	+0.002	+0.192	+0.191	-0.000	+2.686	+2.706	+0.020	+2.877	+2.897	+0.020
4522	4.50	+0.340	+0.340	+0.001	+0.186	+0.186	-0.001	+2.668	+2.683	+0.015	+2.855	+2.869	+0.014
4739	2.50	+0.283	+0.290	+0.007	+0.159	+0.161	+0.002	+2.288	+2.308	+0.020	+2.447	+2.469	+0.022
4891	1.80	+0.258	+0.264	+0.006	+0.140	+0.141	+0.001	+2.183	+2.204	+0.021	+2.323	+2.345	+0.022
4884	2.00	+0.258	+0.261	+0.003	+0.141	+0.140	-0.001	+2.150	+2.184	+0.034	+2.290	+2.324	+0.033
5018	2.50	+0.232	+0.241	+0.009	+0.123	+0.126	+0.003	+1.912	+1.979	+0.067	+2.035	+2.105	+0.070
4978	3.50	+0.241	+0.249	+0.008	+0.126	+0.128	+0.002	+1.934	+1.995	+0.061	+2.060	+2.123	+0.063
5160	4.00	+0.214	+0.222	+0.008	+0.105	+0.107	+0.002	+1.752	+1.820	+0.067	+1.857	+1.926	+0.069
4992	4.50	+0.247	+0.251	+0.003	+0.124	+0.124	+0.001	+1.985	+2.021	+0.036	+2.109	+2.145	+0.037
5501	2.50	+0.158	+0.162	+0.005	+0.070	+0.071	+0.002	+1.543	+1.595	+0.052	+1.612	+1.666	+0.054
5535	3.50	+0.158	+0.165	+0.007	+0.068	+0.068	+0.001	+1.430	+1.507	+0.077	+1.498	+1.576	+0.078
5476	4.00	+0.168	+0.180	+0.012	+0.073	+0.077	+0.004	+1.466	+1.562	+0.096	+1.539	+1.638	+0.100
5487	4.50	+0.170	+0.178	+0.008	+0.071	+0.073	+0.001	+1.480	+1.548	+0.069	+1.551	+1.621	+0.070
5789	3.70	+0.127	+0.133	+0.006	+0.044	+0.045	+0.001	+1.288	+1.357	+0.069	+1.332	+1.402	+0.070
5873	3.50	+0.116	+0.120	+0.004	+0.036	+0.036	-0.000	+1.271	+1.331	+0.060	+1.307	+1.367	+0.060
5846	4.00	+0.122	+0.128	+0.006	+0.039	+0.040	+0.000	+1.247	+1.321	+0.074	+1.286	+1.361	+0.075
5920	4.50	+0.116	+0.123	+0.006	+0.033	+0.033	-0.000	+1.198	+1.269	+0.071	+1.232	+1.302	+0.071
6306	3.50	+0.064	+0.068	+0.004	-0.001	+0.000	+0.002	+1.150	+1.192	+0.042	+1.149	+1.193	+0.043
6269	4.00	+0.074	+0.078	+0.004	+0.003	+0.004	+0.000	+1.102	+1.157	+0.055	+1.105	+1.161	+0.055
6242	4.00	+0.077	+0.080	+0.003	+0.006	+0.005	-0.000	+1.109	+1.157	+0.048	+1.115	+1.162	+0.048
6273	4.50	+0.078	+0.084	+0.006	+0.004	+0.004	+0.000	+1.058	+1.125	+0.067	+1.062	+1.130	+0.068
6408	4.00	+0.059	+0.063	+0.005	-0.008	-0.007	+0.001	+1.070	+1.109	+0.039	+1.062	+1.102	+0.040
6550	4.50	+0.049	+0.053	+0.004	-0.018	-0.018	-0.000	+0.986	+1.038	+0.051	+0.969	+1.020	+0.051